



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

582.


C 219

30/2

11-1-100
Vol. 1 - VI (156 -) MR

The Hopkins Library
presented to the
Yeland Stanford Junior University
by Timothy Hopkins.







PRODROMUS
SYSTEMATIS NATURALIS
REGNI VEGETABILIS.

PARS I.

Apud eosdem Bibliopolas venundatur :

- Regni vegetabilis Systema naturale : sive Ordines, genera et species plantarum secundum methodi naturalis normas digestarum et descriptarum, auct. de Candolle. In-8., tom. I et II. Paris. 1818 et 1822..... 27 fr.
- Icones selectæ plantarum quas in Systemate universali descripsit A. P. de Candolle, ex archetypis specimenibus à P. J. F. Turpin delineatæ et editæ à Benj. Delesert. Vol. primum exhibens *Ranunculaceas, Dilleniaceas, Magnoliaceas, Anonaceas et Menispermeas* cum 100 fig. gr. in-4. 1820..... 35 fr.
- Idem, charta velina, gr. in-4. 5ofr. — Idem, gr. in-fol. 70 fr.
- Exposition des Familles naturelles et de la Germination des Plantes, par Jaume-Saint-Hilaire ; contenant, 1°. la description de 2337 genres de botanique, et d'environ 4000 espèces les plus utiles et les plus intéressantes ; 2°. 117 planches, dont les figures dessinées par l'auteur et gravées par Sellier, représentent les caractères des familles naturelles et les différens modes de germination. 4 vol. grand in-8. pap. grand raisin, fig. en noir. 36 fr.
- Le même, 2 vol. in-4. les planches en couleur, papier ordinaire..... 96 fr.
- Le même, 2 vol. in-4. les planches en couleur, pap. vél. 192 fr.

**PRODROMUS
SYSTEMATIS NATURALIS
REGNI VEGETABILIS,**

SIVE

ENUMERATIO CONTRACTA

ORDINUM GENERUM SPECIERUMQUE PLANTARUM

**HUC USQUE CONITARUM, JUXTA METHODI NATURALIS NORMAS
DIGESTA**

AUCTORE ALG. PYRAMO DE CANDOLLE.

PARS PRIMA

Sine Thalamiflorarum Ordines LIV.

PARISIIS,

**Sumptibus Sociorum TREUTTEL ET WÜRTZ,
rue de Bourbon, n° 17.**

VENITQUE IN EORUMDEM BIBLIOPOLIIS

ARGENTORATI ET LONDINI.

M. DCCC. XXIV.

2 ✓



HB 601

IRÆFATIO.

VEGETABILIMUM nunc cognitorum immensus numerus, qui ad quinquaginta milli et amplius pervenit et in dies rapidè crescit; necessitas omnia hæc inventa uno in opere disponendi et ordinandi juxtà leges methodi naturalis, cujus commoda atque utilitates quotidie plenius in aperto sunt, me nuper induxerant ad aggrediendum *Systema universale Regni vegetabilis*, cujus duo jam volumin in lucem prodire. Ex hoc tempore multi meorum amicorum et nonnulli inter clarissimos artis herbariæ studiosos, sæpè mihi exhibuerunt, quod operis tanti necessaria diuturnitas et quotidiana plantarum repertarum auctio injiciebant cupidinè maturius fruendi opere multò breviorè, sed in brevitate nā completo (si hæc vox in Historiæ naturalis linguâ admitti aieat) quod offerret synopsis statûs hodierni scientiæ, et quæ uterentur quasi duce, aut saltem quasi indice; Viatores Hortorum Præfecti, Herbariorum Collectores, imò omnes Botanici qui inter innumeros libros à paucis annis editos, sæpè incerti vagantur. Quùm igitur in redigendo Systemate dudum omnia collecta et omnia documenta tali labori idonea percepissem, in mente habui interrumpere librum susceptum, ut hujus operis *Prodromum* conficerem, cupiens tamen Systema resumere statim ac, Prodromo edito, vota explevissem Botanicorum qui methodicam scientiæ suæ synopsis dudum efflagitat.

Non me fefellit hujus operis moles. Facile sensi quot in errores inciderem, dùm totum Vegetabile Regnum recognoscere et recensere vellem, atque pariter enumerarem, tùm species imperfectè cognitæ tùm eas quæ diligenter descriptæ fuerunt. Speravi tamen utilitatem talis operis, vel manco modo peracti hæc inevitabiliam excusaturam, et Botánicos exercitatos pro gratis habituros esse cautiones à me adhibitæ, ut lector facili examini asertiones meas subjicere posset, et ut ingenuè indicarem omnia quæ mihi videntur dubia et incerta, vel incompletè descripta inter stirpes quas Botanici vulgò ut cognitæ admittunt. Cupidus quàm citissimè in lucem emittendi hoc opus quod solummodò absolutum in pretio erit, libenter pro adjutoribus aliquot doctos Botánicos accepi: sic clarissimum *SERINGE*, laboribus circa *Salices* et *Cerealia Helvetiæ* nec non

circa Potentillas et Aconita editis notum, non piguit ordinem herbariis meis afferre et sparsas adnotationes ab eo et à mè ipso excerptas colligere; ipse insuper nonnullos ordines vel nonnulla genera, congruenter proposito à me inito, digessit. Clarissimi DUNAL, CHOISY, DE GINGINS, JTH, mihi hoc in primo volumine doctas aliquot monographias largiti sunt; paribus monographiis ditabunt, ut spero futura volumina, clarissimi COULTER, GUILLEMIN et DUBY; licce prætereà benevolè excerpserit omnia quæ recentiores scripsrunt circa Americæ borealis plantas. Clar. RICHARD et GUILLEMIN miro obsequio operam impenderunt in hujus libri typis corrigendis. Speciales animi grates ago clar. RICHARD qui, ut audivit me tali opere detentum, amicitiae perspicuum testimonium dedit, simile opus à se inchoatum prorsus derelinquendo. Ceterum quaecunque fragmenta mihi data sunt vel in posterum dabuntur, præfixum gerent nomen auctoris; horumque fragmentorum duntaxat particeps fui, communicando eorum auctoribus mea herbaria, nec non aliquot notulas pridem à mè collectas, et suscipiendo leviusculum laborem singulas partes harmonicè toto cum opere ordinandi.

Omnes species ex speciminibus in herbario meo asservatis descriptæ, designantur solito signo (v. s.); quâ cautione quisque poterit agnoscere quænam fiducia cuilibet articulo tribuenda sit, secernere ea quæ imputari mihi debeant iis ab in quibus merus narrator alienas observationes refero, agnoscere deniquè plantas quarum in herbario meo typis inveniet; poterunt et omnes quorum in animis esset specimina sicca mihi mittere, cognitis plantis quas possideo, statuere quæ mittere aut vicissim poscere velint.

Quotiescunque Peregrinator ad me misit plantam adhuc ignotam, hanc sub ejus nomine memoravi & totum inventi sui honorem ei impertitus sum. In genere enixè conatus sum cuique meorum antecessorum meorumque coëvorum debitam laboribus suis gloriam tribuere. Si cui tamen aliquoties ab hac normâ deflexisse videor, affirmo hanc omissionem nunquam ex professo factam, me libenter agnituram et reparaturam esse. Cum hoc opus tantum sit (ut ex titulo patet) *Prodromus* operis longioris et majoris momenti, pergratè accipiam omnes additiones, omnes emendationes, imò censuras, tum manuscriptas tum impressas, præsertim si non dedignentur Botanici opinionis suæ evidentia et corporea, ut ità dicam, argumenta subjungere.

Genève, 25 novembris 1823.

PRODROMUS

SYSTEMATIS UNIVERSALIS

REGNI VEGETABILIS.

I. PLANTÆ VASCULARES seu COTYLEDONEÆ. *DC. syst. 1. p. 121.*

CAR. Plantæ contextu celluloso vasisque lymphaticis constantes ; stomatibus donatæ ; partibus, aliis radicalibus deorsum directis, aliis caulinis sursum directis, compositæ ; foliis veris instructæ. Flores sæpius distincti et symetrici. Embryo intrâ spermodermium inclusus, cotyledonibus instructus.

Classis Prima. DICOTYLEDONEÆ seu EXOGENÆ. *DC. syst. 1. p. 123.*

CAR. Truncus constans corporibus duobus, ligneo nempe et corticali, juxtâ leges inversas crescentibus ; Corpus ligneum exhibens in centro medullam e quâ productiones medullares radiantes trans strata lignea annua, alia (lignum) interiora seniora duriora, alia (album) exteriora juniora molliora albida. Cortex constans complexu celluloso sub epidermide sito, et stratis corticalibus, aliis exterioribus senioribus, aliis interioribus junioribus nomine Libri insignitis. Folia nervis ramosis anastomosantibus. Flores distincti, symetrici, sæpius numero quinario dispositi. Embryo cotyledonibus, sæpius 2 oppositis, rarius plurimis verticillatis, constans.

Subclassis I. THALAMIFLORÆ.

Calyx poly-sepalus. Petala plurima distincta, staminaque thalamo inserta nec calyci accreta.

Ord. I. RANUNCULACEÆ. *Juss.* 231. *DC. syst.* 1. p. 127.

Perigonium duplex, utrumque liberum hypogynum. Sepala 3-6. Petala sepalorum numero æqualia dupla triplave, libera, æstivatione imbricatâ, rariùs abortâ nulla, nunc ex filamentis dilatatis facta et tunc plana, nunc ex antheris transformatis orta cucullata. Stamina hypogyna, libera, numero indefinita. Antheræ adnatæ, in R. veris extrorsæ. Pistilla ∞ toro inserta, abortu aut coalitione rariùs solitaria. Carpella alia pseudosperma, alia baccata, alia capsularia aut follicularia, 1- ∞ -sperma. Semina nunc solitaria erecta aut pendula, nunc plurima secus carpelli margines utrinquè seriâtîm disposita. Albumen corneum magnum. Embryo minimus, intrâ albuminis foveolam locatus. — Herbæ, suffrutices aut frutices sarmentosi. Radices fasciculatæ grumosæ aut fibrosæ. Folia alterna aut in Clematideis opposita, basi in vaginam semi-amplexicaulem dilatata, simplicia, sæpè variè secta. Pili o aut simplices.

Trib. I. CLEMATIDÆ. *DC. Syst.* 1. p. 131.

Calycis æstivatio valvata aut induplicata. Petala o aut plana. Antheræ lineares extrorsæ. Carpella 1-sperma indehiscencia stylo accreto caudata. Semen pendulum. — Caules sæpè sarmentosi. Folia opposita. Radices fibrosæ.

I. CLEMATIS. *DC. syst.* 1. p. 131.

Involucrum o aut calyciforme sub flore. Sepala 4-8 colorata. Petala o aut sepalis breviora. Cariopsides ∞ in caudam sæpius barbato-plumosam productæ. — Radices perennes. Folia exactè opposita.

Sect. I. FLAMMULA. *DC. l. c.*

Involucrum o. Petala o. Carpellorum caudæ longæ barbato-plumosæ. Embryones in semine cotyledones distantes.

§. 1. *Floribus paniculatis, foliis pinnatim bipinnatimve sectis.*

1. C. ERRECTA, (All! ped. n. 1078) Caule erecto, foliis pinnatisectis, segmentis petiolulatis ovato-lanceolatis integerrimis. 4 in collibus et sylvis Europæ australis. C. recta Lin. spec. 767. Jacq. fl. austr. t. 291. DC. syst. 1. p. 135. (v. v.)

β. C. surrecta præcocior, etc. Tourn. inst. 394.

γ. C. Hispanica surrecta, etc. Tourn. inst. 394.

δ. C. bracteosa Banks! herb.

2. C. FLAMMULA. (Lin! spec. 766.) Caule scandente, foliis pinnatisectis, segmentis glabris integris trilobiave, orbiculatis ovalibus oblongis sublinearibusve acutiusculis. 3 in Europæ australis siccis et sepibus. DC. syst. 1. p. 134. (v. v.)

a. rotundifolia, fol. segmentis suborbiculatis. C. fragrans Ten! fl. neap. prod. 32. vol. 1. t. 48.

β. vulgaris, fol. segmentis ovali aut oblongo-lanceolatis. C. Flammula Lin. spec. 766.

γ. maritima, fol. segmentis linearibus. C. maritima Lin. sp. 767 DC. fl. fr. 4. p. 873. 5. p. 562*.

- d. rubella*, Pers. ench. 2. p. 100.
s. caespitosa, segmentis minutis integris incisive. *C. caespitosa* Scop. fl. carn. edl. 2. v. 1. p. 389. *C. flammula* B. Bertol. auncn. ital. 256
3. *C. MARSONIANA* (DC. syst. 1. p. 135.) caule scandente, foliis pinnati-sectis, segmentis glabris subglaucais ovatis inciso-dentatis, trilobis trisectisve. $\zeta \cup$ in Africâ australiori. Flores ignoti. Pericarpia pubescentia, elliptica. (v. s. in herb. Banks.)
4. *C. ORIENTALIS* (Lin. spec. 765.) caule scandente, foliis pinnati-sectis, segmentis glaucais glabris cuneiformibus trilobis, lobis dentatis acuminatis. $\zeta \cup$ in Oriente et in Caucasi virgultis. Dill. elth. 144. t. 119. f. 145.* DC. syst. 1. p. 135. — Variat pedunculis 3-fidis aut 3-chotomo-paniculatis. (v. v. et s. in h. Tourn.)
5. *C. GLAUCA* (Willd! arb. 65. t. 4. f. 1.) caule scandente, foliis pinnati-sectis, segmentis glaucais glabris cuneiformibus sublobatis, lobis integris subobtusis, pedunculis 3-fidis. $\zeta \cup$ in Sibiria australis aridis et in Iberiâ. Foliorum segmenta rarò 3-loba obtusa aut submucronata. DC. syst. 1. p. 136. (v. v. et s.)
6. *C. PANICULATA* (Thunb. act. soc. Lin. 2. p. 337.) caule scandente, foliis pinnati-sectis, segmentis ovato-cordatis acutis integris, pedunculis paniculatis co-floris $\zeta \cup$ in Japoniâ propè Nagasaki. *C. Vitalba Japonica* Houtt. Pflanz. 7. p. 309. f. 2. DC. syst. 1. p. 136. (v. s.)
7. *C. LONGILOBA* (DC. syst. 1. p. 136) caule scandente, foliis pinnati-sectis, segmentis oblongo-lanceolatis acuminatis integerrimis glabris. \cup in Chinâ — Media videtur inter paniculatam et minorem. (v. s. in h. Lamb.) †
8. *C. MINOR* (Lour. fl. coch. 1. p. 422.*) caule scandente, foliis pinnati-sectis, segmentis ovato-lanceolatis integerrimis, pedunculis multifloris folii longitudine, ovariis quaternis. ζ in suburbiis Cantonienſibus Sinarum. DC. syst. 1. p. 136. Flores totius generis minimi. (v. s.)
9. *C. CHINENSIS* (Retz. obs. 2. p. 18. n. 53. t. 2.*) caule scandente, foliis pinnati-sectis, segmentis ovato-lanceolatis integerrimis, pedunculis paucifloris folio longioribus, ovariis subquaternis, caudis subnudis. $\zeta \cup$ in Chinâ et insulis Danes dictis. DC. syst. 1. p. 137. Bractea parvæ sub pedicellis et secus eos sitæ. (v. s. in h. Banks.)
10. *C. TENUIFLORA* (DC. syst. 1. p. 137.) caule scandente, foliis pinnati-sectis, segmentis ovatis obtusis integerrimis, pedunculis 3-fidis 3-floris, ovariis subquinis, caudis subbarbatis. ζ in Chinâ. *C. Flammula* var. Lin! herb. Pedicellus medius nudus, laterales infrâ medium bracteolas subulatas duas gerentes. (v. s. in h. Banks.)
11. *C. BREVICAUDATA*. (DC. syst. 1. p. 138) caule scandente, foliis pinnati-sectis, segmentis ovato-lanceolatis acuminatis grossè serratis, pedunculis patentibus folio brevioribus confertè paniculatis $\zeta \cup$ in Chinâ inter Peking et Jehu. Flores ignoti. (v. s. in h. Banks.)
12. *C. HERACLEOPOLIA* (DC. syst. 1. p. 138*) caule scandente, foliis pinnati-sectis, segmentis ovato-lanceolatis latè inciso-dentatis, ultimis 3-lobis, pedunculis erectis ramisque cinereo-velutinis. $\zeta \cup$ in Chinâ. Ovaria 15-20 in caudas barbato-plumosas breves desinentia. (v. s. in h. Banks.)
13. *C. GOURIANA* (Roxb! ined. in h. Lamb.) Caule scandente, foliis pinnati-sectis, segmentis ovato-lanceolatis basi cordatis sub 5-nerviis acuminatis integris subdentatisve; pedunculis pubescentibus folio sublongioribus. $\zeta \cup$ in Indiâ orientali. DC. syst. 1. p. 138.* Bractea ovata, basi in petiolum ferè angustata. (v. s.)
14. *C. VITALBA* (Lin. spec. 766.) caule scandente, foliis pinnati-sectis, segmentis ovato-lanceolatis dentato-incisis acuminatis, basi truncato-cordatis, pedunculis folio subbrevioribus. $\zeta \cup$ vulgarissima ad sepes et dumeta Europæ mediæ, Græciæ et Tauriæ. Jacq. fl. austr. t. 308. DC. syst. 1. p. 139. Ovaria glabra compressa. (v. v.)

β. integrata, segmentis paucissimis dentatis integerrimisve. *Vitis nigra* Fuchs. hist. 97. ic.

15. *C. GREWIAEFLOHA* (DC. syst. 1. p. 140.*) caule scandente, foliis pinnati-sectis utrinque velutino-pubescentibus, segmentis cordatis acutis acutè serratis, pedunculis folio brevioribus 3-5-floris. \S in Napauliâ. *C. Vitalba*? Buchan! ined. in h. Lamb. Pedunculi oppositi foliis breviores. (v. s.)
 16. *C. BUCHANIANA* (DC. syst. 1. p. 140) caule scandente, foliis pinnati-sectis subtus velutinis, supernè glabris, segmentis subcordatis crenatis, pedunculis folii longitudine 7-9-floris. \S in Napauliâ. *C. bucamara* var. Buchan! (v. s. in h. Lamb.)
 17. *C. LOSÆFOLIA* (DC. syst. 1. p. 140) caule scandente, foliis utrinquè villosissimis, inferioribus cordatis palmatis 3-5-fidis, superioribus trisectis, segmentis subcordatis sub 3-fidis. \S in Napauliâ. *C. bucamara* var. Buchan! ined. in h. Lamb. (v. s.) †
 18. *C. CONNATA*, caule scandente, foliis glabris pinnati-sectis, segmentis ovato-lanceolatis acuminatis grossè serratis, petiolis basi dilatatis connatis. \S in Napauliâ. Wallich.
 19. *C. PERUVIANA* (DC. syst. 1. p. 141*) caule scandente, foliis bipinnati-sectis villosis, segmentis basi ovatis, apice 3-fidis incisissimis, floribus subpaniculatis polygamis, pedunculis folio longioribus bracteosis. \S in Peruviâ. Flores *C. Virginianæ*, majores. (v. s. in h. Mus. Paris.)
 20. *C. CORDATA* (Pursh! fl. bor. amer. 2. p. 384.) caule scandente, foliis pinnati-sectis, segmentis cordatis acuminatis lobatis inciso-dentatis, pedunculis elongatis 00-floris, floribus dioicis racemoso-paniculatis. \S in Virginie montibus excelsis. Sims. bot. mag. t. 1816. Habitu *C. Vitalbæ* æmulans. DC. syst. 1. p. 141. (v. s. in h. Lamb.)
 21. *C. CARIPENSIS* (H. B. et K. nov. gen. 5. p. 36*) caule scandente, foliis pinnati-sectis, segmentis ovatis acuminatis 5-nerviis integerrimis glabris, pedicellis bracteisque pubescentibus, floribus paniculatis dioicis. \S in umbrosis propè Caripe Cumanensium alt. 400 hex. *C. Caracasana* DC. syst. 1. p. 141.* Differt ab omnibus ferè speciebus dioicis ob folia pinnatim nec ternatim secta. (v. s. \S in h. H. et B.)
- \S . 2. *Floribus paniculatis, foliis ternatim biternatimve sectis.*
22. *C. VIRGINIANA* (Lin! amœn. 4. p. 275*. spec. 766.) floribus paniculatis dioicis, foliis ternatim sectis, segmentis cordatis acutis grossè dentatis lobatisve. \S in Americâ septentrionali. Albin. acad. ann. 1. p. 79. t. 7. DC. syst. 1. p. 142. Habitus *C. Vitalbæ*? sed foliorum segmenta constanter ternata. (v. v. et s.)
 - β. bracteata*, foliorum segmentis ovato-lanceolatis integris. *C. bracteata* Mœnch. suppl. 103*.
 23. *C. CATESBYANA*. (Pursh! fl. bor. amer. 2. p. 736.) floribus paniculatis subdioicis, foliis biternatim sectis, segmentis subcordatis 3-lobis. \S in Carolinâ merid. — Similis *C. Virginianæ*. DC. syst. 1. p. 142. (v. s. in h. Lamb.)
 24. *C. BRASILIANA* (DC. syst. 1. p. 143.*) floribus paniculatis dioicis, foliis 3-lobis, aut ternatim pinnatimve sectis, segmentis petiolulatis ovali-lanceolatis integris aut vix dentatis. \S in Brasiliâ. Deless. ic. sel. 1. t. 1. Species *C. Virginianæ* et dioicæ affinis. (v. s. in h. Deless.)
 25. *C. DIOICA* (Lin. amœn. 5. p. 398. spec. 765.) floribus paniculatis dioicis, foliis ternatim sectis, segmentis subglabris ovatis subcordatis acuminatis triplinerviis integerrimis, pericarpis ovalibus, pedicellis pubescentibus. \S in Jamaicâ. Sloan. hist. 1. p. 199. t. 128. f. 1. Flores abortu subdioici DC. syst. 1. p. 143.* (v. s. in h. Deless. et Banks.)
 26. *C. GLABRA* (DC. syst. 1. p. 143) floribus paniculatis dioicis, foliis ternatim sectis, segmentis glaberrimis ovatis acuminatis triplinerviis integerrimis, pericarpis ovali-oblongis, pedicellis glabris. \S in Sancto-Domingo. Affinis *C. dioicæ* et forsân ab eâ non meritò segregata. (v. s.)

27. *C. AMERICANA*. (Mill. dict. n. 14.) floribus paniculatis dioicis, foliis ternatim pinnatimve sectis, segmentis glabris ovato-cordatis 5-nerviis integerrimis acuminatis, pericarpis lanceolatis. ♀ in Americâ equatoriali. *C. Guadelupæ* Pers. ench. 2. p. 99. *C. Americana*. DC. syst. 1. p. 144.* (v. s.)
28. *C. LOURIERIANA*. (DC. syst. 1. p. 144.)* floribus paniculatis dioicis, foliis ternatim sectis, segmentis ovatis acutis integerrimis subquinqnerviis, pericarpis 80 obtusè trigonis. ♀ in Cochinchinâ. *C. dioica* Lour. fl. coch. 1. p. 421.* A Clem. dioicâ diversa videtur et ob patriam diversissimam et ob pericarpia duplò triplòve numerosiora. †
29. *C. SERICEA* (H. B. et K. nov. gen. 5. p. 37.)* floribus paniculatis dioicis, foliis ternatim pinnatimve sectis, segmentis ovatis 5-nerviis grossè 3-5-dentatis subtus sericeo-villosis, pedunculis folio brevioribus multibracteatis ♀ in Americâ merid. DC. syst. 1. p. 144. (v. s. in h. H. et B.)
- α. striatula*, caule striis 10-12. Mascula ad Santa-Fez.
- β. tereticaulis*, caule tereti non striato. Pericarpia 20-30 pubescentia.
- γ. costata*, caule costis vix prominulis. Ad ripas fluminis Rio-major. — fl. et fr. ign.
30. *C. HOLOSERICA* (Pursh! fl. bor. amer. 2. p. 384.) floribus paniculato-corymbosis dioicis, foliis ternatim utrinque pubescentibus, segmentis oblongo-lanceolatis integris. ♀ in Carolinâ. DC. syst. 1. p. 145. Tota holosericeo-pubescentia; sepala linearia staminibus longiora. (v. s. in h. Lamb.)
31. *C. BONARIENSIS* (Juss! herb. et DC. syst. 1. p. 145.) floribus paniculatis polygamis, foliis ternatim sectis subglabris, segmentis ovato-lanceolatis integerrimis, pedunculis petiolo brevioribus ramisque pubescentibus. ♀ in Circâ Buenos-Ayres. — Ovaria villis longis sericeis onusta. (v. s. in h. Juss.)
32. *C. GLYCINOIDES* (DC. syst. 1. p. 145.) floribus paniculatis dioicis, foliis ternatim sectis glabris, segmentis ovato-lanceolatis acuminatis integerrimis, basi triuerviis, pedunculis folio dimidio brevioribus, ramis glabris. ♀ in Novâ-Hollandiâ. Glycinen trifoliolatam habitu æmulans. (v. s. in h. Banks.)
33. *C. HEXASEPALA* (DC. syst. 1. p. 146.)* floribus paniculatis dioicis hexasepalis, foliis ternatim sectis, segmentis ovato-cordatis latè crenatis reticulatum venosis glabris coriaceis. ♀ in Novâ-Zelandiâ. *C. Forsteri* Gmel. syst. 873. Ic. ined. in bibl. Banks t. 170: *C. hexapetala*. Lin. f. suppl. 271. Bracteæ oblongo-lineares, flores patentes. (v. s. in h. Mus. Par. et Banks.)
34. *C. CORIACRA* (DC. syst. 1. p. 146.)* floribus paniculatis dioicis 4-sepalis, foliis ternatim sectis, segmentis cordatis grossè crenato-incisis glabris coriaceis, antheris ovato-oblongis. ♀ in Novâ-Hollandiâ. Valdè affinis *C. hexasepalæ*.
- α. obtusa*, segmentis cordatis obtusis (♂ in h. Mus. Par.)
- β. acuta*, segmentis ovato-acuminatis (♀ in h. Lamb.)
35. *C. STENOSEPALA* (DC. syst. 1. p. 147.)* floribus paniculatis dioicis 4-sepalis, foliis ternatim sectis, segmentis cordatis acuminatis 3-nerviis integris aut subdentatis, antheris ovato-oblongis. ♀ in Novâ-Hollandiâ. *C. stenopetala* Brown! ined. Bracteæ oblongæ integerrimæ, utrinque acutæ. (v. s.)
36. *C. MICROPHYLLA* (DC. syst. 1. p. 147.) floribus paniculatis dioicis, foliis ternatim sectis, segmentis oblongis sub-uni-nerviis integerrimis glabris coriaceis. ♀ in Insulis sterilibus circâ Novam-Hollandiam. Affinis *C. stenosepalæ*. (v. s. in h. Mus. Par.)
37. *C. ARISTATA* (Brown! prodr. 2. ined.) floribus paniculatis dioicis 4-sepalis, foliis ternatim sectis, segmentis ovato-subcordatis acutis grossè acutèque dentatis 3-nerviis, antheris apice aristatis. ♀ in Novâ-Hollandiâ. Ker. in bot. reg. t. 238.* DC. syst. 1. p. 147. (v. s. in h. Mus. Par. et Banks.)
38. *C. INDIVISA* (Willd. spec. 2. p. 1291.) floribus paniculatis, foliis ternatim sectis, segmentis ovatis integerrimis mucronatis coriaceis glabris. ♀ in Novâ-Zelandiâ. *C. integrifolia* Forst! prod. 42. n. 23r. non Lin. DC. syst. 1. p. 148.* (v. s. sine fl. et fr. in h. Mus. Par. Banks et Lamb.)

39. *C. MEDYSAEIFOLIA* DC. syst. 1. p. 148.* floribus paniculatis, foliis ternatim sectis, segmentis ovato lanceolatis acuminatis subintegris glabris imâ basi 5-nerviis. $\frac{3}{2}$ in rupibus Indîæ orient. Ker. bot. reg. t. 599. Ovaria villosissima in caudam barbatam brevem desinentia. (v. s. in h. Banks.)
40. *C. ACUMINATA* (DC. syst. 1. p. 148.*) floribus subpaniculatis, foliis ternatim sectis, segmentis ovato-lanceolatis acuminatissimis medio subserratis 3-5-nerviis glabris. $\frac{3}{2}$ in Napauliâ circa Narain-Hetty. Ovaria oo in stylos barbato-plumosos desinentia. (v. s. in h. Lamb.)
41. *C. APIFOLIA* (DC. syst. 1. p. 149.*) floribus paniculatis hexagynis, foliis ternatim sectis, segmentis ovatis acuminatis inciso-dentatis subtus nervosis. $\frac{3}{2}$ in Japoniâ. flores ignoti. Pericarpia 5-7 stellatim patentia. (v. s. in h. Lamb.)
42. *C. TRIFOLIATA* (Thunb. act. soc. Lin. 2. p. 337.) floribus paniculatis, foliis ternatim decompositis glabris, segmentis ovatis obtusis repando-dentatis. $\frac{3}{2}$ in Japoniâ. DC. syst. 1. p. 149. folia patentissima; petioli sesqui-polliares; segmenta petiolulata ovata obtusa repando-dentata subtus pallida $\frac{3}{2}$.
43. *C. BITERNATA* (DC. syst. 1. p. 149.*) floribus paniculatis hermaphroditis hexagynis, foliis glabris biternatim sectis, segmentis subcordatis acutis. $\frac{3}{2}$ in Insulâ Timor, Japoniâ et Cochinchinâ. *C. Virginica* Thunb. fl. jap. 240. non Lin. flores albi, hermaphroditi. (v. s. in h. Mus. Par.)
44. *C. TRITERNATA* (DC. syst. 1. p. 150.) floribus., foliis 2-3-ternatim sectis subglabris, segmentis ovali-cuneatis trinerviis acutè trifidis. $\frac{3}{2}$ patr. ign. — *Atragene triternata*. Desf. hort. par. Caules teretes, rami striis sex sulcati glabri, nascentes tenuissimè velutini. (v. v. in h. par. sub dio.) $\frac{3}{2}$
- §. 3. *Pedunculis 3-fidis 3-floris seu ternis 1-floris, foliis ternatim (rarius pinnatim) sectis.*
45. *C. DOMINICA* (Lam! dict. 2. p. 45.) pedunculis 3-fidis 3-floris folio longioribus, foliis ternatim sectis, segmentis ovatis subcordatis acutis subtus pubescentibus subdentatis. $\frac{3}{2}$ in Sancto-Domingo, Martinicâ et Dominicâ. *Atragene polygama* Jacq. amer. ed. picta 2. t. 261. f. 244. — DC. syst. 1. p. 150.* Flores parvi extus ferrugineo-pubescentes. (v. s. in h. Lam.)
46. *C. BRACHYLATA* (Ker! in bot. reg. t. 497.*) pedunculis ternis 1-floris seu 3-fidis paniculatisve foliis longioribus, foliis ternatim pinnatimve sectis, segmentis ovatis grossè dentatis, alabastris globosis. $\frac{3}{2}$ ad prom. Bonæ-Spei. DC. syst. 1. p. 150. Stamina filamenta basi præsertim hispida (v. s. in h. Banks.)
47. *C. GRANDIFLORA* (DC. syst. 1. p. 151.*) pedunculis 1-3-floris folio brevioribus, foliis pinnati-sectis glabris, segmentis ovato-lanceolatis acuminatis subtus reticulatim nervosis subserratis. $\frac{3}{2}$ in Sierrâ-Leonâ. Flores totius generis maximi. (v. s. in h. Banks et Smith.)
48. *C. LECHENAUTIANA* (DC. syst. 1. p. 151.*) pedunculis 3-fidis 3-floris folio brevioribus, foliis ternatim sectis adpressè sericeis, segmentis ovato-lanceolatis 5-nerviis exsertè serratis. $\frac{3}{2}$ in Javâ. Calycia sepala oblonga subacuta intus extusque velutina. (v. s. in h. Mus. Par.)
49. *C. NORONHIANA* (DC. syst. 1. p. 151.*) pedunculis 1-3-floris folio brevioribus, foliis ternatim sectis pilosis, segmentis ovato-lanceolatis 3-nerviis exsertè serratis. $\frac{3}{2}$ in Javâ. Trigula trifoliata. Noronha ic. in ed. in bibl. Juss. Antheræ villosæ; styli albi; fructus ignotus. (Descript. ex ic.) $\frac{3}{2}$
50. *C. MAURITIANA* (Lam! dict. 2. p. 42.*) pedunculis 3-fidis, foliis ternatim sectis glabris, segmentis ovatis acuminatis grossè serratis. $\frac{3}{2}$ in insulis Mauritiî et Madagascar. Deless! ic. sel. 1. t. 2. *C. triflora* Vahl! symb. 3. p. 74. *C. triflora* et *Sonneratii* Pers. ench. 2. p. 99. — DC. syst. 1. p. 152. (v. s.)
51. *C. HAVANENSIS* (H. B. et K! nov. gen. et spec. amer. 5. p. 38.*) pedunculis 1-3-floris, foliis ternatim sectis subtus sericeo-pubescentibus, segmentis ovato-lanceolatis grossè serratis. $\frac{3}{2}$ circâ Havanam. Pericarpia pubescentia

desinentia in caudam barbatam. DC. syst. 1. p. 152.* (v. s. sine fl. in h. H. et B.)

52. C. JAVANA (DC. syst. 1. p. 152.) pedicellis 1-3-floris, foliis ternatim biter-natimve sectis pubescentibus, segmentis ovato-lanceolatis acuminatis integris 3-fidisve subdentatis. ♂ in Javā. Sepala 4 oblonga patentia margine subvelutina, staminibus paululum longiora. (v. s. in h. Deless.)

53. C. DAHURICA (Perslensch. 2. p. 99.) pedunculis ternis 1-floris, foliis ternatim sectis glabris, segmentis ovatis basi obliquis integerrimis mucronatis, floribus hermaphroditis natantibus. ♂ in Dahuriā. Ex flore species videtur viti-cellæ affinis. DC. syst. 1. p. 153. (v. s. in h. Juss.)

54. C. PLUCKNETII (DC. syst. 1. p. 153.) pedunculis 1-floris, foliis ternatim sectis glabris, segmentis ellipticis obovatisve obtusis integerrimis mucronatis, floribus dioicis erectis. ♂ in America? Pluck. alm. 109. flores dioici; alabastra subglobosa. (v. s. in h. Banks et Catesby.)

§. 4. *Pedunculis solitariis, 1-floris, foliis pinnatim (rarius ternatim) sectis.*

55. C. ANGUSTIFOLIA (Jacq. enum. 310. coll. 1. p. 137* ic. rar. 1. t. 104.) pedun-culis unifloris, sepalis 6-8 obtusissimis, foliis pinnatim sectis, segmentis lanceolato-linearibus acuminatis integris 3-lobisve, caulibus erectis. ♀ ad littora septentrionalia maris Adriatici, in collibus Dahuriæ, (nec Tauriæ Fisch.) C. lasiantha. Fisch. h. gorenk. ex litt. DC. syst. 1. p. 154.* (v. s.)

56. C. SCABIOSÆFOLIA (DC. syst. 1. p. 154.) pedunculis 1-floris, sepalis 4 ovali-oblongis, foliis pinnati-sectis, segmentis petiolulatis 3-partitis incisive pube-scentibus, lobis cuneiformibus, caule erecto velutino. ♀ in Indiā? Tota velu-tino pubescens. Stamina filamenta basi hispida. (v. s. in h. Mus. Par.)

57. C. VILLOSA (DC. syst. 1. p. 154.) pedunculis 1-floris, sepalis 4 ovali-oblongis, foliis villosis pinnati-sectis, segmentis sessilibus ovalibus apice grossè dentatis, terminali 3-fido, caule erecto villoso. ♀ in Indiā? Adeò similis C. scabiosæfoliæ ut vix ab eâ primo intuitu separanda. (v. s. in h. Mus. Par.)

58. C. LINEARILOBA (DC. syst. 1. p. 155.) pedunculis 1-floris, sepalis acutissi-mis, foliis pinnatim sectis glabris, segmentis integris aut 3-partitis, lobis linearibus. ♂ ♀ in Carolinā inferiore. Deless! ic. sel. 1. t. 3. Sepala acuta margine pubescentia, staminibus duplò fere longiora. (v. s.)

59. C. WALTERI (Parsh. fl. bor. amer. 2. p. 384.) pedunculis unifloris, sepalis ellipticis, foliis pinnatim sectis, segmentis divaricatis lineari-lanceolatis acu-tis integerrimis subius glaucis. ♀ in Carolinā. Folia cirrhosa, 3-juga; flores albi. DC. syst. 1. p. 155.

60. C. DIVERSIFOLIA (DC. syst. 1. p. 155.*) pedunculis 1-floris, sepalis acutissimis subpatentibus, foliis glabris, nunc integris, nunc 3-lobis, nunc 3-sectis, segmentis lanceolatis acutis, lateralibus sessilibus, mediò petiolato. ♂ habi-tat. Deless! ic. sel. 1. t. 4. Antheræ filamentaque villosa. (v. v. in hort. Parmensi.)

61. C. VIORNA (Lin! sp. 765.) pedunculis 1-floris, sepalis conniventibus crassis apice reflexis acuminatis, foliis glabris pinnatim sectis, segmentis integris 3-lobisve ovatis acutis, floralibus integris. ♂ ♀ in Americā boreali, Virginiā, Carolinā inferiore. Jacq. fil. eclog. 1. p. 50. t. 32. Flores violacei, pedicello apice inflexo natantes. DC. syst. 1. p. 156. (v. v. et s.)

62. C. CYLINDRICA (Sims. in bot. mag. t. 1160.) pedunculis 1-floris, sepalis acuminatis margine undulatis tenuibus, foliis glabris tenuibus pinnatim de-compositis, segmentis petiolatis ovatis oblongisque. ♂ ♀ in Pensylvaniā, Carolinā et Virginiā. C. crispa Lam! dict. 2. p. 44*. C. Viorna Andr. bot. rep. t. 71. non Lin. Affinis C. Viornæ reticulatæ et crispæ. DC. syst. 1. p. 156.* (v. s. in h. Lam. Rich.)

63. C. RETICULATA (Walt. fl. car. 156.) pedunculis 1-floris, sepalis conniventi-

- bus, foliis coriaceis reticulatim nervosis glabris pinnatim sectis, segmentis petiolatis trilobis aut integris ovatis. ♂ in Georgia, Carolinā inferiore. *C. rosea* Abbot. insect. am. ic. t. 101. ex Roem. coll. 1. p. 401. Flores ferè Viornæ; DC. syst. 1. p. 157.* (v. s.)
64. *C. JAPONICA*. (Thunb. fl. jap. 240.*) pedunculis 1-floris, sepalis conniventibus, foliis villosis ternatim sectis, segmentis ellipticis acuminatis apice serratis. ♂ in Japoniā. Pedunculi uniflori longitudine foliorum. DC. syst. 1. p. 157*.
65. *C. COMOSA* (DC. syst. 1. p. 157.*) pedunculis unifloris, foliis ternatim sectis velutinis, segmentis ovali-oblongis acuminatis 3-nerviis integerrimis. ♂ in Indiā orientali. Flores ignoti; pericarpia ovata, subcompressa, pubescentia, in caudam ferè bipollicarem barbato-plumosam desin. (v. s. in h. Mus. Par.)
66. *C. TRILOBA* (Roth. nov. pl. spec. 251*) pedunculis unifloris subcorymbosis, caule petiolisque tomentosis, foliis pubescentibus, inferioribus ovato acuminatis, mediis 3-lobis, superioribus ternatim sectis, sepalis oblongo-lanceolatis mucronatis extus sericeis. ♂ in Indiā orientali. Affinitates quasdam præbet cum *Cl. orientali* et *Vitalbæ*. †
67. *C. CLITORIOIDES*. (DC. syst. 1. p. 158.*) pedunculis 1-floris axillaribus tortilibus, foliis glabris pinnatim sectis, segmentis oblongis acutis integerrimis. ♂ in Novā-Hollandiā, Glabra, Clitoriam Kennediamve habitu omnino æmulans, (v. s. in h. Banks.)
- §. 5. *Pedunculis solitariis 1-floris, foliis indivisis, caulibus herbaceis erectis.*
68. *C. INTEGRIFOLIA* (Lin! spec. 767.) pedunculis unifloris, floribus nutantibus, foliis integris ovato-lanceolatis glabris. ♀ in Hungariā, Carnioliā, Tatariā, Sibirīā, Pyrenæis. Jacq. fl. aust. t. 363. Tratt. tab. 4. t. 179. Clus. hist. 1. p. 123. fig. 2. Sepala cœrulea subcoriacea. DC. syst. 1. p. 158. (v. v. et s.)
- β *angustifolia*. (Fisch! in litt.) foliis angustissimis acuminatis 3-5-nerviis, sepalis subinæqualibus, staminibus sublongioribus.
- γ. *elongata*. floribus erectiusculis, foliis acuminatis pilosiusculis. *C. elongata* Tratt. tab. 4. t. 178. in Sibirīā.
69. *C. OCHROLEUCA* (Ait. h. kew. ed. 1. v. 2. p. 260.) pedunculis 1-floris, floribus suberectis, foliis integris ovatis, junioribus sericeis. ♀ Secus rivos in montibus Carolinæ, Virginie, Pennsylvania. Sims. Bot. mag. t. 1175. *C. sericea* Michx! fl. bor. amer. 1. p. 319. Affinis *C. integrifoliæ*, sed differt foliis ovalibus ovatisve obtusis. DC. syst. 1. p. 159.* (v. s.)
70. *C. OVATA*. (Pursh. fl. bor. amer. 2. p. 736.) pedunculis unifloris, floribus erectis, foliis ovatis acutis glabris utrinque reticulato-venosis, infimis subcordatis. ♂ in Carolinā merid. — Pluk. mant. t. 379. f. 4? Similis *C. integrifoliæ*. Pericarpiorum caudæ plumosæ longissimæ. DC. syst. 1. p. 159.
71. *C. GENTIANOIDES* (DC. syst. 1. p. 159.*) pedunculis 1-floris, floribus erectis dioicis, foliis ovatis integris dentatisve glabris 3-nerviis. ♀ in Novā-Hollandiæ insulā Mariā, etc. Deless! ic. sel. 1. t. 5. Sepala 4 oblonga, tenera, callo minimo terminata, (v. s. ♂ in h. Mus. Par. ♀ in h. Lamb.)

Sect. II. VITICELLA. DC. syst. 1. p. 160.

Involucrum o. petala o. caudæ breves ebarbatæ. — Folia ternatim secta seu decomposita; caules scandentes.

72. *C. FLORIDA* (Thunb. fl. jap. 240.) pedunculis 1-floris folio longioribus, foliis ternatim decompositis, segmentis ovatis acutis integerrimis, sepalis ovali-lanceolatis acuminatissimis. ♂ in Japoniā. Sims. bot. mag. t. 834. *Atragene Indica* Desf. tab. par. ed. 1. p. 123. Fructus ignotus. Sæpius flore pleno in hortis herbariisque occurrit. DC. syst. 1. p. 160. (v. v. et s.)

73. *C. VITICELLA* (Lin! spec. 762.) pedunculis 1-floris folio longioribus, foliis integris aut ternatim decompositis, lobis segmentisve integerrimis, sepalis patentibus obovatis. $\frac{5}{2}$ ad sepes et dumeta Europæ calidioris, etc. *C. altera* Clns. hist. 1. p. 122. f. 1. Variat caule prostrato aut scandente, etc. DC. syst. 1. p. 160. (v. s.)

* *flore pleno*. *C. pulchella* Pers. ench. 2. p. 99. (v. v.)

℞. *tenuifolia* foliorum segmentis oblongo-lanceolatis. *C. tenuifolia lusitana*, etc. Tourne! inst. 299.

γ. *baccata*. fructibus subbaccatis. *C. baccata*. Pers! ench. 2. p. 99.

74. *C. CAMPANIFLORA* (Brot. fl. lus. 3. p. 359. *) pedunculis 1-floris folio longioribus, foliis biternatim decompositis, segmentis integris 3-lobisve, sepalis semipatentibus apice dilatatis undulatis. $\frac{5}{2}$ $\frac{4}{2}$ in Lusitaniâ ad sepes. Media inter *C. Viticellam* et *crispam*. DC. syst. 1. p. 161.

75. *C. CRISPA* (Lin! spec. 765.) pedunculis 1-floris folio brevioribus. foliis integris 3-lobis 3-sectisve acutissimis, sepalis basi conniventibus apice patentibus reflexis. $\frac{5}{2}$ in Americâ boreali. Sims. bot. mag. t. 1892. * *C. flore crispo*. Dill. elth. 1. t. 73. f. 84. * *Pericarpia* in caudam subulatam crassam brevem desinentia. DC. syst. 1. p. 162. (v. s.)

76. *C. PARVIFLORA* (DC. mem. soc. h. n. Gen. 1. p. 433.) pedunculis 1-floris, sepalis ellipticis apice subdilatatis, foliis pinnatim sectis, petiolis cirrhoriformibus, segmentis petiolulatis 3-partitis integrisve, lobis ovatis mucronulatis integris. $\frac{5}{2}$ hab. *Carpella* glabra ecaudata, flores albi. (v. v.)

Sect. III. CHEIROPSIS. DC. syst. 1. p. 162.

Involucrum calyciforme ex duabus bracteis coalitis sub flore ad apicem pedunculi situm; petala 0. caudæ barbatae. — Frutices sarmentosi.

77. *C. CIRRHOSA* (Lin. spec. 766.) pedunculis 1-floris involucreatis, foliis ovato-subcordatis dentatis fasciculatis. $\frac{5}{2}$ ad sepes et dumeta in Ætica, Majorcâ, Siciliâ, Calabriâ. Clns. hist. 123. f. 1. Smith. fl. Græc. t. 517. *Atragene cirrhosa* Pers. ench. 2. p. 98. Frutex elegans sempervirens super arbores sepesque scandens. DC. syst. 1. p. 162. (v. v.)

℞. *pedicellata*, flore minore pallido intrâ involucrum pedicellato. *C. cirrhosa* Sims. bot. mag. t. 1070. *C. Balearica* Pers. ench. 2. p. 99. (non Rich.) Insulâ Majorcâ. (v. s.)

78. *C. SEMITRILoba* (Lag. cat. h. madr. 17.) pedunculis 1-floris sub flore involucreatis, foliis cordatis serratis plerisque trifidis. $\frac{5}{2}$ in Hispaniâ merid. Similis *C. cirrhosæ*; sepala ovata extus tomentosa. DC. syst. 1. p. 163.

79. *C. BALEARICA* (Rich! in Journ. phys. 1779. févr. 127. ic*. non Pers.) pedunculis unifloris sub flore involucreatis, foliis ternatim sectis, segmentis petiolatis 3-lobis dentato-incisis. $\frac{5}{2}$ in insulâ Minorcâ. *C. calycina* Ait! h. kew. ed. 1. p. 259. Schrank. hort. mon. t. 15. *Atragene Balearica* Pers. ench. 2. p. 98. Involucrum campanulatum subhilobum obtusum florem arcuè cingens. DC. syst. 1. p. 163. (v. v. et s.)

80. *C. NAPAULENSIS* (DC. syst. 1. p. 164.) pedunculis 1-floris sub flore involucreatis, foliis 3-sectis, segmentis oblongis acuminatis basi subdentatis, lateralibus subsessilibus 1-nerviis, terminali 3-nervio. $\frac{5}{2}$ in Napauliâ. *C. montana* var. ? Buchan! ined. in h. Lamb. *Pericarpia* ∞, compressa, ovato-acuminata. (v. s. in h. Lamb.)

81. *C. MONTANA* (Buchan! ined. in h. Lamb.) pedunculis 1-floris, involucreo nullo, foliis ternati-sectis, segmentis oblongis acuminatis subtrifidis grosse serratis. $\frac{5}{2}$ in Napauliæ montanis; habitus *Atragenes* sed petala 0; forsâ sectionem propriam conficiens. DC. syst. 1. p. 164. (v. s. in h. Lamb.)

Sect. IV. ATRAGENE. — *Atragenes* spec. Lin. gen. 615.

Involucrum 0. Sepala 4, æstivatione subinduplicata. Petala ∞. Cariopsidum caudæ barbatae. *Cotyledones* in semine approximatae. Caules sarmentosi.

82. *C. ALPINA* (Mill. dict. n. 9.) pedunculis unifloris folio longioribus, foliis biternatim sectis, segmentis ovato-lanceolatis acuminatis serratis, petalis subspatulatis obtusis. $\frac{1}{2}$ in dumetis asperis montium præsertim calcareorum Europæ à 400-1000 hexap. alt. Atragene Alpina Lin! spec. 764. Jacq. fl. aust. t. 241.* Flores cærulei, variantes albi nunquam flavescentes. DC. syst. 1. p. 165. (v. v.)
83. *C. SIBIRICA* (Mill. dict. n. 12) pedunculis 1-floris folio subæqualibus, foliis biternatim sectis, segmentis oblongo-lanceolatis acuminatis serratis, petalis apice emarginatis. $\frac{1}{2}$ in Sibiria. Atragene Sibirica Lin. spec. ed. 1. p. 343. Sims. bot. mag. t. 1951. Variat, ex Fisch., in uno eodemque caule floribus sordide et virescenti-ochroleucis et floribus rubore lilæino suffusis. DC. syst. 1. p. 165. (v. v. et s.)
84. *C. OCHOTENSIS* (Poir. suppl. 2. p. 298.) pedunculis unifloris, foliis biternatim sectis, segmentis oblongo-lanceolatis acuminatis serratis, petalis paucis linearibus. $\frac{1}{2}$ in Sibiria versus mare Ochotense. Atragene Ochotensis Pall. ross. 2. p. 69*. Petala o aut interdum stamina exteriora sub abortiva elongato-linearia. DC. syst. 1. p. 166. (v. s. in h. Lamb.)
85. *C. VERTICILLARIS* (DC. syst. 1. p. 166.) pedunculis 1-floris, foliis quaternatim verticillatis ternatisectis, segmentis petiolulatis cordatis lanceolatis integerrimis, petalis acutis. $\frac{1}{2}$ in America bor. Atragene Americana Sims. bot. mag. t. 887. Habitus *C. Alpinae*; flores intensè purpureo cærulei.
86. *C. OCCIDENTALIS* pedunculis..... foliis oppositis ternatim sectis, segmentis subintegerrimis lucidis, sepalis ovato-lanceolatis campanulato-patentibus. $\frac{1}{2}$ Hab..... Atragene occidentalis. Horn. h. hafn. 2. p. 520.

†. *Clematides non satis notæ.*

87. *C. TOURNEFORTII* (DC. syst. 1. p. 167.) *C. orientalis latifolia* semine brevissimis pappis donato. Tourn. cor. 20. Hab. in Oriente. (Deest in herb. Tourn.)
88. *C. P. POLYPETALA* (Poir. suppl. 2. p. 296.) in Japonia. Atragene Japonica Thunb. fl. jap. 259.* DC. syst. 1. p. 167. ex descriptione videtur *Anemone*.
89. *C. CANALICULATA* (Lag. cat. h. madr. 17.) in Murcia et Granatæ regnis, locis saxosis petrosisque. Suffruticosa scandens; folia decomposita, linearia, canaliculata; pedunculi axillares subtriflori. (Lag.) DC. syst. 1. p. 167.
90. *C. SMILACIFOLIA* (Wall. in Asiat. res. 13. 1819. p. 414. n° 2.) $\frac{1}{2}$ in Neapoli. Scandens, foliis simplicibus cordato-ovatis, racemis axillaribus paucifloris (Wall.).

II. *NARAVELIA*. DC. syst. 1. p. 167.

Petala 6-12, calyce longiora; carpella stipite crasso cavo instructa. — Habitus *Clematidum* sed folia segmenta tantum duo opposita connervia gerentia.

1. *ZEYLANICA*. (DC. syst. 1. p. 167.) $\frac{1}{2}$. Pericarpia in caudam longam harbatam apice producta. (v. s.)

a. *Hermanni*, foliis subtus velutino-tomentosis — in Ceylonæ nemorosis. Atragene Zeylanica Lin! amæn. 1. p. 405.

ß. *Roxburghii*, foliis subtus velutino-pubescentibus. Atrag. Zeylanica Roxb! corom. 2. p. 47. t. 188* in sepibus humidis Coromandelie.

Trib. II. *ANEMONEÆ*. DC. syst. 1. p. 168.

Calycis corollæque æstivatio imbricativa. Petala o aut plana. *Carpella* 1-sperma indehiscencia in caudas mucronesve sæpius

desinentia. Semen pendulum. Caules herbacei nunquam scandentes; folia nunc radicalia, nunc caulina alterna.

III. THALICTRUM. *Lin. gen. n. 697. DC. syst. 1. p. 168.*

Involucrum sub flore o. Petala o. Carpella sicca in caudas non desinentia, nunc stipitata, nunc longitudinaliter striato-sulcata. — Herbes perennes; caulis annuus; flores corymbosi, paniculati aut subracemosi, herbacei, albi vel flavi.

Sect. I. TRIPTERIUM. *DC. syst. 1. p. 168.*

Fructus tri-queter, angulis alatis, substipitatus.

1. *T. AQUILEGIFOLIUM* (Lin! spec. 770.) stipellis ovatis binis ad basim petioli ramificationum, panicula corymbosa, fructibus 3-quetris. \mathcal{U} in montanis subeylvaticis Europæ. Sims. bot. mag. t. 1818. — J. Bauh. hist. 3. p. 2. p. 487. f. 2. Pericarpia glabra, basi angustata ex thalamo longè pedicellata pendula obtusa. DC. syst. 1. p. 169.

β . *atro-purpureum* Murr. syst. veg. 513. Jacq. h. vind. 3. t. 81.

γ . *formosum*, staminibus intense purpureis superne dilatatis. Sims. bot. mag. t. 2025.*

2. *T. CRELIDONI*, stipellis nullis, foliorum axillis gemmiferis, floribus laxè paniculatis, fructibus stipitatis pedicello inflexo pendulis. — In Napauliâ. Flores majusculi; sepala videntur in sicco sub purpurascentia; pistilla 8; stamina 8-12 (v. s. in h. Mus. Par.)
3. *T. CONTORTUM* (Lin! spec. 770.) stipellis nullis, floribus laxè corymboso-racemosis, fructibus 3-quetris pendulis. \mathcal{U} in Sibiria. Gmel. fl. sib. 4. p. 198. simile *T. aquilegifolii* sed humilius. Ovaria immatura *T. aquilegifolii*. DC. syst. 1. p. 170. (v. s. in h. Lin.)

Sect. II. PHYSOCARPUM. *DC. syst. 1. p. 171.*

Fructus inflatus vesicarius stipitatus. — Flores dioici, aut polygami. Folia 2-3-ternatim secta.

4. *T. CLAVATUM* (DC. syst. 1. p. 171.) floribus monoïcis, filamentis clavatis, pericarpis compressis stylo brevissimo vix rostellatis, foliis glabris exstipellatis. \mathcal{U} in America boreali. Deless! ic. sel. 1. t. 6. Folia petiolo ramoso biter-natim secta. (v. s. in h. Mus. Par.)
5. *T. PODOCARPUM* (H. B. et Kl nov. gen. et sp. amer. 5. p. 38.*) floribus polygamiis, filamentis filiformibus, antheris longis apice subaristatis, pericarpis ovali-oblongis lævibus stylo persistente longè aristatis, foliis glabris exstipellatis. \mathcal{U} in Novâ-Granadâ. Folia petiolo ramoso quadripinnatim seu quadri-ternatim secta. DC. syst. 1. p. 171. (v. s. in h. H. B.)
6. *T. LONGISTYLM* (DC. syst. 1. p. 171.) floribus polygamiis, filamentis filiformibus, antheris longè aristatis, pericarpis ovali-oblongis stylo longè aristatis, petioli ramis stipellatis, segmentis ad nervos subtus puberulis. \mathcal{U} in America merid. Deless! ic. sel. 1. t. 7. Folia petiolo ramoso 3-4 ternatim secta, segmentis ovato-orbiculatis. (v. s. in h. Mus. Par.)
7. *T. PELTATUM*, floribus polygamiis, filamentis filiformibus, antheris longè aristatis, carpellis sessilibus stylo longè aristatis, segmentis foliorum inferiorum peltatis. \mathcal{U} in Mexico. (v. s.)
8. *T. RUTIDOCARPUM* (DC. syst. 1. p. 172.) floribus polygamiis, filamentis filiformibus, pericarpis ovato-subcompressis rugis elevatis ramosis notatis, foliis exstipellatis, segmentis ad nervos subtus puberulis. \mathcal{U} in America merid. Affine *T. podocarpum* et *longistylum*. (v. s. in h. Mus. Par.)

Sect. III. EUTHALICTRUM. DC. syst. 1. p. 172.

Fructus ovato-oblongus striatus sessilis. — Folia varia, fere semper exstipellata.

§. 1. Heterogama. *Floribus dioicis polygamisve.*

9. T. CORYNELLUM (DC. syst. 1. p. 172.) floribus dioicis, filamentis apice clavatis, pericarpis oblongis sessilibus striatis, foliorum segmentis obtusiusculè 3-lobis, subtus glaucis pilosque parcos in nervis gerentibus. \mathcal{U} in Amer. boreali. T. Cornuti. Lin. spec. 768? — Moris. hist. 3. p. 325. s. g. t. 20. f. 15? (v. s. in h. Mus. Par.)
10. T. REVOLUTUM (DC. syst. 1. p. 173.) floribus dioicis polygamisve, filamentis filiformibus, foliorum segmentis ovatis subtrilobis, margine subtus revolutis, subtus velutino-pubescentibus. \mathcal{U} in America boreali. T. pubescens Pursh. fl. bor. amer. 2. p. 288 — Hern. mex. 235. ic? (v. s.)
- β . *subglabrum* à priori differt tantum foliis magis reticulato-venosis subtus vix ac ne vix pubescentibus, et panicula pauciflora. (v. s.)
11. T. DIOICUM (Lin. spec. 768.) floribus dioicis, filamentis filiformibus, foliorum segmentis subrotundis cordatis obtusè lobatis glabris, pedunculis axillaribus folio brevioribus. \mathcal{U} in Canadâ. Deless. ic. sel. 1. t. 8. — T. lævigatum. Michx! fl. bor. amer. 1. p. 322. Pericarpia ovata mutica. DC. syst. 1. p. 173. (v. s.)
12. T. CAROLINIANUM (Bosc! ined. non Walt.) floribus dioicis, filamentis filiformibus, foliorum segmentis ovatis 3-5 dentatis glabris subtus glaucis, pedunculis axillaribus folio longioribus. \mathcal{U} in Pensylvaniâ et Carolinâ. T. rugosum Pursh. fl. bor. amer. 2. p. 383. non Ait. Affine Thal. dioico, sed differt foliorum segmentis ovalibus minùs rotundis cordatisve. DC. syst. 1. p. 174. (v. s.)
- β . *subpubescens*. foliis subtus in nervis tenuissimè pubescentibus in New-Jersey.
13. T. PURPURASCENS (Lin. spec. 769.) floribus dioicis monoicisve, filamentis filiformibus coloratis, foliorum segmentis suborbiculatis grossè dentatis glabris subtus glaucis, paniculâ contractâ subaphyllâ. \mathcal{U} in Canadâ, Pensylvaniâ, et Virginîâ — Moris. oxon. 3. p. 324. Panicula femina erecta; ovaria

● 7-8, stylo longo acuto. DC. syst. 1. p. 174. (v. s. in h. Mus. Par.)

β . *monoicum*, panicula pauciflora basi feminea apice mascula.

§. 2. Genuina. *Floribus hermaphroditis, foliis ∞ -sectis, radice fibrosa.*

14. T. FOLIOLOSUM (DC. syst. 1. p. 175.) caule tereti lævi ramoso, floribus paniculato-corymbosis erectis, petiolis quadrinerviatis decompositis, segmentis glabris orbiculatis obtusi 3-dentatis. \mathcal{U} in Napauliâ. T. Dalingo Buchan! ined. in h. Lamb. Species forsan polygama et ideo ad priorem divisionem cui habitu accedit transferenda. (v. s. in h. Lamb.)
15. T. SANICULÆFORME caule subtereti ramoso, floribus paucis paniculatis, pedunculis rigidis divaricatis, sepalis reflexis, carpellis striatis stylo apice inflexo longe acuminatis. \mathcal{U} in Napauliâ. (v. s.)
16. T. PETALOIDEUM (Lin. spec. 770.*) caule tereti subnudo, floribus corymbosis, staminum filamentis apice dilatatis, foliorum segmentis glabris ovatis obtusis integris 3-lobisve. \mathcal{U} in montibus Dahuriæ et in monte Odon-Tchelen ad Sinenses limites. Deless! ic. sel. 1. t. 9. Antheræ lineares flavæ; pericarpia 6-8 sessilia oblonga striata. DC. syst. 1. p. 175. (v. s.)

β . *stamineum* Lin. f. suppl. 271. (ex Retz et Willd.)

17. T. ALPINUM (Lin! spec. 767.) caule simplicissimo subnudo digitali, racemo

- simplici terminali, floribus ntantibus, fol. segmentis glabris. \mathcal{L} in pascuis editioribus montium Kamschatkæ, Laponiæ et Pyrenæorum — Moris. oxon. 3. p. 325. s. 9. t. 20. f. 14. — Species humillima et digiti altitudinem vix adæquans. DC. syst. 1. p. 175. (v. v.)
18. *T. FORTIUM* (Lin. spec. 768?) caule simplici, basi nudo, medio folioso, apice paniculato, foliis pubescenti-viscidis, segmentis dentibus obtusis. \mathcal{L} in Alpium vallibus, Monte-Cenisio, Valle Binn. Helvetiæ. Lam! illustr. t. 497. f. 3. *T. styloideum* Lin. f! suppl. 271. ex h. Lin! caulis pubescenti-viscidus, basi vaginis tectus. DC. syst. 1. p. 176. (v. s.)
19. *T. PUBESCENS* (Schl! pl. exsic.) caule simplici foliis sparsis obauto apice paniculato, foliis subpubescenti-viscidulis. \mathcal{L} in lapidosis calcareis Occitanis circa Monspelium, Avenionem, et in Vallesia inferiori. *T. Cornuti* Poir! dict. 5. p. 319.* excl. syn. et patriâ. Videtur species à Linnæo pro *T. fœtido* habita. DC. syst. 1. p. 176. (v. v.)
20. *T. ACUTILOBUM* (DC. syst. 1. p. 177.) caule simplici basi nudo, medio folioso, apice paniculato, foliis pubescenti-viscidulis, segmentorum lobis acutis. \mathcal{L} in Sibiria — Deless! ic. sel. 1. t. 10. *T. fœtido* simillimum, sed segmentorum lobi paulò profundiores et omnes acuti nec obtusi. (v. s. in h. Deless.)
21. *T. SQUARROSUM* (Steph. in Willd. spec. 2. p. 1299.*) caule tereti, floribus paniculatis cernuis, petiolis amplexicaulibus alatis. \mathcal{L} in Sibiria. *T. Sibirico* et minori simile, sed diversum petioli structura. DC. syst. 1. p. 177.
22. *T. SIBIRICUM* (Gaertn. fruct. 1. p. 355. t. 74.) caule subtereti, floribus paniculatis cernuis, foliorum segmentis glabris ovato-cuneatis trifidis, lobis acutis integris aut argutè incisis. \mathcal{L} in Armenia, Sibiria. *T. Sibiricum* Lin. spec. 769? *T. acuminatum* Spreng! pug. n. 130* Planta spontanea est duplò minor quam *T. minus*. DC. syst. 1. p. 177. (v. s.)
23. *T. MINUS* (Lin! spec. 769.) caule tereti polline glauco, floribus laxè paniculatis cernuis 4-sepalis, foliorum segmentis glabris sub-rotundis apice dentatis subtus glaucis, carpellis utrinque acutis striato-sulcatis. \mathcal{L} in Europæ ferè totius pascuis submontosis et in Sibiria. Jacq. fl. aust. 5. p. 9. t. 419. — J. Bauh. hist. 3. p. 487. f. 3. DC. syst. 1. p. 178. (v. v.)
24. *T. COLLINUM* (Wallr! sched. crit. 259.) caule basi folioso internodiis approximatis intercepto, foliis utrinque viridibus glaberrimis, radicalibus sessilibus confertis, segmentis è basi ovatâ subcordatave subrotundis obtusè trifidis, floribus paniculatis cernuis, carpellis ovatis æqualiter costatis concoloribus stigmate recto coronatis. \mathcal{L} in agro Halensi. *T. minus* Spreng. fl. hal. 156. excl. syn. an satis à sequente distinctum? (v. s.)
25. *T. SAXATILE* (Schl! pl. exsic.) caule tereti polline destituto, floribus paniculatis erectis foliorum segmentis glabris subrotundis apice dentatis, subtus glaucis, carpellis utrinque acutis. \mathcal{L} in Europæ mediæ collibus sylvaticis montiomque vallibus. *T. minus* Poll. pal. n. 522*. *T. minori* valde affine. DC. syst. 1. p. 178. (v. v.)
26. *T. CALABRICUM* (Spreng. pug. 1. p. 37. n. 67*) caule tereti usculo polline destituto, florum panicula subracemosa contracta aphylla, foliorum segmentis subrotundis obtusè 3-lobis subtus subglaucis, carpellis profundè sulcatis basi attenuatis. \mathcal{L} in monte Leonis Calabriae et Sicilia. Icon Morisoni 1. s. 9. t. 20. f. 16. non malè plantam refert. (v. s. in h. Moric.)
27. *T. ELATUM* (Murr. syst. ed. 14. p. 513.) caule tereti usculo polline glauco destituto, floribus paniculatis erectis, foliorum segmentis glabris ovatis subcordatisve subtrifidis, carpellis basi obtusis. \mathcal{L} in Hungaria. Jacq. h. Vind. 3. t. 95.* Ambigit inter *T. saxatile* et majus. DC. syst. 1. p. 179. (v. v.)
- β . *T. ambiguum* Schl! pl. exsic. ad margines agrorum in valle Binn Helvetiæ. (v. s.)
28. *T. MAJUS* (Murr. syst. 513.) caule tereti polline glauco destituto, floribus

- laxè paniculatis cernuis, foliorum segmentis glabris subtus glaucis 3-fidis, lobalis ovatis submacronatis, carpellis basi obliquè rotundatis. \mathcal{U} in collibus siccis dumosis Helvetiæ, Palatinatûs, Angliæ, etc. Jacq. fl. aust. 5. t. 420. Engl. bot. t. 611. DC. syst. 1. p. 179. — Conf. Wallr. sched. cr. p. 263. (v. v.)
29. *T. NUTANS* (Desf! tabl. mus. ed. 1. p. 123.) caule tereti polline glauco destituto, floribus laxè paniculatis cernuis, foliorum segmentis subtus glaucis apice 3-5 dentibus acutè incisæ, carpellis basi obliquis obtusis. \mathcal{U} in Alpibus Sabandis. DC. syst. 1. p. 179. (v. v.)
30. *T. MEDIUM* (Murr. syst. 513.) caule tereti polline destituto, floribus laxè paniculatis erectiusculis, foliorum segmentis glabris oblongis cuneiformibus acutè 3-fidis, summis indivisis lanceolatis. \mathcal{U} in collibus Hungariæ, Tauriæ. Jacq. h. vind. 3. t. 96. DC. syst. 1. p. 179. (v. s.)
31. *T. TRIGYNUM* (Fisch. in litt.) caule sulcato erecto, floribus laxissimè paniculatis erectis 3-gynis, foliorum segmentis glabris subtus glaucis ovali-cuneatis acutè trifidis. \mathcal{U} in Dahuriâ. Carpella 1-3, oblonga, striata. (v. v.)
32. *T. CONGENNUM* (Willd. enum. 584.) caule tereti striato, floribus cernuis in panicula patentissimâ amplam dispositis, foliorum segmentis glabris cuneiformibus 3-fidis acutis, lobo medio sæpè 3-dentato. \mathcal{U} patr. ign. Affine *T. minori* et *T. medio* sed habitu strictiore jam è longinquo distinctum. DC. syst. 1. p. 180. (v. v.)
33. *GLAUDESCENS* (DC. syst. 1. p. 180.) caule tereti striato, floribus subnutantibus laxè paniculatis, stipellis minimis ad basin præcipuorum petioli ramificationum, segmentis glabris ovatis apice 3-dentatis. \mathcal{U} in Imperio Ruthenico? (v. v.)
34. *T. GALIÖIDES* (Nestl! in Pers. ench. 2. p. 101.) caule tereti subsulcato stricto, radice repente, panicula stricta, floribus nutantibus, foliorum segmentis linearibus angustissimis integerrimis margine revolutis, ultimo subsessili. \mathcal{U} in pratis subarenosis Alsatiæ circa Argentoratam. Deless! ic. sel. 1. t. 11. *T. angustifolium* β . galiöides. DC. fl. fr. n. 4601.* florens è longinquo adeò simile Galio vero ut vix discerni possit. In h. Linnæano mixtum cum *T. angustifolio* adest. DC. syst. 1. p. 180. (v. v.)
35. *T. ANGUSTIFOLIUM* (Jacq. h. vind. 3. t. 43.) caule stricto tereti subsulcato, radice fibrosa, panicula multiplici erecta, floribus erectiusculis, foliorum segmentis lineari-lanceolatis integerrimis, ultimo subinciso remoto. \mathcal{U} in Germaniâ et Galliâ. *T. angustifolium* var. Lin! spec. 769. Carpella ovata striata brevissima. DC. syst. 1. p. 181. (v. v.)
36. *T. LUCIDUM* (Lin! spec. 770.) caule ramoso tereti subsulcato, radice fibrosâ, paniculâ multiplici erecta, floribus erectiusculis, foliorum segmentis lineari-lanceolatis integris, ultimo lobato. \mathcal{U} in pratis circa Parisios, in Hispaniâ. *T. medium* Poir! dict. 5. p. 316.* excl. syn. Clus. hist. 1. p. 234. — medium inter *T. angustifolium* et *T. flavum*. DC. syst. 1. p. 181. (v. s.)
37. *T. NIGRICANS* (Jacq! fl. austr. 5. t. 421.) caule stricto ramoso sulcato, radice fibrosâ, panicula multiplici erecta subcorymbosâ, foliorum radicalium segmentis cuneiformibus 3-fidis, caulinarum oblongo-linearibus. \mathcal{U} in Austriâ, Italiâ sept., Galliâ merid. — Moris. hist. 1. s. 9. t. 20. f. 3. *T. rugosum* Poir. dict. 5. p. 317. exclus. syn. et patr. — DC. syst. 1. p. 182. an differt à *T. simplicii*. ex cl. Requier? (v. v.)
38. *T. FLAVUM* (Lin! spec. 770. var. α .) caule erecto ramoso sulcato, radice fibrosâ, paniculâ multiplici erectâ subcorymbosâ, foliorum omnium segmentis cuneiformibus 3-fidis acutis. \mathcal{U} in Europâ, Sibirîâ et agro Bizantino. *T. pratense* Lin. fl. lap. 224. — J. Bauh. hist. 3. p. 487. f. 1. (v. s.)
- β . *pauperculum*, panicula pauciflora. DC. fl. fr. n. 4603.* (v. s.)
- γ . *vaginatum*, petiolorum vaginis in auriculas expansis. — *T. vaginatum* Desf! cat. h. par. ed. 2. suppl. 274. DC. syst. 1. p. 182. (v. v.)

39. *T. THUNBERGII* (DC. syst. 1. p. 183.) caule erecto ramoso tereti, panicula multiplici erecta, foliorum segmentis cuneiformibus apice acutè 3 fidis glaucescentibus. \mathcal{U} in Japoniâ. Deless! ic. sel. 1. t. 12. *T. flavum* Thunb! fl. jap. 241. non Lin. Affine *T. flavo* sed distinctum, flores paulò minores, ovaria 3-5 nec 10-15, stamina 10-12 nec 24. DC. syst. 1. p. 183. (v. s.)
40. *T. SIMPLEX* (Lin! mant. 78.) caule erecto simplici angulato, radice repente, paniculâ erectâ racemosâ pauciflorâ, foliorum segmentis linearibus aut inferioribus oblongis acutis 3-fidis. \mathcal{U} in Sueciâ, Norvegiâ, Helvetiâ. *T. angustifolium*. Vill. Dauph. 3. p. 722.* exclus. synonym ad. fl. dan. t. 244. DC. syst. 1. p. 183. (v. s.)
41. *C. CINEREUM* (Desf! cat. h. par. ed. 2. p. 146.) caule erecto tereti substriato, paniculâ ramosissimâ laxâ rigidâ, floribus erectis, foliorum segmentis ovalibus basi cuneatis, apice 3-5 dentatis. \mathcal{U} Hab. ign. Ovaria 7-8 oblonga apice acuta. DC. syst. 1. p. 183. (v. s. in h. Desf.)
42. *T. DENSAIFLORUM* (H. B. et Kl nov. gen. et spec. amer. 5. p. 38.) caule erecto striato, paniculâ multiplici erectâ, floribus congestis subcapitatis, foliorum segmentis ovalibus acutè 3-fidis 3-dentatisve subtus glaucis reticulato-venosis. \mathcal{U} in Americâ merid. Ovaria pauca, forsan interdum nulla aut abortiva ut in plerisque speciebus americanis. DC. syst. 1. p. 184. (v. s. in h. H. et B.)
43. *T. GLAUCUM* (Desf! cat. h. par. ed. 2. p. 146.) caule erecto tereti striato poliline glauco, panicula multiplici erectâ confertâ, foliorum segmentis subcordato-ovatis obtusè 3-fidis subtus glaucis. \mathcal{U} in Hispaniâ, Lusitaniâ, Pyrenæis et Helvetiâ. Schrad. h. gött. 1. 14. t. 8*. — Moris. hist. 1. s. g. t. 20. f. 1. *T. speciosum* Poir. dict. 5. p. 315*. Ovaria 4-6, ovata. DC. syst. 1. p. 184. (v. v.)
44. *T. RUGOSUM* (Ait. h. Kew. ed. 1. v. 2. p. 262.) caule erecto teretis triato virecente, paniculâ erectâ multiplici, floribus confertis, foliorum segmentis ovatis subcordatis grossè 3-5-crenatis, subtus glaucis, supernè lucidis. \mathcal{U} Secus rivos Americæ bor. à Pennsylvania ad Carolinam. *T. discolor* Willd enum. app. 46. Valdè simile *T. glauco*. DC. syst. 1. p. 185. (v. v.)
- β . *umbelliferum*, caule crassiore elatiore apice in pedunculos longos umbellatos desinente, floribus ad apicem cujusque pedunculi plurimi confertis. An species propria?

§. 3. Indivisa. *Foliis indivisis.*

45. *T. ROTUNDFOLIUM* (DC. syst. 1. p. 185.) foliis radicalibus petiolatis reniformibus suborbiculatis sinuato crenatis co-nerviis. \mathcal{U} in Napauliâ. *T. Batula* Buchan! ined. Ovaria sessilia oblonga stigmate acutiusculo terminata, in capitulum globosum collecta; carpella substriata. (v. s.)
46. *T. RANUNCULINUM* (Mühl. in Willd. enum. 585.) foliis simplicibus 5-lobis serratis. — in Amer. bor. Folia Ranunculi vel Geranii 5-lobo-palmata; flores corymbosi *T. speciosi*. DC. syst. 1. p. 186. †

§. 4. Grumosa. *Radicibus grumosis, sepalis petaloideis stamina superantibus.*

47. *T. TURKOSUM*. (Lin! spec. 768.) radice grumosâ, floribus laxè corymbosis solitariisve, involucro nullo, bracteis sessilibus. \mathcal{U} in pascuis sterilibus et petrosis collium Aragoniæ et Pyrenæorum. Moris. hist. 2. p. 438. s. 4. t. 28. f. 13. Flores plurimi pedunculati laxè corymbosi. Sepala 5 alba petalis Parnassæ ferè similia. DC. syst. 1. p. 186. (v. v.) β . caule unifloro. DC. fl. fr. 5. p. 623.
48. *T. ANEMONOIDES* (Michx! fl. bor. amer. 1. p. 322.) radice grumosâ, floribus umbellatis, foliis floralibus petiolatis biternatis sectis involucrum constituentibus. \mathcal{U} in Americâ sept. *Anemone thalictroides* Lin. spec. 763. Juss! ann. mus. 3. p. 249. t. 21. f. 2. Habitus et frondescentia *Isopyri*, florum dispositio *Anemones*, fractus *Thalictri*.

β. uniflorum (Pursh. fl. bor. amer. 2. p. 387.)

†. *Species non satis notæ.*

49. *T. MEXICANUM* (DC. syst. 1. p. 187.) Coztipatli n. Hern. mex. 236. ic. \mathcal{U} in Mexico. Radices crassæ fasciculato-fibrosæ, amethystini coloris, fibrillis luteis; caulis luteus; folia biternatim secta, segmenta petiolulata ovata denticulata. (Hern.)
50. *T. LASERPITTIPOLIUM* (Willd. ex. Horn. h. hafn. suppl. 62.) foliolis inferioribus cuneatis 3-fidis, caulinis linearibus, terminali 3-partito, floribus erectis. \mathcal{U} hab. ign.
51. *T. DIVARICATUM* (Horn. h. hafn. add. 967.) caule folioso stricto sulcato, foliolis ternatis linearibus 3-fidis divaricatis, paniculæ strictæ floribus erectis. \mathcal{U} hab. vign.
52. *T. ? SINENSIS ?* (Lour. fl. coch. 1. p. 423.) \mathcal{J} \mathcal{U} in China. Radix tuber subrotundus solidus albidus; caulis pedalis teres rectus suffruticosus simplicissimus 1-florus; flos albus terminalis; calyx o. petala ? 5, rotundo-patentia; stamina polyandra breviter; semina plurima mucosa, folia lineari-lanceolata. (Lour.) An *Thalictrum* ? an *Ranunculacea* ? DC. syst. 1. p. 187.

IV. TETRACTIS. *Spreng. neu. entd.* 3. p. 53.

Invol. sub flore o. Cal. 4-sepalus obtusus. Pet. 'o. Stam. 4. Antheræ oblongæ basi insidentes. Carpella caryopsidea 4 acuta. — Fruticulus foliis alternis oblongis subintegerrimis. — Genus mihi ignotum huc ex auctoritate cl. Sprengelii relatum.

1. *T. CAPENSIS* (Spreng. l. cit.) 5 ad Cap. Bonæ-Spei. Pedunculi capillares, versus ramulorum apicem conferti. Calyces rubri.

V. ANEMONE. *DC. syst.* 1. p. 188.

Involucrum 3-folium à flore distans; foliolis incis. Calyx petaloideus, 5-15-sepalus. Petala o.

Sect. I. PULSATILLA. *C. Bauh. pin.* 177.

Cariopsides in caudas longas barbatas productæ; involucri folia sessilia in lobos lineares palmati-partita.

1. *A. VERNALIS* (Lin! sp. 759.) foliis pinnati-sectis, segmentis cuneato-lanceolatis trifidis glabriusculis, flore erecto, involucri villosissimo, sepalis 6 rectis elliptico oblongis. \mathcal{U} in scopulis et pratis montium excelsorum Europæ ad nivem deliquescentem. — *Æd. fl. dan. t. 29.* — *Pulsatilla vernalis.* Mill. dict. n. 3. Variat foliis radicalibus subvillosis aut glabris. (v. v.)
- β . flore luteo — *Pulsatilla lutea apii hortensis folio.* C. Bauh. pin. 177.
- γ ? *Pulsatilla apii folia autumnalis.* C. B. prod. 94.
2. *A. HÄLLERI* (All! pedem. n. 1922. t. 80. f. 2.) foliis pinnati-sectis villosissimis, segmentis 3-partitis, lobis lobatis, lobulis lanceolato-linearibus acuminatis, flore erecto, sepalis 6 ovali-lanceolatis. \mathcal{U} . ad nivem deliquescentem in summis Alpibus. Pili totius plantæ longi sericei argentei; flos magnus erectus intus purpurascens. DC. syst. 1. p. 190. (v. v.)
3. *A. CERNUA* (Thunb. fl. jap. 238) foliis pinnati-sectis subtus villosis-subtomentosis, segmentis pinnatifidis, lobis incis. oblongis, flore subcernuo, sepalis 6 patentibus elliptico-oblongis. \mathcal{U} in Japoniâ, juxta Jedo. DC. syst. 1. p. 190. (v. s. in h. Lamb.)
4. *A. PATENS.* (Lin! sp. 759) foliis ternati-sectis hysteroantheis, segmentis tripartitis, lobis apice dentato-incis. flore erecto-patente. \mathcal{U} in campis sylvestribus Sibiriae. — Breyn. ic. t. 61. Species distinctissima in eo quod flore

nascuntur antè folia, et sic planta florens aphylla, et fructifera foliis radicalibus donata. DC. syst. 1. p. 191. (v. s.)

β. *ochroleuca*, foliis isochronis, flore ochroleuco. Sims. bot. mag. t. 1994. in toto imperio Ruthenico orientali. (Fisch.)

5. A. *PULSATILLA* (Lin! sp. 759.) foliis pinnati-sectis, segmentis multipartitis, lobis linearibus, flore subnutante, sepalis 6 patentibus. 2 in campis sylvestribus collibusque apricis Europæ et Sibirizæ. — Clus. hist. 1. p. 246. f. 1. DC. syst. 1. p. 191. (v. v.)

β. *rubra* (Lam! dict. 1. p. 163. *) staturâ nanâ, flore erecto, sepalis obtusioribus.

γ. *lilacina*. A. intermedia Schult. obs. 101. A. longipetala Schl! exsic.

δ. *Dahurica*, pumila, villosissima, flore erecto, sepalis oblongis. (v. s. in h. Lamb.)

6. A. *PRATENSIS* (Lin! sp. 760.) foliis pinnati-sectis, segmentis multipartitis, lobis linearibus, flore pendulo, sepalis 6 erectis apice reflexis. 2 in campis pratensibus apricis Galliæ, Germaniæ, Daniæ. — *Pulsatilla nigricans* Stork. lib. puls. ic. Ab. A. *Pulsatilla* differt flore minore pendulo nec suberecto, colore saturatiore, sepalis angustioribus acutioribus basi erecto-conniventibus apice summo reflexis. DC. syst. 1. p. 192. (v. v.) †

β. *obsoleta*, flore majore pallido, pinnularum lobis latioribus aristatis. Sims. bot. mag. t. 1863. — Helw. puls. p. 65. t. 11.

7. A. *ALBANA* (Stev. mem. soc. nat. mosc. 3. p. 264) foliis pinnati-sectis, segmentis multipartitis, lobis oblongo-linearibus, flore nutante, pedicello vix involucri longiore, sepalis 6 erectis apice subreflexis. 2 in excelsis montibus Caucasii orientalis. Affinis A. *pratensis*, sed differt flore albedo aut ochroleuco nec purpureo. DC. syst. 1. p. 545. (v. s.)

8. A. *DAHURICA* (Fisch! in litt.) foliis ternati-sectis, segmentis 3 lateralibus sessilibus inæqualibus 3-lobis, terminali petiolato 3-partito, lobis 2 et 3-fidis omnibus linearibus acutis, sepalis 6 erecto-conniventibus. 2 in Dahuriâ ad flumen Ingodam. Flos carneus, magnitudine A. *pratensis*, sed pallidus; fructus A. *Pulsatillæ*. (v. s. sp. comm. à cl. Fisch.)

9. A. *NUTTALLIANA* (DC. syst. 1. p. 193.) foliis ternati-sectis, segmentis tribus multifidis, lobis linearibus elongatis, involucralibus in lobos lineares multi-sectis; flore erecto, sepalis 6 erecto-conniventibus. 2 in Louisiana. Clematis hirsutissima Pursh. fl. bor. amer. 2. p. 385. DC. syst. 1. p. 193. A. *Ludoviciana* Nutt. gen. am. 2. p. 20.* Flos purpureus erectus, fructus A. *Pulsatillæ*. (v. s. in h. Lamb.)

Sect. II. PREONANTHUS. DC. syst. 1. p. 193.

Cariopsides in caudas longas barbatas productæ. Involucri folia trisecta petiolata.

10. A. *ALPINA* (Lin! spec. 760.) foliis petiolo ramoso biternatim sectis, segmentis pinnatim inciso-serratis, involucralibus conformibus, sepalis 6 patentibus. 2 in pratis montium Europæ mediæ. DC. syst. 1. p. 193. (v. v.)

α. *major*, floribus magnis albis extus subpurpurascentibus, sepalis ellipticis; foliis glabratiss aut parcè villosis antè florem expansis. Sims. bot. mag. t. 2007. A. *apiifolia* Hop! h. viv. DC. fl. fr. 4. p. 881. — variat flore pleno.

β. *millefoliata*, floribus magnis albis extus purpurascentibus, sepalis ellipticis, foliis synanthesis, lacinii lanceolatis acutis. A. *millefoliata* Bertol. amœn. 374.*

γ. *micrantha*, floribus parvulis albis extus subpurpurascentibus, sepalis ellipticis, foliis glabriusculis ante florem expansis. A. *Alpina*. Jacq. fl. austr. T. 85.

δ. *flavescens*, floribus magnis flavis extus pallidoribus, sepalis ellipticis, foliis parcè villosis ante florem expansis. A. *apiifolia* Scop. fl. car.

- n. 663.* Jacq. misc. 2. p. 47. t. 4.* *A. myrrhidifolia*, β . Vill. Dauph. 4: p. 727.
8. *nivalis*, floribus mediocribus albis subtus subpurpurascens, sepalis ellipticis, foliis villosissimis cum flore aut post florem expansis, scapo nano.
9. *intermedia*, floribus magnis flavis extus pallidioribus, sepalis ellipticis, foliis villosissimis ante florem aut cum flore nascentibus.
- n. *sulphurea*, floribus mediocribus flavis extus subpallidioribus, sepalis ovatis, foliis villosissimis ante florem aut cum flore expansis. *A. sulphurea* Lin. mant. 78. *A. apiifolia* Willd. sp. 2. p. 126. *Pulsatilla lutea* Cam. epit. 393. ic.

Sect. III. PULSATILLOIDES. DC. syst. 1. p. 195.

Cariopsides hirsutissimæ. Calyces 15-20-sepali. Involucrum folia 2-3-sessilia apice incisa.

- ✓ 11. *A. CAPENSIS* (Lin! sp. 764.) foliis biternatim sectis rigidis glabris, segmentis cuneiformibus apice dentatis. \mathcal{L} ad Caput Bonæ-Spei. Sims. bot. mag. t. 716. *Clematis Capensis* Poir. Dict. suppl. 2. p. 296. Sepala intus glabra purpurascens, extus sericea. Fruct. ign. DC. syst. 1. p. 195. (v. v.)
12. *A. TENUIFOLIA* (DC. syst. 1. p. 196) foliis triternatim sectis glabris, segmentis pinnatipartitis, lobis lineari-filiiformibus acutis integris. \mathcal{L} ad C. B.-Spei. *Atragene tenuifolia* L. fil. suppl. 270. Flos dimidio minor quam in *A. capensis*. (v. s. in h. Deless.)

Sect. IV. ANEMONANTHEA. DC. syst. 1. p. 196.

Cariopsides ecaudatæ ovoidæ. Pedicelli solitarii aut bini, omnes aphylli uniflori. Involucrum foliola sessilia aut petiolata.

§. 1. *Involucrum foliis sessilibus; radice truncata ovoidæ tuberosa.*

- ✓ 13. *A. CORONARIA* (Lin! spec. 760.) foliis ternatim sectis, segmentis multifidis, lobulis linearibus mucronatis, involucralibus sessilibus multifidis, sepalis 6 ovalibus approximatis. \mathcal{L} in pascuis subhumidis Europæ. In hortis ornamentati gratia. — bot. mag. t. 841. Variat flore purpureo cærulescente violaceo lilacino flavido aut albo aut variegato, simplici aut pleno. DC. syst. 1. p. 196. (v. v.)
14. *A. PUSILLA* (DC. syst. 1. p. 197.) foliis ternatim sectis, segmentis multifidis, lobulis linearibus mucronatis, involucralibus sessilibus apice incisis, sepalis 6 oblongis distantibus. \mathcal{L} in insula Cypro. *A. pusilla* Deless! ic. sel. 1. t. 12. Nimis affinis *A. coronariæ* et forsitan varietas. Carpella lanata aggregata in capitulum ovali-oblongum. (v. s. in h. Desf.)
- ✓ 15. *A. PAVONINA* (DC. syst. 1. p. 197.) foliis tripartitis, lobis cuneatis inciso-dentatis, involucralibus sessilibus oblongis integris subincisive, sepalis 10-12 lanceolatis acutissimis. \mathcal{L} in Gallo-provincia. *A. latifolia*, etc. Moris. oxon. s. 4. t. 25. f. 10. Hanc ex synonymia ad sequentem retulit Linnæus, sed in ejus herbario inter *A. coronariæ* varietates reperi. DC. syst. 1. p. 197. exclusis syn. Loisel. et Thor. (v. v. et s.)
- ✓ β ? *fulgens*, foliis 3-partitis, lobis cuneatis inciso-dentatis, involucralibus sessilibus oblongis integris subincisive, sepalis oblanceolatis apice latioribus basi attenuatis. \mathcal{L} in vineis Vasconiarum circa Dax. *A. fulgens*. Gay. ined. *A. hortensis* Thor. chl. land. 238. *A. pavonina*. Lois. not. 87. Flores ampliores quam var. α . (v. s.)
- ✓ 16. *A. STELLATA* (Lam. dict. 1. p. 166.) foliis 3-partitis, lobis cuneatis inciso-dentatis, involucralibus sessilibus oblongis integris subincisive, sepalis 10-12 oblongis obtusiusculis. \mathcal{L} in sepiibus et dumetis Europæ. *A. hortensis* Lin! spec. 761. Curt. bot. mag. t. 123. Δ . versicolor Salisb. prod. 371. Rarius in hortis colitur et sæpius flore simplici donatur quam *A. pavonina* quæcumdiu fuit confusa. DC. syst. 1. p. 198. (v. v. et s.)

- ✓ 17. *A. PALMATA* (Lin! spec. 758) foliis cordatis suborbiculatis obsolete 3-5-lobis dentatis; involucralibus sessilibus trifidis, sepalis 10-12 oblongis obtusis. \mathcal{L} in locis humidis Algeriæ, Lusitanicæ, Hispaniæ. Ker. bot. reg. t. 200.*
 β ? *flore pleno*. *A. hortensis-latifolia*, etc., Clus. hist. 1. p. 249. f. 1. — Ab *A. stellatâ* differt ob *florem flavum*, etc. DC. syst. 1. p. 199. (v. s.)
- ✓ 18. *A. DECAPETALA* (Lin! mant. 79.*) foliis tripartitis, lobis grosse dentatis aut subincisis, involucralibus sessilibus bis-trifidis, lobis linearibus, sepalis 10-12 oblongis obtusis. \mathcal{L} in Brasiliâ, Peruvîâ et regni Chilensis sylvis. *A. trilobata* Juss! ann. mus. 3. p. 247. t. 21. f. 3.* *Flos solitarius* 10-12 sepalis albus extus cærulescens. DC. syst. 1. p. 200. (v. s.)
- ✓ 19. *A. PARVIFLORA* (Michx! fl. bor. amer. 1. p. 319.) foliis 3-partitis, lobis cuneatis apice truncato-crenatis, involucralibus sessilibus 3-partitis, lobis apice subdentatis, sepalis 6 ovali-oblongis. \mathcal{L} in Americâ septentr. *A. cuneifolia*. Juss! ann. mus. 3. p. 248. t. 21. f. 1. Affinis *A. decapetalæ* et *palmatæ*; fructus lanati acuminati in capitulum globosum aggregati. DC. syst. 1. p. 200. (v. s.)
20. *A. CAROLINIANA* (Walt. fl. car. 157.) foliis ternatim partitis, lobis 3-partitis incisive acutè dentatis, involucralibus trifidis, lobis incisis, sepalis 10-12 linearibus. \mathcal{L} in Carolinâ et ad ripas Missouri. *A. tenella* Pursh! fl. bor. am. 2. p. 386. fructus lanati mucronati. DC. syst. 1. p. 201. (v. s. in h. Lamb.)
21. *A. TRITERNATA* (Vahl. symb. 3. p. 74. t. 65.) foliis petiolo ramoso triternatim sectis, segmentis tridentatis, involucralibus sessilibus in lobos setaceos multifidis, sepalis 10-12 oblongis obtusis. \mathcal{L} in Brasiliâ et Peruvîâ. *A. fumaricæ-folia* Juss! ann. mus. 3. p. 247. t. 20. f. 2.* Radix et habitus *A. decapetalæ*, fructus lanati in capitulum oblongum digesti. DC. syst. 1. p. 201. (v. s.)
22. *A. BIFLORA* (DC. syst. 1. p. 201.) foliis ternatim sectis, segmentis partitis in lobos lineares obtusos subincisos, involucralibus sessilibus multipartitis, pedicellis geminis, altero involucellato. \mathcal{L} in Oriente.
 α. *bifoliata*, involucro involucelloque bifoliolatis. — flores subcernui flavi 5-sepali.
 β. *trifoliolata*, involucro trifoliolato. — flores longius pedicellati ex albido subpurpurascens. Carpella lanata ut in *A. Baldenai*. An ab hac satis differt? *A. formosa* Spreng. neu. entd. 3. p. 207 (v. s. in h. Deless.)
- §. 2. *Involucri foliolis petiolatis; radice trunco ovoideo tuberoso.*
- ✓ 23. *A. APENNINA* (Lin! spec. 762.) foliis petiolo ramoso triternatim sectis, segmentis lanceolatis inciso-dentatis acutis, involucralibus petiolatis trisectis inciso-dentatis, sepalis 12-14 oblongis obtusis. \mathcal{L} in Belgio, montibus Apenninis, Romanis, Neapoli. — Engl. bot. t. 1060. Curt. Lond. 6. t. 35. *A. geraniifolia* J. Bauh. hist. 3. p. 405. f. 2. DC. syst. 1. p. 202. (v. s.)
 β. *Ranunculus nemorosus*, etc. flore albo. Tournef. cor. 20.
 γ. *parvula*, foliis ferè sessilibus. — in Iberiâ Caucasos et ad ripas Simois.
- §. 3. *Involucri foliolis petiolatis, radice trunco cylindræco tenui elongato.*
24. *A. CŒRULEA* (DC. syst. 1. p. 203.) foliis , involucralibus breviter petiolatis trisectis, segmentis inciso-dentatis, sepalis 4-5 ovalibus. \mathcal{L} in Sibirîâ Altaicâ. Deless! ic. selec. 1. t. 14. Variat caule 1-2-floro, floribus cæruleis aut albis. (v. s.)
25. *A. URALENSIS*, foliis , involucralibus breviter petiolatis trisectis, segmentis linearibus inciso-dentatis, sepalis 5-6 ovali-oblongis. \mathcal{L} in montibus Uralensibus. Comm. a cl. Fischer. Planta parva habitu et magnitudine ferè *A. cœruleæ*. (v. s.)
- ✓ 26. *A. BALDENSI* (Lin. Mant. 78.) foliis petiolo ramoso bitermatim sectis, segmentis multipartitis, lobis linearibus, involucralibus breviter petiolatis mul-

- tifidis, sepalis 8-10 oblongo-subovalibus. \mathcal{L} in asperis editis montis Baldi, Alpium Helvetiæ, Pedemontii, etc. A. fragifera. Murr. syst. 510. Jacq! ic. rar. 1. t. 103. Sepala alba, extus adpressè pubescentia et subrubentia. DC. syst. 1. p. 203. (v. v.)
- ✓ 27. A. NEMOROSA. (Lin! spec. 762.) foliis ternatim sectis, segmentis 3-fidis inciso-dentatis lanceolatis acutis, involucralibus petiolatis conformibus, sepalis 6 ellipticis. \mathcal{L} vulgaris in nemoribus sepius umbrosis Europæ et Sibiria. OED. fl. dan. t. 549. Variat fl. roseo et purpureo.
- β . *quinquefolia*, foliis sub-5-partitis. \mathcal{L} in Americâ boreali et Sibiria. A. quinquefolia. Lin. spec. 762. Nullo modo ab Europæâ differt; hujus foliorum lobi laterales sunt sæpè profundè bipartiti, quod etiam in Europæanis speciminibus observatur. DC. syst. 1. p. 203. (v. v.)
28. A. ISOPYROIDES (Juss! ann. mus. 3. p. 249. t. 20. f. 3.) foliis petiolo ramoso biternatim sectis, segmentis inciso-3-dentatis subcuneatis, involucralibus petiolatis ternatim sectis, segmentis lateralibus bifidis, sepalis 5 oblongis. \mathcal{L} Hab. ign. — Valdè similis A. nemorosa. DC. syst. 1. p. 205. (v. s. in h. Juss.)
29. A. FISCHERIANA, foliis petiolo ramoso biternatim sectis, involucralibus brevissimè petiolatis, lobis elongatis acuminatis, pedicellis 2 pubescentibus, sepalis 5 ellipticis. \mathcal{L} in Sibiria, propè Salair. Fisch! in litt. flores albi ferè omninò A. isopyroidis. Carpella villosa-pubescentia. (v. s.)
30. A. LANCEIFOLIA (Pursh! fl. bor. amer. 2. p. 386.) foliis omnibus petiolatis ternatim sectis, segmentis lanceolatis crenato-dentatis, sepalis 5 ovatis acutis. \mathcal{L} in Pennsylvania et Virginia. Ab A. trifolia vix ac ne vix differre videtur foliorum segmentis lanceolatis, nec ovato-lanceolatis, floribus paulo majoribus; fructus ovati, stylo brevi uncinati. DC. syst. 1. p. 205. (v. s. in h. Lamb.)
- ✓ 31. A. TRIFOLIA (Lin! spec. 762.) foliis omnibus petiolatis ternatim sectis, segmentis ovato-lanceolatis acutis dentatis, sepalis 5 ellipticis obtusis. \mathcal{L} in Gallia, Pedemontio, Carniola, Sibiria. Sturm. deutschl. fl. ic. Moris. oxon. 2. p. 424. s. 4. t. 25. f. 1. Flos A. nemorosa similis. DC. syst. 1. p. 205. (v. s.)
32. A. MINIMA (DC. syst. 1 p. 206.) foliis , involucralibus petiolatis 3-partitis, lobis ovatis acuminatis extus et ad apicem serratis, sepalis 5 ovali-oblongis obtusis. \mathcal{L} in Virginia. Omnium tenuissima. A. trifolia affinis sed triplò minor; ovaria pauca pubescentia. DC. syst. 1. p. 206. (v. s. in h. Beauv.)
- ✓ 33. A. RANUNCULOIDES (Lin! spec. 762.) foliis radicalibus 3-5 sectis, segmentis sub-3-fidis inciso-dentatis, involucralibus subpetiolatis tripartitis inciso-dentatis, sepalis 5-6 ellipticis. \mathcal{L} in Europâ bor. et mediâ, nec non in Sibiria. fl. dan. t. 140. Engl. bot. t. 1484. Lodd. cab. t. 556. A. lutea. Lam! fl. fr. 3. p. 318. DC. syst. 1. p. 20. 6. (v. v.)
34. A. REFLEXA (Steph. in Willd. spec. 2. p. 1282*) foliis ternatim sectis, segmentis subtrifidis apice dentatis, involucralibus petiolatis conformibus, sepalis 5-6 linearibus obtusis reflexis. \mathcal{L} in Sibiria. Deless! ic. sel. 1 t. 15. flos triplò minor quàm in A. ranunculoide. DC. syst. 1. p. 207. (v. s.)
- §. 4. *Involucri foliolis petiolatis, radicibus fibroso-fasciculatis.*
35. A. SYLVESTRIS (Lin! spec. 761.) foliis ternatim quinatimve sectis, segmentis apice inciso-dentatis, involucralibus petiolatis conformibus, pedicello solitario, sepalis 6 ellipticis, fructibus hirsutissimis. \mathcal{L} in Gallia, Italia septentr. Germania et Sibiria. — Cart. bot. mag. t. 54. (v. v.)
- β . A. sylvestris alba minor C. Bauh. pin. 176. Propè Herbiopolim. DC. syst. 1. p. 207. (v. s.)
- ✓ 36. A. ALBA (Juss! ann. mus. 3. p. 248. t. 20. f. 1.*) foliis ternatim quinatimve sectis, segmentis apice inciso-dentatis, involucralibus petiolatis conformibus, pedicello solitario, sepalis 5 obovatis, fructibus hirsutissimis. \mathcal{L} in Da-

huria, Sims. bot. mag. t. 2167. *A. Ochotensis*. Fisch. h. gor. 47. *Simillima A. sylvestri* et forsan pro hujus merâ varietate habenda. DC. syst. 1. p. 208. (v. s.)

Sect. v. ANEMONOSPERMOS. DC. syst. 1. p. 208.

Cariopsides ecaudatæ subcompressæ. Pedicelli ex involucrio oo, 1 aphyllus 1-florus, 2-3 involucrellum bifolium gerentes.

37. *A. VIRGINIANA* (Lin! spec. 761.*) foliis ternatim sectis, segmentis 3-fidis acuminatis dentato-incisis, involucri involucrellorumque petiolatis conformibus, sepalis 5 ellipticis. \mathcal{L} in Americâ boreali. — Herm. par. 18. ic. Gaert. fruct. 1. p. 357. t. 74. Carpella compressa hirsuta, matura in lanæ filis divaricatis è loco suo extracta et dispersa. DC. syst. 1. p. 208. (v. v. s.)

38. *A. MULTIFIDA* (DC. syst. 1. p. 209.*) foliis radicalibus ternatim sectis, segmentis multifidis, lobis linearibus, involucri involucrellorumque multifidis breviter petiolatis, sepalis 5-10 ellipticis obtusis. \mathcal{L} in Americâ bor. ad fretum Magellanicum.

a. Magellanica, pedicellis ternis. *A. multifida* Poir! suppl. 1. p. 364.

Deless! ic. sel. 1. t. 16. (v. s.)

ß. Hudsoniana, pedicellis binis. *A. multifida* Banks! herb. (v. s.)

γ. uniflora, *A. multifida* var. γ Deless. ic. sel. 1. t. 17.

39. *A. PENNSYLVANICA* (Lin. mant. 247.*) foliis 3-partitis subtrisectis, lobis inciso-dentatis acuminatis, involucrellis sessilibus conformibus, sepalis 5 ellipticis, carpellis villosis. \mathcal{L} in Americâ boreali. *A. aconitifolia* Michx! fl. bor. amer. 1. p. 320. (v. s.)

ß. ? Laxmanni. Involucrio bifolio, pedicellis lateralibus basi imâ involucrellatis. \mathcal{L} in Sibirîâ. Verosimiliter species distincta. (v. s. in h. Pall.)

40. *A. DICHOTOMA* (Lin. amœn. 1. p. 155.*) foliis 3-partitis, lobis oblongis apice inciso-dentatis, involucrellis sessilibus conformibus, omnibus bifoliis, sepalis 5 ellipticis, carpellis glabris. \mathcal{L} in Sibirîâ, Dahuriâ et Canadâ. Gmel! sib. 4. p. 1797. n. 37. exclus. syn. Lin. f. decad. 29. t. 15. Similis *A. Pennsylvanica*, sed differt staturâ graciliori. An planta Americana omninô eadem ac Sibirica? DC. syst. 1. p. 210. (v. s.)

41. *A. MEXICANA* (H. B. et Kl! nov. gen. et spec. amer. 5. p. 41.*) foliis 3-sectis, segmentis ovalibus subcuneatis apice inciso-dentatis, involucrellis duobus sessilibus incisis dentatis, ovariis pubescentibus. \mathcal{L} in Mexico. Herba ex omni parte subvillosa; flores albi sat similes *A. Pennsylvanica*. DC. syst. 1. p. 210. (v. s. in h. H. et B.)

42. *A. HELLEBORIFOLIA* (DC. syst. 1. p. 211.*) foliis pedatis sectis, segmentis glabris subcoriaceis subpetiolulatis basi cuneatis trifidis, lobis serratis acutis, involucri omnibus trifoliis subsessilibus, ovariis glabris. \mathcal{L} in Americâ merid. Planta speciosa distinctissima. (v. s. in h. Mus. Par.)

43. *A. VITIFOLIA* (Buchan! in DC. syst. 1. p. 210*) foliis . . . involucrellis petiolatis subtus albo tomentosissimis, lobis obtusè cordatis 5-lobis, involucrellisque 3-foliis, ovariis glabris. \mathcal{L} in Napauliâ. Sepala ovali-oblonga extus villosa albida, intus purpurascens (v. s.)

44. *A. RIVULARIS* (Buchan! in DC. syst. 1. p. 211.*) foliis subvillosis 3-partitis, lobis cuneatis 3-fidis, lobulis incisis acutè dentatis, involucrellis sessilibus 3-partitis, lobis apice subpinnatifidis. \mathcal{L} in Napauliâ. (v. s. in h. Lamb.)

Sect. VI. OMALOCARPUS. DC. syst. 1. p. 212.

Cariopsides compresso-planæ ovali-orbiculatæ glabræ, ecaudatæ; pedicelli oo umbellati aphylli 1-flori.

45. *A. NARCISSEIFLORA* (Lin! spec. 763.) foliis radicalibus subvillosis palmatim 3-5-partitis, lobis inciso-dentatis, lobulis linearibus acutis, involucrellis apice. 3-5 fidis, floribus umbellatis. \mathcal{L} in pascuis montanis præsertim calcareis he-

mispheeræ borealis. Clas. hist. 1. p. 235. f. 1. Jacq. fl. aust. t. 159. DC. syst. 1. p. 212. (v. v.)

β. *fasciculata*, floribus umbellatis subsessilibus fasciculatis. Tourn. cor. 20. voy. 2. p. 245. ic. A. fasciculata Lin. spec. 763. non Vahl. (v. s.)

γ. *monantha*, floribus. solitariis aut binis. A. dubia Bell. app. fl. ped. 232. t. 7. (v. s.)

δ? *pedicellaris*, pedicellis plurimis elongatis, foliis radicalibus trisectis?

ε? *frigida*, pedicellis paucis elongatis, foliis radicalibus glabris in lobos augustissimos lineares partitis.

ζ. *villosissima*, floribus paucis subsessilibus, caule villosissimo. — in Unalaschka. Fischer. (v. s.)

✓ 46. A. UMBELLATA. (Willd. spec. 2. p. 1284*. non Lam.) foliis radicalibus 3-5-partitis, laciniis 3-fidis integerrimis ad margines dense villosis, involucrialibus indivisis, floribus umbellatis. ♀ in Cappadocia montibus. Deless! ic. sel. 1. t. 18. DC. syst. 1. p. 213. (v. s. in h. Tourn. et Desf.)

47. A. SIBIRICA (Lin! spec. 763.) foliis ternatim sectis, segmentis inciso-dentatis ciliatis, involucrialibus breviter petiolatis trisectis conformibus sepalis 6 orbiculatis. ♀ in Sibiria. — Gmel. sib. 4. p. 199. n. 41. Ovaria glabra! hoc caractere ad Omalocarpus accedit. sed fructus maturus desideratur. An potius A. Albæ affinis? DC. syst. 1. p. 214. (v. s. in h. Lin. et Deless.)

†. *Anemones non satis notæ.*

48. A. WALTERI (Pursh. fl. bor. amer. 2. p. 387.) in Carolina. Thalictrum Carolinianum Walt. fl. car. 157. Radix tuberosa; caulis 1-florus nudus; folia palmata longius petiolata; sepalia 5. Deest in herb. Walteriano. ex Pursh. verosimiliter A. parvifloræ affinis? DC. syst. 1. p. 214.

49. A. PEDATA (Rafin. Schmalz. in Desv. journ. bot. 1808, 1. p. 230.) in New-Jersey. — Caulis brevis 1-florus; folia pedatim 5-partita, lobis laciniatis, sepalia 6. DC. syst. 1. p. 214.

VI. HEPATICA. Dill. giess. p. 108. t. 5. Lin. h. clif. 223. DC. syst. 1. p. 215.

Involucrum 3-foliatum flori approximatum 1-florum calyciforme; Foliis integris; Sepala petaloidea 6-9, duplici triplicive serie disposita. Stamina et ovaria ∞. Carpella ecaudata. Scapi plurimi radicales 1-flori.

1. H. TRIBOLA (Chaix in Vill. Dauph. 1. p. 336) foliis cordatis 3-lobis, lobis integerrimis ovatis acutiusculis, petiolis scapisque subpilosis. ♀ in Europâ. — *Anemone Hepatica* Lin! spec. 758. Gd. fl. dan. t. 612. Harb. Sam. t. 61. A. præcox Salisb. prodr. 371. Variat flor. albo, cinereo, carneo, rubro, purpureo et pleno. DC. syst. 1. p. 216. var. α. (v. v.)

2. H. AMERICANA (Ker. bot. reg. t. 387.) foliis cordatis 3-lobis, lobis integerrimis rotundatis obtusis, petiolis scapisque pilosissimis. ♀ in Americâ boreali. — Ray hist. 581. — H. triloba β. DC. syst. 1. p. 216. (v. s.)

3. H. ANGULOSA (DC. syst. 1. p. 217.) foliis palmato-5-lobis, lobis serratis. ♀ olim in h. pharm. Par. culta. *Anemone angulosa* Lam. dict. 1. p. 169*. sepalia 8-9 elliptica patentia.

4. H. ACUTIFLORA, foliis cordatis 3-lobis, lobis integerrimis acutis, petiolis pilosis. ♀ in Americâ sept. *Anemone Hepatica* var. *acuta* Bigelow! an spec. propria? (v. s.)

5. H. INTEGRIFOLIA (DC. syst. 1. p. 217.) foliis ovatis integerrimis, scapis petiolisque villosissimis. ♀ in Americâ merid. alt. 1850 hexap. *Anemone integrifolia* H. B. et K. nov. gen. et sp. am. 5. p. 40*. An involucrum potius 1-phyllum, foliolis ad basin usque 3-partitis? An folia decidua? et tunc ad Pulsatillas accederet hæc species anemones hepaticasque fortè conjungens. (v. s. in h. H. et B.)

VII. HYDRASTIS. *Lin. gen. n. 704. Juss. gen. 232. DC. syst. 1. p. 217. Warneria Mill. ic. t. 285.*

Calyx 3-sepalus, sepalis ovatis; Petala 0. Stamina ovariaque ∞. Cariopsides baccatæ, plurimæ in capitulum aggregatæ, stylo terminatæ, 1-loculares, 1-2-spermæ, semina obovoïdea lævigata.

1. *H. CANADENSIS* (Lin! spec. 784.) ♀ in Canada, Pennsylvania. — Michx! fl. bor. amer. 1. 317. *Warneria Canadensis* Mill. dict. n. 1^o. Fructus carnosus ruber fructui Rubi similis, carpellis ovato-acutis stylo coronatis baccatis constans. DC. syst. 1. p. 218. (v. s.)

VIII. KNOWLTONIA. *Salisb. prod. 372. DC. syst. 1. p. 218. — Anamenia Vent. malm. 12.*

Calyx 5-sepalus. Petala 5-15, ungue nudo. Stam. ∞. Ovaria ∞ receptaculo globoso imposita; cariopsides ∞ 1-spermæ baccatæ, stylo deciduo muticæ. — Herbæ ♀, habitu umbelliferas plurimas æmulantes.

1. *K. RIGIDA* (Salisb. prod. 372.) foliis biternatim sectis, segmentis subcordatis coriaceis glabriusculis, lateralibus basi obliquè truncatis, umbellâ supra-decompositâ patentissimâ. ♀ ad prom. Bonæ-Spei. DC. syst. 1. p. 219. *Adonis capensis* Lin! spec. 772. *Anamenia coriacea* Vent! malm. 1. 22. t. 22.* (v. v.)
2. *K. VESICATORIA* (Sims. bot. mag. t. 775.) foliis biternatim sectis, segmentis subcordatis rigidis glabriusculis, lateralibus basi obliquè truncatis, umbellâ subsimplici pauciflorâ. ♀ ad prom. Bonæ-Spei. DC. syst. 1. p. 219. *Adonis vesicatoria* Lin. fl. supp. 272. (v. s. et v.)
3. *K. GRACILIS* (DC. syst. 1. p. 219.) foliis biternatim sectis, segmentis ovatis profundè serratis rigidis pilosis, scapis apice ramosis, ramis erectis paucifloris ♀ ad Cap. B.-Spei. Deless! ic. sel. 1. t. 19. *Anamenia gracilis* Vent. malm. 1. 22. in obs. (v. s. in h. Juss. missum à Cl. Thunberg sub nomine *Adonidis Capensis*.)
4. *K. HIRSUTA* (DC. syst. 1. p. 220.) foliis biternatim sectis, segmentis lanceolatis profundè serratis hirsutis, scapis basi ramosis, ramis decumbentibus paucifloris. ♀ ad prom. Bonæ-Spei. Burm! afr. 145. t. 51.* *Adonis hirsuta* Poir. supp. 1. p. 147. (v. s. in h. Deless.)
5. *K. DAUCIFOLIA* (DC. syst. 1. p. 220.) foliis ternatim sectis, segmentis pinnatifidis, lobis linearibus acutis, umbella composita. ♀ ad prom. Bonæ-Spei. *Adonis daucifolia* Lam. dict. 1. p. 46. Pedunculi pubescentes. (v. s. in h. Deless.)

IX. ADONIS. *Dill. giess. nov. gen. 109. t. 4. Lam. ill. t. 498.*

Calyx 5-sepalus adpressus, sepalis interdum basi solutis; petala 5-15 ungue nudo; stam. ∞ ad basin gonophori inserta; ovaria ∞; cariopsides ∞, 1-spermæ, secus gonophorum spicatæ, ovatæ, stylo brevi subaccreto acuminatæ; embryo ovatus, cotyledonibus subdistantibus. — Folia caulina pinnati-partita, lobis multifidis. DC. syst. 1. p. 220.

Sect. 1. ADONIA. *DC. syst. 1. p. 221. — Adonis C. Bauh. pin. 178.*

Cariopsides stylo recto uncinato recto acuminatæ; petala 5-10. Radices annuæ. — Species hujus sectionis sunt forsàn unius speciei varietates.

1. *A. AUTUMNALIS* (Lin! spec. 771.) calyce glabro, petalis 6-8 concavo-conniventibus calyce vix majoribus, carpellis subreticulatis in capitulum ovatum aggregatis stylo brevissimo coronatis, caule ramoso. ① inter segetes Europæ. Engl. bot. t. 308. Clus. hist. 1. p. 336. f. 1. Flos intensè sanguineo-ruber, rariùs pallidior. DC. syst. 1. p. 222. Wallr. sched. p. 226. (v. y.)

2. *A. FLAVA* (Vill. cat. strash. 247.) calyce glabro basi soluto, petalis planis oblongis calyce duplò longioribus, carpellis sublaevibus in capitulum oblongum aggregatis, caule sub-simplici. ① inter segetes et in vineis Galliarum. Weinm. phyt. t. 28. f. a et b. exclus. syn. — DC. syst. 1. p. 222. (v. v.)
3. *A. PARVIFLORA* (Fisch! in litt.) calyce basi soluto glabro, petalis planis oblongis calycis longitudine, carpellis in capitulum ovatum collectis paucis stylo conico recto coronatis, caule simplici. ① ad ripas flum. Rhynni propè lacum salsum Inderiensem. Petala pallidè coccinea in sicco albida. (v. s.)
4. *A. MIGRANTHA* (DC. syst. 1. p. 222.) calyce glabro basi non soluto, petalis planis oblongis calyce paulo longioribus, carpellis subreticulatis in capitulum subovatum aggregatis, caule subramoso. ① in segetibus Galliarum australis. Species dubia; flos parvus flavus seu flammeus; ovaria pauca 7-10 in capitulum brevissimum disposita (v. v.) †
5. *A. MICROCARPA* (DC. syst. 1. p. 223.) calyce glabro, petalis planis oblongis calyce duplo longioribus, carpellis reticulatis in capitulum oblongum dispositis, caule subsimplici. ① inter segetes Hispaniarum. Variat flore flavo-citrino et subflammeo. (v. s.) †
6. *A. ANOMALA* (Wallr! sched. 273.) calyce extus piloso acutiusculo, petalis 3 oblongis planis obtusiusculis patentibus ungue concoloribus, carpellis ovatis sexfariis, stylo apice sphacelato brevi in gibbum internum inclinato coronatis, demum in spicam oblongam laxè irregulariterque dispositis, caule ramoso. ① in apricis agri Halensis. (v. s.)
7. *A. CITRINA* (Hoffm. fl. germ. 1. p. 251.) calyce basi hispido, petalis planis oblongis calyce longioribus, carpellis in capitulum ovato-oblongum dispositis, caule subsimplici, flore parvo inter folia subsessili. ① inter segetes Galliarum, Germaniarum, Teneriffarum — *A. maculata* β Wallr! sched. p. 271. — *A. citrina* DC. syst. 1. p. 223. (v. v.)
8. *A. FLAMMEA* (Murr. syst. 514.) calyce basi hispido, petalis planis oblongis acutis calyce longioribus, carpellis in capitulum cylindricum dispositis, caule ramoso, flore magno pedunculato. ① inter segetes Austriarum, Brunsvigiarum. Jacq. fl. aust. 4. t. 355. — Besl. h. Eyst. æst. ord. 5. t. 11. f. 3. Flores pedunculati magni flammei. DC. syst. 1. p. 223. (v. v.)
9. *A. ÆSTIVALIS* (Lin! spec. 772.) calyce basi hispido, petalis planis oblongis obtusis calyce duplò longioribus, carpellis reticulatis, in spicam oblongam laxam dispositis, conoideo-prismaticis, stylo recto concolori rostratis, caule subsimplici. ① inter segetes Galliarum australis, Italiarum. *A. miniata*. Jacq. fl. aust. t. 354. *A. maculata* α . Wallr! sched. p. 270. — *A. æstivalis* DC. syst. 1. p. 224. *A. citrina* et forsàn flava huc forsàn ut varietates spectant. (v. v.)
10. *A. DENTATA* (Del fl. Æg. ill. 17. Desc. Eg. t. 53. f. 1.) calyce basi hispido, petalis planis ovali-oblongis calyce paulò longioribus, carpellis basi tuberculato-dentatis reticulatis in spicam elongatam continuam digestis. ① DC. syst. 1. p. 224.
 - α . *Orientalis*, petalis flavis. *A. dentata* Del. l. c. Inter segetes et in aridis Ægypti. (v. s.)
 - β . *Provincialis*, petalis flammeis. in Gallo-provinciâ inter Digne et Colmars. (v. v.)

Sect. II. CONSILIGO. DC. syst. 1. p. 224.

Cariopsides stylo uncinato recurvo uncinatæ; petala 8-15; radices perennes.

11. *A. VERNALIS* (Lin! spec. 771.) foliis radicalibus infimisque abortivis ad squamas subvaginantibus reductis, mediis et summis sessilibus multifidis, lobulis integerrimis, carpellis velutinis, petalis 10-12 oblongis subdenticulatis. ② in Sibiria, Tauria, Oelandia, Germania, Gallia, Helvetia, Italia. DC! syst. 1. p. 224. quoad var. confer Wallr. sched. p. 277. Gaert de fruct. 1. p. 355. f. 74. *A. Apennina* Jacq. fl. aust. t. 44. (v. v.)

β. *Mentzelii*. *Helleborus niger ferulaceus*, etc. *Mentz.* pag. 1. 3. f. 1. *A.*

Apennina *Lin!* spec. 772. *A. vernalis*, β. *Lam.* dict. 1. p. 45.

γ. *Sibirica* (*Patrin.*) *A.* n. 43. *Gmel. fl. Sib.* 4. p. 200. non differt à nostrâ nisi flore majore. *Circâ Barnaoul.* (v. s.)

12. *A. Volænzis* (*Steven!* ined. *DC. syst.* 1. p. 545.) foliis radicalibus infimivæ abortivis ad squammas subvaginantes reductis, mediis et summis sessilibus multifidis, lobulis margine dentatis, carpellis subvelutinis, calyce extus pubescente, petalis 10-12 oblongis. ♀ ad flumen Volgam et Rhynum. *Deless!* ic. sel. 1. t. 20. *A. chærophylla* *Fisch!* ined. *Media inter A. vernaalem et Pyrenaicam.* (v. s.)

13. *A. IRCUTIANA*, foliis radicalibus. . . . mediis summivæ sessilibus pinnatim multisectis, lobulis dentatis, calyce glaberrimo, petalis obovatis. ♀ circâ *Irkoutsk.* *Fisch!* in litt. *Species non satis nota.* (v. s.)

14. *A. PYRENAICA* (*DC. fl. fr.* 5. p. 635.*) foliis radicalibus longè petiolatis, petiolo trifido, segmentis multipartitis, summis sessilibus multifidis, lobulis linearibus integerrimis, carpellis glabris, petalis 8-10 oblongo cuneatis integris. ♀ in *Valle Eynes Pyrenæorum orientalium* et prope *Edinburg* in *Hungariâ.* *Deless!* ic. sel. 1. t. 21. *Species priori affinis sed certissimè distincta.* *DC. syst.* 1. p. 226. exclus. synonymis *Gmel. Pallas. et Fischerii.* (v. v.)

β. *minor.* (*A. Apennina* *Schouw!* in litt. non *Lin.*) in *Apennini australis monte Velino.*

X. *HAMADRYAS.* *Commers. in Juss. gen.* 232. *DC. syst.* 1. p. 226.

Flores abortu dioici. Calyx 5-6 sepalus. Petala 10-12 linearia longa. ♂ *stam.* ∞ breviora. ♀ *ovaria* ∞ congesta; *stigmata* sessilia. *Carpella* 1-sperma ovata. — *Fructus non satis notus et ideò locus generis in ordine incertus.*

1. *H. MAGELLANICA* (*Lam!* dict. 3. p. 67.*) calyce glabro, foliis superne glabris 5-partitis, lobis inciso-partitis. ♀ in *memoribus montosis ad fretum Magellanicum.* *Deless!* ic. sel. 1. t. 22. *Scapus foliis paulò longior hirsutocanus.* *DC. syst.* 1. p. 226. (v. s.)

2. *H. TOMENTOSA* (*DC. syst.* 1. p. 227.) calyce tomentoso, foliis superne arachnoideo-villosis 5-7-fidis, lobis oblongis aut subtrifidis. — In *declivibus montium Americæ australioris in Statenland.* *Herba erecta, tota albedo tomentosa.* *Semen ovatum.* (v. s. ♀ in *h. Lamb.*)

Trib. III. *RANUNCULÆ.* *DC. syst.* 1. p. 228.

Æstivatio calycis et corollæ imbricata. Petala bilabiata aut intus squamulâ ad basin aucta. *Carpella* 1-sperma sicca indehiscentia. *Semen erectum. Folia radicalia aut alterna.*

XI. *MYOSURUS.* *Dill. giess. p.* 106. t. 4. *Lin. gen. n.* 394. *DC. syst.* 1. p. 229.

Calyx 5-sepalus, sepalis basi solutis seu ultrâ insertionem deorsum longè productis. Petala 5 ungue filiformi tubuloso. *Stamina* 5-20. *Ovaria* et dein *cariopsides plurimæ triquetræ confertissimæ, receptaculo longissimo spicatim dispositæ, stylo recto subacuminatæ.*

1. *M. MINIMUS* (*Lin!* spec. 407.) scapis folio æqualibus vel longioribus calycis appendicibus subfoliaceis. ☉ in *apricis pinguibus Europæ.* — *Fl. dan.* t. 406. — *Ranunculus Myosurus Afz. in Liljebl. sv. fl. p.* 230.

2. *M. SHORTII* (*Raf. in Sillim. am. journ. sc.* 1819. v. 1. n. 4. p. 379.) scapis folio brevioribus, calycis appendicibus membranaceis, ☉? circâ *Hopkinsville in Kentucky occid.* — *An prior. var.?*

XII. CERATOCEPHALUS. *Manch. meth. p. 218. DC. syst. 1. p. 230.*

Calyx 5-sepalus subpersistens, sepalis basi non productis; Petala unguiculata; Stamina 5-15. Carpella ∞ secus rachin persistentem longè spicata, basi utrinquè gibba, apice producta in cornu compressum semine sextuplò longius; Semen 4-gonum. Embryo erectus.

1. *C. FALCATUS* (Pers. ench. 1. p. 341.) pericarpiorum cornubus falcatis subascendentibus \odot inter segetes totius regionis Mediterraneæ. DC. syst. 1. p. 230. Ranunculus falcatus. Lin! spec. 781. Jacq. fl. aust. t. 48. (v. v.)
2. *C. ORTHOCERAS* (DC. syst. 1. p. 231.) pericarpiorum cornubus rectis. \odot frequens in campis incultis Tauriæ et Sibiria australis. Deless! ic. sel. 1. t. 23. Ranunculus falcatus Bieb. fl. cauc. 2. p. 29. exclus. syn. Certò à *C. falcato* differt, nec in sexcentis speciminibus ullum transitum ad hunc demonstrat (Fisch.). (v. s.)

XIII. RANUNCULUS. *C. Bauh. pin. 180. Pers. ench. 2. p. 102.*

Calyx 5-sepalus, sepalis basi non solutis deciduis. Petala 5, rarius 10, intus basi squamulâ foveolari nectariferâ instructa. Stam. ovariaque ∞ ; cariopsides ovatae subcompressæ, in mucronem aut cornu semine vix longius desinentes, læves striatæ aut iuberculatæ, in capitulum globosum cylindraceumve dispositæ. (DC. syst. 1. p. 231.)

Sect. I. BATRACHIUM. *DC. syst. 1. p. 233.*

Pericarpia transversè rugoso-striata. Petala alba ungue flavo foveâ nectariferâ notato.

1. *R. HEDERACEUS* (Lin! spec. 781.) caule repente, foliis subreniformibus sub 3-5-lobis, lobis latis integris obtusissimis, petalis oblongis calyce vix longioribus, staminibus 5-12, carpellis glabris. \mathcal{L} ad scaturigines et fontes in aquis vadosislocisque inundatis Europæ occident. Engl. bot. t. 2003. Schlecht. anim. Ran. 1. p. 7. DC. syst. 1. p. 233. (v. v.)

2. *R. TRIPARTITUS* (DC! ic. gall. rar. 1. p. 15. t. 49.) caule natante, foliis submersis capillaceo-multifidis, emersis 3-partitis, lobis cuneatis apice dentatis, carpellis glabris. \mathcal{L} in fossis scaturiginibus inundatisque Gallia occidentalis. Species omninò media inter *R. hederaceum* et *aquatilem*. DC. syst. 1. p. 233. (v. v.)

a. micranthus, petalis oblongis acutiusculis calycis longitudine. *R. tripartitus* DC. ic. gall. rar. 1. p. 15. t. 49. et syst. l. c.

β . obtusiflorus, petalis obovatis obtusis calyce longioribus. — Petiv. engl. herb. t. 59. f. 1.

3. *R. AQUATILIS* (Lin. sp. 781.) caule natante, foliis submersis capillaceo-multifidis, emersis 3-partitis, lobis cuneiformibus apice dentatis, petalis obovatis calyce majoribus. \mathcal{L} in Europâ, Asiâ occidentali, Africâ et Americâ.

a. heterophyllus, foliis emersis 5-partitis. *R. heterophyllus* Hoffm. fl. germ. 197. *R. fluviatilis* Tabern. ic. 54. f. 2. Variat foliis emersis peltatis aut sæpius non peltatis, carpellis setis rigidis hispidis aut rarius glabris ut me docuit cl. Wallroth (Sched. p. 282.). Huc etiam forsan spectat prioris var. *β* ? (v. v.)

β . capillaceus, foliis petiolatis, omnibus immersis in lacinias filiformes tenues dissectis. — Barr. ic. 57. t. 566. *R. trichophyllus*. Chaix in Vill. Dauph. 1. p. 335. Variat flore majore et minore, carpellis hispidis aut rarius glabris. *R. Pantothrix* var. *a*. DC. syst. 1. p. 235. (v. v.)

γ . cæspitosus, foliis petiolatis, omnibus emersis, circumscriptione suborbiculatis in lacinias divergentes rigidulas dissectis, petiolis basi latè vagiuato-auriculatis. *R. aquatilis* var. *B*. Schk. handb. 2. n. 1527. t. 152. *R. rigidus* Pers. in Hoffm. fl. germ. 4. p. 257. (v. v.)

3. *stagnatilis*, foliis sessilibus omnibus immersis capillaceo-multifidis circinnatis, laciniis abbreviatis, vaginis obsolete anniculatis, carpellis acutiusculis glabriusculis. R. *stagnatilis* Wallr! sched. 285.

4. *peucedanifolius*, foliis petiolatis omnibus immersis in lacinias longas parallelas dissectis. R. *fluitans*, OEd. fl. dan. t. 376. R. *peucedanifolius* All! ped. n° 1469. (v. v.)

Sect. II. RANUNCULASTRUM.

Carpella lævia compressa in spicam digesta. Radix grumosa.

4. R. *BULLATUS* (Lin! spec. 774.) foliis omnibus radicalibus petiolatis ovatis dentatis scapis nudis unifloris. 2/ in arvis incultis olivetiisque Europæ australioris. Flores lutei odorati. DC. syst. 1. p. 253. (v. s.)

α. *parviflorus*, flore parvo, scapo patentim piloso, foliis parum bullatis. — Clus. hist. 1. p. 238. f. 2.

β. *grandiflorus*, flore magno, scapo adpressè piloso, foliis magis bullatis. — Clus. hist. 1. p. 238. f. 1.

5. R. *CHÆROPHYLLUS*. (Lin! spec. 780.) foliis radicalibus petiolatis subvillosis 3-sectis, segmentis semel aut bis tripartitis acutis, primordialibus ovatis dentatis 3-lobisve, caule erecto 1-2-floro, calyce patente subreflexo. 2/ in collibus totius Europæ meridionalis, Africæ borealis et Archipelagi. Carpella lævia compressa stylo acuminata, in capitulum ovato-oblongum dense conferta. DC. syst. 1. p. 254. (v. v.)

α. *vulgaris*, foliis ferè omnibus 3-sectis multifidis. R. *chærophyllus*. Lin! l. c. — Barr. ic. t. 581.

β. *gregarius*, foliis aliis 3 fidis incisis, aliis 3 sectis multifidis. R. *gregarius* Brot. fl. lus. 2. p. 369. * R. *Thomasii* Ten. prod. fl. neap. supp. 2. p. 68. an potius ad millefoliatum, ex Schlecht. anim. 2. p. 37?

γ. *flabellatus*, foliis aliis ovalibus dentatis, aliis 3-sectis multifidis. R. *flabellatus* Desf! fl. atl. 1. p. 438. t. 114.

6. R. *AGERII*. (Bertol. opusc. sc. 3. 1819. p. 182. t. 6.*) foliis glabris, primordialibus reniformibus 3-lobis crenatis succulentis 3-partitis ultimisque 3-sectis, lobis multifido-linearibus, caule bifolio hirsuto, carpellis in acumen subulatum desinentibus. 2/ Bononiæ. (v. s.)

7. R. *PALUDOSUS* (Poir! voy. 2. p. 184. dict. 6. p. 108.) foliis pubescentibus, radicalibus 3-sectis, lobis multifidis, superis linearibus integerrimis, caule erecto dichotomo, calyce adpresso. 2/ in Barbariâ. R. *chærophyllus* var. D. Biria ren. 42. Corolla R. acris; carpella compressa submutica lævia in capitulum ellipticum aggregata. DC. syst. 1. p. 255. (v. s.)

8. R. *ADSCENDENS*. (Brot. fl. lus. 2. p. 370. *) foliis lanuginosis, radicalibus 3-partitis, laciniis cuneatis 3-fidis inciso-dentatis, caulibus ascendentibus dichotomis, pedunculis sulcatis angulosis, calyce reflexo. 2/ in Lusitaniâ. Variat staturâ maximâ, caulibus ramosioribus, foliis magnis ad sinus sæpè maculatis. DC. syst. 1. p. 256.

9. R. *MILLEFOLIATUS* (Vahl! symb. 2. p. 63. t. 37. *) foliis decompositis multifidis, lobis linearibus glabris, caule subaphyllo erecto adpressè villosa 1-floro, calyce adpresso. 2/ in cultis regni Tunetani, ad montes Sibiæ et Peloponesi, in Siciliâ. Affinis R. *chærophyllus*, sed differt radicem tuberculis brevioribus crassioribusque, etc. DC. syst. 1. p. 256. (v. s.)

10. R. *GRACILIS* (DC. syst. 1. p. 256. *) foliis glaberrimis, aliis orbiculatis 3-lobis apice dentatis, aliis 3-sectis lobis oblongis inciso-lobatis, caule subaphyllo erecto adpressè villosa 1-floro, calyce reflexo subpiloso. 2/ in Archipelagi insulâ Cois. R. nov. spec. Clark. trav. 2. p. 723. R. *gracilis* Spreng. neu. cutd. 3. p. 164. sepalâ ovalia reflexa extus subpilosa, petalâ magnitudine R. *acris*. (v. s. in h. Lamb.)

11. R. *FUMARIÆFOLIUS* (Desf. in ic. pict. h. par. 37. t. 74.) foliis glaberrimis pinnatisectis multipartitis, lobis oblongis, scapis plurimis 1-floris adpressè

villosis, calyce patente glabro. \mathcal{L} olim in hortis cultus. R. rutæfolius Mill. dict. n. 6. non Lin. Species affinis hinc R. gracili et millefoliato, illinc R. asiatico.

12. R. MYRIOPHYLLUS (Russel! gew. alep. in Schrad. journ. 1799, p. 424.) foliis decompositis ∞ -fidis villosis-canescens, lacinii linearibus, caule ramoso adpressè villosis, calyce patente villosis. \mathcal{L} in Aleppo. Deless! ic. sel. 1. t. 31. Herba tota villis albidis canescens; DC. syst. 1. p. 257. (v. s. in h. Banks, Olivier, Labill.)

13. R. ORIENTALIS (Lin! spec. 781.) foliis decomposito-multifidis pubescentibus, caule ramoso adpressè villosis, calyce reflexo, carpellis longè cornutis. \mathcal{L} in Oriente, in insulâ Lesb. Deless! ic. sel. 1. t. 32. DC. syst. 1. p. 257.* (v. s. in h. Tourn. et Lin.)

β . R. *orientalis* pulsatillæ folio parvo flore Tourn! cor. 20. Smyrnæ. — vix a priori differre videtur nisi petalis angustioribus.

14. R. LEPTALEUS (DC. syst. 1. p. 258.) foliis decomposito-multifidis glabris, caule patulè piloso, calyce patente, carpellis longè cornutis. \mathcal{L} in insulâ Cypro. Deless! ic. sel. 1. p. 33. Carpella ut prioris apice uncinata, Cerato cephalos ferè adæquantia. (v. s.)

15. R. CICUTARIUS (Schlecht. anim. ran. 25. t. 4. f. 2.*) foliis pinnati-sectis et pinnati-partitis, segmentis cuneatis incisive dentatisque glabris, caule subramoso adpressè villosis, calyce patente adpresso-pubescente. \mathcal{L} in Sibiria orientali.

16. R. OXYSPERMUS (Willd. spec. 2. p. 1328.*) foliis villosis, radicalibus petiolatis ovatis 3-partito-incisive dentatis, floralibus 3-partitis, lobis elongatis apice dentatis, caule erecto patulè villosis dichotomo, calyce reflexo, carpellis muricatis aristatis. \mathcal{L} in Tauriæ et Caucasi etiam Iberiæ campestribus. an sectio propria? DC. syst. 1. p. 258.* (v. s.)

17. R. PEDATUS (Waldst et Kit. plant. rar. hung. 2. p. 112. t. 108.*) foliis glabris, radicalibus petiolatis 3-partitis pedatisve, lobis linearibus integris bifidisve, caulinis sessilibus partitis, summis linearibus, caule erecto paucifloro calyce adpresso. \mathcal{L} in Hungaria prope Budam, Tataria, propè Volgam flumen. Lodd. bot. cab. 4. p. 351. Variat caule pubescente aut glabro aut basi glabro apice pubescente, 1-5 floro. DC. syst. 1. p. 258.* (v. v.)

18. R. PETROSELINUS (Biria! ren. 43. t. 2.*) foliis glabris ad basim petioli villosis, 3-sectis, segmentis 3-partitis, lobis subtrilobatis inciso-dentatis, caule erecto ∞ -floro. \mathcal{L} in insulâ Borboniæ. Carpella ovata compressa lævia, stylo brevissimo apiculata, in capitulum ovato-oblongum aggregata. DC. syst. 1. p. 259.* (v. s.)

19. R. ILLYRICUS (Lin! spec. 776.) foliis sericeo-lanuginosis, primordialibus integris lineari-lanceolatis, ceteris 3-partitis, lobis integris 3-partitisve linearibus caule erecto ∞ -floro, calyce subreflexo. \mathcal{L} in pascuis Illyriæ, Anstræ, in Tauriâ et circa Odessam. Jacq. fl. aust. t. 222. R. sericeus Willd! enum. 589. Carpella glabra lævia compressa acuminata in capitulum ovato-oblongum aggregata. DC. syst. 1. p. 259. (v. s.)

20. R. MONSPELIACUS (Lin. spec. 778.) foliis lanuginosis, radicalibus 3-lobis, lobis cuneatis 3-fido-dentatis, superioribus 3-partitis, lobis integris linearibus, caule erecto paucifloro, calyce reflexo, carpellorum spica ovata. \mathcal{L} in pascuis apicis saxatilibusve regionis Mediterraneæ. DC. syst. 1. p. 260.* (v. v.)

α . *angustilobus*, lobis foliorum angustis elongatis. — in apricis. R. Illyricus. Besl. syst. vern. ord. 1. t. 13. f. 1.

β . *cuneatus*, lobis foliorum cuneiformibus apice 3-fido-dentatis. — in herbosis. R. monspeliacus DC. ic. gall. rar. t. 50. R. apiifolius Desf. in pict. hort. par. 37. t. 73. non Pers.

γ . *rotundifolius*, foliis rotundatis 3-fidis, lobis dentatis obtusis. — in saxosis. R. monspeliacus. Gou! fl. monsp. 279. Poir! dict. 6. p. 111.

21. *R. SPICATUS* (Desf! fl. atl. 1. p. 438. t. 115.) foliis subhirsutis, radicalibus petiolatis orbiculatis 3-lobis dentatis, summis 3-partitis lobis integris linearibus, caule erecto paucifloro, calyce reflexo, carpellorum spicâ cylindricâ. DC. syst. 1. p. 260. (v. s.)
 - a. Foliis trilobis dentatis, caule 1-2-floro. *R. spicatus*. Desf! l. c. Bir! ren. 42. in paludibus Algeriæ.
 - β. Foliis crenatis, caule 1-2-floro. *R. Lusitanicus* grumosa radice, etc. Tourn! inst. 286. *R. Olyssiponensis* Pers! ench. 2. p. 106. circa Olyssiponam (v. s. h. Juss.)
 - γ. Foliis 3-lobis crenatis, caule 2-5-floro. *R. asphodeli* radice, etc. Tourn! herb. ad Carthusiam Laverno.
22. *R. ASIATICUS* (Lin! spec. 777.) foliis ternatim biternatimve sectis, segmentis dentatis aut inciso-3-fidis, caule erecto simplici aut infernè ramoso, calyce patente demùm reflexo, carpellorum spicâ cylindricâ. DC. syst. 1. p. 261. (v. v.)
 - a. *vulgaris*, caule infernè ramoso, foliis 3-sectis, segmentis 3-fidis, incisius acutis — in Oriente. Clus. hist. 1. p. 241. f. 2. *R. asiaticus* Mill. ic. t. 216. dict. n. 11.
 - β. *sanguineus*, caule simplici, foliis 3-sectis, segmentis dentatis obtusis in Archipelago, Cariâ, Ciliciâ, Syriâ. Clus. hist. 1. p. 242. ic. *R. sanguineus* Mill. dict. n. 10.
 - γ. *tenuilobus*, caule subramoso, foliis multifidis, lobis linearibus acutis. In insulâ Cypri. *R. creticus* albo flore, etc. J. Bauh. hist. 3. p. 862. f. 2. (ex h. Vaill.)
23. *R. JAPONICUS* (Thunb. act. soc. Lin. 2. p. 337.*) foliis hirsutis, radicalibus rotundatis 3 fidis, lobis incisius dentatis, supremis fissis, caule erecto hirsuto apice ramoso. ʒ in Japoniâ. — *R. asiaticus* Thunb. fl. jap. 241. Radix fructusque ignoti. DC. syst. 1. p. 263.
24. *R. KRAPPÆ* (DC. in Deless. ic. sel. 1. t. 35.) villosus, foliis radicalibus longè petiolatis ovatis subtrifidis dentatis, caulinis parvis basi attenuatis apice 3 fidis, caule 1-floro. ʒ in Chili. *Krapfia ranunculina*. DC. syst. 1. p. 228*. Petala obovata retusa calyce æqualia. (v. s. in h. Mus. Par.)
25. *R. GUZMANNI* (Humb. tabl. reg. eq. 69. nov. gen. 5. p. 43.*) foliis superne petiolisque villosis, radicalibus suborbiculatis 3-fidis, lobis apice grossè dentatis, superioribus 3-fidis lobis oblongis integris, caule erecto piloso paucifloro, calyce adpresso villosissimo. ʒ in summis Andium cacuminibus saxosis. Deless! ic. sel. 1. t. 34. DC. syst. 1. p. 263. (v. s. in h. Juss.)
26. *R. MACROPETALUS*, petiolis basi membrana auctis, foliis radicalibus orbiculatis crenatis glabris pilosisve, caulinis ovato-lanceolatis sessilibus, caule glabro paucifloro, calyce adpresso petalis multò minore. ʒ in Peruvix montibus. *R. cochlearifolius* Ruiz et Pav! in h. Lamb. non Horn. (v. s.)
27. *R. CRETICUS* (Lin! spec. 775.) foliis cauleque molliter subhirsutis, radicalibus petiolatis cordatis orbiculatis subincisius dentatis, caulinis sessilibus 3-partitis, lobis oblongis obtusis apice subdentatis, caule ramoso oo-floro, calyce adpresso. ʒ in Cretâ. *R. Creticus latifolius* Clus. hist. 1. p. 239. f. 1. DC. syst. 1. p. 264. (v. v. et s.)
 - β. *macrophyllus* (Desf! fl. atl. 1. p. 437.) foliis minus villosis majoribus, profundius lobatis, dentibus paulò rotundioribus — in Barbariâ.
28. *R. CORTUSÆFOLIUS* (Willd. enum. 588.) foliis cauleque subpilosis, radicalibus subcordato reniformibus sublobatis latè crenatis, caulinis subsessilibus 3-partitis, floralibus lanceolatis, caule ramoso corymboso, calyce patentissimo. ʒ in rupibus insulæ Teneriffæ. Deless! ic. sel. 1. t. 36. DC. syst. 1. p. 264. (v. s.)
 - β. *Teneriffæ* (Pers! ench. 2. p. 103.) *R. creticus*. var. b. Bir! ren. 45. folia radicalia reniformia basi subtruncata vix cordata, caulinum in lobos integros fissum. (v. s.)

Sect. III. THORA.

Carpella lævia subglobosa. Radices grumosæ.

29. R. THORA (Lin! spec. 775.) foliis glaberrimis, radicalibus petiolatis caulinoque sessili reniformibus crenatis, floralibus incisissimis, caule 2-3-floro glabro. \mathcal{L} in pascuis Jurassi, Alpium, Apenninorum, Carpathorum. Jacq. fl. austr. 5. t. 442. DC. syst. 1. p. 264. (v. v.)
30. R. SCUTATUS (Waldst. et Kit. pl. rar. hung. 2. p. 205. t. 187. *) foliis glaberrimis, radicalibus nullis, caulino sessili reniformi crenato, floralibus incisissimis caule 2-3-floro basi subpiloso. \mathcal{L} in Hungariâ. R. Thora var. scutatus Wahl. fl. carp. n. 551. * simillimus R. Thoræ et forsitan cum eo à veteribus nonnullis non immerito conjunctus. DC. syst. 1. p. 265.
31. R. BREVI-FOLIUS (Ten. fl. neap. prod. supp. 2. p. 68.) foliis glaberrimis, radicalibus petiolatis apice 3-lobis, lobis lateralibus subcrenatis, intermedio usque ad folii tertiam partem producto, caulino sessili cuneiformi 3-partito, caule 1-floro incurvo. \mathcal{L} in saxosis Magellæ in regno Neapolitano. Aff. R. Thoræ. Radix fructusque ignoti. DC. syst. 1. p. 266. (v. s.)
32. R. HYBRIDUS (Bir! ren. 38.) foliis glaberrimis, radicalibus longè petiolatis reniformibus lobato-crenatis, caulinis 2 breviter petiolatis apice lobatis, caule 1-3-floro glabro. \mathcal{L} in Alpibus Austriæ. R. Thora. Sturm. deutsch. fl. ic. R. Thora var. petiolatus Wahl. fl. carp. 170. Aff. R. Thoræ ob habitum totum et R. auricomum ob radicem. DC. syst. 1. p. 266. (v. s.)

Sect. IV. HECATONIA.

Carpella lævia ovato-subrotunda in capitulum subrotundum digesta. Radix fibrosa.

§. I. Floribus albis, foliis dissectis.

33. R. BITERNATUS (Smith. in Rees cycl. n. 48.) caule repente, foliis 3-partitis, partitionibus 3-lobis, lobis oblongis acutis integris subdentatisve, petalis oblongis calyci æqualibus. \mathcal{L} ? in fretum Magellanicum. Schlecht. anim. ran. 1. p. 11. Deless! ic. sel. 1. t. 24. Herba pusilla multicaulis tenera glaberrima. DC. syst. 1. p. 236. (v. s.)
34. R. RUTÆ-FOLIUS (Lin! spec. 777.) foliis pinnatim sectis, lobis trilobato-incisissimis multifidis, caule sub-1-floro, calyce glabro, petalis 8-10, ungue discolori. \mathcal{L} inter scopulos summorum Alpium prope nives deliquescentes. Wulf. in Jacq. coll. 1. p. 186. t. 6 et 7* folia juniora lobis intus (ut in Thalictris) replicatis. DC. syst. 1. p. 237. (v. s.)
35. R. ISOPYROIDES (DC. syst. 1. p. 238.) foliis radicali pinnati-secto, segmentis petiolulatis 2-3-fidis, caulinis 3-sectis, calyce glabro, petalis 5 ovalibus. \mathcal{L} in Sibiria. Schlecht. anim. ran. 1. p. 11. R. fumarifolius h. Gorenk. ex Fisch. litt. Nimis affinis R. rutæfolio; petala interdum ultra 5 sed folii limbus potius in R. isopyroide ellipticus et in R. rutæfolio ovatus. (v. s.)
36. R. GLACIALIS (Lin! spec. 777.) foliis radicalibus petiolatis palmatim 3-partitis 3-sectisve, lobulis subobtusis crassiusculis, caule subunifloro, calyce hirsutissimo, carpellis compressis marginatis. \mathcal{L} inter scopulos ad nives perpetuas in Alpibus summis. Wulf. in Jacq. coll. 1. p. 189*. t. 8, 9. f. 1. 2. (v. v.)
 \mathcal{L} ? aconitoides, foliorum segmentis cuneatis apice acutè incisissimis. R. ex glaciali et aconitifolio hybridus. Thomas! exsic. — in M. Sti. Bernardi inter duas species citatas.
37. R. CHAMISSONIS (Schlecht. anim. ran. 1. p. 12. t. 1*) foliis radicalibus petiolatis 3-partitis, partitionibus cuneatis 3 lobis, lobis integris obtusiusculis, intermedio majore, caule 1-floro, calyce hirsutissimo, carpellis ovalibus basi curvatis utriculosis acuminatis. — ad pedem montium glacialium in sinu Sti. Laurenti.
38. R. SGUIERI. (Vill! Dauph. 4. p. 737. t. 49.) foliis 3-partitis, partitionibus acutis 3-fidis aut integris, caule subunifloro, calyce glabro, petalis 5 integris.

¶ inter scopulos et fissuras rupium propè nives perpetuas in Alpibus. R. Columnæ All! pedem. 1453. t. 67. fig. 3. 4. Aff. R. glaciali et R. alpestri. DC. syst. 1. p. 238. (v. v.)

39. R. ALPESTRIS (Lin! spec. 778.) foliis orbiculatis 3-lobis, lobis apice lobulato-crenatis obtusiusculis, caule subunifloro, calyce glabro, petalis 5 obcordatis 3-lobatis. ¶ inter scopulos et in pascuis editioribus Carpathorum, Alpium. etc. — Seg. ver. 1. p. 489. t. 12. f. 1. DC. syst. 1. p. 239. (v. v.)

β. *Magellensis* Ten. cat. h. app. 1. p. 53.

γ. *Traunfellneri* Hopp. Jahrg. 1819. 731. ex ic. Schlechtend. anim. 2. p. 6.

40. R. CRENATUS (Waldst. et Kit. pl. rar. hung. 119. t. 10*) foliis cordato-subrotundis apice crenatis, caule 1-floro, calice glabro, petalis 5 obovatis apice crenatis. ¶ in vallibus Hungariæ. Habitus ferè R. alpestris. Petala basi poro mellifero nec squamulâ instructa. DC. syst. 1. p. 240.

41. R. ACONITIFOLIUS (Lin! mant. 79.) foliis palmatis 3-7-partitis, partitionibus inciso-dentatis, supremis sessilibus in lobos lineari-lanceolatos fissis, caule ramoso ∞-floro, calycibus adpressis glabris. ¶ in montibus Europæ mediæ. DC. syst. 1. p. 240. (v. v.)

α. *humilis*, foliis radicalibus 3-5-partitis, bracteis lanceolatis serratis, caule gracili glabro. R. aconitifolius. Lin. l. c. Moris. hist. 3 p. 462. sect. 12. t. 2. f. 5.

β. *crassicaulis*, foliis radicalibus 3-partitis, bracteis lanceolatis serratis, caule crassiusculo apice adpressè villosa. R. heterophyllus. Lapeyr. abr. 316 non Smith.

γ. *intermedius*, elatior, foliis radicalibus 3-partitis, bracteis linearibus integris.

δ. *platanifolius*, elatior, foliis radicalibus 5-7-lobis, lobis acuminatis, bracteis linearibus integris. R. platanifolius. Lin! mant. 79. fl. dan. t. III. — * flore pleno. R. aconitifolius β. Curt. bot. mag. t. 204.

ε. *dealbatus*, foliis radicalibus 7-lobis dentatis subtus candicantibus, supremis 3-lobis. R. dealbatus Lapeyr. abr. 315.

42. R. TERNATUS (Thunb. fl. jap. 241.) foliis omnibus 3-sectis, segmentis ovatis 3-fidis dentatis, caule pilosiusculo ∞-floro, calycibus reflexis. ¶ in insulâ Nipon Japoniæ. Deless! ic. sel. 1. t. 25. Affinis R. aconitifolio et præsertim var. β. DC. syst. 1. p. 242. (v. s. in h. Deless.)

43. R. APHIFOLIUS (Pers. each. 2. p. 105. non Desf.) foliis 5-lobis, lobis sinuato-serratis, supremis sublinearibus, caule glabro ∞-floro, calycibus patentibus, carpellis muticis. ¶ in paludosis Bonariæ. Deless! ic. sel. 1. t. 26. Affinis R. scelerato sed flores ex albo incarnati, carpella minima in capitulum ovatum aggregata. DC. syst. 1. p. 242. (v. s. in h. Juss. et Deless.)

44. R. LACERUS. (Bell! act. taurin. 5. 233. t. 8*) foliis cuneiformibus apice irregulariter acutè incisis, caule glabro ramoso ∞-floro, calycibus adpressis. ¶ Rarissimus in pascuis fertilibus Alpium Vallesiar, etc. R. Vallesiacus Sut. fl. helv. 1. p. 325. Hybridus omninò videtur ex R. aconitifolio et R. Pyrenæo. Flores albi. DC. syst. 1. p. 242. (v. v.)

β? *pallidus*. R. frigidus Schrank. hort. monac. t. 57*. non Wild. — Differt à vero R. lacero floribus pallidè flavis nec albis. Hybridus R. graminei et R. aconitifolii ex cl. Stoffels, in Ann. gen. sc. ph. 8. p. 352. t. 129. (v. v.)

§. 2. *Floribus albis, foliis integris.*

45. R. PYRÆNEUS. (Lin! mant. 248.) foliis linearibus lanceolatisve integerrimis, scapo pedunculisve apice tomentosis. ¶ in Alpium et Pyrenæorum pascuis. Deless! ic. sel. 1. t. 27. f. B. DC. syst. 1. p. 245. (v. v.)

β. *buplevriifolius*, caule 1-floro, foliis lanceolatis. R. Pyrenæus Jacq. misc. 1. p. 154. t. 18. f. 1. R. buplevriifolius Lapeyr. abr. 314.

γ. *plantagineus*, caule ∞-floro, foliis lanceolatis. R. plantagineus All. ped. n. 1445. t. 76. f. 1. non Pers.

46. *R. ANGUSTIFOLIUS* (DC. rapp. voy. 1. p. 74.) foliis linearibus sublan-
 ceolatis glaucescentibus, non amplexicaulis, scapo pedunculisque glabris. \mathcal{L} in
 Pyrenæis circa Mont-Loup. Deless! ic. sel. 1. t. 27. f. A. Omnino medius
 inter *R. Pyrenæum* et amplexicaulem. DC. syst. 1. p. 243. an huc referendus
R. alismoides Bory. ann. gen. sc. ph. 3. 1820. p. 13, in montibus Granaten-
 sibus lectus? (v. s.)
47. *R. AMPLEXICAULIS* (Lin! spec. 774.) foliis ovali-lanceolatis acuminatis am-
 plexicaulis, scapo pedunculisque glabris. \mathcal{L} in locis excelsis udis scopu-
 losis Alpium, Apenninorum, etc. Deless! ic. sel. 1. t. 27. f. c. DC. syst. 1. p.
 244. *R. amplexicaulis* var. α Goul! ill. 34. (v. v.)
48. *R. PARNASSIFOLIUS* (Lin! spec. 774.) foliis radicalibus petiolatis subcordatis
 ovato-subrotundis, caulinis sessilibus ovato-lanceolatis, pedunculis hirsutis.
 \mathcal{L} in Alpium Pyrenæorumque scopulis nivi æternæ contiguus. Wulf. in
 Jacq. coll. 1. p. 171. t. g. f. 3*. DC. syst. 1. p. 244. *R. cordatus*. guid. herb.
 Valais. 177. (v. v.)
- ß. parviflorus*, foliis acutiusculis floribus minoribus, — in Alpibus Vallesiæ.

§. 3. *Floribus flavis, foliis indivisis.*

49. *R. GRAMINEUS* (Lin! spec. ed. 1. p. 549. ed. 2. p. 773.) foliis lanceolatis linea-
 ribusve integerrimis, caule erecto glaberrimo ad collum fibrilloso, squamulâ
 petalorum tubulosâ, radice fasciculatâ. \mathcal{L} in pratis interdum inundatis cam-
 pisque subargillosis Galliæ, Vallesiæ et Lusitaniæ. DC. syst. 1. p. 245. (v. v.)
- α. linearis*, foliis linearibus. — Tab. ic. 51. f. 1. — Moris. hist. 2. p. 244.
 s. 4. t. 30. f. 38.
- * *ß. phœnicifolius*, foliis lanceolatis. — Moris. hist. 2. p. 445. s. 4. t. 30. f. 39.
γ. hortensis, foliis lineari lanceolatis, flore pleno. — Moris. hist. 2. p. 445.
50. *R. BUPLEVROIDES* (Brot. fl. lusit. 2. 365*) foliis radicalibus petiolatis ovato-
 lanceolatis 3-5-nerviis integerrimis, caule erecto inferuè lanuginoso, squa-
 mula petalorum tubulata, radice fasciculata. \mathcal{L} in Lusitaniæ collibus. — *R.*
plantagineus. Pers. ench. 2. p. 102. non All. — DC. syst. 1. p. 246.
51. *R. LINGUA* (Lin! spec. 773.) foliis lanceolatis subseratis sessilibus semi-
 amplexicaulis, caule erecto glabro. \mathcal{L} in palustribus inter arundines Eu-
 ropæ, Sibiriae et Americæ. — Weinm. phyt. t. 854. f. c. — *R. longifolius* Lam!
 fl. fr. 3. p. 189. Affinis flammulæ, sed statura major rigidior. DC. syst. 1. p. 246.
 (v. v.)
52. *R. FLAMMULA* (Lin! spec. 772.) foliis glabris linearibus lanceolatis subova-
 tisve subintegris, inferioribus petiolatis, caule declinato pleno basi subradi-
 cante, pedunculis oppositifoliis, carpellis lævibus. \mathcal{L} in Europæ totius, Asiæ,
 Africæ et Americæ locis udis. fl. dan. t. 575. engl. bot. t. 387. DC. syst. 1.
 p. 247. (v. v.)
- ß. serrata*, foliis sublan-
 ceolatis serratis. — Moris. hist. 2. p. 442. s. 4. t. 29.
 f. 35.
- γ. ovata*, foliis fere omnibus ovatis petiolatis. — *R. ovatus*. Pers. ench.
 2. p. 102.
- δ. arenaria*, foliis radicalibus ovato-lanceolatis rigidulis, caulinis lineari-
 bus. — in udis arenosis circa Baïonam. *R. flammuloides* Raf! in litt. ab
 hac var. vix differt. *R. lanceolatus* Pall! herb.
53. *R. REPTANS* (Lin! spec. 773.) foliis linearibus utrinque æqualibus integerri-
 mis glabris, vaginis carpellisque glaberrimis, caule repente ad quemque
 nodum radicales agente. \mathcal{L} in glareosis udis secus ripas lacuum Helvetiæ,
 Germaniæ, Angliæ, Sibiriae, etc. fl. lapp. 236. t. 3. f. 5. *R. flammula* var.
 Smith! fl. brit. 587. Biria! ren. 37. an prioris var?
- ß? filiformis*, foliis lineari-subulatis obtusis, caulibus repentibus geniculatis.
 — in Americâ boreali. *R. filiformis* Michx! fl. bor. Amer. 1. p. 320.
54. *R. FUSILLUS* (Poir! dict. 6. p. 99*) foliis omnibus longè petiolatis, infe-

- rioribus ovatis subdentatis, superioribus lineari-lanceolatis, caulibus plurimis erectis, pedicellis oppositifoliis, petalis calycis longitudine. \mathcal{L} in Carolinâ. Deless! ic. sel. 1. t. 28. *R. flammula* Walt. in Carol. 158. *R. humilis* Pers. ench. 2. p. 102. Species *R. flammulæ* affinis, sed. ex cl. Pursh. certò distincta. DC. syst. 1. p. 249. (v. s.)
55. *R. BONARIENSIS* (Poir! dict. 6. p. 102*) foliis omnibus petiolatis cordato-ovatis subcrenatis, pediculis oppositifoliis, petalis calyce vix longioribus, carpellis lævibus. (C) ? in Americæ locis humidis circa Buenos-Ayres. Deless! ic. sel. 1. t. 29. *R. heterophyllus* Smith! in Rees cycl. n. 12. Carpella in capitulum globosum aggregata. DC. syst. 1. p. 250. (v. s. in h. Mus. Par.)
56. *R. POLYPHYLLUS* (Willd. spec. 2. p. 1331.*) foliis caulinis numerosissimis in verticillos aggregatis, emersis petiolatis ovatis subtrilobis, immersis subfiliiformibus. (I) in Hungariæ turfosis humidis, ad Tanaim versus Tscherkawk. Planta glaberrima ad *R. flammulam* consistentiâ accedens. DC. syst. 1. p. 250. (v. s.)
57. *R. FLAGELLIFORMIS* (Smith! in Rees cycl. n. 13.*) foliis longè petiolatis orbiculatis cordatis integris glabris, caule repente, pediculis oppositifoliis, carpellis muticis sublævibus. \mathcal{L} in inundatis regni Chilensis et in Novâ Granadâ. *R. hydrocotyle* Domb! ined. herb. Carpella parva, ovata, mutica, lævia aut subpunctulata. DC. syst. 1. p. 251. (v. s. in h. Mus. Par.)
58. *R. SALSUGINOSUS* (Pall! itin. ed. 3. v. 3. p. 173.) foliis radicalibus petiolatis ovalibus subcordatisve apice 3-5-dentatis, scapis nudis 1-floris erectis, petalis oblongo-cuneatis calyce longioribus. \mathcal{L} in locis salsis udis imperii Ruthenici. *R. Ruthenicus* Jacq. hort. vind. 5. p. 19. t. 31.* Variat scapo erecto aut adscendente, foliis longiore aut brevior, (Fisch.) DC. syst. 1. p. 251. (v. s.)
59. *R. CYMBALARIÆ* (DC. syst. 1. p. 252.) foliis radicalibus petiolatis glabris subcarnosis ovato-orbiculatis grossè crenatis, flagellis è collo ortis, scapis nudis 1-3-floris ascendentibus, petalis linearibus calycis longitudine. \mathcal{L} in palustribus salsis Sibiricæ. — Amm. ruth. 81. n. 107. t. 13. f. 2. *R. nanus* Fisch! in litt. — DC. syst. 1. p. 252. (v. s.)
- β. Americanus*, *R. cymbalaricæ* Pursh! fl. bor. am. 2. p. 392. Smith. in Rees cycl. n. 14. in Noveboraco circa Onondago ex. Pursh.
60. *R. TRIDENTATUS* (H. B. et K! nov. gen. et spec. amer. 5. p. 42.*) foliis radicalibus petiolatis glabris-subcarnosis ovatis apice 3-dentatis, scapis 1-2-floris glabris petioli longitudine, petalis linearibus calyce longioribus. \mathcal{L} in Americâ. Planta parva, glabra, carnosa, valde similis hinc *R. cymbalaricæ* illinc *R. nubigeno*. DC! syst. 1. p. 252. (v. s. in h. H. et B.)
- α. major*. juxta Carpio in Mexico.
- β. minor*. in Latacungâ.
61. *R. NUBIGENUS* (H. B. et K! nov. gen. et spec. amer. 5. p. 42.*) foliis radicalibus petiolatis suborbiculatis 7-dentatis glabris subcarnosis, scapo subnudo subbifloro apice pubescente, bracteis linearibus subtridentatis. \mathcal{L} in monte Antisanâ Peruvicæ. Herba pygmæa, flores minimi lutei, calyx adpressus. DC. syst. 1. p. 253. (v. s. in h. H. et B.)
62. *R. BREVICAPUS* (DC. syst. 1. p. 253.) foliis radicalibus petiolatis orbiculato-cordatis 3-5 fidis, flagellis è collo ortis, scapis 1-floris petiolo dimidio brevioribus. \mathcal{L} in Peruvâ. Herba (quæ paludosa videtur) animalibus infesta et à Dombeyo Centella dicta. (v. s. in h. Mus. Par.)
- §. 4. *Floribus flavis, foliis dissectis.*
63. *R. AURICOMUS* (Lin! spec. 775.) foliis glabris, radicalibus petiolatis cordatis plerisque 3-partitis lobatisve, caulinis in lobos lineares integros subdentato-ve partitis, calyce pubescente petalis brevior. \mathcal{L} in Europâ, Sibirîâ, Japoniâ et Pensylvaniâ. fl. dan. t. 665. DC. syst. 1. p. 266. *R. polymorphus* All. pedem. n. 1449. t. 82. f. 2. (v. v.)

ℓ. *procerior*. in Hungariâ. R. Cassubicus Gener. elench. scepus. ex Wahlenb.
 γ. *apetalus*, petalis nullis, calyce colorato corollino. Wallr! in Mitt. in Thuringiâ.

64. R. *CASSUBICUS* (Lin! spec. 775.) foliis glabris, radicalibus petiolatis reniformibus crenatis, caulinis in lobos lineares serratos partitis, calyce pubescente petalis brevioribus. ♀ in pratis humentibus Borussiae orientalis, Russiae et Sibiriae. Sims. bot. mag. t. 2267. R. auricomus b. Biria ren. 39. Schlecht. anim. ran. 2. p. 7-10. Priori minus affinis. DC. syst. 1. p. 267.
65. R. *ABORTIVUS* (Lin! spec. 776.) foliis glabris, radicalibus petiolatis cordato-orbiculatis crenatis, nonnullis 3-partitis 3-sectisve, caulinis in lobos oblongo-lineares 3-5 partitis, calyce glabro petalis sublongiore. ♀ in Virginia et Carolina. R. auricomus var. c. Bir! ren. 39. Carpella ovata compressa parva laevia mutica in capitulum ovatum aggregata. DC. syst. 1. p. 268. (v. v.)
 ℓ. *nitidus* Walt. fl. car. 159. R. auricomus var. d. Bir! ren. 39. duplò fere major. (v. s.)
66. R. *SCLERATUS* (Lin! spec. 776.) foliis glabris, radicalibus petiolatis 3-partitis, lobis 3-lobatis obtusè subincisis, summis 3-partitis lobis oblongo-linearibus integris, floralibus oblongis, calyce glabro, carpellis minimis in spicam oblongam dispositis. (I) in fossis et paludibus totius Europae, Sibiriae, Cochinchinae, etc. OEd. fl. dan. t. 371. Hecatonia palustris Lour! fl. coch. 371. R. digitatus Gilib. exerc. phyt. 1. 373 ex Schlecht. — DC. syst. 1. p. 227 et 268. (v. v.)
 ℓ. *umbellatus*, (Roxb. in Willd. enum. 588.), pericarpiorum spicis magis oblongis. In horto Bengalensi à Roxburgio lecta. (v. s.)
 γ. *minimus*, horte nano, foliis radicalibus 3-fidis. in Tauriâ. A vero R. sclerato differt tantum ut R. parvulus Lin. à R. philonoti. (v. s.)
67. R. *INUNDATUS* (R. Brown! prod. nov. holl. 2. ined.) foliis glabris, radicalibus petiolatis 5-partitis, lobis 3-fidis, lobulis linearibus apice 3-dentatis, caule erecto 2-4-floro subnudo, pedunculis oppositifoliis, calycinis sepalis glabris rotundatis. (O) in Nova Cambria australi ad Guillielmi rivum. — Flores minimi, calycis sepalia glabra rotundata margine membranacea. DC. syst. 1. p. 269. (v. s. comm. à cl. R. Brown.)
68. R. *LANGSDORFII*, foliis subpubescentibus palmato-∞-fidis, laciniis linearibus tenuissimis, caule 1-floro, sepalis oblongis subreflexis petalis dimidio brevioribus. — propè Ochotsk. (Langsdorf ex Fisch.) Herba pusilla, folia fere ut in Batrachii aut in R. multifido palmato-laciniata sub lente pubescentia. Flos flavus terminalis parvus. (v. s.)
69. R. *MULTIFIDUS* (Parsh! fl. bor. am. 2. p. 736.) foliis glabris 3-partito-multifidis, laciniis incisis tenuibus acutis, caule natante, calycinis sepalis subrotundis petalis paulò brevioribus. ♀ nataus in aquis Louisianae superioris. Species notabilis R. aquatilis et speciatim hujus var. capillaceam omnino refert, sed floribus luteis primâ fronte distinctissima. DC. syst. 1. p. 270. (v. s.)
70. R. *RIVULARIS* (Banks et Sol! mss.) foliis glabris longè petiolatis 3-partitis, partitionibus cuneiformibus apice incisis, caule natanti-reptante, pedunculis oppositifoliis 1-floris, petalis lanceolatis remotis. (I) in rivulis Novae-Zelandiae. — Carpella ovata subcompressa laevia. DC. syst. 1. p. 270. (v. s. comm. à cl. Banks.)
71. R. *ACAULIS* (Banks et Sol! mss.) foliis glabris, radicalibus longè petiolatis 3-partitis, lobis ovalibus integerrimis, scapo 1-floro nudo petiolis brevioribus, calyce patente, petalis lanceolatis remotis erectis. (O) in fossis limosis Novae-Zelandiae. An hujus et antecedentis petala sint pallidè flava aut alba dubito. DC. syst. 1. p. 270. (v. s. comm. à cl. Banks.)
72. R. *COLLINUS* (R. Brown! prod. fl. nov. holl. 2. ined.) foliis pubescentibus, omnibus radicalibus petiolatis 3-sectis, segmentis ovalibus apice subdentatis, scapo nudo 1-floro foliis longiore, calyce patente, petalis stipitatis ovali-oblongis. (O) in collibus terrae Van-Diemen. — Flos minimus erectus, calyx

- caducissimus, ovaria pauca glabra. DC. syst. 1. p. 271. (v. s. comm. à cl. Brown.)
73. *R. PUMILIO* (R. Brown! prod. nov. holl. 2. ined.) foliis subvillosis 3-partitis, lobis 3-fidis integrisve oblongis, caulibus erectiusculis, pedunculis oppositifoliis, carpellis villosis. ☉? in terrâ Van-Diemen. Priori habitu paululum affinis, forsân tamen *R. parvifloro* et vicinis affiniôr? DC. syst. 1. p. 271. (v. s. comm. à cl. R. Brown.)
74. *R. LAPPONICUS* (Lin! spec. 778.) foliis glabris, radicalibus longè petiolatis 3-partitis, lobis dilatatis obtusis grossè dentatis, scapo nudo 1-floro foliis longiore, calyce 3-sepalo reflexo. 2℥ in Lapponiâ. Smith in fl. lapp. ed. 2. p. 194. t. 3. f. 4. Stips rariissima tenella glabra repens. Carpella 6-10 in capitulum subrotundum aggregata. DC. syst. 1. p. 271. (v. s.)
75. *R. Gmelini* (DC. syst. 1. p. 303.) foliis 3-5-partitis oo-fidisve ceterum integerrimis, caulibus declinatis, floribus pedunculatis oppositifoliis, calyce reflexo, carpellis subrotundis stylo recto terminatis. — in Sibiariâ et Americâ valdè boreali. Schlecht. anim. 2. p. 35.* — Gmel. sib. 4. p. 203. t. 83. β.
76. *R. HYPERBOREUS* (Rothb. act. hafn. 10. p. 458. t. 4. f. 16.) foliis glabris petiolatis 3-fidis, lobis ovali-oblongis divaricatis, lateralibus subfidis, medio integro, vaginis basi bianiculatis, caule filiformi repente. 2℥ in Islandiâ, Groenlandiâ, Alpibus jugis centralis Norvegiæ, etc. — fl. dan. t. 331. *R. montanus* α Bir. ren. 39. Species *R. cymbellarie* affinis, sed foliis 3-fidis facîle distinguenda. DC. syst. 1. p. 272. (v. s.)
77. *R. PYGMÆUS* (Walh. fl. lapp. n. 286. t. 8. f. 1*.) foliis glabris 3-5-fidis, radicalibus petiolatis, caulino sessili, caule 1-floro, calyce glabro petalis longiore subreflexo; carpellis subrotundis stylo adunco brevi mucronatis. 2℥ in Alpibus Norvegiæ, Finmarkiæ, Lapponiæ etc. — *R. nivalis pygmæus* Lin. fl. lap. n. 232. var. γ. t. 3. f. 3. Herba minima glabra Saxifragam rivularem æmulans et inter *R. hyperboreum* et nivalem media. DC. syst. 1. p. 273*.
(v. s. comm. à cl. C. Smith.)
78. *R. NIVALIS* (Gunn. norv. 627.) foliis glabris, radicalibus petiolatis 5-fidis, lobis integerrimis ovatis, caule 1-floro, calyce hirsuto petalis obtusis dimidio brevior. 2℥ in Lapponiâ, Norvegiâ, Sibiariâ, etc. *R. nivalis* α. Lin! spec. 778. *R. caule uniflora*, etc. Lin. fl. lapp. ed. 1. n. 232. t. 3. f. 2. excl. var. γ. Carpella subrotunda lævia stylo longitudine ipsorum pericarpiorum recto rostrata. DC. syst. 1. p. 273. (v. s.)
β. *sulphureus* (Phipps. voy. 202.) *R. nivalis* β. Wahl fl. lapp. 157. *R. primus* Mart. spitzb. t. 6. f. c. ex Schelecth. anim. 2. p. 14*.
79. *R. FRIGIDUS* (Willd! spec. 2. p. 1312.) foliis glabris, radicalibus cuneatis obovatis orbiculatisve apice grossè dentatis incis, caulinis in lobos 5-7 integros palmatifidis, caule 1-floro, calyce hirsuto petalis emarginatis brevior. 2℥ in Asiâ boreali, Sibiariâ. *R. Altaicus* Laxm! nov. comm. acad. Petrop. 1774. v. 18 p. 533. t. 8.* *R. sulphureus*. DC. syst. 1. p. 274. exclusis syn. Mart. Phipps et Wahl. Herba glabra digitalis *R. nivali* et *R. glaciali* affinis. (v. s.)
80. *R. ESCHENHOLZII*. (Schlecht. anim. ran. 2. p. 16.* t. 1.) foliis ciliatis, radicalibus petiolatis 3-partitis, partitionibus lobatis, caule subbifloro, calyce hirsuto petalis brevior, carpellis obliquè ovatis stylo ipsis brevior terminali. — in Insulis Unalaska et Sti Georgii (Cham. ex Schlecht.)
81. *R. PERUVIANUS*. (Pers! ench. 2. p. 103.) foliis subpilis, radicalibus longè petiolatis semi-orbiculatis crenatis, caulinis sessilibus in lobos lineares integros oo-partitis, calyce villosissimo. 2℥ in Peruviâ et in humidis Antisanæ. H. B. et K! nov. gen. 5. p. 44. Deless! ic. select. 1. t. 37. Carpella ovato-compressa minima lævia. DC. syst. 1. p. 274. (v. s. in h. Juss. H. B. et K.)
82. *R. POLYRHIZOS*. (Teph. in Willd. spec. 2. p. 1324*.) foliis glabris, radicalibus petiolatis 3-5-fidis, lobis ovatis integris, caulinis sessilibus in lobos lineares integros oo-partitis, caule subprostrato 1-3-floro, calyce patente glabro. 2℥

in regione Caucasicâ, desertis Caspici et collibus Altaici jugi. Deless! ic. sel. 1. t. 38. Schlecht. anim. 2. p. 17. Bieb. pl. rar. ross. 1. t. 19. Carpella lævia stigmate coronata. DC. syst. 1. p. 275. (v. s.)

83. *R. DEMISSUS* (DC. syst. 1. p. 275.) foliis glabris, radicalibus petiolatis 3-5-partitis, lobis in lacinias lineares fissis, scapo subnudo demisso 1-2-floro, calyce patente pubescente. \mathcal{U} in monte Libano. — Carpella suborbiculata compressa submutica. (v. s.)

84. *R. PEDATIFIDUS* (Smith! in Rees cycl. n. 72^a.) foliis subpuberulis, radicalibus petiolatis palmatim pedatimve in lobos lineares integros partitis, scapo subnudo erecto 1-2-floro, calyce patente glabro. \mathcal{U} in Sibiria et in sinu Sti Laurentii. Schlecht. anim. 2. p. 18^a. Carpella ovata stylis subrecurvis superata. DC. syst. 1. p. 275. (v. s. in h. Smith.)

85. *R. MONTANUS* (Willd! spec. 2. p. 1321.) foliis radicalibus glabris 3-partitis orbiculatis, laciniiis 3-fidis obtusis, caulinis sessilibus in lobos lineares integerrimos 3-5-partitis, caule 1-floro apice adpressè pubescente, calyce glabriusculo. \mathcal{U} in pratis asperis Alpium Austriæ, Helvetiæ, Galliæ et Pyrenæorum. *R. trilobatus* Kit. ex Schlecht. anim. 2. p. 21. — DC. syst. 1. p. 276. (v. v.)

β . *tenuifolius*, foliorum laciniiis acutis. *R. gracilis* Schlecht! pl. exsic. — in Jurasso, Alpibus Appuanis. (v. s.)

86. *R. VILLARSI* (DC! fl. fr. 4. p. 876^a.) foliis pubescentibus, radicalibus suborbiculatis 3-partitis, laciniiis 3-fidis dentatis acutis, caulinis sessilibus in lobos lineares integros 3-5-partitis, caule 1-floro apice pubescente, calyce patentim villosulo. \mathcal{U} in Alpibus Sabaudia, Delphinatû et Gallo-provinciæ. *R. Breynianus* Crantz. aust. 2. p. 91. t. 4. f. 2^a? *R. oreophilus* Bieb. suppl. 383. ex Schlecht. anim. 2. p. 37. (v. \mathcal{U} .)

87. *R. GOUANI* (Willd. spec. 2. p. 1322.) foliis radicalibus orbiculatis 5-lobatis, lobis inciso-dentatis, caulinis sessilibus palmato-partitis, lobis dentatis, caule pubescente, calyce subvillosulo. \mathcal{U} in pratis Pyrenæorum. — *A. R. montano* distinctissima. *R. Pyrenæus* Gouan! ill. 33. t. 17. f. 1. 2^a. non Lin. *R. montanus* var. *g.* Bir! ren. 40. — DC. syst. 1. p. 277. (v. v.)

88. *R. ACRI* (Lin! spec. 779.) foliis pubescentibus subglabrisve palmato-partitis, lobis inciso-dentatis acutis, summis linearibus, caule erecto ∞ -floro subpubescente, pedunculis teretibus, calyce subvillosulo, carpellis mucrone subrecto terminatis. \mathcal{U} in pratis Europæ. Smith. engl. bot. t. 652. variat fl. pleno. Curt bot. mag. t. 215. DC. syst. 1. p. 277. (v. v.)

β . *sylvaticus*, petiolis foliisque subtus velutino-villosis. *R. sylvaticus*. Thuill! fl. par. ed. 2. p. 276. *R. lanuginosus* β . DC. fl. fr. 4. p. 899 ex Schlecht. anim. (v. s.)

γ . *multifidus*, foliorum lobis inciso-multifidis. *R. polyanthemus* Lob. ic. 686. f. 1^a (v. s.)

89. *R. BATTIUS* (Tenor. fl. neap. prod. suppl. 1. p. 61 flor. 1. t. 50.) foliis 3-5-partitis, partitionibus 3-partitis, lobis lanceolatis bifidis utriusque pubescentibus, caule erecto villosulo ∞ -floro, pedunculis teretibus calyce adpresso petalis dimidio brevioribus. \mathcal{U} in monte Pollino regni Neapolitani. Affinis *R. acri* et polyanthemo. DC. syst. 1. p. 278. (v. s.)

90. *R. STEVENI* (Bess. enum. Volhn. n. 683.) foliis 3-5-partitis, laciniiis cuneatis trifidis inciso-dentatis, summis linearibus, radice multicipite, caule subnudo, pedunculis teretibus, calycibus hirsutis patulis, stigmatibus patentibus. \mathcal{U} in pratis Volhyniæ et Podoliæ. Aff. *R. acri*.

91. *R. CAUCASICUS*. (Bieb! fl. cauc. 2. p. 27^a.) foliis radicalibus longè petiolatis subpubescentibus 3-partitis, lobis 3-fidis inciso-serratis, caulinis petiolatis 3-sectis, summorum lobis linearibus, caule erecto ad collum fibrilloso, pedicellis teretibus, calyce patente. \mathcal{U} in Caucaso subalpino. Carpella compressa lævia, stylo brevissimo subuncinata in capitulum subrotundum aggregata. DC. syst. 1. p. 279. (v. s.)

92. *R. RUFULUS* (Brot. fl. lus. 2. p. 367*.) foliis radicalibus 3-5-partitis, lobis oo-fidis, caulinis inferioribus petiolatis 3-partitis, summis linearibus, caule erecto pedunculisque teretibus fæctis villosis, calyce patente. \mathcal{U} in Lusitaniâ. *R. chærophyllos* var. e. Bir. ren. 42. nimis affinis *R. acris*. DC. syst. 1. p. 279.
93. *R. POLYANTHEMOS* (Lin! spec. 779.) foliis 3-5-palmato-partitis inciso-multi-fidis, lobis sublinearibus, caule erecto oo-floro petiolisque patentim pilosis, pedunculis sulcatis, calyce hirsuto, fructibus submuticis. \mathcal{U} in sylvis Belgii Germaniæ, Hungariæ, etc. *R. Napellifolius* var. Crantz. aust. 2. p. 90. t. 4. f. 1. sup. Similis *R. acris* et præsertim hujus var. \mathcal{E} . DC. syst. 1. p. 279. (v. s.)
94. *R. NEMOROSUS* (DC. syst. 1. p. 280.) foliis radicalibus ultra medium 3-5-fidis. lobis cuneiformibus 3-fidis, lobulis apice dentatis, caule erecto petiolisque patentim pilosis, pedunculis sulcatis, carpellis stylo uncinatis. \mathcal{U} *R. polyanthemus* \mathcal{E} . Wallr! sched. 291. (v. v.)
 - a. *multiflorus*, foliis superioribus in lobos lineares integros partitis. *R. polyanthemus*. Sut. fl. helv. 1. p. 341. — in montanis ad radices Jurassi. (v. v.)
 - β . *pauciflorus*, caule 1-3-floro, foliis superioribus sublobatis inciso-dentatis, summis minimis. In subalpinis Vallesis, Cebennorum, Pyrenæorum. *R. aureus* Schleich! pl. exsic. *R. villosus* St. Am. fl. ag. 227. bouq. t. 5. (v. s.)
95. *R. LANUGINOSUS* (Lin! spec. 779.) foliis 3-fidis holosericeis, lobis latis dentatis subincisis, caule erecto oo-floro petiolisque reflexo-pilosis, pedunculis teretibus, carpellis plano-compressis stylo uncinatis. \mathcal{U} in sylvis montanis Europæ. — Oeder fl. dan. t. 397. *R. magnus* valde hirsutus flore luteo J. Bauh. hist. 3. p. 417. f. 3. DC. syst. 1. p. 281. (v. v.)
 - β . *geraniifolius*, foliis superioribus 3-partitis, lobis oblongis acuminatis grossè serratis. *R. montanus* subhirsutus, etc. C. Bauh. pin. 182. Tourn! inst. 291.
 - γ . *parvulus*, caule 1-2-floro, foliis minutissimis. in Jurasso. (v. s. in h. Ser.)
 - δ . *Constantinopolitanus* (D'Urv! enum. pl. arch. n. 476.) foliis radicalibus amplissimis longissimè petiolatis trifidis lobis latis subincisis, supremis lacinatis, lacinis linearibus, cunctis sericeo-lanuginosis, caule erecto lanuginoso multiflora, calyce reflexo, carpellis planis suborbiculatis stylo uncinatis. \mathcal{U} circa Byzantium ad sepes frequens. — *R. lanuginosus* γ . DC. syst. 1. p. 281. — ic. Aubr. in Bibl. Juss.
96. *R. TUBEROSUS* (Lapeyr! abr. pyr. 320.) foliis radicalibus amplis 3-lobis, lobis distantibus, lateralibus bilobis, intermedio cuneiformi 3-lobo, lobulis incisis, caule subadscendente pubescente, pedunculis divaricatis teretibus, calyce adpresso. \mathcal{U} in Pyrenæorum pratis. Schlecht. anim. 2. p. 26. Affinis *R. lanuginoso*. An satis à *R. nemoroso* distincta. DC. syst. 1. p. 281. (v. s. in h. Mus. Par.)
97. *R. ANEMONIFOLIUS* (DC. syst. 1. p. 282.*) foliis radicalibus 3-partitis pubescentibus basi cuneatis, lobis inciso-dentatis, caulino 3-secto, lobis summisque foliis lougè-linearibus integris, caule erecto 1-2-floro basi glabro apice pubescente, calyce reflexo. \mathcal{U} in Cappadociâ. (v. s. in h. Mus. Par.)
98. *R. NAEPELLIFOLIUS* (DC. syst. 1. p. 282.*) foliis radicalibus 3-partitis, partitionibus oo-fidis, lobis linearibus, floralibus 3 partitis, lobis linearibus, caule erecto 1-2-floro ad collum fibrilloso adpressè villosa, calyce adpresso. \mathcal{U} in Cappadociâ et in agro Constantinopolitano. Tourn! cor. 20. Fructus ignotus. (v. s. in h. Mus. Par. et Desf.)
99. *R. PLATYSPERMUS* (Fisch! in litt.) foliis radicalibus oo-fidis, caulinis 3-partitis, lobis linearibus, caule erecto, carpellis compressissimis orbiculatis membranaceis stigmate brevi uncinatis. \mathcal{U} ad lacum salsum Inderiensem deserti Kirghisorum. Species distinctissima, *R. oxyspermo* et *R. dissecto* affinis. (v. s.)

100. *R. DISSECTUS* (Bieb. fl. taur. 2. p. 25.* suppl. 381.) foliis radicalibus subpinnatim ∞ -partitis, lobulis linearibus acutis, caulinis in lobos lineares integerrimos partitis, caule erecto 1-2-floro pubescente, calycibus hirsutis patentibus. \mathcal{L} in subalpinis Caucasi orientalis, Sibiriae. *R. nivalis* Pall. ind. taur. ex Bieb. DC. syst. 1. p. 282.* (v. s.)
101. *R. VILLOsus* (DC. syst. 1. p. 283.*) foliis villosis 3-sectis, segmentis ovatis inciso-dentatis, floralibus 3-partitis, lobis oblongis subintegris, caule erecto dichotomo pilis subdeflexis hispido, calyce hispido patente. \mathcal{L} in Persia. Ovaria in capitulum subrotundum aggregata stylo uncinato superata. (v. s. in h. Lamb.)
102. *R. CAPPADOCIcus* (Willd. spec. 2. p. 1326.) foliis pubescentibus; radicalibus cordatis 3-fidis, lobis dentatis acuminatis, caulino inferiore petiolato, superiore sessili integro, caule erecto subbifloro, calyce patente, radice praemorsa. \mathcal{L} in Cappadocia. *R. chærophyllus* var. g. Bir. ren. 43. Carpella ovata subcompressa obtusè subuncinata lævia. DC. syst. 1. p. 283. (v. s. in h. Mus. Par.)
103. *R. SERICEUS* (Poir! dict. 6. p. 109.*) foliis villososericeis, radicalibus petiolatis 3-5-partitis, caulinis 3-sectis, segmentis longè petiolatis 3-partitis, omnium lobis inciso-serratis, caule erecto villososo ∞ -floro, calyce patente. \mathcal{L} in insulis Mauritanis. Deless! ic. sel. 1. t. 39. Carpella compressa lævia stylo brevissimo acuminata. DC. syst. 1. p. 284. (v. s.)
104. *R. RHEADIPOLIUS* (DC. syst. 1. p. 284.) foliis subglabris pinnatim 3-sectis, inferioribus petiolatis longis, segmentis ovalibus 3-lobis grossè dentatis, superiorum lobulis linearibus, caule erecto subramoso subglabro, calyce reflexo. \mathcal{L} in agro Magadoriensi et forsàn in Canariis. Deless! ic. sel. 1. t. 40. Fructus ignotus. (v. s. in h. Desf.)
105. *R. REPENS* (Lin! spec. 779.) foliis pinnatim 3-sectis, segmentis cuneatis 3-lobatis inciso-dentatis, caule flagella prostrato-repentia è collo agente, florifero suberecto, calyce adpresso, carpellis acuminè recto. \mathcal{L} in Europà, Pennsylvania. OEd. fl. dan. t. 795. engl. bot. t. 516.* Variat flore pleno. DC. syst. 1. p. 285. (v. v.)
- β . *erectus*, caulibus floridis erectis, flagellis repentibus nullis.
- γ . *glabratus*, foliis cauleque glaberrimis. *R. lucidus*. Poir! dict. 6. p. 113.*
- δ . *linearilobus*, flagellis reptantibus longissimis floriferis, lobis foliorum angustissimis. (v. s. in h. Ser.)
106. *R. MARGINATUS* (D'Urv. enum. pl. arch. n. 479.) radice fibrosà, caule basi ramoso patulo subhirsuto, foliis margine calloso cinctis, inferioribus petiolatis trisectis, segmentis crenatis obtusis, caulinarum laciniis linearibus, calyce reflexo, carpellis rotundatis planis marginatis stylo subuncinatis. \mathcal{L} in collibus circa Trapezum. Petala parva.
107. *R. JAPONICUS* (Langsd. ex Fisch! in litt.) foliis palmatim 3-sectis, segmentis subrotundis obtusè inciso-dentatis, caule repente, calyce reflexo. \mathcal{L} ? in Japonià prope Nagasaki. Carpella compressa lævia dorso subalata stylo apice subuncinata in capitulum subrotundum aggregata. Japonicè *Kimbu nohana* dicta.
108. *R. DIFFUSUS*, villosus, foliis basi cordatis subrotundis 3-lobatis dentatis, caule elongato diffuso, pedicellis 1-floris oppositifoliis, calyce adpresso, carpellis lævibus acuminè recto. \mathcal{L} in Napaulià. Wallich. Flores parvi verosimiliter flavi. (v. s.)
109. *R. GERANIoidES* (H. B. et K! nov. gen. et spec. amer. 5. p. 44.*) villosus, foliis radicalibus 3-sectis, segmentis petiolulatis basi cuneatis 3-fidis inciso-dentatis, caule adscendente paucifloro, pedicellis 1-floris oppositifoliis, petalis oblongis, calyce subreflexo. \mathcal{L} in Novà Granadà. Carpella compressa lævia stylo brevi recto aristulata. DC! syst. 1. p. 286. (v. s. in h. H. et B.)
110. *R. CHILENSIS* (DC. syst. 1. p. 286.*) caulibus procumbentibus petiolisque *hispidis*, foliis subvillosis cordato-orbiculatis 3-5-fidis, lobis grossè dentatis,

- calycibus villosissimis. \mathcal{L} in Chili, propè Talcaguanho. Schlecht. anim. 2. p. 27. Affinitas dubia ob carpella ignota. (v. s.)
111. *R. LAFFACEUS* (Smith! in Rees cycl. n. 61.*) foliis villosis petiolatis pinnatisectis pinnatifidisve, segmentis ovatis 3-fidis acutè inciso-lobatis, caule erecto oo-floro, calyce patente, carpellis stylo uncinato superatis. \mathcal{L} in Novâ-Hollandiâ. Species affinis *R. pinnato*, sed carpella non tuberculosa. DC! syst. 1. p. 286. (v. s. in h. Mus. Par. et Sm.)
- β . *obtusatus*, foliis brevioribus obtusioribus 3-partitis, caulinis subnullis, caule 2-floro.
- γ . *pubescens*, caule petiolisque adpressè pubescentibus, calyce hispido.
112. *R. PILOSUS* (H. B. et K! nov. gen. et spec. amer. 5. p. 45.*) caule ascendente 2-chotomo, petiolis patentim pilosis, foliis adpressè subvillosis pinnatisectis, segmentis 3-fidis grossè dentatis, summorum linearibus oblongis, calyce reflexo subglabro, carpellis stylo recto. \mathcal{L} in Americâ meridionali. — Carpella in capitulum globosum disposita. DC! syst. 1. p. 287.* (v. s. in h. H. et B. et Deless.)
113. *R. NAPAULENSIS*, caule erecto 2-chotomo foliisque pilosis, foliis inferioribus 3-sectis, supremis 3-partitis, lobis oblongo-cuneatis dentatis, calyce villosi reflexo petalis obtusis ferè longiore, carpellis stylo recto. \mathcal{L} in Napauliâ. Wallich. (v. s.)
114. *R. PETIOLARIS* (H. B. et K! nov. gen. et spec. amer. 5. p. 45.*) foliis radicalibus longissimè petiolatis adpressè sericeo-pubescentibus, partitionibus 3-fidis inciso-dentatis, floralibus 3-partitis, lobis linearibus, caule dichotomo erecto, calyce reflexo. \mathcal{L} in Mexico propè Santa-Rosa. Carpella ovata glabra, stylo brevissimo recto subaristata. DC! syst. 1. p. 287. (v. s. in h. H. et B.)
115. *R. DICHOTOMUS* (Moq. et Sessé pl. mexic. ined.) foliis glabrinusculis, radicalibus longissimè petiolatis bipinnatisectis, caule erecto dichotomo, calyce reflexo. \mathcal{L} in Mexico. Flores lutei *R. bulbosi*, fructus apice in acumen rectum desinentes. DC. syst. 1. p. 288. (Descr. ex ic.)
116. *R. PERSICUS* (DC. syst. 1. p. 288.*) foliis glabris, radicalibus petiolatis pinnatisectis, segmentis 3-lobis, lobis ovatis basi cuneatis apice regulariter dentatis, caulinis 5-sectis, summis 3-partitis subpubescentibus, caule erecto 2-chotomo adpressè pubescente, calyce patente extus villosi. \mathcal{L} in Persiâ. Flores lutei magnitudine *R. repentis* et paulò majores; fructus ignotus. (v. s. in h. Lamb.)
117. *R. PLEBEIUS* (R. Brown! prod. nov. holl. vol. 2. ined.) caule erectiusculo dichotomo, basi petiolisque patentim pilosissimo, apice subglabro, foliis 3-sectis villosis, segmentis petiolulatis ovatis inciso-dentatis, summis linearibus integris, calyce subreflexo. \mathcal{L} in Novâ-Hollandiâ. Carpella compressa lævia stylo subuncinato terminata. DC. syst. 1. p. 288.* (v. s. comm. à cl. R. Brown.)
118. *R. HIATUS* (Banks et Sol! mas.) caule erecto dichotomo, petiolis pedunculisque pilosissimis, foliis caulinis petiolatis, inferioribus 3-sectis, segmentis petiolulatis ovatis dentatis, mediis 3-lobatis, summis ovali-oblongis, calycibus retroflexis. \mathcal{L} in graminosis Novæ-Zelandiæ. Aff. *R. plebeio*. Carpella compressa stylo brevissimo recto apiculata. DC. syst. 1. p. 289. (v. s. comm. à Cl. Banks.)
119. *R. HISPIDUS* (Michx! fl. bor. amer. 1. p. 321.) caule erecto ramoso petiolisque patentim pilosissimis, foliis 3-sectis 3-partitisve, segmentis ovalibus acutis inciso-dentatis, pedicellis adpressè pubescentibus, calyce adpresso. \mathcal{L} in sylvis Carolinæ inferioris, Virginie. Petala magnitudine *R. acris*, carpella compressa lævia, stylo brevissimo. DC! syst. 1. p. 289.* (v. s. in h. Mus. Par.)
120. *R. RECURVATUS* (Poir! dict. 6. p. 123.*) caule erecto petiolisque patentim pilosissimis, foliis 3-partitis adpressè villosis, partitionibus ovalibus subincisè dentatis, calyce reflexo, carpellis stylo uncinatis. \mathcal{L} in Americâ septentrionali.

Deless! ic. sel. 1. t. 41. *R. saniculæformis* Muhl. ex Raf! in litt. Affinis *R. Pensylvanico* sed evidenter differt. Petala linearia ex Pursh. DC. syst. 1. p. 290.* (v. s.)

β. *Nelsonii*, foliorum lobis approximatis, pedicellis adpressè pilosis folii longitudinis. in insulâ Unalaschkâ.

121. *R. PENNSYLVANICUS* (Lin. suppl. 272.) caule erecto petiolisque rigidè patentimque pilosis, foliis 3-sectis adpressius villosis, segmentis subpetiolatis acutè 3-lobis inciso-serratis, calyce reflexo, carpellis stylo recto. 2/ in Pennsylvania et Canadâ. *R. Canadensis* Jacq. misc. 2. p. 343.* ic. rar. 1. t. 105. Carpella in capitulum ovatum oblongumve conferta. DC. syst. 1. p. 290. (v. v.)

122. *R. FASCICULARIS* (Muhlenb. in Big. fl. bor. 137.) caule erecto ramoso, foliis adpressè pubescentibus trisectis segmento medio 3-partito subpinnatisectis, lateralibus remotè 3-lobis, calyce patente subtus villosus petalis brevior. 2/ in Americâ boreali. Schlecht. anim. 2. p. 30. t. 2.* Dux species hic videntur confusæ. DC. syst. 1. p. 291.

123. *R. BELVISII* (DC. syst. 1. p. 291.*) caule erecto ramoso petiolisque rigidè parèque pilosis, foliis glabris 3-sectis, segmentis 3-lobis, lobis oblongis apice acutè incis, calyce glabro patente petalis brevior. 2/ in Americâ boreali. Carpella compressa lævia stylo brevi recto aristata. (v. s.)

124. *R. MARYLANDICUS* (Poir! dict. 6. p. 126.) caule erecto subramoso petiolisque basi molliter hirsutis, apice adpressè pubescentibus, foliis glabriusculis 3-sectis, segmentis 3-lobatis, lobis oblongis acutis inciso-dentatis, calyce glabro patente petalis brevior. 2/ in Marylandiâ, Pennsylvania, Virginia. Carpella compressa stylo recto acuminata. DC. syst. 1. p. 291.* (v. s.)

125. *R. CAROLINIANUS* (DC. syst. 1. p. 292.*) caule erecto subramoso, petiolisque adpressè pubescentibus, foliis glabriusculis 3-sectis 3-lobisve, lobis ovatis subincisis dentatis, calyce glabriusculo reflexo petalis paulò brevior. 2/ in Carolinâ inferiori. *R. hispidus* var. Michx! fl. amer. bor. 1. p. 321. valdè aff. *R. Marylandico*. (v. s.)

126. *R. TOMENTOSUS* (Poir! dict. 6. p. 127.*) caule patentim villosissimo ascendente 1-2-floro, foliis petiolatis tomentosius 3-sectis, summo sessili ovato integro, calyce villosissimo subreflexo. 2/ in Carolinâ editiore. *R. lanuginosus* var. Pursh. ex h. Lamb. Petala calyce paulò longiora; fructus ignotus. DC. syst. 1. p. 292. (v. s. in h. Bosc.)

127. *R. PRÆMORSUS* (H. B. et K! nov. gen. et spec. amer. 5. p. 47*) caule erecto subbifloro petiolisque adpressè villosis, foliis 3-sectis, segmentis obtusè inciso-dentatis, lateralibus bifidis, medio 3-fido petiolulato, calyce reflexo, petalis 10-15 cuneatis. 2/ in Americâ meridionali. Affinis ex habitu *R. Marylandico*. Carpella stylo brevi recto aristulata. DC. syst. 1. p. 293.* (v. s. in h. H. et B.)

128. *R. BONPLANDIANUS* (H. B. et K. nov. gen. et spec. amer. 5. p. 46.*) caulibus erectis 2-3-floris pilosiusculis, foliis 3-fidis 3-partitis 3-sectisque suprâ glabris, subtus pubescenti-pilosis, segmentis crenato-serratis 2-3-lobis, calycibus pilosis reflexis, petalis 10-12 apice rotundatis. 2/ in regno Novo-Granatensi alt. 1000-1200 hex.

129. *R. GROÏDES* (H. B. et K! nov. gen. et spec. amer. 5. p. 47*) caule erecto 1-3-floro petiolisque patentim villosissimis, foliis adpressè villosis 3-fidis tripartitisve, lobis ovatis dentatis, calyce reflexo, petalis 10 oblongo-linearibus. 2/ in Mexici Montosis. Herba Geum montanum habitu æmulans. Petala 10 oblongo-linearia, basi vix angustiora, calyce duplò longiora. DC. syst. 1. p. 293. (v. s. in h. H. et B.)

130. *R. SIBBALDIOIDES* (H. B. et K! nov. gen. et spec. amer. 5. p. 48*) caule subnullo 1-floro, foliis pilosis pinnatisectis, segmentis ovatis dentatis, terminali 3-lobis, floralibus pinnatifidis, calyce patente petalis oblongis brevior.

¶ in America meridionali. Herba humillima Sibbaldiam procumbentem æmulans, DC. syst. 1. p. 293* (v. s. in h. H. et B.)

131. *R. DELPHINIIFOLIUS* (H. B. et K. nov. gen. et spec. amer. 5. p. 48*) caulibus erectis paucifloris foliisque pilosis 3-sectis, segmentis lateralibus biterminalibus 3-partitis, laciniis 2-3-fidis inciso-serratis, lobis lanceolatis, calycibus pilosis reflexis, petalis 15. ¶ in regno Mexicano, alt. 1430 hex.

132. *R. PEDUNCULARIS* (Smith! in Rees cycl. n. 49*) foliis subglabris radicalibus petiolatis 3-sectis, segmentis 3-lobatis apice acutis incis, caulinis 1-2 linearibus 3-fidisve, caule erecto 1-folio 1-2-floro, calyce demum reflexo, petalis 8-10 oblongis. ¶ ad sylvarum margines freti Magellanici. Deless! ic. sel. v. t. 42. Habitus *R. bulbosi*. Carpella globosa lævia stylis uncinatis superata. DC! syst. 1. p. 294* (v. s. in h. Mus. Par. et Juss.)

133. *R. GRANDIFLORUS* (Lin. spec. 781.) foliis subvillosis radicalibus petiolatis 3-sectis, segmentis inæqualiter laciniatis, caulinis linearibus aut in lobos lineares partitis, caule erecto villosulo paucifloro, calyce reflexo. ¶ in Cappadocia. Desf. choix. 57. t. 44* — flores lutei majores quàm *R. bulbosi*. DC. syst. 1. p. 294. (v. s. in h. Deless.)

134. *R. PALUSTRIS*. (Lin! mss. Smith in Rees cycl. n. 52.*) foliis subhirsutis petiolatis ultrà mediam 3-fidis, sinibus obtusis, lobis ovatis incis dentatis, caule erecto ramoso subnudo, carpellis omnino lævibus stylo brevi recto acuminatis. ¶ DC. syst. 1. p. 294.*

α. *Orientalis*, caulibus 3-chotomis, foliis radicalibus brevius petiolatis *R. orientalis palustris* apii folio, etc. Tourn. cor. 20. in Oriente. (v. s. in h. Smith.)

β. *Corsicus*, caulibus 2-chotomis, foliis radicalibus longissimè petiolatis. *R. Corsicus*. DC. fl. fr. 5. p. 637* in Corsicà. (v. s. in h. Desf.)

135. *R. BULBOSUS* (Lin! spec. 778.) foliis radicalibus petiolatis 3-sectis subpinatisectis, segmentis 3-fidis inciso dentatis, medio petiolulato, caule erecto ad collum bulboso, calyce reflexo. ¶ in Europà et Americà boreali. Fl. dan. t. 551. engl. bot. t. 515. — Variat flore pleno. *R. maximo tubere*, etc. J. Bauh. hist. 3. p. 418. f. 1. DC. syst. 1. p. 295.

β. *bracteatus*, *R. bracteatus* Scheich! pl. exsic. seu brachiatus Schl. cat.

Sect. v. ECHINELLA.

Carpella tuberculis aculeisve scabra.

§. 1. *Foliis dissectis lobatisve.*

136. *R. PHILONOTIS* (Retz. obs. 6. p. 31*) foliis 3-lobatis 3-sectisve, lobis obtusis inciso-dentatis, medio pedunculato, calyce reflexo, caule basi non bulboso, carpellis ad margines tuberculorum minimorum seriem unicam gerentibus. (1) in Europà medià et australi, Taurià. DC. syst. 1. p. 296. (v. v.)

β. *intermedius*, foliis subglabris. *R. intermedius* Poir! dict. 6. p. 116* *R. pumilus* Thuill! fl. par. ed. 2. 1. p. 277. non Poir! — in humidioribus.

γ. *parvulus*, caule nano subunifloro. *R. parvulus* Linn! mant. 79. *R. parviflorus* Gouan. fl. monsp. 270* non Linn. — in siccis lapideis apricis aridis.

137. *R. TUBERCULATUS* (Kit. ex Balb. cat. h. taur. 1813. p. 64.) foliis glabris inciso-multifidis, lobis linearibus acutis, caule erecto oo-floro, carpellis compressis utrinquè tubercula obtusissima gerentibus. (2) ¶ inter segetes Hungariæ, in Iberià et Taurià merid. Schlecht. anim. 2. p. 35. *R. arvensis* valde similis staturâ, foliatione, ramificatione et florescentiâ. DC. syst. 1. p. 297. (v. v.)

138. *R. ARVENSIS* (Lin! spec. 780.) foliis glabris, primordialibus apice dentatis, radicalibus 3-partitis, caulinis oo-fidis, lobis linearibus, caule erecto oo-floro, carpellis utrinquè longè aculeatis. (1) in Europà medià et australi. Ed. fl. dan. t. 219. engl. bot. t. 135. *R. echinatus* Crantz austr. 2. p. 118. non Vent. — DC. syst. 1. p. 297. (v. v.)

- β. *orientalis echinatus arvensis* crassiore fructu. Vaill! herb. fructibus-duplò ferè majoribus, aculeis longioribus.
139. *R. MURICATUS*. (Lin! spec. 780.) foliis glabris petiolatis suborbiculatis 3-lobis grossè dentatis, caule erectiusculo aut diffuso, pedunculis oppositifoliis, calyce patente, carpellis utrinquè tuberculoso-aculeatis in cornu acuminatum rectum desinentibus. ① in totà Mediterraneâ regione. Lam! ill. t. 498. *R. lobatus* Moench. meth. 214. — DC. syst. 1. p. 298. (v. v.)
- β. *Creticus*, caule erecto. — Moris. hist. 2. p. 440. s. 4. t. 29. f. 24. in insulâ Cretâ.
- γ. *Carolinus*, petalis calyce longioribus. — *R. muricatus* Michx. fl. bor. amer. 1. p. 321. *R. echinatus* Vent! h. cels. p. et t. 73. in Carolinâ et Virginiâ.
- δ. *Tucumanicus*, carpellis minùs acuminatis. *R. palustris echinatus* Feuill. obs. 3. p. 58. t. 18. Propè rivum Plata.
- ε. *Brasilianus*, petiolorum vaginis dilatatis. *R. ventricosus* Vent! h. Cels. 73. in maritimis Brasiliæ.
140. *R. CHRUS* (DC. syst. 1. p. 299*.) foliis subvillosis cordatis orbiculatis latè dentatis, caulibus erectis villosis, pediculis oppositifoliis, calyce reflexo, carpellis utrinquè tuberculosus in cornu acuminatum uncinatum productis. ① in insulâ Scio archipelagi Græci. Radix calyx et petala *R. philonotis*, folia ferè *R. parviflori* et carpella ferè *R. muricati*. (v. s. in h. Oliv.)
141. *R. CORNUTUS* (DC. syst. 1. p. 300*.) foliis glabris 2-ternatim sectis, segmentis 3-lobis, lobis oblongo-linearibus acutis, pediculis oppositifoliis, calyce reflexo, carpellis utrinquè tenuiter tuberculosus in cornu longum acuminatum desinentibus. ① in Oriente. Species fructu echinato media inter *Ranunculos* et *Ceratocephalos*. (v. s. in h. Labill.)
142. *R. PINNATUS* (Poir! dict. 6. p. 126.) foliis villosis subpinnatisectis, segmentis petiolulatis oblongis acute subpinnatifidis, pediculis oppositifoliis, calyce reflexo, carpellis ovalibus stylo brevi recto submucronatis utrinquè tenuiter tuberculosus. ②? Nullum discrimen video inter specimina duarum varietatum, sed has disjungo ob patrias diversas.
- α. *Sonneratii*. *R. pinnatus*. Poir. l. c. *R. parviflorus* var. C. Bir. ren. 46. in Indiâ. (v. s. in h. Lam.)
- β. *Hermanni*. *R. Africanus pratensis* hirsuto similis. Herm. cat. pl. Afr. p. 18. ex h. Burm. nunc Deless. Ad Cap. Bonæ-Spei. (v. s. in h. Deless.)
143. *R. PARVIFLORUS* (Lin! spec. 780.) foliis villosis orbiculatis 3-lobis grossè dentatis incisive, caulibus molliter pilosis decumbentibus, pediculis folio oppositis, calyce demùm reflexo petala æquante, carpellis granulato-tuberculosus. ① in Europâ occident. australique. Smith. engl. bot. t. 120. *R. muricatus* Moench. meth. 215. Species distinctissima ob petala lutea oblouga, interdum numero 3-4. — DC. syst. 1. p. 300.
- β. *acutilobus*, foliis profundius 3-lobis, lobis apice incisis, dentibus acutis. in Teneriffâ. (v. v.)
- γ. *erectus*, caule erecto, foliis 5-lobis seu grossè 5-dentatis. in Oriente. (v. s. in h. Oliv.)
144. *R. TRILOBUS* (Desf! fl. atl. 1. p. 437. t. 113*.) foliis subvillosis glabratissive, primordialis subrotundis crenatis, ceteris 3-partitis, lobis divergentibus basi cuneatis apice dentatis, caule erecto, pediculis oppositifoliis, calyce demùm subreflexo petalis brevioribus, carpellis granulato-tuberculosus. ① in humidis regionibus Mediterraneæ. Aff. hinc *R. parvifloro*, illinc *philonoti*. *R. ammoides* Pall. herb. ex Fisch! in litt. — DC. syst. 1. p. 301. (v. v.)
145. *R. SESSILIFLORUS* (*R. Brown!* prod. fl. Nov.-Holl. v. 2. ined.) foliis subvillosis reniformibus 3-partitis aut grossè dentatis, caulibus procumbentibus, floribus sessilibus oppositifoliis, carpellis granulato-tuberculosus. ① in Novâ Hollandiâ. Dispositione florum affinis *R. nodifloro*, sed foliis lobatis nec integerrimis, cauleque prostrato nec erecto. DC. syst. 1. p. 302. (v. s.)

§. 2. *Foliis integris.*

146. *R. OPHIOGLOSSIFOLIUS* (Vill! dauph. 4. p. 732. t. 49*) foliis inferioribus petiolatis cordatis obtusis, superioribus oblongis sessilibus, caule erecto fistuloso, pedunculis oppositifoliis, carpellis granulato-tuberculosis. ① in fossis Europæ australis. *R. flammula* var. f. Bir! ren. 37. Carpella ovata compressa. DC. syst. 1. p. 248.* (v. v.)
147. *R. UIGINOSUS* (Willd. enum. 586.) foliis ovatis lanceolatisque serratis glabris, caule ascendente fistuloso? pedunculis oppositifoliis, petalis calycis longitudine. ① in udis Teneriffæ. Affinis *R. flammulæ* et *ophioglossifolio*; flores parvi ferè *R. abortivi*; fructus (immaturi visi) in capitulum sphericum aggregati, granulato-tuberculati? DC. syst. 1. p. 249. (v. s.)
148. *R. NODIFLORUS* (Lin! spec. 773.) foliis radicalibus petiolatis ovali-oblongis, floribus sessilibus oppositifoliis, carpellis granulato-tuberculosis stylo vix rostellatis. ① in Galliâ. Vaill! bot. par. 168. Carpella in capitulum globosum aggregata. DC. syst. 1. p. 250.* (v. v.)
ß. dentatus, foliis apice dentatis 3-nerviis. — *R. nodiflorus* *ß.* Sims. bot. mag. t. 2171. *R. nodiflorus* Waldst et Kit. pl. rar. hung. 2. p. 192. t. 176*. *R. sculus*, etc. Petiv. gaz. 39. t. 24. f. 9.
149. *R. LATRIFLORUS* (DC. syst. 1. p. 251.) foliis radicalibus petiolatis ovali-oblongis, floribus sessilibus oppositifoliis axillaribusve, carpellis granulato-tuberculatis stylo elongato rostellatis. ① in Oriente. Deless! ic. sel. 1. t. 50. *R. alpestris palustris*, etc. Cnp. h. cath. 185. ex h. Vaill. et Juss. (v. s.)

†. *Ranunculi non satis noti.*

§. 1. *Foliis integris.*

150. *R. OBTUSISCULUS* (Raf. in Journ. bot. 1808. v. 1. p. 225.) in Americâ boreali. Caulis erectus simplex; folia petiolata lanceolata obtusiuscula, flores pauci terminales. (Raf. l. c.)
151. *R. OVALIS* (Raf. in Journ. bot. 1814. v. 2. p. 268.) in Canadâ. — folia radicalia longè petiolata ovalia integra acuta villosa, caulina rara sessilia lanceolata; flores pauci terminales. (Raf. l. c.)
152. *R? KAMCHATICUS.* (DC. syst. 1. p. 302*.) Hab. in Kamchatkâ. Stirps non satis nota, calyce persistente ab omnibus ranunculis distincta, forsân cum *Ficariâ* glaciali Fisch. et etiâm cum *Calthæ* sectione primâ genus proprium inter *Ficariam* et *Caltham* medium conficiens. (v. s. in h. Pall. nunc Lamb.)

§. 2. *Foliis multifidis.*

153. *R. DEBILIS* (Raf. in Journ. bot. 1808. v. 1. p. 225.) Hab. propè Germatown in Pensylvaniâ. Caulis debilis; folia pauca longa petiolata semigeniculata bifida incisa serrata; flores terminales solitarii. DC. syst. 1. p. 303.
154. *R. FORSKÆHLII* (DC. syst. 1. p. 303.) *R. multifidus* Forsk. fl. aeg. 102. Smith. in Rees cycl. n. 71. *R. chærophyllus* var. *α* Bir. ren. 42. — in Arabiâ. folia oo-fida, inferiora pinnatim laciniata; caulis hepedalis erectus oo-florus, infernè teres, supernè striatus pilosus; calyx corollam æquans. (Forsk.) DC. syst. 1. p. 303.
155. *R. CANTONIENSIS.* *Hecatonia pilosa* Lour. fl. coch. 371. DC. syst. 1. p. 228. ① propè Cantonem Sinarum. — Caulis erectus ramosus pilosus, folia pilosa ternata lanceolata serrata; flos luteus terminalis; pedunculi solitarii; ovaria et carpella acuminata in capitulum globosum acuminata (Lour. l. c.) Com *Hecatonia palustris* sit certè *Ranunculi* species, hanc etiâm ad *Ranunculum* referendam judico.
156. *R. LAGASCANUS.* *R. pubescens.* Lag. cat. h. madr. p. 19. DC. syst. 1. p. 303. 2/ in Hispaniâ. Hirtus; pedunculi elongati teretes uniflori; calyces pilosi reflexi; folia 3-partita 3-loba crenata; summa lanceolata integerrima. (Lag.)

157. *R. HORNEMANNI*. (Schlect. anim. 2. p. 36.) *R. tuberosus* Horn. h. hafn. 2. p. 527. in Americâ bor. Calyces reflexi pilosi, pedunculis sulcati, folia 3-nata, foliola 3-lobata hirsuta, Radix tuberoso-bulbosa. Affinis. *R. philonoti*.
158. *R. OBTUSIFOLIUS*. (Horn. h. hafn. 2. p. 528. Schlect. anim. 2. p. 36.) in Hispaniâ. Similis *R. lanuginoso* sed lobi foliorum obtusi.
159. *R. LACINIATUS*. (Baumg. enum. st. trans. 2. p. 131. Schlect. anim. 2. p. 36.) hab. in montibus siccioribus Transylvaniæ.

XIV. *FICARIA*. *Dill. nov. gen. p. 108. t. 5.*

Calyx 3-phyllus caducus; petala 9 intus basi foveolata; stamina ovarique ∞ ; carpella lævia compressa obtusa. — Herba glabra perennis; radix grumosa. numerus partium variabilis; sepala interdum 5 et petala 8-12. DC. syst. 1. p. 304.

1. *F. RANUNCULOIDES* (Mönch. meth. 215.) radice grumosa, caule folioso, foliis cordatis. $\frac{1}{2}$ in sepibus dumetisque totius Europæ. *Ranunculus ficaria*. Lin! spec. 774. fl. dan. t. 479. DC. syst. 1. p. 304. (v. v.)
2. *F. GLACIALIS* (Fisch! in litt. icon.) radice fibroso-fasciculatâ, foliis ovatis, scapis nudis 1-floris. $\frac{1}{2}$ in Dahuriâ ad apicem montis Tchockondo: Stamina ovarique ∞ , calycis circiter longitudine, fructus ignotus. — Petala ficariæ sed calyx 5-sepalus. Habitus omnino *Calthæ* appendiculatæ, sed petala 13 nec o; an *Psychrophilis* aut potius *Ranunculo Kamchatico* affinium, ipso non repugnante Fischero. DC. syst. 1. p. 305. (Desc. ex ic.)

Trib. IV. *HELLEBOREÆ*. DC. syst. 1. p. 130 et 306.

Æstivatio calycis et corollæ imbricata. Petala nunc o, nunc irregularia bilabiata nectarifera. Calyx petaloideus. Carpella capsularia intus dehiscentia ∞ -sperma.

XV. *CALTHA*. *Pers. ench. 2. p. 107.*

Calyx coloratus 5-sepalus, sepalis suborbiculatis petaloideis. Petala o. Stamina ∞ . Ovaria 5-10. Capsulæ 5-10 compressæ patentēs 1-loculares ∞ -spermæ. — Herbae perennes glaberrimæ. DC. syst. 1. p. 306.

Sect. I. *PSYCHROPHILA*. DC. syst. 1. p. 307.

Calycis sepala persistentia. Scapo 1-flori nudi. Folia radicalia sagittata, appendicibus sursum erectis.

1. *C. APPENDICULATA* (Pers. ench. 2. p. 107.) scapo 1-floro brevissimo, foliis 3-fidis cuneiformibus, sepalis 5, pistillis 8. $\frac{1}{2}$ in montibus humentibus freti Magellanici. *C. paradoxa* Sol! mss. in bibl. Banks.* Forst! trans. Lin. soc. 8. p. 34. Scapus foliorum longitudine, sepala oblongo-lanceolata persistenti-marcescentia. DC. syst. 1. p. 307. (v. s.)
2. *C. SAGITTATA*. (Cav. ic. 5 n. 456. t. 414*) scapo 1-floro, foliis sagittatis auriculatis sursum inflexis, sepalis 9-10, pistillis 25. $\frac{1}{2}$ in aquis portus Egmont, in Americâ meridionali. *C. multicapsularis* Soland! mss. in bibl. Banks.* Forst! trans. Lin. soc. 8. p. 324. Stamina sepalis dimidio breviora.

Sect. II. *POPULAGO*. *Tourn. inst. 273. t. 145. DC. syst. 1. p. 308.* — *Caltha* Lin. gen. n. 703.

Calycis sepala decidua. Caules foliosi. Folia cordata aut reniformia, auriculis sursum non inflexis.

3. *C. PALUSTRIS* (Lin! spec. 784.) caule erecto, foliis cordatis suborbiculatis rotundè crenatis, auriculis rotundatis. $\frac{1}{2}$ in Europâ, secus sepes fossas et rivulos, in Asiâ occidentali et Americâ boreali. Ed. fl. dan. t. 668. Smith. engl.

bot. t. 506. Caulis dichotomus pedalis erectus, pedunculi sulcati, flores amplii. — variat flore pleno.

ℓ. minor, caule subunifloro ascendente. Populago minor. Tab. ic. 750. f. 2.

C. minor Mill! dict. n. 2 Auriculæ folia floresque dimidio minores. cl.

Miller assertit hanc per plurimos annos cultam characteres servasse! In Angliâ præcipuè adhucdum reperta.

4. *C. RADICANS* (Forst! in trans. Lin. soc. 8. p. 321. t. 17.) caule repente, foliis 3-angulari-cordatis serrato-crenatis. \mathcal{U} secus lacus et rivulos Scotiæ. Differt à *C. palustri* caule basi repente apice ascendente paucifloro, foliis ferè triangularibus, etc. DC. syst. 1. p. 309. (v. s.)

5. *C. INTEGRERRIMA* (Pursh! fl. bor. amer. 2. p. 390.) caule erecto corymboso, foliis orbiculato cordatis sinu clauso integerrimis, floralibus sessilibus reniformibus basi obsolete crenatis, sepalis ovalibus. \mathcal{U} secus rivulos Novæ-Angliæ et Virginie. Flores corymbosi, lutei, paulò minores quàm in *C. palustri*, sepala subobovata obtusissima. DC. syst. 1. p. 309. (v. s.)

6. *C. ASARIFOLIA* (DC. syst. 1. p. 309.) caule suberecto 1-floro, foliis cordato-reniformibus crenatis sinu obtusis, sepalis 6-7 ovalibus. \mathcal{U} in insulâ Unalashkâ, una ex ins. Aleuterianis. Folia radicalia petiolata, petiolo limbo duplò longiore basi in vaginam membranaceam amplam dilatato, *C. palustri* similia sed paulò minora. (v. s. in h. Banks.)

7. *C. PARNASSIFOLIA* (Raf. New-York. med. rep. 2. p. 351. n. 28. et in Journ. bot. 1808. 1. p. 229.) caule 1-floro 1-folio, foliis radicalibus petiolatis lanceolatis cordatis obtusissimis multinerviis, sepalis ellipticis. \mathcal{U} in New-Jersey et Carolinæ paludibus umbris. *C. ficarioides* Pursh. fl. bor. amer. 2. p. 389. DC. syst. 1. p. 310. Flores intensè lutei, magnitudine Ficariæ.

8. *C. LEPTOSEPALA* (DC. syst. 1. p. 310.) caule 1-folio 1-2 floro, foliis radicalibus petiolatis ovatis cordatis crenatis oo-nerviis, sepalis 10 linearibus. \mathcal{U} in orâ occidentali Americæ borealis. Capsulæ 8-10 brevissimè substipitatæ stylo acuminatæ. (v. s.)

9. *C. BIFLORA* (DC. syst. 1. p. 310.) caule 1 folio 2-floro, foliis radicalibus petiolatis reniformibus crenatis, sinu latissimo, sepalis oblongis. \mathcal{U} in Americâ boreali. Petioli basi in vaginam membranaceam dilatati. (v. s. in h. Banks.)

10. *C. FLABELLIFOLIA* (Pursh! fl. bor. amer. 2. p. 390. t. 17.) caule procumbente oo-floro, foliis dilatato-reniformibus, lobis patentissimis argutè acutèq; dentatis, sepalis obovatis, capsulis rostro adnato. \mathcal{U} in monte Pokono Pennsylvania. Flores lutei, magnitudine Ranunculi arvensis; aff. hinc *C. natanti*, illinc *C. radicans*. DC. syst. 1. p. 310. (v. s. in h. Lamb.)

11. *C. NATANS*. (Pall! itin. ed. min. 3. p. 248.) caule procumbente natante, foliis cordato-reniformibus crenatis, lobis subapproximatis posticè obsolete crenatis antice dentatis, sepalis ovalibus, capsulis rostro recto. \mathcal{U} in aquis stagnantibus Sibirie orientalis. Gmel. fl. Sib. 4. p. 192. t. 82. Flores albi, dimidio quàm in *C. palustri* minores. DC. syst. 1. p. 311. (v. s. in h. Lin. Lam.)

XVI. TROLLIUS. Lin. gen. n. 700.

Calyx coloratus 5-10-15 sepalus, sepalis deciduis petaloideis. Petala 5-20 parva basi tubulosa 1-labiata. Stamina ovarique oo. Capsulæ oo sessiles subcylindraceæ polyspermæ. — Herbæ glabræ erectæ perennes Ranunculos Chrysanthemos habitu et Helleboros characteribus, referentes DC. syst. 1. p. 311.

1. *T. EUROPEUS*. (Lin! sp. 782.) sepalis 15 in globum ferè conniventibus, petalis 5-10 longitudine staminum. \mathcal{U} in pratis montanis subfrigidis totius Europæ. OEd. fl. dan. t. 133. DC. syst. 1. p. 312. Herb. amat. t. 69. *T. altissimus* Wend. flor. 1818. p. 578. (v. v.)

ℓ. humilis, minor 1-florus, flore suprà folium vix pedunculato. — *T. hu-*

- millis Crantz austr. 2. p. 124. excl. syn. Buxb. — T. minimus Wend. 1. c. 579.
- γ. *napellifolius*, (Rœpp. in flor. 1820. p. 105.) foliis sub-7-partitis, sepalis 12, stylis recurvis. ♀ variat caule ramisque rectis et caule patulo ramis recurvis. T. medius Wend. in flor. 1818. p. 579.
2. T. ASIATICUS (Lin! sp. 782. excl. Buxb. et Tourn. syn.) sepalis 10 patentibus, petalis 10 staminibus longioribus. ♀ in sylvis pratisque Sibiriae. Curt. bot. mag. t. 225. herb. amat. t. 88. Simillimus T. Europæo. DC. syst. 1. p. 313. (v. s.)
3. T. CAUCASICUS (Stev. in mem. nat. mosc. 3. p. 265.) sepalis 10 patentibus, petalis 10 staminibus brevioribus. ♀ in Caucasi orientalis alpe Schahdaghad. Aff. T. asiatico, sed petala staminibus breviora.
4. T. PATULUS (Salisb. trans. lin. soc. 8. p. 303.) sepalis 5 patentibus, petalis 1-5 stamina æquantibus. ♀ DC. syst. 1. p. 313.
- α. *pedunculatus*, flore supra folium superius pedicellato. Deless! ic. sel. 1. t. 44. in Sibiria.
- β. *subinvolucratus*, flore supra folium superius subsessili. in Cappadociae montibus altissimis. Buxb. cent. 1. p. 15. t. 22. Digito vix longior.
5. T. AMERICANUS (Muhlenb. et Gaissenb. in Donn. cat. h. cant.) sepalis 5-10 patentibus, petalis 10-15 staminibus brevioribus. ♀ in Pennsylvania. Lodd. bot. cab. t. 56. Valdè similis T. laxo et T. Asiatico. DC. syst. 1. p. 313. (v. s. in h. Banks, Lamb. et Bosc.)
- α. T. *pentapetalus*, herb. Banks!
- β. T. *decapetalus*, herb. Bosc!

XVII. ERANTHIS. Salisb. trans. soc. Lin. 1807. v. 8. p. 303.

Involucrum in segmenta plurima fissum sub flore; flos sessilis 5-8-sepalus, sepalis coloratis petaloïdeis oblongis deciduis. Petala 6-8 tubulata ore inæqualiter 2-labiata brevissima. Stamina 20-30. Ovaria 5-6. Capsulæ pedicellatæ. Semina globosa simplici serie disposita. DC. syst. 1. p. 314.

1. E. HYEMALIS (Salisb. tr. Lin. soc. v. 8. p. 303.) sepalis 6-8 oblongis. ♀ ad radices montium Galliae, Helvetiae, Italiae, Austriae. Helleborus hyemalis. Lin! spec. 783. Jacq. fl. austr. t. 202. herb. amat. t. 42. DC. syst. 1. p. 314. (v. v.)
2. E. SIBIRICA (DC. syst. 1. p. 315.) sepalis 5 ovalibus. ♀ in Sibiria orient. Habitus omnino E. hyemalis; capsulæ pedicellatæ ovato-oblongæ, semina globosa subcompressa. (v. s. in h. Lamb.)

XVIII. HELLEBORUS. Adans. fam. 458.

Calyx persistens 5-sepalus, sepalis subrotundis obtusis magnis sæpè viridibus; petala 8-10 brevissima tubulata infernè angustiora nectarifera; stam. 30-60. Ovaria 3-10. Stigmata terminalia orbiculata; capsulæ coriaceæ; semina duplici serie disposita elliptica umbilicata. DC. syst. 1. p. 315.

§. 1. *Scapo aphyлло 1-2-floro.*

1. H. NIGER. (Lin! spec. 783.) foliis radicalibus pedatisectis glaberrimis, scapo aphyлло 1-2-floro bracteato. ♀ in montosis sylvaticis subalpinis Europæ. Jacq. fl. austr. t. 201. Curt. bot. mag. t. 8. flos amplus calyce petaloïdeo albo sæpius roseo. DC. syst. 1. p. 316. (v. v.)

§. 2. *caule paucifloro subdichotomo folia floralia subsessilia fissa aut incisa gerente.*

2. H. ORIENTALIS (Gars. exot. t. 19. f. B. ex Lam! dict. 3. p. 92.) foliis radicalibus subtus pubescentibus pedatisectis, floralibus subsessilibus palmatipar-

titis, pedunculis subbifidis, calycinis sepalis ovalibus coloratis. \mathcal{U} in montanis et asperis Orientis. Desf! choix pl. cor. p. 58. t. 45. Species media fere inter *H. nigrum* et *H. viridem*. DC. syst. 1. p. 317. (v. s.)

3. *H. PURPURASCENS*. (Waldst. et Kit. pl. rar. 2. p. 105. t. 101. *) foliis radicalibus subtus subpubescentibus palmatisectis, segmentis basi cuneatis apice 3-5-lobis, caule bifloro, foliis floralibus subsessilibus, calycis sepalis subrotundis coloratis. \mathcal{U} in sylvis Hungariæ, Podoliæ, Vohlynæ. DC. syst. p. 318.

β . *Bocconi*, caule foliis longiore. — Boccon. mus. 2. p. 26. t. 11. f. R. R. — in Apenninis Etruriæ. An species propria?

4. *H. ODORUS* (Waldst. et Kit. ex Willd. enum. p. 592.) foliis radicalibus subtus pubescentibus palmatisectis; segmentis oblongis indivisis basi integerrimis apice serratis, caule bifido, sepalis ovato-oblongis acutiusculis virescentibus. \mathcal{U} in Hungariâ. Affinis *H. purpurascenti* et viridi. DC. syst. 1. p. 318. (v. s.)

5. *H. VIRIDIS*. (Lin! spec. 784.) foliis radicalibus glaberrimis pedatisectis, canlinis subsessilibus palmatipartitis, pedunculis sæpè bifidis, sepalis subrotundo-ovatis viridibus. \mathcal{U} in sylvis et dumetis Europæ. Jacq. fl. austr. t. 106. variat foliis sanguineo-maculatis et foliorum segmentis basi integerrimis. DC. syst. 1. p. 318. (v. v.)

6. *H. ATRO RUBENS* (Waldst. et Kit. pl. rar. 3. p. 301. t. 271. *) foliis radicalibus glaberrimis pedatisectis subtus pallidioribus nitidis, canlinis subsessilibus palmatipartitis, caule subangulato bifidè ramoso, sepalis subrotundis coloratis. \mathcal{U} in sylvaticis et fruticetis Croatiae. DC. syst. 1. p. 319.

7. *H. DUMETORUM*. (Waldst. et Kit. ex Willd. enum. 592.) foliis radicalibus glaberrimis pedatisectis, canlinis subsessilibus palmatisectis, caule tereti bifidè ramoso, sepalis subrotundis viridibus. \mathcal{U} in Hungariâ. Folia cum caule nascentia. DC. syst. 1. p. 320.

§. 3. *Caule folioso multifloro, pedunculis bracteatis.*

8. *H. FÆTIDUS* (Lin! spec. 784.) caule multifloro folioso, foliis pedatisectis glaberrimis, segmentis oblongo-linearibus. \mathcal{U} in pascuis glareosis Europæ occidentalis. Smith. engl. bot. t. 613. Bractæe sunt folia floralia ad petiolum dilatato-membranaceum reductæ. DC. syst. 1. p. 320. (v. v.)

9. *H. LIVIDUS* (Ait. h. Kew. ed. 1. v. 2. p. 272.) caule multifloro folioso, foliis trisectis glaberrimis subtus glaucis, segmentis ovato-lanceolatis. \mathcal{U} in Corsicâ. Curt. bot. mag. t. 72. flores lividi paulò majores et magis patentes quàm subpurpurascens *H. fætidus*. DC. syst. 1. p. 321. (v. v.)

β . *integrilobus*, segmentis foliorum integerrimis. *H. trifolius* Mill. dict. n. 4. non Lin.

XIX. COPTIS. *Salisb. tr. lin. soc.* 8. p. 305.

Calyx 5-6-sepalus, sepalis coloratis petaloideis deciduis. Petala parva cucullata. Stam. 20-25. Capsulæ 6-10 longè stipitatae substellatae membranaceæ ovato-oblongæ stylo acuminatæ 4-6-spermæ. DC. syst. 1. p. 321.

1. *C. TRIFOLIA* (Salisb. tr. soc. Lin. 8. p. 305.) foliis 3-sectis, segmentis obovatis obtusis dentatis vix subtrilobatis, scapo 1-floro. \mathcal{U} in humidis palustribus sylvaticis submontanis regionum borealium Americæ, Asiæ et Europæ. Hel-leborus trifolius Lin! amœn. 2. p. 355. t. 4. f. 18. * spec. 784. DC. syst. 1. p. 322. Big. med. bot. 1. p. 60. t. 5. * (v. s.)

2. *C. ASPLENIIFOLIA* (Salisb! tr. Lin. soc. 8. p. 306.) foliis biternatim sectis, segmentis subpinnatifidis, scapo bifloro. \mathcal{U} ad littora occidentalia Americæ borealis. Thalictum Japonicum Thunb. act. soc. lin. 2. p. 337. Willd. spec. 2. p. 1505. DC. syst. 1. p. 187. ex Smith in Rees cyclop. vol. 19. qui specimina Thunbergiana vidit. DC. syst. 1. p. 322. (v. s. in h. Banks.)

XX. ISOPYRUM. *Lin. gen.* 701. *Gærtn. fruct.* 1. p. 312. t. 65. f. 5.

Calyx 5-sepalus, sepalis petaloïdeis deciduis; petala 5 æqualia tubulata bilabiata, labio exteriori bifido; stamina 15-20, Ovaria 2-20, Styli intus longitudinaliter stigmatosi, capsulæ totidem sessiles 1-loculares oblongæ compressæ membranacæ ∞-spermæ. Semina minuta punctata. DC. syst. 1. p. 323.

1. I. THALICTROIDES (Lin. spec. 783.) capsulis 1-3, sepalis obtusis, radice repente fasciculatâ aut grumosâ, petiolis basi in auriculas membranaceas dilatatis. \mathcal{L} in sylvaticis Pyrenæorum, Delphinatûs, Sabandiar, etc. Jacq. fl. austr. t. 105. Capsulæ ovatæ subcompressæ stylo elongato aristatæ. DC. syst. 1. p. 323. (v. v.)
2. I. GRANDIFLORUM (Fisch! in litt.) capsulis 4-5, scapo radicali unifloro, sepalis ovalibus segmenta foliorum magnitudine multo superantibus. \mathcal{L} in Alpibus Altaicis. Stamina sepalis dimidio breviora (v. s.)
3. I. FUMARIODES (Lin! spec. 783.) capsulis 10-20, sepalis acutis, radice exili simpliciter perpendiculari, petiolis basi non dilatatis. ① in nemoribus Sibiriar, Dahuriar. Schkuhr. handb. 2. n. 153. t. 153. Capsulæ compressæ stylo breviter aristatæ. (v. v.)
4. I. ADOXOIDES (DC. syst. 1. p. 324.) capsulis 2-3, sepalis obtusiusculis, radice tuberosa, petiolis basi non dilatatis in Japoniâ propè Nangasaki. I. capnoides Fisch! in litt. Capsulæ 3-4 oblongæ, acutæ polyspermæ. (v. s. ex Jap.)

XXI. P ENEMION. *Raf. in journ. phys.* 1820. 91. p. 70.

Calyx 5-sepalus petaloïdeus deciduus. Stam. 20-30, filamentis clavatis, antheris subrotundis bilobis. Pistilla 2-6. sæpius 4, stylo longitudine ovarii. Capsulæ 2-6 stellatæ ovatæ compressæ stylis apiculatæ 2-spermæ. Semina ovalia — An satis ab Isopyro differt.

1. E. BITERNATUM (Raf. l. c.) \mathcal{L} in Kentucky ad Lexington. Flos magnitudine Anemones 5-foliar.

XXII. GARIDELLA. *Tourn. inst.* 655. t. 43.

Calyx 5-sepalus deciduus vix petaloïdeus; petala 5 bilabiata bifida; stam. 10-40; ovaria 3 connexa; styli brevissimi; capsulæ 2-3 polyspermæ connatæ in unicam 2-3-locularem vix 2-3 cornutam. DC. syst. 1. p. 325.

1. G. NIGELLASTRUM (Lin. spec. 753.) petalis sessilibus patentibus, staminibus 10-12. ① in apricis inter vineas et oliveta Gallo-provinciar. Sims. bot. mag. t. 1266. Folia tenuissimè divisa ferè Delphinii Consolidæ. DC. syst. 1. p. 325. (v. v.)
2. G. UNGUICULARIS (Lam! ill. t. 379. f. 2.) petalis erecto-conniventibus longè unguiculatis, staminibus 40. ① circâ Aleppum. Ovaria 2-3 connexa ovata erecta scabra. DC. syst. 1. p. 325. (v. s.)

XXIII. NIGELLA. *Tourn. inst.* 258. t. 134. *Lin. gen.* n. 685.

Calyx 5-sepalus coloratus petaloïdeus patens deciduus; petala parva 5-10 bilabiata, ungue foveolato nectarifero. Stam. ∞-Ovaria 5-10 plus minusve basibus concreta, in stylos simplices longos desinentia. Capsulæ totidem plus minusve concretæ, stylis elongatis rostratæ, intus dehiscentes polyspermæ; embryo linearis. DC. syst. 1. p. 326.

Sect. I. NIGELLASTRUM. *Mœnch. meth.* 313 et 311.

Sepala flava; stam. unica seriè disposita; capsulæ basibus præsertim connexæ compressæ; semina plana orbicularia. DC. syst. 1. p. 326.

1. *N. CORNICULATA* (DC. syst. 1. p. 326.) capsulis 3-5 glabris basi imâ connexis divergentibus apice reflexis, stylis extûs arcuatis. ① hab. (v. s. in h. Desf. olim in h. Paris. culta.)
2. *N. CILIARIS* (DC. syst. 1. p. 327.) capsulis 8-10 hispidulis imâ basi connexis stellatim patentibus, stylis rectis. ① in Oriente. Deless! ic. sel. 1. t. 45. (v. s. in h. Deless. et Jus.)
3. *N. ORIENTALIS*. (Lin. spec. 753.) capsulis 5-10 glabris ad medium ferè connexis erectis vix divergentibus, stylis rectis. ① inter segetes circa Aleppum, in Caucasi orient. et Iberiæ mediæ campis. Sims. bot. mag. t. 1264. Semina plana margine membranaceo cincta. DC. syst. 1. p. 327. (v. v. et s.)

Sect. II. NIGELLARIA.

Sepala patentia cœrulescentia, stamina multiplici serie in 8-10 plalanges disposita; semina ovato-angulata.

4. *N. HISPANICA*. (Lin. spec. 753.) antheris apiculatis, stylis 8-10 erectis, capsulis lævibus dorso uninerviis in fructum obconicum ultrâ medium connexis, caule erecto glabro, ramis erectis. ① in arvis Hispaniæ, Bœotiæ, etc. Sims. bot. mag. t. 1265. in herb. Linnæi sub nomine *N. arvensis* — DC. syst. 1. p. 328. (v. v.)
5. *N. FOENICULACEA*. (DC. syst. 1. p. 328.) antheris apiculatis, stylis 5-6 erectis, capsulis lævibus dorso 3-nerviis in fructum obconicum ultrâ medium connexis, caule erecto glabro, ramis subdivaricatis. ① inter segetes Tauriæ. — Moris. hist. 3 p. 516. s. 12. t. 18. f. 6? Semina triquetra. (v. s.)
6. *N. DIVARICATA* (Beaupré in DC. syst. 1. p. 329.) antheris apiculatis, stylis 5 erectiusculis, capsulis lævibus in fructum obconicum ultrâ medium connexis, caule glabro, ramis divaricatissimis. ① in Tauriâ et Ægypto. Deless! ic. sel. 1. t. 46. (v. s.)
7. *N. ARVENSIS* (Lin. spec. 753.) antheris apiculatis, stylis 5-7 extûs circinnatim revolutis, capsulis lævibus in fructum obconicum basi angustatum ultrâ medium connexis, caule glabro, ramis subdivergentibus. ① in arvis inter segetes Europæ mediæ et meridionalis. Schkuhr handb. 2. p. 92. t. 146. DC. syst. 1. p. 329. (v. v.)
8. *N. ARISTATA* (Sibth. et Sm. fl. græc. t. 510* ex Smith prodr. 1. p. 373.) antheris apiculatis, capsulis in fructum turbinatum connexis, caule glabro, floribus involucri folioso cinctis. ① propè Athenas. Habitus ferè *N. Damascenæ*. DC. syst. 1. p. 330. (Descr. ex ic. Sibth.)
9. *N. SATIVA* (Lin. spec. 753.) antheris muticis, capsulis muricatis in fructum ovatum ad apicem usque connexis, caule erecto pilosiusculo, floribus nudis. ① in arvis Monspelii, Barbariæ, etc. Sibth. et Sm. fl. græc. t. 511. DC. syst. 1. p. 330. (v. v.)
 - β. *Cretica*, stylis flore longioribus. — Clus. hist. 2. p. 108.
 - γ. *citrina*, seminibus luteis, floribus plenis. — Moris. hist. 3. p. 516. s. 12. t. 18. f. 5.
 - δ. *Indica*, caule foliis subglabris. *N. Indica* Roxb! mss. ex Flemming in Journ. bot. 1814. 4. p. 203.

Sect. III. EROBATUS.

Sepala alba aut cœrula; stamina ∞ multiplici serie disposita, carpella 5 ad apicem coalita in capsulam 10-locularem, loculis 5 interioribus seminiferis, 5 externis vacuis; involucrium multifidum foliosum sub flore.

10. *N. DAMASCENA*. (Lin. spec. 753.) antheris muticis, carpellis 5 lævibus bilocularibus in capsulam ovato-globosam ad apicem usque connexis, floribus involucri folioso cinctis, sepalis patentibus. ① inter segetes totius regionis Mediterraneæ. Curt. bot. mag. t. 22. Variat sepalis albis cœruleisve, simpliciter multiplicative serie dispositis. DC. syst. 1. p. 331. (v. v.)

II. N. COARCTATA. (Gmel. fl. bad. 2. p. 502. in adn.) antheris muticis, floribus involucreatis, sepalis erecto-conniventibus. ① Hab..... An satis à N. Damascenâ distinctâ? DC. syst. 1. p. 332. (v. v.)

XXIV. AQUILEGIA. Tourn. inst. t. 242. Lin. gen. n. 275.

Calyx 5-sepalus deciduus colorato-petaloidæus; petala 5 supernè hiantia bilabiata, labio exteriori magno plano, interiori minimo, deorsum producta in calcaria totidem cava apice callosa inter sepala exserta. Ovaria 5. Capsulæ totidem erectæ oo-spermæ styli sacuminatæ. DC. syst. 1. p. 333.

1. A. VULGARIS (Lin! spec. 752.) calcaribus incurvis, capsulis villosis, caule folioso multifloro, foliisque glabriusculis, stylis stamina non superantibus. ½ in sylvis subumbrosis Europæ. engl. bot. t. 297. DC. syst. 1. p. 334. (v. v.)

*. *corniculata*, flore pleno, petalis calcaratis, calcaribus deorsum productis. — Clus. hist. 2. 204. f. 1.

** *inversa*, flore pleno, petalis corniculatis, calcaribus sursum directis. — Clus. hist. 2. p. 204. f. 2.

*** *stellata*, flore pleno, petalis planis ecalcaratis coloratis. — Clus. hist. 2. p. 205. f. 1.

**** *degener*, flore pleno, petalis sepalisve planis ecalcaratis virescentibus. Clus. hist. 2. p. 205. f. 2.

2. A. VISCOSA. (Gouan! fl. Mousp. 267. illustr. 33. t. 19.) calcaribus incurvis, capsulis villosis, caule pauci aut unifloro subnudo foliisque pubescenti viscosis, stylis stamina non superantibus. ½ in apricis regionis Mediterraneæ. Vix à priore distincta. DC. syst. 1. p. 336. (v. v.)

3. A. GLANDULOSA (Fisch! ined. Link. enum. 2. p. 84.) calcaribus incurvis petalorum limbo duplò brevioribus, herbæ parte superiore carpellisq. villosoglandulosis. ½ in Alpibus Altaicis. (v. s.)

α. *discolor*, petalis albis, sepalis cæruleis.

β. *concolor*, petalis sepalisque cærulescenti-violaceis.

4. A. SIBIRICA (Lam! dict. 1. p. 150. *) calcaribus incurvis, capsulis glaberrimis, caule 1-2-floro subnudo glabro, sepalis obtusissimis. ½ in sylvis Dahuriæ. A. vulgaris Dahurica. Willd. spec. 2. p. 1246. — Deless! ic. sel. 1. t. 47. DC! syst. 1. p. 336. (v. s.)

5. A. ALPINA (Lin. spec. 752.) calcaribus rectis apice subincurvis limbo petalorum dimidio brevioribus, caule 2-3-floro folioso, foliorum segmentis in lobos lineares profundè multifidis. ½ in locis umbrosis Alpium Pedemontii, Sabaudix, Helvetiæ. Deless! ic. sel. 1. t. 48. A. Alpina var. α. DC. fl. fr. 4. p. 912. syst. 1. p. 336. * (v. v.)

β. *grandiflora*, sepalis ovalibus, calcaribus limbi longitudine. A. grandiflora Patr! ined. An A. glandulosa concolor?

6. A. PYRENAICA (DC. fl. fr. 4. p. 912. syst. 1. p. 337.) calcaribus rectis limbo vix brevioribus, caule subnudo subunifloro, foliorum segmentis in lobos lineares profundè multifidis. ½ in pratis elatis Pyrenæorum et Apennini Appiani. A. viscosa Trev. delph. p. 23. t. 2. A. Alpina var. β. DC. fl. fr. 4. p. 912. * (v. v.)

7. A. CANADENSIS (Lin! spec. 752.) calcaribus rectis, stylis staminibusque exsertis, sepalis subacutis petala paulò superantibus, foliorum segmentis tripartitis apice subobtusis inciso-dentatis. ½ in Americâ boreali. Curt. bot. mag. t. 246. herb. amat. t. 305. * DC. syst. 1. p. 337. (v. v.)

8. A. FORMOSA (Fisch! in litt. cum ic.) calcaribus rectis limbo petalorum brevissimis staminibusque multò longioribus, stylis non exsertis, sepalis lanceolatis petalis multò longioribus. ½ in Kamchatkâ. Affinis A. Canadensi et habitu et floris colore et formâ. (v. s. comm. à cl. Fisch.)

9. A. VIRIDIFLORA. (Pall. act. petrop. 1779. p. 260. t. 11.) calcaribus rectis

limbo longioribus staminibus petala æquantibus, stylis longè exsertis, sepalis ovali-oblongis petalis brevioribus. \mathcal{L} in Sibiria. Jacq. ic. rar. 1. t. 102. A. flava Lam! dict. 1. p. 149* excl. syn. DC. syst. 1. p. 338. (v. v.)

10. A. ATROPURPUREA. (Willd. enum. 577.) calcaribus rectis limbo æqualibus, stylis staminibusque sepalis subæquantibus, sepalis longitudine petalorum. \mathcal{L} in Sibiria. DC. syst. 1. p. 338. mem. soc. h. n. Gen. 1. p. 455. (v. s. com. à cl. Fisch.)

a. *brevistyla*, (Willd. l. c.) stylis stamina non superantibus.

β. *Dahurica*, (Patr. in Deless! ic. sel. 1. t. 49.) stylis longè exsertis, foliis glabris.

γ. *Fischeriana*, stylis longè exsertis, foliis subtus villosis. A. atropurpurea Fisch!

11. A. HYBRIDA. (Sims. bot. mag. t. 1221.) calcaribus rectis vix apice subincurvis limbo obtusissimo longioribus, stylis stamina petalque vix superantibus, sepalis acutis longitudine petalorum, caule foliisque subtilissimè pubescentibus. \mathcal{L} in Sibiria. A. speciosa var. a. DC. syst. 1. p. 336. A. hybrida DC. syst. 1. p. 339.* (v. s.)

12. A. PARVIFLORA (Ledeb. act. petrop 1815. 5. p. 514. n. 32.) calcaribus rectis brevibus limbo obtuso subæqualibus, genitalibus recurvis sepalorum longitudine, caule 2-3-floro foliisque glabris. \mathcal{L} in Sibiria. — Gmel. fl. sib. 4. p. 186. t. 74. — DC. syst. 1. p. 339.* (v. s.)

13. A. ANEMONOIDES (Willd. gess. naturf. Berl. 1811. t. 9. f. 6.) calcaribus rectis brevissimis limbo æqualibus, petalis calyce triplò brevioribus, pedunculis radicalibus unifloris subnudis. \mathcal{L} in Sibiria Altaicâ. An var. A. glandulosæ ex Fisch. — DC. syst. 1. p. 339

XXV. DELPHINIUM. Tourn. inst. 426. t. 241. Lin. gen. n. 681.

Calyx deciduus petaloideus irregularis, sepalo nempè superiore in calcar deorsum producto. Petala 4; 2 superiora basi in appendicibus intrâ calcar contentis producta. DC. syst. 1. p. 340.

Sect. I. CONSOLIDA. DC. syst. 1. p. 341.

Ovarium 1. Petala 4 in 1 coalita; calcar internum 1-petalum. Species annuæ.

1. D. AXILLIFORUM. (DC. syst. 1. p. 341.*) caule subsimplici subglabro, floribus axillaribus subsessilibus, bracteolis sub flore sitis multifidis. ① in Oriente. Deless! ic. sel. 1. t. 50. Sepala pallidè cærulescentia oblonga acuta. (v. s. in h. Labill.)

2. D. OLIVERIANUM (DC. syst. 1. p. 341.*) caule glabro subramoso, ramis vix divergentibus, floribus paucis laxè racemosis, pedicellis bractæ longitudine, capsulis glabris. ① in agris cultis inter Bagdad et Kermancha. Deless! ic. sel. 1. t. 51. Flores cæruleo-purpurei, ampli, D. Ajacis floribus duplò majores. (v. s. in h. Olivier.)

3. D. AJACIS. (Lin! spec. 748.) caule erecto subglabro subsimplici, ramis vix divergentibus longe floriferis, floribus densè racemosis, pedicellis bractæ longitudine, capsulis pubescentibus. ① in Tauria. — Clus. hist. 2. p. 206. f. 1. DC. syst. 1. p. 341. Variat floribus nempè sepalis albis, cæruleis, roseis, incarnatis, purpureis, et ex iis coloribus variegatis. (v. v.)

4. D. CONSOLIDA (Lin! spec. 748.) caule erecto subglabro divaricatim ramoso, floribus paucis laxè racemosis, pedicellis bractæ longioribus, capsulis glabris. ① in agris Europæ. OEd. fl. dan. t. 683. Variat omnino ut D. Ajacis. DC. syst. 1. p. 343. (v. v.)

5. D. PUBESCENS (DC. fl. fr. 5. p. 641*. syst. 1. p. 343.) caule recto pubescente apice divaricatim subramoso, floribus laxè racemosis, pedicellis bractæ longioribus, foliis capsulisque pubescentibus. ① in agris regionis Gallo-Mediterranæ. — Sibth. et Sm. fl. græc. t. 504. D. ambiguum. Lois! not. 85. non Lin. (v. v.)

6. *D. RIGIDUM*. (DC. syst. 1. p. 344.) caule erecto ramosissimo rigido, foliis capsulisque velutino-canescens, pedicellis bractea triplo longioribus. ① in Oriente. Deless! ic. sel. 1. t. 52. Medium inter *D. pubescentem* et *pusillum*. (v. s. in h. Labill. et Deless.)
7. *D. FUSILLUM* (Labill. Syr. dec. 4. p. 5. t. 2. f. 2.*) caule erecto tenuissimo 2-4-floro foliisque pubescentibus subpedatim 5-fidis, pedicellis bractea lineari paulo longioribus. ① propè Damascum. *D. pygmaeum* Poir! suppl. 2. p. 458*. Calcar subulatum rectum. — DC. syst. 1. p. 344. (v. s.)
8. *D. TENUISSIMUM* (Sibth. et Sm! fl. græc. t. 505. prod. 1. p. 370.) caule erecto gracili subramoso, apice subpubescente, pedicellis bractea subulata multo longioribus. ① in Græcia circa Athenas. Herba tenuis gracilis vix palmaris. DC. syst. 1. p. 345.* (v. s. in h. Banks.)
9. *D. ACONITI* (Lin! mant. 77*) caule erecto ramosissimo subpubescente, foliis pedato-multifidis, pedicellis longissimis, calcare apice incurvo horizontali supernè fisso. ① ad Dardanellas. Vahl! symb. 1. p. 40. t. 13*. *Aconitum monogynum* Forsk. cat. pl. Const. p. 27. n. 248 ex Vahl. Statura ferè *D. consolidæ*. DC. syst. 1. p. 345.* An ab hac satis differt *D. incanum* Clark. in Spreng. neu. eutd. 3. p. 162? (v. s.)
10. *D. EXSERTUM* (DC. syst. 1. p. 345.*) caule erecto ramoso glabro, foliis 3-sectis, segmentis petiolulatis palmato-multifidis, pedicellis bracteis vix longioribus, calcare recto, staminibus exsertis. ① in Oriente. Deless! ic. sel. 1. t. 53. (v. s.)
11. *D. FLAVUM* (DC. syst. 1. p. 346.*) caule erecto ramoso supernè pubescente, foliis palmato-multipartitis, pedicellis distantibus floris longitudine, calcare recto patente, staminibus subexsertis, capsula cylindracea pubescente. ① in Oriente. Deless! ic. sel. 1. t. 54. (v. s. in h. Oliv.)
- a. velutinum*, totum pubescens, bracteolis in medio pedicelli.
β. glaucum, infernè glabrum, bracteolis ad basin pedicelli.
- Sect. II. DELPHINELLUM. DC. syst. 1. p. 346.
- Ovaria 3. Petala libera glabra; calcar interdum 1-petalum. Species annuæ.
12. *D. AMBIGUUM* (Lin! spec. 749.*) caule erecto subvelutino, foliis 3-5-partitis, partitionibus pinnatifidis, racemis laxis, calcare recto pubescente calyce breviorè, ovaris villosis. ① in Mauritania. DC. syst. 1. p. 346.* (v. s. in h. Lin.)
13. *D. CARDIOPETALUM* (DC. syst. 1. p. 347*) caule erecto subramoso, foliis glabris membranaceis 3-sectis, segmentis oo-fidis, lobis linearibus, rameis bracteisque inferioribus multifidis, racemis confertis, petalis stipitatis, limbo orbiculato basi cordato. ① in Pyrenæorum vallibus. *D. peregrinum* Lam! dict. 2. p. 264.* (v. v.)
14. *D. GRACILE* (DC. syst. 1. p. 347*) caule erecto subramoso, foliis subglabris rigidulis 3-sectis oo-fidis, lobis linearibus, rameis bracteisque linearibus, racemis laxis, petalis stipitatis limbo ovato basi cordato. ① (v. s.)
- a. glabrum*, caule foliisque glabris. — in Andalusia.
β. velutinum, caule foliisque velutino-pubescentibus. circa Aranjuez. Pourr.
15. *D. JUNCHEUM* (DC. fl. fr. 5. p. 641. syst. 1. p. 348*) caule erecto ramosissimo, foliis glabris rigidis, inferioribus multifidis, rameis bracteisque linearibus integris, racemis laxis, petalis stipitatis limbo elliptico. ① in asperis apricis et agris Italiae, etc. All! pedem. n. 1508. t. 25. f. 3. (v. s. et v.)
- β. subvelutinum*, caule foliisque velutino-pubescentibus. — in monte Libano.
16. *D. NANUM* (DC. syst. 1. p. 349*) caule suberecto nano foliisque velutino-pubescentibus, inferioribus 3-5-lobatis, superioribus linearibus, racemis laxis 2-3-floris, petalis inferioribus limbo orbiculato. ① circa Alexandriam. *D. peregrinum* Delile! ill. fl. æg. 17. non Lin. Affine var. *β. D. junciei*. (v. s.)

17. *D. HALTERATUM* (Sibth. et Sm. fl. græc. t. 507.) caule erecto ramoso, foliis glabris oo-partitis, floralibus inferioribus 3-fidis, summis integris, racemis laxis, petalis patentibus, inferioribus stipitatis limbo orbiculato (I) in Sicilia DC. syst. 1. p. 349.* (Descr. ex. ic. Sibth.)
18. *D. VIRGATUM* (Poir! suppl. 2. p. 458*.) caule erecto à basi ramoso, foliis glabris, inferioribus 3-fidis, lobis 3-dentatis, rameis floralibusque integris acutis, racemis laxis, petalis inferioribus stipitatis limbo ovato. (I) in Syria. Deless! ic. sel. 1. t. 55. DC. syst. 1. p. 349* (v. s. in h. Desf.)
19. *D. MACROPETALUM*. (DC. syst. 1. p. 350*.) caule erecto ramosissimo, foliis glabris membranaceis, inferioribus palmato-multifidis, summis linearibus acutissimis, racemis laxis, petalis inferioribus stipitatis limbo ovato. (I) in Africa boreali. Deless! ic. sel. 1. t. 56. (v. s.)
20. *D. OBCORDATUM* (DC. syst. 1. p. 350*.) caule erecto ramoso, foliis glabris, caulinis 3 lobatis, lobis subbifidis, rameis floralibusque integris acutis, racemis laxis, petalis inferioribus stipitatis limbo obcordato. (I) in Barbaria. (v. s. in h. Bouch.) †.
21. *D. TRIBRACTEOLATUM* (DC. syst. 1. p. 350*.) caule erecto ramoso, foliis inferioribus palmato-lobatis, racemis laxis, pedicellis bracteolas 3 gerentibus, petalis inferioribus stipitatis limbo orbiculato basi cordato. (I) in Barbaria. Calcar rectum 4. lin. longum. (v. s. in h. Bouchet.)

SECT. III. DELPHINASTRUM. DC. syst. 1. p. 351.

Ovaria 3-5; petala libera, inferiora in disco barbata bifida; calcar elongatum internum dipetalum. Species perennes.

§. 1. *Petalorum inferiorum limbo integro.*

22. *D. GRANDIFLORUM*. (Lin! spec. ed. 1. p. 531. ed. 2. p. 749.) foliis palmatim oo-partitis, lobis linearibus distantibus, pedicellis bractea longioribus, petalis calyce brevioribus, 2 inferioribus limbo obliquè inflexo ovato integro. 2/ in Sibiria. Sims. bot. mag. t. 1686. — Gmel. fl. sib. 4. p. 187. t. 78. DC. syst. 1. p. 351.* (v. s.)

ℓ. *Chinense*, Fisch! in litt. Lodd. bot. cab. t. 71. — An duæ species hic confusæ? à *D. grandifloro* distinguitur *D. Chinense* caule rectiore rigido floribus facilius versicoloribus florescentiâque seriore. (v. v.)

23. *D. CHEILANTHUM* (Fisch. in litt.) caule erecto ramoso, foliis 5-partitis, lobis oblongis acuminatis subtrifidis subdentatis, petalis calyce brevioribus, 2 inferioribus limbo obliquè inflexo ovato integro, capsulis reticulatim pictis pubescentibus. 2/ in Dahuria. — Schrank. pl. rar. 52. ic. — *D. Dahuricum* Stev! ined. — Gmel. fl. sib. 4. p. 187. t. 76. ex Fisch. — DC. syst. 1. p. 352* (v. s.)

24. *D. VIRESCENS* (Nutt. gen. amer. 2. p. 14*.) caule simplici pubescente, foliis longè petiolatis 3-5-partitis, segmentis linearibus subtrifidis, racemo paucifloro, calcare flore longiore. 2/ in planitiibus Missouriensibus.

§. 2. *Petalorum inferiorum limbo bifido.*

25. *D. PUNICEUM* (Pall! voy. 8. p. 327. u. 336. Lin. f. suppl. 267*.) petiolis basi dilatato-vaginantibus, foliis in lobos lineares ad basin usque oo-partitis, racemo elongato, calcare recto obtuso pedicello sublongiore flore brevior. 2/ in aridissimis deserti Tatarici circa Volgam. DC. syst. 1. p. 352. (v. s.)

ℓ. *Damæcornu*, segmentis petiolulatis, lobis lanceolatis divergentibus. — Gmel. fl. sib. 4. t. 77? — videtur è priore à cultura orta?

26. *D. ALBIFLORUM* (DC. syst. 1. p. 353*.) petiolis basi dilatato-vaginantibus, foliis oo-partitis, lobis linearibus, racemo elongato conferto, bracteis membranaceis lato-lanceolatis, calcare recto obtuso pedicello sublongiore. 2/ in Armenia. Deless! ic. sel. 1. t. 58. Flores *D. puniceo* similes, sed albi extus glabri. (v. s. in h. Mus. Par.)

27. *D. OCHROLEUCUM* (Stev. diss. ined. DC. syst. 1. p. 353 et 546.) petiolis basi dilatato-vaginantibus, foliis oo-partitis, lacinii linearis-subulatis, racemo brevi, floribus (albis) extus pubescentibus, calcare flore longiore acuto, capsulis glabris. \mathcal{U} In Iberiâ. *D. puniceum* var. Bieb. fl. taur. 2. p. 13. (v. s.)
28. *D. HYBRIDUM* (Willd. spec. 2. p. 1229*.) petiolis basi dilatato-vaginantibus, foliis oo-partitis, lobis linearibus, racemo conferto, calcare recto flore longiore. \mathcal{U} In Tauriâ et Caucaso. *D. hirsutum* Pers. ench. 2. p. 82. — DC. syst. 1. p. 353. (v. s.)
ℓ. D. fissum (Waldst. et Kit. pl. rar. hung. 1. p. 83. t. 81*.) racemo elongato. — in Hungariâ.
29. *D. VULUTINUM* (Bertol. exc. p. 12.) petiolis basi dilatato-vaginantibus, foliis 5-partito-multifidis, segmentis linearibus, racemo laxo cauleque mollioribus villosis, calcare curvo, bracteis inferioribus flore longioribus. \mathcal{U} in Italiâ montibus. *D. hybridum* γ . DC. syst. 1. p. 554.
γ. D. orientale perenne aconiti folio flore cœruleo. Tourn! cor. 3o. in Armeniâ. (v. s.)
30. *D. PENTAGYNUM* (Lam! dict. 2. p. 264*.) petiolis basi vix dilatatis, foliis inferioribus 5-lobis, lobis apice subobtusis, superioribus 5-partitis in lobos lineares oo-fidis, petalis calice brevioribus. \mathcal{U} inter segetes Algeriæ. Desf! fl. atl. 1. p. 427. t. 111*. DC. syst. 1. p. 354* (v. s. et v.)
31. *D. MENSIESII* (DC. syst. 1. p. 355*.) petiolis basi vix dilatatis, foliis 5-partitis, lobis 3-fidis linearibus integris, bracteis 3-fidis, radice grumosa. \mathcal{U} in Novâ-Georgiâ. Flores cœruleo-violacei magnitudine *D. grandiflori*, calcar rectum. (v. s. in h. Banks.)
32. *D. ELEGANS* (DC. syst. 1. p. 355*.) petiolis basi vix dilatatis, foliis glabris 5-partitis, lobis 3-5-fidis, lobulis lanceolato-linearibus acutis, racemo laxo paucifloro, petalis calyce brevioribus, calcare curvo sepalis brevioribus. \mathcal{U} in Americâ boreali? Habitus ferè *D. grandiflori* sed structura floris diversa. (v. s. in h. Delile.)
33. *D. AMOENUM* (Stev! ined. DC. syst. 1. p. 546*.) petiolis basi vix dilatatis pilosiusculis, foliis subtus pubescentibus 5-partitis, lobis lanceolatis pinnatifidis, lobulis linearibus acutis, racemo ramuloso, petalis calyce brevioribus, calcare recto. \mathcal{U} in Sibirîâ. — Gmel. fl. Sib. 4. p. 187. t. 77. excl. syn. Amman. ex Stev. (v. s.)
34. *D. TRICORNE* (Michx! fl. bor. amer. 1. p. 314.) petiolis basi glabris vix dilatatis, foliis 5-partitis, lobis 3-5-fidis, lobulis linearibus, petalis calyce brevioribus, capsulis à basi patulo-reflexis arcuatis. \mathcal{U} in montibus à Carolinâ ad Virginiam. Deless! ic. sel. 1. t. 59. DC. syst. 1. p. 356* (v. s. in h. Mus. Par.)
ℓ. multiflorum, majus pubescens, floribus 15-20 dense racemosis. — in Louisiana superiore. (v. s. in h. Lamb.)
35. *D. AZURRUM* (Michx! fl. bor. amer. 1. p. 314.) petiolis basi vix dilatatis, foliis 3-5-partitis oo-fidis, lobis linearibus, racemo stricto, petalis omnibus apice barbatis, inferioribus villosissimis. \mathcal{U} in Americâ. Deless! ic. sel. 1. t. 60. DC. syst. 1. p. 356. (v. s. in h. Mus. Par.)
36. *D. EXALZATUM* (Ait. h. Kew. ed. 1. v. 2. p. 244.) petiolis basi non dilatatis, foliis planis ultra medium 3-fidis, lobis cuneiformibus apice 3-fidis acuminatis, lateralibus septè bilobis, racemo stricto, calcare recto longitudine calycis. \mathcal{U} in Americâ. DC. syst. 1. p. 35. * *D. tridactylum* Michx! fl. bor. amer. 1. p. 314. (v. s. in h. Mus. Par. et Rich.)
37. *D. URCEOLATUM* (Jacq! coll. 1. p. 153* ic. rar. 1. t. 101.) petiolis basi non dilatatis, foliis concavis ultra medium 3-fidis, lobis cuneiformibus apice incisis acuminatis, racemo stricto, calcare recto calyce sublongiore. \mathcal{U} in Americâ boreali? Sims. bot. mag. t. 791. DC. syst. 1. p. 357* (v. s. in h. Bosc.)
38. *D. REVOLUTUM*. (Desf! cat. h. Par. ed. 2. p. 148.) petiolis basi non dilatatis,

- foliis orbiculatis cordatis 5-fidis, lobis incisibus acutis deflexis, pedicellis 3-bracteatis, ovario glabris. $\frac{1}{2}$ hab. DC. syst. 1. p. 357.* (v. v. in h. Paris.)
39. *D. MESOLEUCUM* (Link. enum. h. Ber. 2. p. 80.) foliis basi subdilatatis, laciniis cuneiformibus anticè serrato-incisis, caule supernè pedunculisque pubescentibus. $\frac{1}{2}$ patr. ign. — folia *D. exaltati*. calyx cœruleus; petala dilute flava.
40. *D. PALMATIFIDUM* (DC. syst. 1. p. 358.*) petiolis basi non dilatatis, foliis ciliatis basi subtruncatis 5-fidis, lobis apice incisibus, superioribus 3-lobatis, lobis 3-fidis integris, nervis utrinque pilosiusculis, bracteolis capsulis calycibusque glabris. $\frac{1}{2}$ in Sibiria. (v. v.)
- a. *hispidum*, caule pilis patentibus hispidum, pedicellis subglabris. — Gmel. fl. sib. 4. p. 187. t. 79?
- β. *glabellum*, caule subglabro, pedicellis pilis patentibus hispidulis. — Gmel. fl. sib. 4. t. 75?
41. *D. INTERMEDIUM* (Ait. h. Kew. ed. 1. v. 1. p. 243.) petiolis basi non dilatatis, foliis cordatis 5-7-fidis, superioribus 3-lobis, omnium lobis inciso-serratis, pedicellis bracteolis calycibus ovarisque glabris. $\frac{1}{2}$ in vallibus Alpibus, Silesiæ, Hungariæ, Helvetiæ et Pyrenæorum. DC. syst. 1. p. 358.* (v. v.)
- a. *pilosissimum*, caule pilosissimo, foliis villosis, nervis supernè pilosis, racemo ramoso. — Gmel. fl. sib. 4. p. 167. t. 80?
- β. *Alpinum*, Waldst. et Kit. pl. rar. Hung. 3. p. 273. t. 246*. caule subglabro, foliis pubescentibus, racemo ramoso. — Clus. hist. 2. p. 94. f. 2.
- γ. *leptostachyum*, caule petiolisque glabris, foliis pubescentibus, racemo simplici brevi. *D. intermedium*. Lapeyr. abr. 364?
- δ. *ranunculifolium*, caule glabro, petiolis pilosiusculis, foliis tenuibus pubescentibus, racemo simplici laxo.
- ε. *laxum*, caule folisque glabris, lobis elongatis acutius incisibus, racemo laxo ramoso. — Mill. ic. t. 119?
42. *D. CUNEATUM* (Stev! diss. ined. DC. syst. 1. p. 359.*) petiolis basi non dilatatis, foliis basi cuneatis 5-7-lobatis, lobis incisibus acutis, racemo laxo ramoso, bracteolis calycibusque glabris, capsulis subpubescentibus. $\frac{1}{2}$ *D. hybridum*. Lin. herb. *D. Cuneatum*. Deless! ic. sel. 1. t. 61. (v. v.)
- β. *pubiflorum*, pedicellis bracteolis calycibus imò imprimis capsulis junioribus pube densè tenui velutinis. — circa Saratof. (Stev.)
43. *D. VILLOSUM* (Stev! in litt. h. Dorpat.) petiolis basi vix dilatatis, foliis glabris basi cuneatis 3-5-lobatis, lobis incisibus acutis, racemo laxo, pedicellis bracteatis capsulisque villosis, calcare recto. $\frac{1}{2}$ hab. medium inter *D. cuneatum* et *D. dyctiocarpum*. DC. syst. 1. p. 546.* (v. a.)
44. *D. DYCTIOCARPUM* (DC. syst. 1. p. 360.*) petiolis basi non dilatatis, foliis 3-7-lobatis, lobis oblongis acutis inciso-pinnatifidis, superioribus subtripartitis, lobis angustis subintegris, racemo laxo ramoso, bracteolis calycibusque glabris, capsulis reticulatis carinà et margine ciliatis. $\frac{1}{2}$ in Sibiria. Habitus *D. elati* aut *cuneati*, capsula fere *D. Cheilanthes*. (v. v. in h. Monsp.)
45. *D. LAXIFLORUM* (DC. syst. 1. p. 360.*) petiolis basi non dilatatis, foliis 3-7-lobatis, lobis oblongis acutis inciso-pinnatifidis, superioribus subtripartitis, lobis angustis integris, racemo laxo ramoso, bracteolis ovarisque pubescentibus. $\frac{1}{2}$ in Sibiria.
46. *D. MONTANUM* (DC. fl. fr. 5. p. 641*. syst. 1. p. 360.*) petiolis basi non dilatatis, foliis pubescentibus 5-lobatis, lobis basi cuneatis trifidis incisibus, racemo simplici, bracteis calycibus capsulisque pubescentibus, calcare inflexo. $\frac{1}{2}$ in Pyrenæis et Alpibus Galloprovinciæ Pedemontis et Helvetiæ. *D. intermedium*. Lois. not. p. 86. (v. v.)
- β. *bracteosum*, bracteis inferioribus flore longioribus, petalis 4 omnibus apice barbatis. — in Alpibus Barcinouensibus. prioris var. *luxurians*? (v. a.)
47. *D. DASYCARPUM* (Stev! ined. — DC. syst. 1. p. 547.*) petiolis basi non

dilatatis, foliis pubescentibus 5-lobatis, lobis lanceolatis subtrifidis apice inciso-dentatis, racemo simplici pubescente, bracteis pedicellò triplò brevioribus, sepalis subvillosis petala duplò superantibus, calcare recto. 2/ circa Nartschan in Caucaso cisalpino. (v. s.)

48. *D. SPECIOSUM* (Bieb. fl. taur. cauc. 2. p. 12*.) petiolis basi non dilatatis, foliis pubescentibus 5-lobatis, lobis inciso-serratis, bracteis lanceolatis villosis-viscosis, calcare curvulo, capsulis glabris. 2/ in regione Caucasicà. Deless! ic. sel. 1. t. 62. DC. syst. 1. p. 361*.

49. *D. FLEXUOSUM* (Bieb. fl. taur. cauc. 2. p. 12*.) petiolis basi non dilatatis, foliis 5-lobatis, lobis incisis, caule flexuoso petiolisque pilosiusculis, bracteis linearibus, capsulis glabris. 2/ in Caucaso Iberico. Trev. delph. obs. p. 15. t. 1 et t. 2. f. a. b. c. d. — *D. ciliatum* suppl. enum. h. Dörp. 1811. ex Stev. — DC. syst. 1. p. 362*.

50. *D. TRISTE* Fisch! in litt. — DC. syst. 1. p. 362*.) petiolis basi non dilatatis, foliis 3-5 partitis, lobis angustis inciso-subpinnatifidis acutis, summis 3-partitis lobis integris, racemo laxo, capsulis pubescentibus. 2/ in Dahurià et Sibirià. *D. obscurum* Stev. diss. ined? (v. s.)

SECT. IV. STAPHISAGRIA. *C. et J. Bauh.*

Ovaria 3-5. Petala libera, calcar breve, internum dipetalum. Capsulæ ventricosæ; semina pauca magna. Species biennes. DC. syst. 1. p. 362.

51. *D. BEQUIENII* (DC. fl. fr. 5. p. 642*.) syst. 1. p. 362*.) calcare calycis fere longitudine, bracteolis medio pedicelli insertis, petiolis pilosis. ② in insulis Stæchadum. Deless! ic. sel. 1. t. 63. (v. s.)

52. *D. PICTUM* (Willd. enum. 574*.) calcare calycis vix brevior, bracteolis ad basin pedicelli insertis, petiolis pubescentibus, pedicellis flore vix longioribus. ② in Europà australi? *D. Staphysagria* var. B. Lam! dict. 2. p. 265. — Woodv. med. bot. t. 154. DC. syst. 1. p. 363* (v. v. et s.)

53. *D. STAPHYSAGRIA* (Lin. spec. 750.) calcare brevissimo, bracteolis ad basin pedicelli insertis, petiolis pilosis, pedicellis flore duplò longioribus. ② in ruderatis Teneriffæ, Monspelii, etc. Sibth. et Sm. fl. græc. t. 508. DC. syst. 1. p. 364. (v. v.)

XXVI. *ACONITUM*. Tourn. inst. 1. p. 424. t. 239 et 240. Lin. gen. n. 682. DC. syst. 1. p. 364. Ser. mus. helv. 1. p. 115. (1)

Calyx petaloideus irregularis deciduus vel marcescens, sepalò nempe superiore concavo cassidæformi. Petala 2 superiora (cuculli, nectaria) longè unguiculata apice in saccum expansa sub galeâ recondita. — Folia palmatisecta. DC. syst. 1. p. 364.

Sect. 1. *ANTHORA*. DC. syst. 1. p. 364. Ser. mus. helv. t. 15. f. 1: 2, 3, 4 et 47.

Sepala persistentia, galea conica vel subsemicircularis, ovaria 5. — Folia in lobos lineares palmatisecta. flores ochroleuci quandoque cæruleo-variegati; radix napiformis.

1. *A. ANTHORA* (Lin. spec. 751.) floribus paniculatis, sepalis petalisque persistentibus, sacco cucullorum fere nullo, calcare crasso spirali, labio longissimo, staminum alis evanescentibus, foliis multifidis, lacinia linearibus acutis, folliculis viridi-atris (glabris vel pilosis), seminibus. 2/ in Europà. Ser. mus. helv. 1. p. 129*.

a. *vulgare*, (Ser. mus. helv. 1. p. 130. t. 15. f. 1. 3. 4 et 47*.) paniculâ floribus fructibusque pubescentibus, galeâ subconicâ suprâ rostrum vix emarginatâ, floribus luteis. 2/ in Caucaso, Pyrenæis et Jurasso. (v. v.)

(1) *Genus elaboratum à G. Sæmiger.*

- A. *Anthora* β . DC. syst. 1. p. 366. Reichenb. acon. 61. t. 1. f. A. Ac. Pallasii Reichenb. acon. 72. t. 6. f. A. B. (v. s. in herb. DC. et Ser.)
- β . *grandiflorum* (Reichenb. ueb. p. 15. acon. p. 63. t. 1. f. B.), paniculâ floribus fructibusque pubescentibus, galeâ subconicâ amplâ, floribus luteis. — in Jurasso. Ser! mus. helv. 1. p. 130. (v. s. in herb. Ser.)
- γ . *Jacquinianum* (Ser. mus. helv. 1. p. 131.) floribus glabriusculis, galeâ sub conicâ inrostrum elongatum productâ, floribus luteis. — in Austriâ et Pyrenæis. A. anthora Jacq. fl. austr. t. 382. A. Jacquini Reichenb. ueb. p. 17. acon. 65. t. 2. *
- δ . *latilobum* (Ser. mus. helv. 1. p. 131.) paniculâ floribusque pubescentibus, galeâ subconicâ inclinatâ, rostro brevi, lobis foliorum latis, floribus luteis. — in Sibiriâ, A. nemorosum Rieb. ex Reichenb. acon. p. 71. t. 6. f. * A. Decandollii. Reichenb. ueb. 16. acon. 67. t. 3. (v. s. in herb. DC.)
- ϵ . *inclinatum* (Ser. mus. helv. 1. p. 131. t. 15. f. 2.) paniculâ floribus fructibusque pubescentibus, galeâ elatâ conicâ suprâ rostrum inclinatâ, floribus luteis. — in Jurasso. A. anthoroideum Reichenb. acon. 68. t. 4. non DC. — A. anthora Koell. spic. 24. (v. s. in herb. Ser.)
- ζ . *eulophum* (Ser. mus. helv. 1. p. 131.), paniculâ floribusque puberulis, galeâ conicâ suprâ rostrum incumbente, floribus luteis. — in Caucaso et Jurasso. A. anthoroideum DC! syst. 1. p. 366 et herb. Lamb! A. eulophum Reichenb. ueb. 15. acon. p. 69. t. 5. * (v. s. in herb. DC. et Ser.)
- η . *versicolor* (Stev! in herb. DC.), floribus glabriusculis luteis cœruleo-variegatis, galeâ humili subconicâ. — ex Iberiâ. Ser. mus. helv. 1. p. 132. * *Anthora versicolor* Stev! iped. (v. s. comm. à cl. Stev.)
- θ *multicucullatum* (Ser. mus. helv. 1. p. 132. t. 16. *), petalis lateralibus cucullatis, sepalis lateralibus galeatis. — in Jurasso. (v. s. in h. Oth.)

Sect. II. LYCOCTONUM. DC. syst. 1. p. 367. Ser. mus. helv. t. 15. f. 5. 6. 7. 8. 9. 10. 11. 48.

Sepala decidua, galea conico-cylindracea; ovaria 3. — Lobi foliorum cuneiformes pinnati rarò bipinnati-secti, radix fibrosa; flores ochroleuci, albi rarò purpurei aut variegati.

2. A. LYCOCTONUM (Lin! spec. 750.) galeâ conico-cylindraceâ, calcare gracili spiraliter contorto, labio divaricato, filamentis staminum basi alatis, alis submuticis, foliis palmatis ultrâ medium 3-5 lobatis, lobis cuneatis trifidis nervis frequenter anastomosantibus, seminibus 3-quetris transversim rugosis. \mathcal{V} in Europâ. Ser. mus. helv. 1. p. 132. *

α . *vulgare* (Ser. mus. helv. t. 15. f. 5. 6. 48. *) pilosum, floribus spicatis vel paniculatis luteis, caule pedunculisque pilis arcuatis obsitis, galeâ conico-cylindraceâ compressâ, ovarii glabriusculis, foliis ciliatis. — in Alpibus et Jurasso. Gaertn. de fruct. 1. p. 311. t. 65. DC. syst. 1. p. 369. var. α . (v. v.)

β . *Pyrenaicum* (Ser. mus. helv. 1. p. 133. *), totum densè velutino-pubescentis, pilis caulinis pedunculorumque arcuatis, floribus paniculatis luteis numerosis, galeâ conico-cylindraceâ compressâ, calcare laxè spiraliter contorto, foliis amplissimis ovarisque pubescentibus. — in Pyrenæis. A. Pyrenaicum Lam. dict. 1. p. 33. * DC! syst. 1. p. 368. (v. s. in herb. DC. et Ser.)

γ . *grandiflorum* (Ser. mus. helv. 1. p. 134.), glabriusculum, floribus paniculatis luteis, galeâ magnâ apice subinflatâ, ungue cucullorum filamentisque staminum glabris vel pilosis, ovarii glabris, foliis ciliatis. — in Alpibus. A. Lycotonom glabriflorum. DC! syst. 1. p. 269. A. tragocotnum, A. theriophonum, A. lupicida, A. theliophonum, A. vulparia Reicheub. loc. cit. (v. s. in herb. DC. et Ser.)

δ . *Penninum* (Ser. plant. sel. cent. 1. et mus. helv. 1. p. 134.), pubescens, floribus magnis spicatis vel subpaniculatis luteis, galeâ conicâ, rostro

magno, ovariis 3-5 villosulis. — in Monte Pennino. (v. s. in herb. Ser. comm. à Rever. Darbellay et Lamou.)

1. *puberulum* (Ser. mus. helv. r. p. 134.), floribus spicatis vel subpaniculatis luteis, galeâ magnâ, caule foliis floribusque pilis horizontalibus tectis. — in Monte-Cenisio, Stokhorn. (v. s. in herb. DC. et Ser.)

2. *altissimum* (Ser. mus. helv. r. p. 135.) floribus paniculatis numerosis luteis, galeâ magnâ apice subventricosâ, caulibus floribusque glabrin-sculis, bracteis elongatis, foliis plus minusve dissectis. — A. lycocto-num altissimum. DC. syst. r. p. 369? A. rectum. Reichenb. ueb. 59. A. cynoctonum Reichenb. ueb. 73. — in Galliâ, propè Mimatem. (v. s. in herb. DC.)

3. *laxiflorum* (DC! syst. r. p. 369.), paniculâ amplâ, ramis divergentibus, floribus laxis pubescentibus, germinibus villosis, lobis foliorum sub-quadratis divaricatis. — Ser. mus. helv. r. p. 135. (v. s. ex hort. Monspel.)

4. *latiflorum* (Ser. mus. helv. r. p. 135.), glabriusculum, floribus spicatis vel paniculatis luteis, galeâ brevi latâ, pilis pedunculorum arcuatis, ovariis glabris. — in Alpibus Bernensibus. (v. s. in h. Ser. et DC.)

5. *ramosissimum*, caule ramosissimo pedunculisque glabriusculis, floribus luteis, galeâ conico-cylindraceâ compressâ, ovariis glabris. — in Alpibus Bernensibus. (v. v. et s. in herb. Ser.)

6. *rubicundum* (Ser. mus. helv. r. p. 135.), floribus paniculatis rubicundis luteo-variegatis, galeâ conico-cylindraceâ compressâ, calcare arcuè spirali-ter contorto, floribus pedunculisque villosis, pilis rectis horizonta-libus, ovariis elongatis pilosiusculis. — A. rubicundum Fisch! in h. DC.

7. *Carpathicum* (Ser. mus. helv. r. p. 136.), floribus paniculatis lurido-purpureis luteo-variegatis, galeâ conico-cylindraceâ compressâ, calcare arcuè spiraliter contorto, pedunculis caulibusque glaberrimis, foliis profunde sectis. — A. septentrionale *Æ. carpathicum* DC. syst. r. p. 370. Sims. bot. mag. t. 2196.

8. *septentrionale* (Ser. mus. helv. r. p. 136.*), floribus paniculatis cæru-leis, galeâ conico-cylindraceâ compressâ, calcare spiraliter contorto, caule pedunculis floribusque villosis, pilis rectis vel arcuatis, ovariis glabris vel pilosis. — in Lapponiâ, Rossiâ. A. septentrionale Koell. spic. 22. et DC! syst. r. p. 370. A. Lycoctonum A. *Æ. Wahlenb. A. australe* Reichenb. ueb. 71. OEd. fl. dan. t. 123. (v. s. in herb. DC. et Ser.)

9. Floribus paniculatis luteis, galeâ elongato-cylindricâ inclinâtâ deformi, caule pedunculisque glabriusculis, ungue cncallorum brevissimo, cal-care lato, foliis amplis glabriusculis. — in Alpibus Bernensibus. Ser. mus. helv. r. p. 137. (v. v. et s. in h. Ser.)

3. A. *ochroleucum* (Wild. spec. 2. p. 1233.), floribus spicatis vel panicu-latis numerosis, galea conico-cylindraceâ elongatâ, calcare gracili recto apice curvato, labio ungui subparallelo, filamentis staminum basi alatis, alis cuspidatis, foliis profunde 3-5-lobis, lobis cuneatis trifidis, nervis fre-quenter anastomosantibus, seminibus. . . . in Caucaso. Ser. mus. helv. t. 15. f. 7. 8. 9.* A. album Ait. hort. Kew. ed. r. v. 2. p. 246. (v. s. in herb. DC. et Ser.)

4. *puberulum*, pilis rectis patulis. — circa Doroninsk. A. galeatum Stev. in h. DC!

4. A. *barbatum* (Patr! in Pers. ench. 2. p. 83.) floribus paniculatis, galeâ conicâ, calcare crasso obtuso brevissimo, labio ungui parallelo, filamentis staminum basi alatis, alis cuspidatis, mucronibus divergentibus, foliis pro-fundè lobatis, laciniis angustis divergentibus rectis vel arcuatis acutis vel obtusis, nervis divergentibus, seminibus. . . . 2? in Sibiriâ. DC! syst. r. p. 367. Ser. mus. helv. r. p. 138.* t. 15. f. 10. 11. f. Deless. ic. sel. r. t. 64. A. squarrosum. Lin. spec. 757. DC. syst. r. p. 368.* ex syn. (v. v. et s.)

5. *hispidum* (Ser. mus. helv. r. p. 139.), pilis rectis patentibus, lobis foliorum obtusis. — in Sibiriâ. A. hispidum. DC! syst. r. p. 367.*

Sect. III. CAMMARUM. DC. syst. 1. p. 374. Ser. mus. helv. 1. t. 15. f. 12-40 et 49.

Sepala decidua, galea conica compressa; ovaria 3-5. — Lobi foliorum trapeziformes pinnatisecti; radix tuberosa; flores cærulei albo variegati, raro carnei.

5. A. VARIEGATUM (Lin. spec. 751.) paniculâ divaricatâ (glaberrimâ) ramis tortuosis, galeâ conicâ elongatâ inclinâtâ vel rectâ anticè abruptè rostratâ, calcare crasso subspirali, ovariis 3-5 elongatis breviter mucronatis, foliorum lobis rhomboideis. (2) in Carniolâ, Italiâ, Bohemiâ. A. hamatum Reichenb. in cat. hort. Berol. 1818. Ser. mus. helv. 1. p. 139.* t. 15. f. 12. 13. (v. v. et s.)

a. pauciflorum (Ser. mus. helv. 1. p. 140.), galeâ rectâ, floribus pallidè cyaneis. — A. lycocotnum X Thora Italica Clus. hist. 2. p. 98. f. 1. — A. rhyncanthum f. bicolor. Reichenb. ueb. p. 56. ex Schleich! (v. s.)

β. pallidiflorum (Ser. mus. helv. 1. p. 140.) galeâ inclinâtâ, floribus albis cyaneo-variegatis. — A. variegatum Lin. spec. 751. DC. syst. 1. p. 376. A. variegatum bicolor Reichenb. ueb. 57. ex Schleich! (v. s.)

γ. cæruleum (Ser. mus. helv. 1. p. 140.), galeâ inclinâtâ, floribus cæruleis; A. cammarum Schleich! cat. 1821. p. 5. ex herb. DC. (v. s.)

δ. angustilobum (Ser. mus. helv. 1. p. 140.), galeâ inclinâtâ, foliorum lobis angustissimis, floribus cæruleis.

ε. albiflorum (Ser. mus. helv. 1. p. 140.), floribus parvis albis, galeâ rectâ. (v. v. et s. in h. Ser.)

ζ. grandiflorum (Ser. mus. helv. 1. p. 140.) floribus cæruleis, galeâ maximâ. — A. variegatum DC. herb. à cl. Bernhardi missum.

η. falcatum (Ser. mus. helv. 1. p. 141.), galeâ subconicâ falcato-naviculatâ. — A. Rhynchanthum Reichenb. ueb. 56. ex Schleich! exs. 1822. herb. DC.

6. A. GIBBOSUM (Ser. mus. helv. 1. p. 141.* t. 15. f. 14. 15.) paniculâ contractâ (glaberrimâ), galeâ conicâ elongatâ inclinâtâ vel rectâ, rostro inclinato, sacco cucullorum labium æquante anticè gibbosq., calcare elongato arcuato (non spirali) ungue brevi, alis staminum mucronatis, ovariis 3-5 foliorum venis flexuosis. (2) in sylvis umbrosis Caucasi. A. cammarum Bieb. fl. taur. et suppl. n. 1083. (v. s. in h. DC.)

7. A. ROSTRATUM (Bernh. monog. ined. ex Schleich! cat. 1815.) paniculâ laxiusculâ, galeâ conicâ elongatâ anticè abruptè mucronatâ, sacco cucullorum cylindraco, calcare crasso spirali, labio longitudine sacci, filamentorum alis evanescentibus, ovariis 3-5 glabris vel pilosisculis. (2) Ser. mus. helv. 1. p. 141. t. 15.* f. 16. 17. (v. s.)

a. Judenbergense (Ser. mus. helv. 1. p. 142.) galeâ rectâ humili, pedunculis ovariisque glabris. — in Alpibus Salisburgensibus et Helveticis. A. rostratum Bernh. mon. ined. ex Schleich! cat. 1815. DC. syst. 1. p. 376.* A. gracile Reichenb. ueb. 55. A. Bernhadianum Wallr. sched. crit. 1. p. 250.* t. 2 (v. s.)

β. Jacquintinum (Ser. mus. helv. 1. p. 142.), galeâ elatâ rectâ vel inclinâtâ, pedunculis ovariisque glabris. — in Alpibus Bernensibus. A. cammarum Jacq. fl. austr. 5. p. 11. t. 224. (v. s.)

γ. pilosiusculum (Ser. mus. helv. 1. p. 142.), galeâ inclinâtâ, pedunculis ovariisque pilosisculis. — e Gemmaio. A. cammarum. Wahlenb. fl. Carp. n. 533. (v. s.)

δ. latiflorum (Ser. herb. et mus. helv. 1. p. 142.), galeâ conicâ dilatâtâ rostro inclinato vix prominulo. — in Alpibus Bernensibus. (v. v. et s. s.)

8. A. HABEETUM (DC. syst. 1. p. 376. excl. syn. Jacq. et Wahlenb.) paniculâ rigidâ subpatente, ramis multifloris (puberulis) galeâ conicâ inclinâtâ, rostro compresso inclinato, sacco cucullorum conico, calcare crasso brevissimo curvo, labio longitudine sacci, alis staminum evanescentibus, ovaria

3-5 (puberulis) ② in montibus Gemmio-Lavaraz. Ser. mus. helv. 1. p. 143. t. 15. f. 18. 19. (v. s.)

β. *multifidum* (Schleich! ex h. DC.), foliorum lobis multifido-laciniatis caule monstroso, pedunculis agglutinatis. (v. s.)

9. A. *PANICULATUM* (Lam. fl. fr. ed. 1. suppl. 1224. ex DC. syst. 1. p. 375.*) paniculâ divaricatâ, ramis tortuosis flexuosis (puberulis), galeâ conico-semicirculari, rostro inclinato, sacco cucullorum subconicoideo, calcare crasso brevi spirali, labio longitudine sacci, alis staminum angustis evanescentibus, ovariis 3 rariis 4, folliculis elongatis. ② in Helvetiâ. Ser. mus. helv. 1. p. 144.* t. 15. f. 20. 21.

α. *Penninum* (Ser. mus. helv. 1. p. 144.*), caule subflexuoso, paniculâ pauciflorâ subcorymbosâ, galeâ conico-semicirculari, rostro elongato. — in monte Pennino et Cenisio. A. cernuum β. pauciflorum Reichenb. uebers. 44. ex charact. (v. s.)

β. *acuminatum* (Ser. mus. helv. 1. p. 144.), floribus subspicatis laxis, galeâ conico-semicirculari, ovariis pubescentibus. — A. acuminatum Reich. neb. p. 46. ex Schleich! cat. 1821. p. 5. (v. s.)

γ. *Storkianum* (Ser. mus. helv. 1. p. 145.), caule flexuoso, paniculâ laxâ debili, rostro brevi. — in Alpibus Helveticis. A. Napellus officialis Stork. libel. de stram. etc. p. 69. icon. A. paniculatum α. DC. syst. 1. p. 373.

δ. *flexicaule* (Ser. mus. helv. 1. p. 145.), caule flexuoso, paniculâ pyramidalî, galeâ latâ adscendente, rostro brevi. — in Alpibus Salisburgensibus. A. flexicaule, Hoppe exs. dec. 2. A. cernuum α et molle. Reichenb. neb. 43 et 47. (v. s.)

ε. *lacinosum* (Reichenb. ex Schleich! cat. 1821. p. 5.), caule subflexuoso, galeâ conico-semicirculari inclinatâ, rostro longissimo tenui, foliorum lobis laciniatis, ovariis glabriusculis. Ser. mus. helv. 1. p. 145. (v. s.)

ζ. *patentissimum* (Ser. mus. helv. 1. p. 145.), caule flexuoso, galeâ conicâ inflexâ, rostro brevi. — in Sibirîâ. A. paniculatum γ. DC! syst. 1. p. 375. A. cernuum flexicaule, Reichenb. ex Schleich! cat. 1821. p. 5. (v. s. in h. DC.)

η. *cernuum* (DC! syst. 1. p. 375.), caule flexuoso, paniculâ laxâ multiflorâ nutante, galeâ subconicâ inclinatâ. — circa Ratisbonam. Clus. hist. 2. p. 97. f. 1. A. Clusii Reichenb. neb. 22?

10. A. *JAPONICUM* (Thunb. fl. jap. 234.*) glaberrimum, paniculâ laxiusculâ, ramis adscendentibus, galeâ exactè conicâ abruptè mucronatâ, cucullorum sacco amplo ventricoso, calcare crasso inclinato subinvolutò, alis filamentorum evanescentibus, ovariis 3. ② in Japoniâ. Ser. mus. helv. 1. p. 146. t. 15. f. 22. 23.*

α. *carneum* (Ser. plant. sel. cent. 1. et mus. helv. 1. p. 146.), floribus carneis. A. album DC. syst. 1. p. 377? (v. s.)

β. *cæruleum* (Ser. mus. helv. 1. p. 146.), floribus cæruleis. — A. glabrum DC. syst. 1. p. 379?

11. A. *UNCINATUM* (Lin. spec. 750.) paniculâ laxiusculâ, ramis divergentibus, galeâ exactè conicâ, calcare inclinato subspirali labio ungui piloso subparallelo, alis filamentorum evanescentibus, ovariis villosis 3-5 (floribus cæruleis) foliis 3-lobis, lobis subintegris. ②

α. *Linnaeanum* (Ser. mus. helv. 1. p. 147.) — propè Philadelphiam A. uncinatum Lin. spec. 750. A. uncinatum var. α. DC. syst. 1. p. 379.*

β. *Michauxianum* (Ser. mus. helv. 1. p. 147. t. 15. f. 24. 25.*) — A. uncinatum Michx! fl. bor. amer. 315. Sims. bot. mag. t. 1119.* A. uncinatum. β. DC. syst. 1. p. 379. (v. s. in h. DC.)

12. A. *SPECIOSUM* (Otto cat. hort. berol. ex Brunner!) paniculâ laxiusculâ, galeâ exactè conicâ, rostro inclinato, ungue cucullorum apice incurvo, sacco amplo, calcare crassissimo obtuso brevissimo, alis filamentorum apice mucronatis (filamentis pilosis) ② Patr. ign. Ser. mus. helv. 1. p. 147.* t. 15. f. 26. 27. (v. s.)

13. *A. TORTUOSUM* (Willd. enum. 1. p. 576.) paniculâ laxiusculâ, ramis 1-4-floris, galeâ subconicâ, sacco cucullorum inflato, calcare crasso longo abrupte geniculato (non arcuato, nec convoluto) alis filamentorum evanescentibus, filamentis pilosiusculis, ovariis 3-5 glabris. ② Patr. ign. Ser. mus. helv. 1. p. 147. t. 15. f. 28. 29.
- α. *napellifolium* (Ser. mus. helv. 1. p. 148.), paniculâ laxâ simpliciusculâ, ramis pedunculisque pubescentibus, ovariis 5 glabris lobis foliorum angustè cuneiformibus, lobulis acutis. — *A. tortuosum* Willd. enum. 1. p. 576. DC. syst. 1. p. 378.
- β. *illinitum* (Ser. plant. sel. cent. 1. et mus. helv. 1. p. 148.). paniculâ ramosissimâ laxissimâ, ramis pedunculisque glaberrimis, ovariis 3 glabris, lobis foliorum latè cuneiformibus, lobulis obtusis. — *A. illinitum* Reich. neb. 54. *A. neomontanum* Panz! fl. norimb. ined. (v. v.)
14. *A. CILIARE* (DC. syst. 1. p. 378.*) caule patentim piloso, paniculâ laxâ, ramis adscendentibus, galeâ exactè conicâ anticè mucronatâ, sacco cucullorum . . . , calcare apice adunco, staminibus glabris, alis filamentorum evanescentibus, ovariis 5-7, lobis foliorum pinnatisectis, lobulis linearibus. ②†
- α. *oligotrichum* (DC. syst. 1. p. 378.), caule parè piloso. — in Sibiria. *A. volubile* Willd. spec. 2. p. 1237.
- β. *polytrichum* (DC. syst. 1. p. 378.). *A. ciliare* Deless. ic. sel. 1. t. 65. — in Sibiria. Ser. mus. helv. 1. p. 149. t. 15. f. 30. *A. flaccidum* Reichenb. neb. 39. (v. s. in h. DC.)
15. *A. MAXIMUM* (Pall. herb. ex. DC! syst. 1. p. 380.) paniculâ laxâ, ramis paucis longis distantibus pubescentibus, pilis curvatis, galeâ subconicâ, sacco cucullorum inflato, calcare brevi incurvo, filamentis staminum glabris, alis longè mucronatis, ovariis . . . , bracteolis flore approximatis. ② in Kamtschatka. *A. abbreviatum* Laugsd. ex herb. DC. — Ser. mus. helv. 1. p. 149. t. 15. f. 31. 32. (v. s. in h. DC.)
16. *A. PROUCTUM* (Reichenb. acon. p. 75. t. 7. f. 3.*) racemo paucifloro, galeâ conicâ longè mucronatâ, sacco cucullorum subnullo, labio longissimo, calcare obtuso crasso brevissimo, staminibus glabris alis evanescentibus, ovariis 4-5 pubescentibus, foliorum lobis 3-partitis. ② in Sibiria. *A. Delphinii-folium* β. Sibiricum DC. syst. 1. p. 380. ex Reich. l. c. — Ser. mus. helv. 1. p. 150.* t. 15. f. 33. 34.
17. *A. MOLLE* (Reichenb. neb. 47. ex Schleich! cat. 1821. p. 5. h. DC. †) floribus paniculato-spicatis (cæruleis) galeâ conico-rotundatâ amplâ, cucullis maximis, sacco longo, calcare brevi inclinato, alis staminum angustis evanescentibus, filamentis pilosis, ovariis 3-4 glabris ② patr. ign. Ser. mus. helv. 1. p. 150. t. 15. f. 35. 36. †.
18. *A. EXALTATUM* (Bernh. ex specim. hort. goett. 1818.) paniculâ laxiusculâ, ramis adscendentibus rigidiusculis, galeâ conicâ, rostro elongato adscendente, sacco cucullorum subconico inclinato, calcare crasso incurvo, filamentis staminum pilosis vel glabris, alis mucronatis vel evanescentibus, ovariis 3-5 glabris. ② Ser. mus. helv. 1. p. 151.* t. 15. f. 37. 38. †.
19. *A. INTERMEDIUM* (DC! syst. 1. p. 374.*) paniculâ laxiusculâ, ramis adscendentibus rigidiusculis, galeâ valdè convexâ subconicâ, rostro brevi horizontali, sacco cucullorum subconico, calcare crasso incurvo, filamentis staminum pilosis rarè glabris, alis mucronatis vel evanescentibus, ovariis 3-5 (rariùs 7) glabris. ② patr. ign.
- α. *glabrum* (Ser. mus. helv. 1. p. 152. t. 15. f. 39. 40. 49.), caule foliis pedunculisque glaberrimis, floribus cæruleis. — *A. intermedium* DC. syst. 1. p. 374. *A. Napellus* Sturm. deutschl. fl. heft. 6. ic. (v. s.)
- β. *versicolor* (Ser. mus. helv. 1. p. 152.), caule foliis pedunculisque glaberrimis, floribus variegatis. *A. Stœrkianum* γ. *bicolor*. Reichenb. neb. 49.
- γ. *latilobum* (Ser. mus. helv. 1. p. 152.), foliorum lobis latis. — *A. tauricum* Rœm! in herb. DC. (v. v.)

δ. *pubescens*, (Ser. mus. helv. 1. p. 152.), caule pedunculisque puberulis floribus cæruleis.

Sect. IV. NAPELLUS. DC. *systema*. 2. p. 371.

Sepala decidua, galea semicircularis rarò navicularis, ovaria 3-5-7 (rarissimè pilosa). Foliorum lobi cuneati bipinnati-secti; radix tuberosa; flores cærulei albi albo-cærulei vel cæruleo-flavi.

20. A. NAPELLUS (Lin. spec. 751.) floribus densè spicatis vel laxè paniculatis, galeâ semicirculari rarò naviculari, sacco cucullorum subconico, calcare brevi crasso inclinato, alis staminum cuspidatis vel evanescentibus, foliorum lobis cuneatis pinnatisectis, ovariiis 3 rarò 5, glabris vel pilosis. ② in Europâ. Ser. mus. helv. 1. p. 162*. t. 15. f. 41. 42. 50.

* *Varietates ex inflorescentiâ dispositæ.*

- α. *Tauricum*, (Ser. mus. helv. 1. p. 153*.) floribus densè spicatis intensè cæruleis, bracteis longis, galeâ subnaviculari. — in Germaniâ. A. Tauricum Wulf. in Jacq. col. 2. p. 112.* ic. rar. 3. t. 49. 2. DC. *systema*. 1. p. 374. A. Tauricum Reich. acon. 87. t. 12. f. 2-3. A. Schleicheri Reichenb. ueb. 35. (v. s.)
- β. *spicatum* (Ser. mus. helv. 1. p. 154.), floribus laxè spicatis intensè cæruleis, bracteis brevibus, galeâ semicirculari sepala lateralialia amplectente, limbo subemarginato. — A. Kœlleianum Reichenb. acon. t. 11. f. 1. — A. Braunii Reichenb. ueb. 29. (v. s.)
- γ. *bracteosum*, (Ser. mus. helv. 1. p. 154.), floribus laxè spicatis intensè cæruleis, bracteis longissimis, galeâ semicirculari, limbo emarginato. — A. commutatum Reichenb. ueb. 36. acon. t. 18. f. 3. (v. s.)
- δ. *pygmæum*, (Ser. mus. helv. 1. p. 154.), floribus 3-8 spicatis intensè cæruleis, foliis congestis. — A. Kœlleianum pygmæum. Reich. acon. t. 12. f. 1.
- ε. *macrostachyum*, (Ser. mus. helv. 1. p. 154.), floribus numerosis longè spicatis intensè cæruleis, galeâ semicirculari limbo integro sepala lateralialia amplectente, foliorum lobulis divaricatis. — in Moraviâ. A. callybotrion. Reichenb. acon. 98. t. 16. f. 1.
- ζ. *ramosum* (Ser. mus. helv. 1. p. 154.), floribus spicato-paniculatis intensè cæruleis, galeâ semicirculari, limbo emarginato. — A. Halleri Reichenb. uebers. p. 27. (v. s.)
- η. *virgatum* (Ser. mus. helv. 1. p. 155.), floribus paniculatis distantibus, ramis virgatis longissimis, galeâ integrâ sepala lateralialia amplectente vel falcata. A. eriostemum. DC. *systema*. 1. p. 377. A. virgatum. Reichenb. uebers. 28. A. volubile Kœll. spic. 21.* (v. s.)

** *Varietates ex galeæ formâ, floris colore et magnitudine dispositæ.*

- θ. *amplexum* (Ser. mus. helv. 1. p. 155.). floribus laxè spicatis, intensè cæruleis, bracteis brevibus, galeâ semicirculari, limbo integro sepala lateralialia amplectente. — A. anœnum Reichenb. ueb. 23. acon. 93. t. 14. f. 1.
- ι. *Hoppeanum* (Ser. mus. helv. 1. p. 155.*), floribus laxè spicatis cæruleis, galeâ falcato-naviculatâ rostellatâ, foliorum lobis obtusiusculis. — in Carinthiâ. A. Hoppii Reichenb. ueb. 24. et Hoppe! exsic. (v. s.)
- κ. *semigaleatum* (Ser. mus. helv. 1. p. 155.), floribus pallide cyaneis, galeâ naviculatâ. — in Kamtschatkâ. A. delphinifolium γ Kamchaticum. DC. *systema*. 1. p. 380. A. semigaleatum. Pall. et Reichenb. acon. 77. t. 8.
- λ. *falcatum* (Ser. mus. helv. 1. p. 155.*), floribus laxè spicatis cæruleo-purpureis, galeâ falcato-naviculatâ. — A. formosum Reichenb. acon. t. 18. f. 2. A. Eustachium. Reichenb. acon. t. 15. f. 3. etc.
- μ. *rostellatum* (Ser. mus. helv. 1. p. 156.), floribus spicatis cæruleis, galeâ erectâ subconicâ abruptè rostellatâ. — A. acutum. Reichenb. acon. t. 14. f. 2.
- ν. *multicucullatum* (Ser. ic. ined. et mus. helv. 1. p. 156.). (v. s.)
- ξ. *Neubergense* (Ser. mus. helv. 1. p. 156.), caule stricto, floribus laxè

- paniculatis cæruleo-purpureis, laciniis foliorum brevibus obtusiusculis. — in Sibiria. A. strictum Bernh. et DC. syst. 1. p. 373. Reichenb. acon. t. 17 f. 1. A. Neubergense. DC. syst. 1. p. 273. A. callybotryon Reichenb. acon. t. 16 f. 2. etc.
- o. *grossum* (Ser. mus. helv. 1. p. 157.*), floribus laxè spicato-paniculatis amplissimis cæruleo-purpureis, foliorum laciniis subparallelis amplis. — A. autumnale Reichenb. acon. t. 17 f. 2. A. Tauricum Willd. spec. 2. p. 1236. A. laxum Reichenb. acon. 97. t. 15 f. 4. etc. (v. s.)
- π. *lætum* (Ser. mus. helv. 1. p. 157.), floribus laxè spicato-paniculatis amplis cyaneis, laciniis foliorum approximatis subparallelis acutiusculis. — A. lætum, Reichenb. acon. 89. t. 13 f. 2. A. Koellæanum firmum Reichenb. acon. t. 14 f. 2.
- ρ. *viridiflorum* (Ser. mus. helv. 1. p. 158.), floribus spicatis laxiusculis, galeâ semicirculari vix emarginatâ cæruleâ lineis viridibus notatâ (v. s. in h. Ser.)
- σ. *maculatum* (Ser. mus. helv. 1. p. 158.), floribus spicatis cæruleis albo maculatis (v. s. in herb. Ser.)
- τ. *rubellum* (Ser. mus. helv. 1. p. 158.), floribus spicatis cæruleo-rubellis, lobis foliorum angustissimis. — A. Napellus floribus rubellis Schleich! cat. 1821. p. 5.
- υ. *bicolor* (Ser. mus. helv. 1. p. 158.), floribus spicatis vel paniculatis, albis cyaneo-variegatis. — in Helvetiæ monte Stokhorn. A. bicolor Schult. obs. bot. 101.* A. vulgare γ. bicolor. DC. syst. 1. p. 372. A. Halleri bicolor Reichenb. ueb. 28. (v. v.)
- φ. *albiflorum* (Ser. mus. helv. 1. p. 158.), floribus laxè spicatis niveis, galeâ semicirculari, limbo subintegro. (v. s. in herb. Ser.)

***. *Varietates ex foliis dispositæ.*

- χ. *anthoræfolium* (Ser. mus. bot. 1. p. 159.), floribus laxè spicato-paniculatis intensè cæruleis, pedicellis longis, galeâ semicirculari, limbo integro amplo, foliis A. anthoræ. — A. angustifolium Pall. ex Reichenb. acon. 95. t. 15 f. 2. A. eustachyum Reichenb. acon. t. 15 f. 3.
- ψ. *microphyllum* (Gaud. ined.), floribus paucis in spicam laxam dispositis, lobis foliorum linearibus obtusis brevibus. — in Helvetiâ. A. laxiflorum. Schleich! cat. 1821. p. 5 — Ser. mus. helv. 1. p. 159. (v. s.)
- ω. *lacinosum* (Ser. mus. helv. 1. p. 159.), floribus laxè spicatis paniculatisque cæruleis amplis subconicis, laciniis foliorum profundis linearibus acutis. (v. s.)
- αα. *delphinifolium* (Ser. mus. helv. 1. p. 159.), floribus paucis laxè spicatis amplissimis, pedunculis elongatis intensè cæruleis, ovariis 4-6, foliorum lobis 2-3-partitis, lobulis indivisis. — in Americâ bor. A. delphinifolium α. americanum. DC! syst. 1. p. 380. Reichenb. acon. t. 9 f. 1. 2. 3.
- ββ. *tenuifolium* (Ser. mus. helv. 1. p. 159.), floribus spicatis cæruleis, lobis foliorum tenuibus obtusiusculis vix divergentibus. β A. venustum Reichenb. ueb. p. 28. (v. s.)

****. *Varietates ex superficie dispositæ.*

- γγ. *pubescens* (DC. syst. 1. p. 372.), caule floribus pedicellis foliisque pubescentibus. — in Pyrenæis. A. Napellus β. DC. A. fr. 5. p. 642. A. Schleicheri Reichenb. ueb. 35. (v. s.)

****. *Varietates ex statu caulis.*

- δδ. *bulbiferum* (Ser. mus. helv. 1. p. 160.), floribus spicatis, galeâ sepala lateralâ amplectente, foliis longè petiolatis bulbos subrotundos in axillis gerentibus, lobulis obtusis. (Ser. ic. ined. et herb.)
- εε. *volubile* ? floribus. foliorum laciniis profundis linearibus acutis
†. A. ciliare α. oligotrichum. DC. syst. 1. p. 379?

21. *A. FEROX* (Wallich! in litt.) floribus laxè spicato-paniculatis, galeâ semicirculari, cucullorum sacco longo angusto, calcare crasso inclinato, labio angustissimo divaricato, lobis foliorum cuneatis, lobulis acutis divaricatis, ovariis 5 villosis. ② in Nepauliâ. Ser. mus. helv. 1. p. 160.* t. 15. f. 43. 44. (v. s. Wall. communic.)
22. *A. BIFLORUM* (Fisch! in litt. cum ic. DC! syst. 1. 380.) caule 2-3-floro, cucullorum sacco amplo subrotundo obtusissimo ecalcarato, labio retroflexo, staminum filamentis basi alatis, alis latius cuspidatis, foliis profundè pinnatisectis, lobis linearibus divergentibus simplicibus, radice tuberosâ, ovariis 3 pilosis. ② in Sibirîâ. Ser. mus. helv. 1. p. 161.* t. 15. f. 45. 46.
- α. *luteo-cæruleum*, A. biflorum Fisch! in litt. cum ic. DC! syst. 1. p. 380.* A. grandiflorum Fisch! cat. hort. gor. 1808. p. 77. Ser. mus. helv. 1. p. 161. (v. s. in h. DC.)
- β *cæruleum*, A. biflorum Reichenb. acon. 74. t. 7. f. 1. 2. Ser. mus. helv. 1. p. 161.

Trib. V? *PÆONIACEÆ*. — Ranunculaceæ spurîæ. DC. syst. 1. p. 281.

Antheræ introrsæ. — an ordo proprius?

XXVII. ACTÆA. Lin. spec. ed. 2. p. 722. DC. syst. 1. p. 381.

Calyx deciduus 4-sepalus. Petala 4. Carpella oo-sperma. — Herbæ perennes.

Sect. I. CIMICIFUGA. DC. syst. 2. p. 282.

Flores polygyni. Carpella sicca dehiscentia.

1. *A. CIMICIFUGA* (Lin! amœn. 2. p. 354.* spec. ed. 2. p. 722.) Ovariis 4 subsessilibus villosissimis, racemis paniculatis, foliis ternatim biternatimve sectis, segmentis ovato-lanceolatis inciso-dentatis. ½ in sylvis montanis Galliziæ et in Sibirîâ orient. Cimicifuga foetida. Lin! syst. nat. ed. 12. p. 659. Gaertn. de fruct. 2. p. 275. t. 140. Herba foetidissima ad cinices fugandos in Sibirîâ adhibita. (v. s.)
- β? *simplex* ½ in Kamschatkâ frequens. Cimicifuga simplex (Wormsk ined. ex Fisch. in litt.) caulis simplex indivisus, racemi solitarii aut rarè bini recti altero, dum adsit minuto laterali rarissimè terno. An spec. propria?
2. *A. PODOCARPA* (DC. syst. 1. p. 282.) ovariis 4-5 glabris stipitatis, racemis paniculatis, foliis decompositis. ½ in opacis sylvaticis montium Carolinæ. Cimicifuga Americana Michx! fl. bor. am. 1. p. 316. A. podocarpa Deless! ic. sel. 1. t. 66. Habitus A. racemosæ. (v. s. in h. Mus. Par.)
3. *A. CORDIFOLIA* (DC. syst. 1. p. 383.*) ovariis 2-3 glabris sessilibus, racemis paniculatis, foliis biternatim sectis, segmentis basi cordatis 5-7 lobis serratis. ½ in sylvis umbrosis montium Carolinæ. Cimicifuga cordifolia Pursh! fl. bor. am. 2. p. 373. excl. syn. Similis A. racemosæ et A. podocarpæ. (v. s. in herb. Lamb.)
4. *A. PALMATA* (DC. syst. 1. p. 383.*) ovariis 12-15 in capitulum subrotundum collectis, racemis dichotomè subpaniculatis, foliis palmatifidis. ½ secus rivulos montium Carolinæ. Cimicifuga palmata Michx! fl. bor. am. 1. p. 316. Planta omninò glabra. (v. s. in h. Mus. Par. et Lam.)

Sect. II. MACROTYS. Raf. in Journ. bot. 1808. v. 2. p. 170.

Flores monogyni. Carpella sicca dehiscentia. DC. syst. 1. p. 384.

5. *A. RACEMOSA* (Lin. spec. 722.) monogyna, racemis longissimis, carpellis siccis dehiscentibus, foliis ternatim sectis, segmentis serratis subincisive. ½ in sylvis montium à Canadâ ad Floridam. A. monogyna Walt. fl. car. 151. Cimicifuga racemosa Bart. phil. 2. p. 12. Herba A. spicata sed major, florescentia A. Cimicifugæ sed monogyna, fructus A. cordifoliæ sed unicapularis. (v. s.)

7. *A. JAPONICA* (Thunb. jap. 221.*) monogyna, spicis longissimis; foliis tri-sectis, segmentis cordatis palmatim 5-7-lobatis. $\frac{1}{2}$? in Japoniâ Willd. spec. 2. p. 1140. Flores sessiles. DC. syst. 1. p. 384.*

Sect. III. CHRISTOPHORIANA. Tourn. inst. 299. t. 154.

Carpellum 1 indehiscens. DC. syst. 1. p. 384.

8. *A. SPICATA* (Lin! spec. 722.) monogyna, baccis subglobosis, petalis staminum longitudine, racemo ovato, foliis bi-triternatimve sectis, segmentis ovato-lanceolatis serratis incisive. $\frac{1}{2}$. in Europæ ferè totius. OEd. fl. dan. t. 589. Gaertn. de fruct. 2. p. 154. t. 114. DC. syst. 1. p. 384. (v. v.)
9. *A. BRACHYPETALA* (DC. syst. 1. p. 385.*) monogyna, baccis ovatis oblongis, petalis staminibus brevioribus, racemo ovato, foliis bi-triternatimve sectis, segmentis ovato-lanceolatis serratis incisive. $\frac{1}{2}$. in Americâ boreali.
a. *alba*, baccis albis, segmentis foliorum oblongioribus acutioribus. — in Canadâ.
ß. *rubra*, baccis rubris, foliorum segmentis latioribus obtusioribus. — in Canadâ. — Flores fructusque majores, semina majora. videtur species propria.
γ. *cærulea*, baccis cæruleis. — in Floridâ.
δ. *microcarpa*, baccis parvis albis subrubellis, pedunculis incrassatis. Circa Bostoniam.

XXVII. ZANTHORHIZA. Marsh. arb. 167. Lam. ill. t. 854. DC. syst. 1. p. 386.

Calyx deciduus 5-sepalus. Petala 5. Carpella 2-3 sperma, abortu 1-sperma.

1. *Z. APHIFOLIA* (L'her! st. nov. p. 79. t. 38.*) $\frac{1}{2}$. in Virginiâ. Duham. abr. ed. nov. 3 p. 181. t. 37.* Flores atro purpurei, parvi, sæpè abortu unisexuales.

XXVIII. PÆONIA. Lin. gen. n. 678.

Cal. 5-sepalus foliaceus inæqualis. Pet. 5-10 suborbiculata. Stam. ∞. Discus carnosus ovaria cingens. Carpella 2-5 grossa stigmatibus bilamellatis crassis instructa, in folliculos capsulares couversa. Semina subglobosa nitida. — Radices fasciculatæ. Folia caulina biternatim secta. Flores ampli albi aut purpurascens. DC. syst. 1. p. 386.

Sect. I. MOUTAN.

Caulis fruticosus. Discus expansus in urceolum membranaceum carpella plus minus involentem.

1. *P. MOUTAN*. (Sims bot. mag. t. 1154.*) caule fruticoso, foliorum segmentis ovali-oblongis, subtus glaucis, carpellis villosis 5. — in Chinâ. DC! syst. 1. p. 387. Anders. tr. Lin. soc. 12. p. 252.* ex quo.
a. *papaveracea* (bot. rep. t. 463.) petalis 8-13 albis basi macula purpurea notatis, capsulis omnino in urceolum inclusis.
ß. *Bankii*, floribus plenis, petalis medio rubicundis, segmentis apice fissuris obtusis. — bot. rep. t. 448.
γ. *rosea*, floribus subplenis roseis, segmentis apice fissuris obtusissimis. bot. rep. t. 373. (v. v.)

Sect. II. PÆON.

Caulis herbaceus. Discus vix aut non expansus et carpella imâ basi tantum circumdantia.

* *Folliis glabris.*

2. *P. CORALLINA* (Retz. obs. 3. p. 34.*) herbacea, carpellis tomentosis, foliorum segmentis ovatis integris glabris. $\frac{1}{2}$. in sylvaticis asperis Europæ. — Blackw. herb. t. 245. DC! syst. 1. p. 388. And. l. c. p. 268. (v. s.)
3. *P. OFFICINALIS* (Retz. obs. 3. p. 35.) herbacea, carpellis tomentosis rectiusculis, foliorum segmentis inæqualiter laciniatis glabris, laciniis ovato-lanceolatis, $\frac{1}{2}$. in Europæ sylvaticis montosis. — bot. mag. t. 1784. —. Var. floribus purpureis roseis carneis imo albis, simplicibus aut plenis. DC. syst. 1. p. 339. And. l. c. p. 264 et 286. (v. v.)
4. *P. TRITERNATA* (Pall. nov. act. petr. v. 10.) herbacea, carpellis tomentosis
I.

- erectis, foliorum segmentis subtus glaucis glabris sublobatis, lobulis obovatis obtusis. *Æ.* in Tauriâ nec in Daburiâ monente cl. Fischer. *P. Daurica* Andr. bot. rep. t. 486. — DC! syst. 1. p. 391. And. l. c. p. 270.* (v. s.)
5. *P. LOBATA* (Desf. cat. h. par. 126.) herbacea, carpellis tomentosis erectiusculis, foliorum segmentis glabris decurrentibus pinnatipartitis apice trilobatis. *Æ.* in Lusitaniâ ex Juss. DC. syst. 1. p. 391. excl. forsan veterum syn. (v. s.)
6. *P. TENUIFOLIA* (Lin! sp. 748.) herbacea, carpellis tomentosis patentibus, foliorum segmentis glabris multipartitis, laciniis linearibus. *Æ.* in Ukranîâ, Sibirîâ, Tauriâ. bot. mag. t. 926. DC. syst. 1. p. 394. And. l. c. p. 263. (v. v.)
Æ. laciniata (Willd. en. 573, non Pall.), caule elatiore, foliorum laciniis subtilioribus.
7. *P. HYBRIDA* (Pall. fl. ross. 2. p. 94 t. 86.*) herbacea, carpellis pubescentibus, foliorum segmentis glabris multipartitis, laciniis linearibus. *Æ.* Spontanea inter Volgam et Tauriam ex Fischerò; proles ex priore et posteriore hybrida ex Pall. — prioris mera varietas ex And. et Sab. tr. lin. soc. 12. p. 263 (v. s.)
8. *P. ANOMALA* (Lin! mant. 247.) herbacea, carpellis 5 glabris depressis obtusis, foliorum segmentis glabris pinnatipartitis, lobis lanceolatis acuminatis. *Æ.* in Sibirîâ. DC. syst. 1. p. 393. And. l. c. p. 261. bot. mag. t. 1754. *P. laciniata* Pall. ross. 2. t. 85. (v. s.)
9. *P. ALBIFLORA* (Pall. ross. 2. t. 84.) herbacea, carpellis glabris recurvatis, foliorum segmentis glabris nitidis tripartitis, lobis ovato-lanceolatis. *Æ.* in Sibirîâ. bot. reg. t. 42 DC. syst. 1. p. 392. And. l. c. p. 256. — Variat floribus 1-5 albis roseis ve, simplicibus aut plenis; foliorum segmentis planis aut concavis, angustioribus latioribusve. (v. v.)
- ** *Foliis subtus puberulis.*
10. *P. RUSSI* (Biv. man. sic. 4. p. 12.*) herbacea, carpellis pilosis recurvatis, foliorum segmentis ellipticis integris subtus vix puberulis. *Æ.* in montibus Panormitanis Siciliæ. Radix fusiformis. Petala chermesina. Sequenti affinis. (Biv.)
11. *P. HUMILIS* (Retz. obs. 3. p. 35.) herbacea, carpellis subpilosis erectiusculis, foliorum segmentis 3-5-partitis subtus villosis, lobis oblongis integris. *Æ.* in Hispaniâ. bot. mag. t. 1422. DC. syst. 1. p. 392. And. l. c. p. 271.* stigmata erecta; foliorum lobi canaliculati. (And.) (v. s.)
12. *P. DECORA* (And. l. c. p. 273.) herbacea, carpellis pubescentibus, patentibus, foliorum segmentis tripartito-laciniatis oblongis obtusis subtus pilosis. *Æ.* in Oriente. *P. Byzantina*. Clus. hist. 276. Stigmata recurvata. fol. lobicanaliculati.
a. Pallasii, segmentis angustè oblongis.
Æ. elatior, segmentis lato-oblongis.
13. *P. ARIETINA* (And. l. c. p. 275.) herbacea, carpellis tomentosis arcuato-patentibus, fol. segmentis trilobatis pinnatifidisque decurrentibus ovali-oblongis planiusculis subtus pilosis. *Æ.* in Oriente?
a. Andersonii, petalis saturatè roseis minus crispis. — J. Bauh. hist. 3. p. 493.
Æ. carnea, petalis carneis lacero-crispis. *P. cretica*. Clus. rar. 281. — Besl. syst. vern. ord. 6. p. 11.
14. *P. PEREGRINA* (Mill. dict. n. 3.) herbacea, carpellis tomentosis rectis, fol. segmentis tripartito-laciniatis integrisque ovato-lanceolatis planiusculis subtus pilosis. *Æ.* in Oriente? bot. mag. t. 1050. And. l. c. p. 277. — *P. peregrina*. *γ.* DC. syst. 1. p. 390. (v. s. in h. Banks.)
15. *P. PARADOXA* (And. l. c. p. 288.) herbacea, carpellis tomentosis rectis, fol. segmentis multipartitis obtusis subundulatis subtus glaucis pilosis. *Æ.* in Galliæ austr. montibus apricis. *P. promiscua* Lob. ic. 683. f. 2. *P. peregrina* var. *a.* DC. syst. 1. p. 390. (v. v.)
Æ. leiocarpa, carpellis glaberrimis. *Æ.* hanc cum priore commixtam sed rarissimam reperi in Monte Serane. an potius *P. humilis* var. ut vult cl. Anderson. (v. v.)
γ. Tatarica (Mill. dict. n. 5. ic. t. 199.), eadem? flore-pleno ex And. tr. lin. soc. 12. p. 289.
16. *P. MOLLIS* (And. l. c. p. 282.) herbacea, carpellis tomentosis rectis, fol. segmentis ovali-lanceolatis planis lobatis imbricatis subtus cæsiò-pilosis. *Æ.* in Sibirîâ? *Calyx* extus pubescens.

Ordo II. DILLENACEÆ. DC. syst. 1. p. 395.

Partes floris numero quinario dispositæ; tegumenta per æstivationem imbricata. Sepala persistentia, 2 exteriora, 3 interiora. Petala decidua 1-serialia hypogyna. Stamina ∞ , toro inserta, filamenta filiformia, plana, basi aut apice dilatata, libera aut polyadelpa, regulariter circà carpella sita aut unilateralia; antheræ adnatæ, loculis rimâ longitudinali sæpius dehiscentibus, introrsis lateralibusve nunquàm extrorsis. Carpella plurima (sæpius 2-5) interdum abortu solitaria aut in unicum coalita, monostyla, stigmatibus simplicibus, 1-locularia, demum membranacea, baccata aut bivalvia, libera coalitave, stylo acuminata; semina ovata, duplici serie ad carpellorum loculorumve angulum interiorem adfixa, plurima aut dua, abortu tantum solitaria, nunc nuda nunc arillo pulposo cincta. Spermoderma duriusculum; albumen carnosum subcartilagineum semini conforme; embryo minutus erectus inferus ad basim albuminis. — Arbores frutices suffrutesve. Folia ferè semper alterna, rarissimè opposita simplicia, penninervia, integra aut dentata, sæpius coriacea et persistentia nunc petiolata, nunc sessilia basi angustata, sæpè supra basim amplexicaulem transversè secta, parte inferiore persistente. Flores solitarii racemosi aut paniculati, terminales, sæpius flavi et Cistorum flores subæmulantes.

Tribus I. *DELIMEÆ*. DC. syst. 1. p. 397.

Staminum filamenta apice dilatata utrinque antheræ loculos discretos subrotundosque gerentia; styli filiformes acuti. Carpella capsularia utriculiformia aut baccata. — Arbores fruticesve interdum scandentes; flores racemosi aut paniculati.

I. TETRACERA. *Lin. gen. n. 683, Gærtn. fruct. 1. p. 336. t. 69.*

Flores sæpius dioici aut polygami. ♂ Stam. ∞ . ♀ Carpella 3-5, capsularia, sepalis imbricatis cincta; semina 1-2-ovata nitida arillata.

1. *T. volubilis* (Lin! spec. 617.) foliis obovatis obtusis apice subsinuatis scaberrimis. floribus paniculatis laxiusculis, pedunculis velutinis. 5 in insulis Barbadosibus, Brasiliâ et in Americâ merid Gaertn. de fruct. 1. p. 336. t. 69. f. 3. Capsulæ ad apicem pilis nonnullis vestitæ. DC. syst. 1. p. 398. (v. s. in h. Juss.)
2. *T. oblongata* (DC. syst. 1. p. 399.*) foliis oblongis obtusis utrinque scabris sinuato-dentatis, floribus in racemum simplicem densum congestis, pedunculis glabris. 5 in Brasiliâ. Deless! ic. sel. 1. t. 67. Aff. *T. volubili*, sed distincta ramis pedunculisque glaberrimis. (v. s. in h. Juss. Thouin.)
3. *T. Lima* (Willd! ined. in herb. Bonpl.) foliis oblongis obtusis basi cuneatis utrinque scabris integris, pedunculis paniculatis pubescentibus. 5 in Brasiliâ. Petala oblonga obtusa angusta calyce paulò longiora. DC. syst. 1. p. 399.* (v. s. in h. Bonpl.)

4. *T. ACUMINATA* (DC. syst. 1. p. 399.*) foliis ovali-oblongis acuminatis serrulatis utrinque scabris, pedunculis racemosis pubescentibus. 5 (In Américâ merid? Sepala 2 exteriora brevissima, 3 interiora orbiculata. (v. s.)
5. *T. JAMAÏCENSIS* (DC. syst. 1. p. 399.*) foliis ovali-oblongis acutiusculis basi secus petiolum longè productis utrinque sublævibus subserrulatis, pedunculis racemosis glabris subasperis. 5 in Jamaicâ. Flores ignoti. (v. s. in h. Lamb.)
6. *T. OVALIFOLIA* (DC. syst. 1. p. 400.*) foliis ovalibus obtusis integerrimis utrinque scabriusculis, pedunculis racemosis subvelutinis. 5 in Cayennâ. Deless! ic. sel. 1. t. 68. Sepala subrotunda, exteriora dorso subvelutina, interiora margine subciliata. (v. s.)
7. *T. ROTUNDIFOLIA* (Smith. in Rees cycl. vol. 35.) foliis rotundato-ellipticis integris utrinque scaberrimis, paniculis terminalibus, floribus hermaphroditis 4-gynis intus lævibus. 5 in Guyanâ. Capsulæ 3-4 ovales læves lincidæ compressæ minùs quàm *T. volabilis* inflatæ. — An satis a *T. ovalifolia* differt?
8. *T. MULTIFLORA* (DC. syst. 1. p. 400.*) foliis ovalibus subobtusis apice sinuato-dentatis secus petiolum subproductis lævibus, paniculâ multiflorâ, pedicellis glabris. 5 in Pará. Deless! ic. sel. 1. t. 69. flores ignoti. (v. s. in h. Mus. Par.)
9. *T. ALNIFOLIA* (Willd. spec. 2. p. 1243.*) foliis ovali-oblongis obtusis aut subacutis glabris supernè subasperis subtus lævibus apice subdenticulatis, pedunculis paniculatis subpubescentibus. 5 (in sylvis Guineæ. Petala obovata calyce vix longiora. DC. syst. 1. p. 401*. (v. s. in h. Lamb.)
10. *T. OBOVATA* (DC. syst. 1. p. 401.*) foliis obovatis obtusis basi attenuatis lævibus subtus rufo-velutinis integris, floribus paniculatis. 5 in Senegaliâ. An forsân Curatelæ species? (v. s. in h. Juss. et Deless.)
11. *T. SENEGALENSIS* (DC. syst. 1. p. 401.*) foliis ovalibus subacutis integerrimis lævibus, pedunculis paniculatis multifloris. 5 (in Senegaliâ. Calyx 5-sepalus glaber, sepalis suborbiculatis; petala 3. (v. s. in h. Juss.)
12. *T. LÆVIS* (Vahl. symb. 3. p. 71.*) foliis oblongis acuminatis subintegerrimis lævibus, racemis terminalibus simplicibus. 5 in Indiâ orientali. Capsulæ 4 subrotundo-ventricosæ mucronatæ glaberrimæ nitidæ. DC. syst. 1. p. 401*. (v. s. in h. Deless.)
13. *T. EURYANDRA* (Vahl. symb. 3. p. 71.) foliis ovalibus obtusis integerrimis lævibus, pedunculis paniculato-racemosis pubescentibus. 5 (in novâ Caledoniâ. *Euryandra scandens* Forst! prod. n. 228. gen. n. 41. t. 41. Deless! ic. sel. 1. t. 70. Petala 3. oblonga, calyce longiora. DC. syst. 1. p. 402*. (v. s.)
14. *T. RHEEDI*. (DC! syst. 1. p. 402.*) foliis oblongis utrinquè acuminatis integerrimis lævibus paniculâ ramosâ subdichotomâ, floribus 4-sepalis. 5 (in Malabariâ. *Acara-pats jotti*: Rheed. malab. 5. p. 15. t. 8.* Aff. videtur *T. Assæ*, sed differt foliis magis oblongis et integerrimis.
15. *T. ASSA* (DC. syst. 1. p. 402.*) foliis ovalibus utrinquè acuminatis subdenticulatis supernè lævibus, subtus pedunculisque pubescentibus, paniculâ 4-5-florâ, sepalis 4 glabriusculis. 5 in Indiâ orient. *T. Malabarica* Lam. ill. t. 485. f. 1. (v. s.)
16. *T. WAHLBOMIA* (DC. syst. 1. p. 403.*) foliis ellipticis acuminatis apice serratis basi stipulatis? subtus pedunculisque pubescentibus, paniculâ 4-5-florâ, sepalis 4 extus villosis. 5 in Javâ. *Wahlbomia Indica* Thunb. act. holm. 1790. p. 215*. t. 9.
17. *T. TIGAREA* (DC. syst. 1. p. 403.*) foliis subrotundis subrepandis utrinquè ramulisque scabris, paniculis ramosis multiflogis. 5 (in sylvis Guianæ et Cayennæ. *Tigarea aspera* Aubl! Guian. 2. p. 920. t. 350.* Calyx persistens sepalis exterioribus demum reflexis, interioribus conniventibus. (v. s.)
18. *T? TOMENTOSA* (Willd. spec. 2. p. 1241.) foliis ovatis acuminatis dentatis,

supernè glabris, subtùs ramulisque tomentosis. 5 in Cayennæ sylvia. Tigræa dentata Aubl! Guian. 2. p. 920. t. 351*. Flores ignoti. DC. syst. 1. p. 404. (v. s. in h. Banks.)

19. *T. CUSPIDATA* (Mey. prim. fl. esseq. p. 205*) foliis ovalibus cuspidato-serratis subtùs tomentosis, pedunculis fasciculatis axillaribus 1-floris. 5 in sylvia siccis insulæ Arowabiah Guianæ.

20. *T. RECTA* (Sessé et Moc. fl. mex. icon. et DC. syst. 1. p. 404.*) foliis obovatis, apice subobtusis acutè dentatis, basi attenuato-cuneatis, paniculâ ramosissimâ. 5 in Mexico. Flores in iconè videntur hermaphroditi aut polygamo-monoici. (Descr. ex. ic. pict.)

21. *T? LUTEA* (Spreng. nove entd. 1. p. 164.*) foliis elliptico-oblongis nitidis integerrimis, pedunculis lateralibus racemosis. 5 in Brasiliâ. Calyx coloratus; corollâ latea; flos monogynus! capsulæ 3. (Spreng.)

22. *T? PERUVIANA* (Spreng. nove entd. 1. p. 164.*) foliis ellipticis utrinquè nitidis obsolete serrulatis, corymbis axillaribus. 5 in America merid? — Calyx 4-5-sepalus; corollâ albida; flos monogynus! capsulæ 4. (Spreng.)

23. *T. ARBORESCENS* (Mal. misc. 1. n. 2. p. 45.) foliis obovatis integerrimis glabris, floribus paniculatis axillaribus et terminalibus. 5 in Sumatrâ. Arillus laciniatus flavescens.

II. DAVILLA. *Vand. fl. lus. et bras. prod.* 115. t. 2. f. 14.

Stam. ∞. Carpellum 1 testaceum 1-2-spermum sepalis 2 interioribus accretis concavis oppositis valvæformibus inclusum. Semen subglobosum. — Arbor Tetracerae aut Delimæ habitu valdè affinis et inter eas quasi media. DC. syst. 1. p. 404.

1. *D. BRASILIANA* (DC. syst. 1. p. 405.) 5 in Brasiliâ. Deless. icl. sel. 1. t. 71. *D. rugosa* Poir. suppl. 2. p. 456*. folia alterna oblonga secus petiolum subdecurrentia. Flores virescentes aut rufescentes. (v. s. in h. Juss.)

III. DOLIOCARPUS. *Roland. act. holm.* 1756. p. 249. t. 9.

Stam. ∞. Carpellum 1 baccatum, 1-2-spermum. — Frutices sæpius sarmentosi Tetracerae facie. DC. syst. 1. p. 405.

1. *D. ROLANDI* (Gmel. syst. 805.) caule scandente, foliis oblongis acuminatis apice dentatis, pedunculis lateralibus 1-floris, floribus 3-petalis. 5 in Surinamo. — Roland. act. holm. 1756. p. 260. t. 9. f. 1. 2. 3. Tetracera Dolio-carpus Willd. spec. 2. p. 1241. — DC. syst. 1. p. 405.

2. *D. STRICTUS* (Poir. suppl. 2. p. 498.) caule stricto, foliis ovato-lanceolatis dentatis deflexis, floribus terminalibus 3-petalis. 5 in Surinamo. Tetracera stricta Willd. spec. 2. p. 1241. — DC. syst. 1. p. 405.

3. *D. CALINEA* (Gmel. syst. 805.) caule scandente, foliis oblongis acuminatis integerrimis, pedunculis lateralibus ∞-floris, floribus 3-petalis. 5 in sylvia Guianæ. Calinea scandens. Aubl. Guian. 1. p. 556. t. 221.* — DC. syst. 1. p. 406.* (v. s. in h. Desf.)

4. *D? SORAMIA* (DC. syst. 1. p. 406.*) caule scandente, foliis obovatis integerrimis, pedunculis lateralibus subcorymbosis, floribus 5-petalis. 5 in Guianâ. Soramia Guianensis Aubl. Guian. 1. p. 552. t. 219.* (v. s. in h. Desf.)

IV. DELIMA. *Lin. amæn.* 1. p. 403.

Stam. ∞. Carpellum 1 capsulare 1-2-spermum. — Flores interdum abortu dioici.

1. *D. SARMENTOSA* (Lin! spec. 736.) foliis ovatis oblongis serratis rigidis scabris, floribus hermaphroditis paniculatis, ovarii capsulisque glabris. 5 in Ceylonâ. DC. syst. 1. p. 407. Burm. ind. 122. t. 37. f. 1. Tetracera sarmentosa Vahl. symb. 3. p. 70.* (v. s.)

2. *D. NEGUNGA* (DC. syst. 1. p. 407.) foliis obovatis subcrenulatis scabris, floribus hermaphroditis paniculatis, ovaria capsulisque pubescentibus. 5 in Javâ. Deless. ic. sel. 1. t. 72. Semina semi-arillata. (v. s.)
3. *D. MEXICANA* (Sessé et Moc. fl. mex. ic. ined. et DC. syst. 1. p. 407.*) foliis ovalibus obtusiusculis basi subattenuatis glabris serratis, floribus dioicis paniculatis secus ramulos sessilibus fasciculatis. 5 in Mexico. Semen ovatum crassum reticulatum. (Descr. ex ic. pict.)
4. *D. GUIANENSIS* (Rich! in DC. syst. 1. p. 408.) foliis oblongis utrinque acuminiatis glabris lævibus subserratis, floribus dioicis axillaribus breviter pedicellatis, fructibus globosis pubescentibus. 5 in Guianâ. Flores masculi ignoti. An potius *Doliocarpi* species? (v. s. ♀ in h. Rich.)
5. *D. NITIDA* (Vahl. symb. 3. p. 70.*) foliis lanceolato-oblongis scabris integerrimis, floribus hermaphroditis paniculatis. 5 in insulâ Trinitatis? stylus longitudine staminum; stigma subpeltatum. DC. syst. 1. p. 408.
6. *D. PIRIPU* (DC. syst. 1. p. 408.*) foliis ovali-oblongis margine undulato-crenatis mollibus, floribus paniculatis hermaphroditis pentandris. 5 in Malabarâ culta circa Pananie. Piripu. Rheed. mal. 7. p. 101. t. 54.*

V. CURATELLA. *Lin. gen. n.* 679. *Lam. ill. t.* 479.

Stam. ∞. Carpella 2 capsularia 1-2-sperma. Semina ovata nitida.

1. *C. AMERICANA* (Lindl. spec. 248.) foliis ovatis subrepandis subdenticulatis asperimis secus petiolum decurrentibus, racemis ramosis lateralibus. 5 in Americâ merid. Aubl! Guian. 1. p. 579. t. 232.* *Lam. ill. t.* 479. DC. syst. 1. p. 409* (v. s.)
2. *C. ALATA* (Vent! choix. p. 49.*) foliis ovalibus integerrimis lævibus utrinque obtusis, petiolis alatis, paniculâ terminali. 5 in Guianâ. An forsan, suspicante cl. R. Brown, *Wormiæ* species? DC. syst. 1. p. 409.* (v. s. in h. Deless.)

VI. TRACHYTELLA. *DC. syst. 1. p.* 410. †

Stam. ∞. Carpella 1-2-baccata ∞-sperma. — Genus omnino dubium et soli Loureiro cognitum.

1. *T. ACTÆA* (DC. syst. 1. p. 410.*) foliis lanceolatis subserratis, racemis spicatis linearibus interruptis. 5 inculta prope Cantonem Sinarum. *Actæa aspera* Lour. fl. coch. ed. Willd. 1. p. 405*.
2. *T. CALLIGONUM* (DC. syst. 1. p. 410.) foliis ovatis integerrimis, racemis subdivisis. 5 in sylvis Cochinchinæ. *Calligonum asperum* Lour. fl. coch. ed. Willd. 1. p. 418.* — Hujus cum *Delima* affinitatem *Loureirus* ipse memorat.

VII. RECCHIA. *Sessé et Moc. fl. mex. ic. ined. et DC. syst. 1. p.* 411. †

Stamina 10. Ovaria 2. †

1. *R. MEXICANA* (Sessé et Moc. l. c. DC. syst. 1. p. 411.) 5 in Mexico.

Trib. II. *DILLENEÆ*, *Salisb. par. lond. n.* 73.

Staminum filamenta apice non dilatata sed subattenuata; antheræ elongatæ adnatæ; carpella sæpius 2-5 distincta, rarius solitaria, aut 5-20 subcoalita. — Arbores frutices suffruticesve rarissimè scandentes. DC. syst. 1. p. 411.

VIII. PACHINEMA. *R. Brown. in DC. syst. 1. p.* 411.

Petala 5 ovata. Stamo. 7-10, filamentis basi crassissimis erectis apice attenuatis, antheris ovatis apice filamento angustato adnatis distinctis

conniventibus parallelisve. Ovaria 2 3 ovata in stylos subulatos desinentia. Fructus,...

1. *P. COMPLANATUM* (R. Brown! in DC. syst. 1. p. 412.) $\frac{5}{5}$ ad Carpentariam. Deless! ic. sel. 1. t. 73. Folia præter denticulos nulla; flores ex denticulorum axillis orti solitarii aut gemini. (v. s.)

IX. HEMISTEMMA. Juss. ined. DC. syst. 1. p. 412.

Stam. ∞ unilateralia, exteriora sterilia squamiformia; ovaria 2; styli filiformes. Semina arillo membranaceo cincta, albumen carnosum.

\S . 1. *Oppositifolia Madagascariensis, staminibus sterilibus apice spathulatis, petalis emarginatis.*

1. *H. COMMERSOHN* (DC. syst. 1. p. 413*) foliis oppositis ovali-oblongis mucronatis, pedunculis tomentosis. $\frac{5}{5}$ in Madagascar. Deless! ic. sel. 1. t. 74. Helianthemum coriaceum var. α . Pers! ench. 2. p. 76. Calyces tomentoso-lanati. (v. s.)
2. *H. AUBERTII* (DC. syst. 1. p. 413*) foliis oppositis oblongo lanceolatis basi attenuatis apice acutis, pedunculis glabrisculis. $\frac{5}{5}$ in Madagascar. Deless! ic. sel. 1. t. 75. Helianthemum coriaceum var. β . angustifolium Pers! ench. 2. p. 76. Calyces adpressè sericeo-pubescentes. (v. s.)

\S . 2. *Alternifolia Australasica, staminibus sterilibus linearibus, petalis obtusis.*

3. *H. DEALBATUM* (R. Brown! in DC. syst. 1. p. 413*) foliis alternis obovato-oblongis basi attenuatis apice obtusis mucronatis, subtus pube adpressissimè candicantibus. $\frac{5}{5}$ in Novæ-Hollandiæ terrâ Arnheimicâ. Deless! ic. sel. 1. t. 76. (v. s.)
4. *H. BANKSII* (R. Brown! in DC. syst. 1. p. 414*) foliis alternis oblongis basi angustatis apice obtusis subtus tomentosis. $\frac{5}{5}$ in Novâ Hollandiâ. Flores sessiles secundi. (v. s. comm. à cl. R. Brown.)
5. *H. ANGUSTIFOLIUM* (R. Brown! in DC. syst. 1. p. 414*) foliis alternis longis linearibus acutis subtus candicantibus. $\frac{5}{5}$ in Novâ-Hollandiâ. Deless! ic. sel. 1. t. 77. Pedunculi foliorum circiter longitudine. (v. s.)
6. *H? LECHERNAULTII* (DC. syst. 1. p. 414*) foliis alternis oblongis basi attenuatis apice truncatis subemarginatis subtus candicantibus. $\frac{5}{5}$ in Novâ-Hollandiâ et in insulis Sti. Francisci. Flores minimi axillares solitarii brevissimè pedicellati. (v. s.)

X. PLEURANDRA. Labill. Noy. Holl. 2. p. 5. t. 143 et 144.

Stam. 5-20 unilateralia, omnia fertilia. Styli filiformes. Carpella 2 membranacea 1-2-sperma. — Flores flavi ad apices ramulorum solitarii. DC. syst. 1. p. 415.

\S . 1. *Daphnoideæ; staminibus basi liberis, foliis oblongis obovatisve majusculis planis glabris.*

1. *P. BRACKETATA* (R. Brown! in DC. syst. 1. p. 415*) foliis oblongis subglabris mucronulatis, bracteis circa flores sessiles confertis extus sericeo-villosis, ovariiis hirsutissimis. $\frac{5}{5}$ in Novâ-Hollandiâ circa Port-Jackson. Deless! ic. sel. 1. t. 78. Variat foliis latioribus et angustioribus. (v. s.)
2. *D. MITIDA* (R. Brown! in DC. syst. 1. p. 416*) foliis oblongis glabris mucronatis, bracteis paucis subsarciosis glabris, ovariiis tomentosis. $\frac{5}{5}$ in Novâ-Hollandiâ. Priori affinis sed ex omni parte (ovariorum exceptis) glabra. (v. s.)
3. *P. CNEORUM* (DC. syst. 1. p. 416*) foliis oblongis basi angustatis obtusis bracteis calycibusque glabris, ovariiis subglabris. $\frac{5}{5}$ in Novâ-Hollandiâ. Flores ad apices ramulorum sessiles solitarii. (v. s. in h. Lamb.)

§. 2. *Alysoideæ; staminibus basi liberis, foliis oblongis obovatisve utrinque aut subtus pube stellatâ velutinis.*

4. *P. SERICEA* (R. Brown! in DC. syst. 1. p. 416.*) foliis oblongis margine revolutis supernè subpilosis subtus velutinis, ramis hirsutis, floribus sessilibus. §. in Novâ-Hollandiâ. Deless. ic. sel. 1. t. 79. Flores solitarii terminales inter folia sessiles. (v. s.)
5. *P. CINEREA* (R. Brown! in DC. syst. 1. p. 417.*) foliis oblongis margine revolutis utrinque velutinis, ramis subvillosis, floribus breviter pedicellatis. §. in Novâ-Hollandiâ. Flores solitarii, breviter pedicellati. (v. s.)
6. *P. FURFURACEA* (R. Brown! in DC. syst. 1. p. 417.*) foliis oblongis planis obtusis supernè subscabris subtus tomentoso-velutinis, ramis velutinis, floribus pedicellatis. §. in Novâ-Hollandiâ. Deless. ic. sel. 1. t. 80. Pedicelli pollicem ferè longi pilis stellatis velutini. (v. s.)
7. *P. PARVIFLORA* (R. Brown! in DC. syst. 1. p. 417.*) foliis obovatis basi cuneatis supernè scabris subtus ramisque velutinis, floribus pedicellatis. §. in Novâ-Hollandiâ. florès parvi 3 lin. diam. (v. s.)

§. 3. *Hibbertiæ; staminibus basi liberis, foliis linearibus oblongisve parvulis patentibus glabris aut pilis non stellatim scabris.*

8. *P. OVATA* (Labill! nov. holl. 2. p. 5. t. 143.*) foliis ovatis obtusis supernè subscabris subtus lævibus, floribus pedicellatis, calycibus pilosiusculis, ovariis pilosis. §. in insulâ Van-Diemen. Pedicelli terminales gracillimi nudi 1-flori. DC. syst. 1. p. 418.* (v. s.)
9. *P. SCABRA* (R. Brown! in DC. syst. 1. p. 418.*) foliis oblongis obtusis supernè scabris subtus subcanescentibus, floribus pedicellatis, calycibus subglabris, ovariis tomentosis. §. in Novâ-Hollandiâ. (v. s.)
 - α. foliis oblongo-linearibus.
 - β. foliis oblongo-ovalibus. *P. ovata* var. Labill. nov. holl. 2. p. 5. t. 143? media ferè inter *P. scabram* et *ovata*. An species propria?
10. *P. RIPARIA* (R. Brown! in DC. syst. 1. 419.*) foliis linearibus obtusiusculis margine subrevolutis subglabris, floribus sessilibus, calycibus glabris, ovariis hirsutis. §. in Novâ-Hollandiâ. Flores ad apices ramulorum sessiles solitarii. (v. s.)
11. *P. PEDUNCULATA* (R. Brown! in DC. syst. 1. p. 419.*) foliis linearibus obtusis glabris margine subrevolutis, floribus pedicellatis, calycibus ovariisque glabris. §. in ins-Van-Diemen. Stamina 7-8 lateralialia, filamentis longissimis. (v. s.)
12. *P. EMPETRIFOLIA* (DC. syst. 1. p. 420.*) foliis linearibus obtusis margine subrevolutis supernè subnitidis subtus velutinis, floribus pedicellatis, calycibus subglabris, ovariis hirsutis, staminibus 5. §. in Novâ-Hollandiâ. Pedicelli gracillimi foliis paulò longiores. (v. s. in h. Lamb.)
13. *P. INTERMEDIA* (DC. syst. 1. p. 420.*) foliis linearibus obtusis margine revolutis, supernè subnitidis subtus subpubescentibus, floribus pedicellatis, calycibus glabris, ovariis hirsutis, staminibus 10-15 subsparsis. §. in montibus Novæ-Hollandiæ. Flores parvi, sepala interiora margine membranacea. (v. s. in h. Lamb.)
14. *P. ERICAFOLIA* (DC. syst. 1. p. 420.*) foliis linearibus obtusiusculis margine revolutis supernè scabris subtus subpubescentibus, floribus sessilibus, calycibus subvillosis, ovariis hirsutis. §. in Novâ-Hollandiâ. Affinis *Hibbertiæ pedunculatæ*, sed notis genericis et flor. sessilibus differt. (v. s. in h. Lamb.)
15. *P. HYPERICOIDES* (DC. syst. 1. 421.*) foliis linearibus obtusis margine revolutis supernè scabris subtus incanis, floribus pedicellatis, calycibus subpubescentibus, ovariis velutinis. §. in Novâ-Hollandiâ. Deless. ic. sel. 1. t. 81. Suffrutex ramosissimus Rosmarinum Hypericæ angustifoliæ æmulans. (v. s.)
16. *ANERVIA* (DC. syst. 1. p. 421.*) foliis linearibus subscutis enerviis glabris,

floribus sessilibus, calycibus obtusis mucronulatis ovariisque glabris. ζ . in Novâ-Hollandiâ. Caulis humiles ramosissimi. (v. s. in h. Lamb.)

17. *P. ACICULARIS* (Labill. nov.-holl. 2. p. 6. t. 144.*) foliis linearibus apice acuminato-aristatis lævibus, floribus pedicellatis, calycibus subhispidis mucronatis, ovariis hirsutis. ζ . in capite Van-Diemen Novæ-Hollandiæ. DC. syst. 1. p. 421.* (v. v.)

18. *P. ACEROSA* (R. Brown! in DC. syst. 1. p. 422.*) foliis lineari-subulatis margine revolutis supernè scabris subtus lævibus, floribus pedicellatis, calycibus subscabris subacutis, ovariis villosis. ζ in Novâ-Hollandiâ. Suffrutex humilissimus ramosissimus. (v. s.)

\S . 4. *Candolleæ*; *staminibus imâ basi monadelphis, foliis linearibus erectis vix patentibus*. — *An genus proprium?*

19. *P. STRICTA* (R. Brown! in DC. syst. 1. p. 422.*) foliis linearibus subobtusis erectis supernè scabridis subtus lævibus, floribus sessilibus, calycibus subscabris, ovariis velutinis. ζ . in Novâ-Hollandiâ. Suffrutex erectus strictus ramosus. (v. s.)

20. *P. CALYCINA* (DC. syst. 1. p. 422.*) foliis linearibus acutis erectiusculis supernè scabris subtus subpubescentibus, floribus sessilibus, calycibus sericeo-velutinis, ovariis hirsutis. ζ . in Novæ-Hollandiæ montibus. Caulis erectus ramosus. (v. s. in h. Lamb.)

XI. CANDOLLEA. Labill. nov.-holl. 2. p. 33. t. 176.

Stam. ∞ polyadelphe. Styli filiformes. Carpella 2-5 ovata intus dehiscentia. — Suffrutices è Novâ-Hollandiâ orti, Pleurandræ et Hibbertiæ facie. DC. syst. 1. p. 423.

1. *C. CUNEIFORMIS* (Labill! nov.-holl. 2. p. 34. t. 176.*) foliis glabris obovato-cuneatis apice obtusis integris. ζ in Novâ-Hollandiâ occid. Hibbertia cuneiformis Smith. in Rees cycl. vol. 17. Frutex sesquiorgyalis, ramis cinereis rugosis. DC. syst. 1. p. 423. (v. s.)

2. *C. PEDUNCULATA* (R. Brown! in DC. syst. 1. p. 424.*) foliis glabris linearicuneatis apice truncatis 3-dentatis. ζ in Novâ-Hollandiâ. Suffrutex ramosus, rami juniores parce villosi, adulti glabri. (v. s.)

3. *C. FASCICULATA* (R. Brown! in DC. syst. 1. p. 424.*) foliis villosis linearibus integris. ζ in Novâ-Hollandiâ. Rami teretes juniores villosi.

XII. ADRASTAEA. DC. syst. 1. p. 424.

Stam: 10 libera æqualia; filamenta plana margine loculos oblongatos gerentia. Ovaria 2, styli recti conico-subulati. — Suffrutex paludosus Hibbertiæ ferè facie.

1. *A. SALICIFOLIA* (DC. syst. 1. p. 424.*) foliis linearibus subintegris punctis nempe utrinque 3-5 callosis dentiformibus ad apicem instructis. ζ in paludibus circa Botany-bay. (v. s. in h. Lamb.)

XIII. HIBBERTIA. Andr. bot. rep. t. 126.

Stam. ∞ libera filiformia æqualia; antheræ ovali-oblongæ. Ovaria 1-15; styli filiformes inflexi. Carpella membranacea dehiscentia, sæpius 1-2-sperma. Semina exarillata. DC. syst. 1. p. 425.

\S . 1. *Carpellis 10-15, basi glabris apice subpilosis*. (Burtonia Salisb.) — *An genus proprium?*

1. *H. GROSSULARIÆFOLIA* (Salisb. par. lond. n. 73. t. 73.*) foliis suborbiculatis crenato-dentatis, floribus pedunculatis oppositifoliis, caulibus procumbentibus. ζ \mathcal{H} . in Novâ-Hollandiâ. Sims. bot. mag. t. 1218. *H. crenata* Andr. bot. rep. t. 472. — DC. syst. 1. p. 425.* (v. v.)

§. 2. *Carpellis* 1-8 *glabris*.

2. *H. VOLUBILIS* (Andr. bot. rep. t. 126.*) foliis obqvato-lanceolatis subintegerrimis mucronatis subtus pubescentibus, floribus sessilibus 5-8-gynis, caule subvolubili. ζ in Novâ-Hollandiâ. *Dillenia speciosa* Curt. bot. mag. t. 449. excl. syn. Semina nigra pisiformia compressa dura exarillata. DC. syst. 1. p. 426.* (v. s. et v.)
3. *H. DENTATA* (R. Brown! in DC. syst. 1. p. 426.*) foliis oblongis acuminatis glabris aristulato serratis, floribus pedunculatis 3-gynis, caule volubili. ζ in Novâ-Hollandiâ, ad montes cæruleos. Ker. bot. reg. t. 282.* Lodd bot. cab. t. 347. Sims. bot. mag. t. 2338. (v. v. in h. Kew.)
 β . pedicellis calycibusque glabris aut vix pilosis, foliis magis serratis.
4. *H. SALICNA* (R. Brown! in DC. syst. 1. p. 427.*) foliis oblongo-linearibus apice mucronato-acuminatis integerrimis subtus villosis, floribus sessilibus 2-4-gynis, caule erecto. ζ in Novâ-Hollandiâ. (v. s.)
5. *H. PROCUMBENS* (DC. syst. 1. p. 427.*) foliis linearibus acuminatis crassiusculis glabris aveniis, floribus subsessilibus 3-5-gynis, caulibus procumbentibus. ζ in Capite Van-Diemen. *Dillenia procumbens*. Labill! nov.-holl. 2. p. 16. t. 156.* Semina globosa castanea in pulpâ carnosâ immersa. — An genus proprium? (v. s.)
6. *H. VIRGATA* (R. Brown! in DC. syst. 1. p. 428.*) foliis linearibus subobtusis glabris, floribus sessilibus 7-8-andris 3-4-gynis, caule erecto. ζ in Novâ-Hollandiâ. Fruticulus gracilis erectus glaber. (v. s.)
7. *H. FASCICULATA* (R. Brown! in DC. syst. 1. p. 428.*) foliis lineari-subulatis glabris subfasciculatis, ramulis subpilosis, floribus sessilibus 3-4-gynis, caule erecto. ζ in Novâ-Hollandiâ orâ orient. — Habitus ferè Hyperici fasciculati. (v. s.)
8. *H. LINEARIS* (R. Brown! in DC. syst. 1. p. 428.*) foliis linearibus acutis integerrimis glabris, floribus sessilibus 2-gynis 20-andris, caule erecto. ζ in Novâ-Hollandiâ orâ orient. Suffrutex erectus ramosissimus. (v. s.)
9. *H. OBTUSIFOLIA* (DC. syst. 1. p. 429.*) foliis linearibus obtusis apice angustatis glabris, floribus sessilibus 20-andris 2-gynis, caule erecto. ζ in arenosis siccis terræ Van-Diemen. *H. lineari* simillima et forsan hujus varietas? (v. s.)
10. *H. DIFFUSA* (R. Brown! in DC. syst. 1. p. 429.*) foliis cuneatis apice dentatis glabris, floribus sessilibus 2-3-gynis 20-andris, caule diffuso. ζ in Novâ-Hollandiâ orâ orient. (v. s.)
 β . *oblongata*, foliis oblongo-cuneatis. Differt à priori foliis suboblongis integris emarginatis aut lateraliter dentes 2-3 gerentibus. (v. s. in h. Lamb.)
11. *H. MONOGYNA* (R. Brown! in DC. syst. 1. p. 429.*) foliis spathulatis basi cuneatis apice 2-3-dentatis glabris, floribus sessilibus 1-gynis 12-andris, caule erecto. ζ in Novâ-Hollandiâ montibus. Carpellum unicum globosum membranaceum indehiscens 1-spermum. (v. s.)

§. 3. *Carpellis* 2-4 *pube brevi velutinis squamulisve lepidotis*.

12. *H. PEDUNCULATA* (R. Brown! in DC. syst. 1. p. 430.*) foliis linearibus obtusiusculis margine subrevolutis, floribus pedicellatis 2-gynis, ovariis subincanis. ζ in Novâ-Hollandiâ. Ovaria 2 globosa subincana, styli filiformes stamina vix superantes. (v. s.)
13. *H. SERPYLLIFOLIA* (R. Brown! in DC. syst. 1. p. 430.*) foliis ellipticis obtusis integerrimis glabris, floribus pedicellatis 2-gynis 8-10-andris, ovariis villosis. ζ in Novâ-Hollandiâ orâ orient. Antheræ oblongæ filamentis longioribus. (v. s.)
14. *H. ASPERA* (DC. syst. 1. p. 430.*) foliis oblongis obtusis margine subrevolutis superne subcæbris subtus cinereo-velutinis, floribus pedicellatis 2-gynis, ovariis hirsutis. ζ in Novâ-Hollandiâ. An potius *Pleurandæ* species?

sed adeo ex habitu *H. tomentosa* similis ut separare noluerim. (v. s. in h. Lamb.)

15. *H. HERMANNIIFOLIA* (DC. syst. 1. p. 431.*) foliis obovato-cuneatis obtusis utrinque villosis-tomentosis, villis supernè simplicibus subtus stellato-ramosis, floribus pedicellatis 2-gynis, ovariis tomentosis. ♂ in montibus Novæ-Holland. (v. s. in h. Lamb.)
16. *H. OBLONGATA* (R. Brown! in DC. syst. 1. p. 431.*) foliis oblongis integerrimis supernè glabris subtus pilis stellatis velutinis, nervulis ad marginem confluentibus, floribus pedicellatis 2-gynis, ovariis lepidotis. ♂ in Carpentariâ. Carpella globosa subtriquetra. (v. s.)
17. *H. CISTIFOLIA* (R. Brown! in DC. syst. 1. p. 431.*) foliis oblongis integerrimis uninerviis, pilis stellatis supernè pubescentibus subtus albedo-velutinis, floribus pedicellatis digynis, ovariis lepidotis. ♂ in Novæ-Holland. terrâ Arnheimicâ. (v. s.)
18. *H. TOMENTOSA* (R. Brown! in DC. syst. 1. p. 432.*) foliis oblongo-linearibus uninerviis integerrimis utrinque tomento brevi velutinis, floribus pedicellatis digynis, calycibus ovariisque lepidotis. ♂ in Carpentariâ. Ovaria globosa, styli glabri filiformes. (v. s.)
19. *H. LEPIDOTA* (R. Brown! in DC. syst. 1. p. 432.*) foliis linearibus acuminatis integerrimis utrinque lepidotis, pedicellis confertis unifloris, floribus digynis, ovariis lepidotis. ♂ in Carpentariâ. Styli arcuati filiformes glabri. (v. s.)

XIV. WORMIA. Rottb. nov. act. hafn. 1783. v. 2. p. 522. t. 3.

Stam. ∞ libera filiformia æqualia; styli 5 filiformes; stigmata emarginata. Carpella 5 capsularia intus dehiscentia 8-12-sperma; pulpa seu arillus ad basim seminum. DC. syst. 1. p. 433.

1. *W. MADAGASCARIENSIS* (DC. syst. 1. p. 433.*) foliis ovalibus obtusè sinuatis, pedunculis paniculato-ramosis. ♂ in Madagascar. Deless! ic. sel. 1. t. 82. *Lenidia Madagascariensis* Poir. suppl. 3. p. 330.* (v. s. in h. Mus. Par. et Juss.)
 - a. *rotundifolia*, foliis orbiculatis basi subcordatis.
 - β. *oblongifolia*, foliis oblongis basi non cordatis.
2. *W. DENTATA* (DC. syst. 1. p. 434.*) foliis ovatis retusis grossè et subacutè dentatis, pedunculis triquetris 3-6-floris. ♂ in Ceylonâ. *Dillenia dentata* Thunb! act. soc. lin. 1. p. 201. t. 20. Petala 5 orbiculata, ovaria 5. (v. s. in h. Deless.)
3. *W. TRIQUETRA* (Rottb. nov. act. hafn. 2. p. 532. t. 3.*) foliis ovatis subobtusis obtusè subsinuatis, pedunculis triquetris racemosis. ♂ in Ceylonâ. Ovaria trigona approximata. — An satis a priore distincta? DC. syst. 1. p. 434.* (v. s.)
4. *W. ALATA* (R. Brown! in DC. syst. 1. p. 434.*) foliis ovalibus integerrimis, petiolis alato-marginatis. ♂ in Novâ-Holland. Styli subulati divergentes staminibus longiores. (v. s.)

XV. COLBERTIA. Salisb. parad. n. 73.

Stamina ∞ libera, 10 cæteris multò longiora. Carpella 5 coalita? stigmata capitata. Semina in quoque loculo plurima reniformia in pulpâ pellucidâ glutinosa nidulantia. DC. syst. 1. p. 435.

1. *C. COROMANDELIANA* (DC. syst. 1. p. 435.*) ♂ in vallibus montium Coromandelicæ. *Dillenia pentagyna* Roxb. fl. corom. 1. p. 21. t. 20.* Fructus penduli. (v. s. sine flor. in h. Lamb.)

XVI. DILLENIA. Lin. gen. n. 688.

Stam. ∞ libera æqualia. Carpella 10-20 coalita in baccam spuriam multilocularem stigmatibusque radiantibus coronatam. DC. syst. 1. p. 435.

1. *D. SPECIOSA* (Thunb. act. soc. lin. p. 1. 200.) foliis elliptico-oblongis simpliciter serratis, pedunculis unifloris. \S in Malabariâ. Syalita Rheed. mal. 3. p. 39. t. 38. 39.* Rami cinerei expansi rugosi. DC. syst. 1. p. 436.* (v. s. in h. Mus. Par. Deless! etc.)
 2. *D. AUREA* (Smith. exot. bot. t. 92. 93.*) foliis elliptico-oblongis serratis, floribus sæpius solitariis pedunculatis autè folia ortis. \S in Indiâ orient. Semina margine non fimbriata. DC. syst. 1. p. 436.
 3. *D. INTEGRÆ* (Thunb. act. soc. Lin. 1. p. 199. t. 18.*) foliis oblongo-obovatis obtusis subintegris, pedunculis unifloris. \S in Zeylonâ. — An Wormia species? DC. syst. 1. p. 437.
 4. *D. ELLIPTICA* (Thunb. act. soc. Lin. 1. p. 200.) foliis elliptico-ovatis acutis serratis, pedunculis unifloris. \S in Celebe orientali prope Tambocco. Semina plana fusca, 8 in quoque loculo. DC. syst. 1. p. 437.*
 5. *D. SERRATA* (Thunb. act. soc. Lin. 1. p. 201.) foliis elliptico-ovatis acutis serratis, pedunculis trifloris. \S in Javâ. Sangius Rumph. amb. 2. p. 142. t. 46.* Fructus aurantii formâ et magnitudine. DC. syst. 1. p. 437.*
 9. *D. RETUSA* (Thunb. act. soc. Lin. 1. p. 200. t. 19.*) foliis obovatis apice truncatis remotè dentatis, pedunculis unifloris. \S in Zeylonæ sylvis. Lam. ill. t. 492. f. 2. Petala obovata calyce tertiâ parte tantum longiora. DC. syst. 1. p. 437.
-

Ord. III. MAGNOLIACEÆ. DC. *theor.* 213. *Ess. prop. ed.* 2. p. 72. *syst.* 1. p. 439.

Partes floris numero ternario dispositæ; sepala 3-6, decidua; petala 3-27, multi-serialia hypogyna; stamina ∞ libera, toro infra ovaria inserta; antheræ filamentis adnatæ, elongatæ; ovaria ∞ toro supra stamina inserta, sæpius spicatum disposita, monostyla; styli breves et stigmata simplicia; carpella tot quot ovaria 1-locularia, 1- ∞ -sperma, nunc capsularia rimâ superâ dehiscentia, nunc capsularia bivalvia rimâ inferâ dehiscentia, nunc follicularia, nunc subcarnosa indehiscentia, nunc samaræformia, aggregata subcoalitave in strobilum laxum densumve; semina angulo carpelli interno adnata; albumen carnosum; embryo rectus parvus inferus. — Arbores fruticesve elegantes. Folia alterna pennatim nervosa. Flores pulchri sæpius intensè odorati.

Trib. I. *ILLICIEÆ*.

Carpella verticillatim disposita, rarissimè abortu solitaria. — Folia pellucido-punctata.

I. *ILLICIUM*. Lin. *gen.* 611. *Gertn. fruct.* 1. p. 339. t. 96.

Sepala 3-6 petaloidea. Carpella stellatim disposita capsularia, supernè dehiscentia 1-sperma. Semen nitidum. — Frutices sempervirentes glaberrimi. DC. *syst.* 1. p. 440.

1. I. *FLORIDANUM* (Ellis. act. angl. 1770. p. 524. t. 12. *) petalis 27-30 purpureis, exterioribus oblongis, interioribus lanceolatis. ♂ in Floridâ occid. juxta flumen Mississipi. Lam. ill. t. 493. f. 1. herb. amat. t. 174. Lodd. bot. cab. t. 209. Folia trita anisum redolentia. DC. *syst.* 1. p. 441. * (v. v.)
2. I. *ANISATUM* (Lin. spec. 664) petalis 27-30 flavescentibus, exterioribus oblongis, interioribus lineari-subulatis. ♂ 5 in Japoniâ, et Chinâ. Gaertn. fruct. 1. p. 338. t. 69. — Clus. hist. 2. p. 202. f. 3. Capsulæ è Chinâ sub nomine Anisi Chinensis adportatæ. DC. *syst.* 1. p. 441. * (v. s. in h. Deless. et Lamb.)
3. I. *PARVIFLORUM* (Michx! fl. bor. amer. 1. p. 326.) petalis 9-12 flavescentibus ovato subrotundis. ♂ in Floridâ occidentali. Vent! Cels. t. 22. herb. am. t. 330. — Sepala 3 ovata subciliata. DC. *syst.* 1. p. 442. * (v. v.)

II. *TEMUS*. Mol. chil. 153. Juss. *gen.* 435.

Calyx 3-fidus. Petala 18 linearia longissima. Stam. 26 (27?) petalis breviora, antheræ globosæ. Ovaria 2 monostyla. Carpella 2 baccata (coalita?). Semina arillata — Arbor sempervirens, folia alterna pectiolata ovalia lævia Myristicæ odorem spirantia. Flores terminales. — Cl. Molina dixit ovaria 2 et baccam 1 dicoccam, quod verosimiliter est erroneum; an ex ovariiis 2 unum abortit et alterum in baccam dicoccam mutatum? An potius ovaria 2 grandefacta et carnosa coalescunt in fructum unicum hinc dicoccum — Affinitas cum Illicio et Drimyde manifesta videtur e characteribus classicis et etiam ex habitu et viribus.

1. *T. MOSCHATA* (Mol. chil. 153.) ♂ in Chili. Gmel. syst. 1. p. 831. Poir. dict. 7. p. 595. — Folia in ramis conferta 2 poll. longa viridia lucida; flores suaveolentes; calycis lobi obtusi; petala flava aut alba angusta 2-3. poll. longa; staminum filamenta setacea petalis dimidio breviora; bacca Coffeæ similis sed amarissimâ. (Mol.)

III. DRIMYS. Forst. gen. t. 42. Lam. ill. t. 494.

Carpellâ congesta baccata polysperma. Staminum filamenta apice crassiora loculosque discretos gerentia. — Genus structurâ staminum et calycis subpersistencia paululûm ad Delimaceas spectans. DC. syst. 1. p. 442.

Sect. 1. EUDRYMIS. DC. syst. 1. p. 442.

Calyx integer dehiscens. Flores minimi.

1. *D. AXILLARIS* (Forst! gen. t. 42.) foliis oblongis utrinque acuminatis, pedicellis unifloris. ♂ in Novæ-Zelandiæ nemoribus. Lam. ill. t. 494. f. 2. Ovaria tetrasperma. DC. syst. 1. p. 443.* (v. s. in h. Mus. Par. Banks et Lamb.)

Sect. 2. WINTERA. DC. syst. 1. p. 443.

Calyx 2-3-partitus aut 2-3-sepalus.

2. *D. WINTERI* (Forst gen. p. 84. t. 42.) foliis oblongis obtusis subtus glaucis, pedunculis subsimplicibus aggregatis, aut brevissimis in pedicellos elongatos divis. ♂ in Freti Magellanici vallibus soli expositis. Winterana aromatica Sol. med. obs. 5. p. 46. t. 1. DC! syst. 1. p. 443.* (v. s.)

♂. *punctata*, Lam. dict. 2. p. 330, ill. t. 494. f. 1. Minus glauca, magis distinctè punctata.

3. *D. GRANATENSIS*. (Linn. fl. suppl. 269.) foliis lanceolato-oblongis utrinque acutis subtus glaucis, pedunculis elongatis trifidis. ♂ in Novâ-Granatâ et in regni Santa-Fæ de Bogotâ montibus ad altitudinem 1500 hexap. Humb. et Bonpl! pl. æquin. 1. p. 205. t. 58. DC. syst. 1. p. 444.* (v. s. in h. Mus. Par.)
4. *D. CHILENSIS* (DC. syst. 1. p. 444.*) foliis oblongo-obovatis subtus glaucis, pedicellis confertis 1-floris aut è pedunculo communi ortis, petalis 6-9 oblongis subobtusis. ♂ in regni Chilensis paludosis. Deless! ic. sel. 1. t. 83. Baccæ ovals subcompressæ obtusæ. (v. s. in h. Mus. Par.)
5. *MEXICANA* (Moc. et Sessé pl. mexic. ined. ic. et DC. syst. 1. p. 444.*) foliis oblongo-lanceolatis utrinque acuminatis, pedunculis elongatis apice pedicellos 4 elongatos gerentibus, petalis 20-24 acutis. ♂ in Mexico. Baccæ 4 an abortu 2-3. obovatæ basi attenuatæ cæruleo violacæ. DC. syst. 1. p. 444.* (v. ic. pict.)

IV. TASMANNIA. R. Brown. in DC. syst. 1. p. 445.

Flores dioici vel polygami minimi Drimydis floribus similes. ♂ *stam.* ∞ cum rudimento pistilli. ♀ *sepala* 2 petalque 2-5-decidua; *ovarium* 1 uniloculare; stigma ad latum internum ovarii longitudinaliter adnatum. Fructus membranaceus indehiscens polyspermus 1-locularis; stamina subcurva, testâ nitida fragili albumineque carnosio donata, lateraliter adfixa biserialia? subhorizontalia. — Genus Drimydi valde affine ex cl. R. Brown, sed differt floribus dioicis, fœmineis monocarpis.

1. *T. AROMATICA* (R. Brown! prod. nov.-holl. 2. ined. DC. syst. 1. p. 445.*) foliis oblongis basi in petiolum sensim attenuatis, fructibus globosis subdidymis. ♂ in Novâ-Hollandiæ frigidioribus. Deless! ic. sel. 1. t. 84. Winterana lanceolata Poir! dict. 8. p. 799.* (v. sine flor.)
2. *T. INSIPIDA* (R. Brown! prodr. nov.-holl. vol. 2. ined. DC. syst. 1. p. 445.*) foliis ovali-oblongis ad apicem petioli brevissimi abruptè subauriculatis, fructibus ovali oblongis. ♂ in Novâ-Hollandiâ. T. dipetala R. Br. in litt. (v. s.)

Trib. II. *MAGNOLIEÆ*.

Carpella secus axim spicatim disposita. — Folia non pellucido-punctata.

V. *MAYNA*. *Aubl. guian.* 2. p. 922. t. 352. †.

Flores dioici. *Carpella* plurima? ♀ ignoti. — Genus ex solo *Aubletio* notum, inter *Magnoliaceas* et *Annonaceas* medium, fructibus autem ignotis dubium, ob folia stipulacea *Magnoliaceis* relatum, sed antheris tetragonis, et florescentiâ forsan ad *Annonaceas* referendum. DC. syst. 1. p. 446.

1. *M. ODORATA* (*Aubl. guian.* 2. p. 922. t. 352.) foliis oblongis integerrimis glabris. ♂ in Cayennâ prope hortum prædii Loyola.
2. *M. SERICEA* (*Spreng. neue entd.* 2. p. 153.*) foliis lanceolatis integerrimis subtus serriceis. ♀ in Brasiliâ. — Flos odoratus subhexapetalus; ovaria subæna; flores pedunculati aggregati.
3. *M. BRASILIENSIS* (*Raddi in att. soc. ital.* 18. p. 23. f. 1.*) foliis oblongis subdenticulatis basi leviter attenuatis, pedunculis 3-floris. ♂ in Brasiliâ prope Rio-Janeiro in sylvis densis montis Escorodo. Arb. 30-40 ped.

VI. *MICHELIA*. *Lin. gen.* 691. *Gaertn. fruct.* 2. p. 263. t. 137.

Carpella spicatim laxiusculè disposita subbaccata, apice dehiscentia polysperma. — Flores odorati et ideò arbores in Indiâ ornamentariæ. DC. syst. 1. p. 447.

1. *M. CHAMPACA* (*Lin! spec.* 756.) foliis lanceolatis glabris. ♂ in locis arenosis Indiæ. *Lam. ill.* t. 493. Petala oblonga 6-8 lin. longa, 3. lin. lata. DC. syst. 1. p. 447. (v. s.)
2. *M. KISORA* (*Buchanan! in DC. syst.* 1. p. 448.*) foliis ovato-lanceolatis glabris secus petiolum non attenuatis. ♂ in Napauliâ. Valdè similis *M. Champacæ*, sed petalis 9 oblongioribus et sentioribus. (v. s. in h. Lamb.)
3. *M. DOLTSOPA* (*Buchanan! in DC. syst.* 1. p. 448.*) foliis ovali-oblongis acuminatissimis glabris subtus glaucescentibus. ♂ in Napauliâ. Petala 6-9, exteriora oblonga, interiora angustiora. (v. s. in h. Lamb.)
4. *M. TSJAMPACA* (*Lin! mant.* 78.) foliis lanceolato-ovatis, junioribus subtus pube adpressa sericeis. ♂ in sylvis Amboinensibus. *Sampacca* sylvestris. *Rumph. amb.* 2. p. 202. t. 68.* *M. sericea* *Pers. ench.* 2. p. 94. An plures hic confusæ species? DC. syst. 1. p. 448.* (v. s. in h. Deless. et Lin.)
5. *M. VELUTINA*, foliis elliptico-oblongis acutis supernè glaberrimis subtus etiam adultis tomento albedo velutinis. ♂ in Napauliâ. Alabastra tomentosa, flores axillares solitarii subsessiles. (v. s. comm. à cl. Wallich.)
6. *M. RUFINERVIS* (*DC. syst.* 1. p. 449.) foliis elliptico-oblongis utrinque acuminatis subtus in nervis petiolis turionibusque pube flavo-rufâ velutinis. ♂ culta in insulis Mauritanis-Fol. nervulis reticulatim anastomosantibus. (v. s. sine fl.)
7. *M. PARVIFLORA* (*Rumph. amb.* 2. p. 199? DC. syst. 1. p. 449. n. 8.) foliis ellipticis subacuminatis glabris, ramulis tomentosis. ♂ in Javâ. — Deless! ic sel. 1. t. 85.* (v. s. in h. Deless.)

VII. *MAGNOLIA*. *Lin. gen.* n. 690. *Gaertn. fruct.* 1. p. 343. t. 70.

Carpella spicatim confertèque disposita, angulo externo dehiscentia, 1-2-sperma, persistentia. Semina baccata subcordata, funiculi umbilicalis longissimi ope extrâ capsulam pendula. DC. syst. 1. p. 449.

Sect. I. MAGNOLIASTRUM. DC. syst. I. p. 450.

Species Americanæ, bractea unicâ alabastrum includente, ovariis approximatis, antheris extrorsis.

1. *M. GRANDIFLORA* (Lin. spec. 755.) foliis perennantibus ovali-oblongis coriaceis suprâ nitidis, subtus ferrugineis, floribus erectis 9-12-petalis expansis. ♂ in Americæ boreal. locis pinguibus subudis. Lam. ill. t. 490. Savi bibl. ital. 1819. n. 47. p. 219. DC. syst. I. p. 451.* (v. v.)
 - α. *obovata*, foliis obovato-oblongis, floribus expansis. Ait. h. kew. ed. 2. vol. 3. p. 329.
 - β. *elliptica*, foliis oblongo ellipticis, floribus subcontractis. Ait. l. c.
 - γ. *lanceolata*, foliis oblongo-lanceolatis, floribus subcontractis. Ait. l. c.
2. *M. MEXICANA* (DC. syst. I. p. 451.*) foliis ovalibus basi subattenuatis obtusis, floribus 9-petalis contractis, petalis ovatis planis. ♂ in Mexico. *M. grandiflora* Moc. et Sessé fl. mex. descr. ined. *M. glauca* fl. mex. icon. ined. An potius *Talauma* species altera?
3. *M. GLAUCA* (Lin! spec. ed. 2. p. 755.) foliis ellipticis obtusis subtus glaucis, floribus 9-12-petalis contractis, petalis ovatis concavis. ♂ in New-Jersey et Carolinâ. Schkuhr. handb. 2. n. 1441. t. 148. Lodd. bot. cab. t. 215. Bonpl. nav. p. 103. t. 42*. *M. Virginiana*, α. *glauca*. Lin. spec. ed. 1. p. 538. — DC. syst. I. p. 452.* (v. s. et v.)
 - β. *longifolia*, foliis perennantibus ellipticis utrinquē acutis. Ait. h. Kew. ed. 1. v. 2. p. 251. Pursh! fl. bor. amer. 2. p. 381. — in Georgiâ et Floridâ.
 - γ. *argentea*, foliis oblongis subtus sericeis albido glaucis. Pursh! in herb. Lamb.
4. *M. UMBRELLA* (Lam. dict. 3. p. 673.*) foliis deciduis lanceolatis patentissimis adultis glabris, junioribus subtus pubescentibus, petalis 9-12-exterioribus dependentibus. ♂ in Americâ boreali *M. tripetala* Lin. spec. ed. 2. p. 756. Michx! f. arb. am. 3. p. 90. t. 5.* — DC. syst. I. p. 452. (v. s. et v.)
5. *M. ACUMINATA* (Lin! spec. ed. 2. p. 756.) foliis deciduis ovalibus acuminatis subtus pubescentibus, floribus 6-9-petalis. ♂ in Americâ boreal. Michx. f. arb. am. 3. p. 82. t. 3.* Lodd. bot. cab. t. 418. — DC. syst. I. p. 453.* (v. s.)
 - β. *Candolii*, foliis ovato-oblongis acutis, floribus subvirescentibus. M. De Candolii Savi. bibl. ital. 1819. n. 47. p. 224. cum ic. (v. v. in hort. Lavurne.)
6. *M. AURICULATA* (Lam! dict. 3. p. 673.*) foliis deciduis glabris subtus subglaucescentibus spatulato-obovatis basi cordatis, auriculis obtusis approximatis, sepalis 3 patentissimis, petalis 9 oblongis. ♂ in Americâ boreali. *M. auricularis* Salisb! parad. lond. 43. t. 43.* — Sims. bot. mag. t. 1206. DC. syst. I. p. 454. (v. s. et v.)
7. *M. PYRAMINATA* (Bartr. itin. ed. germ. 390.) foliis deciduis glabris utrinquē concoloribus spatulato obovatis, basi cordatis, auriculis divaricatis, sepalis 3 patentibus, petalis 9 lanceolatis acuminatis. ♂ in Carolinâ et Georgiæ partibus occidentibus. *M. auriculata* var. Michx! fl. bor. amer. 1. p. 328. Mera prioris var. ex Nutall. — DC. syst. I. p. 454. (v. s.)
8. *M. MACROPHYLLA* (Michx! fl. bor. amer. 1. p. 327. Mich! f. arb. amer. 3. p. 79. t. 7.*) foliis deciduis amplissimis oblongo-obovatis subcuneatis basi cordatis subtus albido-glaucis, petalis 6-9-ovatis. ♂ in Americâ boreal. DC. syst. I. p. 454.* (v. s. et v.)
9. *M. CORDATA* (Mich! fl. bor. am. 1. 328. Mich. f. arb. am. 3. p. 87. t. 4.*) foliis deciduis cordiformibus subovatis subtus subtomentosis suprâ lævibus, petalis 6-9-oblongis. ♂ in Americâ boreal. Ker. bot. reg. t. 325.* Lodd. bot. Cab. t. 474. DC. syst. I. p. 455.

Sect. II. GWILLIMIA. Rottler. DC. syst. 1. p. 455.

Species Asiaticæ, bracteis sæpius duabus oppositis alabastrum includentibus, antheris introrsis, ovariis subdistantibus. — An species quædam bractea monophyllâ donatæ Micheliis adnumerandæ. DC. l. c.?

10. *M. YULAN* (Desf! arb. 2. p. 6.) foliis deciduis obovatis abruptè acuminatis, junioribus pubescentibus post anthesin prodeuntibus, floribus erectis 6-9-petalis, stylis erectis. ♂ in Chinâ. Bonpl! nav. p. 53. t. 20.* *M. conspicua*. Salisb. par. t. 38.* — DC. syst. 1. p. 455.* (v. v.)

11. *M. KOBUS* (DC. syst. 1. p. 456.*) foliis deciduis obovatis utriusque acuminatis, post anthesin prodeuntibus, junioribus subtus pubescentibus, adultis glabris, floribus erectis 3-sepalis 6-petalis, stylis reflexis. ♂ in Japoniâ insulâ Nipon. *M. gracilis* Salisb! parad. t. 87. *Kobus*. Banks. ic. Kœmpf. t. 42. (v. s.)

12. *M. OBOVATA* (Thunb. act. soc. Lin. 2. p. 336.) foliis deciduis obovatis acutis reticulato-venosis subglabris, floribus erectis 3-sepalis, petalis 6 obovatis, stylis brevissimis. ♂ in Japoniâ. DC. syst. 1. p. 456.*

α. *denudata*, ramis floridis aphyllis, petalis obovatis. *M. denudata*. Lam. dict. 3. p. 675. — Banks. ic. Kœmpf. t. 43 — in insulâ Nipon.

β. *discolor*, ramis floridis foliosis, petalis obovatis discoloribus. *M. purpurea* Curt. bot. mag. t. 390.* *M. discolor*. Vent. malm. 24. t. 24.* (v. v.)

γ. *liliflora*, ramis floridis foliosis, petalis oblongis utrinque albis. *M. liliflora* Lam. dict. 3. p. 675. — Banks. ic. Kœmpf. t. 44.

13. *M. FUSCATA* (Andr. bot. rep. t. 229.) foliis perennantibus elliptico-oblongis, senioribus glabris, junioribus ramulisque fusco-tomentosis, floribus erectis. ♂ in Chinâ. Sims. bot. mag. t. 1008. DC. syst. 1. p. 458.* *M. fasciata*. Vent. malm. n. 24. adn. 2. (v. v.)

β. *annonæfolia* (Salisb! par. lond. n. 5. t. 5.*) foliis latioribus pollicem nempè latis, pedicello paulò brevioribus, flore magis rubente, antheris numerosioribus.

γ. *hebeclada*, floribus brevius pedunculatis, ramis magis velutino-tomentosis, foliisque brevioribus. — ex Indiâ orta.

14. *M. FUMILA* (Andr. bot. rep. t. 226.) foliis perennantibus glabris reticulato-venosis ellipticis utrinque acuminatis subglaucis, floribus cernuis. ♂ 5 in Amboinæ altis montibus. Vent! malm. 37. t. 37.* *Liriodendron lilifera* Lin! spec. 755. — DC. syst. 1. p. 458.* (v. s.)

15. *M?* *INODORA* (DC. syst. 1. p. 459.*) foliis ovato-acuminatis glabris, pedunculis terminalibus nifloris congestis. ♀ agrestis prope Cantonem Sinarum. *Liriodendron lilifera* Lour. fl. coch. ed. Willd. 1. p. 424.* non Lin. *Sam-paca montana* Rumph. amb. 2. p. 204. t. 69?

16. *M?* *COCO* (DC. syst. 1. p. 459.*) foliis ovatis integerrimis nitidis, floribus solitariis, sepalis 3 oblongis petalisque incurvis, ovariis 8. ♂ in Cochinchinâ. *Liriodendron Coco* Lour. fl. coch. ed. Willd. 1. p. 424.*

17. *M?* *FIGO* (DC. syst. 1. p. 460.*) foliis lanceolatis integerrimis nitidis reflexo-incurvis, floribus solitariis, ovariis 40-50. ♂ Culta Macai et Cantone Sinarum. *Liriodendron Figo* Lour. fl. coch. ed. Willd. 1. p. 424.* An potiùs Micheliæ species?

VIII. TALAUMA. Juss. gen. 281.

Carpella spicatum disposita, 1-2-sperma, inter se coalita in fructum strobiliformem extus valvatis irregulariter dehiscentem; semina in quoque loculo 2 aut abortu solitaria, filo adnexa. — *Magnolia Mexicana* forsàn ad *Talaumam* referenda. DC. syst. 1. p. 460.

1. *T. PLUMIERI* (Swart. prodr. 87. fl. ind. occ. 2. p. 997.*) ♂ in proclivibus tor-

rentum Martinicæ, Guadeloupæ, Sanctæ-Luciæ. *Magnolia fatiscens* Rich! ic. et descr. ined. folia coriacea glabra venoso-reticulata ovato-subrotundata basi subcuneata. (v. s.)

β. *longifolia*, foliis obovato-oblongis. — in Dominicâ.

IX. LIRIODENDRON. *Lin. gen. n. 689. Gaertn. fruct. 2. p. 475. t. 178.*

Carpella spicatis disposita, 1-2-sperma, indehiscentia, decidua, in alam apice producta. — Character è solo *L. Tulipifera* desumptus; species Asiaticæ in Magnoliarum sectione secundâ inquirendæ.

1. *L. TULIPIFERA* (Lin! spec. 755.) ♂ in Americæ borealis locis pinguibus: Michx. f. arb. amer. 3. p. 202. t. 5.* DC. syst. 1. p. 461. (v. s. et v.)

Ordo IV. ANONACEÆ. *Juss. gen.* 283. *DC. syst.* 1. p. 463.
Dunal. monog. 1817.

Sepala 3-4 breviter persistentia rarissimè libera sæpiùs in calycem 3-4-fidum 3-4-partitumve coalita. Petala 6, hypogyna, duplici serie disposita, inter se alternantia, sæpiùs coriacea et calycem interiorem subæmulantia, æstivatione imbricatâ, serie quaque æstivatione valvatâ, serie internâ exteriori nunc majore, nunc minore, rariùs nullâ; stamina numerosa adpressa discum sæpè hemisphæricum obtegentia; filamenta brevissima; antheræ subsessiles, connectivo apice glanduloso subtetragono interdum nectarifero, extrorsum et deorsum loculos antherarum longitudinaliter dehiscentes gerentes. Carpella sæpiùs plurima confertissima monostyla, nunc aggregata aut coalita, nunc rarissimè (abortu?) solitaria, stylis brevibus terminata, demum baccata aut capsularia, 1-∞-sperma, sessilia aut stipitata, interdum coadunata; semina ovata aut ovato-oblonga, 1-2-serialia, angulo carpellorum interiori adfixa; testa membranaceo-crustacea fragilis; endoplevra membranacea introrsum producta seu plicata, processibus plurimis lamellosis subulatis intrâ albuminis rimas foraminave immersis; albumen carnosum durum semini conforme, sulco depresso sæpissimè marginatum, rimis transversalibus aut foraminibus contiguis exsculptum; embryo minutus in regione albuminis umbilicati locatus; cotyledones breves; radícula tertiola. — Arbores aut frutices, è regionibus tropicis utriusque orbis ortæ. Turiones sæpè pubentes. Folia alterna, breviter petiolata, penninervia, integra.

§. 1. *Anonaceæ carpellis unilocularibus in fructum unicum multilocularem coalitis.*

I. KADSURA. *Juss. ann. mus.* 16. p. 340. *Dunal. monog. Anon.* p. 57.

Calyx 3-partitus; petala 6 ovata; stam. ∞ breviter; ovaria 30-40 superiorum carnosum ovato-globosum demum ampliatum aggregata; carpella totidem carnosa agglomerata sessilia 1-locularia 2-sperma. (Kœmpf. Th.) Semina ignota et ideò genus dubitanter ad Anonaceas relatum. *DC. syst.* 1. p. 465.

1. K. JAPONICA (Lin. spec. 756.) §. in Japoniâ. Kœmpf. amœn. 476 et 185. t. 477. * hist. Jap. 458. ic. — frutex parvus, procumbens, verrucosus; folia ovalia vel oblongo-ovalia, crassa, serrata; pedunculi 1-flori solitarii.

II. ANONA. *Adans. fam.* p. 365. *Dun. monog. Adon.* p. 58.

Sepala 3 basi coalita concava subcordata acutiuscula. Petala 6 crassiuscula, interiora minora aut nulla; antheræ ∞ subsessiles apice angulatæ dilatatæ torum obtegentes. Carpella ∞ coalita in baccam

unicam sessilem cortice muricato squamoso aut reticulato, intus pulposam, ad ambitum oo-locularem, loculis 1-spermis. DC. syst. 1. p. 466.

§. 1. *Petalis concavis crassis cordatis ovatisque.*

*. *Petalis exterioribus acutis, interioribus obtusis paulò minoribus.*

1. A. MURICATA (Lin! spec. 756.) foliis ovato-lanceolatis glabris subnitidis, pedunculis solitariis unifloris, petalis interioribus cordatis acuminatis, interioribus obtusis, fructibus muricatis, mucronibus carnosis. §. in insulis Caribæis, et Amer. calid. Jacq. obs. 1. p. 10. t. 5.* excl. Plum. syn. Dunal! monog. anon. p. 62.
 β? fructibus subsphæricis rectis. Dunal. monog. anon. p. 62. — In var. α fructus subconicus incurvus extus virens, in var. β. (forsan specificè distinguendâ) subsphæricus extus aureus. DC. syst. 1. p. 467.* (v. s.)
2. A. PURPUREA (Moq. et Sessè. fl. mex. ined. Dun. mon. anon. p. 64. t. 2.*) foliis subsessilibus lanceolatis subtus subferrugineis, floribus axillaribus subsessilibus, petalis exterioribus cordatis acutis, interioribus subrotundis. §. in Mexico. DC. syst. 1. p. 468.*
3. A. HUMBOLDTII (Dunal! monog. anon. p. 64. t. 3.*) foliis oblongo-acuminatis glabris leviter punctatis, pedunculis brevibus solitariis axillaribus unifloris, petalis exterioribus ovato-subcordatis acutis, interioribus obtusiusculis. §. in Cumanæ provinciâ. — DC. syst. 1. p. 468.* A. Humboldtiana Kunth. nov. gen. 5. p. 56. (v. s. in H. et B.)
4. A. LAURIFOLIA (Dunal. mon. anon. p. 65.) foliis ovato-lanceolatis glabris, pedunculis solitariis unifloris pendulis, petalis exterioribus cordatis acutis, interioribus rotundatis, fructibus mammæformibus lævibus. §. in insulis Caribæis. Cat. car. 2. p. 67. t. 67.* folia ovato-lanceolata Lauri nobilis foliis similia. DC. syst. 1. p. 468.

**. *Petalis exterioribus obtusis.*

5. A. OBTUSIFOLIA (Tussac fl. antil. t. 28.*) foliis oblongo-lanceolatis undulatis acuminatis nervosisissimis, junioribus tomentosis, adultis glabris, pedunculis axillaribus unifloris, petalis exterioribus obtusis. §. culta in Santo-Domingo, forsæn ex Asiâ orta? DC. syst. 1. p. 469.

***. *Petalis omnibus acutis, interioribus paulò minoribus.*

6. A. PALUSTRIS (Lin. spec. 757.) foliis ovato-oblongis coriaceis glaberrimis, floribus solitariis pedunculatis, petalis omnibus acutis, fructibus subareolatis. §. in Jamaicæ depressis paludosis. — Pluk. alm. 32. t. 240. f. 6. — Dunal! mon. anon. p. 65. DC. syst. 1. p. 469. (v. s. in h. Desf.)
7. A. LONGIFOLIA (Aubl. Guian. 1. p. 615. t. 248.*) foliis oblongis apice acuminatis mucronatis glabris, floribus axillaribus pedunculatis, petalis omnibus acutis, fructibus ovato-globosis punctatis reticulatis. §. in Guianâ. Dunal! mon. anon. p. 66.* DC. syst. 1. p. 469. (v. s. sine fl.)
8. A. PUNCTATA (Aubl. Guian. 1. p. 614. t. 247.*) foliis ovato-oblongis acutis glabris, floribus axillaribus subsessilibus, petalis omnibus acutis, fructibus subglobosis leviter punctatis. §. in Cayennæ sylvis. Dunal! mon. anon. p. 67. DC. syst. 1. p. 470.*
9. A. PERUVIANA (H. et B! ined. Dunal! mon. anon. p. 67.*) foliis oblongo-ellipticis acutis subcoriaceis subdecurrentibus, pedunculis axillaribus bracteolatis, petalis omnibus acutis, fructibus globosis reticulatis. §. in Peruvîæ uliginosis. DC. syst. 1. p. 470.* A. uliginosa Kunth. nov. gen. 5. p. 56. (v. s. in h. H. et B.)
10. A. AMBOTAY (Aubl. Guian. 1. p. 616. t. 249.*) foliis oblongo-ellipticis acutis subtus ferrugineo-tomentosis, floribus axillaribus solitariis brevissimè pedunculatis, petalis acutis. §. in Cayennæ sylvis. Dunal! mon. anon. p. 67. fructus ignoti. DC. syst. 1. p. 470.

11. *A. PALUDOSA* (Aubl. Guian. 1. p. 611. t. 246.*) foliis oblongis acutis supernè subtomentosis subtùs tomentoso-sericeis rufescentibus nervosis, floribus breviter pedunculatis, petalis omnibus acutis, fructibus ovatis tuberculatis. §. in Guianæ pratis paludosis. Dunal! monog. anon. p. 68.* DC. syst. 1. p. 471.* (v. s.)

§. 2. *Petalis exterioribus ovatis concavis acutis coriaceis, interioribus nullis. — Fructus non satis noti, et ideo sectio ad Anonæ genus ex habitu tantùm relata.*

12. *A. ECHINATA* (Dunal! monog. anon. p. 68. t. 4.*) foliis ovato-lanceolatis acutiusculis supernè glaberrimis subtùs tomentosis, ramis rugosis, pedunculis solitariis unifloris, floribus 3-petalis, fructibus ovoideis echinatis. §. in Cayennâ. DC. syst. 1. p. 471.* (v. s.)

13. *A. SERICEA* (Dunal! monog. anon. p. 69. t. 5.*) foliis ovato-oblongis acuminatis supernè glabris subtùs ramulisque ferrugineo-tomentosis sericeis, floribus 3-petalis solitariis axillaribus pedunculatis extùs ferrugineis. §. in Cayennâ. flores ferè *A. echinatæ*. DC. syst. 1. p. 471.* (v. s.)

§. 3. *Petalis exterioribus lineari-oblongis angustis apice triquetris basi tantum concavis sæpè clausis genitalia occultantibus, interioribus minutissimis.*

14. *A. SQUAMOSA* (Lin! spec. 757.) foliis lanceolatis glabris pellucido-punctatis, petalis exterioribus subclausis, fructibus ovoideis squamosis. §. culta in Indiâ, verosimiliter ex Americâ orta. Jacq. obs. 1. p. 13. t. 6. f. 1.* Dunal! mon. anon. p. 69.* DC. syst. 1. p. 472.* (v. s.)

15. *A. FORSKAHLII* (DC. syst. 1. p. 472.*) foliis oblongis subellipticis glabris punctatis subtùs glaucis, petalis exterioribus oblongis subclausis. §. in Egypto. *A. glabra* Forsk. Egypt. desc. 102.* ic. t. 15. *A. Asiatica* var. B. Dunal! monog. anon. p. 71. (v. s.)

16. *A. CINEREA* (Dunal! monog. anon. p. 71. t. 8.*) foliis oblongo-ellipticis sublanceolatis pellucido-punctatis subtùs pubescentibus, petalis exterioribus subclausis, fructibus ovato-globosis squamosis. §. culta? in insulâ S. Thomas et propè Cumanam? DC. syst. 1. p. 473.* (v. s.)

17. *A. CHERIMOLIA* (Mill. dict. n. 5.*) foliis ovato-lanceolatis impunctatis subtùs tenuissimè tomentoso-sericeis, petalis exterioribus subclausis extùs tomentosis, fructibus subglobosis subsquamosis. §. in Peruvîâ. et Novâ Granatâ. *A. tripetala* Ait. h. kew. 2. p. 252. Sims. bot. mag. t. 2011.—DC. syst. 1. p. 473. (v. v.)

18. *A. RETICULATA* (Lin. spec. 757. excl. Rumph. syn.) foliis oblongo-lanceolatis acutis glabris subpunctatis, petalis exterioribus oblongis subclausis, fructibus ovato-globosis reticulato-areolatis. §. in ins. Caribæis. DC. syst. 1. p. 474.* Dunal! monogr. 72.*

α. Areolis squamoso-rotundatis Sloan. hist. 2. p. 167. t. 226.

β. Areolis angularibus subpentagonis. Dunal. l. c. *A. reticulata* Jacq. obs. 1. p. 14. t. 6. f. 2.*

19. *A. MUCOSA* (Jacq. obs. 1. p. 16.*) foliis oblongo-lanceolatis glabris, petalis exterioribus apicè patulis, fructibus areolatis, areolis gibbis. §. in Martiniçæ sylvis. Dunal. mon. anon. p. 74. DC. syst. 1. p. 474.—Rumph. amb. 1. p. 136. t. 45.*

§. 4. *Petalis exterioribus oblongo-ellipticis obtusis, interioribus minoribus lanceolatis obtusiusculis, calyce coriaceo trifido magno subcampanulato, fructibus conoideis lævibus?*

20. *A. GLABRA* (Lin. spec. 758.) foliis ovato-lanceolatis glabris, pedunculis oppositifoliis bifloris, fructibus conoideis obtusis lævibus. §. in Carolinâ.

Catesb. car. 2. p. 64. t. 64. — Dunal. mon. anon. 1. p. 74. DC. syst. 1. p. 475.*
(v. s. in h. Mus. Par.)

21. A. GRANDIFLORA (Lam! dict. 2. p. 126.*) foliis ovato lanceolatis glabris coriaceis suprà nitidis, pedunculis axillaribus solitariis, fructibus ovoideis glabris subpunctatis. 5. in insulis Mauritanis. Dunal! mon. anon. p. 75. t. 6 et 6. a.* DC. syst. 1. p. 475.* (v. s.)

ℓ. *Madagascariensis*. Pers. ench. 2. p. 95. in Madagascar.

22. A. AMPLEXICAULIS (Lam! dict. 2. p. 127.*) foliis cordato oblongis amplexicaulis acutis glabris, pedunculis axillaribus solitariis unifloris. 5. in insulis Madagascar et Mauriti. Dunal! mon. anon. p. 76. t. 7. DC. syst. 1. p. 475. (v. s. in h. Juss. et Mus. Par.)

§. 5. *Petalis æqualibus patentibus.*

23. A? AXILLIFLORA, foliis ovali-oblongis subobtusis superne glabris, subtus minutissimè pubescentibus, pedicellis axillaribus 1-floris petiolo quadruplò longioribus. 5. in Guianâ Gallicâ. fructus ignotus. (v. s. comm. à Mus. Par.)

24. A? UNIFLORA (Dunal. mon. anon. p. 76.) foliis oblongis acuminatis glabris subtus glaucis, pedunculis oppositifoliis tomentosis candicantibus. 5. in Parâ Braziliz. Deless! ic. sel. 1. t. 87. species pulcherrima sed genus dubium. DC. syst. 1. p. 476. n. 25.* (v. s. in h. Mus. Par.)

†. *Anonæ non satis notæ.*

25. A. ASIATICA (Lin! spec. 2. p. 758.) foliis oblongis acuminatis impunctatis, junioribus pubescentibus, adultis glabris. 5. in Zeylonâ. A. Asiatica. a Dunal. mon. anon. p. 71. DC. syst. 1. p. 476. (v. s. in b. Lin.) †.

26. A. SENEGALENSIS (Pers! ench. 2. p. 95.) foliis latis ovatis subcordatis coriaceis glabris subtus glaucis, petiolis subvelutinis, pedunculis geminis terminisve intrâ-foliaceis. 5. in Senegaliâ et in Guineâ. Dunal. mon. anon. p. 76. Deless! ic. sel. 1. t. 86. DC. syst. 1. p. 476.* (v. s. in h. Mus. Par.)

27. A? EXSUTICA (Dunal. monog. anon. p. 77.) foliis ovato-oblongis coriaceis superne incidis utrinquè ramisque glaberrimis, pedunculis suboppositifoliis simplicibus aut bipartitis. 5. in sylvis Guianæ. DC. syst. 1. p. 477. (v. s. in h. Lamb.)

28. A. DOLABRIPETALA (Raddi in att. soc. ital. 16. p. 15) foliis oblongo-lanceolatis infernè petiolisque hirsutis, petalis 3 connatis lateraliter compresso-dolabrisformibus tomentosis. 5. in Braziliz monte Cercovado propè Rio-Janeiro.

29. A. AFRICANA (Lin. sp. 758.) foliis lanceolatis pubescentibus. 5. in Americâ (Lin.) Deest in herb. Lin. omninò obscura.

30. A. ATABAPENSIS (H. B. et K. nov. sp. am. 5. p. 58.) foliis oblongis obtusis basi cuneatis coriaceis glaberrimis nitidulis epunctatis, pedunculis solitariis 1-floris, petalis ovatis cordatis acutiusculis, interioribus vix minoribus. 5. in novâ Guianâ ad ripam inundatam fluminis Atabapo. an A. palustris?

31. A. BONPLANDIANA (H. B. et K. l. c. 5. p. 58.) foliis oblongis acuminatis subcoriaceis punctulatis glabris nitidis, pedunculis solitariis 1-floris, petalis exterioribus ovatis tenuissimè tomentosis. 5. in littore Oceani Pacifici propè Guayaquil Quitensium.

32. A. EXCELSA (H. B. et K. l. c. 5. p. 59.) foliis oblongis acuminatis basi acutis subcoriaceis epunctatis glabris conduplicatis. 5. in Novâ Hispaniâ propè la Venta del Exido.

33. A. RIPARIA (H. B. et K. l. c. 5. p. 59.) foliis oblongis acuminatis basi acutis membranaceis punctulatis glabris nitidulis, fructibus pyriformibus obsolete reticulatis. 5. in Peruviâ ad ripam fluminis Guancabamba propè Colazei.

34. A. MANIKOTE (H. B. et K. l. c. 5. p. 59.) foliis obovato-ellipticis breviter acuminatis basi subrotundatis membranaceis punctulatis glabris nervo ve-

nisque pubescentibus, fructibus globosis muricatis. 5 incultis propè urbem Angosturæ proxima A. muricatæ videtur.

35. A. LÆVIS (H. B. et K. l. c. 5. p. 60.) foliis lanceolato-oblongis acuminatis basi acutis membranaceis glabris subtus pilosiusculis punctulatis, pedunculis aggregatis unifloris, petalis exterioribus lanceolatis acutiusculis, fructibus lævibus. 5. in cultis Guyanæ, prope Angosturam, Cumanam. An A. laurifolia aut A. glabra?
36. A. QUINDIENSIS (H. B. et K. l. c. 5. p. 60.) foliis lanceolato-oblongis utrinquè acuminatis subcoriaceis suprâ glabris nitidis subtus adpressopilosiusculis obsolete punctulatis, pedunculis 1-5-unifloris, petalis exterioribus ovato-lanceolatis, fructibus punctatis. 5 in Novo-Granatensium Andibus Quindiensibus alt. 1200 hex.

§. 2. *Anonaceæ fructu solitario uniloculari.*

III. MONODORA. Dunal. mon. anon. p. 79.

Calyx 3 sepalus; petala 6 biserialia, exteriora lanceolata, interiora ovata, basi coalita; antheræ ∞ subsessiles; ovarium 1 ovatum stigmate sessili coronatum; Bacca lævis subglobosa 1-locularis ∞ -sperma; semina in pulpâ nidulantia. DC. syst. 1. p. 477.

1. M. MYRISTICA (Dunal! mon. anon. p. 80.) baccis subglobosis magnis. 5. culta in Jamaica, verosimiliter ex Africâ æquatoriali allata. Brown. cong. p. 56. Anona Myristica Gaert. fruct. 2. p. 194. t. 125. f. 1. Lunan. h. jam. 10. — American nutmeg. Long. hist. jam. 3. p. 735. (v. s. in h. Lamb.)

§. 3. *Anonaceæ carpellis plurimis distinctis.*

IV. ASIMINA. Adans. fam. 2. p. 365. — Orchidocarpum. Michx. fl. am. bor. 1. p. 329.

Calyx 3-partitus. Petala 6 patentia ovato-oblonga, interiora minora; antheræ ∞ subsessiles; ovaria plurima plerumque 3, ovata seu oblonga; carpella totidem baccata sessilia; semina plurima, unicâ? seriè disposita. DC. syst. 1. p. 678.

1. A. PARVIFLORA (Dunal! monog. anon. p. 82. t. 9.*) foliis cuneato-obovatis mucronatis subtus ramulisque rufo-pubescentibus, floribus sessilibus, petalis exterioribus calyce vix duplò longioribus. 5. in Virginia et Carolinâ. Porcelia parviflora Pers. ench. 2. p. 95. — DC. syst. 1. p. 478.* (v. s.)
2. A. TRILORA (Dunal! anon. 83.) foliis oblongo-cuneatis acuminatis ramulisque glabriusculis, floribus breviter pedunculatis, petalis exterioribus calyce quadruplò longioribus subrotundo-ovatis. 3 5. in Pennsylvania et Floridâ. — Anona triloba. Lin. sp. 758. Michx. f. arb. amer. 3. p. 161. t. 9.* Schkuhr. handb. 2. p. 95. t. 149. — DC. syst. 1. p. 479.* (v. v. et s.)
3. A. PYGMÆA (Dunal! monog. an. p. 84. t. 10.*) caule suffruticoso, foliis oblongo-linearibus longè cuneatis obtusis ramulisque glabris, floribus breviter pedunculatis, petalis exterioribus calyce multotiès majoribus obovato-oblongis. 5 in agris arenosis Georgiæ et Floridæ. Anona pygmæa Bartr. itin. ed. germ. p. 21. t. 1. ex Willd. spec. 2. p. 1268.* DC. syst. 1. p. 479.* (v. s.)
4. A. GRANDIFLORA (Dunal! mon. anon. p. 84. t. 11.*) foliis cuneato-obovatis obtusis subtus ramulisque rufo-pubescentibus, floribus sessilibus, petalis exterioribus obovatis calyce multotiès amplioribus. 5. in sylvis arenosis Georgiæ et Floridæ. Porcelia grandiflora. Pers. ench. 2. p. 95. — DC. syst. 1. p. 480.* Herb. amat. t. 263. (v. s.)
5. A. CAMPECHIANA (H. B. et K. nov. sp. am. 5. p. 61.) foliis elliptico-oblongis acuminatis basi acutis membranaceis punctatis suprâ hirtellis subtus molliter ramulisque tomentoso-hirtis, pedunculis brevibus solitariis, petalis oblongis subæqualibus. 5 circa Campeche Mexicanorum.

V. PORCELIA. Ruiz. et Pav. syst. 1. p. 144. prod. 84. t. 16.*

Sepala 3 basi subcoalita. Petala 6, interiora paulò majora. Antheræ ∞ subsessiles. Carpella 3-6 sessilia coriacea vix subcarnosa cylindrica subtorulosa ∞-sperma, semina gemino ordine superimposita. DC. syst. 1. p. 480.

1. P. NITIDIFOLIA (Ruiz. et Pav! syst. 1. p. 144.) 5. in Peruvix nemoribus montanis. Dunal! mon. anon. p. 85.* DC. syst. 1. p. 480*. Pedunculi axillares 1-aut pauciflori, plures aggregati. (v. s.)

VI. UVARIA. Lin. gen. 692. Gaertn fruct. 2. p. 155 et 157.

Sepala 3 basi coalita ovato-cordata. Petala 6 ovata, 3 exteriora minora? Antheræ ∞ receptaculum hemisphæricum obtegentes. Ovaria ∞ sæpè villosa. Carpella ∞ baccata carnosa, ovato-globosa ∞-locularia ∞-sperma. Semine duplici serie disposita. DC. syst. 1. p. 481.

1. U. ZÉYLANICA (Lin. spec. 2. p. 756. exclus. Rheed. et Rumph. syn.) sarmentosa, foliis ovato-lanceolatis glabris, baccis pluribus in stipitem attenuatis ovato-cylindricis, seminum integumentum processibus internis lamellosis. 5 in Ceylonâ. Lam. ill. t. 495. f. 2. Dunal. mon. anon. p. 88. DC. syst. 1. p. 481. (v. s. in h. Juss.)?
2. U. CHAMÆ (Beauv. fl. ow. et ben. 2. p. 43. t. 83. f. 2.) foliis alternis integris parvis ovatis nitentibus acutis, floribus racemosis, sepalis coriaceis, petalis ligulatis. 5 ad ripas flum. Sau. Jago. †.
3. U. GAERTNERI (DC. syst. 1. p. 482.*) baccis ovatis in stipitem attenuatis, seminum integumentum processibus internis subulatis et ferè acicularibus. 5. Verosimiliter in Ceylonâ. U. trifoliata Gaert. fruct. 2. p. 167. t. 114. f. 2. Lam. ill. t. 495. f. 3. Dunal. mon. anon. p. 89. DC. syst. 1. p. 482.* †.
4. U. LUTEA (Roxb! fl. corom. 1. p. 32. t. 36.) arborea, foliis oblongis acuminatis glabris nitidis, pedunculis solitariis 1-6-floris, baccis oblongis 6-spermis. 5. in montibus Coromandelix. Dunal. mon. anon. p. 89. DC. syst. 1. p. 482.* (v. s. in h. Lamb.)
5. U. TOMENTOSA (Roxb. fl. corom. 1. p. 31. t. 35.*) arborea, foliis oblongis acutis tomentosis, pedunculis 1-floris subsolitariis, baccis globosis 4-spermis. 5. in Coromandelix montibus. Dunal. monog. anon. p. 90. DC. syst. 1. p. 483.*
6. U. DULCIS (Dunal! monog. anon. p. 90. t. 13.*) foliis oblongo-ellipticis basi attenuatis cordatis subtus ramulisque velutinis, pedunculis geminis axillaribus aut oppositifoliis, medio bracteatis articulatisve. 5. in Javâ. DC. syst. 1. p. 483.* (v. s. in h. Deless.)
7. U. JAVANA (Dunal! monog. anon. p. 91. t. 14.*) foliis oblongo-ellipticis basi cordatis subtus ramulisque tomentoso-ferrugineis, pedunculis axillaribus aut oppositifoliis paucifloris, pedicellis subumbellatis medio bracteatis 5. in Javâ. Pistilla villosa. DC. syst. 1. p. 483.* (v. s. in h. Deless.)
8. U. VELUTINA (Dunal. monog. anon. p. 91.) foliis subsessilibus ovatis basi cordatis acuminatis ramisque velutinis, pedicellis corymbosis unifloris. 5 in Indiâ orient. Ovaria confertissima subvelutina. DC. syst. 1. p. 484. (v. s. in h. Lamb.)
9. U. ? SPECTABILIS (DC. syst. 1. p. 484.*) foliis oblongis acuminatis subglabris, junioribus ramisque ferrugineo-velutinis, pedunculis 1-floris lateralibus aut oppositifoliis, petalis obovatis, interioribus apice bifidis. 5. in Guianâ. Ovaria confertissima vix distincta; fructus ignotus; an genus proprium? (v. s. in h. Lamb.)

VII. UNONA. Lin. suppl. 270.

Sepala 3 rarissimè 4 basi coalita ovata acutiuscula. Petala 6 biserialia,

3. interiora minora. Stamina ∞ . Carpella ∞ sicca? indehiscentia stipitata ovata aut oblonga, 1- ∞ -locularia, lævia aut torulosa, ∞ -sperma. Semina simplici serie disposita. DC. syst. 1. p. 485.

Sect. I. UNONARIA. DC. syst. 1. p. 486.

Flores patentēs; carpella sublævia aut læviter torulosa.

§. 1. *Petalis ovatis oblongisve subæqualibus*. MARENTERIA.

1. U. NARUM (Dunal. mon. anon. p. 99.) caule sarmentoso, foliis lanceolatis acuminatis, pedunculis lateralibus solitariis unifloris, petalis subrotundo-ovatis subæqualibus inflexis, carpellis longè stipitatis sublævibus. $\frac{5}{2}$ in Malabariâ et forsân in Moluccis. Rheed. mal. 2. p. 11. t. 9.* DC. syst. 1. p. 486.
2. U. MUSARIA (Dunal. mon. anon. p. 100.) caule sarmentoso, foliis elliptico-lanceolatis basi cordatis acuminatis, pedunculis 1-floris solitariis axillaribus, petalis subrotundo-ovatis subæqualibus reflexis, carpellis stipitatis subtorulosis. $\frac{5}{2}$ in Amboinæ, Baleyæ et Soloræ campis. Rumph. amb. 5. p. 78. t. 42.* DC. syst. 1. p. 486.*
3. U. PENDULIFLORA (Moc. et Sessé. in Dunal. mon. anon. p. 100. t. 28.) caule arborescente, foliis subsessilibus subcordatis oblongo-lanceolatis undulatis, pedunculis longis unifloris pendulis, petalis inflexis, carpellis longis substipitatis. $\frac{5}{2}$ in Mexico. DC. syst. 1. p. 487.*
4. U. MARENTERIA (DC. syst. 1. p. 487.*) ramulis scandentibus, foliis ovalibus obtusis glaberrimis, pedunculis erectis unifloris subterminalibus, petalis ovalibus extus velutinis subinæqualibus, carpellis substipitatis ventricosis. $\frac{5}{2}$ in Madagascar. Marenteria Nor. ex Petit-th! gen. nov. Madag. p. 18. n. 60. (v. s.)
5. U. CRASSIPETALA (Dunal! monog. anon. p. 101. t. 24.*) foliis oblongo-ellipticis acuminatis glabris, pedunculis axillaribus unifloris pilosis erectis petiolo longioribus, petalis oblongis crassis. $\frac{5}{2}$ in Cayennâ. fructus ignotus. DC. syst. 1. p. 487.* (v. s.)
6. U. FUSCATA (DC. syst. 1. p. 488.*) foliis glabris oblongis acuminatis, pedunculis axillaribus unifloris pilosis erectis, petiolo sub brevioribus, petalis ovali-oblongis acutiusculis. $\frac{5}{2}$ in Guianâ. Ovaria densè approximata, staminum longitudine æqualia; fructus ignotus. (v. s. in h. Lamb.)
7. U. OBTUSIFLORA (DC. syst. 1. p. 488.*) foliis glabris ovali-oblongis utrinquè acuminatis, pedunculis axillaribus unifloris adpressè vix pubescentibus petiolo longioribus, petalis oblongo-obovatis obtusissimis. $\frac{5}{2}$ in Guianâ. Genitalia ut in U. fuscata cui affinis. (v. s. in h. Lamb.)
8. U. ACUMINATA (DC. syst. 1. p. 488.*) foliis glabris oblongis utrinquè acuminatis, pedunculis axillaribus unifloris vix pubescentibus petiolo multò longioribus, petalis oblongis acutis. $\frac{5}{2}$ in Guianâ. Stamina et ovaria ut in U. fuscata. (v. s. in h. Lamb.)
9. U. MACROCARPA (Vahl! ined. DC. syst. 1. p. 489.) foliis ovalibus utrinquè obtusis glaberrimis coriaceis, pedunculis lateralibus brevibus ramosis 2-3-floris, petalis ovali-oblongis acutiusculis. $\frac{5}{2}$ in Guinæâ. Petala 6, ovali-oblonga, subcoriacea, inter se æqualia, pube tenuissimâ subvelutina. (v. s. in h. Juss.)
10. U. OVATA (Vahl! ined. DC. syst. 1. p. 489.*) foliis ovatis acutis supernè glabris subtus in nervo medio ramulisque ferrugineo-tomentosis, pedunculis bifloris brevissimis oppositifoliis, petalis ovali-oblongis acutiusculis. $\frac{5}{2}$ in Guinæâ. Petala subæqualia acutiuscula purpurea subcoriacea. (v. s. in h. Juss.)
ß. Afzeliana. in Sierrâ-Leonâ. Priori ferè similis, sed folia adulta omninò (excepto nervo longitudinali) glabra fusco-ferruginea. (v. s. in h. Lamb.)

11. *U. GRANDIFLORA* (Lechenault! in litt.) foliis ovali-oblongis subsessilibus basi cordatis supernè lævibus subtùs velutinis, pedunculis 1-floris oppositifoliis, petalis ovali-oblongis $\frac{5}{4}$ in Bengalo. Carpella 30-35 stipitata teretiuscula densè breviterque velutina indehiscencia, intùs transversè loculata, loculis 1-seriatis. (v. s.)

§. 2. *Petalis exterioribus ovato-oblongis acutiusculis, interioribus nullis seu minutissimis.* OETANIA.

12. *U. TRIPETALA* (DC. syst. 1. p. 490.) foliis lanceolatis supernè rugosis subtùs tomentosis, carpellis stipitatis ovatis subtriquetris 3-spermis punctato-granulatis. $\frac{5}{4}$ in planitiibus et collibus Amboinæ. — *U. tripetaloides* Dunal. mon. anon. p. 104. — Rumph. amb. p. 197. t. 66. f. 1. — *Uvaria tripetala* Lam.

§. 3. *Petalis lineari-lanceolatis longis angustis.* CANANGA.

13. *U. VIOLACEA* (Dunal. mon. anon. p. 105. t. 25.*) foliis ellipticis obtusiusculis glabris? pedunculis oppositifoliis cernuis unifloris bractea unica amplexis, petalis lanceolato-oblongis. $\frac{5}{4}$ in Mexico. *U. spec. nov.* Moç. et Sessé. pl. mex. ic. pict. ined. DC. syst. 1. p. 490.*

14. *U. UNCINATA* (Lam! dict. 2. p. 127.*) foliis oblongo-lanceolatis acuminatis glabris nitidis, pedunculis oppositifoliis infrà medium uncinatis, carpellis ovato-rotundatis. $\frac{5}{4}$ in Insula Mauritiï et insulis Indiæ orient. Dunal. mon. anon. p. 105. t. 12. et 12. a.* *Artabotrys odoratissimus* R. Brown. ex Ker. bot. reg. 5. t. 423. — DC. syst. 1. p. 490. (v. s. in h. Juss. et Deless.)

℞. pedunculis simplicibus arcuatis. — An huc referenda t. 16. Braam. ic. Chin. 1821?

15. *U. HAMATA* (Dunal! mon. anon. p. 106. t. 27.*) ramis subscandentibus, foliis oblongis acuminatis glabris nitidis, pedunculis solitariis uncinatis, petalis lanceolatis basi constrictis, carpellis ovatis utrinquè acuminatis. $\frac{5}{4}$ circa Cantonem Sinarum. DC. syst. 1. p. 491. (v. s. in h. Deless. et Lamb.)

16. *U. ESCULENTA* (Dunal. mon. anon. p. 107.) caule scandente, foliis ovato-lanceolatis glabris, ramis apice cirrhosis, pedunculis unifloris, petalis lanceolatis. $\frac{5}{4}$ in Indiâ circa Madras. *Uvaria esculenta* Roxb. Rottl. et Willd. in nov. act. nat. cur. berol. 4. p. 201. — DC. syst. 1. p. 491.*

17. *U. LESSERTIANA* (Dunal. mon. anon. 107. t. 26.*) foliis oblongo-ellipticis, supernè glabris nitidis, pedunculis lateralibus unifloris bracteatis, petalis oblongis subseriatis. $\frac{5}{4}$ verosimiliter in Chinâ aut Indiâ. *Uvaria uncata* Vahl! in h. Deless. — DC. syst. 1. p. 492.* (v. s. in h. Deless.)

18. *U. ODORATA* (Dunal! mon. anon. p. 108.*) foliis oblongo-lanceolatis acuminatis glabris, pedunculis lateralibus 1-4-floris, petalis lineari-lanceolatis, carpellis stipitatis ovato-teretibus apice umbilicatis. $\frac{5}{4}$ in Javâ et Chinâ. *Uvaria undulata* Lam! dict. 1. p. 595.* illust. t. 495. f. 1. — DC. syst. 1. p. 492.* (v. s.)

19. *U. LONGIFOLIA* (Lam! dict. 1. p. 597.*) foliis oblongo-linearibus acuminatis margine undulatis glabris, pedunculis lateralibus confertis subumbellatis, petalis lineari-lanceolatis acutis, carpellis stipitatis ovoideis. $\frac{5}{4}$ in Bengalo, Javâ, etc. DC. syst. 1. p. 492.* Sonn! voy. ind. 4. t. 131. (v. s.)

20. *U. NITIDISSIMA* (Dunal! mon. anon. p. 109. t. 23.*) foliis oblongo-ellipticis utrinquè acuminatis glabris nitidis, pedunculis axillaribus brevibus unifloris, petalis linearibus obtusiusculis. $\frac{5}{4}$ in Novâ-Caledoniâ. *Uvaria lucida* Vent! in h. Deless. fructus ignoti. — DC. syst. 1. p. 493. (v. s.)

21. *U. LIGULARIS* (Dunal. mon. p. 110.) foliis ovato-lanceolatis acutis, pedunculis lateralibus multifloris, petalis longis linearibus, carpellis stipitatis subglobosis. $\frac{5}{4}$ in Amboinâ. *Uvaria ligularis* Lam. dict. 1. p. 597. — Rumph. amb. 2. p. 298. t. 66. f. 2.* — DC. syst. 1. p. 493.*

Sect. II. DESMOS. DC. syst. 1. p. 493.

Petala lanceolata, oblonga vel linearia, interdum subclausa; carpella baccata torulosa subarticulata multilocularia? plus minùve moliniformia.

22. U. DISCRETA (Lin. f. suppl. 270.*) foliis oblongo lanceolatis angustis subtus sericeis, carpellis moniliformibus longè stipitatis. 5 in Surinam. Dunal. mon. anon. p. 110. U. monilifera Gaertn. fruct. 2. p. 156. t. 114? Lam. ill. t. 495. f. 4? — DC. syst. 1. p. 494.*
23. U? UNDULATA (Dunal. mon. anon. p. 111.) foliis ovato-oblongis glabris, pedunculis axillaribus unifloris, petalis oblongo-linearibus longis, exterioribus undulatis, carpellis oblongis submoniliformibus. 5 in regno Owarensi. Xylopi undulata Pal. Beauv! fl. owar. 1. p. 27. t. 16.* An Monodora congener ex R. Brown. qui suspicatur Belvisium ad hunc florem retulisse fructum Unonæ Æthiopice. — DC. syst. 1. p. 494.* (v. s.)
24. U. DISCOLOR (Vahl. symb. 2. p. 63. t. 36.*) foliis ovato-oblongis acuminatis glabris subtus glaucis, pedunculis lateralibus unifloris elongatis, carpellis moniliformibus breviter stipitatis. 5. in Indiâ orient. Dunal! mon. anon. p. 111.* DC. syst. 1. p. 494.* (v. s. sine flor. in h. Deless.)
25. U. CHINENSIS (DC. syst. 1. p. 495.*) foliis ovato-lanceolatis glabris, floribus solitariis patentibus longè pedunculatis, carpellis toruloso-moniliformibus subsessilibus. 5 in Chinâ propè Cantonem. Desmos Chinensis. Lour. fl. coch. ed. Willd. 1. p. 431.* Species non satis nota sed verosimiliter distincta ab U. discolor quâcum conjunxit cl. Willd. †.
26. U. COCHINCHINENSIS (DC. syst. 1. p. 495.*) foliis lanceolatis tomentosis, floribus solitariis clausis longè pedunculatis, carpellis subsessilibus toruloso-moniliformibus. 5. in dnetis Cochinchinæ. Desmos Cochinchinensis Lour. fl. coch. ed. Willd. 1. p. 431.* Unona Desmos Dunal. mon. anon. p. 112.
27. U. AROMATICA (Dun. mon. anon. p. 112.) foliis oblongis acuminatis glabris, pedunculis axillaribus 1-2-floris, carpellis oblongo-teretibus torulosis subsessilibus. 5 in sylvis Guianæ. Uvaria Zeylanica Aubl. Guian. 2. p. 605. t. 243.* excl. syn. — DC. syst. 1. p. 495.*
28. U. ÆTHIOPICA (Dunal! mon. anon. p. 113.*) foliis ovato-lanceolatis acutis glabris subtus glaucis, carpellis teretibus torulosis subsessilibus. 5. in Æthiopiâ et in Sierrâ-Leonâ. Piper Æthiopicum Math. comm. 1. p. 434. ic. DC. syst. 1. p. 496.* (v. s.)
29. U. OXYPETALA (DC. syst. 1. p. 496.*) foliis ovali-oblongis acuminatis supernè lævibus subtus subglaucis, pedunculis axillaribus brevissimis 1-floris, petalis linearibus acuminatis. 5. in Sierrâ-Leonâ. Carpella ignota. An ad sequentem sectionem amendanda? (v. s. in h. Lamb.)
30. U? LEPTOPETALA (DC. syst. 1. p. 496.*) foliis ovali-oblongis acuminatis ramisque glabris, pedunculis axillaribus ramosis, petalis oblongo-linearibus acuminatis. 5 in insulâ Timor. Deless! ic. sel. 1. t. 88. Petala æqualia calyce sextuplò longiora. (v. s. in h. Mus. Par.)

Sect. III. MELODORUM. DC. syst. 1. p. 497.

Flores pyramidati angusti elongati, petala lineari-triangularia acuta, sæpè clausa, basi genitalia tegentia; carpella baccata sublævia vel leviter torulosa.

31. U. LATIFOLIA (Dunal. mon. anon. p. 115.) foliis latè oblongo-lanceolatis subtus laevibus, floribus racemosis, baccis numerosis 2-3-spermis. 5 in Moluccis. Rumph. amb. 2. p. 198.* Semina odorata DC. syst. 1. p. 497.* †.
32. U. SYLVATICA (Dunal. mon. anon. p. 115.) foliis ovato-oblongis subtus tomentosis, floribus solitariis breviter pedicellatis carnosissimis, baccis subsessilibus ovato-oblongis asperis. 5. in sylvis Cochinchinæ. Melodorum arborescens. Lour. fl. coch. ed. Willd. 1. p. 430.* — DC. syst. 1. p. 497.*

33. *U. DUMETORUM* (Dunal. mon. anon. p. 116.) foliis lanceolatis glabris, floribus solitariis, baccis subsessilibus ovato-oblongis asperis. ♂ in dumetis Cochinchinæ. Melodorum fructicorum Lour. fl. coch. ed. Willd. 1. p. 430.* Pulpa gratissima. DC. syst. 1. p. 497.*
34. *U. LUCIDA* (DC. syst. 1. p. 498.*) foliis ovali-oblongis utrinque acuminatis glaberrimis supernè nitidis, pedunculis simplicibus, baccis oblongis obtusis subcompressis subtorulosis breviter stipitatis. ♀ in Peruvâ. Deless! ic. sel. 1. t. 89. (v. s. in h. Juss.)
35. *U. ACUTIFLORA* (Dunal! mon. anon. p. 116. t. 22.*) foliis ovato-lanceolatis acutis rigidiusculis glabris, pedunculis brevissimis unifloris, baccis ovato-oblongis breviter stipitatis. ♀ in Caribæis insulis aut in Sierrâ-Leonâ. Petala extus sericea. (v. s.)
36. *U. XYLOPIOIDES* (Dunal! mon. anon. p. 117. t. 21.*) foliis oblongis acuminatis subtus sericè nitidis, margine ad basim præcipuè revolutis, pedunculis 2-4 axillaribus brevibus. ♀ in Americâ merid. regno Novo-Granatensi. H. B. et Kunth. nov. gen. 5. p. 62. Baccæ siccæ oblongæ subcurvatae multiloculares. DC! syst. 1. p. 498.* (v. s. in h. H. et Bonpl.)
37. *U?* *POLYCARPA* (DC. syst. 1. p. 499.*) foliis oblongis acuminatis subglaucescibus, nervo medio basi subtomentoso, baccis longè stipitatis. ♀ in Sierrâ-Leonâ. An potiùs Guatterizæ species? An genus proprium? (v. s. sine fl. in h. Lamb.)
38. *U?* *SELANICA*, foliis ovatis acutis supernè glabris subtus rugosis, racemis paucifloris terminalibus pendulis. ♀ in Moluccis. Dammara Selandica Rumph. amb. 2. p. 168. t. 56. Lam. dict. 2. p. 259.

VIII. XYLOPIA. Lin. gen. 1027. Lam. ill. t. 495.

Calyx 3-5 lobus, laciniis ovatis coriaceis acutiusculis. Petala 6, exteriora majora. Stam. ∞ receptaculo sæpiùs globoso inserta. Carpella 2-15 breviter stipitata compressa 1-locularia 1-2-sperma nunc dehiscentia, nunc subbaccata. Semina obovata nitida, interdum arillata. DC. syst. 1. p. 499.

1. *X. MURICATA* (Lin! spec. 1367.) foliis lanceolatis acuminatis subtus strigosis apicè barbatis, pedunculis multifloris, carpellis muricatis. ♂ in montibus Jamaicæ. *X. frutescens* Gaertn. fruct. 1. p. 339. t. 69. f. 7. — DC. syst. 1. p. 499.* (v. s. in h. Lin.)
2. *X. FRUTESCENS* (Aubl. Guian. 1. p. 602. t. 292.* excl. syn.) foliis oblongo-lanceolatis acuminatis subtus sericeis, pedunculis 1-3 brevissimis, carpellis lævibus. ♀ in Brasiliâ et Guianâ. Lam. ill. t. 495. Dunal. mon. anon. p. 120.* Flores 4 lin. longi. DC. syst. 1. p. 500.* (v. s.)
3. *X. SALICIFOLIA* (Humb. et B! nov. am. sp. 5. p. 63.) foliis oblongis acuminatis obtusiusculis subtus sericeis pedunculis brevibus unifloris? bracteolatis. ♀ in Americâ merid. propè Espinal. Dunal! mon. anon. p. 121. t. 17.* (v. s. in h. H. et B.)
4. *X. LIGUSTRIFOLIA* (Dunal! mon. anon. p. 121. t. 18.*) foliis oblongis acutiusculis utrinque glabris, pedunculis brevibus paucifloris bracteolatis. ♀ propè Buga Popayanensium Americæ merid. — DC. syst. 1. p. 500.* H. B. et K! nov. am. sp. 5. p. 63. (v. s. in h. H. et B.)
5. *X. GLABRA* (Lin. spec. 1367.) foliis oblongo-ovatis glabris, pedunculis 1-floris solitariis aut geminis, carpellis lævibus. ♀ in insulis Barbadosibus, in Jamaicâ. Dunal! mon. anon. p. 121. t. 19.* — Pluk. alm. 395. t. 238. f. 4. Alabastra oblonga, extus adpressè pubescentia, 4-lin. longa. DC. syst. 1. p. 501.* (v. s.)
6. *X. NITIDA* (Dunal! mon. anon. p. 122. t. 20.*) foliis oblongo-lanceolatis glabris supernè nitidis, pedunculis ramosis multifloris, calyce subintegro.

♂ in Cayennâ. Species refert paululum Unonam xylopioidem. DC. syst. 1. p. 501.* (v. s.)

7. *X. ACUMINATA* (Dunal! mon. anon. p. 122. t. 16.*) foliis oblongo-ellipticis acuminatissimis glaberrimis, capsulis univalvibus dispermis. ♂ in Cayennâ, Portorico. (Bertero.) Semina nigra fœtida obovata extus convexa, intus plana. DC. syst. 1. p. 501. (v. s.)
8. *X. PAINOIDES* (Dunal! mon. anon. p. 122. t. 15.*) foliis oblongo-lanceolatis acuminatis apice obtusiusculis glabris membranaceis, floribus solitariis, capsulis bivalvibus. ♂ in Cayennâ. Semina 2, extus concava, intus plana. DC. syst. 1. p. 501. (v. s.)
9. *X. BRASILIENSIS* (Spreng. neu. entd. 3. p. 50.) foliis lanceolatis acuminatis coriaceis subtus pilosis, ramusculis pedunculisque erectis elongatis 1-floris hirsutis, petalis crassiusculis velutinis apertis. ♂ in Brasiliâ. affinis dicitur *X. frutescenti*.

IX. GUATTERIA. Ruiz. et Pav. prod. p. 85. t. 17. Dunal! mon. anon. p. 50. et 123.

Sepala 3 ima basi coalita, ovata, subcordata, acuta. Petala 6 ovata, aut obovata; antheræ ∞ subsessiles; carpella ∞ subbaccata sicca coriacea ovata aut subglobosa stipitata monosperma. DC. syst. 1. p. 502.

1. *G. ABEREMOA* (Dunal! mon. anon. p. 126.) foliis ovato-oblongis acutis tomentosis, pedunculis axillaribus unifloris baccis substipitatis ovatis mucronatis. ♂ in Guianæ sylvis. Aberemoa Guianensis Aubl. Guian. 1. p. 610. t. 245.* — DC. syst. 1. p. 502.* (v. s.)

♂. *microcarpa*, à priori differt tantum fructibus paulò minoribus; an altera?

2. *G. BREVIPES* (DC. syst. 1. p. 502.*) foliis ovalibus subacuminatis utrinque glabris lucidis, pedunculis axillaribus unifloris, baccis ovatis breviter stipitatis. ♂ in Guianâ. Flores ignoti. (v. s. in h. Lamb.)
3. *G. OURÉGOU* (Dunal! mon. anon. p. 126.*) foliis oblongo-ellipticis acuminatis glabris, pedunculis 1-3 axillaribus, baccis ovatis acutis longè stipitatis. ♂ in sylvis Guianæ. Cananga Ourégou. Aubl. Guian. 1. p. 608. t. 244.* Baccæ siccæ coriaceæ lutescentes ovatae. — DC. syst. 1. p. 503.* (v. s.)

4. *G. PODOCARPA* (DC. syst. 1. p. 503.*) foliis ovali-oblongis abruptè acuminatissimis glabris, pedunculis axillaribus solitariis, baccis ovatis submucronatis stipite bacca multò longiore. ♂ in Cayennâ.

♂. *oligocarpa*, baccis 14-20 subcostatis (v. s. in h. Mus. Par.)

♂. *polycarpa*, baccis circiter 40 non costatis. (v. s. in h. Lamb.)

5. *G. CERASOIDES*. (Dunal! mon. anon. p. 127.) foliis lanceolatis acutis subtus pubescentibus, pedunculis axillaribus solitariis, petalis subæqualibus, baccis ovato-globosis, stipite bacca longiore. ♂ in Coromandelæ montibus. Uvaria cerasoides Roxb! corom. 1. p. 30. t. 33.* — DC! syst. 1. p. 503.* (v. s.)
6. *G. SUBEROSA* (Dunal! mon. anon. p. 128.*) foliis oblongis acutis glabris, pedunculis suboppositifoliis unifloris, petalis exterioribus minoribus, baccis globosis, stipite bacca duplò longiore. ♂ in Coromandelâ. Uvaria suberosa Roxb! corom. 1. p. 31. t. 34.*

♂. foliis basi angustioribus margine crispulis.

♂. foliis exactè oblongis pallidè glaucis, ramulis villosis

7. *G. RUFA* (Dunal! mon. anon. p. 129. t. 29.*) foliis ovalibus acuminatis basi cordatis subtus ramulisque rufo-tomentosis, pedunculis brevissimis lateralibus vel oppositifoliis; petalis æqualibus, baccis stipitatis velutinis. ♂ in Indiâ, insulâ Timor, Javâ. Uvaria tomentosa Vahl! lued. non Roxb! — DC. syst. 1. p. 504.* (v. s.)

8. *G. CORDATA* (Dunal! mon. anon. p. 129. t. 30.*) foliis basi cordatis oblongis

- acutiusculis subtiis ramulisque subtomentosis, racemis oppositifoliis brevibus paucifloris. 5 in Javâ. DC. syst. 1. p. 505. (v. s. in h. Deless.)
9. *G. ERIOPODA* (DC. syst. 1. p. 505.) foliis oblongo-lanceolatis basi subatenuatis acuminatis junioribus villosis, pedunculis lateralibus solitariis unifloris tomentoso-hispidis. 5 in Peruvîâ. Deless! ic. sel. 1. t. 90. (v. s. in h. Mus. Par. et Juss!)
10. *G. HIRSUTA* (Ruiz. et Pav. fl. per. syst. 1. p. 146.) foliis lanceolatis acuminatis, pedunculis axillaribus glabris, floribus hirsutis 5 in Peruvîæ nemoribus. Dunal. mon. anon. p. 131. DC. syst. 1. p. 505.
11. *G. PENDULA* (Ruiz. et Pav. fl. per. syst. 1. p. 146.) foliis lanceolatis, pedunculis axillaribus longissimis pendulis. 5 in Peruvîæ montibus. Dunal. mon. anon. p. 131. DC. syst. 1. p. 506. †.
12. *G. OVALIS* (Ruiz. et Pav. fl. per. syst. 1. p. 146.) foliis oblongis ovalibusque, pedunculis axillaribus. 5 in Peruvîæ montibus. Dunal. mon. anon. p. 131. DC. syst. 1. p. 506. †.
13. *G. GLAUCA* (Ruiz. et Pav. fl. per. syst. 1. p. 145.) foliis oblongis ovatisque acuminatis, pedunculis axillaribus. Dunal. mon. anon. p. 131. DC. syst. 1. p. 506. †.
14. *G. VIRGATA* (Dunal! mon. anon. p. 131. t. 31.*) foliis ovatis acuminatis glaberrimis subsessilibus, pedunculis axillaribus 1-floris, baccis substipitatis coriaceis ovoideis obtusiusculis. 5 in montosis sylvaticis Jamaicæ. Uvaria lanceolata Swartz! prod. 87. (v. s.)
15. *G. LAURIFOLIA* (Dunal! mon. anon. p. 132. t. 32.*) foliis oblongis utrinque acuminatis glabris brevissimè petiolatis, pedunculis axillaribus subconfertis, baccis substipitatis ovoideis mucronatis. 5 in sylvis Jamaicæ. Uvaria excelsa Vest! ex Vahl. in h. Juss. — DC. syst. 1. p. 507.*
16. *G. SEMPERVIRENS* (Dunal. monog. anon. p. 133.) foliis ovato-oblongis glabris nitidis, pedunculis 1-floris, petalis oblongis acutis, baccis stipitatis globosis. 5 Δ in plurimis regni Malabarici locis. Rheed. h. mal. 5. p. 31. t. 16.* — DC. syst. 1. p. 507.
17. *G. KORINTI* (Dunal. mon. anon. p. 133.) foliis ovato-oblongis glabris nitidis, pedunculis axillaribus 1-floris, petalis oblongis obtusiusculis, baccis stipitatis globosis. 5 ubiquè in Malabariâ. — Rheed. h. mal. 5. p. 27. t. 14.* — DC. syst. 1. p. 507.*
18. *G. MONTANA* (DC. syst. 1. p. 508.) foliis lanceolato-oblongis angustis glabris nitidis, pedunculis geminis ternisve 1-3-floris, petalis inflexis. 5 in Malabaricæ locis montanis et petrosis. *G. malabarica* Dunal! mon. anon. p. 134. — Rheed. h. mal. 5. p. 33. t. 17.*
19. *G. ACUTIFLORA* (Dunal! mon. anon. p. 134.*) foliis ovato-oblongis acutis crassis glabris, pedunculis axillaribus brevibus. calycinis lobis obtusis, petalis acutissimis, baccis stipitatis globosis. 5 in Malabaricæ locis montanis. — Rheed. h. mal. 5. p. 35. t. 18.* — DC. syst. 1. p. 508.* (v. s. in h. Deless!)
20. *G. UMBILICATA* (Dunal! mon. anon. p. 135. t. 33.*) foliis oblongo-ellipticis acuminatis glabris, pedunculis 1-floris bracteolatis, baccis stipitatis basi umbilicatis apice depressis. 5 in Americâ calidiore aut in Caribæis? DC. syst. 1. p. 508.* (v. s.)

†. *Species non satis notæ.*

21. *G. WILLEMETIANA* foliis lanceolatis glabris, pedunculis lateralibus compressis bifidis. 5. *Anona distincta*. Will. herb. maur. p. 42. †.
22. *G. MAYPURENSIS* (Humb. B. et K. nov. sp. am. 5. p. 64.) foliis oblongis acuminatis in petiolum decurrentibus subcoriaceis glabris nitidis, pedunculis axillaribus solitariis 1-floris, petalis oblongis obtusis exterioribus paulò minoribus. 5 propè Maypures in Miss. Orenocensibus.

Ordo V. MENISPERMACEÆ. — Menispermæ. *Juss. gen.*
284. *Jaum. fam.* 2. p. 82. t. 86. *DC. syst.* 1. p. 509.

Flores (abortu ?) unisexuales, sæpissimè dioïci minimi ; tegumenta floralia uni aut multiserialia, seriebus omnibus ternario quaternariove numero constantibus, hypogyna, decidua ; petala interdum o. ♂. Stamina monadelpha aut rariùs libera, nunc petalis numero æqualia et iisdem opposita, nunc petalis numero tripla quadruplæ ; antheræ adnatæ extrorsæ, aut basi ad apicem filamentum insertæ. ♀ Ovaria nunc plurima monostyla basi subcoalita, nunc solitaria stigmatibus plurimis coronata intus multilocularia et ideò è plurimis carpellis coalitis constantia, rarissimè et verosimiliter abortu unilocularia. Drupæ ferè in omnibus baccatæ, 1-spermæ, obliquæ aut lunulatae, compressæ ; semen conforme ; embryo curvus aut periphericus ; albumen nullum aut parvum carnosum ; cotyledones planæ, nunc approximatae nunc modò singulari distantes et in loculis duobus seminis locatæ ! radícula supera, interdum verò fructus apice cum basi per accrescendi modum contiguo infera dicenda.

Frutices sarmentosi flexiles tenaces. Folia alterna simplicia aut rariùs composita apice mucronata. Flores parvi sæpiùs racemosi.

Tribus I. LARDIZABALEÆ.

Flores sæpissimè dioïci, masculi quoad partium numerum symetrici. Carpella plurima distincta, polysperma, plurilocularia (an in omnibus ?) folia composita.

I. LARDIZABALA. *Ruiz et Pav. prod.* p. 143. t. 37.

Flores dioïci polygamivæ ; sepala petalæque ordine ternario 2-3-serialia. ♂ stamina 6 monadelpha. ♀ Baccæ 3-6, 6-loculares, loculis oo-spermis. *DC. syst.* 1. p. 511.

1. *L. BITERNATA* (*Ruiz. et Pav! syst.* 286. *prod.* t. 37.) foliis 2-3-ternatis, foliolis oblongis acutis basi inæqualibus hinc inde subdentatis, bracteis 2 magnis inæqualiter cordatis ad basim pedunculorum. \hookcup in regni Chilensis sylvis. *Lapeyr. voy.* 4. p. 265. t. 6. 7. 8. *opt. DC. syst.* 1. p. 512.* (v. s. ♂.)
2. *L. TRITERNATA* (*Ruiz. et Pav! syst.* 287.) foliis 2-3-ternatis, foliolis ovalibus obovatisve obtusis integerrimis, bracteis 2 magnis ovatis ad basim pedunculorum. \hookcup in Conceptionis Chili sylvis. *Deless! ic. sel.* 1. t. 91. Florem non vidi ; fructus 3 oblongo-cylindracei, juniores stigmatibus crasso coronati. *DC. syst.* 1. p. 512.* (v. s. ♀.)
3. *L. TRIFOLIOLATA* (*DC. syst.* 1. p. 513.*) foliis ternatis trifoliolatis, foliolis ovatis, bracteis secus pedunculos minimis. \hookcup in Peruvia. *Deless! ic. sel.* 1. t. 92. Petala 6 ovato-rotunda minima ad basim columnæ staminum sita (v. s. ♂.)

II. STAUNTONIA. DC. syst. 1. p. 513.

Sepala ordine ternario biserialia. Petala 0. ♂ Stamina 6 monadelphæ. ♀.....

1. *S. CHINENSIS* (DC. syst. 1. p. 514. *) 5 in Chinâ. Floris sepala 6 lin. longa; antheræ albidæ; folia petiolata, peltata, 5-foliolata (v. s. ♂ in h. Banks. et Lamb.)

III. BURASAIA. Petit-Th. dict. scienc. nat. ex gen. madag. p. 18.

Sepala et petala ordine ternario biserialia. ♂ stam. 6 basi coalita. ♀ stam. 6 sterilia. Ovarium triplex. Drupæ 3. DC. syst. 1. p. 511.

1. *B. MADAGASCARIENSIS* (Petit-Th. l. c.) 5 in Madagascar. Folia alterna longè petiolata; foliola 3 ovata integra. DC. syst. 1. p. 514.

Tribus II. MENISPERMEÆ.

Flores sæpissimè dioici, masculi quoad partium numerum symetrici. Carpella plurima distincta 1-locularia 1-sperma. — Folia simplicia.

IV. SPIROSPERMUM. Petit-Th. dict. scienc. nat. ex gen. mad. p. 19. n. 63.

Sepala et petala ordine ternario bi-serialia. ♂ Stam. 6, 3 interiora basi coalita. ♀ Carpella 8-9. (DC. syst. 1. p. 514.) †.

1. *S. PENDULIFLORUM* (Petit. Th. l. c.) 5 in Madagascar. DC. syst. 1. p. 514.

V. COCCULUS. C. Bauh. pin. 511. DC. syst. 1. p. 515.

Sepala et petala ordine ternario bi-rarissimè tri-serialia. ♂ Stam. 6. libera petalis opposita. ♀ Carpella 3-6. Drupæ baccatæ 1-6. sæpius obliquæ reniformes subcompressæ 1-sperma; Cotyledones distantes.

§. 1. Foliis peltatis.

1. *C. JAPONICUS* (DC. syst. 1. p. 516. *) foliis peltatis subrotundo-ovatis acuminatis integerrimis glabris, petiolo subvolubili folii longitudine, pedunculis femineis petiolo triplò brevioribus umbelliferis, carpellis binis. 5 in Japoniâ. Menispermum Japonicum Thunb. jap. 195. *

℞? *Timoriensis*, differt à descr. Thunbergianâ pedunculis (masculis equidem) petiolo æqualibus. An diversa species? (v. s.)

2. *C. ROXBURGHIANUS*. (DC. syst. 1. p. 516. *) foliis peltatis ovatis hinc rotundatis illinc subacutis integerrimis glabris, petiolo folio triplò brevior, pedunculis femineis petiolo triplò brevioribus umbelliferis, baccis subsenis. 5 in Indiâ orient. Cissampelos glabra Roxb. ex Wall! (v. s. ♀)

3. *C. PELTATUS* (DC. syst. 1. p. 516. *) foliis peltatis subtriangularibus acuminatis integerrimis subscabris, petiolis pilosis, pedunculis femineis racemosis petiolo vix duplò longioribus. 5 in Malabiâ et Coromandeliâ. — Pluk. phyt. t. 24. f. 6. Menispermum peltatum Lam. dict. 4. p. 96. *

4. *C. BURMANNI* (DC. syst. 1. p. 517. *) foliis peltatis triangulari-oblongis acuminatis integerrimis posticè obtusè truncatis nitidis, pedunculis masculis longissimis racemosis. 5 in Ceylonâ. Burm. Zeyl. 218. t. 101. * ♂ flores sexfidi. ♀ Pistillum 1. (Burm. l. c.)

5. *C. FORSTERI* (DC. syst. 1. p. 517.) foliis peltatis ovatis hinc rotundatis illinc subacutis integerrimis glabris, petiolo folio longiore. 5 Menispermum peltatum Forst! (ved. in h. Lamb. (v. s. in h. Lamb.) †.

§. 2. *Foliis basi cordatis.*

6. *C. ROTUNDIFOLIUS* (DC. syst. 1. p. 517.*) foliis subpeltatis basi subcordatis orbiculatis vix acuminatis mucronatis glabris, pedunculis axillaribus folio brevioribus racemoso-paniculatis. \smile anno 1812. in h. Paris. cultus. Baccæ nigricantes subglobosæ subcompressæ pisi magnitudine. (v. s. in h. Juss.)
7. *C. CORDIFOLIUS* (DC. syst. 1. p. 518.*) foliis orbiculatis cordatis acuminato-cuspidatis 7-nerviis utrinquè glabris, racemis femineis lateralibus folio longioribus simplicibus \smile in Malabariâ, Indiâ orient. Citamerdu Rheed. mal. 7. p. 39. t. 21.* *Menispermum cordifolium* Willd. spec. 4. p. 826.
8. *C. CONVOLVULACEUS* (DC. syst. 1. p. 518.*) foliis cordiformibus acuminato-cuspidatis 7-nerviis utrinquè glabris, pedunculis masculis axillaribus folio brevioribus simplicibus. \smile in Indiâ orient. Priori valdè affinis et forsân hujus individuum masculum? (v. s.)
9. *C. MALABARICUS* (DC. syst. 1. p. 518.*) foliis cordatis ovatis acuminatis subtus villosis, racemis femineis lateralibus? folii longitudine simplicibus. \smile in Malabariâ semper florens et frugens. Pee-amerdu Rheed. mal. 7. p. 37. t. 19 et 20.* *Menispermum Malabaricum* var. α . Lam. dict. 4. p. 96. Ex descriptione flores videntur hermaphroditi! an rectè?
10. *C. POPULIFOLIUS* (DC. syst. 1. p. 519.*) foliis cordiformibus acuminatis integerrimis glabris, paniculis femineis amplis multifloris. \smile in insulâ Timor. Aff. *C. cordifolio* et *Malabarico*. Carpella 1-3 baccata pedicellata subglobosa, pisi magnitudine. (v. s. comm. à Mus. Par.)
11. *C. LACUNOSUS* (DC. syst. 1. p. 519.*) foliis cordiformibus acuminatis superne viridibus glabris subtus flavo-lanuginosis, pedunculis lateralibus compositis racemosis petiolo duplò longioribus. \smile in rupibus et saxis ad littora maritima Celebum et Moluccarum. — Rumph. amb. 5. p. 35. t. 22.* *Menispermum lacunosum*. Lam. dict. 4. p. 98.
12. *C. SUBEROSUS* (DC. syst. 1. p. 519.*) foliis cordatis basi subtruncatis compactis lucidis, paniculis multifloris, caule perenni volubili, cortice suberoso rimoso. \smile in Malabar. Cocci orientales Tab. ic. st. 924. f. 1. *Menispermum cocculus* Lin. spec. 1468? excl. syn. Gaertn. fruct. 1. p. 219. t. 70. f. 1. excl. syn. (v. s. ex hort. Calc. comm. à cl. Lechenault.)
13. *C. PLUKENETII* (DC. syst. 1. p. 520.*) foliis ovatis basi subcordatis apice obtusè truncatis mucronulatis, racemis femineis axillaribus simplicibus folio sublongioribus. \smile in Malabariâ et in Javâ. *C. officinarum* Pluk. mant. 52. t. 345. f. 7. *Menispermum cocculus* Willd. spec. 4. p. 826. ?
14. *C. ARISTOLOCHIE* (DC. syst. 1. p. 520.*) foliis cordatis obtusis mucronatis integris, pedunculis femineis unifloris petiolo brevioribus. \smile in Maderaspatanâ provinciâ. Pluk. alm. t. 13. f. 2. †.
15. *C. FLAVESCENS* (DC. syst. 1. p. 520.*) foliis subcordatis ovatis obtusiusculè acuminatis, junioribus subpubescentibus orbiculatis, racemis femineis paniculatis lateralibus folio longioribus. \smile ad littora maritima insularum Moluccarum. — Rumph. amb. 5. p. 38. t. 24.* *Menispermum flavescens*. Lam. dict. 4. p. 98.
16. *C. GLAUCUS* (DC. syst. 1. p. 521.*) foliis cordiformibus acuminatis integris subtus pubescentibus, pedunculis racemoso-corymbosis petiolo brevioribus. \smile in apriciis vallibus et desertis hortis Amboinæ. — Rumph. amb. 5. p. 40. t. 25. f. 1.* *Menispermum glaucum* Lam. dict. 4. p. 100.
17. *C. CRISPUS* (DC. syst. 1. p. 521.*) foliis cordatis acuminatis glabris, caule subangulato tuberculis subrotundis exasperato, racemis gracilibus lateralibus simplicibus. \smile in Javâ et transportatus in Amboinâ et Baleiâ. — Rumph. amb. 5. p. 83. t. 44. f. 1. †.* *Menispermum crispum*. Lin. spec. 1468.
18. *C. TAMOIDES* (DC. syst. 1. p. 521.*) foliis basi cordatis truncatisve ovatis acuminatis 5-nerviis integerrimis glabris, racemis masculis gracilibus folio

- paulò longioribus. 5? \cup in Cayennâ. Aff. C. Carolino, sed differt foliis utrinquè glabris racemisquè multò longioribus. (v. s. in h. Mus. Par.)
19. C. CHONDODENDRON (DC. syst. 1. p. 522.*) foliis cordatis leviter crenatis cum acumine subtùs tomentosis. 5 \cup in Americâ merid. Chondodendron tomentosum. Ruiz et Pav. prod. fl. peruv. 132. syst. 261. Epibaterium? tomentosum Pers. ench. 2. p. 561. †.
20. C. HASTATUS. (DC. syst. 1. p. 522.*) foliis basi cordatis hastatis subtùs villosis, auricularis obtusis, apice ultrâ aurículas parùm producto obtuso, petiolis villosis folio sextuplò brevioribus. 5 \cup in Indiâ orientali. Menispermum hastatum. Lam. dict. 4. p. 98.* Flores ignoti.
21. C. TRILOBUS (DC. syst. 1. p. 522.*) foliis trilobis nervosis villosis, lobis acutis mucronatis integerrimis, petiolis basi reflexis, pedunculis racemosis petiolo brevioribus. 5 \cup in Japoniâ. Menispermum trilobum. Thunb. fl. jap. 194.* Bacca globosa. 1-sperma pisi minoris magnitudine.
22. C. PALMATUS (DC. syst. 1. p. 522.*) foliis basi cordatis palmatim 5-fidis piloso-hispidis, lobis acuminatis integerrimis. 2 \cup in Africâ australi. Menispermum palmatum Lam! dict. 4. p. 99.* — Berry in Asiat. res. 10. p. 385. t. 5. ♂ — Hæc est planta cujus radix in officinis usitata sub nominibus *Columbo* seu *Calumba*, *Calomba*, *Colomba*. (v. s. sine fl.)
23. C. ORBICULATUS (DC. syst. 1. p. 523.*) foliis orbiculatis subcordatis obtusis 5-7. nerviis mucronulatis subtùs cinereo-pubescentibus, pedunculis maximis 1-3 apice racemosis petiolo brevioribus. 5 \cup in Malabariâ, Tranquebariâ. — Pluk. am. 61. t. 384. f. 6. Flores minimi extùs villosi, ♀ ignoti. — An Cissampelos?
24. C. DIVERSIFOLIUS (DC. syst. 1. p. 523.*) foliis inferioribus cordatis, mediis ovatis, summis oblongis, omnibus truncato-obtusis mucronatis, pedunculis folio brevioribus 1-3-floris 5 \cup in Mexico. Menispermum sp. nov. Mocq. et Sessé fl. mex. ic. ined. C. Carolino valdè affinis sed apice foliorum truncato distinctus.
25. C. CAROLINUS (DC. syst. 1. p. 524.*) foliis cordatis ovatisve integris obtusisve subtrilobis subtùs pubescenti-velutinis, racemis masculis à basi floriferis, femineis 3-floris. 5 et sæpè in hortis septentrionalibus 2 \cup in sylvis et sepibus Carolinæ, Georgiæ et Floridæ. Menispermum Carolinum. Lin. spec. 1468. Wendlandia populifolia Willd. spec. 2. p. 275. — Dill. elth. 223. t. 178. f. 219.* (v. s.)

§. 3. Foliis ovatis ovalibus oblongisve.

26. C. TRUNBERGII (DC. syst. 1. p. 524.*) foliis ovatis obtusis cum acumine subtùs villosis, infimis subtriangularibus, summis orbiculatis, floribus paniculatis. \cup in Japoniâ. Menispermum orbiculatum Thunb. fl. jap. 194. Flores axillares paniculati. — Confer t. 2. f. 1. Braam. ic. Chin. 1821. an. sit hæc aut nova Cocculi species?
27. C. VILLOSUM (DC. syst. 1. p. 525.*) foliis ovatis lanceolatisve 3-5-nerviis, junioribus villosis-tomentosis, adultis pubescentibus, ramulis villosis, pedicellis paucifloris petioli longitudine. 5 seu 2 \cup in Malabariâ, Indiâ. Menispermum villosum. Lam. dict. 4. p. 97.*
- α. foliis oblongo-lanceolatis. — Pluk. amalth. 62. t. 384. f. 3. — Menispermum myosotoides Lin. spec. 1469.
- β. foliis omnibus ovatis. Pluk. am. 61. t. 384. f. 7. — Menispermum hirsutum Lin. spec. 1469.
- γ? foliis superioribus ovatis, infimis cordatis. Pluk. am. 61. t. 384. f. 5.
28. C. COTONEASTER (DC. syst. 1. p. 525.*) foliis ovalibus integerrimis mucronatis subtùs tomentosis, ramis tomentosis, pedunculis racemosis petiolo longioribus. 5 \cup Verosimiliter in Americâ calid. Deless! ic. sel. 1. t. 93. Folia Mespilum cotoneasterem æmulantia, sed paulò longiora. (v. s.)

29. *C. FIBRAUREA* (DC. syst. 1. p. 525.*) foliis ovatis acutis integerrimis glabris longè petiolatis, racemis oblongis lateralibus. 5 in sylvis Cochinchinæ et Chinæ. *Fibraurea tinctoria* Lour. fl. coch. ed. Willd. 2. p. 769.*
30. *C. OVALIFOLIUS* (DC. syst. 1. p. 426.*) foliis ovalibus integris mucronatis trinerviis glabris, ramulis villosis, pedunculis inferioribus axillaribus petiolo vix longioribus, summis in racemum terminalem dispositis. 5 in Javâ, et Chinâ. Deless! ic. sel. 1. t. 94. — *Menispermum ovalifolium* Pers! ench. 2. p. 628. Baccæ 2-3 glabræ compressæ orbiculatæ. (v. s.)
31. *C. ELLIPTICUS* (DC. syst. 1. p. 526.*) foliis ellipticis obtusis integerrimis glabris, racemis axillaribus geminis inæqualibus folio multò brevioribus. 5 in Senegaliâ. *Menispermum ellipticum* Poir. suppl. 3. p. 657.* Flores virentes parvi in racemos parvos breves inæquales dispositi.
32. *C. LIMACIA* (DC. syst. 1. p. 526.*) foliis ovato-oblongis acuminatis integerrimis glabris, floribus masculis subterminalibus congestis, femineis binatis axillaribus. 5 in sylvis Cochinchinæ. *Limacia scandens* Lour. fl. coch. ed. Willd. 2. p. 761.* Flores luteo-virides.
33. *C. CEBATHA* (DC. syst. 1. p. 526.*) foliis ovali-oblongis mucronatis nitidis glabris, pedunculis axillaribus petioli longitudine, masculis capitellatis, femineis unifloris. 5 in Arabiâ. *Cebatha*. Forsk. desc. æg. ar. 171.* *Menispermum edule* Vahl. symb. 1. p. 80.
34. *C. ACUMINATUS* (DC. syst. 1. p. 527.*) foliis ovato-acuminatis integerrimis glabris basi sub-5-nerviis, ceterum penninerviis, racemis axillaribus subvillosis petiolo paulò longioribus. 5 in Indiâ orientali. Deless! ic. sel. 1. t. 95. *Menispermum acuminatum* Lam. dict. 4. p. 101.* Habitu aff. *C. radiato*. Baccæ obovatæ plurimæ stipitatæ glabræ. (v. s.)
35. *C. RADIATUS* (DC. syst. 1. p. 527.*) foliis ovato-oblongis acuminatis vix subcordatis integerrimis penninerviis glabris, pedunculis racemoso-paniculatis petiolo triplò longioribus. 5 in Indiâ orient. Valli-Caurim Rheed. mal. 7. p. 5. t. 3.* *Menispermum radiatum* Lam. dict. 4. p. 100.* An hæc species cum priore genus proprium conficere debeat? (v. s. in h. Mus. Par.)
36. *C. LEPTOSTACHYUS* (DC. syst. 1. p. 528.*) foliis ovalibus acuminatis trinerviis glabris, racemis axillaribus simplicibus gracilibus folii longitudine. 5 in insulâ Timor. Flores immaturi minimi, in sicco fuscii. (v. s. comm. à Mus. Par.)
37. *C. BRACHYSTACHYUS* (DC. syst. 1. p. 528.*) foliis ovatis acutis 3-5-nerviis glabris, racemis femineis axillaribus petiolo brevioribus. 5 in insulâ Timor. Semen intra baccam arcuatum. Flores ignoti. (v. s. comm. à Mus. Par.)
38. *C. DOMINGENSIS* (DC. syst. 1. p. 528.*) foliis ovalibus acuminatis integerrimis vix basi subtrinerviis glabris, pedunculis racemoso-paniculatis axillaribus gracilibus folio subbrevioribus è basi tuberculatâ villosâ ortis. 5 in Sancto Domingo. Deless! ic. sel. 1. t. 96. Aff. *C. brachystachyo*. flores minimi, in specimine immaturi. (v. s. in h. Mus. Par.)
39. *C. LEÆBA* (DC. syst. 1. p. 529.) foliis ovato-oblongis obtusis glaucis subpubescentibus breviter petiolatis, ramis subvolubilibus. 5 in Ægypto superiore. *Leæba* Forsk. fl. æg. 172.* Juss. gen. 285. *Menispermum Leæba* Delil! fl. æg. ill. 30. desc. t. 51. f. 2. 3. (v. s. in h. Delil.)
40. *C. OBLONGIFOLIUS* (DC. syst. 1. p. 529.*) foliis oblongis trinerviis utrinquè obtusis mucronatis glabris, pedunculis folio brevioribus, supremis subracemosis, femineis unifloris. 5 in Mexico. *Menispermum nov. spec. Moc. et Sessé fl. mex. ic. ined. (Descr. ex ic.)*
41. *C. TRIFLORUS* (DC. syst. 1. p. 529.*) foliis ovato-lanceolatis acuminatis basi trinerviis integerrimis subglabris, pedunculis femineis axillaribus trifidis trifloris petioli longitudine. 5 in Javâ. Flores ignoti. (v. s. in h. Mus. Par.)
42. *C. MILLEFLORUS* (DC. syst. 1. p. 530.*) foliis ovatis obtusis penninerviis glabris nitidis, paniculæ terminalis multifloræ ramulis patentibus geminatis.

♂ in Madagascar. Azonn-minti, fruit velouté en poire. Poivre! in h. Juss.
♀ ignota. (v. s. ♂ in h. Juss.)

43. *C. GOMPHIOIDEA* (DC. syst. 1. p. 530.*) foliis oblongis acuminatis basi trinerviis integerrimis glabris nitidis, pedunculis axillaribus folio duplò longioribus. ♂ in Madagascar. Baccæ uvæ minoris magnitudine æquales. (v. s. in h. Mus. Par.)

44. *C. LAURIFOLIUS* (DC. syst. 1. p. 520.*) foliis oblongis utrinquè acuminatissimis glabris nitidis, pedunculis axillaribus apice ramosis petiolo subbrevioribus. ♂ in Indiâ orientali. Deless! ic. sel. 1. t. 97. flores minimi 8-10. glabri. (v. s. ♂)

§. 4. *Floribus monoicis. (An genus?)*

45. *C. EPIBATERIUM* (DC. syst. 1. p. 530.*) floribus monoicis, drupis 1-3, foliis oblongis integerrimis glabris. ♂ in insulâ Sti-Jacobi maris australis. *Epibaterium pendulum*. Forst. gen. t. 54.)

46. *C. NEPHROÏA* (DC. syst. 1. p. 531.*) floribus monoicis, drupis 6, foliis ovatis emarginatis glabris. ♂ in sylvis Cochinchinæ. *Nephroïa sarmentosa*. Lour. fl. coch. ed. Willd. 2. p. 692.* — Drupæ minutæ carnosæ subreniformes monospermæ.

VI. *PSELIUM*. Lour. fl. coch. ed. Willd. 2. p. 762.

♂ Sepala et petala ordine ternario biserialia, stamina 6 libera. ♀ Sepala 4. Petala 0. ovarium unicum, stigma 4-fidum.

1. *P. HETEROPHYLLUM* (Lour. fl. coch. ed. Willd. 2. p. 762.*) ♂ in sylvis Cochinchinæ. folia alterna integerrima glabra. DC. syst. 1. p. 531.

VII. *CISSAMPELOS*. Lin. gen. n. 1138. Lam. ill. t. 830.

♂ Sepala 4. Petala 0. Stamina monadelphæ 4 aut 2. ♀ sepalum unilaterale. Petalum 1 antè petalum. Ovarium 1. Stigmata 3. Drupa obliqua reniformis. Albumen 0. Embryo longus teres periphericus; radícula supera nempè è stigmatum basi orta. — frutices scandentes.

§. *Floribus femineis bracteatis, foliis utriusque uniusve sexûs peltatis.*

1. *C. TROPÆOLIFOLIA* (DC. syst. 1. p. 532.*) foliis peltatis ovato-orbiculatis subacutis pilis sparsis utrinquè pubescentibus, racemis axillaribus geminis. ♂ in Americâ merid. Deless! ic. sel. 1. t. 98. fructus compressi orbiculati, striis radiantibus margine notati. (v. s. in h. Mus. Par.)

2. *C. HERNANDIFOLIA* (Wild. spec. 4. p. 861.*) foliis peltatis ovatis acuminatis obtusis mucronatis glabris, venis subtus subpilosis. ♂ in Indiâ orient. flores ignoti. Baccæ 1-spermæ corymbosæ. DC. syst. 1. p. 533.*

3. *C. OWARIENSIS* (Beauv! ined.) foliis peltatis ovato-rotundatis aristatis sublobatis suprâ glabris subtus tenuissimè pubescentibus, floralibus reniformibus longè ciliatis. ♂ in Africæ equinoxialis regno Owariensi. (v. s. ex herb. Beauv. nunc Deless.)

4. *C. PAREIRA* (Lam. ill. t. 830.) foliis peltatis subcordatis ovato-orbiculatis subtus sericeo-pubescentibus, racemis femineis folio longioribus, baccis hispidis. ♂ in fruticetis montosis Martinicæ, Jamaicæ, etc. *C. Pareira* var. α Lin. spec. 1473. Sw! obs. 380. t. 10. f. 5. — Radix hujus stirpis est vera *Pareira-Brava* officinarum. — DC. syst. 1. p. 533.* (v. s.)

5. *C. GUAYAQUILENSIS* (H. B. et K. nov. sp. am. 5. p. 67.) foliis peltatis subreniformibus utrinquè hirtio-pubescentibus subtus canescentibus, cymis masculis axillaribus geminis petiolo-brevioribus. ♂ ad Guayaquil Quitensiam ♀ ign. an var. *C. pareiræ*?

6. *C. ARGENTEA* (H. B. et K. nov. sp. am. 5. p. 67.) foliis peltatis subrotundo-reniformibus apice emarginatis suprâ pubescentibus subtus sericeo-tomentosis

- et argenteis, cymis masculis axillaribus geminis petiolo brevioribus. $\frac{1}{2}$ ad ripam fl. Magdalensæ propè Mompox \varnothing ign. au. var. *C. pareiræ*.
7. *C. MICROCARPA* (DC. syst. 1. p. 534.*) foliis subpeltatis subreniformibus orbiculatis subtus tomentosis, baccis pubescentibus demum glabris. $\frac{1}{2}$ in insulis Caribæis, Jamaicâ, Martinicâ, Sancto-Domingo. *C. Pareira* var. β . Lin. spec. 1473. *C. Pareiræ* valde affinis, differre tamen videtur. (v. s. in h. Mus. Par.)
8. *C. HETEROPHYLLA* (DC. syst. 1. p. 534.*) foliis caulinis peltatis orbiculatis basi cordatis subtus subpubescentibus, floralibus corlato-reniformibus utrinquè velutinis. $\frac{1}{2}$ in Novâ-Hispaniâ. flores σ minimi hispidi 4-sepali; \varnothing ignoti. (v. s. σ .)
9. *C. DISCOLOR* (DC. syst. 1. p. 534.*) foliis peltatis latè ovatis basi truncatis apice acuminatis 7-9-nerviis, subtus pubescentibus albidis, supernè glabris pedunculis masculis 2-3 petiolo triplo brevioribus. $\frac{1}{2}$ in insulis Moluccanis. flores σ minimi corymboso-ramosi; \varnothing ignoti. (v. s. σ comm. à cl. Lamb.)
10. *C. HIRSUTA* (Buchan! in DC. syst. 1. p. 535.*) foliis peltatis ovato-orbiculatis subaveniis subtus tomentosis supernè pubescentibus, pedunculis masculis petioli longitudine. $\frac{1}{2}$ in Napauliâ. flores σ plurimi minutissimi, villosi; \varnothing ignoti. (v. s. σ in h. Lamb.)
11. *C. MAURITIANA* (Petit-Th. in Journ. bot. 1809. 2. p. 65. t. 3 et 4. opt.) foliis cordato-orbiculatis utrinquè pubescenti-villosis, masculi peltatis, feminei palaribus, racemis masculis axillaribus geminatis pluribusve. $\frac{1}{2}$ in sylvis insularum Mauritanarum ubi pro *Pareirâ-bravâ* habetur. *C. pareiroides* DC. ess. prop. pl. ed. 2. p. 78. — DC. syst. 1. p. 535.* (v. s.)
12. *C. ORINOCENSIS* (H. B. K. nov. sp. am. 5. p. 68.) foliis vix peltatis deltoideo-cordatis obtusis suprâ puberulis subtus sericeo pubescentibus, racemis fructiferis petiolum vix superantibus, fructibus tuberculosi pilosis. $\frac{1}{2}$ in umbrosis ad fl. Orinocum. Affinis *C. tamoidi*.
- \S . 2. *Floribus femineis bracteatis, foliis non peltatis.*
13. *C. TOMENTOSA* (DC. syst. 1. p. 535.*) foliis orbiculatis cordatis utrinquè petiolatis caulibusque tomentosis, pedunculis masculis axillaribus pluribus folio brevioribus. $\frac{1}{2}$ in agro Campechiano. Flores σ villosi minimi; \varnothing ignoti. (v. s. σ .)
14. *C. TAMOIDES* (Willd! ined. DC. syst. 1. p. 536.*) foliis cordiformibus utrinquè pubescentibus acutis basi 5-nerviis, racemis masculis folio longioribus. $\frac{1}{2}$ in Brasiliâ. Affinis *C. Pareiræ*. Flores σ minimi, \varnothing ignoti. (v. s. in h. Desf.)
15. *C. TRILOBA* (Spreng. nene entd. 2. p. 152.) foliis cordatis subtrilobis utrinquè pubescentibus denticulato-ciliatis 5-nerviis, pedunculis femineis aggregatis axillaribus. $\frac{1}{2}$ in Brasiliâ. Bractæ ovatæ extus villosissimæ 9-10-nerviæ, denticulato-barbatæ. An forsâ femineæ *C. tamoidis*?
16. *C. CONVULVULACKA* (Willd! spec. 4. p. 863.*) foliis cordiformibus acuminatis subtus pubescentibus supernè glabris, racemis femineis petiolo paulò longioribus. $\frac{1}{2}$ in Indiâ orient. Flores \varnothing minimi, fructus et σ ignoti. DC. syst. 1. p. 536. (v. s. σ in h. Desf.)
17. *C. CAAPEBA* (Lin. spec. 1473.) foliis suborbiculatis basi cordatis 7-nerviis subpubescentibus, racemis femineis petioli longitudine. $\frac{1}{2}$ in Sancto-Domingo. Plm. ed. Burm. t. 67. f. 2. Baccæ parvæ reniformes, vix compressæ, hispidulæ. Fl. σ ignoti. DC. syst. 1. p. 536.* (v. s. \varnothing)
18. *C. ORBICULATA* (DC. syst. 1. p. 537.*) foliis orbiculatis basi cordatis subtus villosis supernè pubescentibus, pedunculis masculis subternis petiolo brevioribus. $\frac{1}{2}$ in Indiâ orient. *C. tetrandra* Roxb! ined. in h. Lamb. —

Foliatione affinis Menispermis orbiculato. Baccæ parvæ subcompressæ hirsutæ. (v. s. ♂ et ♀.)

19. *C. CRENATA*. (DC. syst. 1. p. 537.*) foliis ovato-orbiculatis subsessilibus 7-nerviis obtusis mucronulatis crenato sinuatis utrinquē subtomentosis, racemis femineis binis folio longioribus. 5 in Brasiliâ. — Fructus ovato-orbiculati pubescentes reticulato-nervosi subcompressi. (v. s. ♀)
20. *C. OVALIFOLIA*. (DC. syst. 1. p. 537.*) foliis ovalibus subacutis coriaceis, subtilis cinereo-incanis, superne glabris lævibus, racemis masculis subbinis hispidis petiolo triplò longioribus. 5 in Brasiliâ. Flores ♂ minimi atropurpurei extus villosi; ♀ ignoti. (v. s. in h. Mus. Par.)
21. *C. OVATA* (Poir. dict. 5. p. 11.*) foliis ovatis obtusis mucronatis superne lævibus subtus in nervis pubescentibus, racemis femineis petiolo longioribus, bracteis subulatis. 5 in Indiâ orient. Baccæ reniformes orbiculatæ subcompressæ glabræ lentis magnitudine. DC. syst. 1. p. 538.
22. *C. ACUMINATA* (DC. syst. 1. p. 538.*) foliis ovali oblongis acuminatis glabris basi trinerviis, pedunculis masculis axillaribus racemoso-paniculatis folio dimidio brevioribus. 5 in Indiâ orient. — ♀ ignoti.
23. *C. CAPENSIS* (Thunb. prod. 110.) foliis ovatis obtusiusculis glabris brevissimè petiolatis, racemis ramosissimis masculis? petiolo vix longioribus. 5 ad Caput B.-Spei. Caules frutescentes volubiles ramosi subvelutini, petioli breves pubescentes. Flores minimi tomentosi. Poir. dict. 5. p. 11.* DC. syst. 1. p. 538.* Burch! cat. n. 757. (v. s.)
24. *C. HUMILIS* (Poir. dict. 5. p. 11.*) foliis ovatis basi truncatis obtusis glabris petiolatis, axillis lanatis. 5 ad Caput B.-Spei. — Specimina 2 habeo, unum hortense suberectum, alterum Capense volubile, an diversa? — An Cissamp. Capensis, *C. humilis* et *C. calcarifera* proprium genus formare debent? DC. syst. 1. p. 538. excl. syn. Thunb., etc. (v. s. sine fl.)
25. *C. CALCARIFERA* (Burch! cat. n. 1795.) foliis elliptico-oblongis utrinquē obtusis brevissimè petiolatis cinereo-subpuberulis ad basim petioli externam calcaratis, floribus masculis ad axillas glomeratis subsessilibus. 5 ad Caput B.-Spei. — *C. fruticosa* Thunb. prod. 110? Lin. f. suppl. 432? — Caulis erectus ramosus. (v. s.)
26. *C. LAURIFOLIA* (Poir. dict. 5. p. 11.*) foliis ovato-oblongis obtusis glaberrimis, petiolo brevi crasso, racemis femineis brevibus, fructu maximo. 5 in insulâ Sancti-Thomasii. Fructus ovati basi angustati pulposi rugosi nucis magnitudine, sicci fusco-nigricantes. DC. syst. 1. p. 539.
27. *C. ANGUSTIFOLIA* (Burch. cat. n. 1717. ex itin. 1. p. 389.) foliis lato-linearibus apice rotundato sæpè emarginato, caule volubili. — ad Cap B.-Spei. Cœt. ign.

§. 3. *Racemis femineis ebracteatis.*

28. *C. ANDROMORPHEA* (DC. syst. 1. p. 539.*) foliis subcordiformibus glaberrimis membranaceis integris mucronatis, pedunculis femineis aggregatis ramosis ebracteatis. 5 in Cayennâ. Deless! ic. sel. 1. t. 99. — Species distinctissima et forsau in genus proprium commutanda? (v. s. ♀.)

VIII. MENISPERMUM. Tourn. mem. acad. par. 1705. p. 237. Lam. ill. t. 824.

Sepala et petala ordine quaternario bi seu triserialia. ♂ Stamina 16-20. ♀ Ovaria 2-4. Drupæ baccatæ subrotundo-reniformes 1-spermæ. — Frutices scandentes.

1. *M. CANADENSE* (Lin. spec. 1468.) foliis peltatis subglabris subcordatis subrotundo-angulatis, angulis obtusiusculis, terminali abruptè aristato mucronato, racemis solitariis compositis, petalis 8. 5 in Americâ boreal. Sims. bot. mag. t. 1910. — *M. Canadense* var. *α*. Lam. dict. 4. p. 95.* ill. t. 824. — DC. syst. 1. p. 540.* (v. v. et s.)
2. *M. DAURICUM*. (DC. syst. 1. p. 540.*) foliis peltatis glabris cordatis angu-

latis, angulis acutis, terminali acuminato vix mucronato, racemis geminis capitellatis) in Dahuriæ collibus saxosis. Deless! ic. sel. 1. t. 100. — *Trilophus ampelisagria* Fisch! in h. Gorenki. (v. s.)

3. *M. SMILACINUM* (DC. syst. 1. p. 541.*) foliis peltatis subglabris cordato-subrotundis obtusè angulatis subtus glaucis, racemis simplicibus, petalis 4. 5 in Carolinâ. — *Cissampelos smilacina* Lin. sp. 1473. Jacq. coll. 4. p. 128.* ic. rar. 3. t. 629. Flores ♀ ignoti.
4. *M. LYONI* (Pursh. fl. bor. am. 2. p. 371.) foliis cordatis palmato-lobatis longissimè pedicellatis, racemis simplicibus, floribus 6-petalis 12-andris. 2 in Kentucky et Tennessee. DC. syst. 1. p. 541.

†. *Menisperma non satis nota.*

5. *M?* *FENESTRATUM* (Gaertn. fruct. 1. p. 219. t. 46. f. 5.) cotyledonibus foraminibus fenestratis in Ceylonâ. Fructus solus cognitus est drupa baccata pubescenti-incana. DC. syst. 1. p. 541.*
6. *M?* *ACUTUM* (Thunb. fl. jap. 193.*) foliis cordatis posticè angulatis apicè acuminatis glabris in Japoniâ circâ Nagasaki. Lam. dict. 4. p. 96.—Flores ignoti. DC. syst. 1. p. 541.

IX. *ABUTA*. Barrer. fr. æq. 1. Aubl. Guian. 1. p. 618.

Flores Baccæ ex eodem receptaculo 2-3 magnæ ovatæ exsuccæ subcompressæ, testâ fragili 1-spermâ fœtæ. (Aubl.) Frutices scandentes. †.

1. *A. RUFESCENS* (Aubl. Guian. 1. p. 618. t. 250.) foliis ovatis acutis integris subtus velutinis cinerascensibus rufisve. 5 in sylvis Cayennæ et Guianæ. A. scandens Barrer. fr. æq. p. 1. — Apud Guianenses planta dicitur *Paireira-brava* et forsan hujus radix cum *Cissampelo pareirâ* confusa in nostris officinis prostat. DC. syst. 1. p. 542. (v. s. in h. Mus. Par.)
2. *A. CANDICANS* (Richard! ined. in h. Juss.) foliis ovatis acuminatis subcrenatis lobulatisve subtus glabris candicantibus. 5 in Cayennâ. Cayennensibus dicitur *Liane amère*. DC. syst. 1. p. 543.* (v. s. sine flore.)

X. *TRICHOA*. Pers. ench. 2. p. 634. Batschia Thunb. nov. act. ups. 5. p. 120. t. 2.

Flores dioici. Calyx 3-sepalus. Petala 3 coriacea villosa medio approximata apice reflexa. ♂ Stamina 6 disco inserta, 3 exteriora sterilia petalis alterna, 3 centralia monodelpha fertilia. ♀ stamina 6 sterilia, filamentis apice bimaculatis; carpella 3 drupacea coriacea, oblonga villosa; semen in concamerationes 2 plicatum. — Frutices scandentes, foliis alternis simplicibus, †.

1. *T. RACEMOSA* (Pers. ench. 2. p. 634.) racemis axillaribus solitariis, floribus raris 5 Mariquitæ Americæ calidioris. Batschia racemosa. Thunb. l. c. p. 123. t. 2. f. 1.
2. *T. CONFERTA* (Pers. ench. 2. p. 634.) spicis axillaribus solitariis, floribus confertis. 5 in Americâ, cum priore. Batschia conferta Thunb. l. c. t. 2. f. 2.

XI? *AGDESTIS*. Moç. et Sessé fl. mex. ic. ined.

Flores hermaphroditi. Sepala 4. Petala 0. Stamina 24. Antheræ utriusque bifidæ. Carpella 4 monostyla coalita in ovarium unicum 4-sulcatum, stigmatibus 4 patentibus apice subreflexis. †.

1. *A. CLEMATIDEA* (Moç. et Sessé fl. mex. ic. ined.) 5 in Novâ-Hispaniâ. Flores rufescentes, magnitudine et numero sepalorum quaterno petalorumque nullo Clematidem flammulam æmulantes. Fructus ignotus. DC. syst. 1 p. 543. (v. s. comm. à cl. Lamb.)

Trib. III. *SCHIZANDREÆ*. DC. syst. 1. p. 543.

Flores monoïci, masculi quoad partium numerum non symmetrici.

XII. *SCHIZANDRA*. Michx. fl. bor. amer. 2. p. 18.

Flores monoïci. Sepala ordine ternario triserialia. Petala 0. ♂ Antheræ 5 apice summo coalitæ. ♀ Ovaria ∞. Baccæ secus receptaculum elongatum spicatæ. — Suffrutex glaber subscandens, foliis ovali-lanceolatis. — Menispermis ex. cl. Rich. et Michx. affinis.

1. *S. COCCINEA* (Michx! fl. bor. amer. 2. p. 219. t. 47. Sims. bot. mag. t. 1413.)
 5 in umbrosis sylvæ Carolinæ et Georgiæ. DC. syst. 1. p. 544. (v. s. ♀.)

Ordo VI. BERBERIDEÆ. *Vent. tabl. 3. p. 83. DC. fl. fr. ed. 3. vol. 4. p. 627. syst. 2. p. 1.*

Sepala 3-4, aut sæpius 6, decidua, oblonga aut ovalia, sæpè subcolorata, duplici serie alternatim sæpissimè disposita, extùs squamis petaloideis uni aut multiseriabilibus stipata. Petala sepalorum numero æqualia et iis opposita, rariùs sepalorum numero dupla, hypogyna, sæpius ad basim intùs glandulis squamisve aucta. Stamina tot quot petala, iis opposita; filamenta brevía; antheræ oblongæ, adnatæ, biloculares, loculis sæpè discretis, à basi ad apicem valvulâ subelastica (ut in Lauro.) dehiscentibus. Ovarium (abortu aliorum) solitarium, ovatum, subobliquum, 1-loculare, stylo sublaterali brevissimo et stigmate suborbiculato coronatum. Fructus baccatus capsularisve. Semina imæplacentæ laterali adfixa, rarissimè solitaria, sæpius 2-3, ovata aut globosa. Albumen carnosum, sæpè subcorneum. Embryo rectus, axilis, radiculâ ad apicem plus minùsve incrassatâ, cotyledonibus planis. — Berberideæ pleræque, in zonâ temperatâ hemisphæræ borealis habitant et in montosis stant: nonnullæ tamen in Americâ australi usque ad plagas Magellanicas repertæ fuerunt. — Frutices aut Herbæ perennes, sæpius glabræ.

I. BERBERIS. *Lin. gen. n. 442. Gærtn. fruct. 1. p. 200. t. 42. f. 6.*

Sepala 6 squamis 3 extùs stipata. Petala 6 intùs biglandulosa. Stamina edentula. Bacca 2-3-sperma. Semina 2, rariùs 3 ad basim lateraliter inserta, erecta, oblonga, testâ crustaceâ, albumine carnosò, cotyledonibus foliaceis ellipticis, radiculâ longâ apice capitellatâ. — Frutices foliis primariis abortivis et in spinam sæpius mutatis, secundariis in axillis fasciculatis. Flores in omnibus flavi.

§. 1. *Foliis simplicibus, pedunculis multifloris racemosis.*

1. B. VULGARIS (Lin. spec. 472.) spinis tripartitis, foliis subobovatis ciliato-serratis, racemis multifloris pendulis, petalis integris. ♂ in incultis sepibus et frutetis apricis, præsertim calcareis, totius Europæ, Asiæ occidentalis, etc. — fl. dan. t. 904. — DC. syst. 2. p. 5.* (v. v.)

β. *lutea*, fructibus luteis. L'Her ined.

γ. *violacea*, fructibus violaceis Willd. arb. 34.* Poit. et Turp. arb. fr. 59.

δ. *purpurea*, fructibus purpureis, foliis angustis vix ciliatis. — B. innominata Kiehm. dec. rar. pl. tub. p. 18.

ε. *nigra*, fructibus nigris, foliis oblongis serrulatis paucis ciliato-serratis.

ζ. *alba*, fructibus candidis. Mill. dict. n. 1.

η. *asperma*, seminibus abortivis. Mill. l. c. B. asperma. Clus. hist. 1. p. 121.

ν? *Iberica*, racemis erectiusculis, foliis oblongis subintegris. B. vulgaris? Iberica Stev. et Fisch. in litt. (v. s.)

2. B. EMARGINATA (Willd! enum. 1. p. 395.) spinis tripartitis, foliis lanceolato-obovatis ciliato-serratis, racemis vix pendulis folio brevioribus, petalis emarginatis. ♂ in Sibiria. Simillima B. vulgaris sed ex cl. Link. differt racemis multò brevioribus paucifloris vix pendulis.

3. *B. CANADENSIS* (Mill. dict. n. 2.) spinis tripartitis, foliis obovato-oblongis remotè serratis, summis subintegris, racemis multifloris nutantibus 3 in colibus fertilibus et inter rupes Americæ borealis. Nutt. gen. am. 1. p. 210.* foras non satis a *B. vulgari* distincta. DC. syst. 2. p. 7.* (v. v.)
4. *B. SINENSIS* (Desf. cat. ed. 1804. p. 150.) spinis tripartitis subnullis, foliis oblongis obtusis integris aut infimis subdentatis, racemis multifloris nutantibus. 3 in Chinâ. Bacca ovalis, intensè rubra, 1-2-sperma. DC. syst. 2. p. 8.* (v. s. et v.)
5. *B. ARISTATA* (DC. syst. 2. p. 8.*) spinis infimis 3-partitis, superioribus simplicibus basi vix bidentatis, foliis oblongis glabris dentibus 4-5 spinulosis utrinquè serratis, racemis patulis erective ∞ -floris. 3 in Napauliâ. Ovaria oblongo-teretia, stylo crasso brevissimo sed tamen distincto et stigmate orbiculato superata. (v. s. comm. à cl. Wallich.)
6. *B. CRETICA* (Lindl. spec. 472.) spinis 3-5-partitis, foliis ovali-oblongis integris aut subserratis, racemis 3-8-floris foliis ferè brevioribus. 3 in montibus Sphacioticis Cretæ et Delphii Eubææ. — Clus. hist. 2. p. 301. P. Alp. exot. p. 21. t. 20.* — Bacca ovata, nigræ, 2-spermæ potius acerbæ quàm acidæ, stigma brevissimè pedicellatum. DC. syst. 2. p. 9.* (v. v.)
ß. serratifolia, foliis serrato-ciliatis. Poir. dict. 8. p. 618.*
7. *B. CRATÆGINA* (DC. syst. 2. p. 9.*) spinis simplicibus, foliis oblongis nervuloso-reticulatis hinc indè vix subserratis, racemis ∞ -floris confertis patulis folio vix longioribus. 3 in Asiâ minore. Affinis *B. Creticæ* et *Sinensi*. Flores 12-18 conferti, *B. vulgari* similes.
8. *B. THUNBERGII* (DC. syst. 2. p. 9.*) spinis infimis 3-partitis, superioribus simplicibus, foliis ovalibus basi attenuatis integerrimis, racemis paucifloris corymbosis folio brevioribus. 3 in Japoniâ. *B. cretica* Thunb. fl. jap. 1. p. 146.* non Lin. — Bacca juniores oblongæ, stigmate lato orbiculato sessili terminatæ et quasi truncatæ. (v. s. in h. Lamb.)
9. *B. GLAUCA* (DC. syst. 2. p. 10.* H. B. K! nov. sp. am. 5. p. 71. t. 433.) spinis 3-partitis, foliis obovatis subtus albido-glaucis subintegris, racemis ∞ -floris erectis ramoso-paniculatis, petalis obovatis. 5 in Americâ merid. circa Santa-Fé de Bogotâ. — Flores paulò majores quàm in *B. vulgari*. Sepala 6 cum squamis 3 accessoriis majusculis. (v. s. in h. Humb.)
10. *B. MONOSPERMA* (Ruiz et Pav. fl. per. 3. p. 52.*) spinis 3-partitis, foliis ovalibus obovatisve mucronatis, inferioribus dentato-spinosis, racemis ∞ -floris nutantibus. 3 in montibus Peruvix Bacca nigræ, monospermæ. DC. syst. 2. p. 10.*
11. *ß. LATIFOLIA*. (Ruiz et Pav! fl. per. 3. p. 52.* t. 282. a.) spinis brevibus tripartitis simplicibusve, foliis obovatis integerrimis mucronatis subtus pallidis, pedunculis 3-6-floris folio brevioribus. 3 in nemoribus altis frigidis Andium Peruvix. Flores ignoti. Bacca ovales, atræ 3-4-spermæ, stigmate pedicellato superatæ. DC. syst. 2. p. 10. (v. s. sine fl. à cl. Pavon. comm.)
12. *B. FLEXUOSA* (Ruiz et Pav. fl. per. 3. p. 52. t. 282. t. 2.) spinis 3-partitis, foliis obovatis glaucis integerrimis apice mucronatis, racemis aggregatis inæqualibus paucifloris. 3 in rupibus aridis calidis Andium Peruvix Bacca juniores ovato-oblongæ, apice in collum attenuatæ, stigmate orbiculato coronatæ, adultæ oblongæ nigræ 4 5-spermæ. DC. syst. 2. p. 11.* (v. s. in h. Mus. Par. et Juss.)
13. *B. RUSCIFOLIA* (Lam! ill. t. 253. p. 2.) spinis 3-partitis, foliis oblongis basi attenuatis mucronatis integris aut hinc indè grossè dentato-spinosis, pedunculis brevibus apice 4-5-floris. 3 in Americâ australi circa Buenos-Ayres. Flores paulò minores quàm in *B. vulgari*. DC. syst. 2. p. 11. (v. s.)
14. *B. PANICULATA* (Juss! in DC. syst. 2. p. 12.*) spinis brevibus 3-fidis, foliis lanceolatis mucronatis integris aut hinc indè spinuloso-dentatis, pedunculis angulatis erectis racemoso-paniculatis. 5 in Peraviâ. Bractæ lineari-subu-

latæ, pedicellis dimidio brevioribus. Calyx basi 3-bracteatus. (v. s. in h. Juss.)

15. *B. ILICIFOLIA* (Forst! comm. goett. 9. p. 28.*) spinis 3-partitis, foliis ovalibus basi attenuatis hinc inde grossè dentato-spinosis, pedunculis brevibus 4-floris, pedicellis elongatis subcorymbosis, baccis ovatis lagenæformibus. ζ in rupium fissuris terræ del Fuego. *B. lagenaria*. Poir! dict. 8. p. 619.* — *Bacca* ovata, in collum apice attenuata et idèò lagenæformis. DC. syst. 2. p. 12.* (v. s.)
16. *B. ASIATICA* (Roxb! in DC. syst. 2. p. 13.*) spinis 3-fidis simplicibusve, foliis ovalibus mucronatis glabris subtus glaucis, integris aut spinoso-dentatis, racemis brevibus multifloris, pedicellis elongatis, baccis subglobosis. ζ in Indiâ orient. et Napauliâ. — Deless! ic. sel. 2. t. 1. *Baccæ* subglobosæ, stylo crasso brevissimo superatæ, polline cæsius. (v. s.)
17. *B. RIGIDIFOLIA* (H. B. et K! nov. sp. am. 5. p. 70. t. 431.) spinis 3-partitis, foliis oblongis spinoso-mucronatis vix 1-2 dentibus auctis glabris, racemis paucifloris folio vix longioribus. ζ in Americâ merid. Aff. *B. luteæ*. Flores magnitudine *B. vulgaris*. Bracteolæ subulatæ, pedicellis triplò breviores. (v. s. in h. Humb.)
18. *B. QUINDIENSIS* (H. B. et K! nov. sp. am. 5. p. 70. t. 432.) spinis nullis, foliis oblongis spinuloso-dentatis mucronatis glabris, racemis erectis pubescentibus multifloris folio longioribus. ζ in Americâ merid. — Flores duplò majores quàm in *B. vulgaris*. Petala 6. Bractææ linearisubulatæ glabræ, pedicellis dimidio triplòve brevioribus. (v. s. in h. Humb.)
19. *B. LAURINA* (Billb. in Flor. 1821. p. 330.) spinis. . . foliis ovatis mucronatis integris, racemis simplicibus pendulis. ζ in Brasiliâ. \dagger .

§. 2. Foliis simplicibus, pedicellis unifloris.

20. *B. WALLICHIANA*, spinis 3-partitis, foliis oblongis utrinquè acuminatis glaberrimis rigidis ciliato-serratis, pedicellis 1-floris 10-15 aggregatis. ζ in Napauliâ. (v. s. comm. à cl. Wallich)
21. *B. TOMENTOSA* (Ruiz et Pav! fl. per. 3. p. 52.* t. 282. b.) spinis nullis, foliis ovalibus subtus densè breviterque tomentosis integris aut dentibus paucis spinosis, pedicellis 1-3 unifloris gracilibus. ζ in Peruviâ. Flores 6-petali circiter magnitudine *B. vulgaris*. *Baccæ* ovales, stigmate pedicellato rostratæ, 2-3-spermatæ. (v. s. comm. à cl. Pav.)
22. *B. LUTEA* (Ruiz et Pav! fl. per. 3. p. 51. t. 280 *) spinis nullis, foliis ovatis mucronatis et tricuspidatis, ramulis subpubescentibus, pedicellis plurimis unifloris folio subbrevioribus. ζ in frigidis rupestribus sylvaticis Andium Peruviz. DC. syst. 2. p. 14.* (v. s. in h. Deless.)
23. *B. CONFERTA* (H. B. et K! nov. sp. am. 5. p. 69. t. 430.) foliis minoribus aliis integris spinulâ mucronatis aut spinulâ utrinquè unicâ auctis, plurimis spinulas utrinquè 2-3 gerentibus. ζ inter Caxamarcum et fl. Magdalenæ. *B. lutea* ϵ . DC. syst. 2. p. 14. (v. s. in h. Humb.)
24. *B. NERVIS* (Pers! ench. 1. p. 387.) spinis nullis, foliis ellipticis integerrimis vix macronulatis glabris, pedicellis solitariis unifloris folio longioribus. ζ ad fretum Magellanicum. Ovarium ovatum stigmate coronatum. DC. syst. 2. p. 15.* (v. s.)
25. *B. BUXIFOLIA* (Lam! ill. t. 253. f. 3.) spinis 3-partitis, foliis ovatis et ovato-lanceolatis glabris integerrimis, pedicellis folio longioribus aut solitariis unifloris aut ternis è pediculo brevissimo ortis. ζ ad fretum Magellanicum. Pedicellus solitarius uniflorus. DC. syst. 2. p. 15.* (v. s. in h. Mus. Par.)
ℓ. microphylla, Forst! comm. goett. 9. p. 29.* Pedunculi 3, uniflori. (v. s. in h. M. Par.)

26. *B. EMPETRIFOLIA* (Lam! ill. t. 253. f. 4.) spinis 3-partitis, foliis linearibus integerrimis margine revolutis, pedicellis 1-2 unifloris ζ ad fretum Magel-

- lanicum. Pedicelli è ramulis inter folia orti. DC. syst. 2. p. 16.* (v. s.)
27. *B. CUNEATA* (DC. syst. 2. p. 16.*) spinis 3-partitis folio ferè longioribus, foliis obovato cuneatis spinoso-trifidis glabris, pedicellis solitariis 1-floris folio subæqualibus. § in Americâ austr. Bacca obovata, subglobosa, intensè cæruleo-purpurea, stigmatè sessili coronata. (v. s. in h. Banks.)
28. *B. HETEROPHYLLA* (Juss! in Poir! dict. 8. p. 622 *) spinis 3-partitis, foliis ovato-lanceolatis glabris, aliis integris, aliis 3-dentato-pungentibus, pedicellis solitariis 1-floris folio vix longioribus. § ad fretum Magellanicum. Bacca rotundata purpurea subcærulescentes, pisi magnitudine, 4-spermae, stigmatè sessili coronata. DC. syst. 2. p. 16. Hook. ex. fl. t. 14. (v. s. in h. Juss.)
29. *B. VIRGATA*. (Ruiz et Pav! fl. per. 51. t. 281. f. B.*) spinis minimis aut nullis, foliis obovatis integris aut apice spinoso-dentatis glabris, pedicellis solitariis 1-floris foliorum longitudine. § in sylvis Peruviae. Bacca parva, oblongo-ovata; semina 2 fusca. Aff. *B. lutea*. DC. syst. 2. p. 17.* (v. s.)
30. *B. SIBIRICA* (Pall! fl. ross. 2. p. 42. t. 67. itin. app. n. 108. t. P. f. 2. ed. gall. 3. p. 211. t. 13. f. 2.) spinis 3-7 partitis, foliis lanceolato-obovatis ciliato-ser-ratis, pedunculis 1-floris folio brevioribus. § in Sibiria Altaica. Flores sat similes *B. vulgari*. Bacca ex. cl. Pall. obovata, rubra. DC. syst. 2. p. 17.* (v. s.)
- §. 3. *Species non satis notæ, foliis abruptè pinnatis et petiolo apice spinescente distinctæ.*
31. *B. TRAGACANTHOIDES* (DC. syst. 2. p. 18.*) spinis 3-partitis parvulis, foliis in axillâ congestis 1-2-jugis, petiolo apice spinescente, § Secus flumen Cur propè Tiflis. Aff. *B. Cretica* et *Sibirica*. (v. s. in h. Mus. Par.) †.
32. *B. CARAGANEFOLIA* (DC. syst. 2. p. 18.*) foliis primordialibus basi spinas stipulares 2 agentibus bijugis, petiolo in spinam apice producto. § in Chinâ. Foliola bijuga ad apicem petioli infrâ partem spinescentem inserta. — Flores ignoti. (v. s. in h. Banks.)

II. MAHONIA. Nutt. gen. amer. 1. n. 307.

Sepala 6, squamis 3 extus stipata. Petala 6 intus eglandulosa. Stamina ad apicem filamenti utrinquè dente aucta. Bacca 3-g-sperma. — Frutices, foliis impari-pinnatis, foliolis sinuato-dentatis, floribus flavis. DC. syst. 2. p. 18.

1. *M. FASCICULARIS* (DC. syst. 2. p. 19.*) foliis 4-5-jugis cum impari, jugo inferiore ad basim petioli approximato, foliolis ovato-lanceolatis repando-dentatis, dentibus utrinquè 4-5, racemis erectis confertissimis. § in Novâ-Hispaniâ et circa Nutka. Berberis pinnata. Lag. elenc. h. madr. 1803. p. 6. Ker. bot. reg. t. 702. H. B. K. nov. sp. am. 5. p. 71. t. 434. *B. fasciculata* Sims. bot. mag. t. 2396. Deless. ic. sel. 2. t. 3. (v. s.)
2. *M. AQUIFOLIUM* (Nutt. gen. am. 1. p. 212.*) foliis 2-3-jugis cum impari, jugo inferiore à petioli basi distante, foliolis ovato-lanceolatis repando-dentatis, dentibus utrinquè 6-9, racemis erectiusculis sæpius confertis. § in Americâ bor. secus rivum Colombia. Berberis aquifolium Pursh! fl. bor. am. 1. p. 219. t. 4. Bacca atro purpurea, stigmatè 3-lobo coronata. — DC. syst. 2. p. 20.* (v. s. in h. Lamb.)
- β. *Nutkana*, foliolis 2 inferioribus intervallo ultrâ pollicari à petioli basi distantibus, utrinquè 9-dentatis. — circa Nutka. An diversa? (v. s. in h. Banks.)
3. *M. NERVOSA*. (Nutt. gen. amer. 1. p. 212.) foliis 5-6-jugis cum impari, jugo inferiore à petioli basi distante, foliis oblongo-lanceolatis repando-dentatis sub-5-nerviis, dentibus utrinquè 12-14, racemis congestis brevibus. § in Americâ boreal. Berberis nervosa Pursh! fl. bor. am. 1. p. 219. t. 5. — DC. syst. 2. p. 21.* (v. s. in h. Lamb.)

4. *M. GLUMACEA* (DC. syst. 2. p. 21.*) foliis 6-8-jugis cum impari, jugo inferiore à petioli basi longè distante, foliolis oblongo-lanceolatis repando-dentatis. dentibus utrinquè 7-9, racemis 1-2-elongatis gracilibus. ♂ in Americà bor. Flores magnitudine *Berberidis vulgaris*; ovarium ovatum apice attenuatum. (v. s. in h. Banks.)
5. *M. NAPAULENSIS*. (DC. syst. 2. p. 21.*) foliis 6-jugis cum impari, jugo inferiore minore ad petioli basim approximato, foliolis ovato-lanceolatis basi 5-nerviis repando-dentatis, dentibus utrinquè 5-10, racemis paucis elongatis gracilibus. ♂ in Napaniâ. — Deless! ic. sel. 2. t. 4. (v. s. comm. à cl. Wallich.)

β. *Roxburghii*, B. pinnata Roxb! ined. DC. syst. 1. c. — habet foliola dentibus paucioribus majoribus donata, pedicellos bracteis paulò longiores.

†. *Species non satù nota.*

6. *M. JAPONICA* (Thunb. fl. jap. 77? ic. jap. t. 22.) *Berberidis species Brown.* cong. p. 22. in adnot. ♂ culta in insulâ Nipon Japoniæ. Ex descriptione Floræ Japonicæ videtur illex, sed icon citata ad Mahoniam quamdam momentè cl. Brownio pertinere videtur. DC. syst. 2. p. 22. †.

III. *NANDINA*. Thunb. nov. gen. 1. p. 14. Gaert. fruct. 2. p. 69. t. 92. f. 3.

Sepala 6 squamis ∞ multiseriis stipata. Petala 6 intùs eglandulosa. Bacca exsucca globosa, stylo coronata. Semina 2 rotunda hinc convexa, illinc concava. — Frutex foliis decompositis, foliolis integris, floribus paniculatis albis. DC. syst. 2. p. 22.

1. *N. DOMESTICA* (Thunb. diss. nov. gen. pl. 1. p. 14.) ♂ in Japoniâ et Chinâ. Lam. ill. t. 261. Herb. amat. t. 281. DC. syst. 2. p. 22. (v. v.)

IV. *LEONTICE*. Lin. gen. n. 423. Lam. ill. t. 254.

Sepala 6 extùs nuda. Petala 6 intùs basi squammulam gerentia. Capsula vesicaria 2-4-sperma. Semina in fondo capsulæ inserta globosa. — Herbæ radice tñerosâ, foliis variè sectis, calycibus sæpè coloratis. DC. syst. 2. p. 23.

Sect. I. *LEONTOPETALUM*. DC. syst. 2. p. 24.

Capsula per maturitatem valdè inflata nunquàm rupta; semina inclusa. — Folia superiora pinnatim ternatimve secta, petiolo simplici aut apice nec à basi ipsâ partito.

1. *L. CHRYSOGONUM*. (Lin. spec. 447.) foliis pinnatisectis, segmentis sessilibus ovali-oblongis apice 3-5-fidis, bracteis parvis scariosis. ♀ inter segetes Græciæ. — Moris. hist. 2. p. 285. s. 3. t. 15. f. 7. — Stamina petalis ferè æqualia. — DC. syst. 2. p. 24.* (v. s.)
2. *L. LEONTOPETALUM*. (Lin! spec. 448.) foliis biternatim sectis, segmentis subpetiolulatis obovatis, bracteis foliaceis pedicello multoties brevioribus. ♀ in arvis et segetibus Apuliæ, Etruriæ, Cretæ. Lam! dict. 3. p. 465.* ill. t. 254. f. 1. — Semina 3-4, fusca, globosa. DC. syst. 2. p. 25.* (v. s.)
3. *L. VESICARIA*. (Pall. act. petrop. 1779. p. 2. t. 9. f. 4.) foliis biternatim sectis, segmentis oblongo-subcuneatis obtusis submucronatis, bracteis foliaceis pedicello vix dimidio brevioribus. ♀ in præruptis limosis ad lacum salisum Inderiensem deserti Kirghisorum. L. incerta. Pall. itin. 3. app. n. 84. t. 5. f. 2. ed. gall. app. n. 321. t. 77. f. 3.* — DC. syst. 2. p. 25.* (v. s. sine fl. comm. à Cl. Fisch.)

Sect. II. *CAULOPHYLLUM*. Michx. fl. bor. amer. 1. p. 204. t. 21.

Capsula vix inflata, per maturitatem demùm rupta. Semina idèò exserta, interdum baccata. — Folium caulinum unicum sub racemo situm, petiolo ab

ipsâ basi 3-partito, ramulis tribus segmenta 3-5-gerentibus. DC. syst. 2. p. 26.

4. *L. ALTAICA* (Pall act. petrop. 1779. p. 257. t. 8. f. 1. 2. 3.*) folio caulino solitario, petioli à basi 3-partiti ramulis segmenta 5 oblonga integra palmatim disposita gerentibus. \mathcal{L} in apricis montium Altaicorum Sibiriz. Lam. ill. t. 254. f. 2. DC. syst. 2. p. 26.* (v. s.)

β . *Odessana* (Fisch! in litt.), — à priori differt tantum ex sicco pedicellis paulò longioribus, staminibus duplò petala superantibus et segmentis foliorum paulò magis petiolulatis.

5. *L. THALICTROIDES* (Lin! sp. 448.) folio caulino solitario, petioli à basi tripartiti segmentis ovatis apice grossè incisis acuminatis. \mathcal{L} in umbrosis montanis frigidis Americæ borealis. — Caulophyllum thalictroides Mich! fl. bor. am. 1. p. 205. t. 21.* — DC. syst. 2. p. 27.* (v. s.)

V. EPIMEDIUM. *Lin. gen. n. 148.*

Sepala 4 extus basi bibracteolata. Petala 4 intus appendice discoloreiuncta. Capsula siliculæformis oo-sperma. Semina obliquè transversim sita, unilateralia. — Herbæ perennes, foliis radicalibus petiolatis multisectis, foliolis serrato-aristatis. DC. syst. 2. p. 28.

1. *E. ALPINUM* (Lin. spec. 171.) foliis radicalibus nullis, caulino biternatim secto, foliolis cordato-lanceolatis acuminatis serratis, serraturis aristatis. \mathcal{L} in umbrosis et dumetis montanis Angliæ, Galliæ, etc. Lam. ill. t. 83. Schk. bot. Handb. 1. p. 81. t. 24.* DC. syst. 2. p. 28. (v. v.)

β . *pubigerum*, petiolis pilosis, nodis hirsutissimis. Circa Constantinopolim. An species propria. (DC. l. c.*)

2. *E. PINNATUM* (Fisch. in litt. cum icone. DC. syst. 2. p. 29.*) foliis radicalibus pinnatim sectis, scapo aphylo. \mathcal{L} in Persiæ, provinciâ Gilan. Flores erecti, fructus apice pedicelli inflexo-penduli. (descr. ex icon.)

VI. DIPHYLLEIA. *Mich. fl. bor. am. 1. p. 203. t. 19 et 20.*

Sepala 3 extus nuda. Petala 6 intus nuda. Bacca sessilis 2-3-sperma. Semina ovato-oblonga. — Herba glabra habitu *Leontices* aut *Podophylli*. DC. syst. 2. p. 30.

1. *D. CYMOsa*. (Michx! l. c.) \mathcal{L} ad rivulos excelsorum montium Carolinæ septentrionalis. — Folia 2 alterna., ampla subreniformia, sæpius apice profundè biloba. Flores albi. Baccæ cæruleo-nigræ subrotundæ. — DC. syst. 2. p. 30.* (v. s. in h. Mus. Par.)

Ord. VII. PODOPHYLLACEÆ. DC. *syst.* 2. p. 31.

Calyx 3-4 sepalus, sepalis deciduis aut persistentibus, sæpè petaloïdeis. Petala simplici duplici triplicive serie disposita, serie quâque sepalorum numero æquali, petalis seriei externæ cum sepalis alternantibus. Stamina hypogyna, petalis numero æqualia et tunc iis opposita, aut dupla tripla multipliciave, seriebus nempè binis ternis plurimisve disposita; filamenta filiformia; antheræ ovales linearesve, terminales, introrsæ, rimâ duplici sæpius longitudinali dehiscentes. Torus non ampliatus et nullo modo carpella involvens. Ovaria plurima aut duo aut (abortu ?) solitaria; stylus subnullus; stigma crassum, subpeltatum. Carpella tot quot ovaria, 1-locularia, nunc subcarnosa indehiscencia, nunc sub apice circumscissè dehiscencia. Semina ∞ aut pauca, imò abortu solitaria, ovato-globosa, placentæ laterali adfixa, inversa. Albumen carnosum. Embryo parvus, ad basin seminis. — Herbæ nunc humidorum umbrosorumque locorum incolæ nunc verè aquaticæ. — Ordo adhuc dubius, paucissimas species amplexans, hinc Ranunculaceis et Berberideis, illinc Papaveraceis et Nymphæaceis affinis.

Trib. I. *PODOPHYLLEÆ*. DC. *Syst.* 2. p. 32.

Stamina numero petalorum dupla. Ovarium unicum. Stigma crassum subpeltatum. Semina plurima. — Herbæ erectæ nec in aquâ natantes.

I. *PODOPHYLLUM*. *Lin. gen. n.* 646. *Lam. ill. t.* 449.

Calyx 3-sepalus. Petala 6-9. Stamina 12 18. Bacca subcarnosa 1-locularis. — Herba perennis, foliis peltato-lobatis oppositis, floribus solitariis albis, habitu Meconopsidis. DC. *syst.* 2. p. 34.

1. *P. PELTATUM* (Lin! spec. 722.) caule erecto bifolio unifloro, fructibus ovatis. \mathcal{L} in Americâ sept. à Novâ-Angliâ ad Carolinam. Bacca ovata magnitudine Pruni domesticæ stigmate persistente coronata. DC. *syst.* 2. p. 34.* (v. s. et v.)
2. *P. CALLICARPUM* (Raf. fl. lud. p. 14. n. 20.) caule furcato, fructibus oblongis. \mathcal{L} in Louisianâ. fructus oblongus avellanæ majoris magnitudine, albus subroseus. DC. *syst.* 2. p. 34. †.

II. *JEFFERSONIA*. *Bart. act. soc. am.* 3. p. 334.

Calyx 4-sepalus. Petala 8. Capsula circumscissè dehiscens. Semina ∞ arillo demum lacero ad basin instructa. — Herba acaulis hinc Podophyllo, illinc Sanguinariæ affinis.

1. *J. DIPHYLLA* (Pers. ench. 1. p. 418.) \mathcal{L} in Tennessee occidentali et in Virginia. Sims. bot. mag. t. 1513.* — Flos albus; calyx deciduus coloratus. Semina lucida, oblonga, spadicea. DC. *syst.* 2. p. 35.* (v. s. in h. Mus. Par. Deless. et Lamb.)

III. ACHLYS. DC. syst. 2. p. 35. †.

Calyx. . . . Petala ∞ . Stamina filamenta apice dilatata, antheræ loculi subglobosi distantes transversè rimâ dehiscentes. — Genus non satis notum. An *Jeffersoniæ*, an *Acteæ* potius affine?

1. *A. TRIPHYLLA* (DC. syst. 2. p. 35. *) 2. in orâ occidentali Americæ borealis. *Leontice triphylla* Smith! in Rees cycl. n. 5. * folium unicum radicale. flores albi parvi. (v. s. in h. Smith.) †.

Trib. II. *HYDROPELTIDÆ*. DC. syst. 2. p. 36. — *Cabomba* Rich. anal. fr.

Ovaria 2 aut ∞ . Semina pauca aut abortu solitaria. — Herbæ natantes. — An potiùs Tribus prima inter *Nymphæaceas*?

- IV. *CABOMBA*. *Aubl. Guian.* 1. p. 321. t. 124. *Rich. Ann. Mus. Par.* 17. p. 230. t. 5. f. 23. — *Nectris Schreb. gen.* n. 610.

Calyx 3-sepalus. Petala 5. Stamina 6. Ovaria 2. Semina plurima globosa inversa. — Herba aquatica, Ranunculum aquatilem habitu æmulans. DC. syst. 2. p. 36.

1. *C. AQUATICA* (Aubl. l. c. *) 2. in stagnis et rivulis Cayennæ et Guyanæ. Pedunculi axillares, 1-flori, longi, solitarii. Flores parvi, latei. DC. syst. 2. p. 36. * (v. s.)

- V. *HYDROPELTIS*. *Michx. fl. bor. amer.* 1. p. 324. t. 29. *Sims. bot. mag.* t. 1147.

Calyx 3-4-sepalus. Petala 3-4. Stamina 18-36. Ovaria 6-18. Semina intrâ pericarpium pendulum ovato-globosum. — Herba aquatica undiquè gelatinâ viscosissimâ lubrica, radicibus fibrosis humo adfixa.

1. *H. PURPUREA* (Michx! fl. bor. amer. 1. p. 324: t. 29.) 2. in Carolinâ. *Brasenia peltata* Pursh. fl. bor. amer. 2. p. 389. — DC. syst. 2. p. 38. (v. s.)

Ordo VIII. NYMPHÆACEÆ. DC. *prop. med. ed.* 2. p. 119.
syst. 2. p. 39.

Torus floris expansus nunc in conum invenum ovaria in eo semi-immersa gerentem, nunc in urceolum ovaria includentem et stigmatibus coronatum. Calyx 4-6-sepalus, sæpius 4-sepalus, sepalis toro insertis, super eum non articulatis, sæpè persistentibus, intus coloratis. Petala seriebus plurimis disposita, toro nunc demissiùs nunc altiùs inserta, oblonga, plana, serie quaque totidem petalis constante quot sunt sepala, petalis seriei externæ sepalis et cæteris inter se alternis. Stamina ∞ , multiserialia, toro exactè ut petala, sed paulò altiùs inserta; filamenta planiuscula, interdum ultrà antheræ loculos producta; antheræ adnatæ, introrsæ, lineares, biloculares, rimâ longitudinali duplici dehiscentes. Ovaria seu carpella 8-24, nunc toro ampliato supernè foveolato semi-immersa, nunc intrâ torum valdè ampliatus et urceolæformem inclusa, monostyla, membranacea, evolvija, 1-2 aut ∞ -sperma; Styli in carpellis liberis distincti stigmate simplici coronati, in carpellis inclusis connati stigmatibus suprâ urceolum peltatim (exactè ut in Papavere) radiatis basi connatis apice liberis. Semina in carpellis liberis verosimiliter 2 et abortu solitaria, in carpellis inclusis innumera, parietibus carpellorum laterilibus adfixa, inversa, ovato-globosa, punctulata, arillo folliculæformi subgelatinoso cincta, loculis pulpâ gelatinosâ ad maturitatem farctis. Albumen nunc nullum? in seminibus solitariis, nunc farinaceum in seminibus carpellorum polyspermorum. Embryo parvus, turbinato-globosus, extrâ albumen ad basin seminis situs, ideò in fructu inversus primâ fronte indivisus, inclusus nempe in sacculo seu membranâ propriâ et intus cotyledones 2 foliaceas exhibens, interdum antè seminum delapsus germinans. — Herbæ aquarum tranquillarum aut lentè fluentium incolæ.

Trib. I. *NELUMBONEÆ*. DC. *syst.* 2. p. 43.

Carpella plurima distincta 1-2-sperma monostyla toro elevato obconico supernè profundè foveolato immersa. Semina in quoque carpello solitaria exarillata exalbuminosa.

1. *NELUMBIUM*. Juss. *gen.* 68. — *Nelumbo*. Tourn. *inst.* 261. Gaert. *fruct.* 1. p. 73. t. 19.

Car. gen. est idem ac trib. — Herbæ nymphæas habitu omninò æmulantes. — Flores ampli, albî rosei aut flavi.

1. *N. speciosum* (Willd. *spec.* 2. p. 1258. var. α β .) corolla polypetala, antheris ultra loculos in appendicem clavatam productis. \mathcal{Y} in aquis lentè fluentibus cænosis et stagnosis Asiæ calidioris. Sims. *bot. mag.* t. 903. *Lam.*

ill. t. 453. *Nymphæa Nelumbo* var. *α*. Lin. sp. 730. *Nelumbo nucifera*. Gaertn. fruct. 1. p. 73. t. 19. f. 2. — Flores pulcherrimi, anisum spirantes. DC. syst. 2. p. 44. (v. s.)

β. *Tamara*, staminibus exterioribus sterilibus apice dilatatis alatis obcordatis, appendice ex emarginaturâ ortâ. — *Tamara Rheed*. mal. xi. p. 59. t. 30.* in Malabarîâ.

γ? *Caspicum*, petalis interioribus vix externis minoribus, obtusis. — ad ostium fluminis Volgæ propè Astracan. N. *Caspicum* Fisch! in litt.

2. *N. LUTRUM* (Willd. spec. 2. p. 1259.) corolla polypetala, antheris ultra loculos in appendicem linearem productis. 2 in lacubus et fossis stagnosis Americæ bor. Turp. ann. mus. 7. p. 210. t. 11. f. 27. *Nymphæa Nelumbo* var. β. Lin. spec. 730. — Valdè simile *N. specioso*. — DC. syst. 2. p. 47.* (v. s.)

Species non satis notæ.

3. *N. CODOPHYLLUM* (Raf. fl. lud. p. 22. n. 64.) foliis subtus tomentosis. 2 in lacubus Louisianæ. Napoleone Rob. voy. louis. 1. p. 355. 2. p. 322. et 441. ex Raf. — Flores lutei, majores quàm *N. lutei*. folia diametro bipedali. DC. syst. 2. p. 47.*

4. *N. PENTAPETALUM* (Willd. spec. 2. p. 1259.) floribus pentapetalis. 2 in Carolinâ. *Nymphæa pentapetala* Walt. fl. car. 155. Poir. dict. 4. p. 458. — Flores albi. — DC. syst. 2. p. 47.

5. *N. JAMAÏCENSE* (DC. syst. 2. p. 47.) . . . in Jamaicâ — *Nymphæa* foliis orbiculatis peltatis subtus radiatis, fructu obversè conico seminibus majoribus nidulantibus. Brown. jam. 343. n. 2. *N. speciosum* γ. Willd. spec. 2. p. 1259.

Trib. II. *NYMPHÆEÆ*. DC. syst. 2. p. 48.

Carpella plurima polysperma intra torum ampliatus inclusa, stigmatibus supra fructum baccæformem radiantibus. Semina ad parietes laterales carpellorum adfixa, arillata. Albumen farinosum.

II. *EURYALE*. Salisb. ann. bot. 2. p. 73.

Sepala petala staminaque cum toro carpella tegente longè adnata et ideò bacca pseudo-infera. — Herba aquatica, tota ferè aculeis horrida.

1. *E. FEROX* (Salisb. l. c.) 2 in Indiâ orient. *Anneslea spinosa* Andr. bot. rep. t. 618.* — Petioli et calyces præsertim aculeis rigidis hispidi. Folia ampla acutata; semina magnitudine Pisi. DC. syst. 2. p. 48.* (v. s.)

III. *NYMPHÆA*. Neck. elem. n. 1828. DC. syst. 2. p. 49.

Sepala ad basin tori; petala staminaque cum toro carpella tegente longè adnata et bacca ideò quasi semi-infera cicatrisata. — Flores ampli, albi, rosei, rubri aut cærulei, nunquàm lutei.

Sect. I. *CYANÆA*. DC. syst. 2. p. 49.

Antheræ apice productæ. Flores cærulescentes aut cærulei. Folia peltata, integra aut obtusè sinuata. — An stirpes hujus sectionis ut species aut ut varietates habendæ, dubito?

1. *N. SCUTIFOLIA* (DC. syst. 2. p. 50.*) foliis peltatis obtusè sinuato-dentatis impunctatis utrinquè glabris, basi bilobis, lobis incumbibus, antheris apice appendiculatis, stigmate 20-radiato. 2 ad. prom. Bonæ Spei. — *N. cærulea* Andr. bot. rep. t. 197. Sims-bot. mag. t. 552. — Differre videtur a sequenti floribus intensius cæruleis, petalis sepalisque obtusioribus. (v. s.)

2. *N. CÆRULEA* (Savigny! decad. egypt. 3. p. 74.) foliis peltatis subintegrè impunctatis utrinquè glabris basi bilobis, lobis liberis, antheris apice appendiculatis, stigmate 16-radiato. 2 in oryztis et canalibus Ægypti inferioris ann. Mus. Par. 1. p. 366. t. 25.* Vent! malm. t. 6. herb. amat. t. 338. DC. syst. 2. p. 50.* (v. v. et s.)

3. *N. MADAGASCARIENSIS* (DC. syst. 2. p. 50.*) foliis peltatis obtusè sinuatis impunctatis utrinquè glabris basi bilobis, lobis divaricatis, antheris apice appendiculatis, stigmate 8-10-radiato. 2 in insulâ Madagascar. — Differt à minoribus Ægypti speciminibus præsertim foliorum lobis magnis divaricatis. — Potius sequentis varietas. (v. s.)

4. *N. STELLATA* (Willd. spec. 2. p. 1153.) foliis peltatis integris impunctatis utrinque glabris basi bilobis, lobis divaricatis, antheris apice appendiculatis, stigmate 8-12-radiato. \mathcal{L} in rivulis lacubusve stagnosis Malabaricæ. Andr. bot. rep. t. 330. — Citambel Rheed. h. malab. 11. p. 53. t. 27. DC. syst. 2. p. 51.* (v. s. *imb.* Deless.)
5. *N. PULCHRELLA* (DC. syst. 2. p. 51.*) foliis peltatis subintegris impunctatis utrinque glabris basi bilobis, lobis subdivergentibus obtusis, antheris apice appendiculatis, stigmate 12-radiato. \mathcal{L} in Peruvia. Flores albi è sicco, dimidiò minores ac *N. albâ*. Petala 8 oblonga, acuminata, calyces breviora. (v. s. in h. Lamb.)

Sect. II. *LOTOS*. DC. syst. 2. p. 52.

Antheræ apice in appendicem non productæ. Flores non cærulei sed albi rosei aut rubri. Folia peltata, rarius integra, sæpius argutè dentata aut subtus pubescentia.

6. *N. EDULIS* (DC. syst. 2. p. 52.) foliis peltatis latè ovalibus integerrimis subtus pubescentibus. \mathcal{L} in paludosis Indiæ orient. Castalia edulis Salisb. ann. bot. 2. p. 73. parad. n. 14. in adnot. — Flores parvi.
7. *N. KURRA* (Roxb! ined. Sims. bot. mag. t. 280.) foliis peltatis argutè dentatis subtus pubescentibus immaculatisque. \mathcal{L} in Indiâ orient. — Castalia magnifica Salisb. parad. t. 14.* — DC. syst. 1. p. 52. (v. s. in h. Lamb.)
Æ. roea, floribus roseis foliis supernè virentibus. Sims bot. mag. t. 1364.
8. *N. PUBESCENS* (Willd. spec. 2. p. 1154.*) foliis peltatis argutè dentatis orbiculato-reniformibus subtus velutino-pubescentibus maculatisque, lobis rotundis. \mathcal{L} in Indiâ orient. et in Africâ. *N. Lotus* Burm! ind. 119. Beauv! fl. Owar. 2. p. 50. t. 83. — Species diù confusa cum *N. Loto* sed diversa. — DC. syst. 2. p. 52.* (v. s.)
9. *N. LOTUS* (Lin. spec. 729. excl. syn. Br. et Sl.) foliis peltatis argutè serratis subtus in nervis pilosiusculis inter nervos pubescentibus. \mathcal{L} in Nilo et aquis lentè fluentibus Ægypti. Delile! fl. Ægypt. ill. p. 16. descr. p. 159.* t. 60. f. 1. opt. — Castalia mystica Salisb. ann. bot. 2. p. 73. — DC. syst. 2. p. 53.* (v. s.)
10. *N. THERMALIS* (DC. syst. 2. p. 54.*) foliis peltatis argutè dentatis utrinque glaberrimis, auriculis approximatis. \mathcal{L} in Hungariâ. *N. Lotus*. Waldst. et Kit! pl. rar. hung. 1. p. 13. t. 15.* Sims bot. mag. t. 797.* Castalia mystica Salisb. parad. n. 14. in adnot. Flores albi, odore subalcoólico donati. (v. s.)
11. *N. AMPLA* (DC. syst. 2. p. 54.*) foliis peltatis argutè dentatis utrinque glaberrimis, nervis subtus valdè prominentibus areolatis reticulatis. \mathcal{L} in aquis Jamaicæ, Sancti-Domingi. Castalia ampla. Salisb. ann. bot. 2. p. 73. parad. n. 14. in adnot. (v. s. var. α sine fl. in h. Banks et Deless.)
Æ. Rudgeana. Mey. prim. fl. esseq. p. 198. ex syn. Distinguitur a *N. Loto* foliis supernè minutè tuberculatis.
12. *N. VERSICOLOR* (Roxb. hort. beng. p. 41.) foliis peltatis margine et intrâ fissuram sinuato-dentatis punctato-pustulatis utrinque glabris. \mathcal{L} in Bengalo. Sims. bot. mag. t. 1189.* Flos albus aut carneus. — DC. syst. 2. p. 55.* (v. s. in h. Lamb.)

Sect. III. *CASTALIA*. DC. syst. 2. p. 55.

Antheræ apice non productæ. Flores albi. Folia cordata nec peltata, glabra, integerrima.

13. *N. RENIFORMIS* (Walt. fl. car. 155.) foliis reniformibus, stigmate 16-radiato. \mathcal{L} in Carolinâ. Nelumbium reniforme Willd. spec. 2. p. 1260. Flores albi. *N. albâ* ferè majores. — DC. syst. 2. p. 55.* Deless! ic. sel. 2. t. 5. (v. s.)
14. *N. ALBA* (Lin. spec. 729.) foliis cordatis integerrimis stigmate 16-radiato, radiis adscendentibus. \mathcal{L} in fossis lacubus fluminibusque lentè fluentibus Europæ. fl. dan. t. 602. Schkuhr handb. 2. t. 142.* Semina horizontalia seu deflexa obovata rubra. DC. syst. 2. p. 56.* (v. v.)
Æ. minor. Besl. hort. cystt. vern. ord. vii. t. 3. f. 1. Weinm. phyt. 3. p. 456. t. 761. f. c. — in Alsatiâ circâ Argentinam.

15. *N. ODORATA* (Ait! h. kew. ed. 1. v. 2. p. 227. ed. 2. v. 3. p. 292.) foliis cordatis integerrimis subtus nervis venisque prominentibus, stigmatibus 16-20-radiato, radiis erectis apice inflexis. \mathcal{L} in Americâ boreal. Sims. bot. mag. t. 819. *N. alba* Walt. fl. car. 155. — Affinis *N. Albæ* et cum eâ à plurimis autoribus confusa, sed certè diversa. DC. syst. 2. p. 57.* (v. s. in h. Mus. Par.)
16. *N. MINOR* (DC. syst. 2. p. 58.*) foliis cordatis integerrimis, subtus nervis venisque prominentibus, pedunculis petiolisque pilosiusculis, stigmatibus 16-20-radiato. \mathcal{L} in Americâ boreali. — *N. odorata* β . *minor*. Sims. bot. mag. t. 1652.
17. *N. MITIDA* (Sims. bot. mag. t. 1359.*) foliis cordatis integerrimis, nervis lateralibus subtus æquatis, petiolis lævibus, petalis obtusis, stigmatibus 12-20-radiato. \mathcal{L} in Sibiria. *N. odorata* Willd. h. berol. t. 39. Affinis *N. odorata* et *albæ*. — DC. syst. 2. p. 58.* (v. s.)
18. *N. PYGMÆA* (Ait. h. kew. ed. 2. v. 3. p. 293.) foliis cordatis integerrimis, nervis lateralibus subtus æquatis, petiolis lævibus, petalis acutis, stigmatibus 8-radiato. \mathcal{L} in Chinâ et Sibiria orient. — Sims. bot. mag. t. 1525.* flores odorem ferè *Pollanthis tuberosæ* spirantes. — DC. syst. 2. p. 59.* (v. s. in h. Lamb.)
19. *N. BLANDA* (Meyer. prim. fl. esseq. p. 201.*) foliis cordatis integerrimis, nervis subtus canaliculatis, petalis 16-acuminatis. \mathcal{L} in Americâ calid. propè Essequibo. — *N. glandulifera*. Rodsch. obs. p. 76. ex Meyer. Aff. *N. odorata*, sed distincta nervis canaliculatis. DC. syst. 2. p. 59.*
20. *N. ACUTILOBA*, foliis ovatis subdenticulatis basi profundè cordatis, lobis acutissimis, petalis 16-acutis, stigmatibus 16-radiato. \mathcal{L} in Chinâ. Braam. ie. Chin. t. 18. Character ex icone. †
- IV. *NUPHAR*. *Sibth. et Smith. prodr. fl. græc. 1. p. 361. DC. syst. 2. p. 59.*

Sepala, petala staminaque ad basin tori inserta et ideò bacca quasi supera lævis. — Herbæ aquaticæ; flores lutei.

1. *N. LUTEA* (Smith. prodr. fl. græc. 1. p. 361.) calyce 5-sepalo, stigmatibus integro 16-20-radiato profundè umbilicato, foliis ovali-cordatis, lobis approximatis, petiolis triquetris acutangulis. \mathcal{L} in fossis et fluminibus lentè fluentibus Europæ. *Nymphæa lutea* Lin. spec. 729. fl. dan. t. 603. Engl. bot. t. 159. Sepala 5 obtusissima. Petala multò minora, truncata. DC. syst. 2. p. 60.* (v. v.)
2. *N. PUMILA* (Smith. engl. bot. t. 2292.) calyce 5-sepalo, stigmatibus lobato 10-radiato, foliis oblongo-cordatis punctatis subpubescentibus, lobis approximatis, petiolis basi semi-teretibus apice subtriquetris. \mathcal{L} in fossis et rivulis lentè fluentibus ducatùs Megapolitani etiamque in lacubus montium Scotiæ et Vogesorum. — *Nymphæa pumila* Hoffm. fl. germ. 1800. p. 241. — Dimidiò minor quàm *N. lutea*. DC. syst. 2. p. 60.* (v. s. comm. à cl. Flörke.)
3. *N. KALMIANA* (Ait. h. kew. ed. 2. v. 3. p. 295.) calyce 5-sepalo, stigmatibus inciso 8-10-radiato, foliis cordatis subemersis, lobis subapproximatis, petiolis subteretibus. \mathcal{L} in fossis Americæ bor. — *Nymphæa Kalmiana*. Sims. bot. mag. t. 1243.* — DC. syst. 2. p. 61.* (v. s. in h. Lamb.)
4. *N. JAPONICA* (DC. syst. 2. p. 62.*) calyce 5-sepalo, stigmatibus 15-16-radiato, foliis oblongo-sagittatis cordatis, auriculis acutis subdistantibus, petiolis teretiusculis. \mathcal{L} hinc inde in aquis Japoniæ. — *Nymphæa lutea* Thunb! fl. jap. 223. excl. syn. — Media ferè inter *N. luteam* et sagittæfoliam. Deless! ic. sel. 2. t. 6. (v. s. in h. Deless.)
5. *N. SAGITTÆFOLIA* (Pursh. fl. bor. am. 2. p. 370.) calyce 6-sepalo, petalis nullis, antheris subsessilibus, foliis sagittato-cordatis oblongatis obtusis. \mathcal{L} in Carolinâ. — *Nymphæa sagittæfolia* Walt. fl. car. 154. Flores parvi lutei. DC. syst. 2. p. 62.* (v. s. in h. Mus. par.)
6. *N. ADVENA* (Ait. h. kew. ed. 2. vol. 3. p. 295.) calyce 6-sepalo, petalis plurimis parvis, pericarpio sulcato, foliis erectis cordatis, lobis divaricatis. \mathcal{L} in fossis et lacubus aquæ dulcis et etiam aquæ salis in Americâ bor. — *Nymphæa advena*. Ait. h. kew. ed. 1. vol. 2. p. 226.* — DC. syst. 2. p. 63. (v. s.)

Ordo IX. PAPAVERACEÆ. DC. *syst.* 2. p. 67. — Papaveracearum gen. *Juss. gen.* 236.

Calyx 2-sepalus, foliaceus, deciduus, sepalis concavis, florem juniorem sæpius includentibus. Petala hypogyna, regularia, eglandulosa, antè evolutionem irregulariter plicata, sæpius 4 unicâ serie regulariter disposita et inter se cruciata, 2 paulò exteriora, 2 interiora, rariùs duplici triplicive serie 8-12, interdum nulla. Stamina hypogyna, inter se non coalita, rariùs serie unicâ 4 petalis opposita, sæpius seriebus duplicatis multiplicatisve 8, 12, 16, 20, 24, et usquè circiter 100, filamentis filiformibus, antheris bilocularibus basi insertis sulco duplici longitudinaliter dehiscentibus. Ovarium 1, liberum, ovatum seu oblongum, constans carpellis ∞ -2 coalitis intrà thalami productionem membranaceam sæpè amplexis. Stylus brevis aut sæpius 0. Stigmata ∞ -2 patentia, sæpè super ovarii apicem sessilia, stellatim disposita. Capsula ovata aut elongata constans carpellis ∞ -2 (in hoc casu Siliqua sæpè dicitur), marginibus seminiferis inter se coalitis, et placentas idèò utrinquè seminiferas nunc bifiliformes nunc septiformes conficientibus, dorsis semina non gerentibus valvatim nunc à basi nunc ab apice dehiscentibus, apicibus carpellorum in stylum aut in stigmata persistentia desinentibus. Semina ∞ (rarissimè et verosimiliter abortu solitaria), placentis intervalvularibus (seu ex placentis carpellorum duorum vicinorum adglutinati factis) inserta, horizontalia, subglobosa, arillo vero destituta. Albumen carnosio-oleosum. Embryo minimus, ad basin albuminis, rectus, dicotyledoneus, cotyledonibus ovato-oblongis hinc planis illinc convexiusculis. — Herbæ annuæ aut perennes, seu suffruticosæ, succo lacteo albo, rariùs flavo croceo seu sanguineo, fœtæ. Radices fibrosæ. Folia alterna, simplicia, penninervia, dentata aut pinnati-lobata. Pedunculi elongati 1-flori. Flores albi flavi aut rubri, nunquàm cærulei.

I. PAPAVER. *Tourn. inst.* 237. t. 119. *Lin. gen. n.* 648. *Gaertn. fruct.* 1. p. 289. t. 60.

Sepala 2, convexa, decidua. Petala 4. Stamina ∞ . Stylus 0. Stigmata 4-20, radiantia, sessilia super discum ovarium coronantem. Capsula obovata, 1-locularis, è carpellis 4-20 in thalami productione membranaceâ inclusis constans, sub stigmatum coronâ valvulis brevibus dehiscens. Placentæ intervalvulares intus in dissepimenta incompleta productæ. — Herbæ perennes succo albo fœtæ. Pedunculi antè florescentiam apice inflexi. DC. *syst.* 2. p. 69.

§. 1. *Capsulis hispidis.*

1. P. NUDICAULE. (*Lin! spec.* 725.) capsulâ hispida obovato-oblongâ, sepalis

setosis, pedunculis radicalibus longissimis, foliis pinnatilobatis, lobis dentatis incisive acutis. \mathcal{L} in Sibiria orient. Sims. bot. mag. t. 1633. — Dill. h. elth. 302. t. 224. f. 291.* — Petala flava rarius albida. DC. syst. 2. p. 70.* (v. v.)

β . *glabratum*, in Sibiria orient. — Ferè glabra in foliis et scapo.

γ . *radicatum*, P. nudicaule OEd. fl. dan. t. 41. opt. P. *radicatum* Rottb. ex Horn. in litt. — in Norvegia. — Minor, hirsutissima, habitu subdi-tincta sed caracteribus non segreganda. (v. s.)

δ ? *rubro-aurantiacum*, (Fisch! in litt. Bot. mag. t. 2344.) in Dahuria. An species propria? Distinguitur non tantum flore rubro-aurantiaco, sed etiam scapi pilis adpressis foliorumque lobulis seta terminatis. (v. s.)

2. P. MICROCARPUM (DC. syst. 2. p. 71.*) capsulâ hispidâ obovato-globosâ, sepalis pilosis, pedunculis radicalibus, foliis pinnati-lobatis, lobis incis dentatisve subacutis. \mathcal{L} in Kamschatkâ. — Species media inter P. nudicaule et Pyrenaicum, nondum satis ritè cognita, sed ut videtur ab utroque capsulâ parvâ obovato-globosâ distincta. (v. s.)

3. P. PYRENAÏCUM (Willd! enum. 563.) capsulâ hispidâ obovatâ, sepalis setosis, pedunculis radicalibus, foliis pilosis pinnatilobatis, lobis incis dentatisve subobtusis. \mathcal{L} in montanis apricis calcareis Europæ australis. DC. syst. 2. p. 71*.

α . *luteum* (DC. syst. 2. p. 71.) Argemone Pyrenaica. Lin. spec. 728. ex h. Banks et ex Willd. P. *aurantiacum* Lois! not. 84.* P. *suaveolens* Lapeyr. suppl. 72.* — Barr. ic. t. 764. (v. v.)

β . *punicum* (DC. syst. 2. p. 72.) P. Pyrenaicum Willd. enum. 563. P. Alpinum B. Lapeyr. abr. 296. in Pyrenæis loco dicto *Port-de-Plan*.

4. P. ALPINUM (Lin! spec. 725.) capsulâ hispidâ obovato-oblongâ, sepalis pilosiusculis, pedunculis radicalibus, foliis glabriusculis bipinnatisectis, lobulis tenuibus subacutis. \mathcal{L} in apricis inter lapides et saxa Alpinum editiorum. Jacq! fl. aust. t. 83. P. *Bursari* Crantz austr. 2. p. 138. t. 6. f. 4.* — DC. syst. 2. p. 72.* (v. v.)

5. P. HYBRIDUM (Lin! spec. 725.) capsulâ hispidâ obovato-globosâ torosâ, sepalis pilosis, caule folioso multifloro, foliis pinnatisectis, lobis multifidis linearibus. ① in agris versuris et oleraceis Europæ occid. et austr. Smith. engl. bot. t. 43. DC. syst. 2. p. 73.* (v. v.)

6. P. ARGEMONÆ (Lin! spec. 725.) capsulâ hispidâ elongato-clavatâ, sepalis glabriusculis, caule folioso multifloro, foliis bipinnatisectis, lobis linearibus. ① in campis arenosis, agris oleraceis Europæ. Smith. engl. bot. t. 643. Petala pallidè coccinea, ad basin maculâ nigrâ notata. DC. syst. 2. p. 74.* (v. v.)

β . caule unifloro. P. *maritimum* With. brit. 486.

§. 2. *Capsulis glabris (aut apice tantum parçè pilosis).*

7. P. DUBIUM (Lin! spec. 726.) capsulâ glabrâ oblongo-clavatâ, sepalis pilosis, caule multifloro setis patentibus hispido, setis pedunculorum adpressis, foliis pinnatipartitis, lobis inciso-dentatis. ① in arvis sabulosis oleraceisque Europæ. Smith. engl. bot. t. 644. Schkuhr handb. 2. p. 69. t. 140. Variat florè intensè coccinea pallido aut albo, ungue concolore aut nigricante. DC. syst. 2. p. 75.* (v. v.)

β . flore albo. Balb. fl. taur. 85. P. *dubium* Jacq. fl. austr. 1. p. 17. t. 25.*

8. P. PAUCIFLORUM (Desf! fl. atl. 1. p. 407.) capsulâ glabrâ obovatâ, sepalis pilosis, caule paucifloro setis patentibus subhispido, foliis bipinnatipartitis, lobis ovatis. ① in Athlante, propè Belide. Aff. P. *dubio* sed capsula dimidiò ferè brevior. Petala rosea magnitudine P. Argemonæ DC. syst. 2. p. 75.* (v. s.)

9. P. RHEAS (Lin! spec. 726.) capsulâ glabrâ obovatâ, sepalis pilosis, caule multifloro setis patentibus scabro, foliis pinnatipartitis, lobis elongatis inciso-dentatis acutis. ① frequentissimum inter segetes. Smith. engl. bot. t. 645.

Variat petalis coccineis absque maculis, carneis, albis, coccineis albo marginatis vel albis coccineo marginatis. DC. syst. 2. p. 77.* (v. v.)

10. *P. trilobum* (Spreng! fl. hal. suppl.) capsulâ glabrâ subrotundâ, sepalis parçè pilosis, caule multifloro glabro, foliis basi cuneatis apice trilobis. (I) in agro Halensi. Aff. *P. Rhæadi* sed distinctum videtur. Flores parvi rubri. DC. syst. 2. p. 78.* (v. s.)
11. *P. lævigatum* (Bieb. suppl. p. 364.*) capsulis glabris obovatis, sepalis glabris, caule uni vel paucifloro subglabro, foliis profundè pinnatifidis, lobis integris linearibus acutis. (I) in collibus circa Odessam. Flores parvi, rubri, magnitudine *P. dubii*. DC. syst. 2. p. 78.* (v. s.)
12. *P. Arenarium* (Bieb. suppl. p. 364.) capsulis oblongis glabris, sepalis hispidis, foliis bipinnatifidis, lacinulis linearibus, setis caulibus multiflori patulis, pedunculorum adpressis, fructiferorum raris caducis. (I) in campis arenosis ad fl. Tereti. Corolla rubra fundo atro.
13. *P. Roublæi* (Vig! diss. 39. n. 4. t. 1. f. 1. opt.) capsulis glabris subrotundis, sepalis pilosis, caule multifloro piloso, foliis bipinnati-partitis villosis, lobis linearibus pilo terminatis. (I) in arenosis Frontignani propè Monspelium. Aff. *P. Rhæadi* sed foliis bipinnatipartitis et habitu distincta. DC. syst. 2. p. 78.* (v. s.)
14. *P. floribundum* (Desf. Choix cor. 62. t. 46.*) capsulis glabris oblongis, sepalis pilosis, caule multifloro hispido, foliis subpilosis, imis pinnatisectis, superis pinnatifidis, lobis incis, lobulis setâ terminatis. (I) in Armeniâ. *P. virgatum* Smith. in Rees cycl. n. 9.* Planta tota pilis rigidis patulis setiformibus subflavescentibus hispida. DC. syst. 2. p. 79.* (v. s. in h. Mus. Par.)
15. *P. horridum* (DC. syst. 2. p. 79.*) capsulis glabris ellipsoideis, sepalis hispidis, caule paucifloro setis rigidis hispidissimo, foliis subamplexicaulibus glaucis sinuato-pinnatifidis lobulis apice et in nervis setigeris. (I) in Novâ-Hollandiâ. Aff. *P. setigero*. Petala rosea (v. s.)
16. *P. Gariepinum* (Burch! cat. geogr. pl. afr. austr. n. 1633.) capsulis glabris obovato-oblongis, sepalis sinuatis, caule multifloro setis innumeris horrido, foliis sessilibus hispidis sinuato-pinnatifidis, lobis distantibus ovatis. (I) in Africâ austr. Aff. *P. horrido*. Petala coccineo-aurantiaca basi concolora. DC. syst. 2. p. 79.* (v. s. in h. Burch.)
17. *P. Orientale* (Lin! spec. 727.) capsulis glabris subglobois, sepalis pilosis, caulibus unifloris scabris foliosis, foliis pinnatipartitis hispidis, lobis oblongis serratis. 2/ in Armeniâ. Curt. bot. mag. t. 57. Flos magnus intensè coccineus seu sanguineus cum ungue petalorum è purpuro nigro. — DC. syst. 2. p. 80.* (v. s.)
18. *P. bracteatum* (Lindl. coll. t. 23.) floribus bracteatis 4-5-petalis, capsulis glabris globoso-obovatis, sepalis pilosis, caulibus unifloris scabris foliosis, foliis bracteisque pinnatipartitis hispidis, lobis oblongis serratis. 2/ in imp. Russico. Priori simillimum sed certè differt calycis pedunculorumque pilis adpressissimis nec patentibus, florescentiâ paulò præcior etc. Ker. bot. reg. t. 658. (v. v. in h. Genev.) Sepala ut in priore sæpè 3.
19. *P. pilosum* (Sibth. et Smith. fl. græc. t. 492. ex Smith. prodr. 1. p. 360.) capsulis glabris obovatis, petalis pilosis, caule multifloro pilis patentibus hirsuto, foliis amplexicaulibus incis, undiquè pilosis. 2/ in Olympo Bithyno monte. — Petala pallidè lurido-coccinea seu intensè aurantiaca cum unguibus albis. — DC. syst. 2. p. 80.*
20. *P. setigerum* (DC. fl. fr. 5. p. 585.*) capsulis glabris obovatis, sepalis sub-setosis, caule glabro paucifloro, foliis amplexicaulibus glaucis inciso-repandis, dentibus setâ terminatis. (I) in insulis Stæchadum. Deless. Icon. sel. 2. t. 7. — An forsàn hæc est stirps sylvestris Papaveris somniferi nigri? (v. s. comm. a cl. Req.)
21. *P. somniferum* (Lin! spec. 726.) capsulis obovatis globosisve calycibusque

glabris, caule glabro glauco, foliis amplexicaulibus inciso-repandis, dentibus subobtusis. (I) in agris Peloponesi, in Ægypto, etc. Lam. ill. t. 451. DC. syst. 2. p. 81.* (v. v.)

α. *nigrum*, capsulis globosis substigmatē foraminibus apertis, seminibus nigris, pedunculis plarimis. — Bull. herb. t. 57. Petala sæpius purpurea, sed ex cl. Lucii Donal interdum alba et ideo stirps oleifera *œillette* dicta huc referenda. An ex eâ species propria?

β. *album*, capsulis ovato-globosis sub stigmatē foraminibus nullis seu oblitteratis, pedunculis solitariis, seminibus petalisque albis. — Woodv. med. bot. t. 185.

22. *P. CAUCASICUM* (Bieb. fl. taur. 2. p. 5.*) capsulis ovato-oblongis glabris, caule ramosissimum pedunculisque setis sparsis evanidis, foliis glaucis pinnatifidis, lobis ovato-oblongis setâ terminatis. (I) in glareosis verus portas Caucasicas. — Sims. bot. mag. t. 1675.* Petala colore Glaucii corniculati, magnitudine ferè P. Rhæadii, fugacissima. DC. syst. 2. p. 83. (v. v.)

23. *P. ARMENIACUM* (Lam. dict. 1. p. 247.) capsulis elliptico-oblongis calycibusque glabris, caule ramosissimo glabriusculo, foliis pinnati-sectis, lobis linearibus setâ terminatis. (I) ? in Armeniâ. Argemone Armeniaea. Lin. spec. 727. Aff. P. Caucasico. DC. syst. 2. p. 83.* (v. s. in h. Mus. Par.)

24. *P. FUGAX*. (Poir. dict. 5. p. 118.*) capsulis ovatis calycibusque glabris, caule ramosissimum virgato pedunculisque ferè glabris, foliis pinnatifidis incisis, lobis lanceolatis pilo terminatis. (I) in Persiâ monte Elwend. — Petala subrotunda, fugacia, pallidè chermesina. DC. syst. 2. p. 84.* (v. s.)

25. *P. TURBINATUM* (DC. syst. 2. p. 84.*) capsulis turbinatis calycibusque glabris, caule paucifloro, setis pedunculorum paucis patentibus, foliis pinnatifidis, lobis ovatis integris apice non setigeris. (I) in Oriente inter Bagdad et Kermancha. — Petala ferè P. dubii.

†. *Species non satis nota.*

26. *P. ? INTEGRIFOLIUM* (Vig. diss. 38.) in Hispaniâ et in agro Romano Bocc. mus. p. 77. t. 65. f. 1. Barr. obs. p. 47. ic. t. 1191. An P. Rhæadii varietas? An potius, jam sagacissimè observante cl. Bertolonio, *Seriola æthnensis* ante florescentiam malè depicta?

II. ARGEMONE. *Tourn. inst.* 239. t. 121. *Gært. fruct.* 1. p. 287. t. 60.

Petala 4-6. Stam. ∞. Stylus vix ullus stigmata 4-7 radiantia concava libera. Capsula obovata, 1-locularis, valvulis apice dehiscens, placentis linearibus. Semina sphaerica, scrobiculata. — Flores flavi aut albi. Pedunculi antè anthesin erecti.

1. *A. MEXICANA* (Lin! spec. 727.) (I) in campestribus montanis Mexici. Lam. ill. t. 452. Curt. bot. mag. t. 243. Folia sessilia, penninervia, repando-sinuata, sinibus dentato-spinulosis, sæpè albo maculata aut picta. DC. syst. 2. p. 85.* (v. v.)

β. *albiflora* (Horn. b. hafn. 469. Bot. mag. t. 2342.) (v. v.)

III. MECONOPSIS. *Vig. diss. p. 20. et 48. f. 3. DC. fl. fr. suppl. p. 586.*

Petala 4. Stamina ∞. Stylus brevis. Stigmata 4-6 radiantia convexa libera. Capsula obovata, 1-locularis, valvulis 4-6 apice dehiscens, placentis tenuibus vix intus in membranulas angustas productis. — Herbae perennes, succo flavescente foetæ. Genus medium inter Papaverem et Argemonem.

§. I. *Capsulis 5-6-valvibus lævibus. MECONOPSIS.*

1. *M. CAMERICA* (Vig! diss. p. 48. f. 3.) capsulis lævibus, foliis plurimis petio-

latis. \mathcal{U} in pratis sepius et dumetis umbrosis Pyrenæorum. Papaver Cambricum Lin! spec. 727. Smith. engl. bot. t. 66. — Flores 2-3 terminales, longè pedunculati, magnitudine ferè P. Rhæadis, è flavo-sulphurei. — DC. syst. 2. p. 87.* (v. v.)

§. 2. *Capsulis 4-valvis echinatis. STYLOPHORUM.*

2. *M. PETIOLATA* (DC. syst. 2. p. 87.*) capsulis echinatis, foliis 2-3-petiolatis glabriusculis. \mathcal{U} in sylvis opacis Americæ bor. Stylophorum petiolatum Nutt. gen. am. 2. p. 8.* Petala rotundata, basi cuneata, intense lutea, ferè magnitudine Glaucoii flavi. (v. s.)

3. *M. DIPHYLLA* (DC. syst. 2. p. 88.*) capsulis, foliis 2 sessilibus pilosiusculis. \mathcal{U} in sylvis umbrosis Americæ bor. Chelidonium diphylum Michx. fl. bor. am. 1. p. 309. Folia lobato-pinnatifida; pedunculi sæpè proliferi.

4. *M?* *NAPAULENSIS*, capsulis valdè echinatis, stylo ovarii ferè longitudine, stigmatè crassissimo, foliis plurimis sinuato-pinnatifidis, summis sessilibus, caule pedunculis sepalisque setosis. — in Napauliâ. Wallich. habitus ferè Glaucoii aut Argemones. An genus? an sectio propria. (v. s.)

IV. SANGUINARIA. Dill. h. elth. 252. Lin. gen. n. 645. Lam. ill. t. 449.

Petala 8-12 Stamina 24. Stigmata 2. Capsula oblonga bivalvis ventricosa utrinquè acuta; valvis deciduis, placentis 2 persistentibus. — Herba Boreali-Americana succo sanguineo foeta foliis longiùs basi cuneatis, paniculâ laxiore cernuâ. — In Novâ-Hispaniâ.

1. *S. CANADENSIS* (Lin! spec. 723.) \mathcal{U} in sylvis siccis fertilibus Americæ bor. Curt. bot. mag. t. 162. Bigel. med. bot. 1. p. 75. t. 7.* folium radicale sæpiùs 1. petiolatum, reniforme. Flos albus, sæpè et facillè plenus. DC. syst. 2. p. 89.* (v. s.)

V. BOCCONIA. Plum. gen. 35. t. 25. Lin. gen. n. 591. Gaert. fruct. 1. p. 204. t. 44. f. 1.

Petala 6. Stamina 8-24. Capsula elliptica 1-sperma. Semen 1. (altero forsàn abortivo) imæ capsulæ adfixum erectum, pulpâ molli ad basin involutum, hylo filiformi, albumine carnoso, embryone minutissimo erecto.

§. 1. *Foliis basi cuneatis, caulibus suffruticosis. Americanæ.*

1. *B. FRUTESCENS* (Lin. spec. 634.) foliis ovali-oblongis basi cuneatis pinnatifidis. \mathcal{F} in Mexico. Lam. ill. t. 394. Lodd. bot. cab. t. 83. Seminum pulpa cinnabarina. DC. syst. 2. p. 90.* (v. v.)

β . *cernua* (Moç. et Sessé ic. fl. mex. ined.), perennis, scapis unifloris.

γ . *subtomentosa* (Lher! in herb. Dombey) foliis subtùs præsertim in nervis subtomentosis, lobis magis crenatis. — In Peruviâ.

2. *B. INTEGRIFOLIA* (H. B. et K. nov. pl. gen. 1. p. 119. t. 35.*) foliis oblongis utrinquè attenuatis integris aut vix crenatis. \mathcal{F} DC. syst. 2. p. 91.

α . *Peruviana*, foliis planis, paniculâ confertiusculâ. H. B. et K. l. c. Poir. suppl. 1. p. 196. — in Andium Peruvianorum frigidis versus Cascas.

β . *Mexicana*, foliis margine subrevolutis, paniculâ laxâ. — B. nov. spec Moç. et Sessé. fl. mex. ined. — In Novâ-Hispaniâ.

§. 2. *Foliis basi cordatis, caulibus herbaceis. Chinenses.*

3. *B. CORDATA* (Willd. spec. 2. p. 841.*) foliis cordatis sublobatis. \mathcal{U} in Chinâ. Jacq. fragm. 63. t. 93. f. 1. Sims. bot. mag. t. 1905. DC. syst. 2. p. 91.* (v. v. sub dio in hortis angl. et nunc Genev.)

VI. ROEMERIA. *Medik. in Ust. ann. 1792. vol. 3. p. 15. DC. syst. 2. p. 92. non Manch, nec Thunb. etc.*

Petala 4. Stamina ∞. Capsula elongata 2-3-4-valvis, valvulis ab apice ad basin dehiscentibus. Semina reniformia scrobiculata, cristâ glandulosâ destituta. — Herbæ annuæ, succo flavo foetæ. Flores violacei.

1. R. HYBRIDA (DC. syst. 2. p. 92.*) siliquis 3-4-valvibus erectis apice rigidè setosis. ① in agris et vineis regionis Mediteraneæ. *Chelidonium hybridum* Lin! spec. 724. Smith. engl. bot. t. 201. *Chelid. violaceum* Lam! fl. fr. 3. p. 169. (v. v.)

β. *eriocarpa*, siliquis à basi ad apicem piloso-setosis. *Chelid. dodecandrum* Forsk! fl. æg. p. 100.* in desertis Kahirinis.

γ. *velutina*, caule molliter villosa. *Chelid. hybridum* Bieb. fl. taur. 2. p. 3. — in Tauria.

2. R. REFRACTA (DC. syst. 2. p. 93.*) siliquis 3-4-valvibus glabris pedicello demum recurvo pendulis. ① in Tauriâ. *Glaucium refractum* Stev! in litt. *Ræmeria refracta*, Deless. Icon. sel. 2. t. 8. Valdè aff. R. hybridæ, sed differt foliis bipinnatis, laciniiis paulò magis obtusis. (v. s.)

3. R. BIVALVIS (DC. syst. 1. p. 93.*) siliquis 2-valvibus setosis subincurvis. ① in Mesopotamiâ. Folia pinnatim multifida, lobis linearibus pinnatipartitis; petala violacea. (v. s. in h. Mus. Par.)

VII. GLAUCIUM. *Tourn. inst. 254. t. 130. Gaertn. fruct. 2. p. 165. t. 115.*

Petala 4. Stamina ∞. Capsula elongata 2-valvis, dissepimento cellulari 2-locularis. Semina ovato-reniformia. — Herbæ biennes, semper-virentes, glaucæ, succo croceo acri scatentes.

1. G. FLAVUM (Crantz fl. austr. 2. p. 141.) caule glabro, foliis caulinis repandis, siliquâ tuberculato-scabrinusculâ. ② in glareosis arenosisve littoralibus secus mare, lacus aut flumina in Europâ austr. et med. *Chelidonium Glaucium* Lin. spec. 724. Smith. engl. bot. t. 8. — DC. syst. 2. p. 94. (v. v.)

2. G. FULVUM (Smith. exot. bot. 1. p. 11. t. 7.) caule glabro, foliis caulinis rotundo-sinuatis, siliquis scabris, floribus subsessilibus. ② in glareosis arenosisve Europæ australis. *Chelidonium fulvum* Poir. suppl. 5. p. 606. Petala lateritio-coccinea, basi cuneata flava, sæpè apice subbifida, obtusissima. DC. syst. 2. p. 95.* (v. v.)

3. G. CORNICULATUM (Curt. lond. 6. t. 32.) caule piloso, foliis caulinis pinnatifidis incis, siliquâ setosâ. ① in Galliâ et Angliâ. DC. syst. 2. p. 96.* (v. v.)

α. *phœniceum*, *Chelidonium corniculatum* Lin! spec. 724. — *Glaucium phœniceum* Smith. engl. bot. t. 1433. — Clus. hist. 2. p. 91.

β. *flaviflorum*. *Glaucium corniculatum* flore flavo Stev! in litt. — Nullo modo à priori differt nisi flore flavo.

4. G. RUBRUM (Sibth. et Smith. fl. græc. t. 488.) caule piloso, foliis caulinis pinnatifidis incis, siliquâ pilosiusculâ. ② ad viam inter Smyrnam et Bursam et in insulâ Rhodo. Petala rubro-fulva nec phœnicea. Siliqua mollissimè pilosa demum glabrata, nunquam setoso-aspera. DC. syst. 2. p. 97.* (v. s. in h. Mus. Par.)

5. G. PRASICUM (DC. syst. 2. p. 97.*) foliis glaucis velutinis, radicalibus pinnatisectis, segmentis inferioribus minoribus, terminali maximo reniformi. ② in Persiâ circâ Teheran. (v. s. sine flore in h. Mus. Par.)

VIII. CHELIDONIUM. *C. Bauh. pin. 144. Gaert. fruct. 2. p. 164. t. 115. f. 5. — Chelidonii sp. Lin.*

Sepala 2, glabra. Petala 4. Stamina ∞. Capsula elongata 2-valvis 1-

locularis, valvulis à basi ad apicem dehiscentibus. Semina cristâ glandulosâ instructa. — Herbæ perennes sempervirentes teneræ, succo croceo acri foetæ. Flores flavi, minores quàm in Glaucio et Roemeria. DC. syst. 2. p. 98.

1. *C. MAJUS* (Mill. dict. n. 1. OEd. fl. dan. t. 676.) pedunculis umbellatis, foliis pinnatisectis segmentis subrotundis dentato-lobatis, petalis ellipticis integris. \mathcal{L} in umbrosis secus muros Europæ totius. Cl. majus. var. α . Lin! spec. 723. Smith. engl. bot. t. 1531. Variat flore simplici et pleno. DC. syst. 2. p. 98.* (v. v.)

2. *C. GRANDIFLORUM*, pedunculis umbellatis, foliis pinnatisectis, segmentis subrotundis dentato-lobatis, petalis subrotundis crenatis \mathcal{L} in Dahuriâ. Fischer. Ch. Dahuricum Hortul. — Cl. Nestler qui vidit vivam me monet in litteris hanc esse speciem propriam et ex eo caract. eru. (v. s.)

3. *C. LACINIATUM* (Mill. dict. n. 2.) pedunculis umbellatis, foliis pinnatisectis, segmentis in lobos lineares laciniatos acutos incis, petalis serratis incisive. \mathcal{L} in Germaniâ et Angliâ. C. majus var. β . Lin! spec. 724. C. quercifolium Will. fl. lorr. 2. p. 613. — Planta caracteribus levibus et vix certis distincta. DC. syst. 2. p. 99.* (v. v.)

β . *fumariæfolium*, lobis in lobulos profundiores magis irregulariter et omnino lineares fissis. Moris hist. 2. p. 258.

†. *Species non satis notæ.*

4. *C. SINENSE* (DC. syst. 2. p. 100.) \mathcal{L} 5 in diversis Sinarum provinciis. C. majus Lour. fl. coch. 1. p. 402.* Descriptio Loureiriana à C. majore differt caule suffruticoso, pedunculis multifloris non umbellatis, radice longâ intus et floris flavâ.

5. *C. JAPONICUM* (Thunb. fl. jap. 221.*) in Japoniâ. Folia petiolata, pinnatisecta cum impari, flores axillares solitarii, pedunculati. Fructus ignotus et ideò genus dubium. DC. syst. 2. p. 100.*

IX. *HYPECOUM*. Tourn. inst. 230. t. 115. Lin. gen. 171. Gaertn. fruct. 2. p. 164. t. 115.

Petala 4, interiora sæpius 3-loba. Stamina 4. Capsula siliquæformis, 2-valvis, transversè nodosa seu articulata. Semina inter articulationes solitaria. — Herbæ annuæ, parvæ, succo flavo foetæ. DC. syst. 2. p. 101.

1. *H. PROCUMBENS* (Lin. spec. 181.) siliquis articulatis compressis arcuatis, petalis 3-lobis, externis dorso glabris. ① in agris præsertim maritimis, et subsabulosis à Lusitaniâ ad Astracan. Lam! ill. n. 1720. t. 88. Schkuhr. handb. 1. p. 90. t. 27. Flores flavi; petala 2, exteriora majora. DC. syst. 2. p. 102.* (v. v.)

2. *H. LITTORALE* (Wulf. in Jacq. coll. 2. p. 205.* ic. rar. 2. t. 309.) siliquis articulatis compressis arcuatis, petalis omnibus integris lineari-spathulatis, externis dorso pubescentibus. ① in maritimis arenosis aridis Cariuthiæ secus Adriaticum. H. procumbenti simillimum, sed ex autoribus differt petalis exterioribus majoribus apice subcochleariformibus, interioribus calycis longitudine lanceolatis. DC. syst. 2. p. 102.

3. *H. IMBERBE* (Sibth. et Sm. fl. græc. 2. p. 47. t. 156.*) siliquis articulatis compressis arcuatis, sepalis dentato-fimbriatis, petalis omnibus imberbibus. ① in insulâ Cypro. Calyx dentato-fimbriatus. Petala interiora semi-trifida nec tripartita, lobis lateralibus oblongis divaricatis. DC. syst. 2. p. 103.*

4. *H. PATENS* (Willd. hort. ber. t. 4.) siliquis articulatis teretibus curvatis, petalis glabris, exterioribus subintegris unguiculatis cordiformibus, interioribus tripartitis. ① in desertis arenosis prope Alexandriam. Mnemosilla

Ægyptiaca Forsk. æg. 122.* Petala flava, 2 exteriora magna unguiculata limbo latissimo acuto utrinquè quasi lobulis aucto et inde fere hastæformi. DC. syst. 2. p. 103.* (v. v.)

5. *H. PENDULUM* (Lin. spec. 181.) siliquis nodosis teretibus pendulis, petalis glabris, 2 exterioribus ovato-oblongis pendulis, 2 interioribus 3-partitis. ① in agris arenosis ab Hispaniâ ad mare Caspium. — Mill. ic. t. 250. f. 1. Petala glabra pallidè flava magna ovali oblonga basi subattenuata integra. DC. syst. 2. p. 103.* (v. v.)

6. *H. ERECTUM* (Lin. spec. 181.) siliquis inarticulatis erectis compressis, petalis glabris, exterioribus cuneiformibus subtrilobis, interioribus 3-fidis, lobis lateralibus subbilobis, medio parvo. ① in Dahuriâ. — Amm. ruth. 58. t. 9. Flores generis ampliores, flavidi, immaculati; sepala minima, lanceolata, acuminata. DC. syst. 2. p. 104. (v. s.)

Ordo X. FUMARIACEÆ. DC. *syst.* 2. p. 105.

Calyx 2-sepalus, membranaceus, deciduus, parvus. Petala 4, inter se cruciata, nunc omnia libera, nunc omnia basibus coalita, nunc inferius liberum et 3 superiora coalita; 2 exteriora cum sepalis alternantia, nunc ambo basi in calcar cavum gibbositateve æqualiter producta, nunc inferius planum ecalcaratum et superius basi calcaratum gibbosumve; 2 interiora cum externis cruciata, oblongo-linearum, apice subcallosa colorata puncto terminali connexa, antheras et stigmata foventia. Glandula nectarifera intrà calcar. Stam. Filamenta 6 disposita in phalanges adelphasve duas oppositas antè petala exteriora sitas et ideò cum petalis internis alternas, nunc rarius omnia libera, cum 3 cujusve phalangis coalita et filamentum sæpius 3-nervium 3-antheriferum constituentia. Antheræ 6 parvæ, laterales cujusve phalangis 1-loculares, media 2-locularis et ideò antherarum loculi 8, seu antheræ 4 strictiori sensu habendæ. Ovarium 1 liberum. Stylus filiformis. Stigma bilamellatum petalis interioribus parallelum. Fructus siccus, formâ varius, nunc siliquosus bivalvis polyspermus dehiscentis, valvulis oppositis, placentis 2 ad suturas nerviformibus persistentibus, nunc 2-valvis 2-spermus valvulis coalitis indehiscentis, nunc evalvis? indehiscentis abortu? monospermus. Semina placentis lateralibus adfixa, horizontalia, ovato-globosa, nitida, nigra, arillo seu carunculo basi instructa. Albumen carnosum. Embryo basilaris, in fructibus indehiscentibus parvus rectus, in dehiscentibus longior subarcuatus. Cotyledones oblongæ, planæ. — Herbæ succo aqueo nec lacteo foetæ, teneræ. Flores purpurei albi aut flavi, Polygaleas Balsamineasve æmulantes.

I. DICLYTRA. DC. *syst.* 2. p. 107.

Petala 4, 2 exteriora basi æqualiter calcarata aut gibbosa. Siliqua bivalvis polysperma. — Herbæ perennes. Flores racemosi, albi aut purpurascens.

1. D. CUCULLARIA (DC. *syst.* 2. p. 108.*) calcaribus 2 rectis acutis, scapo nudo, racemo simplici. \mathcal{L} in America bor. *Fumaria cucullaria*. Lin! spec. 983. Sims. bot. mag. t. 1127. Flores albi, apice flavi. (v. v.)

β . *divaricata*, calcaribus divaricatis. *Fumaria cucullaria* Mill! dict. n. g. ex h. Banks. — An species propria? (v. s.)

2. D. BRACTEOSA (DC. *syst.* 2. p. 109.*) calcaribus 2 rectis acutis, caulē folioso, bracteis incisis. \mathcal{L} in America bor? Aff. D. cucullariæ, sed certè differt staturâ minore, etc. (v. s. in h. Mus. Par.)

3. D. FORMOSA (DC. *syst.* 2. p. 109.*) calcaribus 2 subincurvis obtusis, scapo nudo, racemo subcomposito, stigmate bi-angulato. \mathcal{L} in rupibus umbrosis Virginie, Carolinæ, et Canadæ. *Fumaria formosa* Andr. bot. rep. t. 393. Sims. bot. mag. t. 1335.* (v. v.)

4. *D. EXIMIA* (DC. syst. 2. p. 109.*) calcaribus 2 subincurvis obtusis brevibus , scapo nudo , racemo composito , stigmate quadriangulato. \mathcal{U} in Americâ bor. *Fumaria eximia* Ker. bot. reg. 1. t. 50.* (v. s.)
5. *D. SPECTABILIS* (DC. syst. 2. p. 110.*) calcaribus 2 obtusissimis ventricosis brevibus , caule folioso , foliorum segmentis obovato-cuneatis incis. \mathcal{U} in Sibiria ad fines Chinæ et in Chinâ bor. *Fumaria spectabilis* Lin! amœn. 7. p. 457. t. 7. spec. 953.* Flores purpurei , pollicem longi. (v. s. in h. Juss.)
6. *D. TENUIFOLIA* (DC. syst. 2. p. 110.*) calcaribus 2 brevissimis obtusissimis , scapo nudo 1-3-floro , pedicellis calyce brevioribus , foliis multifidis , lobulis linearibus. \mathcal{U} in Kamtschatka. *Corydalis tenuifolia* Pursh! fl. bor. amer. 2. p. 462. Deless! ic. sel. 2. t. 9. f. B. (v. s.)
7. *D. CANADENSIS* , calcaribus 2 brevibus obtusis , scapo nudo simplici paucifloro , pedicellis brevibus , foliis multifidis , lobis linearibus , stigmate porrecto 4-lobo. \mathcal{U} in Canadâ circa Montreal. *Corydalis Canadensis*. Gold. in Edinb. phil. journ. 1822. p. 330. Folia glauca. Herba 6-8 poll. longa.
8. *D. LACHENALIEFLORA* (DC. syst. 2. p. 111.*) calcaribus 2 brevissimis obtusissimis , scapo nudo 3-4-floro , pedicellis calyce longioribus , foliis multifidis , lobis linearibus acutissimis. \mathcal{U} in Sibiria transbaicalensi. *Corydalis lachenaliæflora* Fisch! in litt. — Rudolphi in mem. sc. Petersb. 1. t. 19. Flores purpurei. *D. tenuifoliæ* valdè affinis , sed distincta floribus minoribus. (v. s.)

II. ADLUMIA. Raf. in Desv. journ. bot. 1809. 2. p. 169.

Petala 4 in corollam monopetalam basi bigibbosam fungosam persistentem coalita. Siliqua bivalvis polysperma. — Herba glabra , et petiolis cirrhosis scandens. DC. syst. 2. p. 111.

1. *A. CIRRHOSA* (Raf. l. c.) ② in Canadâ et Pensylvaniâ. *Corydalis fungosa* Vent! choix. t. 19.* Flores albidæ , pallidè subrosæ. Semina 4-6 lucida nigra. DC. syst. 2. p. 111.* (v. s. et v.)

III. CYSTICAPNOS. Boerh. lugd. 391. Gaertn. fruct. 2. p. 161. t. 115.

Petala 4 , unicum basi gibbum. Capsula vesicaria polysperma , placentis reticulo membranaceo connexis. — Herba scandens , glabra.

1. *C. AFRICANA* (Gaert. l. c.) ① ad caput Bonæ-Spei. *Fumaria vesicaria*. Lin! spec. 985. Corolla roseo-albida. DC. syst. 2. p. 112.* (v. v.)

IV. CORYDALIS. DC. fl. fr. 4. p. 636. syst. 2. p. 113.* — Capnoides. Tourn. inst. 423. t. 237.

Petala 4 , unicum basi calcaratum. Siliqua 2-valvis compressa polysperma. — Herbæ glabræ , sæpè glaucæ \mathcal{U} aut ①.

§. 1. Radice fusiformi , Caule simplici , foliis caulinis 2 oppositis.

LEONTICOÏDES.

1. *C. VERTICILLARIS* (DC. syst. 2. p. 114.*) caule simplicissimo 2-folio , foliis glaucis oppositis biternatis sectis , lobis segmentisve linearibus. ② in Persiâ. Flores 8 lin. longi. Calcar conico-subulatum , obtusum. (v. s. in h. Mus. Par.)
2. *C. OPPOSITIFOLIA* (DC. syst. 2. p. 114.*) caule simplicissimo bifolio , foliis glaucis oppositis trisectis , petioli ramo medio trisecto , segmentis lobisve ovatis? ② in Mesopotamiâ. Flos pollicem longus , calcar 6 lin. (v. s.)

§. 2. Radice tuberosâ , caule simplici , foliis caulinis paucis alternis.

CAPNITES.

3. *C. RUTÆFOLIA* (DC. syst. 2. p. 115.*) caule simplici exsquamato , foliis 2 suboppositis glaucis trisectis , segmentis ovatis aut medio tripartito , bracteis ovatis. \mathcal{U} in insulâ Cypro. *Fumaria rutæfolia* Sibth. et Smith. fl. græc. t. 667.

Calcar horizontale, subadscendens, obtusum, apice callosum, flore longius, ferè semi-pollicare. An ad priorem sect. amandanda? (v. s. in h. Banks.)

4. *C. DECUMBENS* (Pers. ench. 2. p. 269.) caule simplici decumbente, foliis biternatim sectis, lobis cuneiformibus 3-dentatis, bracteis ovatis integris, racemis 3-4-floris. \mathcal{L} in Japoniâ. *Fumaria decumbens* Thunb. nov. act. petrop. 12. p. 102. t. A.* Floris cornu limbi longitudine. DC. syst. 2. p. 115.*
5. *C. PAUCIFLORA* (Pers. ench. 2. p. 69.) caule simplici exsquamato, foliis caulinis 2-3 infrâ medium biternatim sectis, segmentis 3-partitis, lobis obovatis, bracteis ovatis acutis, racemo conferto paucifloro. \mathcal{L} in montibus Altaicis Sibiriz. *Fumaria pauciflora*. Steph. in Willd. spec. 3. p. 861. Calcar crassum, apice incurvum, obtusissimum, tubo ferè longius. — DC. syst. 2. p. 116.* Deless! ic. sel. 2. t. 9. f. A. (v. s.)
- β . *aquilegifolia* (Patr! ined. in h. Deless.) — à priore vix differt. Racemus 7-8-florus. Flores pollicem longi. (v. s. in h. Deless.)
- γ ? *Altaica* (Ledeb. mem. acad. petersb. 5. 1815. p. 551.*) Flores penduli.
6. *C. MARSHALLIANA* (Pers. ench. 2. p. 269.) caule simplici exsquamato, foliis 2 suprâ medium sitis biternatim sectis, lobis ovalibus integris, racemo brevi, bracteis ovatis. \mathcal{L} in Tauriâ. *Fumaria Marshalliana* Pall. nov. act. petrop. 10. p. 315. Bieb. fl. taur. 2. p. 145.* Calcar rectum, aut vix subincurvum, obtusum. DC. syst. 2. p. 116.* Deless! ic. sel. 2. t. 10. (v. s.)
7. *C. LONGIFLORA* (Pers. ench. 2. p. 269.) caule simplici squamifero, foliis biternatim sectis, segmentis tripartitis, lobis ovali-oblongis, bracteis oblongis integris, racemo elongato, calcare pedicello longiore. \mathcal{L} in Alpibus Altaicis Sibiriz. *Fumaria Schangini*. Pall. act. petrop. 1779. 2. p. 267. t. 14. f. 1-3.* Calcar gracile, subulatum, apice subincurvum, acutum, ferè pollicem longum. — DC. syst. 2. p. 116.* (v. s.)

β . *caudata* (Lam! dict. 2 p. 569.*) non differt à priore nisi flore minore et lobis foliorum paulò obtusioribus. (v. s. in h. Juss.)

8. *C. TUBEROSA* (DC. fl. fr. 4. p. 637.* syst. 2. p. 117.*) caule simplici exsquamato, foliis 2 biternatim sectis, segmentis cuneatis inciso-multifidis, bracteis ovatis integris, radice cavâ. \mathcal{L} in sepibus Europæ. *Fumaria cava* Mill. dict. n. 7. Flores horizontales, purpurascens (Curt. bot. mag. t. 232.) aut albi (Bot. mag. t. 234o.) (v. v.)
9. *C. FABACEA* (Pers. ench. 2. p. 269.) caule subsimplici erecto, sub infimo folio squamigero, foliis 3-4 petiolatis biternatim sectis, segmentis oblongis obtusiusculis, bracteis ovatis acutis pedicello longioribus, radice solidâ. \mathcal{L} in umbrosis montanis Sueciæ, Daniæ, Germaniæ, Vallesiz, etc. *Fumaria fabacea* Reiz. prod. ed. 2. n. 859. excl. syn. fl. Dan. — Horn. fl. Dan. t. 1394. — DC. syst. 2. p. 118.* (v. s.)
10. *C. CAUCASICA* (DC. syst. 2. p. 119.*) caule simplicissimo erecto, sub infimo folio squamigero, foliis 2 biternatim sectis, segmentis in lobos lineares incis, bracteis oblongis integris, racemo laxiusculo. \mathcal{L} in montibus sylvaticis Caucasi. *Fumaria fabacea* Bieb. fl. taur. 2. p. 145. Differt à *C. fabacæ* caule simplicissimo bifolio, foliorum segmentis in lobos lineares fissis. (v. s.)
11. *C. BULBOSA* (DC. fl. fr. 4. p. 637.) caule simplici erecto sub infimo folio squamigero, foliis 3-4 petiolatis biternatim sectis, segmentis cuneatis oblongis bracteis apice incis, radice solidâ. \mathcal{L} in sepibus Europæ. *Fumaria solida* Smith. engl. bot. t. 1471. DC. syst. 2. p. 119.* (v. v.)
- β . *viridiflora*. — Moris. hist. 2. s. 3. p. 261. t. 12. f. 7.
12. *C. ANGUSTIFOLIA* (DC. syst. 2. p. 120.*) caule simplici infrâ infimum folium squamigero, foliis 2 petiolatis biternatim sectis, segmentis in lobos lineares fissis, bracteis apice inciso-serratis. \mathcal{L} in Iberiâ. *Fumaria angustifolia* Bieb. fl. taur. 2. p. 146.* Vix distincta à *C. bulbosâ*. (v. s.)
13. *C. INCISA* (Pers. ench. 2. p. 269.) caule simplici erecto, foliis biternatim sectis, segmentis acutè incis, bracteis oblongo-cuneatis incis pedicello

brevioribus. \mathcal{U} in Japoniâ. *Fumaria incisa* Thunb. nov. act. petr. 12. p. 104. t. D. Differt à *C. nobili* et bracteata dentibus foliorum acuminatis, bracteis inciso-dentatis pedicello brevioribus. DC. syst. 2. p. 121.*

14. *C. BRACATEATA* (Pers. ench. 2. p. 269.) caule simplici erecto propè basin squamigero, foliis 2 biternatim sectis, segmentis in lobos lineares fissis, bracteis cuneatis apice profundè incisis pedicello longioribus, calcare recto longo. \mathcal{U} in Sibiriz montibus Altaicis. *Fumaria bracteata* Steph. in Willd. spec. 3. p. 858.* Flores flavo-sulphurei horizontales. DC. syst. 2. p. 121.* (v. s.)

15. *C. NOBILIS* (Pers. ench. 2. p. 269.) caule erecto simplici exsquamato, foliis bipinnatim sectis, lobis cuneatis apice incisis, bracteis acutis integris incisive. \mathcal{U} in Sibiria. *Fumaria nobilis* Jacq! h. vind. t. 116. Sims. bot. mag. t. 1953.* Flores albedo-flavescentes, cum calcare pollicem longi. DC. syst. 2. p. 121.* (v. v.)

β . *odontophylla*, bracteis omnibus inciso-dentatis. A priore differt habitu graciliore, bracteis omnibus ovatis apice inciso-dentatis. (v. s.)

§. 3. *Radice fibrosa, caule ramoso, foliis caulinis plurimis alternis.*

CAPNOIDES.

16. *C. PÆONIEFOLIA* (Pers. ench. 2. p. 269.) caule erecto ramoso, foliis bipinnatim sectis, segmentis ovatis, ultimis lobatis, bracteis oblongo-linearibus acutis, siliquis ovatis. \mathcal{U} ad radices rupium Sibiriz. *Fumaria pæoniæfolia* Steph. in Willd. spec. 3. p. 859.* — Gmel. fl. sib. 4. p. 66. t. 34.* Flores magni semi-nucleares purpurei, calcar conicum, rectum, subacutum. — An hujus sectionis? — DC. syst. 3. p. 122.* (v. s. in h. Lamb.)

17. *C. GLAUCA* (Pursh. fl. bor. am. 2. p. 463.) caule erecto ramoso, foliis glaucis decompositis, segmentis petiolulatis cuneatis trifidis, bracteis linearibus pedicello brevioribus, siliquis linearibus. ① in declivibus montium Canadæ. *Fumaria sempervirens*. Lin! spec. 984. *Fumaria glauca* Curt. bot. mag. t. 179. Calcar corollâ dimidiò triplòve brevius, obtusum. — DC. syst. 2. p. 123.* (v. v.)

18. *C. STRICTA* (Steph. ex Fisch! in litt.) caule erecto subramoso, foliis glaucis pinnatisectis, segmentis pinnatim et acutè incisis, bracteis linearibus pedicello subbrevioribus, siliquis ovato-lanceolatis. \mathcal{U} in Sibiria. Flores mihi ignoti. Pedicelli fructiferi sesquilineares, deflexo-patentes. DC. syst. 2. p. 123. (v. s.)

19. *C. CHÆROPHYLLA*, caule erecto ramoso, foliis bipinnatisectis, segmentis ovato-oblongis pinnatifidis, bracteis parvis acutis, siliquis obovato-oblongis pedicello brevioribus. — Hab. in Napauliâ. (v. s. comm. à cl. Wallich.)

20. *C. SIBIRICA* (Pers. ench. 2. p. 270.) caule suberecto ramoso, foliis subglaucis bipinnatisectis, segmentis in lobos oblongo lineares incisis, bracteis linearibus pedicello subæqualibus, siliquis ovalibus pedicello vix longioribus. \mathcal{U} in Sibiriz Transbaicalensibus regionibus. *Fumaria Sibirica* Lin. f. ! suppl. 314. — Gmel. fl. sib. 4. p. 65. t. 33. — DC. syst. 2. p. 124.* (v. s.)

21. *C. IMPATIENS* (Fisch! in DC. syst. 2. p. 124.*) caule subdiffuso ramoso, foliis glaucis bipinnatisectis, segmentis in lobos oblongo-lineares incisis, bracteis setaceo-linearibus pedicello subæqualibus, siliquis linearibus pedicello duplò longioribus. ① in Dahuriâ. *Fumaria impatiens* Pall. itin. ed. min. 3. p. 233? Proxima *C. glaucæ* et præsertim *C. Sibiricæ*. Differt siliquis in morem cardamine elasticè dissilientibus et revolutis. (v. s.)

22. *C. LONGIPES*, caule ramoso subdiffuso, foliis bipinnatisectis glaucescentibus, bracteis cuneatis apice incisis, siliquis linearibus pedicello dimidiò brevioribus. — In Napauliâ. — Habitus omninò *C. Sibiricæ* et *C. impatiens* sed pedicellis 8-9 lin. longis distincta. (v. s. comm. à cl. Wallich.)

23. *C. AUREA* (Willd. enum. 740.) caule ramoso diffuso, foliis glaucis bipinnatisectis, lobis oblongo-linearibus, bracteis lanceolato-linearibus acuminatis siliquisque linearibus pedicello quadruplò longioribus. ① ② in rupibus umbrosis Pensylvaniæ, Virginiz et Canadæ. *Fumaria aurea* Muhl. in Willd. enum. p. 740. Ker. bot. reg. t. 66.* — DC. syst. 2. p. 125.* excl. syn. Raf. et Poir! (v. s.)

24. *C. FLAVULA* (Raf! in Deav. journ. bot. 1808. t. p. 224.) caule subramoso, foliis glaucis bipinnatisectis, lobis oblongo-linearibus, bracteis ovatis acuminatis, siliquis linearibus torulosis pedicello duplò longioribus. ①② circa Philadelphiam. Calyx parvus foliaceus. (v. s.)
25. *C. RACEMOSA* (Pers. ench. 2. p. 270.) caule ramoso erecto, foliis bipinnatisectis, segmentis ovatis obtusè trilobis dentatis, bracteis pedicello triplò longioribus. ♀ in Japoniæ muris antiquis et rupibus. *Fumaria racemosa* Thunb. nov. act. petr. 12. p. 103. t. B.* Calcar breve obtusum. An fructus monospermus? — DC. syst. 2. p. 125.
26. *C. CAPNOIDES* (Pers. ench. 2. p. 270.) caule ramoso diffuso, foliis bipinnatisectis, segmentis obovato-cuneatis trifidis, bracteis lineari-subulatis pedicello triplò brevioribus, siliquis linearibus vix pedicello longioribus. ♀ in fissuris rupium Europæ mediæ et australis. DC. syst. 2. p. 125.* (v. v.)
a. albida, floribus albis aut albidis. ♀ in fissuris rupium et murorum Mauritaniae Tingitanæ. *Fumaria capnoides*. Lin! spec. 984. — Pluk. alm. t. 90. f. 2.
β. lutea, floribus luteis. In Angliâ, Germaniâ, etc. *Fumaria lutea* Lin. mant. 258. Smith engl. bot. t. 588. *Capnoides lutea* Gaertn. fruct. 2. p. 163. t. 115. f. 3.
27. *C. ACAULIS* (Pers. ench. 2. p. 270.) scapis nudis, foliis radicalibus pinnatisectis, segmentis obovato-cuneatis trifidis. ① Hab. Tergesti. *Fumaria acaulis* Wulf. in Jacq. coll. 2. p. 203.* ic. rar. 3. t. 554. Lam. ill. t. 597. f. 3. Corolla dilutè flavens. Calcar breve. An prioris mera varietas? — DC. syst. 2. p. 127.*
28. *C. URALENSIS* (Fisch! in DC. syst. 2. p. 127.*) caule erecto subramoso vix foliis radicalibus longiore, foliis longè petiolatis trisectis, segmentis 3-partitis, lobis subincisis; racemo paucifloro, bracteis apice incisis. ♀? in montibus Uralensibus. Habitus ferè *C. acaulis*, sed distinctissima. Flores flavi cum calcare, 6 lin. longi. Calcar rectum, tenue. (v. s.)
29. *C. BREVIFLORA* (DC. syst. 2. p. 127.*) caule ramoso diffuso, foliis bitermatim sectis, petiolis subcirrhosis, segmentis paucis in lobos oblongo-lineares 3-partitis, bracteis oblongis pedicello subæqualibus. ① in Kamchatkâ. Aff. hinc *C. Capnoidi*, illinc *C. claviculata*. Flores pallidi. Calcar rectum, latum, subacutum floris dimidiam longitudinem occupans. (v. s.)
30. *C. CLAVICULATA* (DC. fl. fr. 4. p. 638. syst. 2. p. 128.*) caule ramoso diffuso-scandente, foliis bipinnatisectis, petiolis cirrhosis, segmentis ovalibus integris, bracteis oblongis pedicello longioribus. ① in sepibus saxosisque uliginosis Europæ. *Fumaria claviculata* Lin. spec. 985. Smith engl. bot. t. 103. Flores ex albo subflavescentes, per exsiccationem flavi. Calcar obtusum breve. (v. v.)
31. *C. PALLIDA* (Pers. ench. 2. p. 270.) caule ramoso decumbente, foliis bipinnatisectis, bracteis acutè dentatis. ♀? in Japoniâ. *Fumaria pallida* Thunb. nov. act. petr. 12. p. 103. t. C.* Flores pallidi, calcar obtusum, longiusculum.

V. SARCOCAPNOS. DC. syst. 2. p. 129.

Petala 4, unicum basi calcaratum. Capsula 2-valvis indehiscens 2-sperma, valvis 3-nerviis planiusculis.

1. *S. ENNEAPHYLLA* (DC. fl. fr. 5. p. 587.* syst. 2. p. 129.*) foliis petiolo ramoso tritermatim sectis, segmentis ovato-orbiculatis. ♀ ferè ♂ in Lnsitaniâ. *Fumaria enneaphylla* Lin! spec. 984. Lam! ill. t. 597. f. 4. Calcar rectum flore dimidio brevius. (v. s.)
2. *S. CRASSIFOLIA* (DC. syst. 2. p. 130.*) foliis petiolo simplici indivisis aut trisectis, segmentis ovatis. ♀ prope Tlemsen in Mauritanîâ. *Fumaria crassifolia* Desf! fl. atl. 2. p. 126. t. 173.* Flores pallidè flavi. Bracteæ ovatæ, minimæ. (v. s.)

VI. FUMARIA Tourn. inst. p. 422. t. 237. Gaert. fruct. 2. p. 162. t. 115.

Petalum unicum basi gibbum aut calcaratum. Cariopsis indehiscens 1-sperma stylo post anthesin deciduo mutica.

§. 1. Platycapnos. DC. syst. 2. p. 131.

Siliculæ compressæ.

1. *F. SPICATA* (Lin! spec. 985.) siliculis compressis lævibus, racemis spicatis, pedicellis bractea multò brevioribus, caulibus erectis. ① in agris oleraceisque Lusitanie. — Moris. hist. 2. p. 261. s. 3. t. 12. f. 11. Calcar brevissimum. — DC. syst. 2. p. 131.* (v. v.)

β. *aurantiaco-crocea*. — in Hispaniâ. Distinguitur habitu paulò minori.

2. *F. TURBINATA* (Smith! in DC. syst. 2. p. 132.*) siliculis compressis turbinate truncatis striatis, caulibus subscandentibus, petiolis apice cirrhosis. — in Oriente. *F. claviculata* herb. Lin! (sed evidenter errore quodam) Flores racemosi, parvi, vix noti, stylo brevissimo exserto. An hujus generis? (v. s. in h. Lin.)

3. *F. CORYMBOSA* (Desf! act. soc. hist. nat. par. 1. p. 26. t. 6.* fl. atl. 2. p. 124.*) siliculis ovatis compressis punctulatis apiculatis, racemis subcorymbosis, pedicellis fructiferis bractea multò longioribus, caulibus diffusis. ② in rupium fissuris Atlantis. *F. Africana* Lam. dict. 2. p. 56y.* Flores racemoso-corymbosi, conferti. — DC. syst. 2. p. 132.* (v. s.)

Sect. II. SPHÆROCAPNOS. DC. syst. 2. p. 133.

Siliculæ globosæ.

4. *F. CAPREOLATA* (Lin. spec. 985.) siliculis globosis, pedicellis fructiferis recurvatis bractea longioribus, racemis oblongis, caulibus subscandentibus, foliis bipinnatisectis, petiolis subcirrhosis. ① in asperis saxosis et fissuris rapium Europæ occidentalis et australis. DC. ic. rar. 1. t. 34. syst. 2. p. 133. Calcar compressum, obtusum, breve, quasi mitræforme. (v. v.)

β? *Burchellii*, pedunculis patulis, pedicellis fructiferis bractea duplò longioribus. Fumaria n. 1298. Burch! cat. geogr. pl. afr. austr. An species propria. (v. s. in h. Burch.)

5. *F. MEDIA* (Lois. not. p. 101.*) siliculis globosis depressiusculis, pedicellis fructiferis erectis bractea duplò longioribus, racemis laxiusculis, caulibus erectis, foliis suprâ decompositis petiolis subcirrhosis. ① in agris oleraceisque Europæ. *F. capreolata* Lightf. scot. 380. Smith engl. bot. t. 943. Multò *F. officinali* affiniore. DC. syst. 2. p. 134.* (v. s.)

β. *prehensibilis*, (Kit! ind. h. Pesth. 1812. p. 10.) foliis subcrassis ferè succulentis brevioribus. An species propria?

6. *F. OFFICINALIS* (Lin! spec. 984.) siliculis globoso-retusis, pedicellis fructiferis erectis bractea duplò longioribus, racemis laxiusculis, caule erecto, foliis suprâ decompositis, lobis linearibus. ① in arvis, vineis orbis ferè totius. Smith engl. bot. t. 589. Flores rosei apice sanguinei. DC. syst. 2. p. 134.* (v. v.)

β. *grandiflora*, floribus majoribus dilutè purpureis. *F. officinalis* Barm! prod. fl. cap. p. 20.

7. *F. PARVIFLORA* (Lam! dict. 2. p. 567.*) siliculis globosis submucronulatis, pedicellis fructiferis erectis bractea longioribus, racemis laxis, caule diffuso, foliis suprâ decompositis, lobis linearibus canaliculatis. ① in agris oleraceis Angliæ, Galliæ, etc. Smith. engl. bot. t. 590.* Habitus *F. officinalis*, sed foliorum lacinie angustissimæ. DC. syst. 2. p. 136.* (v. v.)

8. *F. VAILLANTII* (Lois. not. p. 102.*) siliculis globosis vix mucronulatis, pedicellis fructiferis erectis bractea longioribus, racemis brevibus, caule erectiusculo, foliis suprâ decompositis, lobis linearibus planis. ① in agris arenosis Galliæ. — Vaill. bot. par. 56. t. 10. f. 6. Similis *F. parvifloræ* sed rami erecti, herba tota magis glauca. DC. syst. 2. p. 137. (v. v.)

9. *F. DENSIFLORA* (DC. cat. h. Monap. 113. syst. 2. p. 137.*) siliculis globosis, pedicellis fructiferis erectis bractea longioribus, racemis densis, calycibus dentatis, caule erectiusculo, foliis suprâ decompositis, lobis linearibus crassiusculis. ① in agris lapidosis et muris circa Teloneum et Monsapeliâ. (v. v.)

†. Species non satis nota.

10. *F. M'CRANTHA*. (Lag. cat. h. Madr. p. 21.) ① in Hispaniâ. Folia pinnato-decomposita, linearia, angustissima. Calyces cordato-rotundati, corollæ tubo latiores. (Lag.) DC. syst. 2. p. 137.

Ordo XI. CRUCIFERÆ. *Juss. gen.* 237. *DC. syst.* 2. p. 139.

Perigonium duplex, utrumque hypogynum liberum, partibus 4 cruciatis sæpius regularibus semper distinctis et liberis constans. Calycis sepala toro inserta, sæpius decidua, patentia aut erecta; 2 *placentaria* (superius nempe et inferius) placentis siliquæ opposita, in æstivatione exteriora, plana, sæpius angustiora; 2 *lateralia* seu *valvaria*, in æstivatione interiora, valvis fructûs opposita, latiora, basi concava aut deorsum gibba imò interdum calcarata, glandulis tunc è toro productis nectariferis et sepalorum basi imminetibus. Petala 4, sepalis alterna, sæpius unguiculata, limbo integro aut apice emarginato seu fisso, sæpius inter se æqualia, interdum 2 exteriora majora, rarissime abortu evanida, decidua. Stamina 6, rarissime et abortu 4 aut 2 evanida tetradynama, 2 ante sepala lateralialia solitaria breviora interdum dente intus aucta, 4 longiora geminata ante sepala placentaria, sæpissime inter se libera, interdum connata aut ad latus interius dentata. Antheræ biloculares, introrsæ. Torus glandulas virides callosas varias inter petala et genitalia gerens. Carpella duo arcuè in pistillum unicum coalita. Ovarium 1, nunc abbreviatum, nunc elongatum. Stylus 1 brevis ubi ovarium longum, elongatus ubi ovarium breve. Stigmata 2 nunc patentia, nunc approximata. Fructus siliquatus, nunc elongatus et tunc *siliqua* dicitur, nunc abbreviatus et tunc *silicula*, rarius 1-locularis evalvis, sæpius bilocularis bivalvis; loculis septo verticaliter sito separatis, valvulis planis concavis carinatisve longitudinaliter dehiscentibus aut rarius transversè dehiscentibus seu imò indehiscentibus; septo membranaceo, utrinquè nervo placentario circumdato, interdum striâ aut rimâ mediâ notato, latiusculo in fructibus valvulis planis donatis, angusto cum valvulis carinatis; fructus ideò compressus si septum latum et valvulæ planæ, depressus si septum angustum et valvulæ carinatæ, teres si septum lineare et valvulæ concavæ semi cylindricæ, quadrangulare si septum lineare et valvulæ angulo circiter recto plicato-carinatæ. Stylus fructum sæpius superans, nunc brevissimus ferè nullus, nunc filiformis, nunc plus minusve turgidus aut dilatatus et basi loculum 1-spermum indehiscentem fovens; inde fructus interdum bi aut abortu unilocularis, supernè loculo altero stylari coronatus. Semina in quoque loculo ad utramque placentam adfixa, sæpissime pendula, rarissime et forsàn tantum abortu solitaria, sæpius in quoque loculo 2, 4, etc., et ideò (exceptis

casibus accidentalibus ex abortu ortis) semper numero pari dispositis. Funiculi umbilicales sæpius liberi, interdum septo adnati. Spermoderma crassiusculum, extus ut videtur pelli-
culâ cinctum nunc adpressissimâ, nunc in alam membranaceam
expansâ, nunc per aquæ imbibitionem reticulatim mucilaginosâ
gelatinosâve factâ. Albumen nullum. Embryo oleosus, curva-
tus. Radicula teres, subconica, ad umbilicum tendens. Coty-
ledones 2 oppositæ, super radiculam diversis modis pronæ,
nempè cum radícula fissuræ cotyledonum adpressa est, tunc
cotyledones semper planæ *accumbentes* (o =) vocantur, cum
radicula dorso cotyledonum incumbit tunc cotyledones *in-*
cumbentes (o ||) dicuntur; cotyledones sunt aut planæ et tunc
strictiore sensu incumbentes dictæ, aut longitudinaliter pli-
catæ, apice sæpius emarginatæ radiculam in sinu foventes et
tunc *conduplicatæ* (o >>) dictæ, nunc lineares spiraliter convo-
lutæ et tunc *spirales* (o || ||) dictæ, nunc tandem lineares bis
transversè plicatæ et tunc *biplicatæ* (o || || ||) aut *bicrures*
dictæ; in embryonibus cotyledonibus accumbentibus donatis
radicula est *lateralis*, in omnibus aliis *dorsalis*. Embryo ipse
ferè semper in situ seminis naturali habet radiculam cotyle-
donesque adscendentes; dicitur *inversus* cum radícula cotyle-
donesque descendentes. Hi characteres etiamsi è partibus minu-
tissimis eruti, optimè videntur in seminibus, præcipuè antè
perfectam maturitatem aut ineunte germinatione. — Herbæ
annuæ aut perennes, sæpè biennes, aut rarius suffrutices ad
summum tripedales.

Subordo I. PLEURORHIZÆ. (o =) DC. *syst.* 2. p. 146.

Cotyledones planæ, accumbentes. Radicula lateralis. Semina
compressa.

Trib. I. ARABIDÆÆ. seu Pleurorhizæ Siliquosæ. DC. *syst.* 2.
p. 146.

Siliqua dehiscens, septo lineari seminibus plus minus latiore.
Semina ovalia, compressa, sæpè marginata. Cotyledones planæ,
accumbentes, septo parallelæ.

I. MATHIOLA. Brown h. Kew. ed. 2. v. 4. p. 119. (non Lin.)

Siliqua teretiuscula. Stigmata conniventia, dorso incrassata aut
cornigera. Calyx basi bisaccatus. Semina compressa, 1-serialia, sæpius
marginata. Cotyledones planæ.

Sect. I. PACHYNOTUM. DC. *syst.* 2. p. 163.

Petala obovata. Stigmatis dorsa crassa non cornigera. Siliqua idè apice cras-
siuscula non tricuspidata.

1. *M. INCANA* (Brown in h. kew. ed. 2. v. 4. p. 119.) caule basi suffruticoso

- erecto ramoso, foliis lanceolatis integerrimis incanis, siliquis subcylindricis eglandulosis. ♂ ♀ in maritimis regionis Mediterraneæ. *Cheiranthus incanus* Lin! spec. 924. — Weinm. phyt. t. 643. f. a-e et 644. f. a. b. Variat flore coccin. alb. et varieg., simpl. et plen. — DC. syst. 2. p. 163.* (v. v.)
2. *M. ANNUA* (Sweet h. suburb. 147.) caule herbaceo erecto ramoso, foliis lanceolatis obtusis incanis, siliquis subcylindricis eglandulosis. ① in maritimis Europæ australis. *Cheiranthus incanus* Lin. spec. 925. — Bauh. hist. 2. p. 875. f. 1. Cl. Brown hanc, forsân rectissime, habet pro *M. incanæ* varietate. — DC. syst. 2. p. 165.* (v. v.)
 3. *M. GLABRATA* (DC. syst. 2. p. 165.*) caule suffruticoso erecto ramoso, foliis lanceolatis glabris, siliquis subcompressis eglandulosis. ♂ hab. . . . *M. incana* Brown in h. kew. ed. 2. v. 4. p. 119. *Leucojum album* Blakw. herb. t. 181. Variat flore purpureo et albo. (v. v.)
 4. *M. GRÆCA* (Sweet hort. suburb. 147.) caule herbaceo erecto ramoso, foliis lanceolatis glabris, siliquis subcompressis eglandulosis. ① in Oriente, in Græciâ. *Cheiranthus Græcus* Juss! in Pers. ench. 2. p. 201. An hæc quatuor stirpes ut unius speciei primitivæ varietates habendæ? — DC. syst. 2. p. 166.* (v. v.)
 5. *M. FENESTRALIS* (Brown in h. kew. ed. 2. v. 4. p. 119.) caule suffrutescente erecto indiviso, foliis confertis obovatis tomentosis, siliquis pubescentibus eglandulosis basi sublatis. ♂ ② in rupibus maritimis Cretæ. *Cheiranthus fenestralis* Lin. f. dec. 31. t. 16.* Flores coccinei aut purpurei, paulò minores quam *M. incanæ* — DC. syst. 2. p. 166.* (v. s.)
 6. *M. ELLIPTICA* (Brown! in Salt. voy. abyss. app. p. LXV.) caule suffrutescente tortuoso ramoso, foliis petiolatis ellipticis incanis, siliquis teretibus tomentosis. ♀ ⑤ in Abyssiniâ. Flores odori, à *M. incanâ* parùm diversi. DC. syst. 2. p. 167.* (v. s. in h. Banks.)
 7. *M. SINUATA* (Brown in h. kew. ed. 2. v. 4. p. 120.) caule erectiusculo herbaceo ramoso, foliis oblongis tomentosis, inferioribus sinuatis, siliquis compressis velutinis et glandulose muricatis. ② in arenosis maritimis Europæ. *Cheiranthus sinuatus* Lin! spec. 926. Smith engl. bot. t. 462. Flores ferè *M. incanæ*, vesperè odorati. DC. syst. 2. p. 167.* (v. v.)
 8. *M. ACAULIS* (DC. syst. 2. p. 168.*) caule subnullo, foliis linearibus sinuato-dentatis pube stellatâ tomentosis, floribus radicalibus. ① in Egypto. *Cheiranthus acaulis* Balb! in litt. Spreng. nov. prov. p. 10. n. 19.* Siliquæ ignotæ. Planta non confundenda cum *Hesperide pygmæâ*. (v. s. comm. à cl. Balb.)
 9. *M. PERSICA* (DC. syst. 2. p. 168.*) caule erecto imâ basi suffrutescente, apice pube velutinâ et glandulosâ cano, foliis canescentibus, radicalibus oblongis integris petiolatis, siliquis pubescentibus eglandulosis. ♀ in Persiâ. *Cheiranthus Persicus* Pall! in h. Lamb. Habitus *Mathiolæ* sed color floris flavus. (v. s. in h. Lamb.)

Sect. II. LUPERIA. DC. syst. 2. p. 169.

- Stigmatis dorsa crassa non cornigera, petala oblonga sordidè è flavo purpur.
10. *M. TENELLA* (DC. syst. 2. p. 169.*) caule herbaceo erecto subsimplici, foliis oblongis sinuato-dentatis basi attenuatis pube stellatâ canescentibus, siliquis pubescentibus eglandulosis. ① in insulâ Cypro. Flores colore ferè *Math. tristis* sed pallidi. (v. s.)
 11. *M. TORULOSA* (DC. syst. 2. p. 169.*) caule erecto subramoso flocculoso-scabro, foliis linearibus subintegris tomentosis, siliquis subtorulosis pube velutinis et glanduloso-scabris. — ad cap Bonæ-Spei. *Cheiranth. torulosus* Thunb. prod. 108. Habitum accedit ad *Hesperidem*. Flores parvi. (v. s. in h. Banks.)
 12. *M. TATARICA* (DC. syst. 2. p. 170.*) caule erecto subsimplici glabrato, foliis cano-tomentosis irregulariter runcinato-dentatis, siliquis glaberrimis teretiusculis subtorulosis. ♀ in saxis montium gypsaceorum Tataricæ australis.

Hesperis Tatarica. Pall! itin. 1. app. 117. t. O.* Flores è livido purpureo-flavescentes. (v. s. ex h. Pall. et Fisch.)

13. *M. ODORATISSIMA* (Brown in h. kew. ed. 2. v. 4. p. 120.) caule erecto ramoso, foliis tomentosis pubescentibusve dentatis pinnatifidisve, siliquis compressis puberulis. 2½ in montibus calcareis Tauriæ et saxosis Caucasi orientalis. Sims bot. mag. t. 1711. *Hesperis odoratissima*. Poir. suppl. 3. p. 195. Flores ochroleuco-sordidi aut senescentes fusco-subpurpurascens. DC. syst. 2. p. 170. (v. v. et s.)

β. *Tanaicensis*, pube adpressâ, siliquis dimidio brevioribus, Fisch! in litt. *Cheiranthus fragrans*. Fisch! cat. h. Gor. 1812. p. 51.

14. *M. VARIA* (DC. syst. 2. p. 171.*) caule erecto subsimplici nudiusculo, foliis linearibus obtusis integerrimis incanis, floribus subsessilibus, petalis ovalibus undatis, siliquis compressis. 2½ in Oriente. *Cheiranthus varius* Sibth. et Smith fl. græc. t. 636. Flores ferè *M. tristis* sed majores. (v. s.)

β. *Cheiranthus tristis* var. *Sabauda*. All! fl. ped. n. 991. *Cheir. Vallesiacus* Gay! ined. in h. Gaud. — in Vallesiâ.

15. *M. TRISTIS* (Brown in h. kew. ed. 2. v. 4. p. 120.) caule basi suffruticoso ramoso erecto, foliis tomentosis linearibus integris dentatisve, floribus subsessilibus, petalis oblongis, siliquis teretiusculis. 2½ 5 in apricis lapidosisque Europæ australioris. *Cheiranthus tristis*. Lin! spec. 925. — Barr. ic. t. 803. Flores sordide flavi aut viridi-rufescentes. — DC. syst. 2. p. 172.* (v. v.)

Sect. III. PINARIA. DC. syst. 2. p. 172.

Stigmatis dorsa in cornua exserta. Petala oblonga sordide è flavo purpurascens.

16. *M. CORONOPIFOLIA* (DC. syst. 2. p. 173.*) caule erecto à basi ramosissimo, foliis linearibus dentato-pinnatifidis incanis, siliquis subtorulosis breviter tricuspidatis eglandulosis. 2½ in Sicilia. *Cheiranthus coronopifolius* Sibth. et Sm. fl. græc. t. 637, ex prod. 2. p. 25. Petala vinoso-purpurea.

β. *Hispanica*, *Cheiranthus parviflorus* Thibaud! ined. — in Hispaniâ. An species propria? (v. s.)

17. *M. OXYCERAS* (DC. syst. 2. p. 173.*) caule erecto ramoso glanduloso, foliis velutinis sinuato-repandis, superioribus integris, siliquis teretibus subvelutinis glandulosisque tricuspidatis, cuspidibus lateralibus subreflexis. ① Deless! ic. sel. 2. t. 11. Aff. *M. lividæ*. Flores sessiles. (v. s.)

α. caule à basi ad apicem glanduloso. Circa Damascum.

β. caule basi tantum subglanduloso. In Persiâ.

γ. foliis inferioribus pinnatifidis. Inter Alep. et Mossul.

18. *M. LIVIDA* (DC. syst. 2. p. 174.*) caule subdiffuso ramosissimo, foliis oblongo-linearibus infimis sinuatis, tomento velutino glandulis interspersis, siliquis teretiusculis pubescentibus glandulosisque tricuspidatis, cuspidibus adscendentibus stigmate longioribus. ① in Ægypto. *Cheiranthus tristis* Forsk. fl. æg. ar. p. 119.* Deless! ic. sel. 2. t. 12. (v. s.)

19. *M. LONGIPETALA* (DC. syst. 2. p. 174.*) caule subdiffuso ramoso subpubente, foliis oblongis sinuato-dentatis, ovarii teretibus pubentibus eglandulosis tricuspidatis, petalorum laminis calyce longioribus. ① in Oriente circa Bagdad. *Cheiranthus longipetalus* Vent! cels. t. 93.* Petalorum laminæ basi flavescentes, apice purpurascens. (v. s. et v. in h. al. Cels.)

Sect. IV. ACINOTUM. DC. syst. 2. p. 175.

Petala obovata obtusa aut emarginata. Siliqua apice tricuspidata, stigmatum dorsis nempe in cornua exsertis. Semina immarginata (ex cl. Adrz. qui hanc sect. pro genere proprio habet.)

20. *M. TRICUSPIDATA* (Brown in h. kew. ed. 2. v. 4. p. 120.) caule suberecto ramoso, foliis sinuato-pinnatifidis, siliqua cuspidibus 3 acutis subæqualibus. ① in arenis maritimis secus mare Mediterraneum. *Cheiranthus tricuspidatus*

- Lin! spec. 926. Schk. hand! 2. p. 250. n. 1846. t. 184. Flores ferè M. sinuatae. Semina ovata compressa. An ab hac satis differt *Cheiranthus humilis* Spreng. neu. ent. 3. p. 164? DC. syst. 2. p. 175.* (v. v.)
21. *M. PARVIFLORA* (Brown in h. kew. ed. 2. v. 4. p. 121.) caule suberecto ramoso, foliis lanceolatis tomentosis repando-dentatis, floribus sessilibus, siliquis teretibus tricuspidatis, cuspidibus acutis mediâ obtusâ longioribus. ① in Hispaniâ meridionali. *Cheiranthus parviflorus* Schomb. in Schrad. journ. 3. p. 369. Flores similes *M. tricuspidatæ* sed dimidio minores. DC. syst. 2. p. 176.* (v. s. et v.)
22. *M. LUFATA* (DC. syst. 2. p. 176.*) caule suberecto ramoso, foliis oblongis repando-dentatis pubescentibus, pedicellis brevibus, siliquis teretibus tricuspidatis, cuspidibus subincurvis mediâ vix ullâ multò longioribus. ① in Hispaniâ. Media videtur inter *M. tricuspidatam* et *parvifloram*. Flores *M. sinuatae*. (v. s. in h. Mus. Par.)
23. *M. PUMILIO* (DC. syst. 2. p. 177.*) caule brevissimo, foliis pinnatifido-sinuatis incanis, siliquis tricuspidatis, cuspidibus obtusis stigmate vix exserto longioribus. ① in insulâ Rhodo. *Cheiranthus pumilio* Sibth. et Sm. fl. græc. t. 638, ex prod. 2. p. 26. Flores pauci, magnitudine ferè *M. sinuatae*. (v. s. in h. Banks sine fructu.)
24. *M. HUMILIS* (DC. syst. 2. p. 177.*) caule brevissimo, foliis oblongis subincanis sinuato-dentatis, floribus subsessilibus, siliquis subteretibus adultis glabris tricuspidatis, cuspidibus brevissimis. ① in Ægypto circa Rosettam. *Cheiranthus* nov. sp. è Rosetta Clark! voy. app. 4. p. 718. Flores subdistantes, ferè sessiles. (v. s. in h. Lamb.)

†. *Mathiolæ non satis notæ.*

25. *M. BICORNIS* (DC. syst. 2. p. 177.) ① vel. ② in Græciâ? *Cheiranthus bicornis* Sibth. et Smith prod. 2. p. 26. Ad sectionem tertiam referenda videtur.
26. *M. CRUCIGERA* (DC. syst. 2. p. 177.*) 2℥ in montibus Siciliæ di Madonia et di Castelbuono dictis. — Bocc. mus. p. 148. t. 111. Flores violacei. Siliquæ erectæ apice tricuspidatæ.
27. *M. RUPESTRIS* (DC. syst. 2. p. 714.) 3 in montibus Siciliæ. Hesperis rupestris. Raf. specch. 2. p. 46. Affinis *M. incanæ*, sed folia obtusa et petiolata, petala emarginata, calyx evidentiùs gibbus. †.
28. *M. FASCICULATA* (DC. l. c.) 5 in maritimis Siciliæ. Cap. panph. 1. t. 144. ed. Raf. t. 14. f. 2. Hesperis fasciculata Raf. specch. 2. p. 7. Affinis *M. tricuspidatæ*, sed folia fasciculata sessilia linearia integra acuta linearia. An *M. crucigera*? †.

II. *CHEIRANTHUS*. Brown in h. kew. ed. 2. v. 4. p. 118.

Siliqua teres aut compressa. Stigma bilobum aut capitatum. Calyx basi bisaccatus. Semina 1-serialia ovata compressa (o=). DC. syst. 2. p. 178.

Sect. I. *CHEIRI*.

Stylus subnullus. Semina immarginata.

1. *C. CHEIRI* (Lin! spec. 924.) foliis lanceolatis integerrimis, pube bipartiâ adpressâ aut nullâ, siliquis linearibus, stigmatis lobis recurvis. 2℥ interdum ② aut 5 evadens. In muris tectis et lapideis Europæ. Schk. handb. 2. n. 1840. t. 184. — Var. *α*. quæ etiam sylvestris reperitur, forsan tamen ex hortis migrata. Var. *β*. *grandiflorus*, *γ*. *maximus*, *δ*. *serratus*, *ε*. *patulus*, *ζ*. *ferrugineus*, *θ*. *varius*, *ι*. *flavescens*, *κ*. *thyrsoides*, *μ*. *fruticulosus*, sunt variationes hortenses parùm notabiles, *λ* *gynantherus*, antheris nempe in carpellâ mutatis.
2. *C. ALPINUS* (Lin. mant. 93. exclus. syn.) foliis lanceolatis subdentatis stellato-pubescentibus, caule simplici stricto, siliquis patulis pedicello sexies

longioribus, seminibus appendiculatis. \mathcal{L} in Norvegiâ et Lapponiâ. Wahlenb. fl. lap. n. 333. t. 12. f. 1.* DC. syst. 2. p. 181.*

3. *C. OCHROLEUCUS* (Hall. fil. ex Schleich. cat. p. 16.) foliis oblongo-lanceolatis subdentatis, pube bipartitâ aut nullâ, caule decumbente ramoso, petalis obovatis, siliquis erectis stylo pedicelli longitudine apiculatis. \mathcal{L} in Jurassi montibus. *C. dubius*. Sut. fl. helv. 2. p. 65. — Hall. hist. n. 449. t. 14.* Flores subodorati. DC. syst. 2. p. 181.* (v. v.)

Sect. II. CHEIROIDES.

Stylus filiformis. Semina marginata. Siliqua tetragona. — Psilostylis Andr.
in litt.

4. *C. TENUIFOLIUS* (L'Her! st. nov. p. 92.) foliis linearibus integerrimis subsericeis, caule frutescente ramoso. \mathfrak{J} in Maderâ. Willd. spec. 3. p. 517.* Flores flavi, stylus filiformis 2 lin. longus. DC. syst. 2. p. 183.* (v. s.)
5. *C. MUTABILIS* (L'Her! st. nov. 1. p. 92.) foliis lineari-lanceolatis acuminatis argutè serratis pube adpressâ bipartitâ subpubescentibus, caule frutescente ramoso: \mathfrak{J} in Maderâ. Curt. bot. mag. t. 195.* Flores versicolores, nascentes ochroleuci. DC. syst. 2. p. 183.* (v. s. et v.)
- β . floribus nascentibus albis. *C. longifolius* Vent! Malm. t. 83.* — in Teneriffâ.
6. *C. SCOPARIUS*. (Willd. enum. p. 681.) foliis lineari-lanceolatis acuminatis integris pube adpressâ bipartitâ subpubescentibus, caule fruticoso ramoso. \mathfrak{J} in Teneriffâ. Hesperis cinerea Poir. suppl. 3. p. 196.* Aff. *C. mutabilis* sed evidenter differt. — DC. syst. 2. p. 184.* (v. s. et v. c.)
- α . *purpurascens*, flores nascentes albi, adulti purpurei.
- β . *ærginosus*, fl. nascentes ærginosi, adulti pallidiores levissimâ rubedine suffusi.
- γ . *chamæleo*, fl. nascentes citrini, dein sordidè armeniacei. *C. Cheiri* var. *Chamæleon* Ker. bot. reg. t. 219.*
7. *C. SEMPAFLORANS* (Schousb. maroc. ed. germ. p. 181.) foliis lineari-lanceolatis integerrimis subsabris, caule fruticoso ramoso, siliquis compressis, pedicellis calyce dimidio brevioribus. \mathfrak{J} in regno Maroccano. Petala alba, unguibus calycis longitudine. DC. syst. 2. p. 184.* (v. s. et v.)
- β . *C. linearis* Vent! malm. p. 83. in adn. non Thunb. *C. frutescens* Pers. ench. 2. p. 201. Dignoscitur caule magis frutescente, foliis brevioribus angustioribus.
8. *C. LINIFOLIUS* (Pers. ench. 2. p. 201.) foliis linearibus integerrimis scabris confertis, caule fruticuloso ramoso, siliquis teretiusculis calyce triplò longioribus, pedicellis calyce duplò brevioribus. \mathfrak{J} in Hispaniâ. Hesperis lini-folia Desf! cat. h. par. ed. 1. p. 129. DC. syst. 2. p. 185.* (v. s.)

†. Species non satis notæ.

9. *C?* *SYRIACUS* (DC. syst. 2. p. 185.*) foliis ovato-oblongis cordato-auriculatis dentatis glabris, caule erecto flexuoso ramoso glabro. (2) in Syriâ. — Moris. hist. 2. p. 282. s. 3. t. 10. f. 4. Flores non vidi. (v. s. in h. C. Bauh.)
10. *C?* *PULCHELLUS* (Willd. spec. 3. p. 523.*) foliis lineari-lanceolatis acutè dentatis pube bipartitâ subpubescentibus. \mathcal{L} in Cappadociâ. Planta pygmæa. Petalorum limbi obovati flavi magnitudine *Erysimi* Helvetici. Fructus ignotus. DC. syst. 2. p. 185.* (v. s. in h. Banks et Mus. Par.)
11. *C?* *SALINUS* (Lin. Mant. 93.) foliis lanceolatis obtusis integerrimis, caule erecto, antheris inclusis. \mathfrak{J} ad Sibiriz, Tatariz salinas. Hesperis salina Lam. dict. 3. p. 324. Simillimus *Mathiolæ* incanæ sed octies minor. DC. syst. 2. p. 186.*
12. *C. PALLASII* (Pursh! fl. bor. am. 2. p. 436.) foliis lanceolato-linearibus attenuatis repando-dentatis glabriusculis, caule simpliciter erecto tereti. (2) in

Americâ bor. Flores ferè Mathiolæ incanæ. Siliqua teretiuscula stigmatè parvo subcapitato. DC. syst. 2. p. 186 * (v. s. in h. Pallas nunc Lamb.)

13. *C. PYGMÆUS* (Adams in mem. soc. mosc. 5. p. 114 *) foliis lineari-lanceolatis integerrimis subhispidis, caule simplicissimo adscendente, racemo corymboso, siliquis longissimis subtetragonis, stigmatè bilobo. 2/ in Sibiria. A. C. Alpino differt foliis hispidulis lineari-lanceolatis, floribus violaceis nec sulfureis. DC. syst. 2. p. 187*. An potiùs *Erysimi* species?

14. *C?* *FLEXUOSUS* (Sibth. et Smith fl. græc. t. 634. ex prodr. 2. p. 24.) ① in insulâ Cypri. Folia obovato-subrotunda. Caulis diffusus, flexuosus. Siliquæ patentissimæ, rigidæ, pungentes. DC. syst. 2. p. 187.

III. *NASTURTIIUM*. Brown in h. kew. ed. 2. v. 4. p. 109. non Hall. nec Mærch.

Siliqua teretiuscula, abbreviata aut declinata. Stigma subbilobum. Calyx basi æqualis, patens. Semina parva, irregulariter biserialia, immarginata (o=).

Sect. I. *CARDAMINUM*. DC. syst. 2. p. 188.

Petala alba calyce majora. Siliquæ teretiusculæ. Glandulæ 4 ad basim staminum. Semina reticulato-rugosa, cotyled. obliquè incumbentibus. (Andrz. in litt.) An genus proprium?

1. *N. OFFICINALE* (Brown in h. kew. ed. 2. v. 4. p. 110.) foliis pinnatisectis, segmentis ovatis subcordatis repandis. 2/ in rivulis orbis ferè totius. *Sisymbrium nasturtium*. Lin! spec. 916. Smith engl. bot. t. 855. DC. syst. 2. p. 188.* (v. v.)

β. *præcocius*, Early water-cress. Pet. h. brit. t. 47. f. 3.

γ. *Chilense*. Dubia tetradynamia siliquosa Ruiz. et Pav! in h. Lamb.

Sect. II. *BRACHYLOBOS*. DC. syst. 2. p. 190.

Petala flava. Siliquæ teretiusculæ aut ellipsoideæ. Glandulæ receptaculi parvæ.

2. *N. SYLVESTRE* (Brown in h. Kew. ed. 2. v. 4. p. 110.) foliis pinnatisectis, segmentis lanceolatis serratis incisive. 2/ in Europâ, Asiâ boreali, Persiâ, Tauriâ, Chinâ, Novâ-Hollandiâ et Americâ. *Sisymbrium sylvestre*. Lin. spec. 916. Smith engl. bot. t. 2324. Petala aurea calyce longiora. DC. syst. 2. p. 190* (v. v.)

3. *N. ANCEPS*, foliis pinnatifidis, auriculis amplexicaulibus glaberrimis, caulibus repentibus, petalis calyce majoribus, siliquis utrinquè acuminatis ancipitibus. 2/ in ripis suprâ aquam et magis arenosis circa Upsaliam. *Sisymbrium anceps*. Wahlenb. fl. ups. p. 223. *S. amphibium terrestre* Ehrh. Beitr. 5. p. 22. Lin. spec. ed. 2. p. 917. — Fl. dan. t. 984.

4. *N. PALUSTRE* (DC. syst. 2. p. 191.*) foliis pinnati-lobatis, auriculis amplexicaulibus ciliatis, lobis confluentibus dentatis glabris, radice fusiformi, petalis calycem æquantibus, siliquis utrinquè obtusis subiurgidis. ① in inundatis Europæ totius, Persiæ, Chinæ et Americæ bor. *Sisymbrium palustre* Lays. fl. hal. n. 679. — engl. bot. t. 1747. Flores exigui, flavi. (v. v.)

β. *pusillum*, *Myagrum pumilum* Lam. dict. 1. p. 572.* — Vill. dauph. 3. p. 341. t. 39.*

γ. *barbareæfolium*, *Brachylobus barbareæfolius* Desv. journ. 1814. v. 3. p. 170. — in Ægypto circa Rosettam.

δ? *tanacetifolium*, *Sisymbrium tanacetifolium*. Walt. fl. Car. ex Bosc. — In Carolinâ.

ε? *brevipes*, à prioribus differt pedicellis brevissimis vix fructiferis lineam longis et stigmatè ferè sessili. — In Porto-Ricco. (Cl. Bertero). An species propriâ?

5. *N. MICRANTHUM*, pubescens, foliis pinnatisectis, segmentis oppositis incisolacinatis, folio terminali subrotundo-ovato petalis calyce majoribus, siliquis

- erectis lineari-oblongis glabris. — In Indiâ orient. *Sisymbrium micranthum* Roth. nov. pl. spec. 324.* Aff. N. palustri.
6. N. MADAGASCARIENSE (DC. syst. 2. p. 192.*) foliis pinnatisectis, segmentis petiolulatis pinnati-lobatis, lobis ovatis dentatis, terminalibus majoribus. (Y) ? in Orâ orientali insulæ Madagascar. Flores minimi non satis noti. (v. s. in h. Mus. Par.)
7. N. MEXICANUM. (Mocq. Sess. et Cerv. fl. Mexic. ic. ined. et DC. syst. 2. p. 193.*) foliis pinnatifidis glabris, lobis oblongis sinuato-dentatis obtusis, petalis calyce æqualibus, siliquis declinatis subortis pedicello triplò longioribus. — In Mexico. An satis à N. palustre distinctum ? differre videtur præcipue siliquis teretiusculis duplò ferè longioribus apice tortis seu curvulis. (v. ic. pict.)
8. N. BONARIENSE (DC. syst. 2. p. 193.*) foliis pinnatipartitis, lobis distantibus linearibus subdentatis glabris, siliquis suberectis tereti-compressis pedicello duplò triplòve longioribus. (I) in Bonariâ. *Sisymbrium Bonariense* Poir! dict. 7. p. 205.* Flores flavi. (v. s. in h. Mus. Par. et Juss.)
9. N. CERATOPHYLLUM (DC. syst. 2. p. 193.*) foliis linearibus pinnatifido-dentatis glabriusculis, siliquis pedicello brevioribus. (I) in arenosis Mauritanie. *Sisymbrium ceratophyllum* Desf! fl. atl. 2. p. 82. t. 155.* Flores flavi. Petala integra calyce duplò longiora. (v. s. in h. Desf.)
10. N. CORONOPHYLLUM (DC. syst. 2. p. 194.*) foliis lanceolatis pinnatifido-dentatis pubescentibus, caule subnudo adscendente, siliquis linearibus incurvatis pedicelli longitudine. — in Africa bor. *Sisymbrium coronopifolium*. Desf! fl. atl. 2. p. 82. t. 154.* Folia inferiora ferè Plantaginis Coronopi. (v. s. in h. Desf.)
11. N. BURSIFOLIUM (DC. syst. 2. p. 194.*) foliis villosis, radicalibus inciso-pinnatifidis, caulinis sagittatis integris acuminatis, caule pedicellisq. hispidis erectis. (I) in Kamtschatka. *Sisymbrium bursifolium* Patr! herb. Petala calyce majora, in sicco ochroleuca. (v. s.)
12. N. SAGITTATUM (Brown in h. Kew. ed. 2. v. 4. p. 111.) pubescens, foliis radicalibus dentato-runcinatis, caulinis sagittatis oblongis obtusis, caulibus erectis à basi ramosis, siliquis deflexis. (Y) in Sibiria? ad mare Caspium. *Sisymbrium molle* Jacq. ic. rar. 1. t. 122.* Flores pallidè flavi. (v. s.)
13. N. GLAUCOPHYLLUM (DC. syst. 2. p. 195.*) glaberrimum, foliis radicalibus petiolatis obovato-orbiculatis crenatis coriaceis, summis linearibus squamiformibus. (Y) ? in Persiâ propè Teheran. Deless! ic. sel. 2. t. 14. An potius ad sequentem sectionem ? species in utrâque distinctissima (v. s. in h. Mus. Par.)
14. N. LIPPIZENSE (DC. syst. 2. p. 195.*) foliis radicalibus petiolatis obovatis dentatis sublyratisve, superioribus pinnatipartitis, lobis linearibus integris, siliquis linearibus declinatis. (Y) in locis rupestribus Croatiae et Dalmatiae. *Sisymbrium Lippizense* Wulf. in Jacq. coll. 2. p. 161.* ic. rar. 3. t. 505. Species media inter N. Pyrenaicum et sylvestre. (v. s.)
15. N. PYRÆNICUM (Brown in h. Kew. ed. 2. v. 4. p. 110.) foliis radicalibus petiolatis obovatis lyratisve, caulinis amplexicaulibus pinnatipartitis, lobis linearibus integris, siliculis ovalibus stylo apiculatis. (Y) in pascuis siccis rupium fissuris collibus Hispaniæ Pyrenæicæ etc. *Sisymbrium Pyrenæicum* Lin. spec. 916. — All. fl. pedem. n. 1013. t. 18. f. 1. DC. syst. 2. p. 196.* (v. v.)
- β. *Brachylobus Domingensis* Desv. journ. 1814. v. 3. p. 183. *Sisymbrium Domingense* Poir. suppl. 5. p. 161. — in Hispaniolâ. (v. s. in h. Juss.)
16. N. AMPHIBIUM (Brown in h. Kew. ed. 2. v. 4. p. 110.) foliis oblongo-lanceolatis pinnatifidis serratisve, radice fibrosâ, petalis calyce majoribus, siliculis ellipsoideis. (Y) in aquosis Europæ ferè totius. *Sisymbrium amphibium* Lin! spec. 917. Smith! engl. bot. t. 1840.
- α. *indivisum*, foliis omnibus integriusculis serratisve, non aut vix basi auriculatis. *Sisymbrium aquaticum* Gars. fig. t. 549.

β. *varifolium*, foliis aliis serratis, aliis pectinato-pinnatifidis, aliis capillaceo-multifidis. — J. Bauh. hist. 2. p. 867. f. 2.

γ. *auriculatum*, foliis indivisis basi auriculatis. *Sisymbrium stoloniferum*. Presl. fl. cech. p. 137.

17. N. *NATANS* (DC. syst. 2. p. 198.*) foliis emersis oblongo-linearibus integris, immersis multifido capillaceis, petalis calyce vix longioribus, siliculis obovato-globosis. 2♂ in Sibiria. Deless. ic. sel. 2. t. 15. *Myagrum natans* Patr! herb. ined. Flores flavi minores quàm N. *Amphibii* (v. s.)

Sect. III. CLANDESTINARIA. DC. syst. 2. p. 198.

Petala nulla aut minima alba. Siliquæ teretiusculæ. — Sectio dubia constans speciebus quoad caracteres genericos non satis notis, forsitan aliis ad *Sisymbrium*, aliis ad *Arabidem* referendis.

18. N. *BENGHALENSIS* (DC. syst. 2. p. 198.*) foliis obovato-cuneatis apice dentatis, siliquis teretiusculis subturgidis, pedicellis bracteatis siliculâ paulò brevioribus. ①? in Bengalo. *Sinapis Benghalensis* Roxb! ined. in h. Lamb. Species et sequens ex dispositione florum Kiberæ (*Sisymbrii* sect. iv.) affines, sed differunt siliculâ brevi, cotyledonibus accumbentibus. (v. s.)

19. N. *DIFFUSUM* foliis glabris petiolatis ovali-oblongis dentatis, inferioribus subpinnatifidis, siliquis teretibus pedicello triplo longioribus stylo distinctis, pedicellis aliis bracteatis aliis nudis. ① in Javâ. Caules multiplices diffusi pedicelli 3 lin. longi. (v. s. in h. Coulon.)

20. N. *MICROSPERMUM* (DC. syst. 2. p. 199.*) foliis glabris, radicalibus petiolatis pinnatipartitis, caulinis amplexicaulibus inciso-serratis, siliquis teretiusculis, pedicellis bracteatis brevissimis. ① in Chinâ. Flores in bractearum axillis sessiles, solitarii, albi, minimi. (v. s. in h. Banks.)

21. N. *CLANDESTINUM* (Spreng! nov. prov. p. 29. n. 63.†) foliis pinnatipartis, lobis cordato-subrotundis petiolulatis repandis, terminali maximo, siliquis teretiusculis elongatis. ① in Brasiliâ. Flores apetalis inconspicui. DC. syst. 2. p. 199.* (v. s. ex h. Spreng.)

22. N. *INDICUM* (DC. syst. 2. p. 199.*) foliis ovato-lanceolatis runcinato-dentatis utriusque acuminatis glabris, siliquis teretiusculis pedicello quadruplo longioribus. ① in Indiâ, in Chinâ. *Sisymbrium Indicum* Lini! mant. 93.* (v. s.)

β. *Sisymbrium Sinapis* Burm! fl. ind. 140. excl. syn. — in Javâ.

γ. *S. apetalum* Desf! cat. h. par. 1804. p. 130. non Lour. *S. dubium* Pers. ench. 2. p. 199.

23. N. *APETALUM* (DC. syst. 2. p. 200.*) foliis tomentosis ad basin hastatis superne pinnatifidis, siliquis teretiusculis declinatis. — in Cochinchinâ. *Sisymbrium apetalum* Lour. fl. coch. ed Willd. 2. p. 456.* †.

24. N? *ARABIFORME* (DC. syst. 2. p. 220.*) foliis glabris obtusè sagittatis oblongo-lanceolatis acuminatis integris, siliquis teretiusculis adscendentibus, petalia oblongis calyce longioribus. — in Americâ calidiori. *Arabis resediflora* H. B. et Kl! nov. spec. am. 5. p. 81.* Petala alba, oblongo-linearia, basi attenuata calyce duplo longiora.

†. *Species non satis notæ.*

25. N. *NEBRODENSE* (Raf. in Desv. Journ. 1814. v. 2. p. 270.) in Siciliæ montibus Nebrodensibus. *Sisymbrium Nebrodense* Poir. suppl. 5. p. 161. Folia radicalia petiolata, oblonga, cuneiformia; caulina sessilia, ovata dentata; siliquæ ovatis, oblonga. DC. syst. 2. p. 201.

26. N. *HISPIDUM* (DC. syst. 2. p. 201.) in Pensylvaniâ. *Sisymbrium hispidum* Poir. suppl. 5. p. 161. non Vahl nec Mœnch. Folia subvillosa, pinnatifido-runcinata, lobis dentatis, dentibus acutis. Siliculæ breves, ellipticæ.

27. N. *ATROVIRENS* (DC. syst. 2. p. 201.) ① in Chinâ. *Sisymbrium atrovirens* Horn. h. bafn. suppl. 72. Habitus N. *amphibii*. Folia inferiora lyrato-pinnatifida, superiora ovato-lanceolata repando-dentata. Siliquæ arcuatæ.

IV. LEPTOCARPÆA. DC. syst. 2. p. 201.

Siliqua teretiuscula, gracillima, axi ferè parallela. Stigma sessile, bilobum. Calyx patens, æqualis. Semina minima uni (aut forsân biserialia.) — Flores flavi inodori. Cotyledones forsân incumbentes et tunc propè *Sisymbrium* genus collocandum?

1. L. LÖESELII (DC. syst. 2. p. 202. *) ① in aggeribus muris mœnibus tectis Germaniæ, etc. *Sisymbrium Lœselii*. Lin! spec. 921. excl. Bauh. syn. Jacq. fl. aust. t. 324. * (v. s.)

V. NOTOCERAS. Brown in h. Kew. ed. 1812. v. 4. p. 117. DC. syst. 2. p. 202. *

Siliqua tetragono anceps, valvis apice in cornu aut mucronem exsertis. Semina ovalia compressa. — Herbæ annuæ, parvæ.

Sect. I. DICERATIUM. Lag. el. hort. Madr. 1815. p. 20.

Siliquæ dehiscentes bicornes; semina compressa; cotyledones septo parallelæ. — Flores lutei, minimi. Folia integræ. Pili strigosi bipartiti adpressi.

1. N. CANARIENSE (Brown in h. Kew. ed. 2. v. 4. p. 117.) Siliquis bicornibus, petalis æqualibus, foliis integris, pilis strigosis centro adfixis bipartitis adpressis sparsis. ① in insulis Canariensibus. Jacq. f. eclog. t. III. optim. *Erysimum bicorne* Ait! h. Kew. ed. 1. v. 2. p. 391. * Semina in quoque loculo 4 suborbiculata, compressa. DC. syst. 2. p. 203. * (v. s.)
2. N. HISPANICUM (DC. syst. 2. p. 204.) siliquis bicornibus, petalis inæqualibus, foliis integris, pilis strigosis centro adfixis bipartitis confertissimis. ① in locis ruderalis arenoso calcareis Orcelis Muriæ et Granatensis regni. *Diceratium prostratum*. Lag! el. hort. Madr. 1815. p. 20. * Deless! ic. sel. 2. t. 17. Priori nimis affinis; differre tamen videtur habitu minore rigidioris magis prostrato, etc. (v. s.)

Sect. II. TETRACERATIUM. DC. syst. 2. p. 204.

Siliquæ quadricornutæ. — Flores minimi, forsân apetal. Folia sinuato-dentata. Pubescentia mollis ramosa.

3. N. QUADRICORNE (DC. syst. 2. p. 204. *) siliquis quadricornibus, foliis sinuato-dentatis, pube molli ramosâ. ① in Sibiria. *Erysimum quadricorne* Steph. in Willd. spec. 3. p. 514. * — Deless! ic. sel. 2. t. 16. (v. s.)

Sect? III. MACROCERATIUM. DC. syst. 2. p. 204.

Siliquæ indehiscentes bicornutæ. Semina oblonga septo contraria. — Flores albi. Folia pinnatisecta. — An genus proprium?

4. N? GARDAMINEFOLIUM (DC. syst. 2. p. 205. *) siliquis bicornibus, foliis pinnatisectis glabris. ① in Ponto et insulâ Cypri. Deless! ic. sel. 2. t. 18. *Lepidium cornutum* Sibth. fl. græc. t. 617, ex prod. 2. p. 6. (v. s. in h. Mus. Par.)

VI. BARBAREA Brown in h. Kew. ed. 2. v. 4. p. 109. DC. syst. 2. p. 205.

Siliqua tetragono-anceps, valvis apice muticis concavo-carinatis. Calyx basi æqualis.

1. B. VULGARIS (Brown in h. Kew. ed. 2. v. 4. p. 109.) foliis inferioribus lyratis, lobo terminali subrotundo, superioribus obovatis dentatis. ② in Europâ. *Erysimum barbarea* Lin! spec. 922. var. α. et γ. OEd. fl. dan. t. 985. variat flore pleno. DC. syst. 2. p. 206. * (v. v.)
β. *gracilis*, caule gracili parcè ramoso, foliis superioribus subincisis — in Sibiria et Kamtschatkâ. An species propria?
2. B. PRÆCOX (Brown in h. Kew. ed. 2. v. 4. p. 109.) foliis inferioribus lyratis, lobo terminali ovato, superioribus pinnatipartitis, lobis lineari-oblongis integerrimis. ② in Angliâ. *Erysimum præcox* Smith! fl. brit. 2. p. 707. *

- engl. bot. t. 1129. Colitur in hortis, sub nominibus trivialibus, gall. *Roque des jardins*, angl. *Early winter-cress*, *Belle-Isle cress*. etc. DC. syst. 2. p. 207.* (v. v.)
3. **B. TAURICA** (DC. syst. 2. p. 207.*) foliis radicalibus et inferioribus pinnatipartito-lyratis, lobis terminalibus folisque summis ovatis dentatis, pedicellis patulis, siliquis adscendentibus. \mathcal{L} in Tauriâ et in Caucasi alpestribus. *Erysimum arcuatum* Presl. fl. cech. 138. ex Andr. *Barbarea arcuata* Andr. in litt. (v. s.)
4. **B. IBERICA** (DC. syst. 2. p. 208.*) foliis radicalibus et inferioribus pinnatipartito-lyratis, lobis lateralibus ovatis terminali cordato integro, foliis superioribus basi obtuse auriculatis. \mathcal{L} in Iberiâ et Podoliâ? *Cheiranthus Ibericus* Willd. enum. 2. p. 611. ex Stev. obs. ined. in h. Willd. Ch. lævigatus Willd. herb. ex eodem. *Barbarea stricta* Besser enum. Volh. n. 1558. Siliquæ mihi ignotæ, ex cl. Besser strictæ adpressæ. (v. s. sine fructu.)
5. **B. PLANTAGINEA** (DC. syst. 2. p. 208.*) foliis inferioribus dentato-lyratis, lobis lateralibus dentiformibus, terminali maximo subcordato, superioribus ovatis, siliquis subincurvis. \mathcal{L} in Oriente. Deless! ic. sel. 2. t. 19. *Sisymbrium Barbarea* Lin. spec. 921. Simillima *B. Vulgari*, sed ex omni parte major. Folia radicalia lyrata *Doronici plantaginei* ferè æmula. (v. s. in h. Tourn. Banks et Deless.)
6. **B. INTEGRIFOLIA** (DC. syst. 2. p. 208.*) foliis radicalibus integris oblongis petiolatis, canlinis auriculato-amplexicaulibus. \mathcal{L} in Cappadociâ. Siliquæ ignotæ. Ex habitu ad *Barbaream* relata (v. s. in h. Mus. Par.)

VII. STEVENIA. *Adams et Fisch. in mem. soc. nat. mosc. 5. p. 84.*

Siliqua oblonga, inter semina sinuato angustata, oligosperma, valvis planis subtorulosis. Calyx basi bisaccatus. — Flores albi aut subpurpurascens. DC. syst. 2. p. 209.

1. **St. ALYSOIDES** (Adams et Fisch! l. c. 5. p. 84.*) caule subascendente ramosissimo siliquis pedicello brevioribus. (1) in montibus Sibiriz. Flores albi; petala ovali oblonga, brevissimè unguiculata integra. DC. syst. 2. p. 209.* Deless! ic. sel. 2. t. 20. (v. s.)
2. **St. CHEIRANTHOIDES** (DC. syst. 2. p. 210.*) caule strictè erecto simplici aut parè ramoso, siliquis pedicello longioribus. \mathcal{L} (2) in Sibiria ulteriore. Deless! ic. sel. 2. t. 21. *Sisymbrium tomentosum* Stev! in litt. Flores ex albo purpurascens, ætate ut in *Cheiranthi* vari.

β . *Arabis incarnata* Pall! in h. Lamb. Differt foliis longioribus ferè pollicaribus, caule magis ramoso.

VIII. BRAYA *Sternb. et Hopp. diss. cum icon. ex Gætt. anzeig. Jan. 1817. p. 155. DC. syst. 2. p. 210.*

Siliqua oblonga, subcylindracea, valvis planiusculis; stigmatè sessili. Semina pauca, ovata. Calyx basi æqualis. — Flores purpurascens.

1. **B. ALPINA** (Sternb. et Hopp. l. c.) foliis spathulato-linearibus integris, racemo fructifero brevi. \mathcal{L} in Alpibus Carinthiæ superioris. DC. syst. 2. p. 211. Deless! ic. sel. 2. t. 22. (v. s.)
2. **B? GLABELLA** (Richards. in Frankl. narr. Journ. p. 743.) foliis linearibus plerumque remotè dentatis, racemo fructifero laxo elongato. \mathcal{L} in Cupreis montibus à Point Lake ad mare arcticum. Caules palmares erecti costati. Siliquæ torulosæ.

IX. TURRITIS. *Dill. nov. gen. in pl. giess. p. 120. t. 6. Gaertn. fruct. 2. p. 297. t. 143.*

Siliqua linearis; valvis planis. Semina in quoque loculo biserialia. — Flores albi aut albidi.

1. *T. GLABRA* (Lin! spec. 930.) foliis radicalibus dentatis pilosis, caulinis amplexicaulibus integerrimis glabris, siliquis strictissimis pedicello sextuplo longioribus. ② in pascuis siccis Europæ. Smith engl. bot. t. 777. *Arabis perfoliata* Lam! dict. 1. p. 219. ill. t. 563. f. 4. — DC. syst. 2. p. 211.* (v. v.)

β. *ramosa*, *Turrita major* Clus. hist. 2. p. 126. f. 2.

2. *T. SALSUGINEOSA* (DC. syst. 2. p. 212.*) foliis glabris, inferioribus dentatis, caulinis amplexicaulibus integris, siliquis erectiusculis pedicello quadruplo longioribus. ① in Sibiriâ. *Sisymbrium salsugineum* Pall! itin. 2. app. n. 114. t. V. ed. gall. min. 8. p. 342. n. 348. t. 64.* — *Flores exigui, albi.* (v. s.)

3. *T. MISPIDULA* (DC. syst. 2. p. 213.*) foliis pube flocculosâ scabris, caulinis amplexicaulibus oblongo-lanceolatis subserratis, siliquis erectiusculis subcabis pedicello triplo longioribus. — in regno Quitensi. *Arabis Andicola* H. B. et K! nov. spec. amer. 5. p. 81.* Petala alba, linearia, obtusa, basi attenuata. Semina innumera, biseriata, parva, rufa. (v. s. in h. Humb.)

X. *ARABIS*. Lin. gen. n. 818. Lam. ill. t. 563.

Siliqua linearis; valvis planis medio uninerviis. Semina in quoque loculo 1-serialia, ovalia aut orbicularia, compressa. Cotyledones planæ o=. Flores albi aut rarius rosei.

Sect. 1. *ALOMATIUM*. DC. syst. 2. p. 214.

Semina immarginata, seu parvâ alâ membranaceâ cincta.

- §. 1. *Foliis caulinis cordato-amplexicaulibus, petalorum limbis obovatis ab ungue distinctis patentibus.*

1. *A. VERNA* (Brown in h. Kew. ed. 2. v. 4. p. 105 non Desf.) foliis caulinis cordatis amplexicaulibus dentatis pube tripartitâ scabris, pedicellis calyce brevioribus, stigmate subemarginato. ① in arvis agris et asperis Europæ australis. *Hesperis verna* Lin! spec. 928. — Moris. hist. 2. p. 241. s. 3. t. 8. f. 5. Petala parva, purpurea, ungue albo. DC. syst. 2. p. 215.* (v. v.)

2. *A. ROSEA* (DC. syst. 2. p. 215.*) foliis caulinis oblongis semi-amplexicaulibus subcordatis dentatis pube ramosâ scabris, pedicellis calyce longioribus, stigmate apiculato. ② in Calabriâ. Petala roseo-purpurascens, oblonga, subcuneata, calyce duplo longiora. Deless! ic. sel. t. 23. (v. s. in h. Reynier.)

3. *A. VISCOSA* (DC. syst. 2. p. 216.*) foliis pauci-dentatis pube ramosâ scabris, radicalibus petiolatis obovato-oblongis obtusissimis, caulinis cordato-ovatis amplexicaulibus, pedicellis calyce hirsuto æqualibus. ③ in Persiâ septentr. *Cardamine viscosa* Gmel. in herb. Pall. (v. s. ex h. Pall.)

4. *A. ALPINA* (Lin! spec. 928.) foliis pluri-dentatis pube ramosâ villosis lanceolatis acutis, radicalibus subpetiolatis, caulinis cordato-amplexicaulibus, pedicellis calyce glabriusculo longioribus. ④ in rupibus saxosisque apricis montium calcareorum Europæ. Curt. bot. mag. t. 226. DC. syst. 2. p. 216.* (v. v.)

β. *clusiana*, foliis percussis et obtusius dentatis. *A. Clusiana* Schranck fl. mon. 2. p. 125. — Clus. hist. 2. p. 125. f. 2.

γ. *nana*, *Arabis Alpina nana* Baumg. fl. trans. 2. p. 263.

5. *A. ALBIDA* (Stev! in cat. h. gor. 1812. p. 51.) foliis pauci-dentatis pube ramosâ cano-villosis subtomentosis, radicalibus obovato oblongis, caulinis cordato-sagittatis amplexicaulibus, pedicellis calyce longioribus. ⑤ in Tauriâ. *A. Caucasica* Willd. enum. suppl. p. 45. Schranck h. mon. t. 24. Valdè affinis *A. Alpinae* sed distincta foliis pauci nec multidentatis. DC. syst. 2. p. 217.* (v. s. et v.)

6. *A. UNDULATA* (Link enum. h. ber. 2. p. 161.) caule erecto foliisque oblongis dentatis hirtis, siliquis patentibus, stylo latitudine siliquæ. ⑥ in Europâ australi. Aff. præc. sed minor, et serius florens.

7. *A. BILLARDIERII* (DC. syst. 2. p. 218.*) foliis pauci-dentatis pube ramosâ cano-

- villosis subtomentosis, inferioribus obovatis cuneatis, summo oblongo amplexicauli, pedicellis calyce longioribus, siliquis demum petalis. — in Syriâ propè Damascum. Flores albi, ferè A. Alpinae. Siliquæ patulae, lineares, glabrae, sesqui pollicem longæ. Deless! ic. sel. 2. t. 24. (v. s.)
8. *A. brevifolia* (DC. syst. 2. p. 218.*) foliis obtusè pauci-dentatis pube ramosâ velutino-tomentosis, radicalibus obovatis basi breviter attenuatis, caulinis cordato-sagittatis ovatis subintegris, pedicellis calyce longioribus, siliquis petalis curvulis. ½ in Syriâ (v. s.)
9. *A. thirsoidea* (Sibth. et Smith fl. græc. t. 642. ex prod. 2. p. 28.) foliis obovatis obtusè dentatis pube ramosâ cano-tomentosis, caulinis ovatis subamplexicaulibus, racemo capitato, pedicellis calyce subbrevioribus, siliquis curvato-ascendentibus. ½ in Olympi Bithyni cacumine. (v. s. in h. Banks.)
10. *A. longifolia* (DC. syst. 2. p. 219.*) foliis dentatis pube ramosâ cano-pubescentibus, inferioribus obovato-oblongis longissimè petiolatis, caulinis oblongis subamplexicaulibus, pedicellis calyce longioribus. — in Persiâ. Nimis A. albidæ affinis sed distincta. Deless! ic. sel. 2. t. 25. (v. s.)
11. *A. mollis* (Stev! mem. soc. nat. mosc. 3. p. 270.) foliis grossè dentatis pube stellatâ minimâ subpubescentibus, inferioribus longè petiolatis cordato-subrotandis, caulinis cordato-ovatis amplexicaulibus. ②? in rupibus umbrosis Caucasii orient. Bieb. fl. taur. suppl. 448.* non Raf. Flores albi magnitudine A. Alpinae. DC. syst. 2. p. 219.* (v. s.)
12. *A. toxophylla* (Bieb. fl. taur. suppl. p. 448.*) foliis pube minimâ stellatâ subpubescentibus, radicalibus oblongis petiolatis subsinnato-dentatis, caulinis sagittato-lanceolatis integerrimis, pedicellis calyce longioribus. ②? in campis apricis ad Wolgam inferiorem. Sisymbrium salsugineum Schlecht. in h. Willd. ex Stev. obs. ined. Species elegans distinctissima. DC. syst. 2. p. 220.* (v. s.)
- § 2. *Foliis caulinis cordato-amplexicaulibus, petalis oblongo aut cuneato-linearibus erectis.*
13. *A. auriculata* (Lam! dict. 1. p. 219.*) foliis subdentatis pube ramosâ scabris, inferioribus ovalibus in petiolum subattenuatis, caulinis obtusè cordato auriculatis, pedicellis calyce vix longioribus, siliquis glabris puberulisve. ① Petala alba, oblonga. DC. syst. 2. p. 220.* (v. v.)
- α. *A. aspera* All! pedem. auct. p. 18. t. 2. f. 2.* in Pedemontio.
- β. *A. recta* Vill. dauph. 3. p. 319. t. 37.* in Hispaniâ, Delphinatu.
- γ. *Turritis patula* Ehrh. Beit. 7. p. 259? Waldst. et Kit. pl. rar. hung. 1. p. 59. t. 59.* in Hungariâ.
- δ. *dasycarpa* (Andrz! in litt.) siliquis pubescentibus. ① in Thuringiâ et Podoliâ australi. A. recta Bangm. fl. trans. 2. p. 267.* A. recta siliq. pubesc. Wallr! (v. s.)
14. *A. saxatilis* (All! pedem. n. 973.) foliis subdentatis pube ramosâ scabris, inferioribus ovalibus in petiolum subattenuatis, caulinis acutiusculè cordato-auriculatis, pedicellis demum calyce quadruplò longioribus. ② in Delphinatu, Sabaudiâ, etc. A. nova Vill. Dauph. 3. p. 319. t. 37.* DC. syst. 2. p. 221.* (v. v.)
- β. *Tourette cotoneuse* Reyn! mem. suiss. 1. p. 168. Paulò magis cano-tomentosa et minor evadit, forsan ex loco magis siccò et aprico orta.
15. *A. crispata* (Willd. enum. 684.) foliis acutè dentatis pube ramosâ scabris lanceolatis amplexicaulibus undulatis, radicalibus in petiolum attenuatis, pedicellis calyce longioribus, siliquis patentibus. ½ in Carnioliâ. Petala obovato-oblonga, obtusa. DC. syst. 2. p. 221.* (v. v.)
16. *A. sagittata* (DC. fl. fr. suppl. 592.* syst. 2. p. 221.*) foliis subdentatis pube sæpius ramosâ scabris, radicalibus ovatis aut oblongatis in petiolum attenuatis, caulinis lanceolatis sagittato-cordatis, pedicellis calycis longitu-

dine, siliquis strictè erectis. ② in asperis campis in totius ferè hemisphæræ borealis parte temperatâ. (v. v.)

α. *Gerardiana*, foliis scabris, caulinis basi in aurículas productis. — In Enropâ austr. *Turritis hirsuta* Linn! herb. *Arabis hirsuta* Smith fl. brit. 2. p. 717. in adnot.

β. *subglabrata*, foliis ferè glabris, caulinis auriculato-sagittatis, racemo composito. — in Vasconiâ.

γ. *ovata*, foliis scabris, radicalibus ovatis dentatis, petiolo limbo longiore, caulinis amplexicaulibus. — in Americâ boreal. *A. ovata* Poir. suppl. 5. p. 557.

δ. *oblongata*, foliis scabris, radicalibus ovato-oblongis dentatis, petiolo limbo brevior, caulinis sagittato-amplexicaulibus. — in Americâ boreal. *Turritis oblongata* Raf! amer. monthl. mag. 2. p. 44.

ε. *longisiliqua*, siliquis pedicello decies longiore. *Arabis longisiliqua* Wallr! sched. p. 359.

17. *A. hirsuta*. (Scop. fl. carn. ed. 2. n. 835.) foliis dentatis pube sæpius ramosâ scabris, radicalibus obovato-oblongis in petiolum attenuatis, caulinis ovato-lanceolatis; pedicellis calycis longitudine, siliquis erectis. ② in Europâ. Schrank fl. monac. 3. t. 248. *Turritis hirsuta* Jacq. coll. 1. p. 70.* ic. rar. 1. t. 126. DC. syst. 2. p. 223. (v. s. et v.)

18. *A. curtisiliqua* (DC. syst. 2. p. 223.*) foliis oblongis subintegris pilis simplicibus ciliatis, radicalibus in petiolum attenuatis, caulinis subamplexicaulibus, pedicellis calycis longitudine, siliquis erectis. ② in Scaniâ. *Turritis curtisiliqua* Fries ex Horn! herb. (v. s. in h. Hornem.)

§. 3. *Foliis caulinis sessilibus, petalis oblongo aut cuneato-linearibus erectis.*

19. *A. ALLIONII* (DC. fl. fr. 4. p. 676.*) foliis glabris, radicalibus ovato-oblongis subdentatis basi attenuatis, caulinis sessilibus ovatis serratis, pedicellis calyce sublongioribus, siliquis erectis. ② in pratis Pedemontii suprâ Casetto. *Turritis stricta* All! auct. p. 18. Flores albi. Calyces pallidi, demùm subpatentes. — DC. syst. 2. p. 224.* (v. s.)

20. *A. muralis* (Bertol! dec. ital. 2. p. 37.*) foliis pube ramosâ hirsutis, radicalibus spatulatis obtusè dentatis, caulinis ovatis acutè dentatis, racemo stricto, pedicellis calyce demùm duplò longioribus. ② in rupibus montanis Etruriæ, etc. *A. humilis* Schleich! pl. exs. Petala alba, oblonga, Siliquæ strictæ. — DC. syst. 2. p. 224.* (v. s.)

21. *A. stricta* (Huds angl. 292) foliis pube sparsâ bifidâ ciliato-scabris, radicalibus obovatis dentatis, caulinis oblongis subintegris, racemo stricto, pedicello calyce vix longioribus. ② in rupibus Europæ. Smith engl. bot. t. 614. *Turritis Rayi* Vill. Dauph. 3. p. 326. t. 38.* Flores albid. — DC. syst. 2. p. 225. (v. v.)

22. *A. ciliata* (Brown in h. Kew. ed. 2. v. 4. p. 107.) foliis subdentatis glabris ciliatis radicalibus subsessilibus ovali-oblongis, caulinis oblongis, racemo stricto, pedicellis calycis longitudine. ② in Alpium Aquilejensium et Hiberniæ. *Turritis Alpina* Willd. spec. 3. p. 545. Sm. h engl. bot. t. 1746. Valdè accedit ad *A. strictam*, sed differt radice bienni nec perenni. — DC. syst. 2. p. 225.* (v. s.)

23. *A. incana* (Roth. cat. bot. 1. p. 79.) foliis omnibus sessilibus subdentatis pilis ramosis incanis, radicalibus obovato-oblongis, caulinis oblongis, racemo erecto, pedicellis demùm calyce duplò longioribus. ② in campestribus siccis Helvetiæ. *Turritis minor* Schleich. pl. exsicc. Petala alba, oblonga, calyce duplò longiora. — DC. syst. 2. p. 225.* (v. s.)

24. *A. thaliana* (Lin! spec. 929.) foliis pilosis subdentatis, radicalibus petiolatis ovato-oblongis, caule ramoso, siliquis adscendentibus, pedicellis calyce multò longioribus. ① in aridis Europæ. Smith engl. bot. t. 901. Vahl fl. dan. t. 1106. DC. syst. 2. p. 226.* (v. v.)

ℓ. *A. parviflora* Rafin! amer. monthl. mag. 1. p. 43. 1819. jan. p. 194. — in Americâ bor.

25. *A. SERPYLLIFOLIA* (Vill. Dauph. 3. p. 318. t. 37.*) foliis subintegris pilis ramosis ciliato-subscabris, radicalibus caulinisque ovalibus basi subattenuatis, racemo laxiusculo, pedicellis calycis longitudine. ② in rupestribus montosis Alpinum Galloprovinciæ, Delphinatûs, Vallesiæ et in Pyrenæis. Species nullq modo *A. Alpinæ*, sed potiùs *A. thalianæ* affinis. DC. syst. 2. p. 227.* (v. v.)

26. *A. SPATHULATA* (DC. syst. 2. p. 227.*) foliis integris ciliato-scabris, inferioribus spathulatis in petiolum attenuatis, superioribus subrotundis parvis, racemo brevi, pedicellis calycis longitudine, siliquis erectis. ①? in agro Bonariensi. *Sisymbrium spathulatum* Poir! dict. 7. p. 221.* — Petala alba, parva, calyce vix longiora. DC. syst. 2. p. 227.* (v. s. in h. Mus. Par.)

27. *A. ? COMMERSONII*. (DC. syst. 2. p. 228.*) foliis subintegris ciliatis subvillosis, radicalibus ovali-subspathulatis petiolatis, caulinis subnullis, racemo brevi, pedicellis calyce longioribus ①? in Bonariâ. *Sisymbrium bellidifolium* Poir! dict. 7. p. 220.* Petala alba, vix calyce longiora. Semina minima rufescentia. (v. s. in h. Mus. Par.)

28. *A. PUBESCENS* (Poir. suppl. 1. p. 413.) foliis pubescentibus grossè dentatis, radicalibus spathulato-lanceolatis in petiolum angustatis, caulinis lanceolatis, pedicellis brevissimis, siliquis pubescentibus erectis. ②? in Algeriæ montibus. *Turritis pubescens* Desf! fl. atl. 2. p. 92. t. 163. Petala linearia calyce duplò longiora. DC. syst. 2. p. 228.* (v. s. in h. Desf.)

29. *A. PARVULA*. (Dufour! in DC. syst. 2. p. 228.*) foliis pubescentibus subdentatis, inferioribus obovatis, caulinis ovatis, pedicellis brevissimis, siliquis pubescentibus subpatulis. ① in Navarrâ Hispaniâ propè Tudelam. *A. Rœmeriana* Andr. ined. Petala in sicco subochroleuca. (v. s.)

§. 4. *Foliis caulinis sessilibus, aut petiolatis; petalorum limbo obovato seu cuneato ab ungue distincto.*

30. *A. PROCURRENS* (Waldst. et Kit. pl. rar. hung. 2. p. 154. t. 144.*) foliis ovatis integerrimis glabris pilis bipartitis ciliatis, radicalibus in petiolum attenuatis, caulinis sessilibus acuminatis, stolonibus reptantibus, siliquis patulis. ½ in rubis calcareis umbrosis Banatûs et Serviæ. *Nasturtium procurrens* Andr. ined. — Petala alba calyce duplò longiora. — DC. syst. 2. p. 229.* (v. s.)

31. *A. PRÆCOX* (Waldst. et Kit. ined. ex Willd. enum. p. 684.) foliis oblongis acutis sessilibus integerrimis glabris, caulibus strigosis, stolonibus repentibus, siliquis patulis. ½ in Hungariâ. Petala obovato-cuneata alba, calyce duplò longiora. DC. syst. 2. p. 229.* (v. s. comm. à cl. Spreng.)

32. *A. SCHWABECKIANA* (Andr. in litt.) foliis integris pube ramosâ subconfertâ scabris, radicalibus rosulatis obovatis, caulinis oblongis erectis sessilibus, caulibus subscabris, siliquis erectis glabris. ½ in Austriâ. Aff. paululùm ex habitu *A. scabræ*, sed siliquæ dimidio angustiores et semina vix marginata. (v. s.)

33. *A. PETRÆA* (Lam! dict. 1. p. 221.) foliis glabris aut pilis simplicibus bifurcive ciliatis scabrisve, radicalibus incisis, caulinis oblongo-linearibus integris, caule sæpiùs ramoso, siliquis erecto-patulis, stigmate capitato. ½ DC. syst. 2. p. 230.* (v. s.)

α. *Crantziana*, siliquâ longâ. *A. Crantziana* Ehrh. herb. 78. — in Austriâ, Transylvaniâ.

β. *hispida*, *Sisymbrium arenosum* Lin. fl. suec. 233. nec. spec. pl. Ed. fl. dan. t. 386. — in Scotiâ, Cambriâ.

γ. *hastulata*, *Cardamine hastulata* Smith! engl. bot. t. 409. Horn. fl. dan. t. 1462. — in Britanniâ, Norvegiâ.

δ. *Færøensis*, *Cardamine Færøensis*. Horn! fl. dan. 1382. — in ins. Færøensibus.

ε. *runcinata*, *A. petræa* Lam! dict. 1. p. 22. — in Sueciâ, Pyrenæis.

34. *A. AMBIGUA* (DC. syst. 2. p. 231.*) foliis subglabris, radicalibus sinuato-

lyratis, mediis oblongo-ovalibus dentatis, summis oblongo-linearibus integris, caule subsimplici, siliquis erectiusculis ② (v. s. comm. à cl. Fisch. et Stev.)

α. *glabra*, foliis cauleque glabris. — in Kamtschatkâ.

β. *intermedia*, foliis glabris, caule basi hispido. — in Unalaskâ.

γ. *scabra*, foliis piloso-scabris, caule glabro. — in Sibirîâ.

35. A. *LYRATA*. (Lin! spec. 929.) foliis radicalibus lyrato-pinnatifidis glabris ciliatisve, caulinis linearibus, caule basi hispido subramoso, pedicellis subpatulis, siliquis erectis. ① in arvis et collibus siccis Americæ borealis. DC. syst. 2. p. 231.* (v. s.)

β. A. *Kamschatika* (Fisch! in litt.) In posterum forsan distinguenda: differt flore minore, petalis nempe calyce duplò nec triplò longioribus et siliquis longioribus erassioribus. (v. s.)

36. A. *ARENOSA* (Scop. fl. carn. ed. 2. n. 837. t. 40.*) foliis pube bifurcâ villosis, radicalibus lyrato-pinnatifidis, caulinis inciso-dentatis, caule ramoso pilis simplicibus hispido, pedicellis siliquisque patulis. ① in arenosis vineis et rupibus Europæ mediæ. *Symbrium arenosum* Lin. spec. 919. — Petala obovata, rosea rarius alba aut cœrulescentia; siliquæ lineares, patulæ. DC. syst. 2. p. 232.* (v. s.)

37. A. *HALLERI* (Lin! spec. 929.) foliis inferioribus petiolatis lyratis, lobo terminali ovato, superioribus lanceolatis iucis, caule ramoso debili molliter villosulo, pedicellis siliquisque patulis. ② à Transylvaniâ ad Pedemontium et ad Hercyniam. Waldst. et Kit. plant. rar. Hung. 2. p. 126. t. 120. DC. syst. 2. p. 232.* (v. s. et v.)

38. A. *STOLONIFERA* (Horn h. hafn. 618.) foliis radicalibus petiolatis sublyratis, lobo terminali cordato, caulinis oblongis sinuato-dentatis, caule basi stolonifero pubescente, pedicellis siliquisque patulis. ① in Carniolâ. *Cardamine stolonifera* Scop. fl. carn. ed. 2. n. 818. t. 39.* malè. Flores albi. Siliquæ tenuissimæ, semipollicares. — DC. syst. 2. p. 233.* (v. s.)

39. A. *OVIENSIS* (Wulf. in Jacq. coll. 1. p. 196.* ic. rap. 1. t. 125.) foliis inferioribus petiolatis ovalibus basi pectinato-dentatis, superioribus oblongis basi attenuatis, caule debili glabro, siliquis subpatulis pedicello vix longioribus. ① in locis nigligiosis petrosis calcareis in Carinthiæ Oviensibus Alpibus. *Sturmdentschl. fl. ic. Nasturtium Oviense*. Andr. ined. — Petala dilutè rubellâ, rarius alba. Semina ovali-oblonga. — DC. syst. 2. p. 233.* (v. s.)

40. A. *CEBENNENSIS* (DC. syst. 2. p. 234.*) foliis omnibus petiolatis ovatis acuminatis grossè dentatis pube minimâ subvelutinis, pedicellis siliquisque patulis. ② in locis subumbrosis montium Cebennorum in horto Dei et Brémabion. *Hesperis inodora* Gouan! fl. Monsp. 167. Flores pallidè violacæ, ferè A. *arenosæ*. semina rufo-fusca. Deless! ic. sel. 2. t. 26. (v. s.)

41. A. *LASIOLOSA*. (Link. enum. h. ber. 2. p. 163.) caule foliis siliquisque stellulato-tomentosis, foliis radicalibus lyratis, superioribus integris. ② in Mexico. Racemi breves. Cal. patens. Stylus ferè lineam longus.

Sect. II. LOMASPORA. DC. syst. 2. p. 234.

Semina marginata seu alâ membranacèâ latè cincta.

§. 1. *Foliis caulinis amplexicaulibus, petalis oblongo-linearibus.*

42. A. *TURRITA* (Lin! spec. 930.) foliis amplexicaulibus acuminatis subdentatis pubescentibus, pedicellis calycis longitudine, siliquis secundis decurvis. ② in sepibus montosis Europæ. Jacq. fl. austr. t. 11. *Turritis ochroleuca* Lam. fl. fr. 2. p. 490.* — Flores conferti, albidò-ochroleuci; petala oblongo-linearit. — DC. syst. 2. p. 235.* (v. v.)

β. *longisiliqua*, A. umbrosa Stev! in litt.

γ. *pendula*, Lach. obs. p. 10. (non L.) Poir. suppl. 2. p. 410. excl. syn. *Turritis pendula* Desf. cat. ed. 2. p. 152.

43. *A. PENDULA* (Lin! spec. 930.) foliis amplexicaulibus dentatis oblongis basi dilatato-cordatis, caule sulcato hispido, pedicellis calyce triplò longioribus, siliquis pendulis. ① in Sibiria et forsan in amer. bor. ad flum. Missouri ex Nutt. — Jacq. h. vind. 3. p. 20. t. 34. — Petala alba, oblongo-lineararia; siliquæ pendulæ laxæ. (v. s.)

44. *A. PATRIANIANA* (DC. syst. 2. p. 236.*) foliis oblongis acuminatis subseratis cordato-amplexicaulibus villosis, caule tereti hispido, pedicellis calyce triplò longioribus, siliquis pendulis. — in Sibiria ad Ufan. Deless! ic. sel. 2. t. 27. *A. pendula* Patr! herb. an prioris mera varietas? Semina alâ angustâ cincta. (v. s.)

45. *A. OXYOTA* (DC. syst. 2. p. 236.*) foliis pube ramosâ scabris, caulinis acutè sagittatis oblongo-linearibus subdentatis, caule pilis simplicibus hispido, pedicellis calyce triplò longioribus. — in Sibiria orient. Ad duas præcedentes valdè accedit. Petala alba oblonga. (v. s.)

§. 2. *Foliis caulinis sessilibus, petalis oblongis aut cuneato-linearibus.*

46. *A. HELIOPHILA* (DC. syst. 2. p. 237.*) foliis sessilibus linearibus integris pube brevè adpressâ bipartitâ canescentibus, pedicellis calyce brevioribus, siliquis erectis. — in India orient. *Heliophila incana* Burm! fl. ind. 140. t. 46. f. 2. Petala lineararia angustissima vix calycem superantia. (v. s. in h. Deless.)

47. *A. LÆVIGATA* (DC. syst. 2. p. 237.*) foliis caulinis linearibus sessilibus glabris, inferioribus subdentatis, radicalibus obovatis, siliquis erectis. ② \mathcal{L} in rupibus à Pennsylvania ad Virginiam, sed rara. *Turritis lævigata* Willd. spec. 3. p. 543. Flores parvi pauci, corymboso-racemosi. (v. s. in h. Lamb.)

48. *A. CANADENSIS* (Lin! spec. 929.) foliis caulinis sessilibus oblongo-lanceolatis acuminatis subdentatis, pedicellis calyce triplò longioribus, siliquis pendulis subfalcatis. \mathcal{L} in rupibus umbrosis Americæ bor. — Pluk. alm. 136. t. 86. f. 8. — Petala linearia, calyce longiora; semina 1-serialia compressa, alâ membranaceâ cincta. DC. syst. 2. p. 238.* Deless! ic sel. 2. t. 28. (v. s.)

a. mollis Rafin. amer. monthl. mag. 2. p. 43.* non Stev.

ß. falcata, Michx. fl. bor. am. 2. p. 31. *

49. *A. FUMILA* (Wulf. in Jacq. coll. 2. p. 59. fl. austr. t. 281.) foliis scabris subintegris, radicalibus obovatis, caulinis ovatis oblongisve, racemo nutante, pedicellis calyce duplò longioribus, siliquis erectis pedicello triplò longioribus. \mathcal{L} in lapideosis à Galloprovincia ad Transylvaniam. *A. bellidifolia* Crantz. austr. 42. t. 2. f. 3. non Jacq. *A. ciliaris* Willd! enum. 2. p. 684. — DC. syst. 2. p. 238.* (v. v.)

50. *A. BELLIDIFOLIA* (Jacq! obs. 1. p. 22. t. 12. fl. austr. 3. t. 280.*) foliis glabris subintegris, radicalibus obovatis; caulinis ovatis, racemo erecto, pedicellis calyce triplò longioribus, siliquis pedicello quadruplò longioribus. \mathcal{L} in graminosis lapideosis Pyrenæorum orient. et Alpium. *Turritis bellidifolia* All. ped. n. 980. t. 40. f. 1. — Petala oblongo-cuneata, calyce duplò longiora. Siliquæ lineares. — DC. syst. 2. p. 239.* (v. v.)

ß. Turritis Alpina Jacq. in Murr. syst. veg. 502. Willd. spec. 3. p. 545.

51. *A. CÆRULEA* (Wulf. in Jacq. coll. 2. p. 56.*) foliis glabris subintegris, radicalibus oblongo-obovatis; caulinis paucis oblongis, racemo nutante, pedicellis calycis longitudine, siliquis erectis. \mathcal{L} Propè nives perpetuas in Alpibus. Sturm. deutsch. fl. ic. *Turritis cærulea* All. pedem. n. 981. t. 40. f. 2. Petala oblonga pallidè et sordidè cærulea. DC. syst. 2. p. 239.* (v. v.)

§. 3. *Foliis caulinis sessilibus aut nullis, petalorum limbo obovato patente.*

52. *A. STELLULATA* (Bertol! in Desv. journ. 1813. v. 2. p. 76. Amœn. ital. p. 101.*) foliis pube stellato-ramosâ scabris, radicalibus obovatis, caulinis oblongis paucissimis, siliquis pedicello duplò longioribus 16-20-spermis. \mathcal{L}

in summis Apenninis Appuanis. Habitu accedit ad *A. petræam*, var. *Crantzianam*, sed seminibus latè marginatis evidenter differt. DC. syst. 2. p. 240.* (v. s.)

53. *A. ? VOCHINENSIS* (Spreng. pug. 1. p. 46. n. 81.*) foliis pube bipartitâ adpressâ ciliatis, radicalibus obovatis, caulinis paucis oblongis, siliquis pedicelli longitudine 4-8 spermis. \mathcal{L} in Carniolæ Alpibus Vochinensibus. *Draba mollis* Scop. carn. n. 789 t. 34.* Petala obovata basi attenuata, calyce duplò triplove longiora. — DC. syst. 2. p. 241.* (v. s.)

54. *A. COLLINA* (Tenor! prod. fl. nap. xxxix. app. cat. h. ræap. 1815. p. 59.) foliis pube stellatâ canescentibus oblongis sinuato-dentatis, radicalibus petiolatis, caulinis sessilibus, siliquis pedicello octies longioribus. \mathcal{L} in collibus circâ Neapolim. Petala alba, limbo obovato patente calyce duplò longiore. DC. syst. 2. p. 241.* (v. s.)

†. *Species non satis notæ.*

55. *A. STELLERI* (DC. syst. 2. p. 242.*) pilis sæpè bifurcis hispida, foliis inferioribus oblongo-spathulatis, superioribus semi-amplexicaulibus oblongis dentatis, racemo corymboso, petalis cuneato-oblongis. \mathcal{L} in Kamtschatkâ. *A. pendula* Steller in h. Pall. ex Fisch! non Lin. Petala alba, calyce duplò longiora. (v. s.)

56. *A. LUCIDA* (Lin. f. suppl. 298.) foliis amplexicaulibus lucidis. \mathcal{L} in Pennoniâ. Petala alba, linearia, integra, calyce duplò longiora, basi angustiora. Specimen h. Willd! valdè mancum videtur *A. ciliatæ* var. *glabra*. DC. syst. 2. p. 242.*

57. *A. REPTANS* (Lam. dict. 1. p. 222.) foliis subrotundis integerrimis hirsutis, surculis reptantibus. \mathcal{L} in arvis arenosis Americæ. — Pluk. alm. 281. t. 51. f. 5. — Habitus ferè *Hieracii auriculæ*. Flores parvi. Siliquæ parvæ erectæ. DC. syst. 2. p. 242.*

58. *A. LAXA* (Smith. prodr. fl. græc. 2. p. 28.) foliis radicalibus obovatis lyrato-dentatis hispidis, caulinis cordatis amplexicaulibus glaberrimis, siliquis latè deflexis. \mathcal{L} in agro Laconico. Flores albi, siliquæ longissimæ et angustissimæ. DC. syst. 2. p. 243. *

59. *A. PETIOLATA* (Bieb. fl. taur. 2. p. 126.) foliis ovatis petiolatis glabris, inferioribus lobatis, superioribus repandis, siliquis striato-angulatis patentibus. — in Iberiâ. Flores exigui magnitudine *Erysimi cheiranthoidis*, albidis. Siliquæ patentes, 2-3-unciales, obscurè tetragonæ. DC. syst. 2. p. 243.*

60. *A. STENOPETALA* (Willd. enum. suppl. 46.) foliis hispidis grossè dentatis, radicalibus oblongo-lanceolatis petiolatis, caulinis ovatis amplexicaulibus, petalis erectis linearibus calyce duplò longioribus. \mathcal{L} ① Hab. Horn. h. hafn. 2. p. 619. Petala linearia obtusa alba. Siliquæ strictissimæ. — An stirpes Willdenowii et Hornemannii omninò eadem?

61. *A. MULTIFLORA* (DC. syst. 2. p. 243.) foliis sessilibus lanceolatis dentatis scabris, pedunculis adpressis multifloris, siliquis erectis tenuibus compressis confertis, pilis furcatis. (Lapeyr.) ② in Pyrenæis ad montem Chatelet. Turritis multiflora Lapeyr. abr. 386.

62. *A. INTEGRIFOLIA* (Lapeyr. abr. 385. suppl. p. 93.) hirsuta, foliis scabris lanceolatis integerrimis, caulinis amplexicaulibus, petalis erectis calyce duplò longioribus. ② ① in Pyrenæis loco dicto Mail du Cristal. DC. syst. 2. p. 244.

63. *A. LYRÆFOLIA* (DC. syst. 2. p. 244.) lævis, caule stricto simplicissimo, foliis radicalibus lyratis, caulinis sessilibus oblongis acutis dentatis, siliquis falciiformibus. ① in sylvis Americæ boreal. Turritis lyrata Rafin. amer. monthl. mag. 2. p. 44.

64. *A. ANGUSTIFOLIA* (Lam. dict. 1. p. 220.*) foliis radicalibus longè petiolatis rhombeis parvis bidentatis, caulinis angusto-lanceolatis integerrimis. — Hab. olim culta in h. Paris. An *Cardamines Alpinæ* varietas?

65. A? *SILICULOSA* (I. am. dict. 1. p. 221. *) foliis radicalibus oblongis subpetiolatis glabris versùs apicem dentatis, caulinis raris angustis integris. — in Sibiria? Siliculæ complanatæ, utrinquè attenuatæ, 4 lin. longæ, 1-2-spermæ. DC. syst. 2. p. 244.

66. A. *DASYCARPA* (Andrz. in DC. syst. 2. p. 244.) foliis pube stellatâ hirsutis, caulinis sagittatis subintegerrimis, lobis divergentibus, siliquis hirsuto-scarbis erectiusculis latitudine stylum duplò superantibus. — in Podoliâ.

XI. *MACROPODIUM*. R. Brown. in h. Kew. ed. 2. v. 4. p. 108. DC. syst. 2. p. 244.

Siliqua linearis pedicellata, valvis planis medio 1-nerviis. Semina orbiculata, compresso-plana, alâ brevissimâ cincta. (0=.)

1. M. *NIVALE* (R. Brown. in h. Kew. ed. 2. v. 4. p. 108.) foliis radicalibus ovatis longè petiolatis inæqualiter serratis, caulinis lanceolatis basi attenuatis acuminatis integris. ♀ in summis montibus Altaicis propè nives perpetuas. *Cardamine nivalis* Pall! it. 2. app. n. 113. t. U* — DC. syst. 2. p. 245. (v. s.)

XII. *CARDAMINE*. DC. syst. 2. p. 245. — *Cardamines* sp. Lin. Brown.

Siliqua linearis, valvis planis enerviis sæpè elasticè dissilientibus. Semina ovata immarginata; funiculi umbilicales tenues.

§. 1. *Foliis ferè omnibus indivisis.*

1. C. *RHOMBOIDEA* (DC. syst. 2. p. 246. *) foliis ovato-rhomboideis obsolete repando-dentatis glabris, inferioribus longè petiolatis, radice tuberosâ et fibrosâ. ♀ Secus rivulos et fontes Americæ bor. *Arabis rhomboidea* et A. *tuberosa* Pers. ench. 2. p. 204. — Pluk. enalth. t. 435. f. 6. Flores albi magnitudine ferè C. *pratensis*. (v. s.)

2. C. *ROTUNDIFOLIA* (Michx! fl. bor. am. 2. p. 30.) foliis orbiculatis subdentatis glabris petiolatis, caulibus debilibus procumbentibus, radice fibrosâ. ♀ ad rivulos montium Carolinæ, Virginæ, Pensylvaniæ. Siliquæ patentès graciles, stylo 2 lin. longo apiculatæ, tenuissimæ, vix pedicello longiores. DC. syst. 2. p. 247. * (v. s.)

3. C. *SPATHULATA* (Michx! fl. bor. am. 2. p. 29.) foliis radicalibus petiolatis spatulatis integris pube trifurcâ pubescentibus, caulinis linearibus, caule decumbente. — in excelsis montibus Carolinæ. Siliquæ lineares, rectæ, patulæ, subreflexæ, stigmate sessili vix acutæ. DC. syst. 2. p. 247. (v. s. in herb. Mus. Par.)

4. C. *ASARIFOLIA* (Lini! spec. 913.) foliis glabris petiolatis cordato-orbiculatis subsinuato-dentatis, caule erecto, siliquis erectis pedicello duplò longioribus. ♀ in rivulis humidis montanis Pedemontii, Galloprovinciæ, Etruriæ, etc. Sims. bot. mag. t. 1735. Petala alba paulò majora quàm C. *amaræ*. Siliquæ pollicem longæ. DC. syst. 2. p. 247. * (v. v.)

℞. *diversifolia*, foliis caulinis pinnatim trisectis radicalibus et summis orbiculatis indivisis. In Pedemontii valle Lanzo.

5. C. *INDICA* (Burm! fl. ind. 140.) foliis ovatis crenulatis petiolatis glabris, summis oblongo-cuneatis, caule erecto, siliquis patulis, stigmate sessili. ① in Javâ. Flores minimi albi. An potiùs inter *Nasturtia* ad sectionem *Clandestinarie* referenda? DC. syst. 2. p. 248. * (v. s. in h. Deless.)

6. C. *STYLOSA* (DC. syst. 2. p. 248. *) foliis caulinis sagittato-semi-amplexicaulibus oblongis acutis denticulatis glabris, caule erecto, siliquis patulis stylo acuminatis. — in Novâ-Hollandiâ. Flores parvi albi. Siliquæ rectangulo-patentes, strictæ, lineares, glabræ, stylo ferè bilineari apiculatæ, pollicem longæ. (v. s. in h. Lamb.)

7. C. *CHENOPODIIFOLIA* (Pers. ench. 2. p. 195.) foliis ovatis sinuato-sublobatis petiolatis glabris, caule procumbente, siliquis erectis, stigmate sessili. — In Americâ austr. Poir. suppl. 2. p. 394. * Flores albi. Siliquæ erectæ, læves,

lineares, compressæ, pollicem longæ. DC. syst. 2. p. 249.* (v. s. in h. Mus. Par.)

8. *C. BELLIDIFOLIA* (Lin! spec. 913.) foliis glabris crassiusculis, radicalibus petiolatis ovatis integris, caulinis paucis integris aut subtrilobis basi non auriculatis, siliquis erectis, stigmatè sessilib. $\frac{1}{2}$ in summis montibus Europæ. DC. syst. 2. p. 249.* (v. v.)

a. petiolaris, petiolis limbo longioribus, limbis integris. OEd. fl. dan. t. 20.—Lim. fl. lapp. 260. t. 9. f. 2. in fissuris rupium humidis Lapponiæ.

β. Alpina, petiolis limbo ferè brevioribus, limbis integris. *C. bellidifolia* Crantz. fl. austr. 43. Wulf. in Jacq. misc. 1. p. 148. t. 17. f. 2.

γ. subtriloba, foliis superioribus subtrilobis, interdum subtrisectis.

§. 2. Foliis plerisque trilobis.

9. *C. RESEDIFOLIA* (Lin! spec. 913.) foliis glabris membranaceis petiolatis, radicalibus indivisis, caulinis in auriculam acutam basi utrinquè productis, inferioribus trisectis, superioribus pinnatim 5-lobis, siliquis erectis stylo brevissimo apiculatis. $\frac{1}{2}$ ① ② in locis umbrosis humidis Alpium. DC. syst. 2. p. 250. Sturm. fl. germ. ic. Arabis resedifolia Lam. fl. fr. 2. p. 511. (v. v.)

β. integrifolia. A. hastulata. Bert. ined. in Apenninis Appuanis.

10. *C. UNIFLORA* (Michx! fl. bor. am. 2. p. 29.) foliis radicalibus 3-lobis glabris, scapis 1-floris. — in rupibus Kentucky. Flores minores quàm *C. amaræ*. Petala calyce longiora. Siliquæ lineares, compressæ. DC. syst. 2. p. 251.* (v. s. in h. Mus. Par.)

11. *C. MICROPHYLLA* (Adams. mém. soc. nat. mosc. 5. p. 111.* Fisch. in litt. ic.) foliis 3-sectis aut quinato-pinnatisectis glabris, florali unico simplici tridentato, scapo paucifloro. — In Sibiria. *C. minuta* Willd. herb. ex Stev. obs. ined. Pedicelli filiformes, ex icone Fischerianâ floræquales, ex cl. Adams flore et siliquâ longiores. DC. syst. 2. p. 251.*

12. *C. BONARIENSIS* (Pers! ench. 2. p. 195.) foliis glabris petiolatis trisectis, lobis petiolulatis subrepandis, medio subtrilobo, summis simplicibus trilobiæve, pedicellis bracteatis. — In america austr. *C. repanda* Smith! herb. Flores parvi, albi. Siliquæ graciles, erectæ, compressæ. DC. syst. 2. p. 251.* (v. s. in h. Mus. Par. et Juss.)

13. *C. ANTEMQUANA* (Burchell! cat. geogr. pl. afr. austr. n. 6043.) foliis supernè pilosiusculis trisectis, segmentis petiolulatis ovatis dentatis, siliquis erectis, caulibus subdecumbentibus. — ad caput Bonæ Spei. Flores albi, magnitudine *C. parvifloræ*; petala obtusa, calyce duplò longiora. DC. syst. 2. p. 252.*

14. *C. BORBONICA* (Pers! ench. 2. p. 195.) foliis utrinquè pilosis subtus canescentibus trisectis, rariùs pinnatisectis, segmentis petiolulatis ovatis acuminatis dentatis, siliquis erectis. — in insulâ Borboniæ. Poir. suppl. 2. p. 394.* DC. syst. 2. p. 252.* *C. rubifolia* Smith! herb. (v. s.)

β. Allevia, foliis interdum pinnatisectis bijugia cum impari. *C. Africana*, ex insulâ Borboniæ Vahl! symb. 2. p. 77.

γ. Arabica, caule subvillosa. *C. Africana*, ex Arabia Vahl! symb. 2. p. 77.

15. *C. AFRICANA* (Lin. spec. 914.) foliis glabris trisectis, segmentis petiolulatis ovatis acuminatis dentatis siliquis patentibus. $\frac{1}{2}$ ad caput Bonæ Spei. — Pluk. alm. 252 t. 101. f. 5. Flores parvi albi. Siliquæ patentæ, lineares. DC. syst. 2. p. 252.* (v. s. in h. Mus. Par.)

16. *C. TRIPOLIA* (Lin! spec. 913.) foliis glabriusculis trisectis, segmentis sessilibus rhomboideo-subrotundis dentatis, scapo nudo, ramis infimis radiciformibus repentibus. $\frac{1}{2}$ in umbrosis montanis Italiæ, Hungariæ, Germaniæ. Sturm. deutschl. fl. ic. Jacq. fl. austr. t. 27. Petala alba, ungue lato cuneato. DC. syst. 2. p. 253.* (v. s. et v.)

17. *C. CHILENSIS* (DC. syst. 2. p. 254.*) foliis supernè pilosiusculis trisectis, segmentis subpetiolulatis ovato-lanceolatis crenatis, caule subadscendente. —

in Chili. Petala oblonga alba, calyce longiora; siliqua linearis, glabra, gracilis, apice in stylum vix à siliquâ distinguendum desinens. (v. s. in h. Lamb.)

18. *C. TUBEROSA* (DC. syst. 2. p. 254.*) foliis glabris, radicalibus longè petiolatis reniformibus, caulinis pinnatim trisectis, radice tuberosa. $\frac{1}{2}$ in Chili. *Sisymbrium tuberosum* Lag! in litt. Petala sepalis triplò longiora. Siliqua junior linearis, stylo brevi acuminata. Deless! ic. sel. 2. t. 29. (v. s.)

§. 3. *Foliis plerisque pinnatipartitis.*

19. *C. GRANULOSA* (All. auct. p. 16.* excl. Dalech. syn.) foliis radicalibus petiolatis ovato-subcordatis, caulinis pinnatipartitis lobis oblongis integris, radice tuberoso-granulosâ. $\frac{1}{2}$ in pratis collium Taurinensium. Petala alba, obovata obtusa, magnitudine *C. amaræ*. (v. s.)

20. *C. AMARA* (Lin. spec. 915.) foliis pinnatisectis, radicalium segmentis subrotundis, caulinorum dentato-angulatis, stylo filiformi acutiusculo, caule basi radicante. $\frac{1}{2}$ ad rivulos Europæ. Vill. Dauph. 3. p. 362. t. 39. Smith. engl. bot. t. 1000. DC. syst. 2. p. 255.* variat fl. pleno. (v. v.)

♂. *trisecta*, foliis trisectis. *C. trifolia* Wahl. fl. lapp. n. 327. ex Wahl. carp. n. 656.

γ? *umbrosa* (Lej. fl. spa. 2. p. 63.) caule foliisque villosis. *C. hirsuta*. OEd. fl. dan. t. 148?

21. *C. ULIGINOSA* (Bieb. suppl. p. 438.*) foliis pinnatisectis, segmentis angulato-dentatis obtusiusculis, radicalium subrotundis, caulinorum oblongis, stylo brevissimo siliquâ vix angustiore, stolonibus repentibus. $\frac{1}{2}$ in sylvis montanis Tauriæ. *C. amara* Bieb. fl. taur. n. 1283. excl. syn. Flores *C. amaræ*; pedicellis siliquis ferè longitudine æquales. DC. syst. 2. p. 256. (v. s.)

22. *C. PROCRÆNE* (Fisch. in litt. 1819.) foliis pinnatisectis, segmentis ovatis subintegris, terminali rotundato subtrilobo, flagellis repentibus, caule adscendente pubescente, siliquis puberulis pedicello brevioribus. $\frac{1}{2}$ in Sibiria Transbaicalensi. *C. hirsuta* Pall. in h. Willd. ex Stev. obs. ined. Siliquæ erectæ tenues, pilis sparsis puberulæ, stylo filiformi lineam longo. DC. syst. 2. p. 256.* (v. s.)

23. *C. PRATENSIS* (Lin! spec. 915.) foliis pinnatisectis, segmentis radicalium subrotundis, caulinorum linearibus lanceolatisve integris, stylo brevissimo vix siliquâ tenuiore, stigmatate capitato. $\frac{1}{2}$ in pratis humidis Europæ totius, Asiæ bor. etc. Lam. ill. t. 562. f. 1. Smith. engl. bot. t. 776. variat fl. pleno. DC. syst. 2. p. 257.* (v. v.)

♂. *grandiflora*, *C. Bauh.* pin. 104.

γ. *debilis*, foliorum superiorum segmentis angustissimis, caule debili.

δ. *stolonifera*, *C. pratensis monstrosa* Naumburg. in Rœm. arch. 2. p. 14. t. 2.*

24. *C. DENTATA* (Schult. obs. n. 968.) foliis pinnatisectis, segmentis petiolulatis, radicalium subrotundis, caulinorum oblongis basi attenuatis, ultimo cuneato, siliquis obliquè erectis latitudine stylum adæquantibus. $\frac{1}{2}$ secus rivulos Galiciæ, Volhiniæ, Podoliæ. *C. Bachtormensis* Willd. h. ex Stev. obs. ined. Flores albi, ferè *C. amaræ* aut uliginosæ. DC. syst. 2. p. 258.* (v. s.)

25. *C. PENSYLVANICA* (Muhl. cat. p. 60.) foliis pinnatisectis aut pinnato-lyratis, lobis ovalibus angulato-dentatis obtusis, caule erecto, petalis oblongo-linearibus. $\frac{1}{2}$ in pascuis Americæ bor. Poir. suppl. 2. p. 395.* Flores minimi albi; petala oblongo-linearia calyce paulò longiora. DC. syst. 2. p. 258. (v. s.)

26. *C. VIRGINICA* (Lin. spec. 915.) foliis pinnatisectis, segmentis lanceolatis, subauriculatis, caule erecto, siliquis longis strictè erectis. $\frac{1}{2}$ in pascuis depressis Americæ bor. Pluk. alm. t. 101. f. 4. — DC. syst. 2. p. 258. (v. s.)

27. *C?* *TERES* (Michx! fl. bor. am. 2. p. 29.) foliis pinnati-partito-lyratis, lobis ovali-oblongis, terminali subtrilobo, siliquis brevibus erectis teretibus. $\frac{1}{2}$ in inundatis à Novâ Angliâ ad Novam Cesaream. Flores ignoti. An potius *Nasturtii* species? DC. syst. 2. p. 259. (v. s. in h. Mus. Par.)

28. *C. HIRUTA* (Lin! spec. 915.) foliis pinnatisectis, radicalium segmentis subrotundis mucronatis petiolulatis, superiorum oblongis subsessilibus, petalis oblongis, siliquis erectis. ① in sylvaticis umbrosis Europæ totius, Tauriæ et Persiæ. Smith. engl. bot. t. 492. DC. syst. 2. p. 259. (v. v.)
β. maxima, *C. hirsuta maxima* Fisch. cat. gor. 1808. p. 81.
γ. prolifera, St. Hil. not. orl. p. 35.
29. *C. SYLVATICA* (Link in Hoffm. phyt. blätt. 1. p. 50.) foliis pinnatisectis, segmentis irregulariter sinuato-dentatis mucronatis, petalis oblongis, siliquis patentibus latitudine stylum æquantibus glabris, rachi hirsutâ. ① in sylvaticis Europæ totius. *C. impatiens*. OEd. fl. dan. t. 735. — DC. syst. 2. p. 260.
30. *C. UMBROSA* (Andrz. cruc. ined. in DC. syst. 2. p. 260.) foliis pinnatisectis, segmentis integris mucronatis, petalis oblongis, siliquis subpatulis stylum crassum latitudine duplò superantibus rachique glaberrimis. ① in sylvaticis Europæ. Huc forsân pertinet *C. tenella* Clark. voy. vol. 2. p. 2 (v. s.)
31. *C. PARVIFLORA* (Lin! spec. 919.) foliis pinnatipartitis, lobis sessilibus oblongis linearibusve integris infimis à caule remotis, petalis oblongo-linearibus, pedicellis subpatulis, siliquis erectis. ① in herbosis humidis Galliæ et in Transsylvania. Gmel. fl. sib. 3. p. 270. n. 37. t. 64.* Wrangel diss. Stockh. 1823.* valdè accedit ad *C. hirsutam*, sed gracilior decumbens. DC. syst. 2. p. 261. (v. v.)
32. *C. IMPATIENS* (Lin! spec. 914.) foliis pinnatisectis, segmentis ovali-oblongis subdentatis, infimis cauli approximatis acutis stipulæformibus. ① in umbrosis sylvaticis montosis Angliæ, Galliæ, etc. Smith. engl. bot. t. 80. Flores parvi, nunc apetalî, nunc petalis albis oblongis parvis donati. DC. syst. 2. p. 261. (v. v.)
β. eriocarpa, siliquis pilosiusculis. *C. dasycarpa* Bieb. suppl. 437. — in Iberiâ occidentali.
33. *C. LATIFOLIA* (Vahl. symb. 2. p. 77.*) foliis pinnatisectis glabris, segmentis 3-7 suborbiculatis dentato-angulatis, siliquis erectis pedicello paulò longioribus stylo acuminatis. ② Secus rivulos in montanis Europæ australis. *C. chelidonia* Lam! dict. 2. p. 183. excl. Barr. syn. — DC. syst. 2. p. 262. (v. v.)
β. Legionensis, foliorum segmentis septem. — in Hispaniâ.
γ. Calabrica, segmentis foliorum minoribus, pedicellis brevioribus. — in regno Neapolitano propè Scyllam. (v. s.)
34. *C. MACROPHYLLA* (Willd. spec. 3. p. 484.*) foliis pinnatisectis subpubescentibus, segmentis 5 ovali-lanceolatis acuminatis inæqualiter serratis. ② in Sibiria. Dentaria hirsutula Andrz. in litt. — Gmel. fl. sib. 3. p. 269. t. 62.* — Flores magnitudine et colore *C. pratensis*. — Habitus ferè Dentariæ et potiùs præeunte à Cl. Andrz. ad hoc genus transferenda? DC. syst. 2. p. 263.
35. *C. CHELIDONIA* (Lin. spec. 914.) foliis pinnatisectis subglabris, segmentis petiolulatis ovatis dentatis, inferioribus pinnatisectis, segmentulis 3-4. ② in sylvis montanis agri Neapolitani, etc. Waldst et Kit. pl. hung. 2. p. 149. t. 140. Petala purpurea unguiculata ovalia. DC. syst. 2. p. 263. (v. s.)
36. *C. PETIOLARIS* (DC. syst. 2. p. 264.*) foliis longè petiolatis glabris pinnatisectis, segmentis 9-10 petiolulatis ovatis dentatis sublobatis, siliquis erectis. — Ad Caucasum septentrionalem propè Mosdock. *C. chelidoniæ* affinis. Goldbl in litt. Flores albi magnitudine ferè *C. thalictroidis*. Deless! ic. sel. 2. t. 30. f. B. (v. s.)
37. *C. PECTINATA* (Pall! herb. et DC. syst. 2. p. 264.*) foliis petiolatis glabris pinnatisectis, segmentis 7-9 petiolulatis ovatis pectinato-lobatis, siliquis patentibus. ①? in Persiâ, Flores albi parvi. Petala oblonga obtusiuscula. (v. s. in h. Lamb.)
38. *C. HEDERACEA* (DC. syst. 2. p. 264.*) foliis petiolatis glabris pinnatisectis, segmentis 5 petiolulatis latè ovatis obtusè trilobis, siliquis erectis ②? in Syriâ. Flores parvi ferè *C. parvifloræ*. Siliquæ tenuissimæ, seminibus subtorulosæ. Deless! ic. sel. 2. t. 30. f. A. (v. s.)

39. *C. GLACIALIS* (DC. syst. 2. p. 264. *) foliis petiolatis pinnatisectis subciliatis, segmentis inferioribus sessilibus bijugis minimis, terminali maximo reniformi petiolato, siliquis erectis. \mathcal{U} in locis ndia Terræ del Fuego ferè sub nive. *Sisymbrium glaciale* Forst! comm. Gætt. 9. p. 32. * Flores albi; siliquæ subfiliformes. (v. s. in h. Banks.)

β . foliis omnibus unijugis cum impari. (v. s.)

40. *C. DEBILIS* (Banks! mss. et herb.) foliis petiolatis glabris pinnatisectis, segmentis petiolulatis ovatis subsinuatis subbijugis cum impari reniformi, caule debili, siliquis erectis. (1) ? in Novâ Zelandiâ. *Sisymbrium heterophyllum*. Forst! prodr. p. 64. n. 250. — DC. syst. 2. p. 265. (v. s.)

41. *C. SARMENTOSA* (Forst! fl. app. p. 92. n. 529.) foliis petiolatis glabris pinnatisectis, segmentis 5 ovatis infernè dentatis, infimis petiolulatis, caule subnudo basi flagellifero, siliquis patentibus. \mathcal{U} in insulâ Teantèa maris Australis. Flores parvi albi, pedicelli filiformes demum patentes. DC. syst. 2. p. 265. (v. s. in h. Lamb.)

42. *C. THALICTROIDES* (All! ped. n. 951. t. 57. f. 1) foliis petiolatis glabris, superioribus pinnatisectis, segmentis 3-5 petiolulatis ovatis trilobatis, caule debili, siliquis patulis. (2) in sylvaticis alpium Delphinatûs et Pedemontii. *C. Plumieri* Vill. fl. Dauph. 3. p. 359. t. 38. * excl. Bocc. syn. Flores vix paulò minores quàm *C. amaræ*, albi cum fauce et ungue subflavescente. DC. syst. 2. p. 266. * (v. s.)

43. *C. GLAUCA* (Spreng! in litt. 1819) foliis petiolatis glabris glaucis subcanosis pinnatisectis, segmentis 5-9 oblongis, terminali trilobo, caule diffuso ramosissimo, siliquis erectis glabris. \mathcal{U} in Calabriâ. *C. Bergeriana*. Andr! in litt. 1820. Flores albi magnitudine ferè *C. amaræ*. DC. syst. 2. p. 266. * Deless! ic. sel. 2. t. 31. (v. s.)

44. *C. MARITIMA* (Portenschlag! in litt.) foliis petiolatis glabris glaucis pinnatisectis, segmentis 5-7 obovato-cuneatis repandis, caule diffuso ramosissimo, siliquis erectis glabris. (1) in Dalmatiæ peninsula Sabioncello (olim Hyllis). Welden. ic. ined. t. 12. Deless! ic. sel. 2. t. 32. Valde accedit ad *Pteuroneurum carnosum* et præsertim ad *C. glaucam*, sed ab utraqûe differt. (v. s.)

45. *C. DIGITATA* (Richards. in Frankl. narr. jour. 743) foliis digitatim pinnatisectis, lobis 7 sessilibus linearibus integerrimis, stylo brevi vix siliquâ tenuiori, stigmate capitato. \mathcal{U} in sterilibus ad mare Arcticum. Glaberrima dodrantalis simplex basi repens ascendens.

†. *Cardamines non satis notæ.*

46? *C? MULTIFIDA* (Banks! herb. Pursh. fl. bor. am. 2. p. 440.) pubescens, ramosa, foliis interruptè pinnatisectis, segmentis bipinnatifidis laciniis rotundatis incisis, siliquis pedicello brevioribus. \mathcal{U} in Floridâ orientali. An potiùs *Brachylobi* species? DC. syst. 2. p. 267. * (v. s. in h. Banks.)

47. *C? MENZIESII* (DC. syst. 2. p. 267. *) pube stellatâ canescens, foliis pinnatisectis, segmentis bipinnatifidis, laciniis parvis acutis, siliquis pedicelli longitudine. — In Americâ bor. Flores ignoti. An potiùs cum priore *Brachylobi* *Sisymbrii* species?

48. *C? GERANIIFOLIA* (DC. syst. 2. p. 268. *) glaberrima, foliis subbipinnatisectis, segmentis inciso-lobatis acutis, siliquis subulatis erectis. — Ad fretum Magellanicum. *Sisymbrium geraniifolium* Poir! dict. 7. p. 218. * Petala alba obovata. (v. s. in h. Mus. Par.)

49. *C. SCUTATA* (Thunb. act. lin. soc. 2. p. 399.) foliis radicalibus petiolatis trisectis, caulinis paucissimis, lobo terminali subrotundo. — in Japoniâ. DC. syst. 2. p. 268.

50. *C. NAMOROSA* (Lejeun. fl. spa. 2. p. 62.) \mathcal{U} villosissima, foliis pinnatisectis, segmentis subrotundo-augulatis. — in sylvis Belgii Leodensis circa Malmedy. Flores terminales, pauci, corymbosi, paulò majores et colore purpureo intensiores quàm *C. pratensis*. DC. syst. 2. p. 268.

51. *C. articulata* (Pursh. fl. bor. am. 2. p. 439.) caule nudo, foliis radicalibus lanceolatis inciso-dentatis glabris. $\frac{1}{2}$ in America bor. Siliquæ turgidæ subarticulatæ. DC. syst. 2. p. 268. *
52. *C. occulta* (Horn. b. hafn. suppl. p. 71.) foliis radicalibus subrotundis, caulinis pinnatis ternatimve sectis, floribus apetalis. (1) in China. An differt à *Nasturtio Bonariensi*. An saltem *Nasturtii* species? DC. syst. 2. p. 268.
53. *C. heterophylla* (Lapeyr. abr. pyr. 377. *) glabra simplex; foliis radicalibus rotundatis petiolis bidentatis. caulinis lyratis. (2) in Pyrenæis ad apicem montis Pic-du-Midi dicti. An *C. heterophylla* Bory ann. gen. sc. ph. 3. p. 6? Petala calyci clauso albido æqualia. Siliquæ obtusæ cum appendice obtusa. DC. syst. 2. p. 268.
54. *C. propinqua* (Carmich. in trans. lin. soc. 12. p. 507.) Glaberrima, foliis pinnatisectis, segmentis obtusis dentatis terminali majore. in insulâ Tristan da Cunha. Proxima *C. glaciali*.
55. *C. nicolor* (Presl. fl. cech. 136.) foliis pinnatisectis stipulatis, segmentis ellipticis lanceolatis sessilibus, sepalis subrotundis. $\frac{1}{2}$ in pratis humidis Bohemiæ.
56. *C. opich* (Presl. fl. cech. 136.) foliis pinnatisectis stipulatis, segmentis repando-angulatis sessilibus, inferiorum subrotundis, superiorum lanceolatis, sepalis lanceolatis. $\frac{1}{2}$ in pratis humidis Bohemiæ. Variat petiolis hirsutis et glabresculis.
57. *C. reflexa* (Raf. fl. lud. p. 84. n. 269.) prope aquas Louisianæ. Caulis pedalis et ultra Flores albi Petala calyce longiora reflexa. Ad *C. pensylvanicam* forsitan referenda. DC. syst. 2. p. 269.
58. *C. angustifolia* (Raf. fl. lud. p. 84. n. 270.) prope aquas Louisianæ. Species parva foliis pinnatisectis glabris, segmentis lineari-filiformibus, siliquis erectis. D. C. syst. 2. p. 269.

XIII. PTERONEURUM. DC. syst. 2. p. 269.

Siliqua lanceolata, valvis planis enerviis sæpè elasticè dissilientibus, placentis alato-nervosis Funiculi dilatati.

1. *P. carnosum* (DC. syst. 2. p. 270. *) foliorum segmentis ovatis subemarginatis glauciscentibus. $\frac{1}{2}$ inter lapides liberos montium calcareorum Croatiae. *Cardamine carnosus*. Waldst. et Kit. pl. hung. rar. 2. p. 137. t. 129. * Calyx patens. Corolla calyce duplò longior, petalis obovatis albis.
2. *P. græcum* (DC. syst. 2. p. 270. *) foliorum segmentis petiolulatis suborbiculatis dentato-lobatis subæqualibus. (1) in umbrosis montosis Corsicæ, Siciliæ, Græciæ. *Cardamine græca*. Lin! spec. 9. 5. Gærtn. fruct. 2. t. 143. Flores albi vix magnitudine *Cardamines amaræ* (v. s.)

β . *eriocarpum*, siliquis pilosis longis rigidis strigosis. In Calabria ceteriora.

XIV. DENTARIA. Fourn. inst. 225. t. 111. Lin. gen. n. 811. Lam. ill. t. 562.

Siliqua lanceolata, valvis planis enerviis sæpè elasticè dissilientibus, placentis non alatis. Funiculi dilatati. Semina ovata immarginata, 1-serialia.

§. 1. *Foliis verticillatis, stylo longè exserto, valvulis ad basin styli terminatis vix acuminatis.*

1. *D. polyphylla* (Waldst. et Kit. pl. rar. hung. 2. p. 174. t. 160. *) foliis ternis verticillatis alternisve petiolatis pinnatisectis, segmentis 7-9 approximatis lanceolatis acuminatis serratis. $\frac{1}{2}$ in locis umbrosis sylvarum Croatiae. Flores ampli pallidè ochroleuci. DC. syst. 2. p. 271. * (v. s.)

β . *D. ochroleuca*, Gaud! herb. — in Helvetiâ Rheticâ.

2. *D. anaphylla* (Lin! sp. 912.) foliis tribus verticillatis petiolatis trisectis,

segmentis ovato-lanceolatis acuminatis serratis, staminibus petalorum longitudine. \mathcal{L} in sylvaticis sterilibus montanis Europæ. Jacq. fl. austr. t. 316.* Flores albi, siccæ subochroleuci. DC. syst. 2. p. 272.* (v. s.)

3. *D. GLANDULOSA* (Waldst. et Kit. pl. rar. hung. 3. p. 302. t. 272.) foliis tribus verticillatis petiolatis trisectis, segmentis ovali-lanceolatis acuminatis serratis, axillis glanduliferis, staminibus petalis dimidio brevioribus. \mathcal{L} in sylvis humidis Hungariæ. Petala purpurascens, calyce duplò et amplius longiora. DC. syst. 2. p. 273.* (v. s.)
4. *D. LACINIATA* (Muhl. cat. 60.) foliis tribus verticillatis breviter petiolatis tripartitis, lobis segmentisve linearibus integris aut grossè dentatis. \mathcal{L} in umbrosis à Novâ-Angliâ ad Canadâ. Flores albi. Petala oblonga. DC. syst. 2. p. 273. (v. s.)

β . *minor*, Pluk. mant. 135. t. 435. f. 2. ex Vaill. herb.

5. *D. HETEROPHYLLA* (Nutt. gen. am. 2. p. 66.*) foliis caulinis 2-petiolatis trisectis, segmentis lanceolato-linearibus integris. \mathcal{L} in Pensylv. occid. Flores pallidè purpurei circiter magn. *Cardamine pratensis*. DC. syst. 2. p. 274.

§. 2. *Foliis caulinis alternis palmatim 3-5-sectis.*

6. *D. TENELLA* (Pursh! fl. bor. am. 2. p. 439.) foliis 2 alternis sessilibus trisectis, segmentis lineari-ellipticis integris. \mathcal{L} in Amer. bor. Flores purpurei magnitudine circiter *Cardamine pratensis*. DC. syst. 2. p. 274.* (v. s. in h. Lambert.)
7. *D. DIPHYLLA* (Michx. fl. bor. am. 2. p. 30.) foliis 1-2 alternis breviter petiolatis 3-sectis, segmentis ovato-lanceolatis grossè et inæqualiter serratolobatis. \mathcal{L} in Amer. bor. Sims. bot. mag. t. 1465. DC. syst. 2. p. 274.* — *D. bifolia* Stok. bot. mat. med. 3. p. 443. (v. s. in h. Lamb. et Mus. Par.)
8. *D. MAXIMA* (Nutt. gen. am. 2. p. 66.*) foliis plurimis alternis petiolatis 3-sectis, segmentis latè ovalibus inciso-dentatis, axillis eglandulosis. \mathcal{L} in Pensylvaniâ. Flores pallidè purpurei. DC. syst. 2. p. 275.
9. *D. TRIFOLIA* (Waldst. et Kit. pl. rar. hung. 2. p. 148. t. 139.*) foliis plurimis alternis petiolatis 3-sectis, segmentis ovato-lanceolatis remotè dentatis, axillis glanduliferis. \mathcal{L} in vallibus umbrosis Croatiæ. Petala alba patentia obovata, calyce triplò longiora. DC. syst. 2. p. 275.*
10. *D. DIGITATA* (Lam! dict. 2. p. 268.*) foliis caulinis plurimis alternis petiolatis palmatim 5-sectis, segmentis oblongo-lanceolatis acuminatis grossè serratis. \mathcal{L} in Galliâ, Helvetiâ, Germaniâ. — Clus. hist. 2. p. 122. fig. 1. *Dentaria pentaphyllos* Lin! sp. 912. var. β et γ . Scop. carn. n. 814. Flores albi aut subpurpurascens. DC. syst. 2. p. 276. (v. v.)

§. 3. *Foliis caulinis alternis pinnatisectis.*

11. *D. PINNATA* (Lam. diet. 2. p. 268.* illustr. t. 562. f. 1.) foliis caulinis alternis petiolatis pinnatisectis, segmentis oblongis acuminatis serrato-dentatis. \mathcal{L} in umbrosis sylvaticis Helvetiæ et Galliæ. A bulbiferâ differt foliis superioribus nunquàm indivisis ex axillis non bulbiferis. DC. syst. 2. p. 277.* — *D. pentaphyllos* a. Lin. sp. 912. (v. v.)
12. *D. QUINQUEFOLIA* (Bieb. fl. taur. 2. p. 109. suppl. 436.) foliis caulinis tribus alternis verticillatisve pinnatisectis, segmentis 5-7 oblongo-linearibus grossè dentatis, terminalibus non confluentibus. \mathcal{L} in sylvis Tauriæ et Caucasi. *D. Caucasica* Willd. herb. ex Stev. obs. ined. Aff. *D. pinnatæ*, Hypanicæ et bulbiferæ. DC. syst. 2. p. 277. Deless! ic. sel. 2. t. 33. (v. s.)
13. *D. HYPANICA* (Besser! in litt.) foliis caulinis 3 alternis verticillatisve pinnatisectis, segmentis 5-7 oblongo linearibus dentatis, terminalibus confluentibus. \mathcal{L} in Bessarabiâ et Podoliâ. Aded accedit ad *D. quinquefoliam* ut facilè cum eâ confusa veniat. DC. syst. 2. p. 278.* (v. s.)
14. *D. BULBIFERA* (Lin. sp. 912.) foliis caulinis alternis pinnatisectis, summis indivisis, plerisque in axillis bulbiferis. \mathcal{L} in Succiâ, Angliâ, Galliâ, etc.

Smith. engl. bot. t. 309. Flores albi aut roseo pallidè suffusi. DC. syst. 2. p. 278.* (v. s.)

β. *ptarmicæfolia*, foliis superioribus argutiùs serratis. — Besl. h. Eyst. æst. ord. 7. t. 12. f. 2.

15. D. *MICROPHYLLA* (Willd! sp. 3. p. 479*) foliis caulinis 3 alternis subverticillatisve, omnibus pinnatisectis, segmentis 7-11 lineari-lanceolatis integris aut subdentatis. ½ in sylvis Caucasi Iberici. Bieb. fl. taur. 2. p. 108.* Flores purpurascens ferè D. pinnatæ. DC. syst. 2. p. 279.* (v. s.)

16. D. *TENUIFOLIA* (Ledebour. mem. acad. peters. 5. 1815. p. 547.*) foliis caulinis breviter petiolatis alternis, aliis trisectis aliis pinna 5-sectis, segmentis linearibus acutis integerrimis, radice fibrosâ tuberes subrotundos gerente. ½ in Sibiria. D. trifida Lam. ill. t. 562. f. 2. — Gmel. fl. sib. 3. p. 272. n. 41. t. 65. Flores magnitudine C. pratensis. DC. syst. 2. p. 279. (v. s.)

β. segmentis incisia. — In montibus Uralensibus.

XV. NEUROLOMA. Andræ. in litt.

Calyx erectus basi æqualis, sepalis margine membranaceis. Petala unguiculata; limbo obovato. Stam. libera edentula vix calyce longiora. Siliquæ lanceolatæ sessiles compressæ, valvis planis. Funiculi umbilicales supernè septo adnati. Semina latè marginata. Cotyl. accumbentes. (0=.)

1. N. *ARABIDIFLORUM*, pedicellis vix calycis longitudine, petalorum laminis obovatis, foliis subradicalibus subcarnosis lanceolatis, scapo simplici. ½ in Sibiria. Hesperis Arabidiflora. DC. syst. 2. p. 454. (v. s.)

α. *glabrum*, Arabis grandiflora Lin! amœn. acad. 2. p. 358. t. L. f. 20.*

β. *asperum*, Cardamine nudicaulis Pall! in herb. Lamb.

2. N. *SCAPIGERUM*, pedicellis calycis longitudine, petalorum laminis submarginatis, foliis subradicalibus petiolatis lanceolatis carnosis, scapo simplici subglabro. ½ in Sibiria. Hesperis scapigera. DC. syst. 2. p. 554. Flores violacei magnitudine H. matronalis ex. cl. Adams.

3. N. *NUDICAULIS*, foliis radicalibus glabris lineari-oblongis, caulinis nullis, siliquis lanceolatis inter semina angustatis. ½ in Sibiria. Arabis nudicaulis. DC. syst. 2. p. 240.

Trib. II. *ALYSSINÆ* seu Pleurorrhizæ Latiseptæ. DC. syst. 2. p. 280.

Silicula longitudinaliter dehiscens, septo lato ovali membranaceo, valvis planis aut concavis. Semina compressa, sæpè marginata. Cotyledones planæ accumbentes, septo parallelæ.

XVI. LUNARIA. Lin. gen. n. 809. Gaertn. fruct. 2. p. 288. t. 124.

Silicula pedicellata elliptica aut lanceolata, valvis planis. Funiculi longi, septo adnati. Calyx subhisaccatus. Pet. subintegra. Stam. edentula.

1. L. *REDIVIVA* (Lin! spec. 911.) siliquis lanceolatis utrinquè attenuatis. ½ in sylvis montanis Galliæ, Helvetiæ, etc. Lam. dict. 3. p. 616.* ill. t. 561. f. 1. Flores odori purpurascens. DC. syst. 2. p. 281.* (v. v.)

β. *purpurea*.

γ. *Alpina*, foliis superioribus minùs cordatis, siliculis angustioribus. Taubern. ic. 312.

2. L. *BIENNIS* (Mœnch. meth. 261.) siliculis ellipticis utrinquè obtusis. ② ③ in sylvis montanis Sueciæ, Germaniæ, etc. L. annua Lin! sp. 911. Lam. dict. 3. p. 616. ill. t. 561. f. 2. Flores violaceo-lilacini, inodori. DC. syst. 2. p. 282. (v. v.)

β. albiflora, Moris. oxon. 2. p. 246. n. 2.

γ. Corcyraea, siliculis suborbiculatis. — in ins. Corcyra.

XVII. SAVIGNYA. DC. *syst.* 2. p. 283.

Silicula sessilis elliptica, valvis planis. Funiculi breves liberi. Calyx basi æqualis. Pet. integra. Stam. edentula. Semina latè marginata compressissima.

1. *S. ÆGYPTIACA*. (DC. *syst.* 2. p. 283.) ① in Ægypti arenis circà pyramides Saqqarah. *Lunaria parviflora* Delil! ill. fl. æg. p. 19. descr. 104. t. 35. f. 3.* Flores parvi pallide violacei. (v. s.)

XVIII. RICOTIA. Lin. *gen. n.* 810. Lam. *ill. t.* 561.

Silicula sessilis elliptica septo evanido 1-locularis, valvis planis. Funiculi longi liberi. Semina juniora 4, demùm solitaria ferè centralia. Calyx basi bigibbus. Pet. integra. Stam. edentula. DC. *syst.* 2. p. 284.

1. *R. LUNARIA* (DC. *syst.* 2. p. 284.*) foliis subbipinnatisectis, lobis oblongis sinuato-angulatis. ① in Syrià. *R. Ægyptiaca* Lin! spec. 912. Lam. *ill. t.* 561. Silicula ovato-lanceolata, acuta (v. v. et s.)
2. *R. TENUIFOLIA* (Sibth. et Smith. fl. græc. t. 630 ex prodr. 2. p. 17.) foliis subbipinnatifidis, lobis linearibus. ① in Cilicià nunc Caramanià. *Peltaria Caramaniensis* Sibth! in h. Banks. Siliculæ compressæ planæ, obovatæ, obtusæ. DC. *syst.* 2. p. 285.* (v. s. in h. Banks.)

†. *Species non satis nota.*

3. *R. ? CANTONIENSIS* (Lour. fl. coch. ed. Willd. 2. p. 482.*) foliis oblongis pinnato-incisis. — Hab. Cantone Sinarum. *Lunaria Ricotia* Desv. *jour. bot.* 3. p. 174. Siliqua oblonga, compressa, tenuis subsessilis. Flore luteo et fructu polyspermo à Ricotià omninò differt. DC. *syst.* 2. p. 285.*

XIX. FARSETIA. Turr. *fars.* 1765. Desv. *jour. bot.* 3. p. 173.

Silicula sessilis, ovalis aut orbiculata, valvis planis. Semina alata. Calyx basi bisaccatus. Petala integra.

Sect. I. FARSETIANA. Farsetia. DC. *syst.* 2. p. 287.

Petalorum laminæ oblongo-lineares, integræ, ex albo purpurascentes. Stamina omnia edentula. Silicula elliptica. Septum basi fenestratum.

1. *F. ÆGYPTIACA* (Turr. diss. fars. 1765. 4º p. 1. t. 1.) caulibus fruticosis erectis, foliis linearibus adpressè incanis. ⑤ in Ægypto. *Cheiranthus Farsetia* Lin! mant. 94. Desf! fl. atl. 2. p. 89. t. 160.—DC. *syst.* 2. p. 287.* (v. s.)

Sect. II. CYCLOCARPÆA. DC. *syst.* 2. p. 287.

Petalorum laminæ oblongæ subemarginatæ. Stamina minora dente aucta. Silicula orbicularis, glabra. Septum integrum.

2. *F. SUFFRUTICOSA* (DC. *syst.* 2. p. 287.*) caulibus basi suffruticosis erectis, foliis lanceolatis tomentosis. ⑤ in Persià. *Lunaria suffruticosa*. Vent! h. cels. t. 19.* Flores subcernui inodori violaceo-lilacini. Sepala velutina (v. v.)

Sect. III. FIBIGIA. DC. *syst.* 2. p. 288. — *Med. gen.* 1. p. 90. t. 2. f. 23.

Petalorum laminæ ovatæ aut ovals flavæ integræ. Stamina minora dente aucta. Silicula elliptica. Septum integrum. DC. l. c.

3. *F. LUNARIOIDES* (Brown. in h. Kew. ed. 2. v. 4. p. 96.) caulibus suffruticosis ascendentibus, foliis oblongo-obovatis petiolatis siliculisque tomentosis incanis. ⑤ in Archipelagi Græci insulis Stenosa Philocandro et Anaso. *Lunaria græca*. Willd. *enum.* 2. p. 675. — Tourn! *itin. ed. gall.* 1. p. 242. t. 30.

Flores lutei, limbo petalorum ovali, sepalis albidis. DC. syst. 2. p. 288.* (v. s. in h. Mus. Par. et Banks.)

4. *F. ERIOCARPA* (DC. syst. 2. p. 288.) caulibus erectis basi suffruticosis, foliis oblongis, siliculis dense sericeo-lanatis. ♂ in insulâ Cypri. Valde aff. *F. clypeatæ*, sed differt siliculis non pube brevi, sed villis longis albis simplicibus confertissimis lanato sericeis. Deless! ic. sel. 2. t. 34. (v. s. sine flore.)
5. *F. CLYPEATA* (R. Brown. in h. Kew. ed. 2. v. 4. p. 96.) caulibus herbaceis erectis, foliis oblongis repandis, siliculis pube brevi velutinis, stigmatè capitato. ② in saxosis collibus Europæ australis. *Alyssum clypeatum* Lin. spec. 909. Schkuhr. handb. 2. n. 1815. t. 181. Petala flava, oblonga, obtusè truncata. DC. syst. 2. p. 289.* (v. s.)
 - a. *ebracteata*. — Boerh. ind. alt. 2. 6. n. 7.
 - β. *bracteosa*. — Boerh. ind. alt. 2. 7. n. 10.
6. *F. CHEIRANTHIFOLIA* (Desv. journ. 3. p. 173.) caule erecto herbaceo, foliis lanceolatis hirtis integerrimis, siliculis pube brevi velutinis, stigmatè bifido. — in Oriente. *Alyssum cheiranthifolium* Willd. spec. 3. p. 468.* Simillima præcedenti sed folia minus incana. DC. syst. 2. p. 290. †.
7. *F. TRIQUETRA* (DC. syst. 2. p. 290.*) caulibus basi suffruticulosus, ramis triquetris ascendentibus, foliis canescentibus, radicalibus obovatis petiolatis, caulinis oblongo-lanceolatis, stylo longo deciduo, stigmatè simplici. ♂ in Dalmatiâ ad rupes arcis Clissa. Welden. ic. ined. t. 11. Silicula elliptica, stylo deciduo mutica. (v. s.)

XX. BERTEROA. DC. syst. 2. p. 290.

Silicula sessilis, elliptica aut obovata, valvis planis concavisve. Calyx basi æqualis. Petala limbo bipartito. Stamina minora dentata. Semina ovalia, plana, brevissimè marginata.

1. *B. INCANA* (DC. syst. 2. p. 291.*) siliculis pubescentibus subventricosis. ② in ruderalis arenosis Europæ. *Alyssum incanum* Lin! spec. 978. *Draba cheirifolia* Berg. phytom. ic. Silicula ovali-oblonga, valvis membranaceis convexis pubescentibus, adultis subglabris. (v. v.)
 - β. *prolifera*. in Tauriâ.
2. *B. MUTABILIS* (DC. syst. 2. p. 292.*) siliculis compresso-planis ellipticis glabris. ♂ ♀ in insulâ Corcyrâ. *Alyssum mutabile*. Vent! h. cels. t. 85.* ad *B. incanam* proximè accedit. (v. s.)
3. *B. OBLIQUA* (DC. syst. 2. p. 292.*) siliculis planis ellipticis pubescentibus ♂ in campestribus Romæ, Neapoli, Siciliæ, etc. *Alyssum obliquum* Sibth. et Smith fl. græc. t. 623. ex prod. 2. p. 12. Inter duas priores omninò media. (v. s.)
4. *B. ORBICULATA* (DC. syst. 2. p. 293.) siliculis planis obovato-orbiculatis pubescentibus. ♂ in arenosis ad Xerxis canalem. Flores minores quàm in prioribus. (v. s.)
5. *B. PERUVIANA* (DC. syst. 2. p. 293.*) siliculis ovato-oblongis pube minimâ scabris, pedicellis bracteatis. ♂ ♀ in Peruvix frigidissimis. Petala calyce duplò longiora, limbo obcordato. (v. s. in h. Lamb.) †.

XXI. AUBRIETIA. Adans. fam. 2. p. 420. DC. syst. 2. p. 293.

Silicula oblonga, valvis convexis. Semina immarginata. Calyx basi bisaccatus. Petala integra. Stamina minora dentata.

1. *A. DELTOIDEA* (DC. syst. 2. p. 294.*) pedicellis calyce longioribus. ♀ in saxosis asperis montosis agri Neapolitani, Græciæ, etc. *Alyssum deltoideum* Lin! spec. 908. Curt. bot. mag. t. 126. Petala calyce duplò longiora, longè unguiculata, limbo purpurascente (v. s.)
2. *A. PURPUREA* (DC. syst. 2. p. 294.*) pedicellis calyce brevioribus. ♀ in Olympi Bithyni cacumine. *Arabis purpurea*. Sibth. et Smith. fl. græc. t. 643. ex

Smith. prodr. 2. p. 28. Flores magnitudine colore et formâ prioris. Eructus desideratur. (v. s. in h. Banks.)

XXII. VESICARIA. Lam. ill. t. 559. DC. syst. 2. p. 295.

Silicula globosa, inflata, valvis hemisphæricis. Semina plurima (ultrâ 8), sæpius marginata. Petala integra.

Sect. I. VESICARIANA. DC. syst. 2. p. 296.

Silicula globosa, valvis membranaceis inflatis.

1. V. UTRICULATA (Lam. ill. t. 559.) calyce bisaccato, foliis oblongis integerrimis glabris, inferioribus ciliatis subspathulatis. ♂ in rupibus calcariis Banatûs, Græciæ, Italiæ, Galliæ. Flores flavi ferè Cheiranthi Cheiri. DC. syst. 2. p. 296. (v. v.)
2. V. RETICULATA (Lam. ill. t. 559. f. 2.) calyce....., foliis oblongo-linearibus glabris, radicalibus rosulatis exsertè dentatis, caulinis integris. ♀ In Armeniâ, et Syriâ. Alyssum Vesicaria Lin. spec. 910. Flores breviter racemosi, flavi, parvi. DC. syst. 2. p. 297. (v. s. in h. Mus. Par.)
3. V. LUDOVICIANA (DC. syst. 2. p. 297.*) calyce æquali subpatente, foliis oblongo-linearibus integris pube stellato-lepidotâ canis, caule ad collum suffruticoso. ♀ ♂ in Louisianâ ad ripas fluminis Missouri. Alyssum Ludovicianum Nutt! gen. am. 2. p. 63.* Flores flavi paulò majores quàm Alyssi saxatilis (v. s. in b. Lamb.)
4. V. SINUATA (Poir. dict. 8. p. 570.*) calyce æquali subpatente, foliisque velutinis oblongis integris aut sinuato-dentatis, caule herbaceo. ①② in Hispaniâ, Italiâ, Illyriâ. — A. sinuatum et Alyssum creticum Lin! spec. 910. Petala flava, post thesium albidâ, emarginata. DC. syst. 2. p. 297.* (v. s. v.)

Sect. II. ALYSSEIDES. DC. syst. 2. p. 298.

Silicula ovata, valvis concavis rigidiusculis.

5. V. CRETICA (Poir. dict. 8. p. 570.*) calyce deciduo, foliis oblongis integris repandisve undulatis cano-tomentosis. ♂ in Cretâ. Alyssum Creticum Lin. spec. 910? ex syn. Willd. spec. 3. p. 460 Siliculæ ovato-globosæ incanæ. Semina margine membranaceo cincta. DC. syst. 2. p. 298.* (v. s. in h. Mus. Par.)
6. V. VESTITA (Desv. journ. bot. 3. p. 171 et 183.*) calyce persistente, foliis linearibus acutis tomentosis integris. ♂ in Persiâ. Petala ungue calyce longiore filiformi. Semina immarginata. DC. syst. 2. p. 299.* Deless! ic. sel. 2. t. 35. (v. s.)

†. Species non satis notæ.

7. V. GLOBOSA (Desv. journ. bot. 3. p. 171 et 184.*) foliis lanceolatis acutis integris glabriusculis, siliculis sphæricis subvillosis. — in Americæ borealis aridis. Siliculæ parvæ, stylo filiformi siliculam superante. DC. syst. 2. p. 299.
8. V. FRUTICULOSA (Desv. journ. bot. 3. p. 171 et 184.*) foliis subspathulatis integerrimis albidio-tomentosis, siliculis subglobosis inflatis glabris. ♂ forsan in Archipelagi insulis. DC. syst. 2. p. 299.*
9. V. PANICULATA (Desv. journ. bot. 3. p. 171.) foliis obovatis integerrimis glabris, siliculis ovatis subinflatis. ♂ in insulâ Cretâ. Alyssum paniculatum. Desf. cor. tourn. p. 67. t. 50.* Flores ignoti, siliculæ ovatæ subinflatæ. DC. syst. 2. p. 299.*
10. V? PHYSIPHORA (Andrz. cruc. ined. in DC. syst. 2. p. 300.) foliis oblongis obtusis confertis utrinquè tomentosis, siliculis subglobosis subpubescentibus, stylo brevissimo. Hab..... Semina biseriata marginata funiculo alato basi septio aduato. Genus proprium ex cl. Andrz.
11. V. ARCTICA (Richards. in Frankl. narr. journ. p. 743.) foliis radicalibus obovato-spathulatis integerrimis tomentosis, siliculis orbiculatis inflatis, caulibus suffruticosis. ♂ ad coloniam Omanak Groenlandiæ. Alyssum arcticum. Horn. fl. dan. t. 1520. DC. syst. 2. p. 324. .

12. *V. ARENOSA* (Richards. l. c.) foliis inferioribus subrhombeis obsolete sinuato-dentatis pube stellatâ canescentibus, caulibus teretibus basi suffruticosis, siliculâ pubescente. ♀ in arenosis ad littus arcticum Saskatchewan. Flores flavi. Caules patentes aut adscendentes.

XXIII. *SCHIVERECKIA*. Andr. *cruc. ined. in DC. syst. 2. p. 300.*

Siliculâ ovata, valvis convexis medio longitudinaliter subdepressis. Semina plurima immarginata. Calyx basi æqualis. Petala integra. Stamina majora dentata.

1. *S. PODOLICA* (Andrz! et Bess! in litt.) ♀ in Podoliâ, Wolhyniâ. DC. *syst. 2. p. 300.* Deless! *ic. sel. 2. t. 36. (v. s.)*

XXIV. *ALYSSUM*. DC. *syst. 2. p. 301.* — *Alyssi* sp. *Lin. gen. n. 805. Lam. ill. t. 559.*

Silicula orbicularis aut elliptica, valvis planis aut centro convexis. Semina in quoque loculo 2-4, compressa, interdum alâ membranaceâ cincta. Calyx basi æqualis. Petala integra. Stamina nonnulla dentata.

Sect. 1. *ADYSETON*. Scop. *carn. 2. p. 13.* DC. *syst. 2. p. 302.*

Flores flavi. Semina in quoque loculo 1-2, interdum alâ membranaceâ marginata. Stamina omnia aut nonnulla dentata.

§. 1. *Species perennes aut fruticulosæ.*

1. *A. SAXATILE* (Lin! spec. 908.) caulibus basi suffruticosis subcorymbosis, foliis lanceolatis integris tomentoso-canis, siliculis obovato-orbiculatis 2-spermis, seminibus marginatis. ♀ in Ruthenico imperio ad cataractas Borysthensis et in Podoliâ. Curt. bot. mag. t. 159. DC. *syst. 2. p. 302.* (v. v.)*
2. *A. GEMONENSE* (Lin! mant. 92.) caulibus basi suffruticosis paniculatis, foliis lanceolatis subintegris cano-velutinis, siliculis suborbiculatis 2-4-spermis, seminibus marginatis. ♀ in muris urbis Gemonæ et in Austriâ, Carinthiâ, etc. DC. *syst. 2. p. 303.* Jacq. *ic. rar. 3. t. 503.* *Vesicaria Gemonensis.* Poir. dict. 8. p. 571.* (v. s.)
3. *A. ORIENTALE* (Ard! specim. 2. p. 32. t. 15. fig. 1.) caulibus basi suffruticosis paniculatis, foliis lanceolatis repando-dentatis undulatis tomentosis, siliculis suborbiculatis transversè ovalibus 4-spermis, seminibus marginatis. ♀ in Crêtæ montibus Sphacioticis, etc. Clypeola tomentosa Lin. mant. 92. DC. *syst. 2. p. 303.* (v. s.)*
4. *A. SPATULATUM* (Steph. in Willd. sp. 3. p. 465.*) caulibus basi suffruticosis, foliis spathulato-obovatis petiolatis cano-velutinis, siliculis orbiculatis stylo paulò longioribus, seminibus marginatis. ♀ in montibus Sibiricæ Altaicæ. *Alyssum cretaceum* Adams. mem. soc. nat. Mosc. 5. p. 109.* Petala calyce duplò longiora, limbo emarginato. DC. *syst. 2. p. 304.** Deless! *ic. sel. 2. t. 37. (v. s.)*
5. *A. ARGENTUM* (Vitm. summ. 4. p. 30.) caulibus basi suffruticosis pube stellatâ canis, foliis oblongo-spathulatis subtus argenteis, siliculis ovato-orbiculatis compressis velutinis, seminibus submarginatis. ♀ in rupestribus apricis Pedemontii. *Lunaria argentea* All! ped. n. 901. t. 54. fig. 3.* Petala flava. DC. *syst. 2. p. 304.* (v. s.)*
6. *A. BARTOLONI* (Desv. journ. bot. 3. 172 et 185.*) caulibus basi suffruticosis pube stellatâ canis, foliis oblongo-obovatis subtus argenteis, siliculis ellipticis compressis subcanescentibus, seminibus immarginatis. ♀ in collibus apricis et saxosis circa Sarzanam. *A. argenteum* Bert! amœn. ital. p. 34. excl. syn. Savi. bot. etr. p. 185. n. 569.* Priori valdè affine. DC. *syst. 2. p. 304.* (v. v.)*
7. *A. MURALE* (Waldst. et Kit. pl. rar. häng. 1. p. 5. t. 6.*) caulibus basi suf-

fruticulosus pube stellatâ canescentibus, foliis oblongis subacutis subtus albidis, siliculis ovatis subvelutinis, seminibus immarginatis. $\frac{1}{2}$ 5 in Transylvaniâ. Duobus prioribus affine et forsâ cum iis non malè conjungendum. DC. syst. 2. p. 305. (v. v.)

8. *A. OBRUSIFOLIUM* (Stev! in litt. DC. syst. 2. p. 305.*) caulibus basi suffruticulosus pube stellatâ pubescentibus, foliis obovato-spathulatis obtusis subtus argenteis, siliculis ovato-orbiculatis subvelutinis, seminibus immarginatis. $\frac{1}{2}$ 2 in Tauriâ merid. et in Sibirîâ. Deless! ic. sel. 2. t. 38. *A. murale* Bieb. fl. taur. 2. p. 103. non Kit. (v. s.)

9. *A. ATLANTICUM* (Desf! fl. atl. 2. p. 71. t. 149.*) caulibus basi suffruticosis incanis erectis, foliis lanceolatis incanis pilosisque, racemis simplicibus, siliculis orbiculatis canis, valvis medio convexis. $\frac{1}{2}$ 2 in Cretæ rupibus, et in Atlantîs cacumine. DC. syst. 2. p. 305.* (v. s.)

10. *A. SERPYLLIFOLIUM* (Desf! fl. atl. 2. p. 70.*) caulibus suffruticosis erectis incanis, foliis incanis inferioribus obovatis superioribus lanceolatis, racemis corymbosis, siliculis obovato-cuneatis incanis, valvis planis. $\frac{1}{2}$ 2 in Atlante, et in Hispaniâ. DC. syst. 2. p. 306.* (v. s.)

11. *A. REPENS* (Baumg! fl. trans. 2. p. 237.) caulibus basi suffruticulosus, junioribus decumbentibus, adultis erectis, foliis pilosiusculis, inferioribus oblongo-obovatis in petiolum attenuatis, superioribus oblongis, pedicellis hispidis, stylo puberulo ovario elliptico biovulato longiore, $\frac{1}{2}$ in calcareo-saxosis Transylvaniæ. Radix repens. Silicula orbiculata x-sperma ex cl. Baumg. (v. s.)

12. *A. VERNALE* (Horn: ex Schrank hort. mon. t. 96.) caulibus adscendentibus, foliis lanceolatis deorsum attenuatis subintegerrimis pube stellata subcanescentibus, petalis emarginatis, siliculis inflatis stellato-pilosis. $\frac{1}{2}$ patr. ign. Ad *A. repens* valdè accedit sed siliculæ dicuntur inflatæ.

13. *A. TORTUOSUM* (Waldst. et Kit. pl. rar. hung. 1. p. 94. t. 91.*) caulibus basi suffruticulosus tortuoso-diffusus canescentibus, foliis incanis sublanceolatis, racemis corymbosis, siliculis ellipticis velutino-canescensibus, valvis planis. $\frac{1}{2}$ 2 in Hungariâ, Transylvaniâ, et Podoliâ austr. DC. syst. 2. p. 306.* (v. s. v.)

ß. Orientale, caulibus basi magis suffrutescentibus, foliisque magis incanis. in Armeniâ et Tauriâ. *A. serpyllifolium* Bieb. fl. taur. 2. p. 103. suppl. 432.

14. *A. ALPESTRE* (Lin. mant. 92.) caulibus basi suffruticosis diffusis canescentibus, foliis obovatis incanis, racemis simplicibus, siliculis obovato-oblongis canescentibus, valvis planis. $\frac{1}{2}$ 2 in scopulis apricis montanis Gallo-provinciæ Delphinatûs, etc. All! fl. ped. n. 888. t. 18. fig. 2. Petala flava. DC. syst. 2. p. 307.* (v. s.)

15. *A. NEBRODENSE* (Tineo! pug. pl. sicil. p. 12. n. 10.) caulibus suffruticulosus adscendentibus canescentibus, foliis obovatis canis, racemis corymbosis, siliculis elliptico-obovatis incanis, valvis planis. $\frac{1}{2}$ 2 in Nebrodibus Siciliæ. Species recognoscenda. DC. syst. 2. p. 307.* (v. s.)

16. *A. MARSCALLIANUM* (Andrz. cruc. in DC. syst. 2. p. 308.*) caulibus suffruticosis erectis foliisque canis, inferioribus obovatis superioribus oblongo-linearibus, racemis simplicibus, siliculis ellipticis canescentibus, valvis subconvexis. $\frac{1}{2}$ 2 in lapidosis Caucasi Orientalis et Tauriæ. *A. alpestre* Bieb. fl. taur. 2. p. 102. suppl. p. 432. ex Stev! in litt. Medium inter *A. Atlanticum* et *alpestre*. (v. s.)

17. *A. LENENSE* (Adams mem. soc. nat. mosc. 5. p. 110.*) caulibus subherbaceis diffusis foliisque canis lanceolatis imbricatis, racemis simplicibus brevibus, siliculis ovatis emarginatis, stylo subsequantibus pubescentibus. $\frac{1}{2}$ in Sibirîâ. *A. imbricatum* Schlecht. in h. Willd. ex Stev! obs. ined. Petala obovata subemarginata. DC. syst. 2. p. 308.* (v. s.)

18. *A. LANIGERUM* (DC. syst. 2. p. 308.*) caulibus basi suffruticosis cespitosus

foliisque tomentoso-lanatis obovatis, racemis corymbosis, siliculis ellipticis velutinis stylo ferè æqualibus. 5 \mathcal{L} in Persiâ. Flores parvi flavi. Petala obovata. (v. s. in h. Mus. Par.)

19. *A. MONTANUM* (Lin! spec. 907.) caulibus subherbaceis diffusis pubescentibus, foliis subincanis, inferioribus obovatis, superioribus oblongis, racemis simplicibus, siliculis orbiculatis submarginatis canescentibus. \mathcal{L} in collibus apricis calcaris Europæ ferè totius. Jacq. h. vind. 358 * Curt. bot. mag. t. 419. Petala calyce duplò longiora flava. DC. syst. 2. p. 309. (v. v.)

β . *A. albescens*, Schlecht. in h. Willd. ex Stev. observ. ined. *A. montanum* Bieb. fl. taur. 2. p. 104. ex Stev! in litt.

γ . *A. arenarium*, Lois! fl. Gall. p. 401.

20. *A. CUNEIFOLIUM* (Ten! fl. neap. prod. p. xxxvii. app. cat. neap. 1815. p. 58.) caulibus subherbaceis diffusis pubescentibus, foliis subincanis obovato-cuneatis, corymbis simplicibus, siliculis ovatis submarginatis stylo subæqualibus. \mathcal{L} in montibus summis Aprutii. Valdè accedit ad *A. montanum* et facillè cum eo confundendum est. DC. syst. 2. p. 310. (v. s.)

21. *A. DIFFUSUM* (Ten! app. hort. neap. 1815. p. 58.) caulibus subherbaceis diffusis canescentibus, foliis incanis, infimis subobovatis, ceteris linearilanceolatis utrinquè attenuatis, racemis simplicibus, siliculis ovatis submarginatis stylo subæqualibus. \mathcal{L} in montibus summis Aprutii et Pyrenæorum. An satis à priori distinctum. DC. syst. 2. p. 310. (v. v.)

22. *A. WULFENIANUM* (Willd. enum. suppl. 44.) caulibus subherbaceis diffusis canescentibus, foliis punctato canis basi attenuatis, infimis obovatis, superioribus lanceolatis, racemis simplicibus, siliculis ellipticis submarginatis stylo duplò longioribus. \mathcal{L} in Carinthiâ. Differt ab *A. montano* siliculis non orbiculatis sed ellipticis. DC. syst. 2. p. 310. (v. s. comm. à cl. Hollandre.)

23. *A. SIBIRICUM* (Willd! spec. 3 p. 465.*) caulibus herbaceis adscendentibus supernè ramoso-corymbosis, foliis siliculisque obovatis obtusis pube stellatâ punctatis, stylo silicula triplò brevior. \mathcal{L} in Sibirîâ. Flores flavi, parvi. DC. syst. 2. p. 311.* (v. s.)

24. *A. FISCHERIANUM* (DC. syst. 2. p. 311.*) caulibus herbaceis adscendentibus foliisque canis, inferioribus obovatis, superioribus oblongis, racemis simplicibus, siliculis orbiculatis incanis stylo duplò longioribus. \mathcal{L} in Sibirîâ. Petala flava calyce duplò longiora (v. s.)

25. *A. GNAPHALODES* (DC. syst. 2. p. 311.*) caulibus suffruticosis demissis, foliis ellipticis velutino-tomentosis, ramis linearibus, racemis simplicibus, calycibus clausis pedicello longioribus, siliculis ellipticis velutinis stylo duplò longioribus. \mathcal{L} 5 in Persiâ. Petala flava, ungue filiformi, limbo obovato. (v. s.)

26. *A. PETALODES* (DC. syst. 2. p. 312.*) caulibus suffruticosis erectiusculis foliisque hirsutis oblongis basi attenuatis, infimis dentatis, racemis simplicibus, calycibus clausis pedicello longioribus, siliculis ovatis margine velutinis. 5 \mathcal{L} in Syriâ. Ungues petalorum calyce longiores. (v. s. in h. Labill.)

§. 2. *Species herbaceæ annuæ aut biennes.*

27. *A. UMBELLATUM* (Desv. journ. bot. 3. p. 173 et 184.*) caulibus herbaceis diffusis, foliis linearibus puberulis, racemis subumbellatis, siliculis ellipticis pube stellatâ punctato-pilosis. (I) in Tauriâ merid. *A. brachystachyum* Bieb. suppl. 434.* Flores parvi, sulphurei. DC. syst. 2. p. 312.* (v. s.)

28. *A. ROSTRATUM* (Stev! mem. acad. Petersb. 3. p. 295. t. 15. f. 1.) caule herbaceo erecto, ramis floriferis paniculatis, foliis lanceolatis puberulis, siliculis orbiculato-ellipticis piloso-canis longè racemosis stylo vix longioribus. (I) in Bessarabiâ et Tauriâ merid. Petala saturè lutea calyce duplò longiora spatulata. DC. syst. 2. p. 313.* (v. s. v.)

β . *A. vernale*, Kit. ex Horn. h. hafn. p. 60r. folia linearia via ultrâ lineam lata.

29. *A. MICROPETALUM* (Fisch! in DC. syst. 2. p. 313.) caule herbaceo erecto, foliis lanceolatis siliculisque hirsutis orbiculatis stylo duplò longioribus longè racemosis. ① in Iberiâ mediâ prope Tiflim. Deless! ic. sel. 2. t. 39. *A. parviflorum* Bieb. suppl. p. 434.* (v. s.)
℔? procumbens (Bieb. l. c.) caulibus diffusis et multò minoribus.
30. *A. MIRSUTUM* (Bieb. fl. taur. 2. p. 106.*) caule herbaceo erecto, foliis lanceolatis hirsutis, siliculis orbiculatis tuberculato-hirsutis longè racemosis stylo triplò longioribus. ① in Tauriæ et Iberiæ campis apricis. *A. Draba* Willd. enum. p. 672. ex Stev. obs. in herb. Willd. *A. campestri* valdè affine et cum eo, ut videtur, a plurimis confusum. DC. syst. p. 314.* Deless! ic. sel. 2. t. 40. (v. s.)
31. *A. CAMPESTRE* (Lin! spec. 309.) caulibus herbaceis diffusis, foliis lanceolatis sublinearibusve hirsutis, siliculis orbiculatis tuberculato-hirsutis longè racemosis stylo sextuplò longioribus. ① in campis arenosis ab Hispaniâ ad Tauriam. Sibth. et Smith fl. græc. t. 626. ex prod. 2. p. 13. *Mönchia campestris* Roth. germ. I. 274. II. 84. Petala flava calyce majora subemarginata. DC. syst. 2. p. 314.* (v. v.)
℔. A. simplex Rud. in Schrad. journ. 2. p. 291.*
32. *A. CALYCINUM* (Lin! sp. 908.) caulibus herbaceis diffusis, foliis lineari-lanceolatis canescentibus, calycibus persistentibus, siliculis orbiculatis submarginatis pubescentibus stylo quadruplò longioribus. ① in campis siccis Europæ mediæ et australis. Jacq. hort. viud. 114. fl. austr. t. 338. Petala senescentia albidâ. DC. syst. 2. p. 315.* (v. v.)
33. *A. MINIMUM* (Willd. spec. 3. p. 464. excl. Lin. syn.) caulibus herbaceis diffusis, foliis lineari-lanceolatis incanis, siliculis orbiculatis emarginatis glabris. ① in agro Vindobonensi, Podoliâ et Tauriâ. Tratt. thes. bot. p. 8. t. 35. affine *A. calycino* sed calyx deciduus. DC. syst. 2. p. 316.* (v. s.)
34. *A. MINUTUM* (Schlecht. in h. Willd. ex Stev! obs. ined.) caulibus herbaceis erectiusculis subhispidis, foliis radicalibus ovalibus petiolatis, caulinis oblongo-linearibus, calyce piloso tardè deciduo, siliculis ovato-orbiculatis glabris. ① ad rupes Sibirix? Flores parvi flavi, petalâ ferè linearia. DC. syst. 2. p. 316.* (v. s.)
35. *A. GLOMERATUM* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 1204.) caulibus herbaceis adscendentibus, foliis lanceolatis obtusis villosis, racemo conferto, calycibus persistentibus, siliculis orbiculatis glabris ① ad cap. B-Spri. Petala oblonga, calyce paulò longiora. DC. syst. 2. p. 316. (v. s. comm. à cl. Burch.)
36. *A. FULVESCENS* (Smith prod. fl. græc. 2. p. 13.*) caulibus herbaceis adscendentibus, foliis radicalibus petiolatis obovatis canescentibus, summis oblongis, calycibus persistentibus, siliculis ovato-globosis glabris. ① in Peloponneso et ius. Cypro et Scio. Petala flava in sicco interdum fulvescentia, limbo obcordato. DC. syst. 2. p. 317.* (v. s. in h. Mus. Par.)

Sect. II. ANODONTEA. DC. syst. 2. p. 317.

Flores flavi. Stamina edentula. Siliculæ subinflatæ. — Species inter *Alyssa* et *Vesicarias* mediæ.

37. *A. EDENTULUM* (Waldst. et Kit. pl. rar. hung. 1. p. 95. t. 92.*) caule herbaceo erecto, foliis velutinis oblongis subsinuatis, summis linearibus, calyce patulo, petalis bifidis, siliculâ ovatâ glabrâ. ② in rupibus calcareis Banatûs. *Vesicaria edentula* Poir. dict. 8. p. 572.* Flores flavi. Petala calyce duplò longiora, apice bifido. DC. syst. 2. p. 317.* (v. s.)
38. *A. DASYCARPUM* (Steph. in Willd. spec. 3. p. 469.*) caulibus herbaceis erectis, foliis oblongis canescentibus, pedicellis calyce brevioribus, siliculis stylo duplò ferè longioribus. ① in Sibirix. Flores parvi flavescentes. *A. Sibiricum* Trevir. in ges. nat. fr. berl. mag. 1816. p. 151. t. 2. f. 14 et 17. non Willd. — DC. syst. 2. p. 318.* (v. s.)

Sect. III. LOBULARIA. DC. syst. 2. p. 318.

Flores albi. Semina in loculis sæpius solitaria. Stamina edentula. Caules fruticulosi saltem ad basim.

39. *A. MARITIMUM* (Lam. dict. 1. p. 98.*) caulibus basi suffruticosis procumbentibus, foliis lineari-lanceolatis acutis subincanis, siliculis ovalibus glabris stylo brevi apiculatis. ζ \mathcal{U} in arenosis maritimis secus mare Mediterraneum totam. *A. halimifolium* Lin. sp. 907. non Willd. Curt. bot. mag. t. 101. *Clypeola maritima* Lin. mant. 426. *A. minimum* Lin! sp. 908. Petala obtusa. DC. syst. 2. p. 318.* (v. v.)

β . *Canariense*, caulibus longioribus, habitu toto majore, siliculæ loculis sæpe dispermis. in ins. Canariis. (v. s.)

40. *A. RUPESTRE* (Tenor. fl. nap. t. 60.) caulibus basi suffruticosis suberectis, foliis radicalibus oblongo-lanceolatis acutis argenteis, caulinis paucis linearibus, siliculis obovatis flocculoso-lepidotis stylo brevi mucronatis. ζ in Italiæ australis rupibus. Flores majores quàm *A. maritimi*. DC. syst. 2. p. 319.* (v. s.)

41. *A. HALIMIFOLIUM* (Willd. spec. 3. p. 460. ex syn.) caulibus suffruticosis adscendentibus, foliis oblongis obtusis basi attenuatis lepidoto-puberulis, siliculis orbiculatis glabris stylo duplò longioribus. ζ in montosis saxosis apricis Italiæ. *Lunaria halimifolia* All! ped. n. 900. t. 54. f. 1. et 86. f. 1.* Flores ferè *A. spinosi*. DC. syst. 2. p. 320.* (v. s.)

42. *A. SPINOSUM* (Lin! spec. 907.) caule fruticoso, ramis pedunculisque senescentibus spinosis, foliis oblongo-linearibus argenteis, siliculis orbiculatis glabris stylo brevi apiculatis. ζ in rupestribus calcareis Arragoniæ. — Barr. ic. t. 808. Flores paulò quàm *A. maritimi* majores. — DC. syst. 2. p. 320.*

43. *A. MACROCARPUM* (DC. syst. 2. p. 321.*) caule fruticoso ramoso subspinescente, foliis oblongis obtusis argenteis, siliculis obovato-orbiculatis subemarginatis glabris stylo apiculatis 8-ovulatis. ζ in rupibus calcareis Galliæ australis. Valdè affine *A. spinoso*, *A. halimifolio* et *A. Pyrenaico*, sed ab iis omnibus differt. Deless! ic. sel. 2. t. 41. (v. v.)

44. *A. PYRENAÏCUM* (Lapeyr. abr. Pyr. 371. supp. 91.*) caulibus fruticosis ramosis inermibus, foliis obovatis basi attenuatis cano-tomentosis, siliculis ellipticis villosulis stylo longè acuminatis. ζ in rupibus Pyrenæorum orient. DC. syst. 2. p. 322. *A. halimifolium* β DC. fl. fr. 5. p. 594. (v. s.)

45. *A. CANESCENS* (DC. syst. 2. p. 322.*) caulibus fruticulosis adscendentibus, foliis oblongo-linearibus tomentoso-canis, siliculis ellipticis velutinis stylo longè acuminatis. \mathcal{U} ζ in rupestribus Sibirici, Dahuriæ et Kamtschatkæ. (v. s.)

α . *abbreviatum*, racemo brevi corymboso. *A. montanum* et *A. minutum* Patrini! herb.

β . *elongatum*, racemo subelongato. *A. Dauricum* Schlect. in h. Willd. ex Stev! obs. ined. Differt foliis magis hirsuto-tomentosis.

46. *A. TENUIFOLIUM* (Steph. in Willd. spec. 3. p. 460.) caulibus fruticulosis erectis, foliis linearibus stellato-villosis acutis, siliculis ovalibus hirsuto-canis stylo apiculatis. \mathcal{U} ζ in Sibiria. Petala ungue filiformi limbo obovato obtuso patente. in herb. sæpe cum priore confusum. DC. syst. 2. p. 322.* (v. s.)

Sect. IV. ODONTOSTEMON. DC. syst. 2. p. 323.

Flores albi. Stamina majora dente aucta. — Caules herbacei.

47. *A. HYPERBOREUM* (Lin. spec. 910.) caulibus herbaceis cespitosis, foliis ellipticis grossè et acutè dentatis subhirsutis, siliculis ellipticis. \mathcal{U} in saxis Americæ borealis. Petala alba subrotunda unguibus flavicantibus. DC. syst. 2. p. 323.*

†. *Species non satis notæ.*

48. *A. DENSIFLORUM* (Desf. cor. Tourn. p. 65. t. 48.*) foliis angusto-lanceolatis,

floribus dense in racemum cylindraceum congestis, siliculis orbiculatis monospermis pubescentibus. — in Armeniâ. Flores parvi albi. DC. syst. 2. p. 323.* (v. s. in h. Mus. Par.)

49. A. STRICTUM (Willd. spec. 3. p. 464.*) foliis lineari-lanceolatis tomentosis, floribus in racemum cylindraceum dense congestis, siliculis ellipticis tomentosis. — in Armeniâ. A priore differre videtur siliculis ellipticis nec orbiculatis, tomentosis nec pubescentibus. DC. syst. 2. p. 324.

50. A. OBCORDATUM (DC. syst. 2. p. 324.) foliis lineari-lanceolatis dentatis tomentoso-pubescentibus, siliculis glabris rotundato-obcordatis. — Hab. Aurinia obcordata Desv. journ. bot. 3. p. 162 et 174.* Semina in quoque loculo solitaria fusca membranâ latâ cincta.

51. A. DENTATUM (Nutt. gen. am. 2. p. 63.*) foliis radicalibus subruncinato-dentatis, caulinis lineari-lanceolatis, siliculis ellipticis pubescentibus. ① in declivibus rupium Virginie. Draba arabisans Pursh fl. bor. am. 2. p. 434. non Michx. Flores ignoti. DC. syst. 2. p. 324.

52. A. STRIGOSUM (Russel. in Schrad. jour. 1. p. 426.) caule herbaceo pilis stellatis hispido, foliis obovatis integerrimis, calycibus persistentibus. — Hab. Aleppo. DC. syst. 2. p. 325.

53. A. ? DUBIUM (Rory. ann. sc. phys. 3. p. 4.) caule fruticoso ramoso, foliis obovato lanceolatis tomentosis, siliculis ovato-acutis monospermis. — Inter rupes in excelsis Sierræ-Nevadæ. A. purpureum Lag. ann. 5. p. 275. poll. 1-4. altum.

XXV. MENIOCUS. Desv. journ. bot. 3. p. 173.

Silicula sessilis elliptica, valvis planis. Semina in quoque loculo 6-8 immarginata. Calyx æqualis. Petala integra. Stamina majora dentata. — Genus ab Alyssso vix distinctum nisi seminibus in quoque loculo 6 nec 2-4, siliculis glabris et habitu ferè proprio. DC. syst. 2. p. 325.

1. M. LINIFOLIUS (DC. syst. 2. p. 325.*) foliis linearibus obtusiusculis integerrimis. ① in Tauriâ, Caucasio, Syriâ, Podoliâ, Illyriâ et Hispaniâ. Deless. ic. sel. 2. t. 42. Alyssum linifolium. Steph. in Willd. spec. 3. p. 467.* Petala calyce paulò longiora. Semina rufa. (v. s.)

XXVI. CLYPEOLA. Gaertn. fruct. 2. p. 283. t. 141. f. 6. DC. syst. 2. p. 326.

Silicula orbicularis unilocularis 1-sperma, valvis planis. Calyx æqualis. Petala integra. Stamina dentata. Semina compressa.

Sect. I. JONTHLASPI. DC. syst. 2. p. 326.

Silicula margine ciliata disco glabra aut subpubescens.

1. C. JONTHLASPI (Lin. spec. 170.) caulibus diffusis aut adscendentibus. ① in sabulosis apricis Europæ australis et in Persiâ. Lam. ill. t. 560. fig. 1. Flores flavi minimi. Ovarium monospermum. Variat fructibus confertis minoribus aut majoribus et magis distantibus. DC. syst. 2. p. 326.* (v. v.)

Sect. II. ORIUM. DC. syst. 2. p. 327.

Silicula margine dentata, villis longis mollibus hirsutissima.

2. C. XRIOPHORA (Cav. ex Lag! in litt. DC. syst. 2. p. 327.*) caule erecto simpliciter, calyce persistente. ① in Hispaniâ. Orium lanuginosum. Desv. journ. 3. p. 162. t. 25. fig. 10. Flores in sicco albidii parvi. Petala patentia persistencia, vix calyci æqualia. (v. s.)

Sect. III. BERGERETIA. DC. syst. 2. p. 328.

Silicula margine dentata, setis rigidis in utroque disco echinata.

3. C. ECHINATA (DC. syst. 2. p. 328.*) caule erecto, calycibus deciduis. ① in

Oriente. *Berberetia echinata* Desv. *jour. bot.* 3. p. 161. t. 25. f. 9. Flores minimi. Semen unicum omnino ut in *Cl. Jonthlaspi.* (v. s.)

β. *ramosissima*, foliis oblongis pube minimâ stellatâ sparâ puberulis circa Bagdad. DC. *synt.* 2. p. 328.*

XXVII. PELTARIA. *Lin. gen. n.* 1083. *Gaertn. fruct.* 2. p. 383. t. 141. — *Bohatschia. Crantz. austr. p.* 5. t. 1. f. 1.

Silicula orbicularis, septo evanido unilocularis, 1-4-sperma, valvis planis. Calyx æqualis. Petala integra. Stamina edentula. Semina pendula abortu sæpè solitaria. — *Herbæ glabræ, foliis sagittato-amplexicaulibus, floribus albis.* DC. *synt.* 2. p. 328.

1. P. ALLIACEA (*Lin! spec.* 910.) foliis caulinis sagittato-amplexicaulibus, siliculis planis lævibus. 2℥ in locis umbrosis Austriæ, Transylvaniæ et Pedemontii. Jacq. *fl. austr. t.* 123. *Clypeola perennis.* Ard. *spec.* 26. t. 6. — DC. *synt.* 2. p. 329. (v. v.)

2. P. ANGUSTIFOLIA (DC. *synt.* 2. p. 329.) foliis caulinis inferioribus petiolatis, supremis linearibus, siliculis planis brevibus. 2℥ circa Damascus. Siliculæ ovato orbiculatæ compressæ bi ovulatæ abortu 1-spermæ. (v. s.)

3. P. GLASTIFOLIA (DC. *synt.* 2. p. 330.*) foliis caulinis sagittato-amplexicaulibus, siliculis planiusculis rugulosis coriaceis ①? in Oriente. Flores parvi. Silicula orbiculata, compressa, subrugulosa, monosperma. Deless! *ic. sel.* 2. t. 43. (v. s. in h. Mus. Par.)

XXVIII. PETROCALLIS. *Brown! in h. Kew. ed.* 2. vol. 4. p. 93.

Silicula sessilis ovalis, valvis planiusculis. Semina immarginata in quoque loculo 2, funiculis septo adnatis. Cotyledones ovaes obliquè accumbentes. DC. *synt.* 2. p. 330.

1. P. PYRENAICA (*Brown! h. Kew. ed.* 2. vol. 4. p. 93.) 2℥ in rupibus et inter saxa montium Europæ australis. *Draba Pyrenaica* *Lin! spec.* 896. Jacq! *fl. austr. t.* 228. Flores rosei elegantes. Folia cuneata apice 3-5-fida. DC. *synt.* 2. p. 330. (v. v.)

XXIX. DRABA. DC. *synt.* 2. p. 331. — *Drabæ sp. Lin. Brown.*

Silicula sessilis, ovalis aut oblonga, valvis planis convexisve. Semina plurima, immarginata. Calyx æqualis. Petala integra. Stamina omnia edentula.

Sect. I. AIZOPSIS. DC. *synt.* 2. p. 332.

Plantæ perennes. Scapus nudus. Folia rigida ciliata. Flores 4-avi. Stylus filiformis longitudine varius. (Genus proprium juxta Andr. *litt.* 1827.)

1. D. AIZOIDES (*Lin! mant. gr.*) scapis nudis glabris, foliis rigidis lanceolato-linearibus carinatis ciliatis, staminibus petala subæquantibus, stylo latitudine siliculæ æquante. 2℥ in rupestribus apricis glareosis Europæ Jacq. *fl. austr. 2. t.* 194 DC. *synt.* 2. p. 333. (v. v.)

β. *minor.* *Draba ciliaris* *Dol fl. fr.* 4. p. 697.*

γ. *diffusa.* *Draba ciliaris* *Lin! mant.* 91. — *Ger! gallopr. p.* 344. t. 13. fig. 1.

2. D. BRACHYSTEMON (DC. *synt.* 2. p. 334.*) scapis nudis glabris, foliis elongato-linearibus carinatis ciliatis, staminibus calyceem vix æquantibus. 2℥ hab. *Draba aizoides* *Curt. bot. mag. t.* 170. (v. s. in h. Banks. *Lin.*)

3. D. AZOON (*Wahlenb. fl. carpath. p.* 193. in *adnot.*) scapis nudis glabris, foliis linearibus carinatis rigidis ciliatis, stylo siliculæ hirtæ latitudine dimidio brevior. 2℥ in rupestribus montium inferiorum Baviæ, Austriæ, etc. Siliquæ pedicelli longitudine, pilis brevibus semper hispida. DC. *synt.* 2. p. 334.* (v. s.)

4. D. CUSPIDATA (*Bieb. suppl.* 424.) scapis nudis villosis, foliis linearibus acutiusculis carinatis ciliatis, siliculis lanceolatis hispidis stylo paulò longio-

- ribus. *℥* in Tauriâ, Transylvaniâ? Hispaniâ. *Draba aizoides* Pall! ind. taur. Bieb. fl. taur. 2. p. 92. non Lin. — DC. syst. 2. p. 334. (v. s.)
5. *D. RIGIDA* (Willd! spec. 3. p. 425.*) scapo nudo glabro, foliis oblongis rigidis obtusis carinatis parçè ciliatis, senioribus subreflexis, petalis longitudine calycis, stylo siliculæ glabræ latitudinem non æquante. *℥* in rupestribus Iberiæ. Petala subrotunda calycis ferè longitudine. DC. syst. 2. 335.* †. (v. s. in h. Mus. Par.)
6. *D. BYTOMES* (DC. syst. 2. p. 335.*) scapis nudis glabris, foliis oblongis parvis carinatis parçè ciliatis confertissimis, petalis calyce staminibusque duplò longioribus, stylo siliculæ glabræ latitudinem vix æquante. *℥* in rupestribus Caucasi et Iberiæ. *Draba rigida* Bieb. fl. taur. 2. p. 93.* suppl. 426. Petala oblonga, calyce duplò longiora. DC. syst. 2. p. 335.* (v. s.)
7. *D. OLYMPICA* (Sibth! in h. Banks. et DC. syst. 2. p. 336.*) scapis nudis velutinis, foliis linearibus carinatis confertissimis ciliatis, siliculis ovatis canovillosis, stylo brevissimo. *℥* in monte Olympo. *D. Aizoides* Sibth. et Smith prodr. fl. græc. 2. p. 4? non Lin. Flores ignoti. Siliculæ ovatæ, vix compressæ. (v. s. in h. Banks.)
8. *D. BRUNIÆFOLIA* (Stevl in mem. sc. nat. mosc. 3. p. 268.) scapis nudis pubescentibus, foliis linearibus subcarinatis ciliatis acutis laxè rosulatis, petalis calyce et staminibus duplò longioribus, siliculis ovalibus pubescentibus, stylo brevissimo. *℥* in Caucaso orient. Petala calyce duplò longiora. DC. syst. 2. p. 336.* (v. s.)
9. *D. ERICÆFOLIA* (Stevl mem. sc. nat. mosc. 3. p. 268.*) scapis nudis glabris, foliis linearibus subcarinatis acutis laxè rosulatis, petalis staminibusque duplò longioribus, siliculis elliptico-oblongis glabris, stylo brevissimo. *℥* inter saxa Caucasi orientalis. Priori valdè affinis. DC. syst. 2. p. 336. (v. s.)
10. *D. PILOSA* (Adams ex Fisch! in litt.) scapis nudis glabris 2-3-floris, foliis linearibus carinatis laxè rosulatis ciliatis, calycibus pilosis, petalis calyce staminibusque duplò longioribus, siliculis ovato rotundis glabris. *℥* in Sibi-riâ bor. DC. syst. 2. p. 337.* (v. s.)
11. *D. ASPERA* (Adams ex Stevl in litt.) scapis nudis glabris, foliis linearibus carinatis rigidis ciliatis, calycibus glabris, staminibus vix calycis longitudine, siliculis oblongis, stylo brevissimo. *℥* in Sibi-riâ bor. Habitus *D. Aizoidis*, characteres ferè *D. pilosæ* aut *ericæfoliæ*. DC. syst. 2. p. 337.* (v. s.)

Sect. II. CHRYSODRABA. DC. syst. 2. p. 337.

Plantæ perennes. Folia non rigida nec carinata. Flores flavi. Stylus subnullus aut brevis. Silicula ovali-oblonga.

12. *D. ALGIDA* (Adams ex Fisch! in litt.) scapis nudis pilosis, foliis oblongis planis calycibusque pilosis, pilis simplicibus, siliculis oblongis, stylo brevissimo. *℥* in Sibi-riâ bor. Habitus *D. Alpinae* sed ab hac et ab omnibus *Chrysodrabis* distincta pilis omnibus simplicibus nec ramosis. DC. syst. 2. p. 337.* (v. s.)
- f. sabonninata*, foliis nervo medio subtus ad basin prominato subcarinatis.
γ. brachycarpa, siliculâ ovatâ breviorè.
13. *D. ALPINA* (Lin! spec. 896. excl. syn.) scapis nudis pubescentibus, foliis lanceolatis planis pilosis, pilis ramosis, siliculis oblongis, stylo brevissimo. *℥* in Lapponiâ et Norvegiâ. Wahl! fl. lapp. p. 173. n. 316. t. 11. f. 4.* Ed. fl. dan. t. 56. Flores flavi, ferè aurei, paulò minores quàm *D. Aizoidis*. DC. syst. 2. p. 338.* (v. s.)
14. *D. GLACIALIS* (Adams mem. soc. nat. mosq. 5. p. 106.*) scapo nudo stellato-pubescente, foliis lineari-lanceolatis integris pube stellatâ hispidis, siliculis ovatis glabris, stigmate subsessili. *℥* in Sibi-riâ ad littora maris glacialis. Differt à *D. algidâ* foliis angustioribus, siliculâ ovatâ, calyce constanter magisque villosa. DC. syst. 2. p. 338.* (v. s.)

15. *D. REPENS* (Bieb. fl. taur. 2. p. 93*. excl. Gmel. syn. suppl. p. 427.) scapis nudis glabriusculis, foliis lanceolatis integris pube adpressâ bipartitâ pilosis, surculis repentibus, siliculis oblongis glabris. \mathcal{L} in Iberiæ pratis. Petala obovata, obtusa, calyce glabro duplò longiora. DC. syst. 2. p. 338.* (v. s.)
16. *D. GMELINI* (Adams in mem. soc. nat. mosq. 5. p. 107.*) scapis nudis subhispidis, foliis lanceolatis hispidis pube stellatâ pilosis, surculis non repentibus, siliculis oblongis glabris. \mathcal{L} in Sibiria. — Gmel. sib. 3. p. 255. t. 56. f. 2.* Species *D. repenti* affinis et cum eâ à nonnullis confusa. — DC. syst. 2. p. 339.* (v. s.)
- β . *D. porrecta* Trin. ex Fisch. in litt. Surculis longissimis.
- γ . *D. cæspitosa* Willd. herb. ex Stev. obs. ined. Surculis brevioribus, scapisque longioribus.
17. *D. TRIDENTATA* (DC. syst. 2. p. 339.*) scapis nudis glabris, foliis obovatis longè in petiolum attenuatis apice subtridentatis hirtellis, siliculis oblongis glabris. \mathcal{L} in Caucaso Alpino. *D. hispida* Bieb. fl. taur. 2. p. 94.* (excl. syn.) suppl. p. 428. Statura *D. repenti*, sed stolones nulli. Diversa à sequenti, staturâ floribusque majoribus, siliculis longioribus. (v. s.)
18. *D. HISPIDA* (Willd. spec. 3. p. 426.) scapis nudis, foliis obovatis subdentatis hispidis, pilis bifurcatis, siliculis ovalibus. \mathcal{L} in Cappadocia. Petala calyce duplò longiora flava; stylus brevissimus. Habitus *Erophilæ vernæ*. DC. syst. 2. p. 340.*
19. *D. MURICELLA* (Wahlenb! fl. lapp. 178. n. 318. t. 11. f. 2.*) scapo nudo velutino, foliis oblongis integris pube stellatâ cæcio-velutinis, siliculis oblongo-lanceolatis glabris. \mathcal{L} in Norvegiâ, Lapponiâ et Americâ bor. *D. hirta* OEd. fl. dan. t. 142. *D. Liljebladii* Wallm. Flores albidæ, parvi. Siliquæ oblongo-lanceolatæ, utriusque subacutæ, glaberrimæ DC. syst. 2. p. 340.* (v. s.)
- β . *D. cæsia* Adams in mem. soc. nat. mosc. 5. p. 108.* Differt tantum floribus sulphureis nec albidis, siliculisque rarissimè subdentatis.
20. *D. INCOMPTA* (Stev! in mem. soc. nat. mosc. 3. p. 268.*) scapis velutinis foliis triplò longioribus, foliis oblongis pube stellatâ cano-velutinis, pedicellis flore brevioribus. \mathcal{L} in Caucaso orient. Habitus ferè *Androsaces vitalianæ*. Flores flavi. Petala calyce velutinò duplò longiora. DC. syst. 2. p. 340.* Deless! ic. sel. 2. t. 44. (v. s.)
21. *D. MOLLISSIMA* (Stev! in mem. soc. nat. mosc. 3. p. 268.*) scapis nudis glabris foliis multò longioribus, foliis oblongis pube stellatâ cano-velutinis, pedicellis flore longioribus. \mathcal{L} in Caucaso orient. Flores lutei, circiter magnitudine *D. aizoidis*. DC. syst. 2. p. 341.* Deless! ic. sel. 2. t. 45. (v. s.)
22. *D. JORULLENSIS* (H. B. et K! nov. spec. am. 5. p. 78.) caulibus foliosis villosis, foliis radicalibus elliptico-oblongis petiolatis subvelutinis, siliculis ellipticis pubescentibus. \mathcal{L} in monte ignivomò Mexicano Jorullo. Petala obovata calyce vix longiora. DC. syst. 2. p. 341.* (v. s. in h. Humb.)
23. *D. TOLUCCENSIS* (H. B. et K! nov. spec. amer. 5. p. 78.) caulibus foliosis glabriusculis, foliis radicalibus oblongis subcanescentibus, siliculis ellipticis glabris. \mathcal{L} in frigidis regni Mexicani. Aff. *D. Jorullensis*. Flores ignoti. Siliculæ ellipticæ, stylo brevi apiculatæ. DC. syst. 2. p. 341.* (v. s. in h. Humb.)

Sect. III. LRUCODRABA. DC. syst. 2. p. 342.

Plantæ perennes. Folia non rigida. Flores albi. Petala obtusa aut subemarginata.

24. *D. BOREALIS* (DC. syst. 2. p. 342.*) scapo hirsuto folium ovatum gerente, foliis cæteris oblongis basi attenuatis integris pube ramosâ velutinis, siliculis ovatis. \mathcal{L} in insulâ Sancti-Pauli. Petala alba, oblongo-cuneata, calyce duplò longiora. (v. s. comm. à cl. Fisch.)
25. *D. OBLONGATA* (Brown! in Ross. voy. Baffin bay. append.) scapis nudis pube floccosâ subhirsutis, foliis cæspitosis oblongo-linearibus integris ciliatis

velutinisque, siliculis elliptico-oblongis velutinis. \mathcal{L} in Americâ bor. Flores ignoti. Siliculæ, stylo brevissimo. DC. syst. 2. p. 342.* (v. s. in h. Banks.)

26. *D. CORYMOSA* (Brown! in Ross. voy. Baffin bay. app.) scapis nudis hispidulis, foliis dense cæspitosis oblongis basi attenuatis ciliatis subhispidisque, siliculis ellipticis corymbosis hispidulis. \mathcal{L} in Americâ bor. Flores ignoti. Siliculæ ellipticæ stylo brevissimo et stigmate subcapitato terminatæ. DC. syst. 2. p. 343.* (v. s. in h. Banks.)

27. *D. HIRTA* (Lin! spec. 897.) scapis puberulis folia 2 dentata gerentibus, foliis radicalibus subintegris puberulis, siliculis oblongis pedicellis glaberrimis. \mathcal{L} in Lapponiâ Norlandiâ et Grœnlandiâ. Wahlenb. fl. lapp. p. 175. n. 319. t. 11. f. 3.* Planta polymorpha, cum sequentibus sæpè confusa et extricatu difficillima. DC. syst. 2. p. 343.* (v. s.)

β. *alpicola* (Wahlenb. l. c. t. 11. f. 1.) habitu rigidiore humiliore, pilis magis stellatis.

γ. *Norvegica* (Gunn. fl. norv. 2. p. 106. n. 846.) Foliis radicalibus majoribus subcuneiformibus acutis dentatis congestis.

28. *D. RUPESTRIS* (Brown in h. kew. ed. 2. v. 4. p. 91.) scapis nudis 1-phyllisve puberulis, foliis lanceolatis pilosis subintegris, siliculis lanceolatis pedicellis pubescentibus. \mathcal{L} in rupibus montium Scotiæ, Norvegiæ. *D. hirta* Smith fl. brit. 2. p. 677.* engl. bot. t. 1338. non Lin. Silicula lanceolata, pilis brevissimis simplicibus furcatisve tota hirta-pubescentis, interdum, sed rarissimè glabra. DC. syst. 2. p. 344.* (v. s.)

29. *D. NIVALIS* (Willd. spec. 3. p. 427.) scapis nudis 1-phyllisve glabris, foliis oblongo linearibus ciliatis subpilosis, siliculis elliptico-oblongis pedicellis glaberrimis. \mathcal{L} in Lapponiâ, Norvegiâ, Helvetiâ etc. *D. stellata* OED. fl. dan. t. 142. Siliculæ elliptico-oblongæ, glaberrimæ. — DC. syst. 2. p. 344. (v. s.)

30. *D. LAPPONICA* (Willd. herb. ex Stev. obs. ined.) scapis nudis glaberrimis, foliis lanceolatis integerrimis pilosiusculis, siliculis ovato-lanceolatis pedicellis glaberrimis. \mathcal{L} ad rupes Alpium Lapponiæ, Grœnlandiæ. *D. androsacea* Wahlenb. fl. lapp. 174. n. 317. t. 11. f. 5.* excl. syn. *D. Wahlenbergii* Hartm. — DC. syst. 2. p. 344.* (v. s.)

31. *D. HELVETICA* (Schleich. pl. exsic.) scapis nudis glaberrimis, foliis lanceolatis ciliatis, siliculis ovalibus pedicellis glaberrimis. \mathcal{L} in Helvetiâ ad limitem nivalem. *D. ciliaris* Wahlenb. fl. helv. p. 122. n. 671.* non Lin. — DC. syst. 2. p. 345.* (v. s.)

32. *D. FLADNISENSIS* (Wulf. in Jacq. misc. 1. p. 147. t. 17. f. 1.*) scapis 2-3-phyllis glabris, foliis oblongo-linearibus rigidè ciliatis, siliculis oblongis pedicellis glaberrimis. \mathcal{L} in Croatiâ, Carnioliâ, etc. *D. androsacea* Willd. spec. 3. p. 428.* Flores pauci approximati. Sepala viridia, ovalia. — DC. syst. 2. p. 345.* (v. s. comm. à cl. Spreng.)

33. *D. TOMENTOSA* (Clairv. herb. val. p. 217? Wahlenb. fl. helv. p. 123. p. 672. t. 3.*) scapo 1-2-phyll. velutino, foliis ovali-oblongis pube stellatâ brevi tomentosis, pedicellis pubescentibus, siliculis ovalibus ciliatis. \mathcal{L} in rimis rupium sicciorum per Alpes editiores Helvetiæ. Petala alba calyce triplò longiora. Siliculæ stigmate punctiformi apiculatæ. DC. syst. 2. p. 346.* (v. s.)

34. *D. STELLATA* (Jacq. h. viind. 113. obs. n. 54. t. 4. f. 3.) scapis 1-phyllis pubescentibus, foliis ovali-oblongis pube stellatâ brevi tomentosis, pedicellis puberulis, siliquis oblongis. \mathcal{L} in fissuris rupium Pyrenæorum, Helvetiæ, Austriæ etc. *D. hirta* Jacq. fl. austr. t. 43a. Flores albi, parvi. Silicula oblonga sæpius glaberrima. — DC. syst. 2. p. 346.* Deless! ic. sel. t. 46. f. B. (v. v.)

β. *hebecarpa*, siliculis undiquè velutino-pubescentibus.

35. *D. LÆVIPS* (DC. syst. 2. p. 346.*) scapis nudis 1-phyllisve pubescentibus, foliis ovalis pube stellatâ brevi tomentosis, siliculis elongato-linearibus pedicellis glaberrimis. \mathcal{L} in rupibus Pyrenæorum. Duabus prioribus valde affinis, sed ab utraq. differt. Deless! ic. sel. 2. t. 46. f. A. (v. s.)

36. *D. SILIQUOSA* (Bieb. fl. taur. 2. p. 94.* suppl. p. 428.) scapis 1-phyllis subpuberulis, foliis lineari-lanceolatis subdentatis pube ramosâ villosiusculis, siliculis linearibus pedicellisque glabris. ② in Alpibus Caucasi orient. Petala alba oblonga vix emarginata. Siliculæ lineares. DC. syst. 2. p. 347.* (v. s.)
37. *D. LACTEA* (Adams mem. soc. nat. mosc. 5. p. 194.*) scapis nudis glabris, foliis oblongo-linearibus acutis integerrimis piloso-puberulis, siliculis elliptico-oblongis stylo brevi acuminatis pedicellisque glabris. ② in Sibiria bor. Petala lactea breviter unguiculata ovalia. DC. syst. 2. p. 347.* (v. s. in h. Mus. Par.)
38. *D. CALYCINA* (Desv. journ. bot. 3. p. 185.*) scapis nudis glabriusculis 1-floris, foliis oblongis basi attenuatis pilosis integris, siliquis linearibus glabris. ② in Peruvia. Petala alba, oblonga, calyce paulò longiora. DC. syst. 2. p. 347.* (v. s. in h. Mus. Par.)

Sect. IV. HOLARGES. DC. syst. 2. p. 348.

Plantæ annuæ aut biennes. Stylus brevis. Flores albi, rarissimè flavi.

39. *D. CONTORTA* (Ehrh. beit. 7. p. 155.) caule folioso ramoso pube stellatâ velutino, foliis ovatis dentatis, siliculis oblongis glabris subcontortis. ① ② in rupibus calcareis montanis Angliæ, Scotiæ, Pyrenæorum, Caucasi et amer. borealis, etc. *D. incana* Lin! spec. 897. Smith engl. bot. t. 388. Flores albi, petala obtusè emarginata. — DC. syst. 2. p. 348.* (v. v.)

β. *linearifolia*, caule ramosissimo, foliis linearibus vix dentatis, siliculis paulò longioribus. In Grœnlandiâ. An species propria?

40. *D. CONFUSA* (Ehrh. beit. 7. p. 155.) caule folioso subramoso pube stellatâ brevi velutino, foliis oblongis subdentatis, siliculis oblongis pubescentibus pedicello longioribus. ① ② in Nordlandiâ, Finmarkiâ, Sueciâ, Vallesiâ, etc. *D. incana* (Ed. fl. dan. t. 130. Differt à priori siliculis stylo paululùm longiore terminatis, necnon caule minùs folioso, etc. — DC. syst. 2. p. 348.* (v. s.)

β. *paucifolia*, in Kamtschatkâ et Labrador.

41. *D. CINEREA* (Adams mem. soc. nat. mosc. 5. p. 103.*) caule simpliciori folioso subpubescente, foliis oblongo-linearibus integris, siliculis oblongis pubescentibus pedicello brevioribus. ① ② in montosis sterilibus Sibirie bor. Valdè affinis *D. confusæ*, sed differt caule laxiore debiliore minùs folioso, foliis caulinis integris 5-6 sparsis, etc. DC. syst. 2. p. 349. (v. s.)

42. *D. MAGELLANICA* (Lam. dict. 2. p. 328.*) caule simpliciori folioso velutino, foliis oblongis integris, siliculis oblongis velutinis pedicello longioribus. ① ② ad fretum Magellanicum. Proximè etiam accedit ad *D. confusam*, sed differt caule simplici et foliis integris. Flores albi. DC. syst. 2. p. 349.* (v. s.)

43. *D. ARABISANS* (Michx! fl. bor. am. 2. p. 28.*) caule folioso subramoso subpuberulo, foliis lanceolatis acutis dentatis, siliculis linearibus glabris pedicello longioribus. ② in Americâ bor. *D. arabis* Pers. ench. 2. p. 190. Habitus *Arabidis*. Siliculæ erectæ acuminatæ stylo apiculatæ, contortæ, 5 lin. longæ. — DC. syst. 2. p. 349.* (v. s. in h. Mus. Par.)

44. *D. DAURICA* (DC. syst. 2. p. 350.*) caule folioso ramoso subpuberulo, foliis ovatis dentatis pubescentibus, siliculis oblongis glabris pedicelli longioribus. ② in Dauriâ. Petala alba basi attenuata subemarginata calyce duplò longiora. Stigma capiteellatum. (v. s. comm. à cl. Fisch.)

45. *D. UNALASCHIANA* (DC. syst. 2. p. 350.*) caule folioso simpliciori pubescente, foliis ovato-oblongis integris pubescentibus, siliculis oblongis hirsutulis. ②? in insulâ Unalashkâ inter Kamtschatkam et Americam bor. Petala calyce duplò longiora, limbo obovato emarginato. (v. s. comm. à cl. Fisch.)

46. *D. AUREA* (Vahl ex Horn! fl. œcon. ed. 2. p. 599. fl. dan. t. 1460.) caule folioso subramoso velutino, foliis oblongo-linearibus acutis integris pubescentibus, siliculis linearibus puberulis pedicello duplò triplòve longioribus.

(2) in Groenlandiâ. Petala flava unguiculata, limbo obovato obtusissimo. DC. syst. 2. p. 350. (v. s.)

Sect. v. DRABELLA. DC. syst. 2. p. 351.

Plantæ annuæ aut biennes. Stylus nullus. Flores minimi, flavi aut albi.

47. *D. LUTEA* (Gilib. fl. lith. in Ust. del. op. 2. p. 357.*) caule ramoso folioso pubescente, foliis ovalibus denticulatis pubescentibus, siliculis elliptico-oblongis glabris 20-40-spermis. (1) Flores racemosi, parvi, flavi. Petala calyce longiora, obtusa ferè elliptica. DC. syst. 2. p. 351.* (v. s.)

a. brevipes, pedicellis siliculâ vix duplò longioribus. in Armeniâ seu Capadociâ. D. Pontica Desf. cor. tourn. p. 67. t. 51. ann. mus. 11. p. 381. t. 38.*

β. longipes, pedicellis siliculâ triplò quadruplòve longioribus. in Caucaso sept. in montibus Uralensisibus et in Dahuriâ. D. lutea Gilib. l. c. D. muralis Steph. fl. mosc. p. 29. Bieb. fl. taur. n. 1246. suppl. p. 428.

48. *D. MEMORALIS* (Lini spec. ed. 1. p. 643. Montt. pl. syst. 4. t. 60. f. 1.) caule ramoso folioso pubescente, foliis ovatis dentatis pubescentibus, siliculis elliptico-oblongis 32-36-spermis pube minimâ velutinis. (1) in Japoniâ, Tauriâ, Pyrenœis et Americâ bor. etc. A *D. luteâ* tantum differt pube silicularum. DC. syst. 2. p. 351.* (v. s.)

49. *D. MURALIS* (Lini spec. ed. 1. p. 643.) caule ramoso folioso pubescente, foliis ovatis dentatis subcordato amplexicaulibus subhirsutis, siliculis elliptico-oblongis glabris 12-16 spermis. (1) in umbrosis montium Europæ à Lusitaniâ ad Sueciam, etc. Smith engl. bot. t. 912. Lam. ill. t. 556. f. 2. Flores parvi, albi. Petala obovata, integra. DC. syst. 2. p. 353.* (v. v.)

50. *D. CAROLINIANA* (Walt. fl. carol. 174.) caule basi folioso hispido, apice glabro nudo, foliis ovato-rotundis integris hispidis, siliculis linearibus glabris pedicello longioribus. (1) in Americâ bor. *D. hispidula* Michx. fl. bor. amer. 2. p. 28. Flores parvi, albi. DC. syst. 2. p. 353.* (v. s.)

†. *Species quoad characteres genericos dubiæ aut non satis notæ.*

51. *D. PUMILIO* (Brown! in DC. syst. 2. p. 353.*) scapis nudis 1-floris, foliis radicalibus ovatis integris petiolatis, siliculis ovatis. (1) in terrâ van Diemena. Flores albi minimi. Calyx patens. (v. s.)

52. *D. VESICARIA* (Desv. journ. bot. 3. p. 186.*) scapis nudis stellato-hirsutis, foliis confertis oblongis pube stellatâ velutinis pilis simplicibus ciliatis, siliculis ovatis inflatis hirsutis. 2/ in rupibus montium Palestinæ, Syriæ. Flores flavi. DC. syst. 2. p. 354.* (v. s.)

53. *D. PULCHELLA* (Willd. herb. ex Stev. obs. ined. et DC. syst. 2. p. 354.) scapo nudo, foliis lauceolatis integerrimis utrinquè stellato-pubescentibus, siliculis ovatis glabris. — In Persiæ montibus.

54. *D. ARETIOIDES* (H. B. et K. nov. spec. amer. 5. p. 77. t. 435.) scapis nudis, foliis oblongis obtusis villosis imbricatis, siliculis ovatis villosis. 2/ in devexis montis Antisanæ alt. 1800 hexapod. in Andibus Quitensium. Aretia cana Willd. herb. ex Kunth. Androsace cana. Rœm. et Schult. syst. 5. p. 786. Flores minimi flavi. — DC. syst. 2. p. 354.*

55. *D. VIOLACEA* (DC. syst. 2. p. 354.*) caule suffruticoso ramoso diffuso, foliis oppositis ovatis tomentosis, siliculis oblongis. 2/ in frigidissimis nivosis montis Assuay regni Quitensis altit. 1980 hexapod. D. Bonplaudiana H. B. et K. nov. spec. amer. 5. p. 78. Petala violacea, ovata.

56. *D. ALYSSOIDES* (H. B. et K. nov. spec. amer. 5. p. 79.) caule subramoso foliisque ovato-oblongis subdentatis hirsutis, siliculis ovato-lanceolatis hispidis. 2/ in provinciâ Pastoensi. Flores albi. DC. syst. 2. p. 355.

57. *D. RAMOSISSIMA* (Desv. journ. bot. 3. p. 186.) caule ramoso humifuso foliisque glabris spathulatis remotè dentatis. 2/ in Amer. bor. Siliculæ glabræ

compressæ, immarginatæ, sæpè obliquæ, stylum longum gerentes. DC. syst. 2. p. 355.

58. D. GLABELLA (Pursh. fl. bor. amer. 2. p. 434.*) scapo subbifolio erecto, foliis spathulato-lanceolatis glabriusculis, racemo densifloro, siliculis ovato-lanceolatis acutis. (1) ad Hudson's bay in Americâ boreali.

XXX. EROPHILA. DC. syst. 2. p. 356.

Silicula ovalis aut oblonga, valvis planis. Semina plurima immarginata. Calyx æqualis. Petala bipartita. Stamina edentula. — Herbæ annuæ, scapis nudis, floribus albis.

1. E. AMERICANA (DC. syst. 2. p. 356.) siliculis oblongis pedicello brevioribus, scapis 5-10-floris (1) in Americæ borealis campis. D. verna Americana Pers. ench. 2. p. 190. Differt ab Erophilâ vulgari siliculis oblongis neo ellipticis, latitudine triplò ferè longioribus. Forsan varietas? (v. s.)
2. E. VULGARIS (DC. syst. 2. p. 356.*) siliculis ellipticis pedicello brevioribus, scapis 5-15-floris. (1) in muris pascuisque aridis Europæ totius. Draba verna Lin. spec. 896. OEd. fl. dan. t. 983. (v. v.)
3. E. PRÆCOX (DC. syst. 2. p. 357.) siliculis suborbiculatis vix pedicello brevioribus, scapis 3-5-floris. (1) in Caucaso orientali. Draba præcox Stev. mem. soc. nat. mosc. 3. p. 269. (v. s.)
4. E. MINUTISSIMA (DC. syst. 2. p. 357.) siliculis....., scapo nudo subtrifloro, foliis ellipticis petiolatis. (1) propè Constantinopolim. Draba minutissima Willd. herb. ex Stev. obs. ined.
5. E. MUSCOSA (DC. syst. 2. p. 358.*) siliculis ellipticis pedicello ferè longioribus, scapo 2-4-floro. (1) in Peruvîâ. Flores 2-4, parvi, albi. Petala semibifida, calyce longiora. (v. s. in h. Lamb.)

XXXI. COCHLEARIA. Tourn. inst. 215. t. 101. Lin. gen. n. 803. Lam. ill. t. 558.

Silicula sessilis, ovato-globosa aut oblonga, valvis ventricosis. Semina plurima immarginata. Calyx æqualis, patens. Petala integra. Stamina edentula. — Flores albi. Folia sæpè subcarnosa. DC. syst. 2. p. 358.

Sect. I. KERNERA. DC. syst. 2. p. 359.

Silicula globosa, basi subtetragona, valvis rigidiusculis. Filam. 4 medio geniculata. Stylus brevis, stigmata bilobo.

1. C. SAXATILIS (Lam! fl. fr. 2. p. 471.) siliculis lentiformibus lævibus, foliis radicalibus oblongis dentatis pilosis, caulinis lineari-oblongis. 2/ in Galliâ, Helvetiâ, Italiâ, etc. Myagrum saxatile Lin! spec. 894. Jacq. fl. austr. t. 128. Petala obovata calyce duplò triplòve longiora. — DC. syst. 2. p. 359.* (v. v.)
ß. incisa, foliis lyratis pinnatifidisve. — Moris. oxon. 2. p. 293. n. 11. s. 3. t. 17. fig. ult.
2. C. AURICULATA (Lam. diet. 2. p. 165.*) siliculis lentiformibus lævibus, foliis radicalibus oblongis dentatis pilosis, caulinis cordato-sagittatis. 2/ in rupestribus calcareis montium Arvernæ, Sabaudiz, Pyreneorum, etc. Myagrum montanum Berg. phyton. 3. p. 140. icon. — DC. syst. 2. p. 360. (v. v.)
3. C? ALYSSOIDES, siliculis globosis cum calyce inflatis, foliis oblongis obtusis sinuato-dentatis crassiusculis, radicalibus petiolatis, supremis semi-amplexicaulibus. — in Indiâ orientali. Alyssum cochlearioides. Roth nov. pl. spec. 322.* An Vesicariæ spec?

Sect. II. ARMORACIA. DC. syst. 2. p. 360.

Silicula ellipsoidea aut oblonga, stylo brevi filiformi, stigmate capitato ferè disciformi.

4. *C. ARMORACIA* (Lin! spec. 904.) siliculis ellipsoideis, foliis radicalibus oblongis crenatis, caulinis elongato-lanceolatis dentatis incisive, radice carnosâ maximâ. 2/ in aquosis submontanis Europæ. Smith engl. bot. t. 2223. Flores calyce patente. DC. syst. 2. p. 360.* (v. v.)
5. *C. MACROCARPA* (Waldst. et Kit. pl. rar. hung. 2. p. 101. t. 184.*) siliculis ellipsoideis, foliis radicalibus oblongis crenatis, caulinis lanceolatis dentatis, dentibus cartilagineis, radice carnosâ, sepalis erectis. 2/ in Hungariâ et Transylvaniâ. *C. crassifolia* Willd. herb. ab hac non differre videtur ex Stev. obs. DC. syst. 2. p. 362.* (v. s.)
6. *C. MICROCARPA* (DC. syst. 2. p. 362*) siliculis globosis, foliis oblongis basi auriculato-amplexicaulibus dentatis. — in Sibirîâ. *C. Altaica* Schlecht. in h. Willd. ex Stev! obs. ined. Flores ignoti. Siliculæ parvæ. (v. s. in h. Stev.)

Sect. III. COCHLEAR. DC. syst. 2. p. 362.

Silicula formâ varia, apice non emarginata, stylo brevissimo aut nullo.

7. *C. GLASTIFOLIA* (Lin! spec. 904.) siliculis subglobosis, foliis caulinis cordato-sagittatis amplexicaulibus acuminatis integerrimis. ① ② circa Ratisbonam, in Lusitaniâ, Delphinatû, etc. Schkuhr handb. 2. n. 1504. t. 181. DC. syst. 2. p. 362.* (v. v.)
8. *C. OBLONGIFOLIA* (DC. syst. 2. p. 363.*) siliculis subrotundis pedicello dimidiò brevioribus, foliis caulinis oblongis integris sessilibus, summis brevissimè auriculatis. 2/? Petala calyce duplò longiora.
- α. *Draba cochlearioides* Langsdorf. ex Fisch. in litt. — in insulâ Sancti-Pauli.
- β. *C. anglica* è Kamtschatka Pall. in h. Willd. *C. Kamtschatika* Schlecht. in h. Willd. ex Stev! obs.
9. *C. ANGLICA* (Lin. spec. 903.) siliculis ovato-subglobosis venoso-reticulatis pedicello dimidiò brevioribus, foliis radicalibus petiolatis ovatis integris, caulinis oblongis. ① in littoribus maritimis Armoraciæ, Angliæ, Norvegiæ, etc. Oed. fl. dan. t. 329. Smith engl. bot. t. 552. Adest in herbario Linneano sed sine nomine. DC. syst. 2. p. 364.* (v. s.)
10. *C. OFFICINALIS* (Lin! spec. 903.) siliculis ovato-globosis pedicello dimidiò brevioribus, foliis radicalibus petiolatis cordatis, caulinis ovatis dentato-angulatis. ② in littoribus maritimis Europæ septentr. Smith engl. bot. t. 551. DC. syst. 2. p. 365* (v. v.)
- β. *minor*, Smith fl. brit. 2. p. 688. *C. Danica* Gunn. norv. n. 197. ex Vahl.
- γ. *rotundifolia*, Smith l. c. *C. Groenlandica* With. brit. 573. non Lin.
11. *C. PYRENAICA* (DC. syst. 2. p. 365.*) siliculis obovatis subglobosis pedicelli longitudine, foliis radicalibus cordato-reniformibus integris, caulinis paucis ovatis dentatis. ② in summis Pyrenæorum centralium. Deless! ic. sel. 2. t. 48. *C. officinalis* Lapey. abr. pyr. p. 368. (v. v.)
12. *C. GROENLANDICA* (Lin. spec. 904.) siliculis ovatis pedicelli longitudine, foliis radicalibus reniformibus integris, caulinis subnullis. ② in Groenlandiâ, Norvegiâ et Islandiâ. Lodd. bot. cab. 1. t. 45. Herba parva, vix digiti longitudine. DC. syst. 2. p. 366.* (v. s. in h. Vahl.)
13. *C. DANICA* (Lin! spec. 903.) siliculis ellipsoideis pedicelli longitudine, foliis omnibus petiolatis subdeltoideis. ① in maritimis Europæ borealis. Smith engl. bot. t. 696. Ed. fl. dan. t. 100. DC. syst. 2. p. 366. (v. v.)
- β. *integrifolia*, foliis caulinis paucissimis, radicalibus reniformibus, omnibus integris. — in Hiberniâ.
14. *C. LENENSIS* (Adams ex Fisch! in litt.) siliculis ovatis pedicello triplò brevioribus, foliis radicalibus petiolatis subreniformibus obtusis, caulinis oblongis subintegris. — in Sibirîâ boreal. *C. Groenlandica* Willd. herb. ex Stev. obs. ined. Flores parvi DC. syst. 2. p. 367.* (v. s.)

15. *C. ARCTICA* (Schlecht. in h. Willd. ex Stev! obs.) siliculis ovato-oblongis pedicello paulò brevioribus, foliis radicalibus petiolatis ovatis subdentatis, caulinis sessilibus utrinquè 1-dentatis. in Sibiria bor? Siliculæ ovatæ sub oblongæ vix pedicello breviores. DC. syst. 2. p. 367. (v. s.)
16. *C. TRIDACTYLITES* (Banks herb. et DC, syst. 2. p. 367.) siliculis ovato-globosis, foliis caulinis dente unico utrinquè grossè subtrilobatis. — in Labrador. aff. *C. Lenensi* et Arcticæ. Siliculæ magnitudine ferè *C. officinalis*. (v. s.)
17. *C. FENESTRATA* (Brown in app. voy. Ross. exlur.) siliculis ellipsoideis pedicello dimidiò brevioribus, septo elongato fenestrato, foliis radicalibus petiolatis ovatis, caulinis oblongis. — in Americæ bor. regionibus polaribus. Flores parvi albi. DC. syst. 2. p. 367.* (v. s. in h. Banks.)
18. *C. SISYMBRIOIDES* (DC. syst. 2. p. 368.*) siliculis oblongis pedicello duplò brevioribus, petalis oblongis, foliis superioribus ovato-lanceolatis amplexicanibus dentatis. — in Sibiria. Flores albi, majusculi, sepalis oblongis patulis margine membranaceis. (v. s.)
19. *C. GRANDIFLORA* (DC. syst. 2. p. 368.*) siliculis oblongis, petalorum limbis amplis obovatis, foliis caulinis ovato-lanceolatis obtusiusculis, grossè dentatis basi obtusè subauriculatis. — in Sibiria. Priori valdè affinis, sed species distinctissima. (v. s.)
20. *C. INTEGRIFOLIA* (DC. syst. 2. p. 369.*) siliculis oblongis, petalis obovato-oblongis, foliis caulinis integris, inferioribus petiolatis ovatis, superioribus subsessilibus lanceolatis. — in jugo Altaico Sibiria. Duabus prioribus affinis. Flores albi minores quàm *C. sisymbrioidis*. (v. s.)
21. *C. SALSA* (Schlecht. in h. Willd. ex Stev! obs. ined.) siliculis lanceolatis stylo conico superatis pedicello dimidiò brevioribus, foliis caulinis oblongo-lanceolatis cordato-amplexicanibus cauleque pubescentibus. — in Sibiria? salsuginosis. Flores albi magnitudine *C. Anglicæ*. DC. syst. 2. p. 369.* (v. s. in h. Stev.)
22. *C. SPATHULATA* (Schlecht. in h. Willd. ex Stev! obs. ined. et DC. syst. 2. p. 369.*) siliculis lanceolatis stylo brevi apiculatis pedicello paulò brevioribus, foliis spathulatis pendentibus pubescentibus. — in insulis Aleutianis inter Kamtschatkam et Americam. Accedit ad *C. Arcticam*, sed pube trifurcâ faciliè dignoscitur. (v. s. in h. Stev.)
23. *C. SEPTENTRIONATIS* (Schlecht. in Willd. herb. ex Stev. obs. ined.) canle ramoso folioso puberulo, foliis puberulis inferioribus ovalibus longè cuneatis grossè dentatis, siliculis ovatis glabris, septo fenestrato. ①? in Sibiria bor. in insulâ Sancti-Pauli inter Kamtschatkam et Americam. *Draba grandis* Langsdorff ex Fisch! in litt. Deless! ic. sel. 2. t. 47. DC. syst. 2. p. 355.* (v. s.)
24. *C? SILIQUOSA* (Schlecht. in h. Willd. ex Stev! obs. ined. et DC. syst. 2. p. 369.*) siliculis oblongo-lanceolatis stylo apiculatis pedicello dimidiò brevioribus, foliis oblongis integris basi attenuatis pubescentibus. 2/ in elatis saxosis Unalaschka. Flores in sicco flavescentes. (v. s.)
25. *C? VELUTINA* (DC. syst. 2. p. 370.*) siliculis ellipsoideis pedicello dimidiò brevioribus, foliis pube ramosâ velutinis, radicalibus pinnatipartitis, caulinis sagittatis. ① in Mesopotamia. Flores ferè *C. saxatilis*. Petala latè obovata. Deless! ic. sel. 2. t. 49. (v. s.)
26. *C? SAXIFRAGAEFOLIA* (DC. syst. 2. p. 370.*) siliculis ovato-oblongis, foliis glabris, radicalibus petiolatis reniformibus palmatim inciso-dentatis, caulinis ovatis trilobis, summis linearibus. 2/ in Persia. Flores omninò prioris. Deless! ic. sel. 2. t. 50. (v. s.)

Sect. IV. IONOPSIDIUM. DC. syst. 2. p. 371.

Silicula subrotundo-compressa, apice emarginata. Flores lilacini. — Sectio omninò media inter Cochleariam et Thlaspidem.

27. *C. ACAULIS* (Desf! fl. atl. 2. p. 69.*) siliculis subrotundis emarginatis, pedicellis petiolisque radicalibus longis, foliis ovato-rotundatis integris. 2/ in

Lusitaniâ frequens. C. pusilla Brot. phyt. p. 100. n. 45. t. 21. f. 2. 3*. Jacq. eclog. t. 132. opt. — DC. syst. 2. p. 371.* (v. s.)

†. *Species non satis notæ.*

28. C. SAGITTÆFOLIA (Desv. journ. bot. 3. p. 182.) ① hab. Flores albi parvi. Silicula oblonga integra, tumida, oligosperma. DC. syst. 2. p. 371.
29. C. LYRATA (Sibth. et Smith fl. græc. t. 619. ex Smith prodr. 2. p. 8.) ① in Siciliâ. Folia caulina oblonga sagittata amplexicaulia. DC. syst. 2. p. 372.
30. C. REPANDA (Med. in Ust. neu. ann. 2. p. 42.) — Hab. folia superiora amplexicaulia sagittata repanda, in cochlearis formâ cavata. DC. syst. 2. p. 372.
31. C. LONGIFOLIA (Med. in Ust. neu. ann. 2. p. 41.) — hab. Folia inferiora longè petiolata, superiora sessilia plerumquè integra. DC. syst. 2. p. 372.
32. C. HETEROPHYLLA (Schlecht. in h. Willd. ex Stev. obs. ined.) 2? hab. Flores albi, magnitudine Iberidis ciliatæ. Silicula elliptica obtusa, glabra. Stigmatè crasso sessili. DC. syst. 2. p. 372.

Trib. III. *THLASPIDÆ* seu *PLEURORHIZÆ* ANGUSTISEPTÆ.

Silicula dehiscens, septo angustissimo, valvis carinatis navicularibus. Semina ovalia, interdum marginata. Cotyledones planæ, accumbentes, septo contrariæ. — Tribus distinctissima, cum Lepidineis solis conferenda, sed distincta seminibus magis compressis et cotyledonibus accumbentibus.

*. *Siliculæ loculis 2-∞-spermis.*

XXXII. *THLASPI*. Dill. fl. guss. gen. nov. p. 123. t. 6. Vent. tabl. 3. p. 110.

Silicula apice emarginata, valvis navicularibus dorso alatis, loculis 2-∞-spermis. Petala æqualia. Calyx basi æqualis. — Flores albi. DC. syst. 2. p. 373.

Sect. I. *PACHYPHRAGMA*. DC. syst. 2. p. 373.

Silicula lata brevissimè emarginata. Stylus o. Septum duplex! plicis 3 longitudinaliter instructum. Semina 4 non striata.

1. T. LATIFOLIUM (Bieb. fl. taur. 2. p. 99.* suppl. p. 430.) foliis radicalibus longè petiolatis cordatis repando-dentatis, caulinis ovato cordatis breviter petiolatis. 2? in Iberiâ et ad Caucasum sept. Lepia latifolia Desv. journ. bot. 3. p. 166. Flores majusculi. Petala cuneata obtusa calyce triplò longiora. DC. syst. 2. p. 374. Deless! ic. sel. 2. t. 51. (v. s.)

Sect. II. *CARPOCERAS*. DC. syst. 2. p. 374.

Valvæ apice expansæ in alam cornu referentem; semina 4 striata. Septum membranaceum oblongum.

2. T. CERATOCARPON (Murr. comm. gœtt. 1774. p. 26. t. 1.*) foliis radicalibus subpetiolatis obovato-oblongis, caulinis oblongis basi auriculis acutis hastato-amplexicaulibus. ① in campis salsuginosis Sibirîæ. Capsella cornigera Medik. in Ust. neu. ann. 2. p. 46. Flores albi minimi. Pedicelli fructiferi filiformes. (v. v.)

Sect. III. *NOMISMA*. DC. syst. 2. p. 375.

Valvæ per totum dorsum alata; semina ∞ striata.

3. T. ARVENSE (Lin! spec. 901.) foliis oblongis dentatis, caulibus erectis, siliculis obqvato-orbiculatis pedicello brevioribus. ② in arvis Europæ. Smith engl. bot. t. 1659. Flores parvi albi. DC. syst. 2. p. 375.* (v. v. et s.)
4. T. BAICALENSE (DC. syst. 2. p. 376.*) foliis oblongis grossè dentatis, caulibus erectis, siliculis orbiculatis pedicello longioribus. ① in arenosis Sibirîæ. Forsan prioris mera varietas. (v. s.)

5. *T. COLLINUM* (Bieb. fl. tanr. 2. p. 99.) foliis oblongis dentatis, caulinis lineari-sagittatis strictis, caulibus diffusis subadscendentibus, siliculis sub-orbiculatis. (1) in Armeniâ, Iberiâ. *T. nemorosum* Adam. Hoffm. cat. h. mosq. 1808. n. 3249. ex Stev. obs. in h. Willd. Affinitas cum *T. arvensi* summa, tamen distinctum videtur. DC. syst. 2. p. 376.* (v. v. comm. à cl. Stev.)

Sect. IV. NEUROTROPIS. DC. syst. 2. p. 377.

Siliculæ orbiculatæ, sinu angusto emarginatæ. Valvæ dorso expansæ in alam latam nervo circumscriptam. Semina oo non striata.

6. *T. ORBICULATUM* (Stev! in litt. et DC. syst. 2. p. 377.*) foliis superioribus amplexicaulibus integerrimis, siliculis orbiculatis, alâ latâ nervo circumscriptâ. (1) in Iberiâ. Semina oblonga, compressa, rufa. (v. siliq. aicc.)
7. *T. UMBELLATUM* (Stev! in DC. syst. 2. p. 377.*) foliis ovatis dentatis, radicalibus subpetiolatis, caulinis semi-amplexicaulibus, siliculis obcordatis alâ latâ nervo circumscriptâ. (1) in Persiâ. (v. s. in h. Stev.)

Sect. V. PTEROTROPIS. DC. syst. 2. p. 377.

Silicula subobovata, emarginata aut truncata. Valvæ per totum dorsum alatae, alâ nervo non emarginatâ; semina non striata.

8. *T. ALLIACEUM* (Lin. spec. 901.) foliis oblongis obtusis subdentatis, inferioribus petiolatis, superioribus auriculis acutis sagittato-amplexicaulibus, siliculis subovatis ventricosis, stigmatibus sessilibus. (1) in agris et arvis Europæ. Jacq. ic. pl. rar. 1. t. 121. Flores paulò minores quàm *T. arvensis*; siliculæ dimidio ferè minores. DC. syst. 2. p. 377.* (v. s.)
9. *T. PERFOLIATUM* (Lin! spec. 902.) foliis subdentatis, radicalibus petiolatis, caulinis cordato-amplexicaulibus, caule ramoso, petalis calyci æquantibus, siliculis obcordatis 8-spermis, stigmatibus sessilibus. (1) (2) in agris et arvis apricis præsertim calcareis Europæ. Smith engl. bot. t. 2354. DC. syst. 2. p. 378.* (v. v.)

℞. *simplicissimum*, caules simplici stricto à basi striato. — in Persiâ bor. (v. s.)

10. *T. MONTANUM* (Lin! spec. 902.) foliis subcarnosulis integris, radicalibus obovatis petiolatis, caulinis oblongis sagittato-amplexicaulibus, petalis calyce majoribus, siliculis obcordatis 4-spermis, stylo filiformi. 2℞ in montosis Europæ, ab Hispaniâ ad Podoliam, à Sicilia ad Belgium. Jacq. fl. austr. 2. p. 237. DC. syst. 2. p. 379.* (v. v.)

℞. *præcox*, Wulf. in Jacq. coll. 2. p. 124. t. 9.* differt tantum herbâ paulò magis glaucescente et foliis radicalibus apice crenatis.

γ. *Alpinum*, Jacq. fl. austr. 3. t. 238, stylo longiore ovarium superante et siliculæ longitudinem ferè æquante.

11. *T. ALPESTRE* (Lin! spec. 903.) foliis integris, radicalibus ovatis petiolatis, caulinis oblongis amplexicaulibus, petalis calyci subæqualibus, siliculis obcordatis 8-12 spermis, stylo filiformi. 2℞ in agris et pratis montanis Pyrenæorum, Alpium, in regni Valentini montibus excelsioribus, etc. Smith engl. bot. t. 81. DC. syst. 2. p. 380.* *T. cærulescens* Presl. fl. cech. p. 1. (v. v.)
12. *T. HETEROPHYLLUM* (DC. fl. fr. ed. 3. v. 4. p. 712.*) foliis radicalibus petiolatis patulis integris dentatis et lyratis, caulinis erectis cordato-sagittatis, petalis calyci subæqualibus. 2℞ in Pyrenæis. Flores *T. alpestris* paulò minores. DC. syst. 2. p. 381.*
13. *T. MAGELLANICUM* (Pers. ench. 2. p. 189.) foliis radicalibus ovalibus petiolatis subintegris, caulinis oblongis sessilibus, petalis calyci subæqualibus, siliculis oblongis truncato-emarginatis 8-spermis, stylo brevi. — ad fretum Magellanicum. Flores albi parvi. Semina ovali-oblonga, rufa, compressa, non striata. DC. syst. 2. p. 381.* (v. s. in h. Mus. Par.)
14. *T. COCHLEARIFORME* (DC. syst. 2. p. 381.*) foliis subcarnosulis, radica-

libus petiolatis ovatis subdentatis, caulinis cordato-amplexicaulibus, petalis calyce majoribus, siliculis oblongis submarginatis 8-spermis, stylo brevi filiformi. \mathcal{Z} in Sibiriâ et Dahuriâ. Deless. ic. sel. 2. t. 52. Flores omninò T. montani. (v. s.)

†. *Species non satis notæ.*

15. T? SAMOLIFOLIUM (DC. syst. 2. p. 382.*) foliis ovatis obtusis integerrimis, inferioribus breviter petiolatis, petalis emarginatis. — In Cappadociâ. Alysum samolifolium. Desf. choix. cor. p. 66. t. 49.* Flores albi, magnitudine Berterose incanæ. (v. s. in h. Mus. Par.)
16. T? TUBEROSUM (Nutt. gen. am. 2. p. 65.) foliis rhombeo-ovatis obsolete dentatis sessilibus, radicalibus longè petiolatis, caule pubescente, radice tuberosâ. — In Pennsylvania occid. Flores majusculi, ferè Arabidis, rosei. Silicula orbiculata brevis. DC. syst. 2. p. 382.
17. T? SCHRANKII (Schult. fl. aust. p. 83.) foliis subhispidis integris, radicalibus petiolatis, caulinis amplexicaulibus, calyce petalis minore, siliculis obcordatis. — In mœnibus Austriæ cum Capsellâ bursâ-pastoris. DC. syst. 2. p. 382.

XXXIII. CAPSELLA. Vent. tabl. 3. p. 110. DC. syst. 2. p. 383.

Silicula triangularis basi cuneata, valvis navicularibus apteris, loculis co-spermis.

1. C. BURSA-PASTORIS (Mœnch. meth. 271 ①) in Europâ, Indiâ, Japoniâ, Dahuriâ, Persiâ, Promontorio Bonæ-Spei, Americâ bor. etc. Thlaspi Bursa-pastoris Lin! spec. 903. Smith engl. bot. t. 1485. Ed. fl. dan. t. 729. DC. syst. 2. p. 383.* (v. v.)

β . minor. Bursa-pastoris minor Tabern. ic. 199.

γ . integrifolia. — Moris oxon. 2. p. 104. s. 3. t. 20. f. 1.

δ . coronopifolia. — Bauh. pin. 108.

ϵ . apetala (Opiz. in bot. zeit. 1821. p. 440.), floribus apetalis decandris.

Potius monstrum quàm varietas. (v. s. comm. à cl. Jacquin.)

XXXIV. HUTCHINSIA. Brown in h. Kew. ed. 1812. v. 4. p. 82. non Agardh.

Silicula elliptica, valvis navicularibus apteris, loculis 2-spermis, rarius co-spermis. Calyx æqualis. Petala æqualia.

Sect. 1. IBERIDELLA. DC. syst. 2. p. 385.

Stylus filiformis. Folia integra aut subdentata.

1. H. ROTUNDIFOLIA (Brown in h. kew. ed. 2. v. 4. p. 82.) foliis subcarnosis integerrimis, infimis petiolatis obovatis, caulinis ovato-oblongis subamplexicaulibus, staminibus petalis et stylo siliculâ dimidiò brevioribus. \mathcal{Z} propè nives inter lapides Alpium. Iberis rotundifolia. Lin! spec. 905. Scop. fl. carn. n. 805. t. 37. — DC. syst. 2. p. 385.* (v. v.)
2. H. CEPEÆFOLIA (DC. syst. 2. p. 386.*) foliis subcarnosis, inferioribus petiolatis oblongo-obovatis apice denticulatis, caulinis oblongis, staminibus corollâ brevioribus, stylo siliculâ triplò brevior. \mathcal{Z} in Carinthiâ, in Apennino. Iberis cepeæfolia Wulf. in Jacq. misc. 2. p. 28. t. 1.* Priori valde affinis et habitu et characteribus. (v. s. comm. à cl. Spreng.)
3. H. PUMILA (DC. syst. 2. p. 386.*) foliis subcarnosis, inferioribus longè petiolatis ovalibus integris, caulinis oblongis basi acutè sagittatis, staminibus corollâ et stylo siliculâ triplò brevioribus. \mathcal{Z} in Caucaso orient. Iberis pumila Stev! mem. soc. nat. mosc. 3. p. 269. Non confundi potest cum H. brevistylâ ob siliculas utrinquè attenuatas. (v. s.)
4. H. STYLOSA (DC. syst. 2. p. 387.*) foliis subcarnosis, inferioribus petiolatis obovato-oblongis subintegris, caulinis oblongis, staminibus petala et stylo siliculam subæquantibus. ② in editioribus montibus regni Neapolitani. Iberis stylosa Tenor! prodr. fl. nap. xxxvii. Thlaspi minimum Arduin. specim. 2. p. 37. t. 15. f. 1? Flores albi corymbosi. (v. s.)

5. *H. BREVISSYLA* (DC. syst. 2. p. 387.*) foliis subcarnosis petiolatis obovatis subdentatis, siliculâ oblongâ truncato-emarginatâ, stylo brevissimo. $\frac{1}{2}$ in Syriæ montibus. Flores parvi, albi. Petala oblongo-cuneata, obtusa, fere truncata. (v. s.)
6. *H. TRINERVIA* (DC. syst. 2. p. 387.*) foliis integerrimis trinerviis ovali-oblongis sessilibus basi subcordato-amplexicaulibus. $\frac{1}{2}$ in Persiâ. Flores ex albo subpurpurascens. Valvæ carinatæ. Ovula in quoque loculo duo. Deless! ic. sel. 2. t. 53. (v. s.)
7. *H. HASTULATA* (DC. syst. 2. p. 388.*) foliis basi cordato-hastulatis ovato-oblongis subdenticulatis semiamplexicaulibus, siliculis elongatis apice truncatis. $\frac{1}{2}$ in Hyrcaniâ et in Persiâ bor. *Thlaspi hastulatum* Stev! in litt. — Gmel. sib. 3. p. 254. t. 56. f. 1. Flores albi cum rubore suffuso, magnitudine et habitu *H. rotundifoliæ*. (v. s.)

Sect. II. NASTURTIOLUM. DC. syst. 2. p. 388.

Folia pinnato-lobata.

8. *H. CALYCINA* (Desv. journ. bot. 3. p. 168.) foliis pinnatipartitis cauleque erecto pubescentibus, calycibus persistentibus, siliculis oblongis utrinquè attenuatis stylo acuminatis. $\frac{1}{2}$ in Alpihus Altaicis Sibiriz. *Lepidium calycinum* Steph. in Willd. spec. 3. p. 433.* Petala alba, oblonga, calyce duplò longiora, rarò decidua. (v. s. comm. à cl. Fisch.)
9. *H. ALPINA* (Brown in h. kew. ed. 2. v. 4. p. 82.) foliis pinnatipartitis glabris, petalis calyce deciduo duplò longioribus, siliculis utrinquè acutis, stylo brevissimo exserto. $\frac{1}{2}$ in rupibus humidiusculis Pyrenæorum, Alpium Apennini et in montibus Carpathicis. *Lepidium Alpinum* Lin! amœn. 4. p. 321.* Jacq. fl. austr. t. 137. — DC. syst. 2. p. 389.* (v. v.)
10. *H. PETRÆA* (Brown in h. kew. ed. 2. v. 4. p. 82.) foliis pinnati-partitis glabris, petalis calyce vix longioribus, siliculis utrinquè obtusis 4-spermis, stigmate sessili. ① ② in petrosis apricis Europæ. *Lepidium petræum*. Lin! spec. 899. Smith! engl. bot. t. 111. (v. v.)
11. *H. PROCUMBENS* (Desv. journ. 3. p. 168.) foliis sinuato-pinnatifidis aut subintegris, petalis vix calyce majoribus, siliculis utrinquè obtusis 10-12-spermis, stigmate sessili. ① in ruderatis maritimis Hispaniæ, Galliæ australis, Pedemontii, Cypri, Tauriæ. *Lepidium procumbens* Lin! spec. 898. Schkuhr handb. 2. n. 1779. — DC. syst. 2. p. 390.* — (v. v.)

β. *erectiuscula*, caulibus erectiusculis. — In Syriâ.

γ. *integrifolia*, foliis integris aut parçè incisis. — Massiliæ.

XXXV. *TEESDALIA*. Brown in h. kew. ed. 2. vol. 4. p. 83. DC. syst. 2. p. 391. — *Guepinia*. Bast. suppl. 35.

Silicula ovalis apice emarginata, valvis navicularibus, loculis 2-spermis. Stamina intus basi squamulâ aucta. — Herbæ annuæ, foliis radicalibus pinnatilobatis, floribus albis parvis.

1. *T. IBERIS* (DC. syst. 2. p. 392.) petalis inæqualibus, exterioribus majoribus. ① in arenosis sylvaticis Europæ. *Iberis nudicaulis* Lin! spec. 903. fl. dan. t. 323. Smith engl. bot. t. 327. (v. v.)

2. *T. LEPIDIUM* (DC. syst. 2. p. 392.) petalis æqualibus. ① in arenosis sterilibus Europæ australis et in Mauritaniâ. *Lepidium nudicaule* Lin. spec. 898. (v. v.)

β. *integrifolia*, folia integra nec pinnatim secta. — in Hispaniâ, Sardiniâ.

γ. *acutiloba*, folia magis oblonga, lobis dentibusve lateralibus 3-4 acutis, terminali elongato acuminato. In insulâ Scio. An forsâ species propria?

**. *Siliculæ loculis monospermis.*

XXXVI. *IBERIS*. Lin. gen. n. 804. Gaertn. fruct. 2. p. 279. DC. syst. 2. p. 398.

Petala 2 exteriora majora. Silicula compressissima truncato-emarginata. Semina ovata pendula.

Sect. 1. IBERIDIUM.

Radícula descendens. Semen immarginatum. Septum simplex.

§. 1. *Pedicellis fructiferis corymbosis, caulibus suffrutescentibus perennibus.*

1. I. CONTRACTA (Pers! ench. 2. p. 186.) frutescens glabra, caule erecto elongato, foliis linearibus subcuneatis dentatis, siliculis umbellatis confertis. 3 in Hispaniâ Corymbi fructiferi contractissimi. DC. syst. 2. p. 405.* (v. s. in h. Juss.)
♀ ciliolata, I. fruticosa, foliis oblongis crassis, etc. Ant. Juss! in h. Juss.
2. I. PRUITI (Tineo! pug. sic. 1. p. 11. n. ix.*) basi suffrutescens glabra, foliis obovato-spathulatis integris aut subdentatis, siliculis subcorymbosis emarginatis. 2 5 in montibus Nebrodensibus Siciliæ. Ad sequentem valdè accedit, sed differt foliis glaberrimis, floribus albis, fructibus potius brevissimè racemosis confertissimis. DC. syst. 2. p. 404. (v. s.)
3. I. TENOREANA (DC. syst. 2. p. 404.*) basi suffrutescens puberula, foliis subcarnosis crenatis, inferioribus obovatis basi attenuatis, superioribus oblongo-linearibus, siliculis subcorymbosis emarginatis. 2 5 in regno Neapolitano. I. cepæfolia Tenor! prod. flor. nap. p. xxxvii. non Lin. Flores purpurascens aut albescentes, umbellati. (v. s.)

§. 2. *Pedicellis fructiferis corymbosis, caulibus herbaceis annuis aut biennibus.*

4. I. SPATHULATA (Berg. phyt. ic. DC. fl. fr. 4. p. 716.*) herbacea glabra, foliis spathulatis integris subcarnosis, siliculis corymbosis sinu angusto acuto emarginatis. ① in Pyrenæorum locis lapideis calcareis schistosisve aquâ nivali irrigatis. I. carnosa Willd. spec. 3. p. 455. Flores purpurascens, semina crassa rufo-fusca. DC. syst. 2. p. 404.* (v. v.)
5. I. NANA (All! auct. p. 15. t. 2. f. 1.*) herbacea glabra, foliis subrotundo-spathulatis integris subcarnosis, siliculis corymbosis emarginatis sinu latiusculo obtuso. ① ② in Pedemontio, Delphinatu, Gallo-provinciâ I. Anrosica Vill. dauph. 1. p. 349. 3. p. 289.* Flores purpur. DC. syst. 2. p. 403.* (v. v.)
6. I. VIOLACEA (Brown! in h. kew. ed. 2. vol. 4. p. 85.) herbacea glabriuscula, foliis perfoliatis spathulatis obtusis dentatis integerrimisq; ciliatis, corymbo subumbellato, calycibus dorso pilosulis. ① hab. . . . Flores violacei, fructus ignoti. DC. syst. 2. p. 403.* (v. s. in herb. Banks.)
7. I. TAURICA (DC. syst. 2. p. 402.*) herbacea glabriuscula, foliis ciliatis subcarnosis, inferioribus spathulatis apice subbidentatis, superioribus linearibus, siliculis corymbosis emarginatis, lobulis obtusis stylo brevioribus. ① ② in Tauriâ. I. ciliata. Willd. spec. 3. p. 445.* non All. (v. s.)
8. I. CILIATA (All. Ped. auct. p. 15.* non Willd.) herbacea glabriuscula, foliis linearibus integris basi ciliatis, siliculis corymbosis emarginatis lobulis obtusis stylo subæqualibus. ② in rupestribus agri Nicæensis et in Galloprovinciâ circa Brignoles. Flores ferè Iberidis pinnatæ. DC. syst. 2. p. 402.* (v. v.)
9. I. LINIFOLIA (Lin! spec. 905.) herbacea glabra, foliis linearibus integerrimis, radicalibus subdentatis, siliculis corymbosis bidentatis. ① ② in agro Nicæensi Gallo-provinciâ, Occitaniâ et Hispaniâ — Garid. Aix 459. t. 105. Valdè affinis I. umbellatæ. Habitus Lepidii Iberidis. — DC. syst. 2. p. 401.* (v. s.)
♀ albiflora, (Desf. in herb. Mus. Par.) in Lusitaniâ.
10. I. UMBELLATA (Lin! spec. 906.) herbacea glabra, foliis lanceolatis acuminatis, inferioribus serratis, superioribus integerrimis, siliculis umbellatis acutissimè bilobis. ① in Italiâ, Insulâ Cretâ et Hispaniâ. Curt. bot. mag. t. 106. Fl. sæpius purpurascens, rarius albi DC. syst. 2. p. 401.* (v. v.)
11. I. LAGASCANA (DC. syst. 2. p. 400.*) herbacea piloso-pubescent, foliis oblongis subspathulatis apice dentatis, siliculis subcorymbosis acutissimè

bilobis. ① in regno Valentino et Murciæ. I. spathulata Lag! fl. hisp. ined. non Berg. (v. s.)

§. 3. *Pedicellis fructiferis racemosis, caulibus herbaceis annuis biennibusve.*

12. I. ODORATA (Lin. sp. 906.) herbacea glabra, foliis linearibus dentatis basi ciliatis apice dilatatis, siliculis subrotundis, emarginaturæ lobis acutis patulis stylo brevioribus. ① in Cretâ.— Clus. hist. 2. p. 132. f. 1. Affinis I. pinnatæ, sed notis indicatis sat distincta. DC. syst. 2. p. 400. (v. s.)

13. I. PINNATA (Gouan! h. monsp. 319.) herbacea glabra, foliis pinnatifidis, racemis corymbosis post anthesin parum elongatis. ① in segetibus Hispaniæ, Galliæ australis, et Italiæ. Lob. ic. t. 217. f. 2. Flores albi, odorati, corymbosi, DC. syst. 2. p. 399.* (v. v.)

β. *crenata*, foliis inciso-crenatis. I. crenata. Lam! dict. 3. p. 213.*

14. I. INTERMEDIA (Guersent! in bull. philom. n. 82. t. 21.*) herbacea, foliis lanceolatis glabris obtusis integris aut radicalibus subdentatis, floribus mox racemosis, siliculis ovatis truncatis sen latissimè emarginatis. ② in rupibus calcareis inter Rothomagum et Duclair. Media inter I. amaram et umbellatam. DC. syst. 2. p. 399.* (v. s.)

15. I. AMARA (Lin! spec. 906.) herbacea, foliis lanceolatis acutis subdentatis, floribus corymbosis mox racemosis, siliculis orbiculatis angustè emarginatis. ① in agris Europæ frequens. Smith engl. bot. t. 52. Flores albi. Sepala marginè membranacea. DC. syst. 2. p. 398.* (v. s.)

β. I. *ruficaulis* Lejeun. fl. spa 2. p. 58, foliis angustioribus ciliatis, habitu minore.

§. 4. *Pedicellis fructiferis racemosis, caulibus frutescentibus.*

16. I. CONFERTA (Lag! varied. 2. n. 22. 1805. p. 213.) caule suffrutescente humillimo, foliis subradicalibus confertissimis sublinearibus acutis glabris, scapo racemifero nudo. 2/3 in Hispaniæ montibus Legionensibus et in dumetis prope Arvas. Flores albi, juniores apice corymbosi, dein mox racemosi. DC. syst. 2. p. 398.* Deless! ic. sel. 2. t. 54. (v. s.)

17. I. GARREXIANA (All! ped. n. 920. t. 40. f. 3. et t. 54. f. 2.) frutescens, foliis oblongis basi attenuatis obtusis integerrimis glabris, floribus corymbosis. 5/6 in saxosis apricis Pedemontii et Pyrenæorum. I. sempervirens β Willd. spec. 3. p. 453. Flores albi. Media inter I. sempervirentem et saxatilem. Characteres prioris, habitum posterioris exhibens. DC. syst. 2. p. 397. (v. v.)

18. I. SEMPERVIRENS (Lin! spec. 905.) frutescens, foliis oblongis obtusis basi attenuatis glabris, floribus longè racemosis, siliculis sinu angustè emarginatis. 5 in rupibus Cretæ. Sibth. et Smith fl. græc. t. 620. ex prodr. 2. p. 9. DC. syst. 2. p. 397.* (v. s.)

19. I. SUBVELUTINA (DC. syst. 2. p. 397.*) frutescens, foliis linearibus acutis integerrimis utrinquè pube brevi subvelutinis, floribus mox racemosis. 3/4 in Hispaniâ. I. sempervirens. Lag! elench. h. madr. p. 19. n. 253. Aff. I. saxatili et Garrexianæ. (v. s.)

20. I. PUBESCENS (Willd! enum. suppl. p. 43.*) frutescens, foliis ciliatis obtusis lineari-spathulatis, inferioribus apice subdentatis, floribus corymbosis. 5 hab. . . . Flores speciosi pallidè violacèi demùm subracemosi: fruct. ign. DC. syst. 2. p. 396.* (v. s. ex h. Willd.)

21. I. SAXATILIS (Lin! amœn. 4. p. 321.) frutescens, foliis linearibus integerrimis subcarnosis subacutis glabris ciliatisve, floribus corymbosis. 3/4 — Moris. oxon. 2. p. 298. s. 3. t. 18. f. 32. DC. syst. 2. p. 396.* (v. v.)

α. *thymbræfolia*, foliis ciliatis acutis. Moris. ox. 2. p. 298. s. 3. t. 18. f. 33.
β. *corifolia*, foliis glabris obtusis. — in monte Ventoso. Clus. hist. 2. p. 132. ic. Sims bot. mag. t. 1642.

- γ. *vermiculata*, foliis subciliatis obtusis, siliculæ sinu paululum obtuso et subdilato. — in Tauriâ. Forsan species propria ex cl. Andrz.
22. I. *CAPPADOCICA* (Willd. spec. 3. p. 452.*) frutescens, foliis strigosis inferioribus spathulatis, superioribus linearibus acutis, floribus corymbois. 5 in Cappadociâ. Petala obovata. Flores nascentes corymbosi. DC. syst. 2. p. 395.* (v. s. in h. Mus. Par.)
23. I. *GIBALTARICA* (Lin. spec. 905.) frutescens, foliis cuneiformibus obtusis apice subdentatis subciliatis, floribus corymbosis. 3 ad Gibraltariam. Curt. bot. mag. t. 124. Flores albi pauco rubore suffuso. DC. syst. 2. p. 395.* (v. v.)

Sect. II. *IBERIDASTRUM*.

Radicula horizontalis. Semen submarginatum. Septum ferè duplex. — Semen ferè ut in *Biscutellâ* et ideò forsan genus (Andrz.), sed habitus inter *Iberides* hanc retinet.

24. I. *SEMPERFLORENS* (Lin! spec. 904.) frutescens, foliis cuneatis spathulatisve obtusis integerrimis glabris, floribus corymbosis, siliculis apice truncato-submarginatis lobulis obsoletis. 3 in Persiâ? et in rupibus Siciliæ. — Weinm. phyt. t. 973. f. c. — DC. syst. 2. p. 394.* (v. s. et v.)

†. *Iberides non satis notæ.*

25. I. *LINKARIFOLIA* (DC. syst. 2. p. 405.*) glabra, caule erecto, foliis linearibus integerrimis, siliculis racemosis bifidis. — in Novâ-Hollandiâ. *Lepia linifolia*. Desv! journ. 3. p. 166 et 181. (v. s. in h. Mus. Par.)
26. I. *PYRENAICA* (Lapeyr. abr. pyr. 370.) herbacea glabra, foliis ellipticis integerrimis, inferioribus oppositis, siliculis racemosis latè emarginatis. ① in Pyrenæicâ valle Gistain propè Sin. Flores racemosi, nudi, albi, numerosi. DC. syst. 2. p. 406.*

XXXVII. *BISCUTELLA*. Lin. gen. n. 808. Gaertn. fruct. 2. p. 278. t. 141.

Silicula plana biscutata, loculis 1-spermis, orbiculatis, lateraliter axi adnatis, stylo longo persistente. Semina compressa. Embryo inversus. — Flores flavi.

Sect. I. *JONDRABA*. DC. syst. 2. p. 407. — Med. gen. nov. t. 1. f. 14.

Calyces basi bicalcarati. Glandulæ tori valdè prominentes.

1. B. *AURICULATA* (Lin. spec. 911.) calycibus obtusè bicalcaratis, siliculis glabris in disco punctis elevatis scabris, lobis apice in stylum coeuntibus. ① in agris cultis Europæ australis et in Mauritiâ. DC. diss. n. 1. t. 1. f. 2. Flores pallidè flavi. DC. syst. 2. p. 407.* (v. v.)
2. B. *ERIGERIFOLIA* (DC. diss. n. 2. syst. 2. p. 408.*) calycibus obtusè bicalcaratis, siliculis glabris lævibus in stylum apice subcoeuntibus. ① inter segetes Hispaniæ. Deless! ic. sel. 2. t. 55. B. auriculata Dufour in ann. gen. 7. p. 298. (v. s.)
3. B. *HISPIDA* (DC. diss. n. 3. t. 1. f. 1. et syst. 2. p. 408.*) calycibus acutè bicalcaratis, siliculis glabris in disco punctis elevatis scabris apice in stylum non coeuntibus, caule hispido. ① in apricis montosis Gallo-provinciæ superioris et Pedemontii. Valdè accedit ad duas priores, sed certissimè differt. (v. s. et v.)
4. B. *CICORIIFOLIA* (Loisel! add. p. 167.) calycibus acutiusculè bicalcaratis, siliculis glabris in disco punctis elevatis scabris in stylum non coeuntibus, caule villosa. 2 ad radices Pyrenæorum propè Baguères de Luchon. DC. diss. n. 4. t. 2.* syst. 2. p. 409.* Flores magni. Folia runcinato-pinnatifida. (v. s.)

Sect. II. *THLASPIDIUM*. DC. syst. 2. p. 409. Med. gen. nov. t. 1. f. 5.

Calyces basi æquales. Glandulæ tori vix prominentes.

§. 1. *Species annuæ.*

5. *B. LYRATA* (Lin! mant. 354.) siliculis in utroque disco piloso-hispidis, foliis radicalibus lyratis. (I) in Beticâ, Hispaniâ, Calabriâ, Africâ bor. — Bocc. sic. 45. t. 23. Species habitu polymorpha, sed in plurimas, meo sensu, non separanda. DC. diss. n. 13. syst. 2. p. 410.* (v. s. et v.)
6. *B. RAPHEPIFOLIA* (Poir. voy. bar. 2. p. 198.) siliculis lævibus glabris, foliis radicalibus lyratis. (I) in Siciliâ, Mauritaniâ. A priore differt siliculâ lævi glaberrimâ et forsan foliis minùs incis. DC. diss. n. 14. syst. 2. p. 410. (v. s.)
7. *B. MARITIMA* (Tenor! prodr. fl. nap. p. 38. fl. nap. t. 61.) siliculis lævibus margine ciliatis, foliis radicalibus lyratis. (I) in maritimis insulæ Caprearum. Duabus prioribus valde similibus. Differt tantum siliculis nec omnino glabris, nec piloso-hispidis, sed in disco utroque glabris lævibus, secus margines et lineam longitudinalem pube confertâ brevi ciliatis. An mera varietas? DC. syst. 2. p. 410. (v. s.)
8. *B. CILIATA* (DC. diss. n. 9. syst. 2. p. 410.*) siliculis lævibus margine ciliatis, caule erecto elongato folioso, foliis sessilibus oblongis remotè dentatis. (I) in Italiâ, Hispaniâ. *B. coronopifolia*. Willd! spec. 3. p. 474. DC. ic. gall. rar. 1. p. 12. t. 39. non Lin. Sub nomine *B. didymæ* cum *Bisc. apulâ* mixta adest in herb. Linnæano. (v. v. et s.)
9. *B. DEPRESSA* (Willd. enum. 2. p. 673.) siliculis lævibus margine ciliatis, caule humili subdiffuso, foliis oblongo-obovatis basi cuneatis remotè dentatis. (I) in Ægypto. DC. diss. n. 10. Siliculæ constanter in margine et secus lineam suturalem ciliatæ. An prioris varietas? DC. syst. 2. p. 411. (v. v.)
10. *B. MICROCARPA* (DC. diss. n. 11. syst. 2. p. 411.*) siliculis in disco scabro-hispidis margine ciliatis, caule subnundo ramosissimo, foliis subradicalibus oblongis dentato-sinuatis. (I) in Beticâ. Flores longè racemosi, flavi, parvi. (v. s. in h. Desf.)
11. *B. ERIOCARPA* (DC. diss. n. 12. t. 9. f. 2. syst. 2. p. 411.) siliculis in disco piloso-hispidis, caule erecto subsimplici, foliis oblongo-cuneatis subdentatis. (I) in itinere trans Hispaniam ad Mogador reperit Broussonet. Siliculæ magnitudine *B. apulæ*. (v. s.)
12. *B. COLUMNÆ* (Tenor! prod. fl. nap. p. 38.) siliculis pube minimâ disco et margine scabris, foliis radicalibus obovato-cuneatis acutis dentatis, caule subnundo basi hispido. (I) in Apuliâ et Lucaniâ. *B. didyma*, ≠ Lin. spec. 911. ex syn. — Moris. oxon. 2. p. 249. s. 3. t. 9. f. 12. Valde aff. *B. eriocarpæ* et *Apulæ*, sed differt siliculis duplò majoribus. Variat foliis sinuato-lyratis. DC. syst. 2. p. 412.* (v. s.)
13. *B. APULA* (Lin. mant. 254.) siliculis pube minimâ disco et margine scabris, foliis lanceolatis serratis, caule folioso ramoso hirt. (I) in montanis Italiæ, in Rhodo et agro Argolico. Lam. ill. t. 560. f. 1. non dict. Flores parvi. Siliculæ approximatæ. DC. diss. n. 15. syst. 2. p. 412.* (v. v.)
14. *B. LEOCARPA* (DC. diss. n. 16. syst. 2. p. 413.) siliculis glabris lævibus, foliis lanceolatis serratis, caule folioso hirt. ramoso. (I) in Oriente. *B. Apula* Gaertn. fruct. 2. p. 279. t. 141. Priori nimis affinis, sed diversa videtur fructu, etiam nascente, glaberrimo nec pube minutâ scabro. (v. s.)
15. *B. OBOVATA* (hort. par. DC. diss. n. 18. syst. 2. p. 413.*) siliculis glabris lævibus, foliis subradicalibus obovatis in petiolum attenuatis grossè dentatis glabriusculis. (I) hab. species similis *B. montanæ*, sed differre videtur radice (I) nec \mathcal{L} etc. (v. s.)

§. 2. *Species perennes.*

16. *B. MONTANA* (Cav. ic. 2. p. 50. t. 177. ic. opt.) siliculis glabris lævibus, foliis subradicalibus obovato-cuneatis dentatis tomentos. \mathcal{L} in rupibus regni Valentini. DC. diss. n. 17. syst. 2. p. 413.*
17. *B. LEVIGATA* (Lin! mant. 225.) siliculis glabris lævibus, foliis subradica-

libus piloso-scabris oblongis in petiolum attenuatis subdentatis integrise, caulinis linearibus paucis integerrimis. \mathcal{U} in apricis montosis Alpium, Pyrenæorum, Jurassi, Hungariæ, etc. Jacq. fl. austr. t. 339.* DC. syst. 2. p. 413.* (v. s.)

β. *B. alpestris* Waldst. et Kit. pl. rar. hung. 3. p. 253. t. 228.* DC. diss. n. 27.

γ. *B. glabra* Clairv. herb. val. 216.

18. *B. LUCIDA* (DC. diss. n. 20. t. 7. syst. 2. p. 414.*) siliculis glabris lævibus, foliis glabris, plerumque radicalibus oblongis. \mathcal{U} in montibus Umbræ et Aprutii. (v. v.)

α. foliis inferioribus dentatis. *B. lucida*. Balb! hort. Taurin.

β. foliis omnibus integris. *B. spathulata*. Lam. dict. 3. p. 620. — Barr. ic. t. 254.

19. *B. CORONOPIFOLIA* (All. ped. n. 907.) siliculis glabris lævibus, foliis piloso-scabris plerumque radicalibus pinnatifidis, lobis utrinque 2-3 remotis. \mathcal{U} in Hispaniâ, Galliâ austr. et Pedemontio. *Sisymbrium* Valentinum. Lin! spec. 920? (excl. syn.) ex herb. — DC. diss. n. 22. t. 18. syst. 2. p. 414.* (v. v.)

20. *B. AMBIGUA* (DC. diss. n. 23. t. 11. f. 1. syst. 2. p. 415.) siliculis glabris lævibus, foliis piloso-scabris, radicalibus sinuato-dentatis basi attenuatis, caulinis paucissimis basi subcordatis semi-amplexicaulibus. \mathcal{U} in sterilibus apricis Niceæ, Galliæ austr. et Regni Valentini. (v. v.)

α. dentium sinubus revolutis. *B. coronopifolia*. Lin! mant. 255?

β. dentium sinubus planis.

21. *B. SAXATILIS* (Schleich! cent. exs. n. 69.) siliculis glabris punctis elevatis in disco scabris, foliis piloso-scabris plerumque radicalibus oblongis. \mathcal{U} in sterilibus montosis Europæ austr. DC. diss. n. 24. t. 10. syst. 2. p. 415.* (v. s.)

α. foliis integris. *B. longifolia*. Vill. Dauph. 3. p. 305.

β. foliis inciso-dentatis sinuato-pinnatifidisve. *B. mollis*. Lois. not. p. 168.

γ. foliis hinc indè dentatis, caule humillimo. *B. intermedia*. Gouan ill. p. 42.

22. *B. SEMPERVIRENS* (Lin! mant. 255.) siliculis glabris punctis elevatis in disco scabris, foliis plerisque radicalibus erectis lineari-lanceolatis tomentoso-incanis subintegerrimis. \mathcal{U} in Hispaniâ et Lusitaniâ. Bocc. mus. 2. p. 167. t. 122. DC. diss. n. 25. syst. 2. p. 416. (v. b.)

23. *B. TOMENTOSA* (Lag! in litt. et DC. syst. 2. p. 416.) siliculis tenuissimè pubescentibus punctato-scabris, foliis oblongis basi angustatis repando-dentatis mollissimè tomentosis, ramis amplexicaulibus. \mathcal{U} in Hispaniâ. Planta subpedalis. (v. s.)

24. *B. STENOPHYLLA* (Duf. in ann. gen. 7. p. 299.) siliculis disco punctato scabris glabris, petalis biauriculatis, foliis radicalibus piloso-hispidis subrigidis lanceolato-linearibus remotè dentatis subpinnatifidis, caulibus subnudis subsimplicibus. \mathcal{U} in regno Valentino. An satis à *B. coronopifoliâ* et à *B. saxatili* differt?

†. *Species vix notæ.*

25. *B. MAJOR* (Schkuhr. handb. 2. n. 1822.) calyces lateraliter gibbosi. Siliculæ læves venosæ.

26. *B. ANGUSTIFOLIA* (Schkuhr. handb. 2. n. 1823.) calyces lateraliter gibbosi. Siliculæ verrucosæ asperæ.

XXXVIII. MEGACARPÆA. DC. syst. 2. p. 417.

Silicula plana biscutata utrinque emarginata, loculis compressissimis margine alatis 1-spermis lateraliter axi adnatis, stylo nullo. Radicula adscendens.

1. *M. LACINIATA* (DC. syst. 2. p. 417.*) foliis villosis, radicalibus petiolatis pinuatipartitis, lobis grossè incisis dentatis. \mathcal{U} ad lacum Inderiensem et ad

colles gypsaceos Arsagar deserti trans-Wolgensis. *Biscutella megalocarpa* Fisch! in litt. DC. diss. bisc. n. 5. t. 3.* Flores parvi mihi non satis noti. (v. s.)

Species non satis nota.

2. *M. ANGULATA* (DC. syst. 2. p. 418.) foliis glabris integris aut repando-angulatis. — In Sibiria. Frustulum valde incompletum habeo, sed evidentissime prioris congener et ab eâ distinctissimum.

XXXIX. CREMOLOBUS. DC. syst. 2. p. 418.

Silicula pedicellata plana biscutata, loculis marginatis ex apice axeos pendulis, stylo subpyramidato. Semina in loculis solitaria. Radicula adscendens.

1. *C. PERUVIANUS* (DC. syst. 2. p. 419.*) caule suffrutescente, foliis ovatis serratis, stylo thecaphoro brevior. § in Peruvia et Chili. *Biscutella Peruviana* Lam! dict. 3. p. 620. DC. diss. bisc. n. 6. t. 4. (v. s.)
2. *C. SUFFRUTICOSUS* (DC. syst. 2. p. 419.*) caule suffrutescente, foliis ovali-oblongis serratis, stylo thecaphoro longiore. § in Peruvia. *Biscutella suffruticosa*. DC. diss. bisc. n. 7. t. 5.* A Cr. Chilensi facillimè distinguitur caule frutescente nec herbaceo, foliis latioribus magis serratis. (v. s.)
3. *C. CHILENSIS* (DC. syst. 2. p. 419.*) caule herbaceo, foliis oblongis subintegris, stylo thecaphoro sublongiore. ① in Chili. *Biscutella Chilensis* Lagascl! in litt. DC. diss. bisc. n. 8. t. 6.* (v. s.)

XL. MENONVILLEA. DC. syst. 2. p. 419.

Silicula substipitata biscutata, loculis margine in alam expansis discos parallelos conficientibus.

1. *M. LINEARIS* (DC. syst. 2. p. 420.*) ¶ in Peruvia et Chili. Folia radicalia linearia integra aut rarius versus apicem grossè et irregulariter serrata. Petala linearia, calyce duplò longiora. Deless! ic. sel. 2. t. 56. (v. s. in h. Mus. Par. et Lamb.)

Tribus IV. *EUCLIDIEÆ* seu *PLEURORHIZEÆ* *NUCAMENTACEÆ*. DC. syst. 2. p. 420.

Silicula indehiscens, valvis concavis indistinctis aut non secedentibus, septo elliptico interdum evanido. Semina ovalia, paucissima. Cotyledones planæ, accumbentes, septo (dum adsit) parallelæ.

XLJ. EUCLIDIUM. Brown in h. Kew. ed. 2. v. 4. p. 74. DC. syst. 2. p. 421.

Silicula drupacea, ovata, suturis manifestis, stylo subulato, loculis 1-spermis. — Flores parvi albi subspicati.

1. *E. SYRIACUM* (Brown in h. Kew. ed. 2. v. 4. p. 74.) siliculis scabris, stylo subulato persistente, foliis caulinis petiolatis lanceolatis. ① in Syria, Tauria, Iberia, Moldavia, etc. *Bunias Syriaca* Gaertn. fruct. 2. p. 290. t. 141. f. 11.* Flores spicati, distantes. — DC. syst. 2. p. 422.* (v. s.)
2. *E. TATARICUM* (DC. syst. 2. p. 422.*) siliculis glabris, stylo demum deciduo, foliis caulinis linearibus sessilibus. ① in collibus Inderiensibus et in desertis circa Astracan. Vella tenuissima Pall! itin. ed. gall. app. n. 344. t. 77. f. 2.* Flores parvi, vix pedicellati. (v. s.)

XLII. OCHTHODIUM. DC. syst. 2. p. 423.

Silicula coriacea subglobosa, stigmatè sessili, septo crasso, loculis 1-spermis. Cotyledones planæ ovali-oblongæ oblique accumbentes.

1. *O. ÆGYPTIACUM* (DC. syst. 2. p. 423.*) ① in Ægypto, et Syriâ. *Bunias Ægyptiaca* Lin! syst. nat. 3. p. 231. Jacq. h. vind. t. 145. Folia caulina petiolata, pinnatipartita anglyrata. Petala flava, calyce majora. (v. v. et s.)

XLIII. *PUGIONIUM*. Gaertn. fruct. 2. p. 291. t. 142. DC. syst. 2. p. 424.

Silicula coriacea, transversè ovalis, utrinquè echinata et in pugionem producta, abortu 1-locularis 1-sperma.

1. *P. cornutum* (Gaertn. l. c.) — in Oriente. *Bunias cornuta* Lin! spec. 935. Flores parvuli, albi, *Erophilæ* vernæ haud majores. Petala angusta integerrima. (v. s. in h. Lin.)

Trib. V. *ANASTATICÆ* seu *PLEURORHIZÆ SEP-TULATÆ*. DC. syst. 2. p. 424.

Silicula longitudinaliter dehiscens, valvis concavis intus transversè septula horizontalia semina separantia gerentibus. Semina immarginata. Cotyledones planæ, accumbentes, septo parallelæ.

XLIV. *ANASTATICA*. Gaertn. fruct. 2. p. 286. t. 141. DC. syst. 2. p. 425.

Silicula ventricosa valvis extus ad apicem appendiculatis. Petala obovata.

1. *A. HIEROCUNTINA* (Lin! spec. 895.) ① in arenosis Ægypti. Jacq. h. vind. t. 58. Schkuhr. handb. t. 179. Herba parva humilis, folia oblonga aut ovata in petiolum longè attenuata. Petala alba. Silicula subpubescens. DC. syst. 2. p. 426.* (v. v.)

XLV. *MORETTIA*. DC. syst. 2. p. 426.

Silicula ovata, valvis non appendiculatis. Petala linearia.

1. *M. PHILEANA* (DC. syst. 2. p. 426.) — in Nubiâ propè insulam Philæ. *Sinapis Phileana* Delil! fl. æg. p. 99. t. 33. f. 3.* Folia alterna ferè sessilia. Flores distantes parvi. Silicula oblonga extus velutina. (v. s.)

Trib. VI. *CAKILINEÆ* seu *PLEURORHIZÆ LOMEN-TACÆ*. DC. syst. 2. p. 427.

Siliqua aut silicula transversè in articulos 1-2-loculares 1-2-spermos secedens. Semina immarginata. Cotyledones planæ, accumbentes, septo (dùm adsit) parallelæ.

XLVI. *CAKILE*. Tourn. inst. 49. t. 483. Gaertn. fruct. 2. p. 287. t. 141. DC. syst. 2. p. 427.

Silicula biarticulata compressa, articulo superiore ensiformi aut ovato. Semina in loculis solitaria; superioris erectum, inferioris pendulum.

1. *C. MARITIMA* (Scop. fl. carn. n. 844.) siliculæ articulo superiore ensiformi. ① in arenosis Europæ secus Oceanum et Mediterraneam. Lam. ill. t. 554. f. 1. — *C. Serapionis* Gaertn. fruct. 2. p. 287. t. 141. f. 12. — DC. syst. 2. p. 428.* (v. s.)

β. *sinuatifolia*, foliis dentato-sinuatis. *C. Ægyptiaca*. Willd. spec. 3. p. 417. — Moris. ox. 2. p. 231. s. 3. t. 7. f. 20.

2. *C. AMERICANA* (Nutt. gen. amer. 2. p. 62.) siliculæ articulo superiore ovato acuto. ① in maritimis Americanis. *C. Ægyptiaca* Tuss. fl. ant. 1. f. 17. Differt à *C. maritimâ*, foliis oblongis non cuneatis, etc. DC. syst. 2. p. 429.* (v. s. in h. Smith et Humb.)

β. *Cubensis* (H. B. et K. nov. gen. et spec. 5. p. 75.) Folia integra aut vix sinuato-dentata, nunquam pinnatifida.

3. *C. ÆQUALIS* (L'her! herb. et diss. Cak. ined. p. 1. cum ic.) siliculæ articulo superiore basi subtetragono apice longè rostrato. ① in Martinicâ; Flores

albi. Siliqua linearis subuniformis. Semina oblonga badia. DC. syst. 2. p. 430.*
Deless! ic. sel. 2. t. 57. (v. s.)

XLVII. CORDYLOGARPUS. Desf. fl. atl. 2. p. 79. t. 152. non Smith. DC. syst. 2. p. 434.

Siliqua teretiuscula torosa, articulis plurimis, terminali crasso globoso echinato. Semina omnia pendula.

1. *C. MUICATUS* (Desf. fl. atl. 2. p. 79. t. 152.*) ① propè Mayane Algeriæ. Folia inferiora ovato-oblonga, in petiolum decurrentia. Petala pallidè lutea. Siliquæ patentiores. DC. syst. 2. p. 435.* (v. s. in h. Deless.)

XLVIII. CHORISPORA. DC. syst. 2. p. 435.

Siliqua teretiuscula, articulis plurimis æqualibus. Semina omnia pendula.

§ 1. *Petalis purpurascens integris.*

1. *C. TENELLA.* (DC. syst. 2. p. 435.*) siliquis foliisque glabris, superioribus lanceolatis dentatis, infimis pinnatifidis ① in ruderalis Tauriæ, in deserto Caspico, ad ripas Wolgæ etc. *Raphanus tenellus* Pall! itin. 3. app. n. 105. t. L f. 3. ed. gall 8°. app. n. 356. t. 102. f. 2.* *Chorispermum tenellum* Brown in h. Kew. ed. 2. v. 4. p. 129. (v. v. et s.)

β. *taraxacifolia.* *Cheiranthus taraxacifolius* Schrank. mem. soc. Ratisb. 1818. p. 164. (ex Bieb.)

γ. *arcuata.* *Hesperis arcuata* Nocca pl. sel. hort. Ticin. 1. p. 3. t. 2.* *Raphanus arcuatus* Willd. spec. 3. p. 562.*

2. *C. STRICTA* (DC. syst. 2. p. 436.*) siliquis caule foliisque inferioribus hispida erectis, foliis linearibus dentatis. ① ad lacum Inder-koe deserti Kirghisorum. *Raphanus strictus* Fisch! in litt. Bieb. suppl. fl. taur. p. 452. Semina ovalia, compressa, alâ membranaceâ cincta. (v. s. comm. à cl. Fisch.)

§. 2. *Petalis flavis emarginatis.*

3. *C. SIBIRICA* (DC. syst. 2. p. 437.*) siliquis foliisque subglabris, foliis omnibus sinuato-pinnatifidis. ① in Sibiria Altaica. *Raphanus Sibiricus* Lin! spec. 925. Murr. comm. goett. 1775. p. 48. t. 11. Flores flavi, ferè magnitudine *Bunadiis Erucaginis* Semina ovalia, rufa, compressa. (v. s.)

4. *C. IBERICA* (DC. syst. 2. p. 437.*) siliquis hirtis vix stylo longioribus, foliis linearibus runcinato-dentatis. ① in Iberiæ lapidosis. *Raphanus Ibericus* Bieb. fl. taur. 2. p. 129.* suppl. p. 451. Planta tota odore nauseoso pollet. Deless! ic. sel. 2. t. 58. (v. s. comm. à cl. Stev.)

Subordo II. NOTORHIZEÆ. (o ||)

Cotyledones planæ incumbentes. Radicula dorsalis nempè medio dorso cotyledonum incumbens. Semina ovata, non marginata.

Trib. VII. SISYMBREÆ seu NOTORHIZEÆ SILIQUOSÆ.
DC. syst. 2. p. 438.

Siliqua bilocularis longitudinaliter dehiscens, valvis concavis carinatisve. Semina ovata aut oblonga, immarginata. Cotyledones planæ, incumbentes, septo contrariæ. — Tribus distinctissima ex caracteribus spermicis, sed habitu ad Arabideas accedens.

XLIX. MALCOMIA. Brown in h. Kew. ed. 2. v. 4. p. 121. DC. syst. 2. p. 438.

Siligna terestiuscula. Stigma simplex acuminatissimum.

1. *M. AFRICANA* (Brown in h. Kew. ed. 2. v. 4. p. 121.) caule ramoso diffuso, foliis lanceolatis subdentatis pube 2-4-partitâ, pedicellis calyce persistente brevioribus, siliquis scabris. ① in arenosis Africæ, Siciliæ, Hispaniæ, Galliæ mediterran., Iberiæ. *Hesperis Africana* Lin! spec. 928. — Bocc. sic. 77. t. 42. f. 1. Fl. purpurascens. Semina oblongo-ovata, rufa. — DC. syst. 2. p. 439.* (v. v.)
2. *M. TARAXACIFOLIA* (DC. syst. 2. p. 439.*) caule erecto simplici, foliis oblongis inciso-dentatis, pube 3-partitâ, pedicellis calyce deciduo brevioribus, siliquis glabris subtetragonis. ① hab. *Cheiranthus taraxacifolius* Balb! cat. h. Taur. app. 1814. p. 10. Flores parvi (v. s. comm. à cl. Balb.)
3. *M. LAXA* (DC. syst. 2. p. 440.*) caule ramoso basi subpiloso, foliis ovatis acutis dentato-angulatis siliquisque glabris, pedicellis calyce brevioribus. ① in Sibiria *Hesperis laxa* Lam. dict. 3. p. 325.* ? Flores parvi, purpurascens. Semina parva oblongo-ovata, rufa. (v. s. in M. Mus. Par.)
4. *M. CHIA* (DC. syst. 2. p. 440.*) caule erecto ramoso, foliis oblongis integris, pube bipartitâ adpressâ, pedicellis calycis longitudine, siliquis teretiusculis pubescentibus, stylo brevissimo. ① in saxosis insulæ Chio. *Cheiranthus Chius* Lin! spec. 924. — Dill. elth. 80. t. 147. f. 178. Flores purpurascens, dimidio quàm *M. maritimæ* minores. (v. s.)
5. *M. MARITIMA* (Brown in h. Kew. ed. 2. v. 4. p. 121.) caule erecto ramoso, foliis ellipticis obtusis integris basi attenuatis, pube adpressâ 2-4-partitâ, pedicellis calyce subbrevioribus, siliquis pubescentibus apice longè acuminatis. ① in arenosis Europæ australis. *Cheiranthus maritimus* Lin. amœn. 4. p. 280. Curt. bot. mag. t. 166. Petala violacea. Siliquæ teretiusculæ. DC. syst. 2. p. 441.* (v. v.)
6. *M. INCRASSATA* (DC. syst. 2. p. 441.*) caule multiplici subdiffuso, foliis ovalibus petiolatis, superioribus oblongis, pube adpressâ bipartitâ, pedicellis calyce brevioribus post anthesin incrassatis, siliquis pubescentibus. ① in rupibus insularum Archipelagi Scio et Tenedos. Flores paulò majores et siliquæ duplò triplòve crassiores quàm *M. maritimæ*. Deless! ic. sel. 2. t. 59. (v. s.)
7. *M. ARENARIA* (DC. syst. 2. p. 442.*) caule erecto ramoso, foliis lanceolatis acutis, inferioribus dentatis sessilibus, pube stellatâ, pedicellis brevissimis, siliquis torulosis subulatis. ① in Mauritaniâ. *Hesperis arenaria* Desf! fl. atl. 2. p. 91. t. 162.* Petala violacea magnitudine circiter *M. maritimæ*. Siliquæ subulatæ torulosæ (v. s. in h. Desf.)
8. *M. PARVIFLORA* (DC. syst. 2. p. 442.*) caule erecto ramoso, foliis oblongis obtusis subintegris, pube tomentosâ stellatâ, pedicellis demùm calycis longitudine, siliquis pubescentibus. ① in arenosis maritimis Corsicæ, Occitanie, Galloprovinciæ et Italiæ. *Hesperis parviflora*. DC. fl. fr. 4. p. 654.* 5. p. 590. ic. gall. p. 10. t. 35. (v. v.)
9. *M. LYRATA* (DC. syst. 2. p. 443.*) caule erecto ramoso, foliis inferioribus lyratis petiolatis obtusis, pube adpressâ bipartitâ, pedicellis calycis longitudine, siliquis pubescentibus. ① in insulâ Cipro. *Hesperis rigida* Sibth! in h. Banks. *Cheiranthus lyratus* Sibth. et Smith fl. græc. t. 635. ex prodr. 2. p. 24. Flores purpurei similes *M. parvifloræ*. (v. s. in h. Banks.)
10. *M. LITTORÆA* (Brown in h. Kew. ed. 2. v. 4. p. 121.) caule multiplici erecto, foliis lanceolato-linearibus subintegris, tomento brevicanis, pedicellis calycis longitudine, siliquis incanis. ② in arenosis maritimis secus Oceanum et mare Mediterraneum. — Clus. hist. 1. p. 298. f. 2. Flores albido-flavi. Semina ovata immarginata. (v. v.)
11. *M. ALYSSOIDES* (DC. syst. 2. p. 444.*) caule erecto ramoso, foliis imbricatis obovatis obtusis subintegris pube stellatâ cano-tomentosis, pedicellis brevissimis. ② in Lusitaniâ. *Hesperis alyssoides*. Pers. ench. 2. p. 203. — Barr. ic. t. 804. excl. syn. in obs. relato. Flores ad apices ramorum 2-3, subsessiles. (v. s. in h. Vaill.)
12. *M. PATULA* (DC. syst. 2. p. 444.*) caule ramoso diffuso, foliis linearibus subintegris, pube brevi tomentosâ stellatâ canis, pedicellis calyce vix bre-

viribus, siliquis tenuibus pubescentibus longo stylo terminatis. ② circa Madritum in arenosis. *Hesperis arenaria* Lag. cat. h. madr. 1814. p. 20. non Desf. Flores M. littoreæ. (v. s.)

13. *M. BROUSSONETII* (DC. syst. 2. p. 445.*) caule erecto ramoso, foliis oblongis obtusis sinuatis, pube stellatâ tomentoso-canis, pedicellis calyce brevioribus, siliquis pubescentibus torulosis stylo longo terminatis. ① in agro Mogadoriensis. Deless! ic. sel. 2. t. 60. *Cheiranthus trilobus* Lin. spec. 925*? Flores M. littoreæ sed paulô majores. (v. s.)

14. *M. LACERÆ* (DC. syst. 2. p. 445.*) caule ramoso diffuso, foliis oblongis obtusè pinnatifidis, pube stellatâ subtomentosâ, pedicellis calyci demum æqualibus, siliquis pubescentibus torulosis stylo longo terminatis. ① in arenosis maritimis Lusitanix, Hispanix, agri Mogadoriensis, Orientis. *Cheiranthus lacerus* Lin! spec. 926. non syst. veg. *Hesperis pinnatifida* Desf. cor. tourn. 63. t. 47. ann. mus. 11. p. 377. t. 34.* (v. s.)

†. *Species non satis nota.*

15. *M. EROSA* (DC. syst. 2. p. 446.) ① in Bæticæ locis sabulosis. *Hesperis erosa* Lag. cat. h. madr. 1814. p. 20. Pubescens. Folia lineari-oblonga, sinuato-pinnatifida. Siliquæ pedunculatæ, subfiliformes, acutæ.

L. HESPERIS. *Lin. gen. n. 817. DC. syst. 2. p. 446.*

Siliqua teretiuscula aut subtetragona. Stigmata duo erecta conniventia. Calyx basi bisaccatus. Semina oblonga subtriquetra. Stamina edentula.

Sect. I. HESPERIDIUM. *DC. syst. 2. p. 447.*

Petalorum limbi lineares tristes, vespere odorati. Siliqua valvis carinatis subanceps, septo fungoso.

1. *H. ALYSSIFOLIA* (DC. syst. 2. p. 447.*) pedicellis villosis calyce villosissimo vix longioribus, petalorum laminis oblongis undulatis, foliis oblongis integerrimis pube stellatâ velutinis. — in Persiâ. Deless! ic. sel. 2. t. 61. Petala oblongo-linearia, sordidè in limbo purpurascientia, acuta. (v. s. in h. Deless.)
2. *H. TRISTIS* (Lin! spec. 927.) pedicellis longissimis patentibus rigidis siliquæ ancipitis margine incrassatæ latitudinem æquantibus, petalorum laminis oblongis obliquis. ② in agros et sylvarum margines in Austriâ, Pannoniâ, Transylvaniâ, Tauriâ, Rossiâ, Italiâ, etc. Curt. bot. mag. t. 730. Jacq. h. vind. 118. *Cheiranthus lanceolatus* Willd. spec. 3. p. 515.* Flores nocte olentes sordidè ex albedo aut ochroleuco fusco-rubentes. — DC. syst. 2. p. 447.* (v. s.)

Sect. II. DELOSMA. *DC. syst. 2. p. 448.*

Petalorum limbi obovati. Siliqua teretiuscula aut subtetragona, septo membranaceo; funiculi alati.

3. *H. LACINIATA* (All! pedem. n. 985. t. 82. f. 1.*) pedicellis calyce brevioribus, petalis obovato-oblongis, foliis obovatis inciso-dentatis, caule hispido. ② in rupibus apricis Pedemontii, Gallo-provinciæ et Delphinatûs. *Cheiranthus laciniatus* Poir. suppl. 2. p. 780.* Flores nunc purpurascens, nunc flavescens. DC. syst. 2. p. 448.* (v. v.)
4. *H. VILLOSA* (DC. syst. 2. p. 449.*) pedicellis calyce brevioribus, petalis obovato-oblongis, foliis oblongis acuminatis inciso-dentatis cauleque simplici hispido-villosis. ②? in Apuliâ. *Cheiranthus villosus* Spreng! in litt. *Hesperidi laciniatæ* nimis affinis. (v. s. comm. à cl. Spreng.)
5. *H. RUNCINATA* (Waldst. et Kit. pl. rar. hung. 2. p. 220. t. 200.*) pedicellis calyce longioribus, petalis obovatis submucronatis, foliis pubescentibus inferioribus lyrato-runcinatis, summis lanceolatis acuminatis, caule erecto apice paniculato. ② in fruticetis Hungariæ. DC. syst. 2. p. 449.* (v. v. in h. monsp.)

β. H. bituminosa Savi! h. pis. ann. 1808. An Willd. enum. suppl. 45?
Deilosma suaveolens Andr! cruc. ined. (v. s.)

6. *H. MATRONALIS* (Lam. dict. 3. p. 321. ill. t. 564. f. 1.) pedicellis calycis longitudine, petalis obovatis, siliquis erectis torosis glabris margine non incrassatis, foliis ovato-lanceolatis dentatis, caule erecto subsimplici. \mathcal{L} in dumetis et sepibus Europæ ferè totius. DC. syst. 2. p. 450.* (v. v.)

α. hortensis, caule firmo recto, floribus sæpius odoratis, petalis frequentius emarginato-mucronatis, foliis ovato-lanceolatis nunquam cordatis. *H. matronalis*. Lin. spec. 927. — Moris. hist. 2. p. 251. s. 3. t. 10. f. 1. var. * *albiflora*. † simplex †† plena. ** *purpurea*. † simplex. †† plena. *** *variegata*. **** *foliiflora*.

β. sylvestris, habitu sæpè laxo, foliis saltem inferioribus basi plus minus cordatis et grossius sæpè dentatis, flores semper purpurascensibus, rarius odoratis, petalorum limbis obtusis. *H. inodora* Lin! spec. 927. Jacq. fl. austr. t. 347. Smith engl. bot. t. 731.

γ. Sibirica, foliis angustioribus oblongo-lanceolatis acuminatissimis. *H. Sibirica* Lin. spec. 927.

7. *H. HETEROPHYLLA* (Tenor! fl. nap. prodr. p. xxxix.) pedicellis calycis longitudine, petalorum laminis obovatis, caule ramoso diffuso siliquisque breviter pubescentibus, foliis ovato-lanceolatis dentatis. — in agris umbrosis regni Neapolitani. Habitu accedit ad varietates ramosas et diffusas *Hesperidis matronalis*. DC. syst. 2. p. 452.* (v. s.)

8. *H. STEVENIANA* (DC. syst. 2. p. 452.*) pedicellis calyce longioribus, petalorum laminis obovatis, caule siliquisque hispidis, foliis radicalibus petiolatis runcinatis, superioribus ovato-lanceolatis inciso-dentatis. — in Tauriâ merid. Petalorum ungues calyce longiores, laminæ obovatæ purpurascens. Siliquæ hispidae (v. s. comm. à cl. Stev.)

9. *H. AFRICA* (Poir. suppl. 3. p. 194.) pedicellis glanduloso-pilosis calycis longitudine, petalorum laminis obovatis, foliis oblongis obtusis cauleque simplici ciliato-hispidis. \mathcal{L} ② in campis apricis Sibiriae et Dahuriæ. *Cheiranthus hirtus* Schlecht. ex Stev. obs. in h. Willd. — DC. syst. 2. p. 453.* Deless. ic. sel. 2. t. 62. (v. s.)

β. foliis grossè dentatis. An species propria?

10. *H. BICUSPIDATA* (Poir. suppl. 3. p. 194.) pedicellis calycis longitudine, petalorum laminis obtusis, foliis lanceolatis acutis subdentatis cauleque simplici pube stellatâ pubescentibus. \mathcal{L} ② in Armeniâ. *Cheiranthus bicuspидatus* Willd. 3. p. 519.* Tota pubescens subcinerea. — DC. syst. 2. p. 453.* (v. s. in h. Mus. Par. et Banks.)

β. Cappadocica, paulò minore, floribus siliquisque magis patulis. in Cappadociâ.

11. *H. RAMOSISSIMA* (Desf! fl. atl. 2. p. 91. t. 161.*) pedicellis calyce subbrevioribus, petalorum laminis elliptico-oblongis, siliquis scabris torosis, foliis oblongis velutinis, caule ramosissimo. ① in arenosis maritimis agri Algeriensis in Ægypto et Galatiâ. DC. syst. 2. p. 455.* (v. s.)

12. *H. PYGMÆA* (Delil! ill. fl. eg. p. 19. n. 596.) pedicellis calyce longioribus, petalorum laminis ellipticis, siliquis tenuissimis glabris, foliis sinuatis pube stellatâ scabris, caule ramoso. ① in Syriâ et prope Alexandriam. Flores parvi purpurascens. DC. syst. 2. p. 455.* (v. s. comm. à cl. Delil.)

13. *H. PULCHRELLA* (DC. syst. 2. p. 455.*) pedicellis calyce longioribus, petalorum laminis obovatis, foliis subradicalibus obtusè sinuato-pinnatifidis glabrisculis, scapis plurimis vix foliis longioribus. ① in Oriente. Flores rosei, minores quàm in Malcomiâ maritimâ sed sat similes. (v. s.)

14. *H. CRENULATA* (DC. syst. 2. p. 456.*) pedicellis calyce brevioribus, petalorum laminis obovatis, foliis oblongis obtusis crenatis glabris, caule ra-

moso. ① in Oriente. Petalorum ungues calycis longitudine, laminæ obovate obtusæ purpurascens. (v. s. in h. Mus. Par.)

†. *Species quoad genericos caracteres subdubiæ.*

15. H² PINNATIFIDA (Michx! fl. bor. am. 2. p. 31.) pedicellis calyce demum longioribus, petalorum laminis obovatis, foliis inferioribus pinnatifido-lyratis, superioribus inæqualiter serratis. ½ in humilibus sylvarum Tennesseæ, in Kentucky. Herba glaberrima. Petalorum laminæ obtusæ integræ pallide purpureæ, parvæ. DC. syst. 2. p. 456.* (v. s. in h. Mus. Par.)
16. H² PENDULA (DC. syst. 2. p. 457.) pedicellis deflexis hispidis, foliis setosis inferioribus sublyratis, superioribus ovatis dentatis, caule ramoso hispido. — in Syriâ. Flores ignoti. (v. s. comm. à cl. Labill.)
17. H² ANGUSTIFOLIA (DC. syst. 2. p. 457.) pedicellis brevissimis crassis, petalorum laminis oblongis, foliis linearibus grossè dentatis subglabris, caule hispido. ① in Oriente inter Alep et Mossul. Flores albi parvi. (v. s. in h. Mus. Par.)

†. *Species non satis notæ.*

18. H. ELATA (Horn. h. hafn. suppl. 74.) ② hab. . . caulis simplex erectus, folia lanceolata acuminata, remotè denticulata. Petala obtusa, integerrima et apice emarginata. (Horn.) DC. syst. 2. p. 457.
19. H. NIVEA (Baumg. fl. trans. 2. p. 278.*) ½ in Transylvaniâ. Hirsutissima. Folia alterna petiolata. Flores corymboso-recemosi. DC. syst. 2. p. 458.
20. H. REPANDA (Lag. cat. h. madr. 1814. p. 20.) ½ in Hispaniâ. Herbacea. Caulis erectus subangulatus. Florum apica terminalis, multiflora, elongata. Calyx atropurpureus. Corolla violacea. DC. syst. 2. p. 458.
- LI. ANDREOSKIA. Hesperidopsis. DC. syst. 2. p. 484. Dontostemon. Andr. in litt.

Calyx suberectus basi æqualis deciduus. Petala unguiculata integra. Stamina 2 lateralia libera edentula, 4 majora, nunc per paria connata, nunc intus ad apicem dente aucta. Siliqua sessilis teretiuscula, valvis subconcavis, stylo brevi gracili, septo membranaceo bilocularis. Semina ovata uniseriata. Cotyledones planiusculæ incumbentes. (o II)

1. A. INTEGRIFOLIA, foliis linearibus integerrimis, ramis pedicellisque glandulosis pilosisque, siliquis glandulosis. ① ② in montibus sabulosisque Sibiriae. Sisymbrium integrifolium Lin! spec. 922. DC. syst. 2. p. 484.* Flores in meis specimenibus albi, variantes purpurei ex Gmelino. (v. s.)
2. A. EGLANDULOSA, foliis linearibus integerrimis cauleque villosis eglandulosis. ① in Sibiria. Sisymbrium eglandulosum DC. syst. 2. p. 485.* S. Sieversianum Redowski ex Fisch! in litt. Differt caule foliisque villosis sed eglandulosis. (v. s. in h. Fisch.)
3. A. PECTINATA, foliis pectinato-pinnatilobatis, lobis linearibus acutis cauleque pilosis et glanduloso-muricatis. ②, Sisymbrium pectinatum DC. syst. 2. p. 485. Flores omnino A. integrifoliæ sed paulò majores.
- a. caule humili, floribus albis. — in Sibiria. Hesperis pinnata Pers! ench. 2. p. 203. (v. s.)
- β. caule elongato, floribus albis. — in Dahuria. Sisymbrium pectinatum Fisch! in litt. (v. s.)
- γ. caule elongato, floribus rubicundis. — in Sibiria. Cheiranthus pinnatifidus Willd. spec. 3. p. 523.* ex Stev. obs. ined. in h. Willd.

LII. SISYMBRIUM. All. ped. 1. p. 274. DC. syst. 2. p. 458. (excl. n. 44-46.)

Siliqua teretiuscula, super torum sessilis. Stigmata 2 subdistincta aut in capitulum connata. Calyx basi æqualis. Semina ovata aut oblonga. Cotyledones planæ, incumbentes, interdum obliquæ. Stam. edentula.

Sect. I. VELARUM. DC. syst. 2. p. 459.

Siliquæ subuliformes basi latiores, apice in stylum brevissimum desinentes, pedicello brevissimo post anthesin incrassato secus axin adpressæ. Flores flavi.

1. *S. OFFICINALE* (Scop fl. car. ed. 2. n. 824.) foliis runcinatis pilosis, caule piloso, siliquis rachi adpressis subulatis. ① in incultis secus muros, sepes et vias Europæ totius et Mauritaniz. Erysimum officinale Lin! spec. 922. Gd. fl. dan. t. 560. Smith engl. bot. t. 725. Flores minimi lutei. Siliquæ erectæ, rachi adpressæ. — DC. syst. 2 p. 459.* (v. v.)

β. *leiocarpum*, siliquis glabris. — in Carolinâ merid. et Teneriffâ. Erysimum officinale. Pursh fl. bor. am. 2. p. 436.

2. *S. COMULATUM* (Cav. præl. n. 3. ex Lag! in litt.) foliis inferioribus sinuato-runcinatis, superioribus lanceolatis dentatis, siliquis compressis subulatis subpilosis, pedicellis crassis brevissimis. — Circa Madritum. Flores ignoti. Siliquæ strictæ, erectæ. Species distinctissima, non satis nota et forsân genus proprium constituens. DC. syst. 2. p. 461.* (v. s.)

Sect. II. NORTA. DC. syst. 2. p. 461.

Siliquæ teretes. Calyx patens. Semina oblonga. Flores flavi. Racemi ebracteati.

3. *S. STRICTISSIMUM* (Lin! spec. 922.) foliis lanceolatis petiolatis dentatis pubescentibus. ½ in montosis asperis Europæ mediæ. Jacq. h. vind. 122. Flores intense flavi. Siliquæ 2 poll. longæ, glaberrimæ, strictæ. DC. syst. 2. p. 461.* (v. v. et s.)

4. *S. JUNCUM* (Rieb. fl. taur. 2. p. 114.*) foliis glabris glaucis, inferioribus petiolatis runcinato-pinnatifidis, superioribus lineari-lanceolatis integris. ½ ② ① in pratis siccis Hungariæ, Transylvaniz, Tauriz, Sibiriz. Siliquæ strictæ erectæ, axi parallelæ. DC. syst. 2. p. 462.* (v. s.)

α. *Brassica polymorpha* Murr. comm. goett. 1776. p. 35. t. 6? Walds. et Kit. pl. rar. hung. 1. p. 93 t. 90.*

β. *Cheiranthus juncus* Walds. et Kit. pl. rar. hung. 3. p. 259. t. 234.*
Erysimum junceum Willd. spec. 3. p. 513.*

Sect. III. PSILOSTYLUM. DC. syst. 2. p. 463.

Siliquæ teretes stylo gracili longo apiculatæ. Calyx clausus. Semina oblonga. Flores flavi.

5. *S. EXACOIDES* (DC. syst. 2. p. 463.*) foliis ovali-oblongis subintegris glabris glaucis, racemis 1-4-floris. ① in Libano. Petala unguibus linearibus calyce longioribus, limbo flavo, elliptico, parvulo, Deless! ic. sel. 2. t. 63. An genus proprium, ex stylo et calyce à Sisymbrio separandum? (v. s.)

Sect. IV. IRIO. DC. syst. 2. p. 463.

Siliquæ teretes. Semina ovata subtriquetra. Flores flavi. — Pedicelli ebracteati.

§. 1. *Foliis dentatis integrise.*

6. *S. HISPANICUM* (Jacq. coll. 1. p. 69.* ic. rar. 1. t. 124.) foliis lanceolatis dentatis sessilibus glabris, caule ramoso divaricato, siliquis erectis teretiusculis glabris. ② in Hispaniâ, in Galliâ merid? Siliquæ erectæ, adpressæ. DC. syst. 2. p. 463.* (v. s.)

7. *S. LINEARE* (DC. syst. 2. p. 464.*) foliis linearibus integris hispidulis, caule hispido, siliquis glabris tenuissimis. — in Sibiria. S. n. 38. var. floribus luteis Gmel. fl. sib. 3. p. 271. t. 62. ex Fisch! in litt. Flores flavi parvi. Siliquæ tennes, teretiusculæ, glabræ. Semina minima. (v. s. ex h. Fisch.)

8. *S. PUMILUM* (Steph. in Willd. spec. 3. p. 507.*) foliis radicalibus sinuato-dentatis, caulinis sagittato-amplexicantibus dentatis, siliquisque pube ramosâ scabris. ① in Persiâ bor. et Iberiâ. Planta nunc digiti minimi longitudine, nunc pedalis imò sesquipedalis! DC. syst. 2. p. 464.* (v. a.)

§. 2. *Foliis pinnatim lobatis, lobis integris dentatisve.*

9. *S. OBTUSANGULUM* (Schleich! cat. p. 48.) foliis pinnatipartitis, lobis ovali-oblongis obtusis sinuato-dentatis, sinubus rotundatis, caule basi retrorsum subhispido. ① ② in arenosis cultis et ruderatis Hispaniæ, Galliæ, Helvetiæ. *Eruca inodora*. J. Bauh. hist. 2. p. 862. f. 3. — DC. syst. 2. p. 465.* (v. s.)

ℓ. caule basi glabriusculo. *S. lævigatum* Willd! spec. 3. p. 500.

10. *S. ACUTANGULUM* (DC. fl. fr. 4. p. 670.*) caule foliisque glabris, radicalibus runcinatis, caulinis pinnatifidis, lobis sinubusque acutis, calyce patentissimo, siliquis scabris. ② in Pyrenæis, Alpibus Delphinensibus, et Sabaudis, etc. *S. Pyrenaicum*. vill. fl. dauph. 3. p. 341. t. 38. non Lin. *Sinapis Pyrenaica*. Lin! spec. 934. Siliquæ graciles pollicem longæ. DC. syst. 2. p. 465.* (v. v.)

- * 11. *S. TARAXACIFOLIUM* (DC. fl. fr. 4. p. 670.* ic. rar. p. 11. t. 37.) foliis runcinato-pinnati-lobatis, lobis sinubusque acutis, siliquis declinatis glabris, calyce erectiusculo. ② in asperis montium Sedenensium Gallo-provinciæ. Flores flavi parvuli petalis tamen calyce longioribus. Siliquæ sæpius declinatæ. DC. syst. 2. p. 466.* (v. s. et v.)

ℓ? *S. contortum* Cav. ex Willd. enum. 678.* Differt caule basi multò magis hispido, foliis radicalibus hispidis, caulinis minùs strictè erectis.

γ. *S. affine* Willd! enum. suppl. 44. Calyx magis patens; folia hujus et calyx sequentis.

12. *S. AUSTRIACUM* (Jacq. fl. austr. 3. t. 262.) caule siliquis foliisque glabris, radicalibus runcinatis, caulinis incisim pinnatifidisve, lobis sinubusque acutis, calyce patente. ② in asperis et ruderatis collium et vallium Galliæ austr., Pedemontii, Helvetiæ, Germaniæ. DC. syst. 2. p. 466.* (v. s.)

ℓ. *S. Eckartsbergense* Willd. spec. 3. p. 502. Siliquis deflexo-patulis.

γ. *S. erysimifolium*. Poirr! acs. toul. 3. p. 329. Siliquis erectis.

δ. *S. Tiliæri* Bellard! ined. Willd. spec. 3. p. 497.* Forsan ab omnibus diversa siliquis longioribus et foliis radicalibus confertioribus, caulinis paucioribus.

13. *S. ISEO* (Lin! amœn. 4. p. 270.*) caule foliisque glabris runcinato-pinnatipartitis, lobis dentatis, terminali elongato, calyce siliquisque patentius erectis. ① in ruderatis, secùs muros et vias Europæ. Jacq. fl. austr. t. 322. Smith engl. bot. t. 1631. *S. glabrum* Willd! enum. suppl. 44. DC. syst. 2. p. 467.* (v. v.)

ℓ. lobo extimo subrotundiore. *S. glabrum* h. vind. ex Schrad! spec. sicc. distinguitur foliis inferioribus sublyratis.

γ. pedicellis brevioribus. *S. pinnatifidum* Forsk! fl. arab. cxvi. Differt siliquis pedicellisque dimidiò brevioribus.

δ. Caule basi pubescente. *S. gallicum* Willd! enum. 678.

14. *S. NITIDUM* (Zea in Desf! cat. hort. par. 1815. p. 153.) glabrum, foliis sinuato-hastatis basi in petiolum decurrentibus, siliquis filiformibus patentibus. ① hab. . . . Flores parvi, flavi nec albi ut ex sicco dictum fuit. Semina parva, pallida, ovalia, compressa. (DC. syst. 2. p. 468.* (v. v. in h. Par.)

15. *S. SUBHASTATUM* (Willd. enum. 679.) glabrum subglaucescens, foliis radicalibus runcinatis, caulinis lanceolatis integris basi auriculis acutis hastatis, siliquis patentibus rigidis, pedicellis crassis brevissimis. ① in insulis Archipelagi. *Brassica subhastata* Willd. spec. 3. p. 550.* — DC. syst. 2. p. 468.* (v. v. in h. Monsp.)

16. *S. COLUMNÆ* (Jacq. fl. austr. t. 323.) caule villosa subincano, foliis runcinatis pubescentibus, lobis dentatis integrisve acutis, siliquis erectiusculis, calyce laxo. ① in ruderatis Europæ austr. et orient. DC. syst. 2. p. 469.* (v. v.)

α. *altissimum*, siliquis glabris, foliorum lobis subintegris. *S. altissimum*.

Liu! spec. 920. — Buxb. cent. 5. p. 26. t. 51.

- β. *leiocarpum*, siliquis glabris, foliorum lobis dentatis. S. Columnæ Jacq. fl. aust. t. 323.
- γ. *villosissimum*, siliquis pubescentibus, caule foliisque villosissimis, foliorum lobis dentatis. S. Loeselii. Thuil. fl. par. ed. 2. v. 1. p. 335.
- δ. *tenuisiliquum*, siliquis pubescentibus tenuissimis, caule petiolisque pubescentibus, foliorum lobis subintegris. S. Columnæ Bieb. fl. taur. 2. p. 114. var. α.
- ε. *Orientale*, siliquis puberulis, herbæ parte inferiore villoso-tomentosâ, superiore glabrâ, foliorum lobis dentatis. S. Orientale Lin. amœn. 4. p. 322. Spec. 921.
17. S. PANNONICUM (Jacq. coll. 1. p. 70.* ic. rar. 1. t. 123.) foliis inferioribus runcinatis hispidis lobis dentatis superioribus pinnatisectis glabris, lobis linearibus integerrimis, siliquis patentibus. ① in sterilibus, præruptis campis et vineis Europæ. S. altissimum. Pall. ind. taur. Habl. taur. p. 158. ex Bieb. Petala pallidè flava, ferè ochroleuca. Siliquæ patulæ. — DC. syst. 2. p. 470.* (v. s.)
18. S. SEPTULATUM (DC. syst. 2. p. 471.*) foliis glabris pinnatipartitis, lobis linearibus anticè subdentatis acutis, siliquis subpatulis, septo in loculis semina impressa gerente. — in Syriâ. Siliquæ pedicello continuæ, teretes, strictæ. An genus proprium? (v. s. in h. Banks.)
19. S. CARTILAGINEUM (Pall. herb. ex Fisch! in litt.) foliis scabris pinnatipartitis, lobis linearibus integris subdentatisve crassiusculis, summis indivisis. ② in montibus cretaceis Tauriæ et Iberiæ. Siliquæ elongatæ basi attenuatæ. Sem. ign. An Dyploxaxis, an Stanleyæ species? DC. syst. 2. p. 471.* (v. s.)
20. S. LYRATUM (Burm. fl. cap. 17.) foliis inferioribus pilosis lyrato-runcinatis lobis dentatis, superioribus oblongis glabris dentatis, siliquis subpatulis glabris. ② ad cap. Bonæ-Spei. S. sylvestre Burm! herb. Semina ovato-compressa, parva, rufa. — DC. syst. 2. p. 471.* Deless! ic. sel. 2. t. 64. (v. s. in h. Deless.)
21. S. BURCHELLII (DC. syst. 2. p. 472.*) foliis pinnatisectis pilosis, lobis oblongis obtusè sinuato-angulatis, siliquis patulis scabris, pedicellis brevibus crassis, caule pilis patulis hispido. ① ad cap. Bonæ-Spei. S. n. 1496 Burch! cat. geogr. pl. afr. austr. Siliquæ ferè S. asperi. (v. s. comm. à cl. Burchell.)
22. S. GARIEPINUM (Burch! in DC. syst. 2. p. 472.) foliis pinnatipartitis, lobis dentatis acutiusculis, pube ramosâ scabris, siliquis scabris erectiusculis, caule adpressè pubescente. ① ad caput Bonæ-Spei trans flumen Gariep. Flores parvi, flavi. Siliquæ teretes pube parvâ stellatâ scabræ. (v. s.)
- α. *apricum*, Burch! cat. geogr. n. 2080.
- β. *nemorosum*, Burch! cat. geogr. n. 2558.
23. S. ASPERUM (Lin. spec. 920. ex syn. non ex herb.) foliis glabris pinnatipartitis, lobis oblongis obtusis subdentatis, pedicellis brevissimis, siliquis muricato-scabris stylo brevi acuminatis. ② in Galliâ austr. Occitaniâ, Hispaniâ. — J. Bauh. hist. 2. p. 858. f. 3. — DC. syst. 2. p. 473.* (v. v.)
- §. 3. *Foliis caulinis bipinnatisectis, lobis incisissimis pinnatifidis multifidisve.*
24. S. SOPHIA (Lin! spec. 922.) foliis bipinnatisectis, lobis oblongo-linearibus incisissimis, pedicellis calyce quadruplò longioribus, petalis calyce minoribus. ① in Europæ maceris. Smith engl. bot. t. 963. S. parviflorum Lam. fl. fr. 2. p. 519. Flores parvi, flavi, subcorymbosi. Siliqua erecta, linearis, glabra. DC. syst. 2. p. 474.* (v. v.)
25. S. PERSICUM (Spreng! nov. prov. p. 39. n. 88.*) foliis bipinnatisectis, lobulis linearibus, pedicellis calyce duplò longioribus, petalis calycem paulò superantibus. ① in Persiâ. Flores flavi, minimi. Siliquæ axi ferè parallelæ. DC. syst. 2. p. 475.* (v. s.)

26. *S. CANESCENS* (Nutt. gen. am. 2. p. 68.*) foliis bipinnatisectis canescentibus, lobulis obtusis dentatis, petalis calyci æqualibus, siliquis clavatis pedicello brevioribus. ① in Americâ bor. à Virginia ad Georgiam. *S. Sophia* Pursh fl. bor. am. 2. p. 440? ex Nutt. Petala parva, obovata, pallide flava, calycis longitudine. DC. syst. 2. p. 475.*
27. *S. BRACHYCARPUM* (Richards. in Frankl. narr. Journ. p. 744.) foliis bipinnatisectis, lobis obtusis integris vel parciter incisis, petalis calyce majoribus, siliquis linearibus subtetragonis pedicello brevioribus. — in regione sylvaticâ Arcticâ inter lat. bor. gradus 54 et 64. Facies *S. Sophiæ*, sed siliquæ dimidiò breviores.
28. *S. TRIPINNATUM* (DC. syst. 2. p. 475.*) foliis tripinnatisectis pube stellatâ velutinis, lobulis oblongo-linearibus subdentatis, siliquis gracilibus elongatis glabris. ① ad Cap. Bonæ-Spei in siccis ad ripas fluminis Gaurits. *Sinapis?* tripinnata. Burch! cat. geogr. pl. afr. austr. n. 1640. voy. 1. p. 318. Medium inter *S. Sophiam* et *Millefolium*. (v. s. in h. Burch.)
29. *S. MILLEFOLIUM* (Ait! h. kew. ed. 1. v. 2. p. 391. ed. 2. v. 4. p. 114.) foliis subtripinnatisectis incanis, lobulis obtusis minimis, caule suffruticoso, petalis calyce majoribus. ⑤ in rupestribus Teneriffæ. *Sinapis millefolia* Jacq. coll. 1. p. 41.* ic. rar. t. 27. Flores flavi, conferto-corymbosi. DC. syst. 2. p. 476.* (v. s. et v.)
30. *S. TANACETIFOLIUM* (Lin! spec. 916.) foliis pinnatisectis, segmentis lanceolatis inciso-serratis, extimis confluentibus, petalis calyce majoribus, siliquis pedicello brevioribus. ④ in apricis frigidis Helvetiæ, Pedemontii, Sabaudiz, Delphinatûs, Gallo-provinciæ et in Pyrenæis. — Moris. hist. 2. p. 231. s. 3. t. 6. f. 19. Flores lutei. Semina parva. (v. v.)
31. *S. MYRIOPHYLLUM* (H. B. et K. in DC. syst. 2. p. 477.*) foliis bipinnatisectis, segmentis obtusis, petalis calyce majoribus, siliquis lanceolatis. Willd. mss. — in regno Quiteni ad radicem mont. Cotopari. Flores pedicellati, erecti, flavi. Semina minuta ovata. (Deest in herb. Humb.)

Sect. v. KIBERA. DC. syst. 2. p. 477.*

Pedicelli basi bracteati. Stylus brevis crassus retusus. Flores parvi, flavi aut albi.

32. *S. PERUVIANUM* (DC. syst. 2. p. 477.*) pedicellis axillaribus glabris calyce longioribus, siliquis hispidis patulis, foliis oblongis inciso-serratis. — in Peruvia. Flores parvi. Semina minima. (v. s. in h. Lamb.)
33. *S. SUPINUM* (Lin! spec. 917.) pedicellis axillaribus brevissimis solitariis, siliquis erectis puberulis, foliis sinuato-pinnatifidis, caule retrorsum pubescente. ① in arenosis humidis circa Parisios, Lugduni ad Rhodanum, ad lacum Jovis in Jurasso, in Hispaniâ et Gothlandiâ. *Arabis supina* Lam. fl. fr. 2. p. 512. Isu. act. ac. par. 1724. t. 18. — DC. syst. 2. p. 477.* (v. s.)
34. *S. RUNCINATUM* (Lag! fl. hisp. ined. DC. syst. 2. p. 478.*) pedicellis axillaribus brevissimis solitariis, siliquis incurvis canleque glabris, foliis oblongis runcinato-dentatis. ① in ruderatis Hispaniæ. Folia floralia sessilia. Flores minimi. (v. s.)
35. *S. HIRsutum* (Lagasc. ex Duf! in litt. DC. syst. 2. p. 478.) pedicellis axillaribus brevissimis solitariis, bracteis oblongo-linearibus subintegris, siliquis erectis canleque pubescentibus. ① hab. Madriti. *A. S. runcinato* differt caule siliquisque non glabris sed puberulis. (v. s.)
36. *S. POLYCRERATUM* (Lin! spec. 3. p. 918.) pedicellis subternis axillaribus brevissimis, siliquis erectis glabris, foliis sinuato-runcinatis, lobis acutis dentatis, inferioribus majoribus. ① in muris humidis et rupium fissuris Europæ australis. Jacq. h. vind. t. 79. Flores parvi, flavi, ad axillas foliorum sessiles, subterni. DC. syst. 2. p. 479. (v. v.)
37. *S. RIGIDUM* (Bieb. suppl. p. 439.) pedicellis brevissimis axillaribus aut nudis, siliquis caulibusque erectis hispidis, foliis glabriusculis oblongis acutè

runcinato-pectinatis. ① in Tauriâ merid. *Erysimum polyceratum* Pall. itin. ed. gall. 8^o app. p. 346. n. 351. t. 107. — DC. syst. 2. p. 480.* (v. s.)

Sect. VI. ARABIDOPSIS. DC. syst. 2. p. 480.

Siliquæ lineares compressæ stigmatē sessili truncato. Florēs albi brevissimæ pedicellati. Pedicelli ebracteati.

38. *S. BURSIFOLIUM* (Lin! amœn. 4. p. 322.*) foliis lyratis pinnatifidis glabris, caule erecto folioso, pedicellis crassis calyce brevioribus. ① in Sicilia. *Arabis bursifolia* Lam. fl. fr. 2. p. 511. — Dill. hort. elth. 179. t. 148. f. 177. — DC. syst. 2. p. 480. (v. v. et s.)

39. *S. PINNATIFIDUM* (DC. fl. fr. 4. p. 667.* syst. 2. p. 481.* non Forsk.) foliis radicalibus lyratis, caulinis pinnatipartitis, lobis linearibus integris, terminali majore, pedicellis gracilibus calyce ferè brevioribus. ② in pascuis montium Europæ. *Arabis pinnatifida* Lam! dict. 1. p. 221. ill. t. 563. f. 3. *S. dentatum* All. ped. t. 57. f. 3. (v. v.)

40. *S. KAYSIMOIDES* (Desf! fl. atl. 2. p. 84. t. 158.*) foliis lyrato-pinnatifidis, lobis inæqualiter dentatis, terminali majore, pedicellis brevissimis, siliquis rectangulo-patentibus. ① in arenosis propè Kervan, insulâ Teneriffæ et Hispaniâ. *S. rigidulum* Lag! gen. et sp. p. 20. — DC. syst. 2. p. 482.* (v. s.)

41. *S. RAMULOSUM* (Delil! eg. ill. 19.) foliis inferioribus pinnatilobatis, lobis paucis acutis integris, terminali oblongo maximo, caulinis oblongo-linearibus subintegris, pedicellis calyce longioribus. ① in Ægypto. Flores parvi, non satis noti. DC. syst. 2. p. 482.* (v. s. in h. Mus. Par.)

42. *S?* *CINEREUM* (Desf! fl. atl. 2. p. 83. t. 157.*) foliis pubescentibus subcarnosius pinnatipartitis, lobis lineari-filiformibus integris, pedicellis calyce longioribus. ① in arenosis Mauritanix. Willd. spec. 3. p. 501. Petala dilutè violacea calyce duplò longiora. Semina minima, quoad structuram ignota et ideo genus dubium? An potiùs *Hesperis* aut *Arabis*? DC. syst. 2. p. 483.* (v. s.)

43. *S. TORULOSUM* (Desf! fl. atl. 2. p. 84. t. 159.*) foliis oblongis, radicalibus subpinnatifidis, caulinis grossè dentatis, pedicellis brevissimis, siliquis hispidis rectis. ① in regno Tauretano et in insulâ Cypri. Sibth. et Smith fl. græc. t. 632. ex prod. 2. p. 20. DC. syst. 2. p. 483.*

44. *S. CONTORTUPLICATUM* (DC. syst. 2. p. 483.*) foliis oblongis, radicalibus pinnatifidis, caulinis dentatis aut integris, pedicellis calycis longitudine, siliquis hispidis sæpiùs contortuplicatis. ① in deserto Cumano, etiam circa Kisjar et Astrakan. Deless! ic. sel. 2. t. 63. *Cheiranthus contortuplicatus* Steph. in Willd. spec. 3. p. 521.* Flores ex albo purpurascences.

β. *rectisiliquum*, (Fisch! in litt.) siliquis rectis aut vix curvis.

† *Species quoad caracteres genericos dubiæ.*

45. *S?* *ALBUM* (Pall. itin. 3. ed. gall. 8^o v. 8. app. n. 349. p. 344. t. 96. f. 1.*) foliis albidis pubescentibus pinnatipartitis, lobis oblongis obtusiusculis cauleque eglandulosis. ② in Sibirix frigidis. Stamina omnia edentula. An Nasturtium ob siliquas breves? An *Erysimum* ob siliquas quadrangulas dictas? An genus proprium cum Nasturtio sagittato? DC. syst. 2. p. 486.* (v. s.)

46. *S?* *NANUM* (DC. syst. 2. p. 486.*) foliis oblongis sinuato-pinnatifidis pubescentibus stellatè velutinis, siliquis torulosis. ① in Sibiria orient. Merk. Petala oblonga, purpurascens. Semina non satis nota. (v. s. comm. à cl. Steven.)

†† *Sisymbria non satis nota.*

47. *S. SINAPIS* (Burm. fl. ind. 140. excl. syn. Barr. ad Arabidem thalianam ref.) — in Javâ. Statura et facies *Sinapis arvensis*, at flores minutissimi albi. Folia sublyrata, denticulata. DC. syst. 2. p. 487.

48. *S. PATENS* (Moench meth. 251.) ① hab. Folia runcinata laciniâ extremâ sagittatâ, suprema lanceolata sagittata. Petala pallidè flava, integra. Siliqua teres, lævis, bifariè pilosa. An *Brassica*? DC. syst. 2. p. 487.

49. *S. CAPENSE* (Thunb. prod. 109.) ad cap. Bonæ Spei. Caulis paniculatus glaber. Folia pinnatifida, glabra. Siliquæ lineares, læves. DC. syst. 2. p. 487.
50. *S. serratum* (Thunb. prod. 109.) ad cap. Bonæ Spei. Caulis subtrigonus. Folia elliptica, argutè serrata, glabra. DC. syst. 2. p. 487.
51. *S. crassifolium* (Cav. proel. p. 437. n. 977. Lag. h. madr. ined. t. 24.) 2 in Hispaniâ. Folia radicalia sinuato-runcinata, subcarnosa, summa linearia integerrima. Spicæ apice nutantes. Corolla pallidè sulphurea. An *Dyplo-taxis*? DC. syst. 2. p. 488.
52. *S. fugax* (Lag. elench. h. madr. 1805 et 1815. p. 20.) 2 in Hispaniâ. Glabrum, læve. Folia lyrata, lobis lateralibus terminalique oblongo-lanceolatis, acutis. Siliquæ filiformes tandem contortæ. DC. syst. 2. p. 488.
53. *S. leptopetalum*. (Rafin. fl. lud. p. 84. n. 268.) — Secus aquas in Loni-sianâ. Folia pinnatisecta, segmentis lobatis glabris, impari majoribus. Flores minimi, flavidi. Petala calyce staminibusque breviora. Stigma sessile. Sili-quæ longæ, teretes.

LIII. *ALLIARIA*. *Adans. fam. 2. p. 418. DC. syst. 2. p. 488.*

Siliqua teretiuscula nervis prominulis subtetragona. Calyx laxus. Semina subcylindrica. Cotyledones lineari-oblongæ, planæ.

1. *A. officinalis* (Andrzejowski cruc. ex Bieb. fl. taur. suppl. p. 445.) foliis cordatis, siliquis prismaticis pedicello multoties longioribus. 2 ad æpes, dumeta et fossas Europæ totius et in Persiâ. *Erysimum Alliaria* Lin! spec. 922. OEd. fl. dan. t. 935. Smith engl. bot. t. 796. — DC. syst. 2. p. 489. (v. v.)

β. foliis altius incis. Mich. hort. flor. p. 49. n. 4.

2. *A. brachycarpa* (Bieb. fl. taur. suppl. 445.) foliis ovato-orbiculatis, siliquis lanceolatis longitudine pedicelli. 2 in Iberiâ. *Raphanus rotundifolius* Bieb. fl. taur. 2. p. 130.* Flores albi. An prioris verè congener?

LIV. *ERYSIMUM*. *Gaertn. fruct. 2. p. 297. t. 143. DC. syst. 2. p. 490.*

Siliqua tetragona. Calyx clausus. Cotyledones planæ, oblongæ.

Sect. 1. *STYLONEMA*. *DC. syst. 2. p. 491.*

Stylus filiformis, longus. Stigmata 2 patentia. Calyx post anthesin subper-sistens. Flores ferè sessiles.

1. *E. siliculosum* (DC. syst. 2. p. 491.*) siliquis stylo brevioribus, junioribus calyce persistente tectis, floribus breviter pedicellatis, foliis linearibus integris. (2) in arenâ mobili deserti Cumani Tauriæ. *Cheiranthus siliculosus* Bieb. fl. taur. 2. p. 121.* suppl. 443. Species ad *E. sessiliflorum* valdè accedens et cum eâ primum confusa. Deless! ic. sel. 2. t. 65. (v. s.)
2. *E. siculum* (Spreng. neu. entd. 3. p. 51.) siliquis stylo brevioribus prismaticis incanis, etiam adultis calyce persistente tectis, floribus breviter pedicellatis, foliis linearibus glabriusculis. (2) in Sicilia.
3. *E. sessiliflorum* (Brown in h. kew. ed. 2. v. 4. p. 116.) siliquis styli longitudine, junioribus calyce persistente tectis, floribus sessilibus, foliis linearibus integris. (2) 2 in desertis salsis Sibirici, in arenâ mobili Tauriæ et montosis Caucasii. *Cheiranthus quadrangulus* L'her! stirp. 1. p. 91. t. 44. Flores sulphurei, odori. — DC. syst. 2. p. 492.* (v. s.)
4. *E. angustifolium* (Ehrh. beit. 7. p. 155.) siliquis stylo multò longioribus junioribus calyce subpersistente instructis, floribus subsessilibus, foliis linearibus integris. (2) in arenosis siccis planitiarum Hungariæ et Transyl-vaniæ. Walds. et Kit. pl. rar. hung. 1. p. 101. t. 98.* *Cheiranthus virgatus* Poir! suppl. 2. p. 781.* Flores similes *E. sessilifloro*, sed paulò minores. — DC. syst. 2. p. 492.* (v. s.)

Sect. II. CUSPIDARIA. DC. syst. 2. p. 493.

Stylus filiformis brevis. Siliqua tetragono-anceps. Calyx cum petalis deciduus. Flores distinctè sed breviter pedicellati.

5. *E. CUSPIDATUM* (DC. syst. 2. p. 493.*) siliquis stylo triplò longioribus ancipitibus nudis, floribus breviter pedicellatis, foliis oblongo-lanceolatis sinuato dentatis. ② in Bithyniâ, Moldaviâ, Tauriâ, Caucaso et Iberiâ. *Cheiranthus cuspidatus* Bieb. fl. taur. 2. p. 120. Buxb. cent. 2. t. 33. f. 1. Petala flava. Semina ovata, apice utroque obliquè truncata. (v. s.)
6. *E. RUPESIÆ* (DC. syst. 2. p. 494.*) foliis subdentatis, radicalibus spathulatis, canlinis oblongis pube 2-3 partitâ pubescentibus, caule suffruticoso styli longitudine latitudinem siliquæ excedente. ④ ⑤ in Olympi Bithyni cacumine. *Cheiranthus rupestris* Sibth. et Smith fl. græc. t. 633. ex prod. 2. p. 23. Flores lutei, pauci, magnitudine *E. cuspidati*. An *Cuspidariæ* species? (v. s. in h. Banks.)

Sect. III. ERYSIMASTRUM. DC. syst. 2. p. 494.

Stylus brevis aut vix ullus. Siliqua tetragona. Calyx deciduus. Folia nec cordata nec amplexicaulia. Flores distinctè pedicellati.

7. *E. LEPTOSTYLUM* (DC. syst. 2. p. 494.*) foliis lanceolatis remotè subdentatis pube 3-4 partitâ pubescentibus, ramis paucis rectiuscolis, petalorum laminis suborbiculatis, siliquis erectis, stylo filiformi longitudinem siliquæ duplò superante. ② in campis elatis ad Tanaim inferiore passim, circa Charkoviam et in Caucaso. *E. grandiflorum* Bieb. fl. taur. 2. p. 117. (excl. syn.) suppl. p. 441. non Desf. Petala limbo luteo, orbiculato. (v. s.)
8. *E. EXALTATUM* (Andrz. in Bess. enum. cont. 2. n. 1554.) canescenti-scabrum, foliis lineari-lanceolatis utrinquè attenuatis patentibus remotè denticulatis, siliquis erecto-patulis, glandulis externis bicornibus. — in Podoliâ australi. Affine *E. leptostylo* ex cl. auct.
9. *E. SUFFRUTICOSUM* (Spreng! nov. prov. p. 17. n. 36.*) caule basi suffruticoso, foliis lanceolatis integris pube bipartitâ pubescentibus, siliquis erectis, styli longitudine latitudinem paulò excedente. ③ hab. . . . *Cheiranthus fruticosus* l'Her! herb. Flores nascentes corymbosi, quadruplò minores quàm *Cheiri*, inodori, pallidè flavi, ferè citrini. DC. syst. 2. p. 495.* (v. s.)
10. *E. STRICTUM* (Gaertn. fl. wetter. 2. p. 451. n. 836.) foliis lanceolatis dentatis pube tripartitâ pubescentibus, caule stricto, siliquis erectis, stylo brevissimo, stigmate bilobo. ② in Volhyniâ, Transylvaniâ, Austriâ, Wetteraviâ, etc. *E. hieracifolium* Jacq. fl. austr. t. 73.* — DC. syst. 2. p. 495.* (v. v.)
β. odoratum, caule ramoso, siliquis laxis. *E. odoratum* Ehrh. beit. 7. p. 157. DC. syst. 2. p. 496.
11. *E. VIRGATUM* (Roth cat. bot. 75 *) foliis oblongo-lanceolatis subdentatis pube tripartitâ pubescentibus, caule stricto tereti, siliquis erectis, styli longitudine latitudinem siliquæ subexcedente. ② in lapidosis glareosis Jurassi Neocomensis, in arenosis Moguntizæ, etc. *E. hieracifolium* Ed. fl. dan. t. 923? DC. syst. 2. p. 496. (v. s.)
β? *Cheiranthus firmus* Willd. enum. suppl. 45.
12. *E. LONGISILIQUOSUM* (Willd. enum. 680.*) foliis oblongo-lanceolatis integris pube tripartitâ subpubescentibus, caule stricto tereti, siliquis erectis stylo brevissimo superatis. ② in glareosis Vallesizæ et Genevæ. *E. virgatum* Schleich! pl. exsic. DC. ic. gall. rar. t. 36. Petalorum unguis calycis longitudine. — DC. syst. 2. p. 496.* (v. s.)
13. *E. CÆSPITOSUM* (DC. syst. 2. p. 497.*) foliis sublinearibus integerrimis pube bipartitâ pubescentibus, caulibus cæspitosi suffruticulosi, siliquis erectis, styli longitudine latitudinem siliquæ duplò superante. ④ ⑤ in Persiâ. Petalorum laminæ obovatæ flavæ. (v. s. comm. à cl. Fisch.)

14. *E. HIERACIFOLIUM* (Lin! spec. 923.) foliis lanceolatis basi attenuatis, notè argutè sinuato-dentatis, caule stricto subramoso tereti, siliquis erectis. ② in montibus lapidosis Hercyniæ, in Sponhemia, Sueciâ, circa Mosquam, in Hæmo monte. — J. Bauh. hist. 2. p. 873. f. 2. Flores inodori. Aff. hic *E. repando*, illinc *E. odorato*. — DC. syst. 2. p. 497.* (v. s.)

ℓ? *patulum*, siliquis patentibus.

15. *E. AUREUM* (Bieb. fl. taur. 2. p. 117.* suppl. p. 440.) foliis lanceolatis acuminatis dentatis pubescentibus viridibus, ramis siliquisque patentibus, stigmate crasso bilobo subpedicellato. ② in dumetis promontorii Caucasic. Flores aurei, subodori. DC. syst. 2. p. 498.* Deless! ic. sel. 2. t. 66. (v. s.)

16. *E. IBERICUM* (DC. syst. 2. p. 498.*) foliis inferioribus runcinatis dentatis, superioribus lanceolatis indivisis ramis floriferis siliquisque compresso-tetragonis erecto-patulis. ② in Caucaso, in montibus Kubensibus et Armenis. *Cheiranthus Armeniacus* Sims bot. mag. t. 835. (v. s.)

17. *E. CHEIRANTHOIDES* (Lin! spec. 923.) foliis lanceolatis subdenticulatis scabriusculis viridibus, siliquis erecto-patulis pedicello duplò longioribus, stigmate parvo subsessili. ① in ruderatis Europæ totius, in Virginia et Louisiana. Jacq. fl. austr. t. 23.* *Cheiranthus torritoïdes*. Lam. dict. 2. p. 716.* Flores lutei, parvi. — DC. syst. 2. p. 498.* (v. v.)

ℓ. *Cheiranthus scapigerus* Willd. prod. n. 663. t. 5. f. 10.*

γ. *Cheiranthus aquaticus* Lejeun! fl. spa. 2. p. 68.*

18. *E. REPANDUM* (Lin! amœn. 3. p. 415.) foliis linearibus lanceolatis repando-dentatis subpubescentibus, siliquis patulis pedicello brevi vix crassioribus torulosis, stigmate sessili. ① in Europâ temperatâ. Jacq! fl. austr. t. 22.* Flores flavi. Semina rufa, oblonga. DC. syst. 2. p. 500.* (v. v. et s.)

ℓ. *simplex*, caule simplicis. Mich. hort. flor. p. 49. n. 9.

19. *E? TENELLUM* (DC. syst. 2. p. 500.*) foliis linearibus repando-dentatis, caule subsimplici, siliquis patulis? teretiusculis, stigmate latè bilobo pedicellato. ① in Oriente. Valdè accedit ad *E. repandi* varietatem ℓ. (v. s. in h. Mus. Par.)

20. *E. HELVETICUM* (DC. fl. fr. 4. p. 658.*) foliis linearibus integris, cauleque erecto vix ramoso, pube adpressâ bipartitâ cinereis, siliquis erectiusculis, stigmate pedicellato emarginato. ② in siccis Vallesis, et in Carpathis. *Cheiranthus Helveticus* Jacq. hort. vind. t. 9.* — DC. syst. 2. p. 501.* (v. s.)

21. *E. CANESCENS* (Roth cat. bot. 1. p. 76.*) foliis linearibus integris aut subdentatis, pube bipartitâ subcanescentibus, unguibus calyce longioribus, laminis obovato-oblongis, siliquis erectis pedicello quintuplò longioribus, stigmate subsessili. ② in collibus Europæ austr. *Cheiranthus Alpinus* Jacq. fl. austr. 1. p. 48. t. 75.* — DC. syst. 2. p. 501.*

22. *E. ANDRZEJOSKIANUM* (Bess! in litt. et DC. syst. 2. p. 502.*) foliis linearibus canaliculatis subdentatis pube bipartitâ subcanescentibus, unguibus calyce paulò longioribus, laminis obovatis, siliquis erectis pedicello duplò longioribus, stigmate subsessili. ② in Tauriæ et Caucasi campis siccis. *E. diffusum* Bieb. fl. taur. 2. p. 116. ex Stev! in litt. (v. s.)

23. *E. COLLINUM* (Andrz. in litt.) foliis inferioribus petiolatis spathulatis angulato-dentatis, superioribus lineari-lanceolatis integris, pube tripartitâ, caule erecto subramoso, siliquis scabris erectis. ② propè Cosaccorum oppidulum Galugai. *Cheiranthus collinus* Bieb! fl. taur. 2. p. 119.* DC. syst. 2. p. 182.* (v. s.)

24. *E. LEPTOPHYLLUM* (Andrz. in litt.) foliis omnibus lineari-lanceolatis integerrimis incanis scabris, caule ramoso siliquisque incanis patulis. ② in Iberiâ. *Cheiranthus leptophyllum* Bieb. fl. taur. 2. p. 119.* suppl. p. 44a. DC. syst. 2. p. 182.* (v. s.)

25. *E. VERSICOLOR* (Andrz. in litt.) foliis incanis, inferioribus runcinatis, superioribus linearibus integerrimis, ramis patulis, siliquis incanis strictis. ②

in Persiâ bor., in Caucaso et in deserto Cumano. *Cheiranthus versicolor* Bieb. fl. taur. 2. p. 119.* suppl. p. 442. DC. syst. 2. p. 182.* (v. s.)

26. *E. LANCEOLATUM* (Brown in h. kew. ed. 2. v. 4. p. 116.) foliis inferioribus lanceolatis dentatis, superioribus sublinearibus integris, unguibus calyce longioribus, laminis orbiculato-obovatis, siliquis erectis, stigmatibus sessilibus. ② in apricis muris Europæ ferè totius. DC. syst. 2. p. 502.* (v. v.)

α. *majus*, caule simplici aut ramoso pedali rigido. *Cheiranthus erysimoides* Lin. spec. 923. Jacq. fl. austr. t. 74.*

β. *minus*, caule simplici, interdum multicauli subadscendente. *E. ochroleucum*, β. DC. fl. fr. 4. p. 658. — All. specim. p. 44. t. g. f. 3.

27. *E. RHÆTICUM* (DC. syst. 2. p. 503.*) foliis lineari-lanceolatis integris aut subdentatis, caulibus subadscendentibus, petalorum unguibus calyce vix longioribus, laminis obovatis, siliquis erectiusculis longissimis, stylo longiusculo. ② in Rhætiâ. *Cheiranthus Rhæticus* Schleich! pl. exsic. Flores flavi. ferè *E. lanceolati*, (v. s.)

28. *E. DUBIUM* (DC. syst. 2. p. 504.*) foliis lanceolatis dentatis, basi attenuatis, petalorum laminis obovato-oblongis, siliquis patentibus, stylo subnullo ② hab. . . . *Cheiranthus dubius* Horn! h. hafn. suppl. p. 73. (v. s.)

29. *E. LONGIFOLIUM* (DC. syst. 2. p. 504.*) foliis lineari-lanceolatis elongatis subdentatis, petalorum laminis obovato-oblongis, siliquis erectis pubescentibus, stylo longiusculo. ② in fissuris rupium Algeriæ. *E. grandiflorum* Desf! fl. atl. 2. p. 85. * non Bieb. Flores vix magnitudine *E. lanceolati*. (v. s.)

30. *E. GRACILE* (DC. syst. 2. p. 504.*) foliis lineari-lanceolatis dentatis acuminatis, petalorum laminis obovato-oblongis, siliquis erectis pube ramosa subscabris, stylo subnullo. — Ad Caucasum sept. et in Iberiâ. Flores flavi. Petalorum unguis calycis longitudine. (v. s.)

31. *E. RIGIDUM* (DC. syst. 2. p. 505.*) foliis lineari-lanceolatis acutè dentatis, siliquis patulis glabris rigidis, stylo vix ullo. — in Oriente. Deless! ic. sel. 2. t. 67. Flores ignoti. Siliquæ tetragonæ rigidæ. (v. s. in h. Deless.)

32. *E. SCABRUM* (DC. syst. 2. p. 505.*) foliis lineari-lanceolatis, inferioribus obtusis subdentatis, racemo abbreviato, siliquis erectis tuberculato-scabris, stylo conico-filiformi. — in Libano. Flores ignoti. (v. s. in h. Labill.)

33. *E. ASPERUM* (DC. syst. 2. p. 505.*) foliis lineari-oblongis, inferioribus dentato-runcinatis cauleque pubescentibus scabris, siliquis patentibus stylo brevissimo crasso. ② in campis Missouriensibus. *Cheiranthus asper* Nutt. gen. am. 2. p. 436. Siliquæ adpressæ pubescentes. (v. s. in h. Banks.)

34. *E. STRIGOSUM* (DC. syst. 2. p. 506.*) undiquè strigosum, foliis oblongo-lanceolatis integerrimis vel denticulatis, siliquis erectis, stigmatibus bipartito sessilibus. ② in Sibirîâ. *Cheiranthus strigosus* Ledeb. in mem. acad. Peters. 5. ann. 1815. p. 549. Flores erecti, pedicellis ipsorum longitudine insidentes.

35. *E. REDOWSKII* (Weinm. cat. hort. Dorp. 1810. p. 65.*) foliis omnibus lineari-canaliculatis integerrimis canescentibus, caule ramoso angulato, petalorum laminis emarginatis, siliquis erecto-patulis obtusangulis. ② in Sibirîâ. Flores pallide lutei, magni, petalis obcordatis. DC. syst. 2. p. 506.*

Sect. IV. CORINGIA. DC. syst. 2. p. 507.

Stylus vix ullus. Petala erectiuscula. Flores pallide flavi aut albi. — Folia cordato-amplexicaulia.

36. *E. ALPINUM* (Baumg. fl. transylv. 2. p. 263.) foliis membranaceis glabris, caulibus cordato-sagittatis amplexicaulibus oblongis, radicalibus petiolatis ovatis. ② in sylvaticis montanis Europæ. *Brassica Alpina* Lin. mant. 95. Vill. dauph. 3. p. 330. t. 36.* — *Arabis brassicæformis* Wallr! sched. 359. Turritis sagittata Schrank mon. t. 99? — DC. syst. 2. p. 507.* (v. v.)

37. *E. PERFOLIATUM* (Crantz fl. austr. 27.) foliis radicalibus obovatis, caulinis cordato-amplexicaulibus, omnibus obtusis glabris glaucis, siliquæ tetragonæ

faciebus enerviis. ① in arvis Europæ et Asiæ temperatæ et in Japoniâ. Brassica orientalis Lin! spec. 931. Jacq. fl. austr. t. 282. Flores albi, exsiccatione subochroleuci. — DC. syst. 2. p. 508.* (v. v.)

38. E. AUSTRIACUM (Baumg. fl. transylv. 2. p. 263.*) foliis radicalibus obovatis, caulibus cordato-amplexicaulibus, omnibus obtusis glabris glaucis, siliquæ subtetragonæ faciebus nervis elevatis striatis. ① in Hispaniâ, Transilvaniâ et Iberiâ mediâ. Brassica Austriaca Jacq! austr. p. 45. t. 283.* Gorinkia Austriaca Presl. czech. p. 141. Simillimum E. perfoliato et cum eo sæpè confusum. (v. s.)

† Species non satis notæ et ad *Erysimastra* verosimiliter omnes referendæ.

39. E. ALTISSIMUM (Lejeun. fl. Spa. 2. p. 70.*) in sylvis montanis inter Verviers et Limbourg. Folia lineari-lanceolata, canaliculata, integerrima. Petala pallide flava, limbo obcordato. Siliquæ tetragonæ stigmatibus bilobis terminatæ. DC. syst. 2. p. 509.

40. E. BICOLOR (DC. syst. 2. p. 509.) ② in Helvetiâ. Cheiranthus bicolor Horn. h. hafn. 2. p. 613. Folia lanceolata, inferiora remotè dentata, superiora integerrima. Siliquæ patulæ, tetraedræ. DC. syst. 2. p. 509.

41. E. RUMILUM (Horn. h. hafn. 2. p. 613.) ② in Helvetiâ. Folia subdentata, inferiora spathulato-ovata, superiora linearia. Siliquæ patentissimæ, tetraedræ. DC. syst. 2. p. 510.

42. E. PATULUM (Horn. h. hafn. suppl. 73.) ② hab. . . . Folia lyrato-pinnatifida subscabra, superiora sinuata. Siliquæ patulæ. DC. syst. 2. p. 510.

43. E. SYLVATICUM (Bieb. fl. taur. suppl. p. 441.) in Ukrainiæ sylvis. Folia lanceolata remotè subdentata. Rami patentes. Corolla pallidior quàm in E. cheiranthoide cui ceterum accedit. Siliquæ patentes. Stylus longitudinem siliquæ duplò superans.

44. E. MARSCHALLIANUM (Andrz! in Bieb. fl. taur. suppl. p. 441.) in Podoliâ australi. Accedit ad E. hieracifolium, sed ut videtur foliis siliquisque strictioribus duplò brevioribus diversum. DC. syst. 2. p. 510. (v. s. sine fr.)

LV. LEPTALEUM. DC. syst. 2. p. 510.

Siliqua teretiuscula sessilis, stigmata duo conniventia. Calyx basi æqualis. Stamina 4-6. Semina plurima 1-serialia. . . .

1. L. FILIFOLIUM (DC. syst. 2. p. 511.) foliis aliis integris aliis parcè lobatis, caule siliquis longiore, siliquis pube minimâ scabris ① in Sibirîâ ad flumen Kumam. Deless! ic. sel. 2. t. 68. Sisymbrium filifolium Willd! spec. 3. p. 496.* Siliquæ axillares erectiusculæ aut basi deflexæ. (v. s.)

2. L. PYGMÆUM (DC. syst. 2. p. 511.) foliis ferè omnibus pinnatifidis, caule siliquis glabris submuricatis brevior. ① in Persiâ. Valdè priori affine, sed differt 1. caule subnullo, 2. foliis ferè omnibus pinnatifidis, lobis filiformibus distantibus. Deless! ic. sel. 2. t. 68. (v. s.)

LVI. STANLEYA. Nutt. gen. am. n. 166. DC. syst. 2. p. 511.

Siliqua teretiuscula super torum pedicellata. Semina oblonga, teretiuscula. Calyx patens.

1. St. PINNATIFIDA (Nutt! gen. am. 2. p. 71*) foliis interruptè pinnatifidolobatis. 2l in Louisianâ superiore. Cleome pinnata Pursh! fl. bor. am. 2. p. 739. Flores flavi, ferè Cleomes. — DC. syst. 2. p. 512.* (v. s. in h. Lamb.)

2. St. GRACILIS (DC. syst. 2. p. 512.*) foliis superioribus oblongis integris sessilibus basi attenuatis. ① in Americâ bor. Cleome lævigata Soland! in h. Banks. Flores parvi, ex sicco flavi, hexandri. (v. s. in h. Banks.)

3. St. AMPLIXIFOLIA (Nutt. in Sillim. am. journ. 5. 1822. p. 297.) foliis integris?

amplexicaulibus, floribus corymbosis, siliquis nutantibus. — in Floridæ orientalis aridis pinetis. Herbacea lævis glauca. Siliq. 2-3-poll. longæ.

Trib. VIII. *CAMELINEÆ* seu *NOTORHIZEÆ LATI-SEPTÆ*. DC. syst. 2. p. 513.

Silicula valvulis concavis, septo elliptico in majore diametro. Semina ovata. Cotyledones planæ, incumbentes, septo contrariæ.

LVII. *STENOPETALUM*. Brown in. DC. syst. 2. p. 513.

Siliqua ellipsoidea, valvulis concavo-planis, loculis polyspermis. Stylus o. Semina in quoque loculo biserialia minima subovata.

1. *St. LINEARE* (Brown! in DC. syst. 2. p. 513.) ① in orâ australi Novæ-Hollandiæ. Herba circiter pedalis, tenuissima. Folia 7-8 lin. longa, semi-lineam lata. Siliculæ 2 lin. longæ, lineam latæ. (v. s. comm. à cl. amic. Brown.)

LVIII. *CAMELINA*. Crantz fl. austr. 1. p. 17. Medik. gen. pl. 1. p. 67. t. 1. f. 11.

Silicula obovata aut subglobosa, valvulis ventricosus cum styli parte dehiscens, loculis polyspermis. Stylus filiformis. Semina oblonga immarginata.

Sect. I. *CHAMÆLINUM*. DC. syst. 2. p. 514.

Siliculæ obovatæ marginatæ. Stylus conicus. Stigma simplex. ①

1. *C. ARMENIACA* (Desv! Journ. 3. p. 182.*) siliculis basi obconicis elongatis stylo brevi apiculatis, foliis lineari-lanceolatis integerrimis. ① in Armeniâ. DC. syst. 2. p. 514. (v. s. in h. Mus. Par.)
2. *C. SATIVA* (Crantz fl. austr. p. 10.) siliculis cuneato-pyriformibus 4-costatis stylo longiusculo terminatis, foliis subintegerrimis lanceolatis. ① inter segetes et lina in Europâ totâ. Myagrum sativum. Lin. spec. 894. Cav. ic. 1. p. 47. t. 66.* — DC. syst. 2. p. 515.* (v. v. et s.)
 α ? *pilosa*, foliis integris pilosis. Semper sylvestris. Moris. hist. 2. p. 315. s. 3. t. 21. f. 2. *C. sylvestris* Wallr. sched. 347. Forsan species propria et var. β cum sequente specie conjungenda ex Wallr. Sched. 347.
 β . *glabrata*, foliis integris subglabris. Sæpius sativa Lind. als. 94. t. 1.
3. *C. DENTATA* (Pers. ench. 2. p. 191.) siliculis subgloboso-pyriformibus 4-costatis stylo longiusculo superatis, foliis repando-dentatis. ① in arvis Alsatiæ, Palatinatûs, Hispaniæ. DC. syst. 2. p. 516. (v. v. et s.)
 α . *dentata*, foliis glabriusculis grossè dentatis aut sinuatis. Myagrum dentatum Willd. phyt. 1. p. 9. n. 30.* — J. Bauh. hist. 2. p. 893. ic.
 β . *pinnatifida*, foliis sinuato-pinnatifidis, pube sparsâ scabris. Myagrum pinnatifidum Ehrh. dec. ex herb. Smith.
4. *C. MICROCARPA* (Andrz! cruc. in DC. syst. 2. p. 517.) siliculis pyriformibus bicostatis stylo longiusculo superatis, foliis lanceolatis denticulatis bispidis. ① in Podoliâ. Accedit ad var. pilosam *C. sativæ*, sed differt fructu dimidiò breviorè, hi nec quadricostato. Deless! ic. sel. 2. t. 69.

Sect. II. *PSEUDOLINUM*. DC. syst. 2. p. 517.

Siliculæ globosæ immarginatæ. Stylus filiformis. Stigma capitatum 2.

5. *C. BARBAREÆFOLIA* (DC. syst. 2. p. 517.*) siliculis globosis, foliis oblongis pinnatifidis basi obtusè auriculatis, caule basi villosa. 2 in Sibiria orient. Deless. ic. sel. 2. t. 70. Flores, fructus, semina imò cotyledones *C. Austriacæ*. (v. s.)
6. *C. AUSTRIACA* (Brown in h. kew. ed. 2. v. 4. p. 93.) siliculis globosis, foliis oblongis serrato dentatis basi obtusè amplexicaulibus, caule glabro. 2 in

pratis humentibus. Austriæ, Tauriæ, etc. *Myagrum Austriacum* Jacq! fl. austr. 2. t. 111.* — DC. syst. 2. p. 517.* (v. s.)

LIX. EUDEMA. *Humb. et Bonpl. pl. equin.* 2. p. 133. t. 123 et 124.

Silicula ovata, loculis polyspermis, valvulis concavis, septo apice fenestrato. Stylus filiformis. Semina ovalia, cotyledonibus (ex icone) incumbens.

1. *E. RUPESTRIS*. (*Humb. et Bonpl. pl. equin.* 2. p. 133. t. 123.*) foliis linearibus, calycibus corollâ brevioribus, siliculæ loculis 10-12-spermis. $\frac{1}{2}$ in Americâ merid. H. B. et K. nov. sp. an. 5. p. 79. *Draba Humboldtii* Desv. journ. bot. 3. p. 171. Flores axillares, solitarii. — DC. syst. 2. p. 519.* (v. s. in h. Humb.)
2. *E. NUBIGENA* (*Humb. et Bonpl!* 2. p. 136. t. 124.*) foliis spathulatis, calycibus corollâ longioribus, siliculæ loculis 4-spermis. $\frac{1}{2}$ in rupibus elatis Andium Quitensium. H. B. et K. nov. spec. am. 5. p. 79. Petala alba. DC. syst. 2. p. 519. (v. s. in h. Humb.)

LX. NESLIA. *Desv. journ.* 3. p. 162. et 163. DC. syst. 2. p. 519.

Silicula subglobosa, valvis concavis, septo evanido 1-locularis indehiscens 1-sperma. Semina subglobosa, pendula. Cotyledones ovatæ crassæ incumbentes.

1. *N. PANICULATA* (*Desv. l. c.*) (I) in arvis et arenosis Europæ ferè totius. *Myagrum paniculatum* Lin! spec. 894. &d. fl. dan. t. 204. *Rapistrum paniculatum* Gaertn. fruct. 2. p. 285. t. 141. (v. v.)

Trib. IX. LEPIDINEÆ seu NOTORHIZEÆ ANGUSTI-SEPTÆ. DC. syst. 2. p. 521.

Silicula septo angustissimo, valvis carinatis aut valdè concavis. Semina in loculis solitaria aut pauca, ovata, immarginata. Cotyledones planæ, incumbentes, septo parallelæ.

LXI. SENEBIERA. *Poir. dict.* 7. p. 75. DC. syst. 2. p. 521.

Silicula didyma, valvis ventricosis aut subcarinatis subindehiscens 1-spermis. Semina subgloboso-triquetra. Cotyledones incumbentes lineares. — Racemi oppositifolii. Flores albi.

Sect. I. NASTURTIOLUM. DC. syst. 2. p. 522. — *Medik. gen. pl.* p. 82. t. 2. f. 21. ex *Ust. neu. ann.* 2. p. 45.

Siliculæ apice emarginatæ, septo nempè valvulis globosis brevioræ.

1. *S. INTEGRIFOLIA* (DC. soc. hist. nat. par. ann. 7. p. 144. t. 8.*) foliis linearibus integris basi attenuatis, siliculis biglobosis fungosis rimoso-areolatis. — In Madagascar. DC. syst. 2. p. 522.* Semina rufa, subrotunda. (v. s.)
2. *S. LINOIDES* (DC. syst. 2. p. 522.*) foliis linearibus integris acutis, siliculis subcompressis biscutatis minutissimè areolato-punctatis. — Ad cap. Bonæ-Spei. *Lepidium linoides* Thunb. prod. 107? Flores albi parvi. Folia ferè *S. integrifoliæ* et siliculæ *S. pinnatifidæ*. (v. s. in h. Burchell.)
3. *S. HELENIANA* (DC. syst. 2. p. 523.*) foliis inferioribus pinnatipartitis, lobis incisis, superioribus linearibus aut in lobos paucos divisus, siliculis biglobosis subareolatis. (I) in insulâ Sanctæ-Helenæ. Flores parvi, albi. (v. s. comm. à cl. Burchell.)
4. *S. PECTINATA* (DC. syst. 2. p. 523.*) foliis pinnatipartitis, lobis linearibus acuminatis integris, siliculis compressis didymis reticulatis transversè nervulosis. — In regno Quitensi. *S. pinnatifida* H. B. et K. nov. sp. 5. p. 76. Nimis affinis *S. pinnatifidæ*, et ab eâ forsitan non satis distincta. (v. s. in h. Humb.)

5. *S. PINNATIFIDA* (DC. mem. soc. hist. nat. par. ann. 7. p. 144. t. 9.*) foliis pinnatilobatis, lobis oblongis dentatis aut subincisis, siliculis compressis didymis reticulatis. ① ② in ruderatis præsertim maritimis, Europæ occid. etc. *Lepidium didymum* Lin! mant. 92. Flores albi, parvi. Petala nunc nulla, nunc 4 oblonga. — DC. syst. 2. p. 523.* (v. v.)

β. *incisa*, foliorum lobis 3-4-partitis. in Pensylvaniâ, Carolinâ, Brasiliâ, etc. *S. incisa* Willd. enum. 2. p. 668.

Sect. II. *CARARA*. DC. syst. 2. p. 524.

Siliculæ apice non emarginatæ, utrinquè subcompressæ, valvulis compressis dorso cristato-rugosis.

6. *S. CORONOPUS* (Poir. dict. 7. p. 76.) foliis pinnatilobatis, lobis integris dentatis pinnatifidisve, siliculis acutiusculis compressis, valvis dorso cristato-rugosis. ① ② Secus vias in calcareis et arenosis Europæ ferè totius. *Cochlearia coronopus* Lin! spec. 904. OEd. fl. dan. t. 202. *Coronopus Ruellii* Gært. fr. 2. t. 242. Sepala subrotunda margine albida membranacea. — DC. syst. 2. p. 525.* (v. v.)
7. *S. SERRATA* (Poir! dict. 7. p. 76.*) foliis oblongis inciso-serratis, siliculis compressis rugosis. — in Montevideo Brasiliæ. *Coronopus serratus* Desv. journ. 3. p. 163. Flores minimi. Siliculæ compressæ, basi suborbiculatæ, valvis reticulatis glabris rugosis. — DC. syst. 2. p. 526.* Deless! ic. sel. 2. t. 71. (v. s.)

Sect. III. *COTYLISCUS*. DC. syst. 2. p. 526.

Siliculæ nec cristatæ, nec apice emarginatæ, hinc concavo-planæ aut ferè planæ.

8. *S. NILOTICA* (DC. syst. 2. p. 527.*) foliis radicalibus pinnatipartitis, lobis inciso-serratis, summis subintegris, siliculis compressis subcymbæformibus. ① in insulis Niloticis Egypti inferioris. *Cochlearia Nilotica* Delil! ill. fl. eg. p. 19. descr. p. 101.* t. 34. f. 2. Siliqua reniformis stigmate sessili subapiculata. (v. s.)

LXII. *LEPIDIDIUM*. Brown in h. Kew. ed. 2. v. 4. p. 85.

Silicula ovata aut subcordata, valvis carinatis aut rariùs ventricosis, dehiscentibus, loculis 1-spermis. Semina subtriquetra aut compressa. — Racemi terminales. Flores albi. DC. syst. 2. p. 527.

Sect. I. *CARDARIA*. DC. syst. 2. p. 528. *Cardiolepis* Wallr. sched. 340.

Silicula ovato-cordata, valvis subturgidis apteris, stylo filiformi exserto superata.

1. *L. DRABA* (Lin. spec. ed. 1. p. 645.) siliculis cordatis subturgidis apice integris stylo superatis, foliis amplexicaulibus lanceolatis dentatis. ① in arvis Europæ præsertim australis. *Cochlearia Draba* Lin! spec. ed. 2. p. 904. Jacq. fl. austr. t. 315. Cotyledones obovato-oblongæ, crassæ. — DC. syst. 2. p. 529.* (v. s.)

Sect. II. *ELLIPSARIA*. DC. syst. 2. p. 530.

Silicula elliptica, valvis carinatis apteris, stylo filiformi longo.

2. *L. CHALEPENSE* (Lin! amœn. 4. p. 321. spec. ed. 2. p. 898.) siliculis ellipticis pedicello duplò longioribus, stylo filiformi, foliis auriculis acutis amplexicaulibus lanceolatis dentatis. — in Oriente. DC. syst. 2. p. 530.* (v. s.)
- α. *glabratum*, foliis à basi ultrà medium acutè dentatis, racemis laxiusculis. Moris. oxon. 2. p. 314. s. 3. t. 25. f. seriei infimæ secunda. — in agro Chalepensi.
- β. *pubescens*, differt racemis fructiferis magis laxis, pedicellisque longioribus. — in Syriâ.
- γ. *canescens*, multò magis villosa-velutina et cinerea quàm priores et racemis magis abbreviatis distinguitur. — Ad montem Libanum.

3. *L. oxyotum* (DC. syst. 2. p. 530.*) siliculis ellipticis puberulis pedicello sublongioribus, stylo filiformi, foliis auriculis acutis subhastatis oblongis subdentatis. — in Syriâ Petala alba ungue filiformi, calycis longitudine. Semina ovata, rufa. DC. syst. 2. p. 530.* (v. s. in herb. Labill.)
4. *L. GLASTIFOLIUM* (Desf! fl. atl. 2. p. 66. t. 147.*) siliculis ellipticis glabris pedicello brevioribus, stylo filiformi, foliis auriculis obtusis amplexicaulibus oblongis obtusè dentatis. — in montibus umbrosis Algeriæ. *Thlaspi glastifolium* Poir. dict. 7. p. 538.* Petala obovata, alba, calyce duplò longiora. — DC. syst. 2. p. 531.* (v. s. in h. Deless.)
5. *L. AMPLEXICAULE* (Willd! spec. 3. p. 436.*) siliculis subrotundis stylo brevissimo terminatis, foliis indivisis oblongo-lanceolatis sagittatis sessilibus apice dentatis. ½ in Sibiariâ. Flores et siliculæ ut in *L. latifolio* quocum in multis convenit. DC. syst. 2. p. 531.* (v. s. ex h. Willd.)

Sect. III. BRADYPIPTUM. DC. syst. 2. p. 531.

Silicula elliptica, valvis carinatis, stylo brevi, calyce persistente. — Folia caulina nec amplexicaulia, nec auriculata.

6. *L. CÆSPITOSUM* (Desv. journ. bot. 3. p. 165 et 178.) siliculis ovatis angustè emarginatis, stylo brevissimo, calycibus persistentibus, foliis linearibus integris persistentibus. ½ in Armeniâ. — Sepala ovali-oblonga, margine membranacea, persistentia, demùm subreflexa. DC. syst. 2. p. 532.* (v. s. in h. Mus. Par.)
7. *L. CORONOPOLIUM* (Fisch! in cat. h. gor. 1808. p. 79.) siliculis ellipticis integris subpuberulis stylo apiculatis, calyce subpersistente, foliis radicalibus pinnati-lobatis, caulinis linearibus integerrimis. ½ Propè Sareptam ad Volgam australem et ad lacum Inderiensem. *L. laciniatum* Willd. herb. ex Stev. obs. ined. Flores parvi albi. — DC. syst. 2. p. 532.* (v. s.)
8. *L? HUMBOLDTI* (DC. syst. 2. p. 532.*) siliculis ovatis emarginatis stylo brevi apiculatis, calycibus persistentibus, foliis radicalibus bipinnatisectis, caulinis pinnatifidis. — Locis aridis propè Chillo Quitensium altit. 1340. hexap. *Senebiera dubia*. H. B. et K. nov. spec. 5. p. 76. Flores pedicellati, minuti, albi. Petala calyce parum breviora. (v. s. in h. Humb.)

Sect. IV. CARDAMON. DC. syst. 2. p. 533.

Silicula suborbiculata emarginata, valvis navicularibus dorso alatis, cotyledonibus partitis.

9. *L. SATIVUM* (Lin! spec. 899.) siliculis orbiculatis alatis, foliis variè divisis incisisque, ramis non spinescentibus. ① in Persiâ et insulâ Cipro, nunc in hortis Europæ frequens cultum. Boiss. fl. eur. t. 440. f. 2. DC. syst. 2. p. 533.* (v. v.)

ℓ. *crispum*, foliis valdè incisis undulato-crispis. *Nasturtium crispum*. J. Rauh. hist. 2. p. 913. f. 1.

γ *latifolium*, foliis planis parùm divisis. — Moris. oxon. 2. p. 300. s. 3. t. 19. f. 2.

10. *L. SPINASCENS* (DC. syst. 2. p. 534.*) siliculis ovalibus subulatis emarginatis, foliis apice incisis, racemorum rachibus apice spinescentibus. — in Syriâ circa Damascus. Deless! ic. sel. 2. t. 72. Flores ignoti. Cotyledones ut in *L. oleraceo* incumbentes, tripartitæ. (v. s. in h. Desfont. et Labill.)

Sect. V. LEPIA. DC. syst. 2. p. 534.

Silicula suborbiculata emarginata, valvis navicularibus alatis, alis stylo adnatis, stylo brevissimo, cotyledonibus integris.

11. *L. CAMPESTRE* (Brown in h. Kew. ed. 2. v. 4. p. 465.) siliculis ovatis alatis emarginatis lepidoto-punctatis, foliis caulinis sagittatis dentatis. ① ② in viis, agris argillosis, apricis arvis Europæ. *Thlaspi campestre*. Lin! spec. 902. Smith engl. bot. t. 1385. Iberis camp. Wallr. sched. 341. — DC. syst. 2. p. 535.* (v. v.)

β. subglabrum, foliis glabris, siliculis ellipticis. — Moris. oxon. 2. p. 293. s. 3. t. 17. f. 13. in mont. Legionensibus Hisp.

12. *L. hirtum* (Smith comp. ed. 3. p. 98. engl. bot. t. 1803.) siliculis ovatis alatis emarginatis hirtis, foliis caulinis sagittatis villosis subintegris. ② in arvis lapidosis Europæ australis. Thlaspi hirtum Lin! spec. 901. Aff. priori. Flores albi paulò majores. — DC. syst. 2. p. 536.* (v. v.)
13. *L. leiocarpum* (DC. syst. 2. p. 536.*) siliculis ovalibus alatis emarginatis glabris cymbæformibus. — in monte Libano. Folia et flores ignoti. Semina *L. hirtum* similia. — Species recognoscenda. †.
14. *L. rotundum* (DC. syst. 2. p. 537.*) siliculis ovalibus alatis emarginatis glabris, foliis caulinis lanceolato-linearibus subcarnosis glabris — in Novâ-Hollandiâ. Lepia rotunda Desv. journ. bot. 3. p. 166 et 181.* Flores ignoti. Semina grossa, rufa. Cotyledones crassæ. (v. s. in h. Mus. Par.)*
15. *L. spinosum* (Lin! mant. 253*) siliculis oblongis alatis emarginatis subbicornibus glabris, foliis radicalibus pinnatisectis, lobis incis. ① ② in Oriente. Ard! specim. 2. p. 34. t. 16. Flores parvi, albi. Sepala margine albida. DC. syst. 2. p. 537.* (v. s.)

Sect. VI. DILEPTIUM. DC. syst. 2. p. 538.

Silicula subelliptica, apice breviter emarginata, valvis carinatis apteris, stylo brevissimo. — Flores minimi interdum 2-4-andri aut rariùs apetal. 1.

16. *L. virginicum* (Lin! spec. 900.) siliculis orbiculatis emarginatis pedicello brevioribus, floribus 2-4-andris, foliis caulinis lineari-lanceolatis incisoserratis glabris. ① in Americâ bor. *L. Iberis* Schkuhr handb. 2. p. 222. t. 180. Flores albi minimi. Petala obovata obtusa. Semina ovalia compressa. — DC. syst. 2. p. 538.* (v. v. et s.)
17. *L. spicatum* (Desv! journ. bot. 3. p. 164 et 178.*) siliculis orbiculatis emarginatis pedicello sublongioribus, foliis linearibus integris glabris. — Ad fretum Magellanicum. Aff. *L. Virginico*. Siliculæ quasi spicatum imbricatæ. DC. syst. 2. p. 539.* (v. s. in h. Mus. Par.)
18. *L. Menziesii* (DC. syst. 2. p. 539.*) siliculis orbiculatis subemarginatis, foliis pubescentibus, radicalibus pinnatipartitis lobis incis. superioribus linearibus integris. ②? in orâ occidentali Americæ bor. Accedit ad *L. Virginicum* sed differt præcipuè foliis pube brevi subvillosis. (v. s. in h. Banks et Smith.)
19. *L. subulatum* (Lin! spec. 899.) siliculis ovatis subemarginatis, foliis subulatis integris, caule suffruticoso. ⑤ in Hispaniæ collibus gypsaceis. Asso. syn. arr. p. 83. t. 6. f. 3. Flores albi, parvi. Calyx albidus, patens. DC. syst. 2. p. 539.* (v. s.)
20. *L. ruderales* (Lin! spec. 900.) siliculis ovatis emarginatis patulis pedicello brevioribus, foliis glabris, radicalibus pinnatilobatis, rameis linearibus integris, floribus diandris apetalis. ① Secùs vias Europæ totius. Smith engl. bot. t. 1595. (Ed. fl. dan. t. 184. DC. syst. 2. p. 540.* (v. v.)
21. *L. incisum* (Roth nov. cat. 1. p. 224.) siliculis orbiculatis levissimè emarginatis, floribus diandris, foliis imis oblongis lyrato-pinnatilobatis, inferioribus lato-lanceolatis inciso-dentatis, summis linearibus. ① in Tauriâ, ad Caucasum et mare Caspium. Bieb. fl. taur. 2. p. 98.* Aff. *L. ruderali*, sed differt foliis caulinis lineari-lanceolatis latioribus. DC. syst. 2. p. 541.* (v. s.)
22. *L. vesicarium* (Lin. spec. 898.) siliculis ellipticis levissimè emarginatis, foliis pinnatisectis, lobis linearibus, geniculis caulinis inflatis. ① in sterilibus per Caucasum orientalem et Iberiam, in Persiâ et Mediâ. — Buxb. cent. 1. p. 17. t. 26. Flores albi. Semina ferè oblongo-triquetra. — DC. syst. 2. p. 541.* (v. s.)
23. *L. angulosum* (D'Urv. enum. n. 578.) siliculis ovatis subemarginatis, foliis omnibus pinnatisectis, segmentis inciso-dentatis, caule glaberrimo flexuoso angulato sulcato. ① in ruderalis urbis Theodosiæ. Diff. à *L. vesicario* gen-

Spei. Deless! ic. sel. 2. t. 73. *Thlaspi Africanum* Burm! fl. cap. p. 17. (v. s. in h. Deless.)

50. *L. BIPINNATUM* (Thunb! prodr. 107.) siliculis foliis radicalibus bipinnatipartitis, lobis filiformibus. \mathcal{L} 5 ad cap. Bonæ-Spei. Species distinctissima primâ fronte sicca referens *Pimpinellam dioicam*. Fructus ignoti. DC. syst. 2. p. 553.* (v. s. in h. Banks et Smith.)

†. *Species non satis notæ.*

51. *L. PINNATUM* (Thunb. prod. 107.) caule fruticoso erecto, foliis omnibus pinnatis. — Ad caput Bonæ-Spei. DC. syst. 2. p. 553.
52. *L. DECUMBENS* (Desv. journ. bot. 3. p. 165. et 176.) hab. ... *L. divaricatum* Willd. h. berol. ex Desv. Folia quasi petiolata, grandidentata. Siliculæ ovato-oblongæ, apice subrotundæ, stylo vix exserto. DC. syst. 2. p. 553.
53. *L. FRUTICOSUM* (Desv. journ. bot. 3. p. 165 et 180.) 5 in Novâ-Hollandiâ. Folia glabra subcarnosa, ovato-lanceolata. Siliculæ oblongæ, apice angustatæ, obtusæ aut subtruncatæ, pedicello breviores. DC. syst. 2. p. 553.
54. *L. TUBEROSUM* (DC. syst. 2. p. 554.*) — in Louisianâ. *Nasturtium tuberosum* Raf. fl. lud. p. 84. n. 271. Folia radicalia pinnata. Siliquæ rotundatæ, emarginatæ. Stigma sessile. An potius *Thlaspi* ad *T. tuberosum* accedens.
55. *L. PRÆCOX* (DC. syst. 2. p. 554.) in Louisianâ. *Dileptium præcox* Raf. fl. lud. p. 86. n. 273. Folia radicalia pinnata, caulina pinnatifida. Flores albi. Petala calyci æqualia. Aff. priori.
56. *L. DIFFUSUM* (DC. syst. 2. p. 554.) in Louisianâ. *Dileptium diffusum* Raf. fl. lud. p. 85. n. 272. Folia parva, pinnatifida, pinnulis dentatis. Petala minutissima calyce breviora.
57. *L. POLLICHI* (Roth germ. II. 91.) (I) in Germaniâ. *L. Iberidis* Poll. pal. n. 607.* ex Roth. Sub hoc nomine recepi nunc *L. sativi* var. laciniatam, nunc *L. Iberidem* foliis radicalibus donatum. DC. syst. 2. p. 554.
58. *L. CORDATUM* (Willd. herb. ex Stev. obs. ined.) — in Sibirîâ. *Glancum*, foliis cordato-oblongis integerrimis amplexicaulibus. *L. amplexicaule*. Stev. herb. non Willd. — DC. syst. 2. p. 554.

LXIII. BIVONÆA. DC. syst. 2. p. 554.

Silicula ovalis, emarginata, valvis carinatis dorso alatis. Semina in quoque loculo 4-6 pendula, ovata. — Flores parvi lutei.

1. *B. LUTEA* (DC. syst. 2. p. 555.) (I) in montibus siccis Panormi. *Thlaspi lateum*. Biv! cent. 1. p. 78.* — Cup. panph. sicil. 2. t. 256. ex Biv. (v. s.)

LXIV. EUNOMIA. DC. syst. 2. p. 555.

Silicula ovalis, valvis carinatis; semina in quoque loculo 2, oblongo-triquetra, funiculis subcoadunatis. — Folia opposita integra. Flores albid.

1. *E. OPPOSITIFOLIA* (DC. syst. 2. p. 556.*) foliis oppositis suborbiculatis margine integerrimis lævibus. \mathcal{L} 5 in monte Libano. *Thlaspi oppositifolium* Poir! suppl. 5. p. 277. *Lepidium oppositifolium* Labill! syr. dec. 5. p. 14. t. 9. f. 2.* (v. s.)
2. *E. CHLORÆFOLIA* (DC. syst. 2. p. 556.*) foliis oppositis suborbiculatis margine crenulato-scabris. \mathcal{L} 5 in Olympi Bithyni cacumine. *Iberis chloræfolia* Sibth! ex Smith prodr. fl. gr. 2. p. 10.* Petala ignota. Priori nimis affinis. (v. s. in h. Banks.)
3. *EP. CORDATA* (DC. syst. 2. p. 557.*) foliis infimis oppositis, ceteris alternis, omnibus cordiformibus. 5 in Armeniâ et Syriâ. *Thlaspi cordatum* Desf. ann. mus. 11. p. 382. t. 39.* An genus proprium? (v. s. in h. Mus. Par.)

LXV. AETHIONEMA. Brown in h. Kew. ed. 2. v. 4. p. 80.

Silicula ovalis, sæpius emarginata, valvis navicularibus dorso alatis,

loculis 1-2-spermis. Stamina majora aut coalita aut dentata. Semina ovato-oblonga, sub lente muriculata.

1. *Æ. SAXATILE* (Brown in h. kew. ed. 2. v. 4. p. 80.) siliculis bilocularibus polyspermis obcordatis, stylo brevissimo, valvulis dorso alatis integris, staminibus majoribus dentatis, racemis fructiferis laxis. 5 2 ① in rupestribus Europæ australis. *Thlaspi saxatile* Lin. spec. 901. Jacq. fl. austr. t. 236. — DC. syst. 2. p. 558. (v. s.)

β. *obtusifolium*. *Iberis parviflora* Lam. dict. 3. p. 221.* — Moris. hist. 2. p. 297. s. 3. t. 18. f. 30.

γ. *ovalifolium*. *Thlaspi marginatum* Lapeyr. suppl. 90.

2. *Æ. GRACILE* (DC. syst. 2. p. 559.*) siliculis bilocularibus oligospermis obcordatis, stylo valvulis æquali, valvulis dorso alatis subdentatis, staminibus majoribus dentatis, racemis fructiferis laxis. 5 in collibus aridis Carniolæ. — Bocc. mus. p. 79. t. 70. Inflorescentia similis *Æ. saxatili*. (v. s.)

3. *Æ. CRISTATUM* (DC. syst. 2. p. 560.*) siliculis bilocularibus 2-4-spermis orbiculatis basi et apice emarginatis, valvulis dorso alatis grossè cristatodentatis. ① in Syriâ. *Thlaspi peregrinum* Poir! dict. 7. p. 541. Flores minimi, erecti, floribus *Æ. saxatilis* sat similes. Deless! ic. sel. 2. t. 74. (v. s.)

β. *Thlaspi Arabicum* Desv. Journ. bot. 3. p. 167. Inter Alep et Mossul.

4. *Æ. BUXBAUMII* (DC. syst. 2. p. 560.*) siliculis bilocularibus 2-spermis orbiculatis basi et apice emarginatis, in valvularum dorso alatis integerrimis, in racemum lupuliformem confertè aggregatis. ① in Cappadociæ et Iberiæ arenosis. *Thlaspi Arabicum* Bieb. fl. taur. suppl. 430.* — Buxb. cent. 1. p. 2. t. 5. f. 1. Flores parvi, purpurascens, omnino *Æ. saxatilis*. (v. s.)

5. *Æ. CORIDIFOLIUM* (DC. syst. 2. p. 561.*) siliculis bilocularibus 2-spermis obcordatis confertis, valvulis dorso alatis integris, foliis linearibus confertis, inferioribus patulis. 5 in monte Libano. Flores roseo-purpurascens, majores quàm in aliis *Æthionematibus*. Deless! ic. sel. 2. t. 76. (v. s.)

6. *Æ. MEMBRANACEUM* (DC. syst. 2. p. 561.*) siliculis bilocularibus 2-spermis obcordatis confertis, valvulis dorso alatis integris, foliis linearibus distantibus strictè adpressis. 5 in Persiâ. Deless! ic. sel. 2. t. 75. Stam. 4 majora dente parvo apice notata, 2 minora integra. (v. s.)

7. *Æ. STYLOSUM* (DC. syst. 2. p. 562.*) siliculis 2-locularibus? 2-spermis laxè racemosis, valvulis dorso alatis, stylo elongato, staminibus majoribus connatis, foliis ovato-lanceolatis. 5 in monte Libano. *Thlaspi carneum* Russel in Schrad. Journ. 1. p. 426. Flores majores quàm in ceteris generis speciebus. (v. s.)

8. *Æ. POLYGALOIDES* (DC. syst. 2. p. 562.*) siliculis 1-locularibus 1-spermis dehiscentibus utrinquè emarginatis, caule basi suffrutesco, foliis oblougolinaribus. 5 in insulâ Scio. Habitus ferè *Polygalæ saxatilis*. Flores ignoti. (v. s. in h. Mus. Par.)

9. *Æ. MONOSPERMUM* (Brown! in h. kew. ed. 2. v. 4. p. 80.) siliculis 1-locularibus 1-spermis indehiscentibus apice emarginatis, caule herbaceo, foliis ovalibus obovatisve. ② in Hispaniâ. Flores paulò majores quàm in *Æ. saxatili*. DC. syst. 2. p. 562.* (v. s. in h. Banks.)

Trib. X. *ISATIDEÆ* seu *NOTORHIZEÆ* *NUCAMENTACEÆ*. DC. syst. 2. p. 563.

Silicula valvis indistinctis aut indehiscentibus carinatis, septo evanido 1-locularis 1-sperma. Semina ovato-oblonga. Cotyledones planæ incumbentes, septo (si adesset) verosimiliter parallelæ.

LXVI. *APHRAGMUS*. Andr. in litt.

Silicula lanceolata acuta, valvis planis nervosis, septo nullo, semina

biseriata pendula, cotyledones incumbentes crassæ. — An potius Camelinea?

1. A. ESCHSCHOLTZIANUS Andr. in litt. — In insulis Alutianis.

LXVII. TAUSCHERIA. Fisch. in DC. syst. 2. p. 563.

Silicula ovalis, ferè cymbæformis, 1-locularis, 1-sperma, valvis navicularibus indehiscentibus. Semen pendulum, oblongum. — Flores minimi, albi.

1. T. LASIOCARPA (Fisch! in litt. cum ic. ined. DC. syst. 2. p. 563.*) siliculis in parte convexâ villosis. ① ad lacum Inderskoe deserti Kirghisorum. Herba, excepto fructu, glabra, subglauca, Isatidem nanam aut Thlaspidem referens. (v. s.)
2. T. GYMNOCARPA (Fisch! in litt. cum. icon. DC. syst. 2. p. 564.*) siliculis glabris. ① Hab. cum priore, hæc ut videtur paulò præcocior. Priori valde similis sed siliquis omninò glabris maturis nigricantibus. (v. s.)

LXVIII. ISATIS. C. Bauh. pin. 113. Lin. gen. n. 824. Lam. ill. t. 554.

Silicula elliptica, plana, 1-locularis, 1-sperma, valvis carinato-navicularibus vix subdehiscentibus. Semen pendulum, oblongum. — Flores lutei, parvi.

Sect. I. SAMERARIA. DC. syst. 2. p. 565. — Desv. journ. 3. p. 161. t. 24. f. 6.

Silicula ovalis aut orbiculata indehiscens, alâ latâ foliaceo-membranaceâ cineta.

1. I. GARCINI (DC. syst. 2. p. 565.*) siliculis latè marginatis obovatis basi cuneatis subpedicellatis apice obtusissimis bituberculatis. ②? in Persiâ. Deless! ic. sel. 2. t. 77. Peltaria Garcini Burm! fl. ind. 139. t. 46. f. 1. — An forsân genus proprium? (v. s. in h. Deless.)
2. I. ARMENA (Lin. spec. 936.) siliculis orbiculatis basi cordatis latè marginatis stylo apiculatis. ① in pratis siccioribus Armeniæ et Iberiæ. Tratt. arch. 2. p. 41. t. 70. ic. Sameraria Armena Desv. journ. 3. p. 161. t. 25. f. 6. DC. syst. 2. p. 566.* (v. s. in h. Mus. Par. et Banks.)
3. I. LATSILIQUA (Stev! mem. soc. nat. mosc. 1812. v. 3. p. 267.) siliculis ellipticis obtusis sessilibus latè marginatis, stigmate sessili, foliorum auriculis obtusis. ② in Cappadociâ. I. Cappadocica. Desv! journ. bot. 1814. 3. p. 174.* — DC. syst. 2. p. 566.* (v. s.)
 ℘. fructu glabro. Stev!
4. I. LEOCARPA (DC. syst. 2. p. 567.*) siliculis ovali-oblongis sessilibus stigmate sessili subtruncatis, foliorum auriculis acutis. — In Libano. Flores paulò minores quàm I. tiuctoriæ. Aff. I. Armenæ et I. latsiliquæ. (v. s. in h. Labill. et Deless.)
5. I. LUSITANICA (Brot. fl. lus. 1. p. 560.* non aliorum.) siliculis latè marginatis obovatis basi cuneatis apice obtusissimis emarginatis, caule folisque glabris. ① in Lusitaniâ. I. glauca. Willd. herb. ex Stev. obs. in h. Willd. — DC. syst. 2. p. 567.* (v. s. in h. Smith.)

Sect. II. GLASTUM. DC. syst. 2. p. 568.

Silicula ovali-oblonga aut ferè linearis margine suberosa, subdehiscens.

6. I. ALPINA (All! ped. n. 944. t. 86. f. 2.*) Siliculis ovali-oblongis utrinquè obtusis margine subfoliaceo-alatis glaberrimis latitudine triplò ferè longioribus ②? in Pedemontii monte Vesulo. Species distinctissima, inter duas generis sectiones media. DC. syst. 2. p. 568.* (v. s.)
7. I. PRÆCOX (Kit. ex Tratt. arch. 2. p. 40. t. 68.* Stev! in Litt.) siliculis ellipticis utrinquè obtusis margine coriaceo-alatis glaberrimis, latitudine triplò

- ferè longioribus. ② in Hungariâ et circa Astrakan. Habitus omninò I. tinctoriæ, sed distincta siliculis basi non aut vix angustioribus et verè ellipticis. DC. syst. 2. p. 568.* (v. s.)
8. *I. LITTORALIS* (DC. syst. 2. p. 568.*) siliculis oblongo-cuneatis obtusissimis truncato-emarginatis basi attenuatis glaberrimis latitudine majore triplò longioribus, loculis utrinquè exsertis et sulco ab alâ distinctis. — In Tauriâ. *I. littoralis* var. *æ.* Stev. in litt. Deless! ic. sel. 2. t. 78. Flores I. tinctoriæ. (v. s.)
9. *I. HEBECARPA* (DC. syst. 2. p. 569.*) siliculis oblongo-cuneatis obtusissimis velutinis basi subattenuatis latitudine vix ultrâ duplò longioribus, loculis sulco laterali destitutis. — In Tauriâ. Deless! ic. sel. 2. t. 79. *I. littoralis* var. *β.* Stev! in litt. (v. s.)
10. *I. TINCTORIA* (Lin! spec. 936.) siliculis cuneatis basi acuminatis apice subspathulatis obtusissimis glabris latitudine majore triplò longioribus, foliis etiam superioribus basi biauriculatis. ② in Europæ australis et temperatæ apricis lapidosis. Smith engl. bot. t. 97. Lam. ill. t. 554. f. 1. DC. syst. 2. p. 569.* (v. v.)
- β. sativa*, foliis glabris latioribus. *I. sativa* Fuchs. hist. 331. ic.
- γ. hirsuta*, foliis hirsutis angustioribus. *I. Alpina* Vill. dauph. 3. p. 308. excl. syn.
- δ. microcarpa*, foliis glabris angustis, siliculis minoribus. *I. Dalmatica* Mill! dict. n. 2?
11. *I. CAMPESTRIS* (Stev! in DC. syst. 2. p. 571.) siliculis oblongis basi angustatis apice obtusiusculis glabris latitudine quadruplò longioribus pedicello ferè brevioribus. — In campestribus Podoliæ austr. Nimis aff. *I. tinctoriæ* (v. s.)
12. *I. BANNATICA* (Link enum. 2. p. 149.) siliculis cuneatis basi acuminatis apice subspathulatis obtusissimis glabris latitudine majore triplò longioribus, foliorum superiorum auriculis nullis. ① in Bannatu, T.
13. *I. MÆOTICA* (DC. syst. 2. p. 571.*) siliculis oblongis basi angustatis apice obtusis emarginatis glabris latitudine quadruplò quintuplòve longioribus. — Ad Mæoticam paludem propè Taganrog. Aff. prioribus sed aliquid apice in adulto statu emarginata. et ante maturitatem omninò lineari et duplò ferè longiore differt. (v. s.)
14. *I. TAURICA* (DC. syst. 2. p. 571.*) siliculis oblongis basi angustatis apice subobtusis glabris latitudine quintuplò longioribus. ② in Tauriæ saxosis calcareis. Accedit ad Mæoticam, sed siliculâ apicem emarginatâ differt. *I. è Tauriâ* Bieb. fl. taur. suppl. p. 422. (v. s. comm. à cl. Beaupré et Stev.)
15. *I. OBLONGATA* (DC. syst. 2. p. 471.*) siliculis elliptico-oblongis utrinquè attenuatis subacutis glabris latitudine quintuplò longioribus, loculo centrali complanato subtrinervio. — In Sibiriâ? Herba glabra ad Isatidem tinctoriam habitu accedens, sed minor. (v. s.)
16. *I. ORIENTALIS* (Willd. enum. 2. p. 663?) siliculis oblongo-linearibus stigmate sessili apiculatis basi vix angustatis pubescentibus latitudine quintuplò longioribus pedicelli longitudinem vix excedentibus. ① in Oriente. Species sæpè confusa cum *I. canescente*, *I. Aleppicâ*, et etiam cum *I. Lusitanicâ*. (v. s.)
17. *I. CANESCENS* (DC. fl. fr. suppl. p. 598.* syst. 2. p. 572.*) siliculis elongato-cuneatis pubescentibus infernè angustatis supernè subspathulatis obtusis, latitudine quadruplò longioribus, pedicelli apice obconici longitudinem duplò excedentibus. ② in apricis maritimis Galloprovinciæ circa Telonem. Habitu accedit hinc ad *I. orientalem*, illinc ad var. *hirsutam* *I. tinctoriæ*. (v. s.)
- β. Iberica*, foliis inferioribus hinc indè grossè dentatis, siliculis apice paulò magis dilatatis et subglabris *I. Iberica* Stev! mem. mosc. 3. p. 267. An species propria?
18. *I. ALEPPICA* (Scop. del. insubrga. p. 31. t. 16.) siliculis linearibus obtusis basi regulariter angustatis pube retrorsâ villosis, latitudine octuplò, pedicello triplò longioribus. ① in Græciæ et Asiæ minoris rupibus maritimis.

I. Lusitanica Lin! spec. 936 excl. syn. Willd. Lam. ill. t. 554. f. 2. — DC. syst. 2. p. 573.* (v. v.)

β. I. dentata. Pers! ench. 2. p. 193. — In Syriâ prope Tripoli.

LXIX. MYAGRUM. Tourn. inst. 211. t. 99. DC. syst. 2. p. 573.

Silicula compressa ferè cuneata, apice lacunas duas vacuas, infernè foculum 1 monospermum gerens. Semen pendulum, oblongum.

1. M. PERFOLIATUM (Lin! spec. 893.) ① in agris et arenosis Europæ mediæ et australis. Schkuhr handb. 2. p. 210. t. 178. opt. DC. syst. 2. p. 574. (v. v.)

LXX. SOBOLEWSKIA. Bieb. fl. taur. suppl. p. 421. DC. syst. 2. p. 575.

Silicula oblonga, compressa, 1-locularis, 1-sperma, evalvis, membranacea. Semen oblongum pendulum. Cotyledones lineares subcurvæ.

1. S. LITHOPHILA (Bieb. cent. pl. rar. ross. 2. t. 59. ex fl. taur. suppl. 421.) 2. ② ① in Tauriâ. Deless! ic. sel. 2. t. 80. Crambe macrocarpa Bieb. fl. taur. 2. p. 90.* Siliculæ plurimæ abortivæ, 3 lin. longæ, lineam ferè latæ.

Trib. XI. ANCHONIEÆ seu NOTORHIZEÆ LOMENTACEÆ. DC. syst. 2. p. 576.

Silicula aut siliqua transversè in articulos 1-spermos secedena. Cotyledones planæ, incumbentes. — Semina ovata.

LXXI. GOLDBACHIA. DC. syst. 2. p. 576.

Stamina libera. Siliqua biarticulata, stylo subnullo. — Flores parvi, ex albo sublilacini.

1. G. LÆVIGATA (DC. syst. 2. p. 577.*) siliquis isthmo biarticulatis lævibus pendulis. ① in arenâ mobili circa Astrakan. Deless! ic. sel. 2. t. 81. Raphanus lævigatus. Bieb. fl. taur. 2. p. 129.* Petala oblonga, integra, calyce duplò longiora. (v. s.)

2. G. TORULOSA (DC. syst. 2. p. 577.*) siliquis subcylindraceis transversè torulosis subadscendentibus. ① in Oriente. Ad priorem valdè accedit et ex herbâ floribus fructibusque junioribus vix distincta. (v. s.)

LXXII. ANCHONIUM. DC. syst. 2. p. 578.

Stamina majora connata. Siliqua biarticulata, stylo compresso rostriformi. Semina 4 in quoque loco solitaria, oblonga, pendula.

1. A. BILLARDIERII (DC. syst. 2. p. 578.*) 2. 5 in Syriæ monte Libano. Deless! ic. sel. 2. t. 82. Flores purpurei, subconferti; sepala linearia, extus villosa. Stigma acutum, subbilobum. (v. s.)

LXXIII. STERIGMA. DC. syst. 2. p. 579. — Sterigmotemon. Bieb. fl. taur. suppl. p. 444. *

Stamina majora ad medium connata. Siliqua teretiuscula in articulos plurimos demum secedens.

1. S. TOMENTOSUM (DC. syst. 2. p. 579.*) foliis omnibus sinuato-pinnatifidis. ② in campis limosis versùs mare Caspium et circa Irtim et Astracan. Cheiranthus tomentosus Willd. spec. 3. p. 523.* — Pall. itin. ed. gall. 8. app. p. 437. n. 352. t. 103. f. 2.* (v. s.)

2. S. SULFUREUM (DC. syst. 2. p. 580.*) foliis inferioribus runcinato-pinnatifidis summis integris. ② in Syriâ prope Aleppum. Deless! ic. sel. 2. t. 83. Cheiranthus sulfureus Russell! in Schrad. journ. 1. p. 426. Flores omninò S. tomentosi. (v. s. in h. Banks et Mus. Par.)

3. S. TORULOSUM (DC. syst. 2. p. 580.*) foliis oblongis, radicalibus sinuato-

dentatis, superioribus integris. (2) in Iberiâ circa Tiflim. Cheiranthus torn-
losus Bieb! fl. taur. 2. p. 121.* Dnobus prioribus valde affine, sed diversum
foliis minùs tomentosis, inferioribus sinuato-dentatis nec pinnatifidis, supe-
rioribus integris, siliquis brevioribus. (v. s.)

4. *S. ELYCHRSISIFOLIUM* (DC. syst. 2. p. 581.*) foliis omnibus oblongo-lineari-
bus integris. 4? (2)? in Armeniâ, Persiâ. Cheiranthus lanatus Gmel. ex
Fisch! in litt. cum ic. Hæc species exhibet ferè folia Anthonii cum fructifi-
catione et floribus Sterigmatis.—Deless! ic. sel. 2. t. 84. (v. s. in h. Mus. Par.)

Subordo III. ORTHOPLOCEÆ (o>>).

Cotyledones incumbentes, conduplicatæ seu longitudinaliter me-
dio plicatæ et radiculam in sinu foventes. Stylus sæpè ampliatus
et basi loculatus seminifer. Semina sæpissimè globosa, semper im-
marginata.

Trib. XII. BRASSICÆÆ seu ORTHOPLOCEÆ SILI- QUOSÆ. DC. syst. 2. p. 581.

Siliqua valvis longitudinaliter dehiscentibus, septo lineari. Se-
mina globosa Cotyledones conduplicatæ.

LXXIV. BRASSICA. Lin. gen. n. 820. excl. spec. plur. DC. syst. 2. p. 582.

Siliqua teretiuscula, stylo parvo brevi obtuso. Semina uniseriata sub-
globosa. Calyx clausus.

Sect. I. BRASSICA. DC. syst. 2. p. 582.

Siliqua sessilis, rostro nullo aut conico aspermo superata.

1. *B. OLERACEA* (Lin! spec. 932.) foliis polline glaucis subcarnosis repandis
lobatisve etiam junioribus glaberrimis. (2) species à vetrissimis temporibus
cultæ et idèò valde polymorpha. DC. syst. 2. p. 583.* diss. in trans. hort.
soc. v. 5. et in ann. agr. franç. v. 19. (v. v.)

A. *SYLVESTRIS*. *B. oleracea* Huds. angl. 289. Smith! engl. bot. t. 637. Caulis
magis elatior quàm in *B. capitata*, humilior et ramosior quàm in *B.*
acephala. Folia non in capitulum verum collecta. Hab. in rupibus mari-
timis Angliæ, etc.

B. *ACEPHALA*, caule tereti elongato, foliis expansis, racemis paniculatis.
B. oleracea viridis Lam. dict. 1. p. 743. n. 2. DC. syst. 2. p. 583.*

a. *ramosa*, caule ramoso, foliis sinuato-pinnatifidis. *B. viridis* procerior
Lam. dict. 1. p. 743. n. 2. var. *ℓ*.

β. *vulgaris*, caule subsimplici, foliis sinuato-pinnatifidis.—Moris. oxon. 3.
t. 1. f. 6.

γ. *quercifolia*, caule subsimplici, foliis pinnatipartitis, lobis incisis, lobu-
lis oblongis.—J. Bauh. hist. 2. p. 832. f. 2.

δ. *Sabellica*, caule subsimplici, foliis sinuato-lobatis, lobis multifidis.—
Lobel. ic. 247. f. 1.

ε. *palmifolia*, caule simplici, foliis sinuato-repandis subbullatis.

ζ? *costata*, caule humiliore subramoso, foliis sinuato-repandis, nervis pri-
mariis crassissimis. *B. oleracea* viridis crassa Lam. dict. 1. p. 743. n. 2. δ.

C. *BULLATA*, caule tereti subelongato, foliis junioribus subcapitatis demùm
patulis bullatis crispatisve, racemis paniculatis.

a. *vulgaris*, capitulis laxis crassis terminalibus subrotundis.—Lobel. ic.
t. 244. f. 1.

β. *oblonga*, capitulo terminali oblongo.—Moris. oxon. s. 3. t. 1. f. 4.

γ. *major*, capitulo terminali maximo. *Brassica* capitata major Lam. dict. 1.
p. 743. n. 3. var. *λ*.

- δ. gemmifera*, capitulis minimis plurimis ex axillis foliorum secus caulem elongatum ortis. — Moris. ox. s. 3. t. 1. f. 3?
- D. CAPITATA, caule tereti abbreviato, foliis concavis non bullatis in capitulum ante florescentiam congestis, racemis papiculatis. — Weinm. phyt. t. 259.
- α. depressa*, foliorum capitulo sphæroideo depresso. — B. capitata compressa Lam. dict. 2. p. 743. n. 3. var. ζ.
- β. sphærica*, foliorum capitulo globoso. — Moris. ox. 2. p. 207. s. 3. t. 1. f. 2.
- γ. obovata*, foliorum capitulo obovato.
- δ. elliptica*, foliorum capitulo elliptico.
- ε. conica*, foliorum capitulo ovato-conico. Br. capitata conica et subacuta Lam. dict. 2. p. 742. n. 3. var. δ et ε.
- E. CAULO-RAPA, caule ad originem foliorum tumido subgloboso.
- α. communis*, foliis planiusculis. — Moris. ox. s. 3. t. 1. f. 12.
- β? crispa*, foliis fimbriato-crispatis.
- F. ROTRITIS, racemorum pedunculis corymbosis confertis ante florescentiam valde carnosus abbreviatis, floribus sæpè abortivis. — Lob. ic. 246. f. 1.
- α. cauliflora*, caule humili, foliis oblongis cæsiis, capitulis crassis terminalibus. — Moris. ox. 2. p. 208. s. 3. t. 1. f. 1.
- β. asparagoïdes*, caule elatiore, foliis cæsiis elongatis, ramulis apice carnosis. (floribus abortivis) alabastra minima gerentibus. — Dalech. Lugd. 522. f. 2.
2. B. CAMPESTRIS (Lin! spec. 931.) foliis polline glaucis subcarnulosis, inferioribus novellis subhispidis ciliatisve lyratis dentatis, cæteris cordato amplexicanalibus acuminatis. ① ② DC. syst. 2. p. 588.* (v. v.)
- A. OLEIFERA*, radice fusiformi exili, caule elongato. — Dalech. lugd. 523. f. 1. — Smith engl. bot. t. 2224. in agris Angliæ, Lapponiæ, etc.
- B. PABULARIA*, radice fusiformi exili, caule abbreviato. Rarius culta in Europâ.
- C. NAPO-BRASSICA*, radice tumida napi-formi. Br. oleracea Napo-Brassica Lin. spec. 932. Frequentius culta in Europâ.
- α. communis*, radice alba, vel subpurpurascente, collo petiolisque subvirescentibus, aut subpurpurascentibus.
- β. Rutabaga*, radice flavescente subglobosa.
3. B. RAPA (Lin. spec. 931.) foliis radicalibus lyratis polline glauco destitutis setoso-scabris, caulinis mediis incisus summis integerrimis lævibus. ② patria dubia. Mart. fl. rust. t. 49-50. DC. syst. 2. p. 590.* (v. v.)
- A. DEPRESSA*, radice sub collo tumida globoso-depressa, in radiculam tenuem subtus abruptè desinente. — Moris. ox. s. 3. t. 2. f. 1.
- B. OBLONGA*, radice oblonga sensim apice attenuata. — Moris. ox. s. 3. t. 2. f. 2.
- C. OLEIFERA*, radice exili. — Lobel. ic. 298. f. 1.
4. B. NAPUS (Lin. spec. 931.) foliis glabris polline cæcio glaucescentibus, radicalibus lyratis, caulinis pinnatifidis crenatisque, summis cordato-lanceolatis amplexicanalibus, siliquis divaricato-patentibus. ② Patria non satis nota. DC. syst. 2. p. 592.* (v. v.)
- A. OLEIFERA*, radice exili. — Lobel. ic. t. 200. f. 2. Colitur in arvis ad oleum è seminibus eliciendum, nunc sub Napellæ hybernæ, nunc sub Colzæ nomine. Sed vera Colza est B. campestris oleifera.
- B. ESCULENTA*, radice fusiformi, sub collo incrassata. — Lob. obs. 99. f. 2. Est Napus vera cum Napo-brassicâ et Rapâ oblongâ sæpè confusa; à priore differt tantum radice crassâ carnosâ esculentâ, sæpius albâ, nunc flavidâ seu nigricante.
- α. alba*. — Napus dulcis Blackw. herb. t. 410.
- β. flava*.
- γ. nigricans*.
5. B. PRÆCOX (Waldst. et Kit. ined. DC. syst. 2. p. 593.*) foliis glabris polline cæcio glaucescentibus, radicalibus caulinisque inferioribus lyratis, summis

- cordato-lanceolatis amplexicaulibus crenatis, siliquis erectis. ① culta in agris montosis Europæ mediæ. Semina minora quàm in Br. Napo. Scitur vere et semina matura æstate post Napum profert. DC. syst. 2. p. 593. (v. s.)
6. *B. CRATICA* (Lam! dict. 1. p. 747.*) caudice fruticoso, foliis ovato-subrotundis crenatis petiolatis lævibus. 5 in fissuris rupium præsertim maritimarum in Cretâ, in insulis Archipelagi et Græciâ. Sibth! et Smith! fl. græc. t. 645 ex prod. 2. p. 30. Flores racemosi Br. oleraceæ similes. Fructus ignoti. DC. syst. 2. p. 594.* (v. s. in h. Mus. Par. et Banks.)
7. *B. CHINENSIS* (Lin. amœn. 4. p. 280.*) herbacea, foliis ovalibus subintegerrimis, floralibus amplexicaulibus lanceolatis, calycibus nungue petalorum longioribus. ② in Chinâ. — Specimen herbarii Linnæani male cum descriptione Linnæi ipsius convenit. DC. syst. 2. p. 594.* (v. s.)
8. *B. VIOLACEA* (Lin! spec. 932.) herbacea, foliis lanceolato-ovatis glabris indivisis dentatis, sepalis ovarisque villosis, pedicellis bracteatis. ② in Chinâ. Petala ex albo purpurascens, reticulata, calyce duplò longiora, ferè Erucæ. DC. syst. 2. p. 595.* (v. s. in h. Lin.)
9. *B. MAGELLANICA* (Juss! ined. Pers! ench. 2. p. 207.) herbacea, foliis glabris pectinato-pinnatifidis, racemis florentibus subcorymbosis. — ad fretum Magellanicum. Petala ex sicco alba, limbo obovato vix magnitudine *B. Cheiranthi*. — DC. syst. 2. p. 595.* Deless! ic. sel. 2. t. 85. (v. s.)
10. *B. HELENICA* (Burch! fl. helen. ined. n. 128.) herbacea, foliis lyrato-pinnatifidis subscabris, radicalibus 9-10-jugis, caulinis 1-2-jugis, lobo terminali maximo. ① ② in insulæ Sanctæ Helenæ aquis. Fructus ignoti et idè species ex habitu solo ad Brassicam relata. An potiùs *Raphani* species? DC. syst. 2. p. 595.* (v. s. in h. Burch.)
11. *B. INCANA* (Ten! app. prim. hort. nap. cat. p. 59.) pube molli canescens, foliis inferioribus petiolatis lyratis, lobo terminali maximo dentato, caule basi suffruticoso, siliquis glabris rostro quintuplò longioribus. 5 in rupibus insulæ Capræarum, propè Neapolim et in Sicilia. DC. syst. 2. p. 596.* (v. s.)
12. *B. BALEARICA* (Pers. ench. 2. p. 206.) glabra, foliis inferioribus petiolatis sinuatis subpandoriformibus subcarnosis, caule basi suffruticoso, siliquis rostro quintuplò longioribus. 5 in insulis Balearicis, in rupe rubrà Nicææ. Deless! ic. sel. 2. t. 86. Br. sempervirens Schrank h. mon. t. 10. Species ex rostri junioris structurâ forsàn ad *Erucastrum* referenda. — DC. syst. 2. p. 596.* (v. s.)
13. *B. GRAVINÆ* (Ten! app. prima cat. h. nap. p. 59. fl. nap. t. 62.*) pubescenti-hispida, foliis inferioribus oblongis runcinatis dentibus lobulise integris acutis, calyce subpatulo, siliquis glabris. 5 in montibus Aprutii ulterioris. Species distinctissima. DC. syst. 2. p. 597.* (v. s.)
14. *B. PINNATIFIDA* (Desf! fl. atl. 2. p. 95. t. 165.*) foliis pinnatifidis glabris lobis lanceolatis serratis, costis petiolisque pilosiusculis, siliquis subtetragonis. — in Mauritaniâ et Hispaniâ. Flores magnitudine Br. oleraceæ. DC. syst. 2. p. 597.* (v. s. in h. Desf.)
15. *B. LYRATA* (Desf! fl. atl. 2. p. 96. t. 166.*) foliis hispidis, radicalibus lyratis, caulinis incisio-dentatis, calyce siliquisque hispidis, caulibus adscendentibus. — in Africâ bor. Habitus ferè *Diotaxis erucoidis*, sed siliqua diversissima. DC. syst. 2. p. 597.* (v. s. in h. Desf.)
16. *B. HUMILIS* (DC. syst. 2. p. 598.*) foliis radicalibus carnosissimis pinnatifidis, lobis integris apice sæpiùs piliferis, scapis nudis, siliquis patulis basi attenuatis stylo crassiusculo terminatis. 5 in Occitaniâ non longè à Monspelio. *Sisymbrium repandum* var. β . DC. cat. h. monsp. 62. (v. v.)
17. *B. REPANDA* (DC. syst. 2. p. 598.*) foliis radicalibus carnosissimis glabris repando-dentatis, scapis nudis, stylo tenui à siliquâ distincto. 5 in aprieis montium Delphinatûs, in Pedemontio et Galloprovinciâ. *Sisymbrium Monense* Vill! dauph. 3. p. 350. t. 39.* non Lin. Ad priorem valdè accedit ex habitu et *panululam* ex foliorum consistentiâ et formâ ad sequentem. (v. v.)

Sect. II. ERUCASTRUM. DC. syst. 2. p. 598.

Siliqua sessilis, rostro conico basi 1-2-spermo superata.

18. B. RICHERII (Vill. Delph. 71. dauph. 3. p. 331. t. 36.*) foliis glabris, inferioribus petiolatis oblongis subdentatis, summis lineari-lanceolatis paucis. 2 in asperis lapideosis Alpium Delphinatûs, Galloprovinciæ et Pedemontii. Flores flavi, sat similes B. oleraceæ. DC. syst. 2. p. 599.* (v. v.)
19. B. MONENSIS (Huds. angl. 291.) foliis glabris subcarnosis glaucescentibus pinnatipartitis, lobis linearibus distantibus subdentatis. 2 in Angliâ et in Scotiâ. Sisymbrium Monense. Lin. spec. ed. 1. p. 658. Smith! engl. bot. t. 962. Flores magni, citrini, venosi, exsiccatione pallescentes. — DC. syst. 2. p. 599.*
20. B. ERUCASTRUM (Lin. spec. 932.) foliis runcinatis subglabris, lobis inæqualibus obtusè sinuatis, caule basi hispido. ① in ruderalis arenosis Galliæ, Italiæ, Helvetiæ, etc. Bull. herb. t. 331. Sisymbrium erucastrum Vill. dauph. 3. p. 342.* Species quoad classificationem subdubia. DC. syst. 2. p. 600.* (v. s.)
21. B. CHEIRANTHOS (Vill. dauph. 3. p. 332. t. 36. ic. fruct.) foliis petiolatis hispidis pinnatipartitis, lobis sinuato-dentatis ovali-oblongis, caule basi hispido, siliquis rostro quadruplò longioribus. ② 2 in arenosis asperis et ad fluviorum ripas Galliæ austr., Pedemontii, Sabaudiz, Helvetiæ. Sinapis Tournefortii All. ped. n. 962. excl. syn. — DC. syst. 2. p. 600.* (v. v.)
 2. Sinapis recurvata All! ped. n. 963. t. 37. Forsan specificè distinguenda ob calycem laxiorem et rostrum sæpius asperum. — in collinis apricis Pedemontii et Mauritaniz.
2. Br. montana. — DC. fl. fr. 4. p. 651.* in montibus excelsioribus Alpium Pyrenæorumque. Humilis, subcaulis, cæspitosa.
22. B. CHEIRANTHIFLORA (DC. syst. 2. p. 601.*) foliis radicalibus petiolatis lyrato-pinnatilobatis subhispidis, caulinis paucis lobis integris acutis, radice exili, siliquis rostro triplò longioribus. ① in arenosis Hispaniæ, Galliæ. Raphanus cheiranthiflorus Willd. hort. berol. 19. t. 19.* omninò media inter B. cheiranthum et Br. Tournefortii (v. v.)
23. B. TOURNEFORTII (Gouan ill. p. 44. t. 20. f. A.*) foliis radicalibus petiolatis hirtis lyrato-pinnatipartitis, lobis ovatis crenatis, siliquis rostro duplò longioribus. ① ad agrorum margines subhumidos in Hispaniâ, in Ægypto. Eruca erecta Lag! cat. h. madr. 1815. p. 20. Flores pallidè flavi, exsiccatione albidii, parvi. — DC. syst. 2. p. 602.* (v. v. et s.)
 2. Br. sisymbroides Fisch! in litt. — in Persiâ. A priori tantùm differt habitu minore et foliis pinnatisectis, lobis subæqualibus, terminalibus nempe vix alios superantibus.
24. B. LÆVIGATA (Lag! varied. p. 40. n. 19. descr. ined. in litt.*) foliis radicalibus subpetiolatis hirtis pinnatifido-runcinatis, lobis acutè dentato-incisis, caule subundo glabro, siliquis rostro quadruplò longioribus. ① ② in arenosis Hispaniæ. DC. syst. 2. p. 602.* (v. s.)
25. B. VALENTINA (DC. syst. 2. p. 603.*) foliis inferioribus petiolatis hirsutis hispidis pinnatipartitis, lobis subdentatis, superioribus glabris subintegris, pedicellis calyce brevioribus, siliquis teretibus rostro triplò longioribus. ① in locis arenosis Hispaniæ circa Madritum. Sisymbrium Valentinum Lin. spec. 920. — Barr. ic. t. 195. f. 1. Specimen herbarii Linnæi valdè mancum et floribus destitutum, forsân ad Biscutellam ambiguum pertinet. (v. s.)
26. B? STRIGOSA (DC. syst. 2. p. 603.*) foliis inferioribus lyrato-pinnatifidis hispidis, lobis ovatis dentatis, terminali majore, caule basi retrorsum hispido, siliquis rostro quadruplò longioribus. — Ad cap. Bonæ-Spei. Sisymbrium strigosum Thunb. prod. 109. (v. s. in h. Banks.)
27. B? FRUTICULOSA (Cycl. pl. rar. 2. p. 7. t. 1.) foliis inferioribus subpilosissimis lyratis, lobis obtusis dentatis, terminali maximo, caule basi suffruticoso hispido, apice glabro, siliquis torosis rostro quadruplò longioribus. 2 5

in collibus aridis et in cultis circa Neapolim in Calabria, Græciâ et Bœticâ. *Sinapis radicata*. Desf. fl. atl. 2. p. 98. t. 167.* Ambigit inter *Sinapim* et *Brassicam* calyce nempe patente ut in *Sinapi*, habitu et fructu *Brassicæ*. — DC. syst. 2. p. 604.* (v. s.)

Sect. III. MICROPEDIUM. DC. syst. 2. p. 604.

Siliquæ suprà torum breviter pedicellatæ, stylo brevi tenui et stigmate capitato terminatæ, torosæ. — Aff. *Dyplotaxi* sed semina in loculis etiam juniora uniseriata videntur.

28. *B. ELONGATA* (Ehrh! Beitr. 7. p. 159.) foliis petiolatis, inferioribus sinuato-pinnatifidis hispidis, superioribus glabris dentatis, caule glabro. ②③ in Hungariâ, Transylvaniâ, Tauriâ et ad Caucasum. Waldst. et Kit. pl. rar. hung. 1. p. 26. t. 28.* Species quoad classificationem difficilis. Calyx nec clausus ut in *Brassicâ*, nec latus ut in *Sinapi*. DC. syst. 2. p. 604.* (v. s.)

29. *B. SABULARIA* (Brot. phyt. p. 97. t. 43.*) foliis radicalibus subpilis pinnatifidis runcinatisve, summis linearibus, caule basi hispido. ① in Lusitaniâ. *Sisymbrium* Parra Lin! mant. 255.* Specimen herb. Linnæi in horto Upsaliensi lectum optimè descriptioni Linnæanæ convenit. DC. syst. 2. p. 605.* (v. s.)

†. Species non satis notæ.

30. *Br. BUNIAS* (DC. syst. 2. p. 606.*) — verosimiliter in Oriente. Hæc stirps cum *Buniade* orientali verâ à Linnæo ipso confusa, et cum eâ in ejus herbario mixta, videtur diversissima. Siliquæ juniores teretes, graciles, glabræ. stylo filiformi; adultæ oblongæ, stylo conico aspermo superatæ. *Bunias* foliis retrorsum sinuatis Lin! hort. ups. 186. (v. s. in h. Lin.)

31. *B? HIRTA* (Moench. suppl. p. 85.*) ① hab. Folia glabra, lævia, subcarnosa. Petala obcordata emarginata, venis fuscis. Siliqua 4-6-sperma, *hirta*, stylo persistenti subpiloso siliquâ longiori. An potiùs *Erucæ* species? DC. syst. 2. p. 606.*

32. *B. FORSKAHLII* (Schnlt. obs. p. 131. n. 1012.) hab. Folia omnia duplicato-pinnatifida strigosa. Caulis calycesque hispidi. Siliquæ erectæ, glabræ. Petala viridi-flavescentia. DC. syst. 2. p. 606.

33. *B? PURPURASCENS* (Russ. in Schrad. journ. 1. p. 426.) Hab. Haleppo. Folia lanceolata, pinnatifida Caulis pilosus. Siliquæ articulatæ glabræ, rostro elongato subulato. An *Erucariæ* species? forsân *Erucaria Aleppica* ipsa? DC. syst. 2. p. 606.

34. *B? PSEUDO-ERUCASTRUM* (Brot. fl. lus. 1. p. 581.*) ① ② in Lusitaniâ bor. Folia radicalia semipedalia, lyrato-pinnatifida. Calyx erecto-clausus apice pilosus. Siliqua glabra, fusco-viridis, sesquunciam longa, stylo ensiformi superata. Semina globosa, nigra. An *Erucæ* species? DC. syst. 2. p. 606.

35. *B. MONTANA* (Raf. specch. 2. p. 69.) ⑤ in Siciliæ montibus Nebrodensibus. Fruticosa, foliis petiolatis lyrato-pinnatifidis oblongis inæqualiter dentatis carnosius glabris acutis, superioribus linearibus integris, petiolis compressis. An eadem ac *B. balearica*? DC. syst. 2. p. 714. †.

36. *B. CRISPA* (Raf. specch. 2. p. 69.) ⑤ in Siciliæ mont. Nebrodensibus. Suffruticosa, foliis petiolatis lyrato-pinnatifidis obovatis sinuatis crispis obtusius glabris, petiolis depressis, racemis simplicibus. DC. syst. 2. p. 714. †

LXXV. *SINAPIS*. Tourn. inst. 227. Lin. gen. n. 821. Gærtn. fruct. 2. p. 299. t. 143. DC. syst. 2. p. 607.

Siliqua teretiuscula, valvis nervigeris, stylo parvo brevi acuto. Semina uniseriata subglobosa. Calyx patens. — Genus auctorum hîc intactum adhuc retinui, sed verosimiliter in posterum dividendum.

Sect. I. MELANOSINAPIS. DC. syst. 2. p. 607.

Siliqua teres aut subtetragona. Stylus brevis parvus nec rostriformis.

25. *S. TURGIDA* (Delill! ill. fl. eg. n. 606.) siliquis glabris ovatis, valvis nervoso-reticulatis rostro conico subæqualibus. ① in Ægypto. Priori valdè aff. et cum ea interdum in hortis confusa. DC. syst. 2. p. 618.* (v. s.)

Sect. III. *HIRSCHFELDIA*. DC. syst. 2. p. 618.

Siliqua teres, bilocularis, loculo utroque sub-4-spermo, rostro ovato 1-spermo lævi indehiscente.

26. *S. INCANA* (Lin. amœn. 4. p. 281.* spec. 934.) siliquis glabris racemo adpressis subtorulosis, caule ramoso infernè scabro, foliis lyratis scabris. ② in Europâ austral. Jacq. hort. vind. t. 169. *Myagrum Hispanicum* Lin. spec. 893. Brassiceas et Raphaneas ferè conjungit, prioribus tamen proxima. — DC. syst. 2. p. 618.* (v. v.)
27. *S. HETEROPHYLLA* (Lag! cat. hort. madr. 1816.) siliquis pubescentibus rachi adpressis subtorulosis, caule subramoso basi hispido, foliis lyrato-pinnatifidis in nervis hispidis. ② in locis ruderalis circa Madritum. DC. syst. 2. p. 619.* (v. s.)

Sect? IV. *LEUCOSINAPIS*. DC. syst. 2. p. 620.

Siliqua hispida aut glabra, rostro ensiformi superata, valvulis torulosis. — An forsam cum *Eruca* conjungenda? An genus proprium à Sinapi ut *Eruca ipsa* à Brassicâ distinctum?

28. *S. ALBA* (Lin! spec. 933.) siliquis hispidis patentibus rostro ensiformi subangustioribus, foliis lyratis cauleque subglabris. ① inter segetes et in ruderalis Europæ austr. Lam. ill. t. 566. Smith engl. bot. t. 1677. Semina grossa, pallida. DC. syst. 2. p. 620.* (v. v.)
29. *S. HISPIDA* (Schousb. maroc. p. 182. t. 4.*) siliquis hispidis patentibus rostro ensiformi angustioribus, foliis lyratis scabris, caule retrorsum hispido. ① in imperio Maroccano, in arvis Teneriffæ. *S. flexuosa* Lam. dict. 4. p. 341.* Valdè aff. *S. albæ*, sed caulis retrorsum hispidus, folia scabra, siliquæ angustiores. DC. syst. 2. p. 621.* (v. s.)
30. *S. DISSECTA* (Lag! cat. hort. madr. 1816. p. 20.) siliquis suberectis torulosis rostro ensiformi subbrevioribus, foliis pinnatipartitis, lobis angustis inciso-dentatis pinnatifidisve. ① in Hispaniâ. DC. syst. 2. p. 621.* (v. s. et v.)
- α. siliquis glabris.
β. siliquis hispidulis.
31. *S. HASTATA* (Desf! cat. hort. par. ed. 2. p. 151.) siliquis suberectis glabris linearibus torulosis rostro ensiformi longioribus, foliis glabris pinnatilobatis, lobis lanceolatis. ① in Novâ-Hollandiâ. An hujus sectionis? DC. syst. 2. p. 622.* (v. s. in h. Desf.)
32. *S. FOLIOSA* (Willd. enum. 688.) rostro compresso scaberrimo siliquâ hispidâ longiore, foliis lyratis repando-angulatis glabris. ① in Oriente. Flores lutei. Fructus ignotus. DC. syst. 2. p. 622.* (v. v. sine fructu.)
33. *S. APULA* (Tenor. app. prima cat. hort. napol. p. 60.) siliquis lævibus patentibus, stylo oblongo compresso, foliis runcinato-pinnatifidis glabris, lacinis omnibus lanceolatis dentatis acutis. — in arvis Apuliæ. DC. syst. 2. p. 623.†.

Sect? V. *DISACCIUM*. DC. syst. 2. p. 623.

Calyx semipatens, basi bisaccatus. Sectio dubia et forsam genus proprium! sed mihi non satis notum ut stabilire queam.

34. *S. FRUTESCENS* (Ait. hort. kew. ed. 1. v. 2. p. 404. ed. 2. v. 4. p. 127.) calyce bisaccato, foliis coriaceis, inferioribus oblongo-lanceolatis basi attenuatis subdentatis. 5 in Madera inter rupes. Flores magnitudine Cheiranthis, forsam versicolores, juniores pallidè flavi. DC. syst. 2. p. 623.* (v. s. in h. Banks.)
35. *S. ANGUSTIFOLIA* (DC. syst. 2. p. 624.) calyce bisaccato, foliis linearibus

integris. $\frac{5}{8}$ in Maderà. *Brassica frutescens* Soland! in h. Banks. An à priori satis distincta? DC. syst. 2. p. 624. (v. s. sine fl. in herb. Banks.) †.

†. *Species non satis notæ.*

36. *S. PEKINENSIS* (Lour. fl. coch. ed. Willd. 2. p. 485.*) siliquis linearibus glabris, foliis obovatis integris crispis, petiolis subplanis latis. (1) Hab. Pekini culta. An *S. Chinensis*, brassicatæ aut junceæ varietas ex Lour? DC. syst. 2. p. 624.
37. *S. POLYMORPHA* (Genera. in Schult. obs. n. 1021.*) siliquis patulis lævibus, rostro subtetragono compresso, foliis lyrato-pinnatifidis glabris, nervis marginæque scabris, caule subtile hispido. Hab. DC. syst. 2. p. 624.*
38. *S. PROCUMBENS* (Poir. suppl. 4. p. 12.*) caule procumbente, foliis radicalibus lyrato-pinnatis expansis subglabris, caulinis superioribus simplicibus petiolatis, pedicellis inferioribus longissimis capillaceis. (1) ? in pratis siccis Mauritaniz. Foliis accedit ad *Brassicam fruticulosam*. Siliquæ ignotæ. DC. syst. 2. p. 624.*
39. *S. NUDICAULIS* (Lag. cat. hort. madr. 1816. p. 20.) scapo nudo, foliis radicalibus hispidis pinnatifidis, siliquis erectis lævibus cornu subulato longioribus. $\frac{2}{3}$ in Hispaniâ merid. Sat diversa ab omnibus congeneribus. (Lag.) DC. syst. 2. p. 625.
40. *S. MESOPOTAMICA* (Spreng. in Schrad. journ. 4. p. 199.) siliquis subtetragonis apice cuspidatis, foliis lyratis hispidis. (1) in Mesopotamiâ. DC. syst. 2. p. 625.

LXXVI. MORICANDIA. DC. syst. 2. p. 626.

Siliqua tetragona subanceps. Semina biseriata, ovata, parva subalata (ex Andr.) Calyx basi bisaccatus. — Flores purpurascens.

1. *M. ARVENSIS* (DC. syst. 2. p. 626.*) siliquis subtetragonis, foliis caulinis cordato-amplexicaulibus integerrimis (2) in Europæ austr. arvis. *Brassica arvensis*. Lin! mant. 95. — Bocc. sic. p. 49. t. 25. f. 3. 4. Flores violacei (v. v.) β . *Brassica suffruticosa*. Desf. fl. atl. 2. p. 94.* in Hispaniâ austr. Mauritaniz, etc. collibus siccis.
2. *M. HESPERIDIFLORA* (DC. syst. 2. p. 627.*) siliquis compressis, valvis planis medio 1-nerviis, foliis caulinis ovato-oblongis dentato-sinuatis. (1) in convallibus circa Kahiram frequens. Flores ferè *Hesperidis*, sordidè purpurascens. (v. s. comm. à cl. Delill.)
3. *M. TERETIFOLIA* (DC. syst. 2. p. 628.*) siliquis compressis, valvulis enerviis, foliis multifidis lobis filiformibus. (1) in Africâ bor. *Brassica teretifolia* Desf. fl. atl. 2. p. 94. t. 164.* Habitu foliorum ferè accedit ad *Erucariam Alepicam*. (v. s.)

LXXVII. DIPLLOTAXIS. DC. syst. 2. p. 628.

Siliqua compressa, linearis. Semina biseriata, ovata. Calyx basi æqualis. — Flores flavi aut albi. Calyces sæpè molliter pubescentes.

Sect. I. CATOCARPUM. DC. syst. 2. p. 629.

Stylus subnullus. Stigma bilobum ad apicem siliquæ sessile. Siliquæ pendulæ, sessiles aut sæpius pedicellatæ.

1. *D. CRASSIFOLIA* (DC. syst. 2. p. 629.*) siliquis pendulis pedicellatis, foliis dentatis ferè omnibus glabris. — In Siciliæ gypsaceis. *Sinapis crassifolia* Rafin. car. nov. gen. p. 78. n. 192.* Petala flava. Semina pallidè rufa, minima. (v. s. in herb. Schouw. et comm. à cl. Moricand.)
2. *D. LAGASCANA* (DC. syst. 2. p. 629.*) siliquis pendulis pedicellatis, foliis hinc indè piliferis pinnatipartitis, lobis distantibus paucis linearibus. (1) in locis siccis montanis Hispaniæ. *Sisymbrium pendulum*. Lag! in litt. Differre videtur et à priori et à posteriore foliis paucis, minoribus, pinnatipartitis, etc. (v. s. comm. à cl. Lag.)

3. *D. PENDULA* (DC. syst. 2. p. 630.*) siliquis pendulis pedicellatis, foliis caulinis oblongis hispidis grossè inciso-dentatis. — In Africâ bor. *Sisymbrium pendulum* Desf. fl. atl. 2. p. 82. t. 156.* Petala lutea calyce longiora.
4. *D. HISPIDA* (DC. syst. 2. p. 630.*) siliquis pendulis sessilibus, foliis obovatis grossè dentatis hispidis. ① in Ægypto. *Sisymbrium hispidum* Vahl. symb. 2. p. 77.* Flores flavi, mediocres, erecti. Deless! ic. sel. 2. t. 89. (v. s.)
β. subglabra, propè Damascum.

Sect. II. ANOCARPUM. DC. syst. 2. p. 630.

Stylus conico-compressus intus aut vacuus aut semina 1-2 gerens stigmatibus bilobis terminatus. Siliquæ erectæ sessiles aut rarius pedicellatæ.

5. *D. ERUCOIDES* (DC. syst. 2. p. 631.*) siliquis sessilibus erectiusculis, stylo ensiformi, foliis sessilibus runcinato-lyratis dentatis. ① in Hispaniâ merid. in ins. Balearicis, Niceæ, in Sicilia et in oriente. *Sinapis erucoides* Linl amœn 4. p. 322.* Jacq. hort. vind. t. 170. Petala alba aut ad unguem purpurascens ampla obtusissima. Semina pallida, minima. (v. v. et s.)
6. *D. VIRGATA* (DC. syst. 2. p. 631.*) siliquis sessilibus erectiusculis, stylo ensiformi, foliis petiolatis pinnatifidis, lobis obtusis dentatis. ① in ruderatis Hispaniæ. *Sinapis virgata* Cav. præl. ex Lag! et Duff! Planta à spithamâ ad 4 et amplius pedes alta, floribus luteis et foliis petiolatis à *D. erucoides* facile distinguenda. (v. s.)
7. *D. CATHOLICA* (DC. syst. 2. p. 632.*) siliquis sessilibus erectiusculis, stylo subtereti 1-2-spermo, foliis pinnatipartitis, lobis dissectis, lacinii linearibus sinuato-dentatis. ① in Lusitaniâ et Hispaniâ. *Sisymbrium catholicum* Linl mant. 93.* Petala lutea, unguibus calyce brevioribus. Semina parva rufa. (v. s.)
8. *D. TENUIFOLIA* (DC. syst. 2. p. 632.*) siliquis subpedicellatis erectis, stylo filiformi brevi aspermo, foliis supremis integris, infimis pinnatipartitis, lobis linearibus integris pinnatifidisve. ② in glareosis apricis Europæ. *Sisymbrium tenuifolium* Lin. spec. 917. Bull. herb. t. 335. Smith engl. bot. t. 525. (v. v.)
9. *D. MURALIS* (DC. syst. 2. p. 634.*) siliquis sessilibus erectis, stylo brevi subfiliformi, foliis radicalibus dentatis lyratisve glabris, caulibus subnudis adscendentibus. ① ② in Angliâ, Galliâ, Germaniâ, Austriâ et Transylvaniâ. etc. *Sisymbrium murale* Linl spec. 918. Smith engl. bot. t. 1090. Flores minores et pallidiùs flavi quàm in priore. Siliquæ ferè similes. (v. v.)
β. foliorum lobis profundioribus acutioribus dentatis. Sisymbrium erucastrum Gouan. ill. p. 42. t. 20.
γ. minor, foliorum lobis profundioribus obtusis. Sisymbrium Barrelieri Thuill! fl. par. ed. 2. v. 1. p. 334.
δ. pygmaeum. Sisymbrium monense Thuill. fl. par. ed. 2. v. 1. p. 333. ex Mer. fl. par. 259.
10. *D. BARRELIERI* (DC. syst. 2. p. 634.*) siliquis sessilibus erectis, stylo brevi subfiliformi, foliis radicalibus runcinatis dentatis hispidis, caulibus nudis erectis superne glabris. ① in Hispaniâ, in Italiâ? *Sisymbrium Barrelieri* Linl spec. 919. — Barr. obs. p. 44. t. 100. bona. Flores flavi, in sicco albescentes. (v. s. in h. Linn.)
11. *D. SCAPOSA* (DC. syst. 2. p. 635.*) siliquis sessilibus erectis, stylo brevi subfiliformi, foliis radicalibus glabris oblongis pinnatifidis, lobis brevibus integris, caulibus nudis brevissimis. — In insulâ Lampedosâ. *Herba pygmae, glabra. Semina orbiculata, parva.* — Deless! ic. sel. 2. t. 90. (v. s.)
12. *D. VIMINEA* (DC. syst. 2. p. 635.*) siliquis sessilibus erectis, stylo brevi filiformi, foliis radicalibus lyratis obtusissimis lævibus, caulibus nudis subdecumbentibus. ① in arenosis Galliæ, in Sicilia. *Sisymbrium vimineum* Linl spec. 919. — Bocc. sic. 19. t. 10. Planta parva, glabra, *D. murali* affinis. Flores lutei, parvi. (v. s.)
13. *D. SAXATILIS* (DC. syst. 2. p. 636.*) siliquis erectis sessilibus basi atte-

nuatis, stylo brevi conico, foliis radicalibus pinnati-lobatis crassiusculis, lobis integris, caulibus subnudis erectis. ♀ in collinis Europæ austr. *Sisymbrium Monense*. Lin! spec. ed. 2. p. 918. excl. syn. Dill. non ed. 1. Species inter *Diploctaxis* et *Brassicam* ferè media. (v. s.)

14. *D. RAMOSISSIMA* (Spreng. neu. entd. 3. p. 52.) siliquis erectis, stylo brevi tereti aspermo, foliis omnibus lanceolatis elongatis subsessilibus glaberrimis integerrimis. ♀ patr. ign. Siliquæ, ex cl. auct., teretes longè pedicellatæ. Semina non descr.

LXXVIII. *ERUCA*. *Tourn. inst. p. 227. t. 111. DC. syst. 2. p. 636.* — *Euzomum*. *Link enum. 2. p. 174.*

Siliqua teretiuscula, stylo amplo conico aut ensiformi superata. Semina 1-seriata globosa. Calyx erectus, basi æqualis. — Petala alba aut flava, venis fuscis anastomosantibus reticulata.

1. *E. SATIVA* (Lam. fl. fr. 2. p. 496.) foliis lyrato-pinnatipartitis, lobis dentatis acutis, caule hirsuto, pedicellis calyce deciduo brevioribus. ① inter segetes Europæ austr. et Africæ bor. *Brassica Eruca* Lin! spec. 932 — Lob. icon. t. 204. f. 1. An plures species hic confusæ? DC. syst. 2. p. 637.* (v. v.)

α. flore albo, fructibus glabris. *Brassica Eruca* Blackw. herb. t. 242.

β. flore albo, fructibus pilosis.

γ. flore flavo, fructibus glabris. *Brassica Eruca* Bull. herb. t. 313. Schkh. handb. 2. n. 1870. t. 186.

δ. flore flavo, fructibus pilosis. *Brassica Eruca* var. β. Sibth. et Smith fl. græc. t. 646 et 647. ex prod. 2. p. 30.

ε. flore pallido, fructibus glabris turgidis. *Brassica turgida* Pers. ench. 2. p. 207.

ζ. caule glabro, siliquis hispidis. *Brassica erucoïdes* Horn. h. hafn. 2. p. 621.

ζ. caule siliquisque glabris. *Sinapis exotica* hortul.

2. *E. HISPIDA* (DC. syst. 2. p. 638.*) foliis lyrato-pinnatipartitis, lobis dentatis, terminali obtuso, caule hispido, pedicellis calyce deciduo longioribus. ① in regno Neapolitano. Petala alba fusco-venosa. Fructum non vidi. — An satis à priore distincta? (v. s.)

3. *E. VESICARIA* (Cav! ex Lag. DC. syst. 2. p. 638.*) foliis pinnatifidis, lobis acutis subintegris, caule hirsuto, calyce persistente subvesicario. ① Aff. *E. sativæ* sed differt foliis magis elongatis, ad costam usque rarè incisis, etc. (v. s.)

α. *albiflora*. *Brassica vesicaria* Asso syn. arrag. 88. t. 4. in Arragoniâ.

β. *flaviflora*. *Brassica vesicaria* Lin! spec. 933. in Hispaniæ ferè totius arvis.

Trib. XIII. *VELLEÆ* seu *ORTHOPELOCEÆ LATI-SETPÆ*. DC. syst. 2. p. 639.

Silicula valvis concavis longitudinaliter dehiscentibus, septo elliptico. Semina globosa. Cotyledones conduplicatæ.

LXXIX. *VELLA*. DC. syst. 2. p. 639. — *Vellæ* sp. Lin.

Stamina majora connata. Stylus ovatus planus ad apicem siliculæ linguiformis.

1. *V. PSEUDOCYTISUS* (Lin! spec. 895.) ♂ in Hispaniæ collibus gypsaceis. Lam. ill. t. 555. f. 2. Petala longè unguiculata flava cum ungue atropurpureo. Stamina majora per paria omnino connata. DC. syst. 2. p. 640.* (v. s. et v.)

LXXX. *BOLEUM*. *Desv. journ. 3. p. 163. t. 26. DC. syst. 2. p. 640.*

Stamina majora connata. Stylus tenuis subconicus ad apicem siliculæ rostriformis.

1. *B. ASPERUM* (Desv! Journ. bot. 3. p. 163 et 175. t. 6.*) ♂ in Hispaniæ asperis. Vella aspera Pers! ench. 2. p. 185. Petala ochroleuca vel albida. DC. syst. 2. p. 641. (v. s. in herb. Juss.)

LXXXI. CARRICHTERA. DC. syst. 2. p. 641.

Stamina omnia libera. Stylus ovatus planus, foliaceus. — Flores parvi ochroleuci. Siliculæ pedicello inflexo pendulæ.

1. *C. VELLÆ* (DC. syst. 2. p. 642.*) ① ad muros Valentiae, in insulis Balearicis, in Sicilia, Græciâ, Syriâ, etc. Vella annua Lin! spec. 895. Gærtn. fruct. 2. p. 886. t. 141. Semina in quoque loculo 4. globosa, parva, fusca. (v. s.)

LXXXII. SUCCOWIA. Medik. gen. pl. 1. p. 64. t. 1. f. 9. DC. syst. 2. p. 642.

Stamina omnia libera. Stylus tenuis conicus. Siliculæ valvulæ echinatae.

2. *S. BALEARICA* (Medik. in Ust. neu. ann. 1. p. 41.) ① in insulâ Teneriffæ, in insulis Balearicis, in Sicilia. Flores parvi, flavi. Semina in quoque loculo solitaria, pendula, globosa, submaculata. DC. syst. 2. p. 643.* (v. v. et s.)

Trib. XIV. *PSYCHINEÆ* seu *ORTHOPLCEÆ* *ANGUSTISEPTÆ*. DC. syst. 2. p. 643.

Silicula valvis carinatis navicularibusve, septo angustissimo. Semina compressa. Cotyledones conduplicatae. — Tribus distinctissima fructum ferè Thlaspeos aut Capsellæ aut Æthionematis cum flore et cotyledonibus Brassicearum gerens.

LXXXIII. SCHOUWIA. DC. syst. 2. p. 643.

Silicula ovalis, valvis ad dorsum per totam longitudinem angustè alatis.

1. *S. ARABICA* (DC. syst. 2. p. 644.*) ① in Arabiæ felicis argillaceis. Subularia purpurea Forsk! fl. ægypt. arab. p. 117. Folia Brassicæ perfoliatae satè referentia. Flores primùm corymbosi, dein racemosi. (v. s. in herb. Forsk.)

LXXXIV. PSYCHINE. Desf. fl. atl. 2. p. 69. t. 148. DC. syst. 2. p. 645.

Silicula triangularis basi angustata, valvis ad apicem tantùm dorso alatis.

1. *P. STYLOSA* (Desf! fl. atl. 2. p. 69. t. 148.*) ① in Mauritaniâ. Thlaspi Psychine Willd. spec. 3. p. 443. Folia caulina basi cordato-semi-amplexicaulia. Silicula magna, excepto stylo 3 lin. longa. DC. syst. 2. p. 645.* (v. s.)

Trib. XV. *ZILLÆ* seu *ORTHOPLCEÆ* *NUCAMENTACEÆ*. DC. syst. 2. p. 646.

Silicula indehiscens, ovata aut globosa, 1-locularis, 1-sperma, valvis indistinctis. Semina globosa. Cotyledones conduplicatae.

LXXXV. ZILLA. Forsk. descr. 121. DC. syst. 2. p. 646.

Silicula bilocularis, loculis 1-spermis. — Fructus ferè Crambes, spinæ ferè Alyssi spinosi.

1. *Z. MYAGROIDES* (Forsk. æg. descr. 121. n. 74 et 75. icon. t. 17. A.) ♂ DC. syst. 2. p. 646.*

a. *macrocarpa*, siliculis lævibus. Bunias spinosa Lin! mant. 96. Gærtn. fruct. 2. p. 290. t. 142. f. 2. — Circa Calhiram.

β. microcarpa, siliculis rugoso-costatis. — in *Ægypto superiore*.

LXXXVI. MURICARIA. *Desv. journ. bot. 3. p. 159. t. 25. f. 2. DC. syst. 2. p. 647.*

Silicula 1-locularis 1-sperma. Semen lateraliter insertum. Petala æqualia. — Generis in ordine affinitas, ignotis cotyledonibus, paulò dubia.

1. *M. PROSTRATA* (Desv! l. c.) — In arenis Africæ bor. *Bunias prostrata* Desf! fl. atl. 2. p. 76. t. 150.* Petala alba calyce duplò longiora. Stylus brevissimus crassiusculus. DC. syst. 2. p. 647.* (v. s. in h. Desf.)

LXXXVII. CALEPINA. *Adans. fam. 2. p. 423. DC. syst. 2. p. 648.*

Silicula 1-locularis 1-sperma. Semen ex apice pendulum. Petala exteriora submajora. Cotyl. incumbentes curvatæ truncatæ marginibus introflexis subconduplicatæ.

1. *C. CORVINI* (Desv. journ. bot. 3. p. 158.) ① in arvis et ruderatis Arragoniæ, Galliæ, Italiæ, Hungariæ, Tauriæ. *Bunias cochlearioides* Murr. nov. comm. goett. 8. 1777. p. 42. t. 3. Flores albi. (v. v.)

β. Myagrum iberioides Brot. phyt. n. 43. p. 95. t. 42. — in Lusitaniâ.

Trib. XVI. *RAPHANÆÆ* seu *ORTHOPELOCEÆ LOMENTACEÆ*. DC. syst. 2. p. 649.

Silicula aut siliqua transversè in articulos loculosve mono aut oligospermos secedens aut divisa. Semina globosa. Cotyledones conduplicatæ.

LXXXVIII. CRAMBE. *Tourn. inst. 211. t. 100. Gaertn. fruct. 2. p. 292. t. 142.*

Silicula biarticulata, articulo inferiore abortivo, superiore globoso 1-spermo. Cotyledones crassæ, subfoliaceæ, profundè emarginatæ. — Flores albi.

Sect. I. *SARCOCRAMBE*. DC. syst. 2. p. 651.

Silicula articulus inferior depressus crassus. Stigma sessile; filamenta majora hinc dentata. — Radix perennis ad collum multiceps.

1. *C. MARITIMA* (Lin! spec. 937.) filamentis longioribus bifurcis, silicula mutica, foliis subrotundis sinuatis undulatis dentatis glaucis cauleque glaberrimis. ½ in arenosis maritimis secus Oceanum, Mediterraneum et Pontum Euxinum. OEd. fl. dan. t. 316. Smith engl. bot. t. 924. DC. syst. 2. p. 651.* (v. v.)

2. *C. PINNATIFIDA* (Brown in hort. kew. ed. 2. v. 4. p. 72.) filamentis longioribus bifurcis, silicula mutica, foliis pinnatifidis, lobis oblongis acutis dentatis cauleque glabris. ½ in campis Hungariæ, Tauriæ et in desertis Caucasi. Cr. Orientalis Jacq. ic. rar. 1. t. 128. Species ærè media inter maritimam et Orientalem. DC. syst. 2. p. 652.* (v. v. et s.)

3. *C. GRANDIFLORA* (DC. syst. 2. p. 652.*) filamentis longioribus bifurcis, silicula mutica, foliis pinnatifidis, lobis oblongis acutè inciso-dentatis, minoribus secus rachin intermixtis, cauleque glabris. ½ in insulâ Taman. Media inter Cr. pinnatifidam et C. asperam. Fructus ignoti et species ideò adhuc recognoscenda; an Cr. maritimæ var? DC. syst. 2. p. 652.* (v. s.)

4. *C. ORIENTALIS* (Lin! spec. 937.) filamentis longioribus bifurcis, silicula mutica lævi, foliis pinnatifidis dentatis scabris, caule glabro. ½ in Oriente. Species forsàn non ritè definita et duas stirpes inter se diversas colligens. DC. syst. 2. p. 653.* (v. v. et s.)

5. *C. TATARICA* (Jacq. misc. 2. p. 274.* icon. rar. 1. t. 129. excl. syn. Clus.)

filamentis longioribus bifurcis, silicula mutica, foliis radicalibus decompositis, pinnulis dentato-incisis, junioribus scabris, adultis cauleque glabriusculis. \mathcal{L} DC. syst. 2. p. 653.* (v. s.)

α . *Hungarica*, foliorum caulinorum lobis oblongo-linearibus subintegrè subpectinatis. C. Tataria Jacq! misc. 2. p. 274.* — In agris Hungariæ.

β . *Taurica*, foliorum lobis sinnatis lobalis. C. Orientalis Reich. syst. 3. p. 290. Falk. it. p. 219. n. 791. t. 14. ex Bieb. — In campis Tauriæ.

6. *C. ASPERA* (Bieb. fl. taur. 2. p. 90.) filamentis longioribus bifurcis, silicula mutica rugosa, foliis pinnatipartitis, lobis oblongo-linearibus dentatis cauleque scabris. \mathcal{L} in campis Tauriæ. Planta etiam fructifera pilis crebris erectis rigidis undiquè hispida. DC. syst. 2. p. 654.* Deless! ic. sel. 2. t. 91. (v. s.)

7. *C. JUNCEA* (Bieb. suppl. ined. ex Fisch! et Stev! DC. syst. 2. p. 654.*) filamentis longioribus bifurcis, silicula mutica lævi, foliis lyratis, lobis dentatis, terminali lobato maximo, petiolis cauleque retrorsum hispidis. \mathcal{L} in Iberiâ circa Tiflim. Silicula pedicello tenui super-imposita, globosa, læves, piso triplò minores. (v. s.)

8. *C. CORDIFOLIA* (Stev! mem. soc. nat. mosc. 3. p. 267.) staminibus longioribus bifurcis, siliculâ submuticâ, foliis petiolatis dentatis, inferioribus cordatis, superioribus ovatis cauleque subglabris. \mathcal{L} in desertis ad promontorium Caucasi sept. Cr. cordata Willd. enum. suppl. 43. Panicula aphylla ramosissima glabra. Flores C. orientalis. — DC. syst. 2. p. 654.* (v. v. et s.)

Sect. II. LEPTOGRAMBE. DC. syst. 2. p. 655.

Siliculae articulus inferior cylindricus elongatus. Stigma sessile; filamenta dentata aut edentula. — Radix annua aut biennis. Caulis solitarius.

9. *C. HISPANICA* (Lin! spec. 937.) filamentis longioribus dente auctis, siliculâ muticâ, foliis lyratis scabris, lobo terminali cordato orbiculato. ① in Hispaniâ. Lam. ill. t. 553. Gaertn. fruct. 2. p. 292. t. 142. f. 4. Petala alba, silicula exactè globosa, omninò mutica. DC. syst. 2. p. 655.* (v. v. et s.)

10. *C. GLABRATA*, filamentis longioribus dente auctis, siliculâ muticâ, foliis cordato-rotundatis cauleque glabris. — In Hispaniâ propè Sanctum Philippum. Cr. cordifolia Dufour in ann. gen. sc. phys. 7. p. 308.* non Stev. An à Cr. Hispanicâ satis differt?

11. *C. RENIFORMIS* (Desf. fl. atl. 2. p. 78. t. 151.*) filamentis longioribus dente auctis, siliculâ muticâ, foliis lyratis scabris, lobo terminali reniformi. ② in fissuris rupium Atlantis. Willd. spec. 3. p. 419. Silicula carnosa, Cr. Hispanicæ similis. DC. syst. 2. p. 656.*

12. *C. FILIFORMIS* (Jacq. coll. suppl. 120.* ic. rar. 3. t. 504.) filamentis longioribus subdentulis, siliculâ muticâ, foliis pinnato-lyratis pilosis, lobo terminali ovato. \mathcal{L} ② in Patagoniâ ad Champion River. Rapistrum filiforme Mœnch. suppl. 69.* Ovarium gracile biarticulatum, articulo inferiore tereti longiore. DC. syst. 2. p. 656.* (v. v.)

Sect. III. DENDROGRAMBE. DC. syst. 2. p. 656.

Siliculae articulus inferior filiformis brevis. Stigma stylo brevi superimpositum. Caulis fruticosus.

13. *C. FRUTICOSA* (Lin. f. suppl. 299.) filamentis longioribus hinc dente auctis, siliculâ stylo mucronatâ, foliis lyrato-pinnatifidis dentatis canis, caule fruticoso. \mathcal{L} in Maderæ rupibus altissimis. Willd. spec. 3. p. 420. Silicula articulata, articulo inferiore abortivo filiformi, stipitata. (v. s.)

14. *Cr. STRIGOSA* (L'her! stirp. 1. p. 151. t. 72.*) filamentis edentulis, siliculâ stylo mucronatâ, foliis ovatis dentatis basi inæqualibus subauriculatis caule fruticoso hispidis. \mathcal{L} in insulis Canariis. Myagræm arboreascenti Jacq. coll. 1. p. 39.* ic. rar. t. 120. — DC. syst. 2. p. 657.* (v. s.)

β . *glabrata*, à priori non differt nisi foliis ferè glabris, strigis paucissimis lepidotis.

LXXXIX. RAPISTRUM. Boerh. lugd. 406. DC. syst. 2. p. 430.

Silicula biarticulata, articulo superiore ovato rugosø. Semina in loculis solitaria, superioris erectum, inferioris pendulum. Cotyledones oblongæ conduplicatæ ex Andr. in litt. — Flores flavi.

1. *R. PERENNE* (Berg. phyton. ic. Desv. journ. 3. p. 160.) siliculis glabris, articulo superiore ovato, stylo longiore, foliis pinnatifidis, lobis dentato-incisis acutis. \mathcal{L} in agris Galliæ orient. Helvetiæ, Pedemontii, Germaniæ, Austriæ, Liburni et Transylvaniæ. Myagr. perenne Lin! spec. 893. Jacq. fl. austr. t. 414. Flores parvi, lutei. — DC. syst. 2. p. 431.* (v. s.)

2. *R. RUGOSUM* (Berg. phyt. 3. p. 171. icon.) siliculis pubescentibus, articulo superiore globoso rugoso stylo brevior, foliis obtusis dentatis radicalibus sublyratis. \textcircled{I} in agris et arenosis Europæ mediæ et austr. Myagr. rugosum Lin! spec. 893. *R. rugosum* All. ped. t. 78. — DC. syst. 2. p. 432.* (v. v.)

β . *stylosum*. *Cakile rugosa* var. *stylosa* DC. fl. fr. suppl. p. 598. in Occitania et Galloprovincia australi.

γ . *venosum*. *Myagr. venosum* Pers. ench. 2. p. 183. Circa Tolosam et Genuam.

† *Species non satis notæ.*

3. *R. ORIENTALE* (DC. syst. 2. p. 433.*) siliculis sulcatis lævibus, foliis oblongis dentato-sinuatis. \textcircled{I} in Oriente. Myagr. orientale Lin! spec. 893. *Bunias raphanifolia* Sibth. et Smith fl. græc. t. 612. ex prodr. 2. p. 2. Flores flavi paulò minores quàm *R. perennis*. (v. s.)

4. *R. CLAVATUM* (DC. syst. 2. p. 433.*) siliculæ glabræ articulo superiore globoso rugoso stylifero, inferiore lævi subcylindraceo pedicello longiore. \textcircled{I} in Syria circa Tripoli. Flores ignoti. Siliquæ etiam nascentes glabræ. (v. s.)

5. *R. COSTATUM* (DC. syst. 2. p. 434.*) siliculæ glabræ articulo superiore ovato costato in acumen conicum desinente, inferiore lævi obconico pedicello subbreviore. \mathcal{L} ? in Oriente ad Tyram. Myagr. perenne Stev! in litt. Fructum tantum vidi missum à cl. Stev. Ex specimine distincta mihi videtur species nisi forsitan sit eadem ac *R. orientale*. (v. s. sine fl. nec fol.)

XC. DIDESMUS. Desv. journ. bot. 3. p. 160. t. 25. f. 11. DC. syst. 2. p. 657.

Silicula biarticulata, articulo utroque 1-2-spermo, inferiore apice truncato, superiore stylifero. Semina in quoque loculo pendula.

1. *D. AEGYPTIUS* (Desv. l. c.) foliis inferioribus ellipticis aut pinnatifido-lyratis, summis oblongis subdentatis. \textcircled{I} in Aegypto, in insulâ Cypro. Myagr. Aegyptium Lin! spec. 895. — Deless! ic. sel. 2. t. 92. DC. syst. 2. p. 658.* (v. s. in h. Lin.)

β . *oblongifolius*, à priori differre videtur foliis inferioribus oblongis grosse serratis non pinnatifido-lyratis. — in campis Astypalæ. (v. s. com. à cl. D'Urville.)

2. *D. TENUIFOLIUS* (DC. syst. 2. p. 659.*) foliis pinnatifidis, lobis linearibus distantibus, summis linearibus, caule glabro. \textcircled{I} in Archipelago Græco et Cypro. *Bunias tenuifolia* Smith prodr. fl. græc. 2. p. 3. — Tourn. voy. 1. p. 254. ic. (v. s. in h. Banks.)

3. *D. BIPINNATUS* (DC. syst. 2. p. 659.*) foliis pinnatifidis, lobis linearibus dentatis aut incisis, caule retrorsum scabro. \textcircled{I} in montibus Cafzæ Mauritanie. *Sinapis bipinnata* Desf! fl. atl. 2. p. 97.* Petala in sioco albida, limbo obovato retuso ferè submarginato. (v. s. in h. Desf.)

† *Species non satis nota.*

4. *D. PINNATUS* (DC. syst. 2. p. 659.) Myagr. pinnatum Russel in Schrad. journ. 1. p. 426. — in Aleppo. Siliculæ biarticulatæ, striatæ; folia pinnata, glabra, laciniis linearibus incisis. An eadem ac *D. tenuifolius*?

XCI. ENARTHROCARPUS. *Labill. syr. dec. 5. p. 4. t. 2. DC. syst. 2. p. 660.*

Siliqua biarticulata, articulo inferiore obconico brevi 1-3-spermo, superiore longo 9-10-spermo, intus loculato. — Flores flavi, venosi aut purpurascens.

1. *E. arcuatus* (Labill! syr. dec. 5. p. 4. t. 2.*) siliquis teretibus torulosis scabris, pedicellis infimis bracteatis. ① in Cretâ, in Libano. Sepala oblongo-linearia, margine hispidula. DC. syst. 2. p. 660.* (v. s.)
2. *E. lyratus* (DC. syst. 2. p. 661.*) siliquis compressis nodulosis longitudinaliter striatis, pedicellis ferè omnibus bracteatis ① inter hordeaceas segetes circa Alexandriam. Raphanus recurvus Pers. ench. 2. p. 209. Delil. fl. ægypt. p. 105. t. 36. f. 1.* Valdè similis *E. arcuato* sed flores flavi purpureo-venosi, nec purpurascens. (v. s.)
3. *E. pterocarpus* (DC. syst. 2. p. 661.*) siliquis compressis alâ marginatis longè acuminatis, pedicellis ferè omnibus bracteatis. ① in Ægypto. Deless! ic. sel. 2. t. 93. Raphanus pterocarpus Pers! ench. 2. p. 209. (v. s.)

XCII. RAPHANUS. *Lin. gen. n. 1098. DC. syst. 2. p. 662.*

Siliqua transversè multilocularis aut in articulos plurimos secedens. Semina 1-serialia, globosa, pendula. Cotyledones crassiusculæ, con-
duplicatæ.

Sect. 1. RAPHANIS. *DC. syst. 2. p. 663.*

Siliqua fungosa bilocularis isthmis rarissimè coarctata.

1. *R. sativus* (Lin! spec. 935.) siliquis teretibus torosis acuminatis vix pedicello longioribus. ① ② in Chinâ, Japoniâ et Asiâ occid. Lam. ill. t. 566. Species in hortis vulgatissima. DC. syst. 2. p. 663.* — Stirpes plurimæ sub hoc nomine apud botanicos ferè omnes collectæ sunt, sed valdè dubium an sint meræ varietates aut species propriæ.

A. radicula, radice plus minùs carnosa, alba rosea aut rubra. *R. radicula* Pers. ench. 2. p. 208. in adnot. DC. syst. 2. p. 663.*

a. rotunda, radice subglobosa. *R. rotundus* Mill. dict. n. 2. — Weinm. phyt. t. 860. f. e. Variat ex radice colore nunc intus extusque albo, roseo et violaceo, pallidè roseo tenerrimo.

β. oblonga, radice oblonga carnosa. *R. sativus* Mill. dict. n. 1. — Weinm. phyt. t. 860. f. a. Variat ex radice forma et colore albo, roseo et purpureo.

γ. oleifera, radice exili elongata vix subcarnosa. *R. Chinensis* Mill. dict. n. 5.

B. niger, radice carnosa compacta duriuscula sapore acerrima. DC. syst. 2. p. 665.*

a. vulgaris, radice extus nigra, intus alba, oblonga. *R. niger* Merat fl. par. 265. — Moris. oxon. 2. p. 265. n. 2. s. 3. t. 13. f. 2. hæc varietas sæpè cum sequente forsitan non immerito confusa.

β. rotundus, radice extus nigra, intus alba, subrotunda. — Moris. oxon. 2. p. 265. n. 1. s. 3. t. 13. f. 1. Au varietas prioris?

γ. griseus, radice extus grisea. *Petit raifort gris* Bosc. dict. agr. 11. p. 40.

δ. albus, radice extus intusque alba depressa-globosa. *R. orbicularis* Mill. dict. n. 3.

2. *R. caudatus* (Lin! mant. 95. Lin. f. dec. 1. t. 10.*) siliquis depressis acuminatis decumbentibus plantâ totâ longioribus. ① in Javâ. Differt à *Raphanide* siliquis unilocularibus, à *Raphanistro* siliquis non torulosis; ab utroque calyce basi magis bisaccato et seminibus oblongis. DC. syst. 2. p. 665.* (v. s. in h. Banks, Lamb. Smith.)

Sect. II. RAPHANISTRUM. DC. syst. 2. p. 666. — Gaertn. fruct. 2. p. 300. t. 143.

Siliqua teres, acuminata, per maturitatem moniliformis unilocularis, articulis 1-spermis.

3. R. ROSTRATUS (DC. syst. 2. p. 666.*) siliquis unilocularibus articulis striatis 3-4-spermis, stylo conico-subulato brevioribus. — In Persiâ bor. R. raphanistrum var. Fisch! in litt. Specimen valdè mancum et incompletum antè oculos habeo, sed nullo modo cum R. raphanistro confundendum. (v. s.) †.
4. R. RAPHANISTRUM (Lin! spec. 953.) siliquis unilocularibus articulatis striatis 3-8-spermis stylo longioribus, foliis simpliciter lyratis. ① inter segetes totius Europæ. OEd. fl. dan. t. 678. excl. nom. Smith engl. bot. t. 856. DC. syst. 2. p. 666.* (v. v.)

α. flore albo, lineis nigricantibus plus minus striato. — Weinm. phyt. t. 862. f. a.

β. flore purpurascens. — Weinm. phyt. t. 862. f. b.

γ. flore flavo. — J. Bauh. hist. 2. p. 844. ic.

5. R. LANDRA (Moretti! fl. insubr. ined.* DC. syst. 2. p. 668.*) siliquis 1-locularibus articulatis substriatis 2-6-spermis stylo subulato longioribus, foliis interruptè lyratis. ② ʒ inter segetes in Italiâ Insubricâ. Corolla tota flava. Deless! ic. sel. 2. t. 94. (v. s.)

6. R. MARITIMUS (Smith engl. bot. t. 1643.) siliquis 1-locularibus articulatis striatis 2-6-spermis, stylo conico articulo ultimo ferè brevior, foliis interruptè lyratis. ʒ ② in maritimis Angliæ et circa Brest. Flores flavi, vix venosi. DC. syst. 2. p. 668.* (v. s.)

†. Species non satis notæ.

7. R? CAKILOÏDEUS (DC. syst. 2. p. 669.*) siliquis submoniliformibus scabris, foliis crassiusculis pinnatipartitis, rachi lineari, lobis linearibus obtusè dentatis. ① in Oriente. Corolla pallide violacea, ferè Erucariz Alepicæ. — An potius Erucariz species? (v. s.)

8. R. LANCEOLATUS (Willd. spec. 3. p. 562.*) siliquis bilocularibus ventricosis lævibus, rostro subtetragono, foliis oblongo-lanceolatis apice subdentatis. ①? in Antillis. Sinapis integrifolia West. st. cruc. p. 222? ex Willd. Flores mediocres flavi. DC. syst. 2. p. 669.* (v. s. sine fruct.)

9. R? BIARTICULATUS (Willd. enum. suppl. p. 46.) siliquis suberosis crassis biarticulatis, articulis 1-locularibus monospermis, foliis inferioribus lyratis, superioribus oblongis dentatis. ① in Chinâ. An potius Rapiatri aut Didesmi species? DC. syst. 2. p. 669.

Subordo IV. SPIROLOBÆ. (o || ||) DC. syst. 2. p. 670.

Cotyledones incumbentes, lineares, spiraliter seu potius circinnatim convolutæ. Semina subglobosa.

Trib. XVII. BUNIADEÆ seu SPIROLOBÆ NUCAMEN-TACEÆ. DC. syst. 2. p. 670.

Silicula nucamentacea indehiscens 2-4-locularis. Cotyledones verè circinnatim spirales.

XCIH. BUNIAS. Brown in hort. Kew. ed. 2. v. 4. p. 75. DC. syst. 2. p. 670.

Character idem ac trib.

Sect. I. ERUCAGO. DC. syst. 2. p. 670. — Tourn. inst. 232. t. 103.

Calyx erectus. Petala limbo obcordato obtusè emarginato. Silicula tetragona angulis cristato-alatis subdentatis, stylo filiformi superata, intùs 4-locularis.

1. *B. ERUCAGO* (Lin! spec. 935.) siliculis tetragonis, angulis cristatis, foliis radicalibus runcinatis. ① in arvis et agris Europæ australis. Jacq. fl. austr. t. 340. *Myagrum erucago* Lam. dict. 1. p. 571. n. 12.* Flores noctu nutantes, flavi. — DC. syst. 2. p. 671.* (v. v.)

2. *B. ASPERA* (Retz obs. 2. p. 21.*) siliculis tetragonis, angulis cristatis, foliis omnibus lanceolatis. ① in Lusitaniâ. *Myagrum asperum* Poir. suppl. 2. p. 47. Descriptiones auctorum omnino cum *B. erucagine* conveniunt nisi quod folia inferiora ferè integra DC. syst. 2. p. 672.*

β. *cristata*, foliis inferioribus sinuato-dentatis. *Bunias aspera* Brot. fl. lus. 1. p. 562.

Sect. II. LAELIA. DC. syst. 2. p. 672.

Siliculis 2-ocularibus ovatis.

3. *B. ORIENTALIS* (Lin. spec. 936. excl. hort. ups. syn.) siliculis ovatis bilocularibus ecristatis subverrucosis. 2 in graminosis Transylvaniæ, Tauriæ, Arabiæ felicis, in Russiâ Europæanâ et in Sibirîâ. Schkuhr handb. 2. n. 1918. t. 189. *Myagrum taraxacifolium* Lam. dict. 1. p. 570. n. 10 *

Trib. XVIII. ERUCARIÆ seu SPIROLOBEÆ LOMENTACEÆ. DC. Syst. 2. p. 673.

Siliqua lomentacea biarticulata, articulo inferiore biloculari, superiore ensiformi. Cotyledones replicatæ subspirales.

XCIV. ERUCARIA. Gaertn. fruct. 2. p. 298. t. 143. f. 9. DC. syst. 2. p. 673.*

Car. id. ac trib.

§. I. *Siliculæ articulo superiore in stylum filiformem productæ.*

1. *E. ALEPICA* (Gaertn. l. c.) siliquâ styliferâ, foliis pinnatipartitis, lobis linearibus inferiorum pinnatifidis superiorum integris ① in insulis Archipelagi græci, in Asiâ minori, circa Alexandriam et in Syriâ. *Cordilocarpus lævigatus* Willd. spec. 3. p. 563.* Flores pallide violacei. — DC. syst. 2. p. 674.* (v. v. et s.)

2. *E. LATIFOLIA* (DC. syst. 2. p. 675.*) siliquâ styliferâ, foliis pinnatipartitis, lobis oblongis inciso-dentatis. ① in Aegypto, Syriâ. Deless! ic. sel. 2. t. 95. *Sinapis Hispanica* Herb. Banks. Flores vix magnitudine prioris. (v. s.)

3. *E. TENUIFOLIA* (DC. syst. 2. p. 675.*) siliquæ articulo inferiore dispermo, superiore 1-spermo, foliis bipinnatipartitis, lobis linearibus. ① in Hispaniâ? *Sinapis Hispanica* Lin! spec. 935. excl. syn. Tourn. (v. s. in herb. Lin. nunc Smith.) †

§. II. *Siliquæ articulo superiore (rostro) stigmate sessili apiculato.*

4. *E. CRASSIFOLIA* (Delill! ill. fl. eg. p. 20. pl. hot. t. 34. f. 1. descr. p. 100.*) stigmate sessili, rostro siliquâ longiore, foliis pinnatipartitis crassis, lobis linearibus. ① in Aegypto frequens circa Pyramides Saqqarah. *Brassica crassifolia* Forsk. fl. æg. ar. descr. p. 118.* Flores vix magnitudine *E. Alepicæ*, ex albo violacei. — DC. syst. 2. p. 675.* (v. s.)

5. *E. HYRCANICA* (DC. syst. 2. p. 676.*) stigmate sessili, rostro siliquæ longitudine, foliis paucis grossè dentatis. ①? in Persiâ bor. *Sisymbrium myagro affine* Stev! in litt. Flores in sicco ochroleuci. (v. s.)

Subordo V. DIPLECOLOBEÆ (o || || ||) DC. syst. 2. p. 776.

Cotyledones incumbentes, lineares, bicrures seu buplicatæ nempè bis transversè plicatæ. Semina depressa.

Trib. XIX. *HELIOPHILÆ* seu *DIPLECOLOBÆ* *SILICULOSÆ*. DC. syst. 2. p. 676.

Siliqua elongata aut rarius oblonga ovalisve; septo lineari aut ovali, valvis planis aut in siliquis elongatis convexiusculis.

XCv. CHAMIRA. Thunb. nov. gen. pl. 2. p. 48. DC. syst. 2. p. 677.

Calyx basi bicalcaratus.

1. *C. CORNUTA* (Thunb. l. c.) ① ad caput Bonæ-Spei. *Heliophila circeoides* Lin. f. suppl. 298. Folia alterna cordata aut ferè reniformia, ferè Cymbalarie sed multò majora. — DC. syst. 2. p. 677.* (v. s. in h. Lin. Banks Burm.)

XCvi. *HELIOPHILA*. N. Burm. in Lin. gen. n. 816. Lam. ill. t. 563. DC. syst. 2. p. 677.

Calyx basi æqualis.

Sect. I. *CARPONEMA*. DC. syst. 2. p. 679.

Siliquæ sessiles, teretes, inter semina vix ac ne vix coarctatæ, utrinque acuminatæ, indehiscentes aut vix dehiscentes. — Herba annua.

1. *H. FILIFORMIS* (Lin. f. suppl. 296.) herbacea glabra, siliquis teretibus utrinquè attenuatis, foliis lineari-subulatis. ① ad caput Bonæ-Spei. Lam. ill. t. 563. f. 3? Flores parvi, purpurascens. DC. syst. 2. p. 679.* (v. v. in h. Kew.)

Sect. II. *LEPTORMUS*. DC. syst. 2. p. 680.

Siliquæ sessiles subcompressæ gracillimæ submoniliformes, inter semina nempe subcoarctatæ, monilibus ovato-oblongis. — Herba annua inter *Carpone-mata* et *Ormiscos* ferè mediæ.

2. *H. DISSECTA* (Thunb. prodr. 108.) herbacea glabra, siliquis tenuissimis submoniliformibus stigmatè crasso nodoso terminatis, foliis filiformibus integris trifidis partitisve. ① ad caput Bonæ-Spei. Flores cærulei. DC. syst. 2. p. 680.* (v. s. in h. Banks.)

℘. *H. PINNATA* var. Vahl! herb. Flores in sicco albi. (v. s. in h. Vahl.)

3. *H. TENELLA* (Banks! herb. DC. syst. 2. p. 680.*) herbacea glabra, siliquis..... foliis filiformibus integris, caule simplici. ☉ hab..... Priori valde affinis sed adhuc siliquis ignotis dubia! (v. s. in h. Banks.) †.

4. *H. TENUISILIQUA* (DC. syst. 2. p. 680.*) herbacea glabra, siliquis tenuissimis compressis submoniliformibus stigmatè acuminato terminatis, foliis linearibus integris. ① ad cap. Bonæ-Spei. Deless! ic. sel. 2. t. 96. *Arabis Capensis* Burm! herb. non vero fl. cap. Flores in sicco albidii, sed verosimiliter subrubentes. (v. s. in h. Deless.)

5. *H. LONGIFOLIA* (DC. syst. 2. p. 681.*) herbacea glabra, siliquis tenuissimis compressis submoniliformibus, stigmatè punctiformi, foliis linearibus integris tripartitisve. ① ad cap. Bonæ-Spei. Flores cærulei omninò *H. pilosæ*. (v. s. in h. Vahl.)

6. *H. SONCHIFOLIA* (DC. syst. 2. p. 681.*) herbacea glabra, siliquis tenuibus subcompressis torulosis, foliis radicalibus pinnatipartitis lobis 2-3-jugis linearibus. ①? ad cap. Bonæ-Spei. Flores cærulei. Sepala oblonga, margine vix membranacea. (v. s. in h. Banks.)

Sect. III. *ORMISCUS*. DC. syst. 2. p. 682.

Siliquæ sessiles compressissimæ moniliformes, marginibus nempe inter semina sinuato-coarctatis, monilibus 1-spermis orbiculatis. Semina orbiculata, compressissima. Stamina edentula. — Herba annua.

7. *H. AMPLEXICAULIS* (Lin. f. suppl. 296.) herbacea glabra, siliquis monili-

- formibus, foliis infimis oppositis superioribus alternis cordato-amplexicaulibus oblongis integerrimis. ① ad cap. Bonæ-Spei. Jacq. fragm. 49. t. 64. f. 2. Flores parvi ex albo subviolacei. DC. syst. 2. p. 682.* (v. s.)
8. *H. RIVALIS* (Burch! cat. pl. afr. extratrop. austr. n. 5496.) herbacea glabra, siliquis moniliformibus erectiusculis, foliis pinnatipartitis, lobis 3-5-jugis linearibus acutis, petalis obovatis. ① ad cap. Bonæ-Spei. Icon Lamarckii ill. t. 563. f. 2. hunc florentem non malè refert. Flores albi, interdum ut in Cheirantho mutabili lilacini. DC. syst. 2. p. 682.* (* s. comm. à cl. Burch.)
9. *H. VARIABILIS* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 1249.) herbacea glabra, siliquis moniliformibus erectiusculis, foliis pinnatipartitis, lobis 3-5-jugis linearibus acutis petalis oblongo-linearibus. ① ad cap. Bonæ-Spei. Flores primum albi, dein dilutè rosei. Habitus ferè *H. chamæmelifoliæ*. DC. syst. 2. p. 683.* (v. s. in h. Burch.)
10. *H. PENDULA* (Willd. spec. 3. p. 529.*) herbacea glabra, siliquis moniliformibus pendulis, foliis pinnatipartitis 3-5-jugis, lobis linearibus integris. ① ad cap. Bonæ-Spei. *H. pinnata* Vent! malm. t. 113. excl. syn. Lam. ill. t. 562. f. 2. Petala ungue brevi flavescente, limbo sordidè albido. — DC. syst. 2. p. 683.* (v. v.)
11. *H. TRIFIDA* (Thunb. prodr. 108.) herbacea glabra, siliquis moniliformibus subpendulis, foliis 3-fidis aut pinnatim 5-fidis, lobis integris subfiliformibus. ① ad cap. Bonæ-Spei. *H. pinnata* Lin. f. suppl. 297. Flores parvi purpurascetes, ferè *Arabidis vernæ* DC. syst. 2. p. 683. (v. s. in herb. Banks.)
12. *H. FUSILLA* (Lin. f. suppl. 297.) herbacea glabra, siliquis moniliformibus patulis, foliis lineari-setaceis integris. ① ad cap. Bonæ-Spei. *Arabis Capensis* Børn! prodr. fl. cap. 18. — Pluk. mant. 135. t. 432. f. 2. Flores parvi albi. — DC. syst. 2. p. 684.* (v. s.)
13. *H. LEPIDIOIDES* (Link enum. 2. p. 174.) herbacea glabra, siliquis erectopatulis oblongo-ovatis margine subrepandis oligo-spermis, foliis pinnatipartitis, lobis linearibus. ① ad cap. B.-Spei. *Trentepohlia lepidioides* Roth. nov. sp. 326?
14. *H. SESELIIFOLIA* (Burch! cat. geogr. pl. afr. austral. extratrop. n. 1318 et 1391.) herbacea glabra, siliquis moniliformibus, foliis pinnatipartitis, lobis lineari-subulatis, staminibus lateralibus dente auctis. ① ad caput Bonæ-Spei. Habitus ferè *H. pendulæ*. Petala alba, obovata. An hujus sectionis? an potiùs *Leptormis* consocianda? DC. syst. 2. p. 684.* (v. s. in h. Burch.)

Sect. IV. SELENOCARPÆA. DC. syst. 2. p. 684.

Siliquæ sessiles compressæ ovales aut orbiculatæ. ①

15. *H. DIFFUSA* (DC. syst. 2. p. 685.*) herbacea glabra, siliculis ovali-oblongis 4-spermis, foliis pinnatipartitis, lobis filiformibus integris. ① ad cap. Bonæ-Spei. *Lunaria diffusa* Thunb. prodr. 107. *Trentepohlia lepidioides* Roth nov. pl. sp. 326? Flores mihi ignoti, ex reliquiis parvi. (v. s. in h. Banks.)
16. *H. PELTARIA* (DC. syst. 2. p. 685.*) herbacea glabra, siliculis ovali-orbiculatis 2-4-spermis, oliis pinnatipartitis, lobis linearibus integris. ① ad cap. Bonæ-Spei. *Peltaria Capensis* Lin. f. suppl. 296* Flores parvi albi. Planta adest in herbario Linnæano sed innominata. (v. s. in h. Lin. et Smith.)

Sect. V. ORTHOSELIS. DC. syst. 2. p. 685.

Siliquæ sessiles compressæ lineares, marginibus rectis.

§. 1. *Caulibus herbaceis annuis.*

17. *H. PILOSA* (Lam. dict. 3. p. 90.*) caule herbaceo pilis patulis hispido, siliquis linearibus foliis pilosiusculis nunc linearibus integris, nunc apice

trifidi⁴ basi cuneatis. ① ad cap. Bonæ-Spei. Petala lætè cærulea, calyce duplò longiora. DC. syst. 2. p. 686.* (v. s.)

a. *integrifolia*, foliis linearibus integris. H. *integrifolia* Lin. spec. 926. excl. Pluk. syn. Jacq. ic. rar. 3. t. 506.

β. *incisa*, foliis basi cuneatis in lobos lineares apice trifidis. H. *arabioides* Sims! bot. mag. t. 496.

18. H. *DIGITATA* (Lin. f. suppl. 296.) caule herbaceo pilis patulis hispido, siliquis linearibus, foliis ovalibus integris aut hinc inde grosse inciso-dentatis. ① ad cap. Bonæ-Spei. Jacq. f. icon. ex herb. Jacq. Forsan prioris varietas! DC. syst. 2. p. 686.* (v. s. in h. Vahl.)

19. H. *INCISA* (herb. Banks! et DC. syst. 2. p. 687.*) herbacea glabra, siliquia linearibus, foliis oblongis basi cuneatis apice lobatis, lobis angustis acutis. ① ad cap. Bonæ-Spei. Flores mediocres, flavi. (v. s. in h. Banks.)

20. H. *DIVARICATA* (herb. Banks! et DC. syst. 2. p. 687.*) herbacea glabra, siliquis, foliis linearibus integerrimis, ramis patentibus. — ad cap. Bonæ-Spei. Ex habitu saltem in herbario *Lepidium graminifolium* ferè referens. Flores parvi, ex sicco flavi. †. (v. s. in h. Banks.)

21. H. *CORONOPHOLIA* (Lin. spec. 927. excl. syn. Pluk.) herbacea glabra, siliquis linearibus, foliis pinuati-partitis, lobis rachique linearibus integerrimis. — ad cap. Bonæ-Spei. Herm. lugd. 366 et 367, ic. Flores patentes, cæruleo-violacei. An hujus sectionis. DC. syst. 2. p. 687.* (v. s.)

22. H. *TRIFURCA* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 1487.) herbacea glabra, siliquis linearibus deflexis, foliis angustissimis trifurcis, lobis linearibus integerrimis. ① ad cap. Bonæ-Spei. H. *tripartita* Thunb. prodr. 108? Petala obovata magna, ex sicco pallidè purpurascentia. — DC. syst. 2. p. 688.* (v. s. comm. à cl. Burch.)

23. H. *PECTINATA* (Burch. cat. geogr. pl. afr. austr. extratrop. n. 1397 et 1362.) herbacea glabra, siliquis linearibus pendulis, foliis pinnati-partitis, lobis 4-5-jugis linearibus approximatis. ① ad cap. Bonæ-Spei. *Farsetia elongata* Desv. Journ. 3. p. 173? Ex habitu accedit ad *Selenocarpæas* sen *Lunarias Thunbergii* et eas cum *Orthoselides* conjungit. Flores minimi, albidii. — DC. syst. 2. p. 688.* (v. s. comm. à cl. Burch.)

24. H. *FÆNICULACEA* (Brown! in hort. kew. ed. 2. v. 4. p. 100.) herbacea pilosiuscula, siliquis linearibus patulis, foliis pinnatim bipinnatimve partitis, lobis filiformibus. ① ad cap. Bonæ-Spei. Flores vix mihi cogniti, at ex sicco videntur parvi purpurei. DC. syst. 2. p. 689.* (v. s. in herb. Banks.)

25. H. *CHAMÆMELIFOLIA* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 1226.) herbacea basi pilosiuscula, apice glabra, siliquis linearibus erectiusculis, foliis pinnati-partitis rachi lobisve linearibus integerrimis acutis. ① ad cap. Bonæ-Spei. Aff. H. *crithmifoliæ*. Flores albidii. DC. syst. 2. p. 689.* (v. s. in h. Burch.)

26. H. *CRITHMIFOLIA* (Willd. enum. 2. p. 682.) herbacea velutino-pubescentis, siliquis linearibus nutantibus, foliis pinnati-partitis subcarnosis, lobis subfiliformibus superne sulcatis. ① ad cap. Bonæ-Spei. Deless! ic. sel. 2. t. 97. *Sisymbrium crithmifolium* Roth nov. pl. spec. 225.* Flores parvi, violacei. (v. s.)

β. H. *parviflora* Burch! cat. geogr. pl. afr. austr. extratrop. n. 1236. Non differre videtur nisi floribus albis.

§. 2. *Caulibus frutescentibus.*

27. H. *ABROTANIFOLIA* (herb. Banks! et DC. syst. 2. p. 690.*) suffrutescens glabra, siliquis linearibus patulis, foliis 3-5-lobatis, lobis subulatis brevibus. ① ad cap. Bonæ-Spei. Flores mediocres. Calyces membranacei. Semina plana, marginata, orbiculata. (v. s. in h. Banks.)

28. *H. GLAUCA* (Burch! cat. geogr. pl. afr. austr. extratrop.) suffruticulosa glabra glauca, siliquis linearibus erectis, foliis oblongis subcarnosis inferioribus obtusis, superioribus acutis. ζ ad cap. Bonæ-Spei. Flores magnitudine Vellæ pseudocytisi, ad quam habitu frutex noster paululum accedit. DC. syst. 2. p. 690.* (v. s. in herb. Burch.)
- a. candida*, floribus candidis. Burch. l. c. n. 4782. Propè Loeri river.
- ß. purpurascens*, floribus purpureo tinctis. Burch. l. c. n. 4969. in regione Langekloof.
29. *H. FASCICULARIS* (herb. Banks! et DC. syst. 2. p. 691.*) fruticulosa glabra, siliquis linearibus suberectis pedicello vix longioribus, foliis filiformibus. ζ ad cap. Bonæ-Spei. (v. s. in h. Banks.)
30. *H. SUAVISSIMA* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 2742.) fruticulosa glabra, siliquis sublinearibus basi attenuatis patulis, foliis lineari-subulatis acutiusculis. ζ ad cap. Bonæ-Spei. *H. juncea* Vahl! herb. Flores odoratissimi, violacei, in sicco sæpè albescentes. DC. syst. 2. p. 691.* (v. s. comm. à cl. Burch.)
31. *H. SUBULATA* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 6214.) fruticulosa brevissimè pubescens, siliquis linearibus patulis basi non attenuatis, foliis lineari-subulatis acutissimis. ζ ad cap. Bonæ-Spei. Hanc primà fronte pro merà prioris varietate habui, sed post iterata examina ab eà separavi. DC. syst. 2. p. 691.* (v. s. in herb. Burch.)
32. *H. PLATYSILIQUA* (Brown. in h. kew. ed. 2. v. 4. p. 99.) suffruticosa glabra, siliquis linearibus erectis pendulisve, foliis carnosis semi-teretibus. ζ ad cap. Bonæ-Spei. *Hesperis cheiranthus carnosus*. Poir. suppl. 3. p. 197. Flores ex sicco purpurascens. DC. syst. 2. p. 692.* (v. s. in h. Banks.)
33. *H. LINEARIFOLIA* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 347 et 793.) suffruticosa glabrescens, siliquis erectis linearibus stylo apiculatis, foliis linearibus acutis integerrimis. ζ ad cap. Bonæ-Spei. Petala cærulea. DC. syst. 2. p. 692.* (v. s. in h. Burch.)
- ß. hirsuta* Burch! l. c. n. 5518. Ad Nysna in regione Anteniqua-land.
34. *H. STYLOSA* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 3291.) fruticulosa glabra, siliquis pendulis linearibus stylo filiformi pedicello longiore apiculatis, foliis linearibus integerrimis. ζ ad cap. Bonæ-Spei. Petala oblongo-linearia in sicco è flavo subrubentia. DC. syst. 2. p. 693.* (v. s. in h. Burch.)
35. *H. VIRGATA* (Burch! cat. geogr. pl. afr. austr. extratrop.) suffruticosa glabra, siliquis subdeflexis linearibus stylo filiformi pedicello longiore apiculatis, foliis ovatis integris dentatisve. ζ ad cap. Bonæ-Spei. DC. syst. 2. p. 693.* (v. s. in h. Burch.)
- a. integrifolia*, foliis integris, floribus è flavido albis. Burch. l. c. n. 4605.
- ß. dentata*, foliis grossè et argutè dentatis, floribus albis odoratissimis. Burch. l. c. n. 3933.
36. *H. SCOPARIA* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 7887 et 8557.) fruticulosa glabra, siliquis erectis linearibus in stylum brevem abeuntibus, foliis lineari-subulatis, racemis paucifloris. ζ ad cap. Bonæ-Spei. *Cheiranthus strictus* Poir! suppl. 2. p. 781. Petala apice rosea. DC. syst. 2. p. 693.* Deless! ic. sel. 2. t. 98. (v. s.)

Sect. VI. PACHYSTYLUM. DC. syst. 2. p. 694.*

Siliquæ sessiles subcompressæ lineares velutinæ, stylo crasso conico brevi.
— Suffrutex pubescens foliis integris.

37. *H. INCANA* (Ait h. kew. ed. 1. v. 2. p. 397. ed. 2. v. 4. p. 99.) siliquis linearibus compressis velutinis stylo crasso conico glabro superatis, foliis oblongis. ζ ad cap. Bonæ-Spei. *H. frutescens*. Lam. dict. 3. p. 91.* Petala cæruleo-purpurea, calyce longiora. DC. syst. 2. p. 694.* (v. s.)

Sect. VII. LANCEOLARIA. DC. syst. 2. p. 695.

Siliquæ sessiles lanceolares. Cotyledones bicornues subspirales. ζ .

38. *H. MACROSPERMA* (Burch! cat. geogr. pl. afr. austr. extratrop. n. 3425.) suffruticosa glabra, siliquis lanceolatis in stylum brevem desinentibus, foliis linearibus acutis. ζ ad cap. Bonæ-Spei. Flores ignoti. DC. syst. 2. p. 695. (v. s. in h. Burch.)

Sect. VIII. CARPOPODIUM. DC. syst. 2. p. 670.

Siliquæ pedicellatæ lineares.

39. *H. CLEOMOIDES* (DC. syst. 2. p. 695.*) siliquis compressis pedicellatis, foliis lineari lanceolatis. ζ ad cap. Bonæ-Spei, in Monomotapa. Cleome Capensis. Lin. spec. 940. Cheiranthus strictus Lin. f. suppl. 296. Petala oblonga calyce paulò longiora, ex sicco ut videtur flava. — Deless! ic. sel. 2. t. 99. (v. s.)

†. *Heliophilæ vix notæ.*

40. *H. LINIFLORA* (DC. syst. 2. p. 696.) ① ad cap. Bonæ-Spei. — Herm. Lugd. 366.
 41. *H. MOLLUGINEA* (DC. syst. 2. p. 696.) — ad cap. Bonæ-Spei. — Pluk. alm. 213. t. 200. f. 3. Weinm. phyt. t. 641. f. c. in icone folia opposita, tripartita, lobis linearibus acutis.
 42. *H. HETEROPHYLLA* (Thunb. prodr. 107.) — Ad cap. Bonæ-Spei. Specimen, ut videtur è Thunbergio, adest sub hoc nomine in herb. cl. Vahlîi, sed à nostrâ *H. linearifoliâ* non differre videtur et folia omnia indivisa gerit.
 43. *H. LYRATA* (Thunb. prodr. p. 108.) — Ad cap. Bonæ-Spei. Folia lyrata; siliquæ teretes.
 44. *H. CALLOSA* (DC. syst. 2. p. 696.) ζ ad cap. Bonæ-Spei. Cheiranthus callosus Lin. f. suppl. 296. Caulis fruticosus angulatus. Folia lanceolata integra, callosa.
 45. *H? LINEARIS* (DC. syst. 2. p. 697.) — Ad cap. Bonæ-Spei. Hesperis cheiranthus linearis Poir. suppl. 3. p. 197. Cheiranthus linearis Thunb. prodr. 108. Caulis erectus, herbaceus. Folia linearia, glabra.
 46? *H. GRAMINEA* (DC. syst. 2. p. 697.) — Ad cap. Bonæ-Spei. Cheiranthus graminens Thunb. prodr. 108. Hesperis cheiranthus gramineus Poir. suppl. 3. p. 167. Caulis herbaceus, simplex, erectus. Folia radicalia, ensiformia.
 47. *H? ELONGATA* (DC. syst. 2. p. 697.) — Ad cap. Bonæ-Spei. Cheiranthus elongatus Thunb. prodr. 108. Hesperis cheiranthus elongatus Poir. suppl. 3. p. 197. Caulis herbaceus, erectus. Folia linearia, integra, glabra.

Trib. XX. SUBULARIÆ seu DIPLECOLOBÆ LATI-SEPTÆ. DC. syst. 2. p. 697.

Silicula ovalis, septo elliptico, valvis convexis, loculis polyspermis, stigmate sessili. Cotyledones bicipitatæ.

XCVII. SUBULARIA. Lin. gen. n. 799. DC. syst. 2. p. 697.

Car. id. ac trib.

1. *S. AQUATICA* (Lin! spec. 896.) ① in Europâ frigidiori. OEd. fl. dan. 35. Smith engl. bot. t. 732. Folia integerrima 8-10 lin. longa. Flores minimi. DC. syst. 2. p. 698.* (v. s.)

Trib. XXI. BRACHYCARPÆ seu DIPLECOLOBÆ ANGUSTISEPTÆ. DC. syst. 2. p. 698.

Silicula didyma, septo angustissimo, valvis valdè ventricosis, loculis 1-spermis, stylo brevi. Cotyledones bicipitatæ?

Car. id. ac trib.

1. *B. VARIANS* (DC. syst. 2. p. 699.*) suffruticosa glabra, siliculis subdidymis, foliis oblongo-linearibus. 3 ad promont. Bonæ-Spei. Deless! ic. sel. 2. t. 200. Flores ampli. (v. s.)

a. flava, floribus flavis, Heliophila flava, Lin. fl suppl. 297.*

β. purpurascens, floribus purpurascentibus. Polygala bracteolata Burm! herb

**CRUCIFERÆ QUÆDAM INQUIRENDÆ, AD GENERA
NONDUM REVOCATÆ.**

NASTURTIUM? *SINENSE* (DC. syst. 2. p. 699.) — In Chinâ. *Lepidium petræum*. Lour. fl. coch. ed. Willd. 2. p. 479. excl. syn. *Lepidium Chinense* Stok. bot. mat. med. 3. p. 429. Caulis annuus, teres, rectus. Folia lato-lanceolata integerrima, parva, pinnata. Flos dilutè luteus, spicis simplicibus erectis longis terminalibus. Silicula ovata, emarginata, compressa, 2-locularis, polysperma.

DIPLOTAXIS? *SETOSA* (DC. syst. 2. p. 699.*) 2 in Pyrenæis. *Turritis setosa* Lapeyr. abr. suppl. 93.* Flores capitati, conferti. Petala flava angusta. Siliqua longa, teres, glabra, truncata. Planta tota petalis siliquisque exceptis setas longas confertas albas pilis intermixtas gerit.

NASTURTIUM? *ROTUNDFOLIUM* (Rafin. fl. lud. p. 86. n. 274.) in Louisiana. Habitus *Arabis rhomboideæ*, Planta magna, caule erecto ramoso, foliis subrotundis sinuato-lobatis crassis succosis edulibus, petalis elongatis calyce longioribus.

DISCOVIUM. Rafin. in *journ. phyt.* vol. 89. ann. 1819. p. 96.

Calyx clausus. Silicula lenticularis, septo integro, valvis carinatis, loculis polyspermis, stylo persistente, stygmate obtuso. — Intermedium inter *Thlaspi*, *Alyssum* et *Lepidium*. (Rafin.)

- D. *OHIOENSE* (Rafin. l. c.) ① ad ripas fluminis Ohio propè Gallipoli. Species unica puberula erecta simplex gracilis, foliis distantibus sessilibus oblongo-linearibus obtusis integris, petalis flavis integris cuneiformibus calyci fere æqualibus. Rafin.

SCHIZOPETALUM. Sims *bot. mag.* t. 2379.

Calyx cylindraceus, basi æqualis, clausus. Petala ovata, inciso-pinnatifida. Stigmata gibbosa approximata. Siliquæ et semina ign. — An *Cardamini*, an *Sisymbrio*, an *Brassicæ* affine?

- S. *WALKERI* (Sims l. c.) ① seu ② in Chili; glaberrimum adsurgens. Folia oblonga, sinuato-pinnatifida, floralia linearia integra. Flores albi.

Ordo XII. CAPPARIDEÆ. *Juss. gen. 242. ann. 18. p. 474.*

Sepala quatuor, nunc ferè libera, æqualia aut inæqualia, nunc basi in tubum coalita, limbo vario. Petala 4 cruciata, sæpè unguiculata et inæqualia. Stamina ferè perigyna imo calyci inserta, rariùs tetradynama, sæpiùs ordine quaternario multiplici disposita, definita aut indefinita. Torus hemisphæricus aut elongatus sæpè glandulifer. Thecaphorum è toro ortum gracile. Ovarium ideò stipitatum, ovato-oblongum carpellis 2 arctè coalitis compositum. Stylus nullus aut filiformis. Fructus varius, siliquosus baccatusve, intùs unilocularis, rarissimè 1-spermus, sæpiùs placentis binis polyspermis in fructibus dehiscentibus intervalvularibus. Semina sæpiùs reniformia, exalbuminosa (endopleurà tamen tumidà.) Embryo incurvus. Cotyledones foliaceæ, planiusculæ, subincumbentes. — Herbæ, frutices aut arbores. Stipulæ nullæ aut spinescentes. Folia alterna, petiolata, simplicia aut palmatim composita. Flores dispositione varii, Thalamifloras inter et Calycifloras structurà medii. Ordo accedens hinc Cruciferis Papaveraceisque, illinc Rosaceis, Passifloreis, Droseraceis et Flacourtianeis.

Trib. I. CLEOMEÆ. — Cleome *Lin.*

Fructus vere capsularis, valvis submembranaceis dehiscentibus. — Herbæ aut suffrutices foliis sæpè compositis, pube sæpè glandulosâ.

I. CLEOMELLA.

Calyx 4-sepalus patens. Petala 4. Torus ovato-globosus. Stam. 6. Siliqua in calyce stipitata latitudine brevior.

1. *C. MEXICANA* (ic. fl. mex. ined.) \mathcal{U} in Mexico. Folia, 3-foliolata, glabra. Fl. flavi.

II. PERITOMA. — *Atalanta Nutt. gen. am. 2. p. 73. non Corr.*

Calyx basi circumscissus, apice 4-dentatus. Petala 4. Torus parvus. Stamina 6 basi monadelpha. Siliqua oblonga in calyce stipitata.

1. *P. SERRULATUM*. Cleome serrulata Pursh! fl. bor. am. 2. p. 441. (1) in Amer. bor. ad ripas Missouri. — Fol. 3-foliolata, glabra. Fl. purpurei. (v. s. in h. Lamb.)

III. GYNANDROPSIS. — Cleomes sp. *Lin.*

Calyx 4-sep. patens. Petala 4. Torus elongatus. Stam. 6 circa torum monadelpha ad ejus apicem libera. Siliqua in calyce ad apicem tori stipitata.

1. *G. SESSILIFOLIA*, glabra, foliis omnibus 3-foliolatis sessilibus, infimis vix subpetiolatis, foliolis oblongis integris. (1) Hab. . . . Cleome triphylla Lin. sp. 938 ex herb. Vahl et Dahl, sed excl. syn. (v. s. in h. Vahl.)
2. *G. TRIPHYLLA*, glabriuscula, foliis omnibus 3-foliolatis floralibus sessilibus,

- ceteris petiolatis, inferiorum petiolo foliis longiore. ① in ins. Caribæis, Senegaliâ. *Cleome-triphylla* Lin. sp. 938. ex Herm. lugd. 565 ic. An sequentis varietas? (v. s.)
3. *G. PENTAPHYLLA*, glabriuscula, foliis mediis 5-foliolatis, infimis floralibusque 3-foliolatis, foliolis integris subserrulatisve. ① in Caribæis et Amer. calid. Cl. pentapphylla Lin. sp. 938. bot. mag. t. 1681. An etiam spontanea in Ægypto et Indiâ? (v. v.)
 4. *G. DENTICULATA*, glabriuscula, foliis infimis 7-foliolatis, mediis 5, summis 3-foliolatis, foliolis serrulatis. ① in Angolâ. (v. s. in h. Mus. Par.)
 5. *G. PALMIPES*, glabriuscula, foliis 5-7-foliolatis, foliolis integris, petiolulis membranula subcoherentibus. ① in Cayennâ. (v. s.)
 6. *G. HETEROTRICHA*, pilis aliis elongatis aliis glanduloso sessilibus pilosiuscula, foliis 3-5-foliatis, foliolis subintegris. ① ad cap. Bonæ-Spei. *Cleome heterotricha* Burch! trav. 1. p. 537. cat. n. 2011. (v. s.)
 7. *G. BRACHYCARPA*, viscoso-pubescent, foliis 3-5-foliolatis, foliolis integris, siliquâ glabrâ ovali-oblongâ thecaphoro brevior. — in Peruviâ. Cl. brachycarpa Vahl. (v. s. in h. Juss.)
 8. *G. HISPIDULA*, hispida, foliis 5-foliolatis, foliolis integris, siliquâ hispidulâ thecaphoro brevior. — in Peruviâ. *Cleome hirsuta* Ruiz et Pav! (v. s. in h. Lamb.)
 9. *G. SPECIOSA*, apice subvelutina, foliolis 5-7-foliolatis, foliolis subserrulatis oblongis acuminatis. 2? in Novâ-Granadâ propè Carthagenam. *Cleome speciosa* Humb. Bonpl. et Kunth! nov. sp. am. 5. p. 84. t. 436. (v. s. in h. Humb.)

IV. CLEOME. — *Cleomes* sp. Lin.

Calyx 4-sepalus patens subæqualis. Petala 4. Torus subhemisphæricus. Stam. 6, rarius 4. Siliqua dehiscens in calyce stipitata aut sessilis.

Sect. I. PEDICELLARIA.

Torus carnosus subglobosus. Tecaphorum elongatum.

1. *C. GIGANTEA* (Lin. mant. 430.) fruticosa velutino-pubescent subviscosa, foliis 7-foliolatis, venis foliorum utriusque 30-40. ♂ in Americâ merid. Cl. viridiflora Schreb. nov. aet. nat. cur. 4. p. 136. t. 3. (v. v.)
 2. *C. ARBOREA* (Humb. Bonpl. et Kunth! nov. sp. 5. p. 86.) fruticosa velutino-pubescent subviscosa, foliolis 7-foliolatis, venis foliorum utriusque 20. ♂ ad Caracas. An satis à *C. giganteâ* differt? (v. s. in h. Humb.)
 3. *C. GLANDULOSA* (Ruiz et Pav.) subarborescent, pilis crassis apice glandulosis hispida, foliis 7-9-foliolatis, floralibus summis simplicibus ovato-cordatis, ovario piloso-glanduloso. ♂ in Peruviâ fruct. ign. Siliqua thecaphori longitudine. (v. s. in h. Mus. Par. et Juss.)
 4. *C. CARDINALIS*, herbacea glabra, foliis 5-foliolatis, floralibus sessilibus foliolisque ovali-lanceolatis. — in Mexico. Vulg. Borla di Cardinal. (ic. fl. mex. ined.) †.
 5. *C. HEPTAPHYLLA* (Lin. sp. 987.) herbacea aculeata hirsuto-viscosa, foliis 7-foliolatis, floralibus simplicibus cordato-subrotundis, siliqua thecaphoro longiore viscoso-pubescente. ① in Indiis. An plures hic confusæ species? An Indica eadem ac Jamaïcensis, aut Essequiboensis. (v. s. in h. Mus. Par.)
 6. *C. ANOMALA* (Humb. B. et K! nov. sp. 5. p. 85.) herbacea inermis velutino-pubescent, foliis 7-foliolatis, floralibus simplicibus subcordato-rotundis, siliquâ ellipsoideâ mucicato-tuberculatâ. — in Americâ calid. Fruct. valvulæ valdè convexæ. (v. s. in h. Humb.)
 7. *C. CHILENSIS*, herbacea inermis glabra, foliis 7-foliolatis floralibus 3-5-foliolatis, sepalis persistentibus, siliquâ pendulâ glabriusculâ thecaphoro vix longiore. ① in Chili. (v. s. in h. Mus. Par.)
- a. *glabra*, foliis 7-foliolatis.

β. pubescens, foliis 3-5-foliolatis.

8. *C. LONGIPES*, herbacea aculeata glabra, foliis 3-foliolatis, siliquâ cylindraceâ tenni thecaphoro (pedali!) dimidio brevior. — in Hoyaquil. (v. s. in h. Lamb.)
9. *C. SPINOSA* (Lin. sp. 939.) herbacea aculeata pubescens, foliis 5-7-foliolatis glabris, floralibus simplicibus petiolulatis ovato-cordatis, siliquâ glabrâ thecaphorò longiore. ① in Americâ calid. — Marcegr. bras. p. 34. ic. (v. s.)
10. *C. PUNGENS* (Willd! h. ber. t. 18) herbacea aculeata pubescenti-viscosa, foliis 5-7-foliolatis glabris, floralibus simplicibus sessilibus cordatis, siliquâ glabrâ thecaphoro brevior. ①? 2 ex Bert. in Amer. calid. et ins. Carib. *C. pungens* Humb. B. K. nov. sp. 5. p. 85. *C. spinosa* bot. mag. t. 1640. (v. s.)
11. *C. PUBESCENS* (Sims. bot. mag. t. 1857.) herbacea inermis pubescens, foliis 5-7-foliolatis, floralibus simplicibus cordatis, siliqua thecaphori longitudine. ① patr. ign.
12. *C. PSORALEÆFOLIA*, herbacea aculeata pubescens, foliis 3-foliolatis petiolis piloso-glandulosis, floralibus ovatis petiolatis, siliquâ demum glabrâ thecaphoro longiore. ①? in Brasiliâ. (v. s. in h. Mus. Par.)
13. *C. LATIFOLIA* (Vahl! ined.) herbacea inermis glabra, foliis infimis simplicibus, mediis 3-foliolatis petiolatis foliolis ovalibus acuminatis, floralibus simplicibus sessilibus ovatis pedicello dimidiò brevioribus, thecaphoro pedicelli longitudine. ①? in Guianâ. (v. s. in h. Vahl.)
14. *C. ROSEA* (Vahl! ined.) herbacea inermis glabra, foliis 5-foliolatis, infimis floralibusque 3-foliolatis, summis ovatis sessilibus, siliquâ glabrâ thecaphori longitudine. ① ad Rio-Janeiro Brasiliæ. (v. s. in h. Juss.)
15. *C. PURPUREA*, herbacea inermis glabra, foliis 3-5-foliolatis, floralibus summis simplicibus ciliatis, ovario glabro thecaphoro duplò longiore. — in Peruvia. An ad sect. sequentem? (v. s. in h. Lamb.)
16. *C. NUMMULARIA*, herbacea inermis pubescens, foliis superioribus simplicibus petiolatis orbiculatis, ovario glanduloso-pubescente, siliqua demum glabrâ thecaphoro longiore. — in Brasiliâ. Pars plantæ inferior ignota. (v. s. in h. Mus. Par.)

Sect. II. SILIQUARIA. — Forsk. æg. 78.

Torus parvus. Thecaphorum breve aut nullum.

* *Foliis simplicibus.*

17. *C. MONOPHYLLA* (Lin. sp. 940.) herbacea pubescens, foliis simplicibus petiolatis lanceolatis basi obtusis, siliquâ puberulâ subsessili ①.
α. Malabarica, Rheed. h. mal. g. t. 34.
β. Zeylanica, Burm. zeyl. t. 100. f. 2. (v. s. in h. Mus. Par.)
18. *C. CORDATA* (Burch! cat. n. 2374.) subherbacea pubescens, foliis simplicibus breviter petiolatis lanceolatis basi cordatis, siliquâ puberulâ subsessili. — Ad cap. B.-Spei. An satis à priore differt? (v. s.)
19. *C. GLAUCA*, herbacea glabra glauca, foliis petiolatis ovatis, floralibus ellipticis, siliquis pendulis levibus breviter spicatis, seminibus hirsutis. 2 in Arabiâ. (v. s. in h. Mus. Par.)
20. *C. GLAUDESCENS*, herbacea glabra glauca, foliis petiolatis ovatis, floralibus ellipticis, siliquis pendulis striatis breviter stipitatis, seminibus glabris. 2 inter Bagdad et Mossul. Olivier et Bruguier. (v. s.)
21. *C. QUINQUENERVIA*, herbacea subpubescens glauca, foliis petiolatis cordatis quinquenerviis, floralibus minimis, siliquis sessilibus ovali-oblongis glanduloso-asperis, seminibus glabris ①? in Persiâ. (v. s. in h. Mus. Par.)
22. *C. SCAPOSA*, herbacea, foliis subradicalibus petiolatis rotundo-ovatis strigoso-pilosis, scapo nudo aspero, bracteis minimis, siliquis erectis sessilibus glabris. 2? in Ægypto. Stam. 4? Habitus ferè Cardamines. (v. s.)
23. *C. PROSERIFOLIA* (Dell fl. eg. t. 36. f. 2.) fruticosa strigoso-pilosa, foliis

- petiolatis obovato-orbiculatis trinerviis piloso-glandulosis, siliquis sessilibus oblongis pilis apice glandulosis hispidis. ♂ in Ægypto. *Roridula* Forsk! æg. 35. *Rorida* Roem. et Sch. syst. 3. p. 13. (v. s.)
24. *C. RAPHANOIDES* (DC. syst. 2. p. 662.) herbacea apice subhispida, foliis lanceolato-linearibus acuminatis basi runcinatis, siliquâ subsessili tereti stylo apiculata glabra. (I) ? in Guineâ. *Raphanus pilosus* Willd! sp. 3. p. 562. *Cleome longifolia* Vahl. herb. *Wormskioldia heterophylla* Thonn. in herb. Vahl. Pili longi rufescentes. (v. s.)
25. *C. GUIANENSIS* (Aubl. guian. 2. t. 273.) herbacea, glabra erecta foliis linearibus, floralibus pedicello longioribus, siliquis subsessilibus levibus. (I) in Guianâ. (v. s. in h. Mus. Par.)
26. *C. PROCUMBENS* (Jacq. am. t. 120.) herbacea glabra procumbens, foliis lanceolato-linearibus, floralibus pedicello ferè brevioribus, siliquis sessilibus teretibus subtorulosis. ♀ in insul. Caribæis. Sloan. jam. t. 123. (v. s.)

** *Folii 3-5-7-foliolatis.*

27. *C. TENELLA* (Lin. f. supp. 300.) herbacea glabra, foliis 3-foliolatis, foliolis linearibus petiolo duplò longioribus, floribus hexandris, siliquis sessilibus tenuibus. (I) in Indiâ. Foliola 4-5 lin. longa. (v. s.)
28. *C. TETRANDRA* (Banks! herb.) herbacea glabra, foliis 3-foliolatis, foliolis linearibus petiolo triplò longioribus, floribus tetrandris, siliquis sessilibus tenuibus. (I) in Novâ-Hollandiâ. Foliola 2-3 lin. longa. (v. s. in h. Banks.)
29. *C. MULTICAULIS* (fl. mex. ic.) herbacea glabra, foliis omnibus sessilibus 3-foliolatis, foliolis oblongo-linearibus obtusiusculis, floribus hexandris, siliquis brevissimè stipitatis. (I) in Mexico.
30. *C. VIOLACEA* (Lin. sp. 940.) herbacea glanduloso-pubescent, foliis 3-foliolatis, foliolis foliisque summis linearibus, siliquis sessilibus puberulis deflexis linearibus stylo brevi acuminatis. (I) in Lusitaniz arenosis. Schkürhr handbt t. 189. f. b. — excl. syn. Goertn. (v. s.)
31. *C. DEFLEXA*, herbacea glanduloso-subpubescens, foliis 3-foliolatis, foliolis foliisque summis oblongo-linearibus, siliquis sessilibus deflexis glabris acuminatis. (I) ? in Peruvîâ. affinis priori. (v. s. in h. Lamb et Ruiz et Pav.)
32. *C. DILLENIANA*, herbacea glanduloso-pubescent, foliis 3-foliolatis, foliolis oblongo-linearibus petioli longitudine, siliquis subsessilibus deflexis puberulis stylo apiculatis oblongo-linearibus. (I) in Oriente. Dill. elth. t. 266. *C. ornithopodioides* auctorum excl. Buxb. syn. (v. s.)
33. *C. ARABICA* (Lin. sp. 939.) herbacea glanduloso-pubescent, foliis 3-foliolatis, foliolis oblongis, siliquis subsessilibus puberulis oblongis deflexis, seminibus hirsutis. (I) ab Arabiâ ad Mauritaniâ. Lin. f. dec. t. 8. *Siliquaria glandulosa* Forsk! descr. 78. (v. s.)
34. *C. IBERICA*, herbacea glanduloso-pubescent, foliis 3-foliolatis breviter petiolatis, siliquis breviter stipitatis puberulis subpendulis torulosis vix pedicello floris longioribus. (I) in Iberiâ circa Tifin. *C. ornithopodioides Ibericag.* Bieb. fl. taur. 2. p. 130. (v. s.)
35. *C. VIRGATA* (Stev! in litt.) herbacea glanduloso-pubescent, foliis 3-foliolatis breviter petiolatis, foliolis oblongo-linearibus, siliquis breviter stipitatis puberulis subpendulis torulosis pedicello floris triplò longioribus. (I) in Persiâ boreali. — Buxb. cent. 1. t. 9. f. 2. ad hanc, aut forsân ad præcedentem. (v. s.)
36. *C. BRACHYCARPA* (Vahl! ined.) suffruticulosa, junior glandulosa, adulta glabra, foliis 3-foliolatis, foliis ovali-oblongis crassis, siliquis sessilibus ovato-teretibus styloiferis. ♀ in Arabiâ. *C. ornithopodioides.* Forsk! fl. arab. n. 402. (v. s. in h. Banks et Vahl.)
37. *C. FOLIOLOSA*, herbacea glabra glauca, foliis subsessilibus 3-foliolatis

- foliolis summisque suborbiculatis, siliquis subsessilibus oblongo-linearibus piloso-glandulosis. ① in Persiâ. (Oliv. et Brug. in h. Mus. Par. v. s.)
38. *C. CANESCENS* (Stew! ined.) herbacea glaucescens substrigoso-pubescent, foliis 3-foliolatis, foliolis foliisque summis oblongo-linearibus, siliquis stipitatis puberulis erectiusculis torulosis pedicello floris æqualibus. ① in Tauriâ merid. (v. s.)
39. *C. PARVIFLORA* (Brown. in Salt. abyss. 65.) herbacea glanduloso-pubescent, foliis 3-foliolatis, foliolis foliisque summis elliptico-ovatis, siliquis sessilibus oblongis glanduloso-pubescentibus styliiferis pedicello floris subbrevioribus ① in Abyssiniâ. (Salt. in h. Banks v. s.)
40. *C. ASPERA* (Kœn! ined.) herbacea piloso-aspera, foliis 3-foliolatis, foliolis oblongis, siliquis sessilibus teretibus glabris stylo acuminatis. ① in Coromandeliâ. Habitus Polanisæ felix sed hexandra. (v. s.)
41. *C. FLAVA* (Banks! herb.) herbacea glanduloso-pubescent subglauca, foliis inferioribus 5-foliolatis, superioribus 3-foliolatis, foliolis oblongis, siliquis sessilibus striatis glanduloso-puberulis. ① in Novâ-Hollandiâ. (v. s. in h. Banks.)
42. *C. RUTIDOSPERMA*, herbacea glabra, foliis omnibus 3-foliolatis, foliolis ovali-oblongis, siliquis glabris stipitatis elongatis apiculatis. ① in Tabago? Habitus Gynandropsidis triphyllæ, sed torus minimus non elongatus. (v. s.)
43. *C. POLYGAMA*, herbacea glabra, foliis petiolatis inferioribus simplicibus, ceteris 3-foliolatis, foliolis ovato-lanceolatis acuminatis subserrulatis, siliquis subsessilibus teretibus glabris. ① in insul. Caribæis.
- α. *C. polygama* Lin. sp. 939. Sloan. jam. t. 124. f. 1. (v. s.)
β. *C. serrata* Lin. sp. 939. Jacq. am. ed. pict. t. 262. f. 73. folium. (v. s.)
44. *C. ACULEATA* (Lin. syst. 3. p. 232.) herbacea glabra, stipulis spinescentibus, foliis 3-foliolatis, floralibus ovatis, siliquis substipitatis teretibus torulosis glabris. ① in Americâ calid. (v. s.)
45. *C. HOUSTONI* (Brown. h. Kew. 4. p. 131.) herbacea glanduloso-velutina, stipulis petiolisque spinescentibus, foliis 3-5-foliolatis, floralibus ovali-oblongis, siliquis breviter stipitatis glanduloso-velutinis. ① in Jamaicâ. — Mart. dec. t. 45. (v. s.)
46. *C. HUMBOLDTI*, herbacea glabra, petiolis longis aculeatis, foliis 3-5-foliolatis, floralibus ovatis breviter petiolatis pedicello brevioribus, siliquis stipitatis glabris stylo apiculatis. — In Americâ calid. *C. parviflora* Humb. Bonpl. et Kunth nov. gen. 5. p. 83. non Br. (v. s. in h. Humb.)
47. *C. DIFFUSA*, (Banks! herb.) herbacea glabriuscula diffusa, stipulis spinescentibus, foliis 3-foliolatis, floralibus ovatis pedicello brevioribus, siliquis oblongo-linearibus thecaphoro sextuplò longioribus. ① in Rio-Janeiro Brasiliæ. (v. s. in h. Banks.)
48. *C. AFFINIS*, herbacea glabriuscula diffusa inermis, foliis 3-foliolatis, floralibus ovatis pedicello brevioribus, siliquis oblongo linearibus thecaphoro decies longiore. ① Hab. cum priore, à quâ vix differt. (v. s. in h. Banks.)
49. *C. RUEBELLA* (Burch! cat. n. 2025. trav. 1. p. 543.) herbacea glanduloso-pubescent, foliis 5-foliolatis, foliolis linearibus glabris glaucis, siliquis sessilibus brevibus subpubescentibus. ① ad cap. B-Spei. (v. s.)
50. *C. ANGUSTIFOLIA* (Forsk! æg. 120.) herbacea glabra apice punctata, foliis infimis 7-foliolatis, summis 3-foliolatis, foliolis lineari-filiformibus, siliquis stipitatis pendulis. ① in Arabiâ felici. *C. filifolia* Vahl! symb. 1. p. 48. (v. s. in h. Banks et Vahl.)

†. *Species non satis notæ.*

51. *C. PARADOXA* (Brown in Salt. abyss. 65.) basi suffruticosa glabra, apice herbacea glanduloso-scabra, foliis 3-5-foliolatis, foliolis glaucis oblongo-linearibus petiolo muricato multò brevioribus. — In Abyssiniâ. (v. s. sine fl. et fr.)

52. *C. CUNEIFOLIA* (Muhl. ex Natt. gen. am. 2. p. 73.) herbacea lævis, foliis simplicibus cuneatis retusis, siliquis stipitatis. ① in Georgiâ. Flores albi, filamentis 6 longis capillaribus.

53. *C?* *ARBOREA* (Schrad. in Goett. anz. 1821. p. 707.) inermis polyandra, staminibus juxta basin stipitis ovarii, foliis ovatis acuminatis. 5 in Paraíba Brasilæ. Arbor 30-pedalis. Flores parvi flavidi. Fruct. ign. An Polanisiæ species?

V. POLANISIA. Rafin. journ. phys. aout 1819. p. 98.

Calyx 4-sepalus patens. Petala 4. Stamina 8-32. Torus parvus. Siliqua in calyce sessilis aut vix stipitata stylo distincto terminata. — Herbar habitu *Cleomis*.

Sect. I. BRACHYSTYLA.

Stylus ovario brevior.

1. *P. DIANTHERA*, glabra subaculeolata, foliis 7-foliolatis, foliolis lineari-filiformibus, staminibus 8, 6 sterilibus, 2 antheriferis, siliquâ stipitatâ pendulâ. ① ad cap. B.-Spei. *Cleome diandra* Burch! trav. 1. p. 548 cat. n. 2103. (v. s.)
2. *P. CHELIDONII*, piloso-hispida, foliis 7-9 foliolatis, foliolis obovato-cuneatis, staminibus 24-32, siliquâ sessili. — In Indiâ orient. *Cleome Chelidonii* Lin. f. suppl. 300. Flores rosei. (v. s. in h. Banks.)
3. *P?* *ANGULATA*, glabra subtriquetra, foliis 5-7-foliolatis, foliolis oblongo-linearibus, siliquâ sessili striatâ. — In Javâ. Flores ignoti, violacei ex Lesch. (in h. Mus. Par. v. s.)
4. *P. OXYPHYLLA*, glandulosa et pilosiuscula, foliis 3-7-foliolatis, foliolis elliptico-oblongis glaucescentibus, staminibus 8-12, siliquâ breviter stipitatâ, striatâ puberulâ pendulâ. — Ad cap. B.-Spei. *Cleome oxyphylla* Burch! cat. n. 1887. (v. s. in h. Burch.)
5. *P. GRAVEOLENS* (Rafin. l. c.) glanduloso-pilosa, foliis 3-foliolatis, foliolis elliptico-oblongis, staminibus 8-12, siliquis oblongis basi attenuatis glanduloso-muricato-pubescentibus. ① in Amer. boreali. *Cleome dodecandra* var. *Canadensis* Lin. sp. 939. — Corn. can. 131. ic. (v. v.)
6. *P. VISCOSA*, piloso-glandulosa, foliis 3-5-foliolatis, foliolis obovato-cuneatis oblongisve, staminibus 8-16, siliquâ oblongâ sessili striatâ glanduloso-hirsutâ. ① in Indiâ orient.
 - a. *Cleome viscosa* Lin. sp. 938. Rheed. mal. 9. t. 23.
 - β. *Cleome icosandra* Lin. sp. 938. Burm! zeyl. t. 99. (v. v.)
7. *P. DODECANDRA*, puberulo-scabra, foliis 3-foliolatis, foliolis glabris elliptico-lanceolatis subserrulatis, staminibus 8-12, siliquâ sessili basi attenuatâ glabrâ. ① in Indiâ orient. Cl. dodecandra. Lin. sp. 939. excl. Sloan. syn. — Burm. zeyl. t. 100. f. 1. (v. s. in Banks.)
8. *P. FELINA*, strigosa, foliis subradicalibus 3-foliolatis, foliolis cuneiformibus, staminibus 28-32, siliquâ subsessili glabrâ. — in Indiâ Orient. *Cleome felina* Lin. f. suppl. 300.

Sect. II. STYLARIA.

Stylus ovario multò longior.

9. *P. UNIGLANDULOSA*, glanduloso-pubescentis, foliis 3-foliolatis, foliolis bracteis-que oblongo-lanceolatis, staminibus 8-16, siliquâ sessili subpuberulâ. ① in Novâ-Hispaniâ ad Acapulco. *Cleome uniglandulosa* Cav. ic. 4. t. 306. (v. v.)

Trib. II. CAPPAREÆ.

Fructus subcarnosus indehiscens. — Frutices aut arborei.

VI. CRATÆVA. Lin. gen. 599.

Calyx 4-sepalus. Petala 4 calyce majora. Stamina 8-28. Torus elon-

gatus aut hemisphæricus. Bacca stipitata tenuiter corticata ovato-globosa, intus pulposa. — Frutices inermes foliis 3-foliolatis.

1. *C. GYANDRA* (Lin. sp. 636.) staminibus 20-24 toro cylindraceo insertis petalis longioribus, bacca ovata, foliolis ovatis acutis. 5 in Jamaicæ fruticetis — Pluk. t. 147. f. 6. (v. s. in h. Banks.)
2. *C. RELIGIOSA* (Forst. prod. 203.) staminibus 20-28 toro annulari insertis petalorum longitudine, bacca ovali, foliolis lanceolato-ellipticis acutis. 5 in Malabariâ, insulis Societatis. — Rheed. mal. 3. t. 42. (v. s. h. Lamb.)
3. *C. LETA*, staminibus 20-24 toro annulari insertis petalis longioribus, bacca ovali, foliolis ovatis acuminatis basi inæqualibus. 5 in Senegaliâ. (v. s. sp. comm. à cl. Gay.)
4. *C. TAPIA* (Lin. sp. 637.) staminibus 8-16 toro cylindraceo insertis petalis ferè et thecaphoro duplò brevioribus, bacca globosa, foliolis ovato-acuminatis basi inæqualibus. 5 in insul. Caribæis, Amer. calidiore. — Plum. gen. t. 21. Pis. bras. t. 69. An stirps Caribæa octandra differt ab Americanâ dodecandra?
5. *C. ADANSONII*, staminibus 12-16 toro abbreviato insertis, bacca globosa, foliolis oblongis acuminatis basi æqualibus. 5 in Senegaliâ. Cr. tapia Adans! in h. Juss. An à prioribus satis differt? (v. s. in h. Juss.)
6. *C. TAPIOIDES*, staminibus 8-16 toro abbreviato insertis petala oblonga stipitata duplò superantibus thecaphorum ferè æquantibus, foliolis ovatis terminali subobtusis, lateralibus acuminatis. 5 in Americâ meridionali. Bertero. Capparis trifoliata Spreng! in h. Balbis. (v. s.)
7. *C. ACUMINATA*, staminibus 12-16 toro abbreviato insertis petala oblonga parum superantibus, foliolis ovatis acuminatissimis, lateralibus basi inæqualibus. 5 in Guianâ Gallicâ. Perrotet. (v. s.)
8. *C. OBOVATA* (Vahl symb. 1. p. 61.) staminibus 12 toro annulari insertis petalis longioribus, bacca cylindraceâ, foliolis obovatis. 5 in Madagascar. — Othrys. Pet. Th. gen. mad. n. 44. (v. s. sine fructu.)
9. *C. FRAGRANS* (Sims bot. mag. t. 596.) staminibus 12-16 toro annulari insertis petalorum longitudine, bacca oblongâ, foliolis ellipticis, petalis longissimis apice undulato crispis. 5 in Sierrâ-Leonâ. C. capparoides Andr. bot. rep. t. 176. Caulis volubilis. (v. s. in h. Lamb. et Banks.)
10. *C. MAGNA*, staminibus 12-24? toro hemisphærico insertis longissimis, bacca ovata, foliolis ovali-lanceolatis, petalis ovatis. 5 in Cochinchinâ. Capparis magna Lour! fl. cochin. ed. Willd. 1. p. 404. (v. s. in h. Mus. Par.)
11. *C. FALCATA*, staminibus 12-16 toro abbreviato? insertis, bacca oblongâ, foliolis lateralibus ad basim falcatis. 5 in Chinâ prope Cantonem. Capparis falcata. Lour. fl. cochin. ed. Willd. 1. p. 405. †
12. *C. RADIATIFLORA*, staminibus ∞ toro annulari insertis, bacca subrotundo-ovali pendulâ, foliolis ovatis obovatisve acuminatis nitidis. 5 in Guyaquil sylvaticis. Capparis radiatiflora fl. per. 5. t. 433. ined. ex Pav. in litt.

VII. NIEBUHRIA.

Calyx 4-sepalus, sepalis æstivatione valvatis. Petala 0 aut calyce breviora. Torus cylindraceus brevissimus. Stam. ∞. Bacca? ovata aut cylindracea, stipitata. — Frutices inermes.

§. 1. Cratævæformes. *Petala nulla. Folia 3-foliolata.*

1. *N. CAFRA*, racemis terminalibus, foliolis oblongis petiolo quadruplò ferè longioribus. 5 ad cap. B.-Spei ad Cafrorum limites. Cratæva cafra. Burch! cat. n. 3678 et 3648. (v. s.)
2. *N. AVICULARIS*, racemis terminalibus, foliolis obovatis petiolo duplò brevioribus.

ribus. ♂ ad cap. B.-Spei. Crataeva avicularis Burch! cat. n. 4445. An satis à priore differt? (v. s. in h. Burch.)

3. N. LINEARIS, racemis terminalibus, foliolis linearibus recurvato-mucronatis utrinque punctato-rugulosis petiolo brevioribus. 5 in Indiâ orient. Capparis apetala Roth nov. sp. p. 238. Bacca globosa.

4. N. OLEOIDES, pedunculis 1-floris axillaribus, foliolis oblongis petiolo quadruplo ferè longioribus. ♂ ad cap. B.-Spei. Olea Burm! in h. Deless. An a N. cafrâ satis differt? (v. s.)

5. N. MADAGASCARIENSIS, pedunculis 1-floris axillaribus, foliolis oblongis petiolo æqualibus. 5 in Madagascar. (Thompson in h. Lamb. v. s.)

§. 2. Capparoïdæ. *Petala parva. Folia simplicia.*

6. N. OBLONGIFOLIA, foliis ovali-oblongis obtusis mucronulatis. 5 in Arabiâ, Indiâ orient. Capparis oblongifolia Forsk! descr. p. 99. Capparis heteroclita Roxb. ined. (v. s.)

7. N. ARENARIA, foliis ovali-oblongis emarginatis mucronulatis glaucis. 5 in Cambay locis arenosis. An satis à priore differt? (v. s. in h. Banks.)

VIII. BOSCIA. Lam. ill. t. 395. non Thunb.

Calyx 4-sepalus. Petala 0. Stamina 12-20 toro brevi inserta. Bacca stipitata globosa 1-sperma. Frutex inermis, floribus parvis corymbosis.

1. B. SENEGALENSIS (Lam. l. c.) ♂ in Senegaliâ. Podoria Senegalensis Pers. ench. 2. p. 5. (v. s.)

IX. CADABA. Forsk. æg. 67.

Calyx 4-sepalus. Petala 4 aut 0. Nectarium ligulæforme e basi tori subcylindracei ortum. Stam. 4-5 basi monadelpha. Bacca stipitata, cylindracea. — Frutices foliis simplicibus.

§. 1. *Floribus apetalis.*

1. C. ROTUNDIFOLIA (Forsk. descr. 68.) inermis apetala foliis orbiculatis ramisque glabris. 5 in Arabiâ, Abyssiniâ. Strœmia rotundifolia Vahl symb. 1. p. 20. (v. s. in h. Lamb.)

2. C. GLANDULOSA (Forsk! descr. 68.) inermis, apetala, foliis subrotundis scabris, ramis apice piloso-glandulosis. 5 in Arabiâ. Strœmia glandulosa Vahl symb. 1. p. 20. — Numerus staminum hujus et prioris ignotus.

§. 2. *Floribus tetrapetalis.*

3. C. LONGIFOLIA, inermis 4-petala 4-andra, foliis oblongo-linearibus glabris. 5 in Abyssiniâ. Strœmia longifolia Brown. in Salt. voy. abyss. 64. (v. s. in h. Banks.)

4. C. INDICA (Lam. dict. 1. p. 544.) inermis 4-petala 4-andra; foliis oblongis glabris mucronatis. 5 in Indiâ orient. Cleome fruticosa Lin. sp. 937. Burm! ind. t. 46. f. 3. (v. s.)

5. C. DUBIA, inermis 4-petala 4-andra, foliis subellipticis pulvere glauco farinosis uninerviis. 5 in Senegaliâ, Abyssiniâ. Strœmia farinosa Brown in Salt. voy. Alys. 64. An satis à sequente differt? (v. s.)

6. C. FARINOSA (Forsk. descr. 68.) inermis 4-petala 5-andra, foliis ovali-oblongis uninerviis pulvere glauco farinosis. 5 in Arabiâ, Senegaliâ. Strœmia farinosa Vahl symb. 1. p. 20. Flores flavi. An hic plures confusæ? (v. s. ex Senegal.)

7. C. TRINERVIA, inermis 4-petala hexandra, foliis trinerviis ovato-orbiculatis. 5 in Persiâ propè Ispahan. (v. s. in h. Mus. Par.)

8. C. CAPPAROIDES, stipulis spinosis, floribus 4-petalis hexandris, foliis oblongo-lanceolatis penninerviis. 5 in Timor et Novâ-Hollandiâ. Lechen. Petala 3-longè unguiculata. (v. s. in h. Mus. Par.)

X. SCHEPPERIA. *Neck. elem. n. 1392.*

Cal. 4-sepalus patens. Petala 4. Torus elongatus. Stam. 8 circa torum monadelphæ apice libera. Nectarium concavum ad basin tori. Siliqua carnosa in calyce stipitata.

1. *S. JUNCEA*. $\frac{5}{2}$ ad cap. Bonæ-Spei. Cleome juncea Lin. suppl. 300. Macromerum junceum Burch! trav. 1. p. 388 et 492. Aphylla aut foliis minimis mox deciduis. (v. s.)

XI. SODADA. *Forsk. descr. 81.*

Calyx 4-sepalus sepalo superiore maximo concavo. Petala 4. Stamina 8. Torus parvus. Ovarium longè stipitatum ovatum 4-sulcatum. — Frutex stipulis spinosis pedicellis plurimis axillaribus.

1. *S. DECIDUA* (Forsk. l. c. Delil! fl. eg. 74. t. 26.) $\frac{5}{2}$ in Arabiâ, Ægypto superiore. — Hombak. Adans. fam. 2. p. 408. (v. s. sine fructu.)

XII. CAPPARIS. *Lin. gen. n. 643.*

Calyx 4-partitus. Petala quatuor. Torus parvus. Thecaphorum gracile. Stamina ∞ . Siliqua subbaccata stipitata. — Frutices foliis simplicibus integris. — Genus dividendum?

Sect. I. EUCAPPARIS. — *Capparis Plum. gen. 39.*

Alabastrum globosum sepalis ovatis concavis obtusis imbricatis inæqualibus. Thecaphorum longum. — Species omnes veteris orbis aut Novæ-Hollandiæ neo Americæ incolæ.

* *Pedicellis axillaribus solitariis aut rarius 2-3 ex eodem puncto ortis, floribus polyandris. PEDICELLARES.*

1. *C. MARIANA* (Jacq. h. Schœnbr. t. 109.) inermis, foliis subrotundis subcordatis glabriusculis, limbo petiolo decies longiore, pedicellis 1-floris solitariis foliorum longitudine. $\frac{5}{2}$ in ins. Marianis, Timor, ins. Maurit? *C. cordifolia* Lam. dict. 1. p. 609. (v. s.)
2. *C. SANDWICHIANA*, inermis, foliis ellipticis glabriusculis, limbo petiolo vix triplò longiore, pedicellis solitariis 1-floris foliorum longitudine. $\frac{5}{2}$ in ins. Sandwich. Menzies. (v. s. in h. Banks.)
3. *C. RUPESTRIS* (Sibth. et Smith fl. gr. t. 487. ex prod. 1. p. 355.) inermis, foliis subrotundis carnosis, pedicellis 1-floris solitariis folio longioribus. $\frac{5}{2}$ in Cretæ Antipari et Myconis rupestribus. (v. s. in h. Mus. Par.)
4. *C. SPINOSA* (Lin. sp. 720.) stipulis spinosis uncinatis, foliis subrotundis retusis, pedicellis solitariis 1-floris. $\frac{5}{2}$ in maris et rupestribus Europæ australioris et Orientis. — Blackw. herb. t. 417. Variat stipulis abortivis, foliis ovatis, plus minusve canis. Alabastra condimentum præbent. (v. v.)
5. *C. FONTANESII*, stipulis spinosis uncinatis, foliis ovatis basi cordatis apice acutiusculis. $\frac{5}{2}$ in fissuris rupium Mauritanie propè Oran. *C. ovata* Desf! fl. atl. 1. p. 404. non Bieb. (v. s. sine fl. in h. Desf!) †.
6. *C. HERBACEA* (Willd. enum 560.) stipulis spinosis, rectis foliis ellipticis mucronulatis glabris, pedicellis solitariis 1-floris folio longioribus, caulibus herbaceis humistratis. $\frac{5}{2}$ in campis Tauricis et Caspico. Caucasica — *C. ovata* Bieb. suppl. 2. p. 1. excl. syn. (v. s.)
7. *C. ÆGYPTIA* (Lam. dict. 1. p. 605.) stipulis spinosis subuncinatis, foliis subrotundo-cuneatis apice mucronatis glabris, pedicellis solitariis unifloris foliorum longitudine. $\frac{5}{2}$ in Ægypti superioris desertis. Delil! fl. eg. p. 93. t. 31. f. 3. (v. s.)
8. *C. ROTUNDIFOLIA* (Rottl. in Willd. berl. mag. p. 185.) stipulis spinosis rectis, foliis sessilibus subrotundo-ovatis subcordatis glabris, pedunculis solitariis 1-floris, capsulis globosis. $\frac{5}{2}$ in Indiâ orient. †.

9. *C. NAPAULENSIS*, stipulis spinosis subuncinatis, foliis ovatis brevissimè petiolatis cum ramis canescentibus, pedunculis axillaribus 1-floris folio longioribus, ovario oblongo. 5 in Napauliâ. Wallich. (v. s.)
10. *C. NUMMULARIA*, stipulis spinosis rectis, foliis ovato-orbiculatis mucronatis emarginatisve glabris petiolo duplò longioribus. 5 in Novæ-Hollandiæ insulis sterilibus. (v. s. sine fl. et fr.)
11. *C. CANESCENS* (Banks! ined.) stipulis spinosis rectis, foliis ovatis subacutis velutino-puberulis petiolo sextuplò longioribus, pedicellis solitariis unifloris foliorum longitudine. 5 in Novâ-Cambriâ australi. Calyx ignotus. (v. s. in h. Banks.)
12. *C. HETERACANTHA*, stipulis spinosis, aliâ uncinatâ, alterâ longiore rectâ, foliis orbiculatis velutinis brevissimè petiolatis 5 inter Bagdad et Alep. Olivier et Brug. Folia ferè 3-nervia. (v. s. sine fl. in h. Mus. Par.) †.
13. *C. LEUCODAPHYLLA*, stipulis spinosis uncinatis, foliis orbiculatis cano-velutinis brevissimè petiolatis pedicellis solitariis 1-floris folio longioribus. 5 inter Bagdad et Alep. Oliv. et Brug. fructus glaber venis 6 longitudinalibus. (v. s. in herb. Mus. Par.)
14. *C. TOMENTOSA* (Lam. dict. 1. p. 606.*) stipulis spinosis uncinatis, foliis ovato-oblongis obtusis tomentoso-velutinis, pedicellis solitariis 1-floris folio brevioribus. 5 in Senegal. Sparmann. (v. s. in h. Juss.)
15. *C. DEALBATA*, stipulis spinosis brevibus uncinatis, foliis ovato-lanceolatis acuminatis, junioribus subvelutinis, demùm supernè glaberrimis, pedicellis solitariis 1-floris petioli longitudine. 5 in Timor. (v. s.)
16. *C. BREVISPINA*, stipulis spinosis brevibus rectis, foliis ovato-lanceolatis mucronatis glabris, pedicellis solitariis 1-floris petiolo triplò longioribus. 5 in Indiâ orientali. Roxburgh. (v. s.)
17. *C. ERYTHROCARPA* (Isert. berl. natur. 9. p. 289. t. 9.) stipulis spinosis reflexis, foliis oblongis acuminatis, junioribus ramisque pube stellata subvelutinis demùm glabris, pedicellis solitariis 1-floris petiolo brevissimo longioribus. 5 in Guineâ (v. s. in h. Desf.)
18. *C. RHEEDI*, stipulis spinosis rectis, foliis ovali-oblongis mucronatis glabris brevissimè petiolatis subtus reticulatis, pedicellis solitariis 1-floris folii circiter longitudine. 5 in arenosis Malabariz. Badukka Rheed. mal. 6. t. 57. Differt à *C. baducca* Lin. in quâ plures sunt confusæ. (v. s. in h. Banks.)
19. *C. AFZELII*, stipulis spinosis uncinatis, foliis ovato-lanceolatis, junioribus subpuberulis, adultis glabris, brevissimè petiolatis, pedicellis solitariis 1-floris. 5 in Africæ orâ occidentali. Afzel. Brass. priori affinis. (v. s. in h. Banks.)
20. *C. PYRIFOLIA* (Lam. dict. 1. p. 606.*) stipulis spinosis brevibus uncinatis, foliis ovali-lanceolatis acutis, junioribus subtomentosis, adultis glabris, pedicellis solitariis 1-floris folio multò brevioribus. 5 in Indiâ orientali. (v. s. in h. Juss.)
21. *C. STYLOSA*, stipulis spinosis patentibus, foliis oblongo-lanceolatis mucronatis glabris, pedicellis solitariis 1-floris petiolo triplò longioribus, sepalis acuminatis, ovario in stylum crassum attenuato. 5 in aridis Coromandeliz Koenig. (v. s. in h. Banks.)
 β. *velutina*, pube brevi subcanescens. In Mysore Buchanan. (v. s. in h. Banks.)
22. *C. HORRIDA* (Lin f. suppl. 264.) stipulis spinosis patentibus, foliis ovatis mucronatis, pedicellis 1-floris suprâ-axillaribus 2-4 aggregatis. 5 in Coromandelâ, Ceylonâ. (v. s. in h. Banks.)
23. *C. PUBIFLORA*, stipulis spinosis minimis rectis, foliis ovali-oblongis acuminatis glabris, pedicellis 1-3 axillaribus petiolo duplò longioribus. 5 in Timor? Java. Petala pubentia. (v. s. in h. Mus. Par.)
24. *C. APHYLLA* (Roth nov. sp. p. 238.) stipulis spinosis, foliis nullis? ap-

bellis in stipularum axillis subsessilibus, caule flexuoso dichotomo cum ramis divaricatis aphylo glaberrimo. ζ in Indiâ orientali.

** *Pedicellis unifloris supra-axillaribus in seriem verticalem dispositis. SERIALES.*

25. *C. ZEYLANICA* (Lin. sp. 720.) stipulis spinosis rectis, foliis ovalibus utrinque acuminiatis glabris, pedicellis 2-3 seriatis petiolo triplò longioribus. ζ in Ceylonâ, Javâ. Flores parvi 12-andri. An *C. Zeylanica* Lour. eadem? (v. s.)

26. *C. BILLARDIERII*, stipulis spinosis rectis, foliis ovali-oblongis venoso-reticulatis ramulisque glabris, pedicellis 3 seriatis, ovario subsessili. ζ in ins. Moluccanis ad fretum Bouton. (v. s. in h. Desf.)

27. *C. ACUMINATA* (Willd. sp. 2. p. 1131.) stipulis spinosis apice uncinatis, foliis ovatis mucronato-acuminatis subtus ferrugineo-tomentosis, pedicellis 2-3 seriatis. ζ in Indiâ orientali. Petala pube ciliata. Huc forsau referenda t. 29. Braam io. chin. 1821.

28. *C. VOLKAMERIE*, stipulis spinosis apice uncinatis, foliis ovatis mucrone calloso apiculatis subtus secus nervos rufo-velutinis, pedicellis 2-3 seriatis. ζ ad cap. B-Spei. Volkameria Capensis Burml prod. cap. 17. Petala villis ciliata. Stam. circiter 30. (v. s. in h. Deless.)

29. *C. LASIANTHA* (Brown! ined.) stipulis spinosis rectiusculis, foliis ovato-oblongis brevissimè petiolatis demum glabris, junioribus ramulis calycibusque velutinis, pedicellis 2-3 seriatis. ζ in Novæ-Hollandiæ orâ orientali inter tropicos. (v. s.)

30. *C. TERNIFLORA*, stipulis spinosis uncinatis, foliis ovatis mucronatis demum glabris, junioribus pubescentibus, pedicellis 3 seriatis, sepalis velutinis. ζ in Coromandeliâ. (v. s. in h. Mus. Par.)

31. *C. QUADRIFLORA*, stipulis spinosis rectiusculis, foliis ovato-acuminatis calloso-apiculatis superne glabris, subtus ramulis calycibusque velutinis, pedicellis 4 seriatis. ζ in Indiâ orientali. *C. pyrifolia* β . Lam. dict. 1. p. 606. — Pluk. t. 107. f. 3. (v. s.)

32. *C. QUINIFLORA*, stipulis nullis, foliis ovatis subacuminatis junioribus pubescentibus demum glabris, pedicellis 4-6 seriatis. ζ in Novæ-Hollandiæ orâ boreali. (v. s. in h. Mus. Par.)

33. *C. MICRANTHA*, stipulis spinosis parvis rectis, foliis ovalibus obtusis mucrone calloso apiculatis, pedicellis 4-6 seriatis. ζ in Javâ. (v. s.)

*** *Pedicellis in corymbum racemumve dispositis, floribus polyandris. CORYMBOSÆ.*

34. *C. CITRIFOLIA* (Lam. dict. 1. p. 606.) stipulis spinosis uncinatis, foliis oblongis mucronatis glabris, pedicellis umbellatis terminalibus. ζ ad cap. B-Spei. Variat ramulis subpuberulis.

35. *C. SEPIARIA* (Lin. sp. 720.) stipulis spinosis uncinatis, foliis ovatis submarginatis subtus ramulisque pubescentibus, pedicellis umbellatis terminalibus. ζ in Indiâ orientali. — Pluk. t. 338. f. 3. (v. s.)

β . *glabrata*, foliis adultis omninò glabris. ζ in Timor. (v. s.)

36. *C. UMBELLATA* (Brown! ined.) inermis, foliis ovalibus obtusis submarginatis glabris, ramulis petiolisque pubescentibus, pedicellis umbellatis terminalibus. ζ in Novæ-Hollandiæ orâ orientali inter tropicos. (v. s.)

37. *C. INCANESCENS*, stipulis spinosis uncinatis, foliis ellipticis mucronulatis, ramulisque pube adpressâ canescentibus, pedicellis umbellato-corymbosis terminalibus. ζ in Mysore. (v. s. in h. Banks)

38. *C. CORYMBOSA* (Lam! dict. 1. p. 605.) stipulis spinosis uncinatis, foliis ovalibus ovobovatis glabris aut subtus pubescentibus, pedicellis corymbosis terminalibus. ζ in Senegaliâ. Specimina foliis subtus glabris forsau specificè distinguenda? (v. s.)

39. *C. ROXBURGHII*, stipulis ramorum spinosis uncinato-reflexis, ramulorum

- nullis, foliis ellipticis glabris, ramulis cinereo-velutinis, pedicellis racemoso-corymbosis. ζ in Indiâ orientali. (C. aguba herb. Banks. v. s.)
40. *C. OBOVATA* (Burch! ined.) inermis, foliis obovatis subtus ramulisque cinereo-velutinis, pedicellis racemoso-corymbosis. ζ in Mysore. (v. s. in h. Banks.)
41. *C. MAXIMA* (Roth nov. sp. 237.) inermis, foliis subrotundo-ovatis utrinque parum attenuatis, supra glabris nitidis, subtus ramulisque pubescentibus cinerascens, racemis terminalibus subpaniculatis. ζ in Indiâ orientali.
42. *C. GRANDIS* (Lin. f. suppl. 263.) inermis, foliis ovatis, ramulisque junioribus subvelutinis, adultis glabris, racemis terminalibus. ζ in Ceylonâ. Fractus globosus pisi magnitudine. (v. s. in h. Deless. Banks.)
43. *C. RAEMIFERA*, stipulis ramorum subspinosi inæqualibus, ramulorum nullis, foliis obovatis supernè glabris, subtus ramulisque junioribus pubescentibus, racemis terminalibus. ζ in Indiâ orientali. (v. s. in h. Lamb.)
44. *C. PUBERULA*, stipulis ramorum spinosis uncinatis, ramulorum punctiformibus, foliis ovato-lanceolatis supernè glabris, subtus ramulisque subvelutinis, racemis terminalibus. ζ in Africæ calidioris orâ occidentali. Brass. (v. s. in h. Banks.)
45. *C. BRASSI*, stipulis spinosis uncinatis, foliis ovato-lanceolatis glabris, ramulis petiolisque pubescentibus, racemis corymbosis axillaribus. ζ Hab. cum priore. (v. s. in h. Banks.)
46. *C. FASCICULARIS*, stipulis ramorum spinosis subuncinatis, ramulorum nullis, foliis ellipticis, adultis glabris, racemis fasciculato-corymbosis axillaribus. ζ cum prioribus. (v. s. in h. Banks.)
47. *C. LANCEOLARIS*, stipulis spinosis uncinatis, foliis lanceolatis acutis, racemis axillaribus subcorymbosis, pedicellis basi utrinque stipulâ spinosâ instructis. ζ in Java. Leschenault. (v. s. in h. Mus. Par.)
- *** *Pedicellis axillaribus sæpius multifloris, floribus octandris.* OCTANDRÆ. — An sectio propria? — Spec. omnes Capenses.
48. *C. CLUTTIFOLIA* (Burch! cat. n. 388.) inermis, foliis oblongo-cuneatis obtusis cum mucrone glabris, pedicellis axillaribus solitariis 1-floris folio dimidio brevioribus. ζ ad cap. B.-Spei. (v. s. in h. Burch.)
49. *C. OLEOIDES* (Burch! cat. n. 4200.) inermis, foliis oblongis basi attenuatis obtusis cum mucrone glabris, racemis axillaribus folio paulò brevioribus, thecaphoro pedicello brevior. ζ ad cap. Bonæ-Spei. Bacca globosa pisi magnit. (v. s. in h. Burch.)
50. *C. CORIACEA* (Burch! cat. n. 2898.) inermis, foliis oblongis obtusis glabris, racemis axillaribus folio paulò brevioribus. ζ Hab. cum priore à quâ vix differt. (v. s. sine fl. et fr. in h. Burch.)
51. *C. ALBITRUNCA* (Burch! cat. n. 1762. trav. i. p. 343.) inermis, foliis oblongis obtusissimis basi attenuatis glabris, racemis axillaribus folio brevioribus, thecaphoro pedicelli longitudine. ζ cum prioribus. (v. s. in h. Burch.)
52. *C. PUNCTATA* (Burch! cat. n. 1891. trav. i. p. 492.) inermis, foliis oblongis basi subattenuatis apice submucronatis glabris venoso-reticulatis, racemis axillaribus folio multò brevioribus. ζ cum prioribus. (v. s. in h. Burch.)
53. *C. RACEMULOSA*, stipulis spinosis minimis, foliis ovali-oblongis glabris, racemis axillaribus folio triplò longioribus. ζ in Africâ australi. Masson. Flores apetalî 8-12-andri. An hujus sectionis? (v. s. in h. Banks.)

SECT. II. CAPPARIDASTRUM.

Alabastrum subovatum sepalis parvis subrotundis æqualibus per æstivationem imbricatis. Thecaphorum mediocre. — Rami inermes. Folia glabra. — Species *Americanæ*.

54. *C. FRONDOSA* (Jacq. pl. am. 162. t. 104.) foliis coriaceis lanceolatis acuminatis basi subcordatis petiolo triplò et ultrà (Balb.) longioribus, pedicellis ad apicem pedunculi corymbosis. ♀ in sylvaticis Carthagenæ, Sti.-Domingi et Novæ-Granadæ (Humb.)
55. *C. CUNEATA*, foliis coriaceis obovato-cuneatis mucronatis petiolo triplò longioribus, floribus racemosis, thecaphoro pedicello longiore. ♀ in insulis Caribæis. (v. s. in h. Banks ubi *C. petiolaris* dicitur.)
56. *C. MACROPHYLLA* (Humb. B. et Kunth! nov. sp. 5. p. 91.) foliis coriaceis ovalibus utrinquè subacutis petiolo duplò longioribus, floribus racemosis longè pedicellatis. ♀ ad ripas flum. Magdalenzæ. (v. s. in h. Humb.)
57. *C. PETIOLARIS* (Humb. B. et Kunth! nov. sp. 5. p. 91.) foliis membranaceis ovali-oblongis obtusis petiolo duplò longioribus, floribus racemosis, thecaphoro pedicello duplò longiore. ♀ prope Guayaquil. (v. s. in h. Humb.)
58. *C. TRIFLORA* (Mill! dict. n. 10.) foliis elliptico-oblongis utrinquè acuminatis petiolo triplò longioribus, pedunculis 3-floris folio multò brevioribus ♀ in Americâ. calid. circa Tolu. (v. s. in h. Banks.)
59. *C. BRASILIANA*, foliis oblongis acuminatis basi attenuatis et subcordatis subsessilibus, pedunculis racemosis, thecaphoro pedicello vix brevior. ♀ in Brasiliâ prope Rio-Janeiro. (v. s. in h. Banks.)

Sect. III. CYNOPHALLA.

Alabastrum globosum, sepalis subrotundis per æstivationem imbricatis ad basim glandulâ foveolâve instructis. Siliqua longa teres carnosâ. Thecaphorum elongatum.—Species Americanæ, glabræ, inermes, in axillis sæpius glandulosæ.

60. *C. SÆSSILIS* (Banks! herb.) foliis subsessilibus ovalibus, basi cordatis, glandulâ axillari nullâ aut concavâ, pedicellis axillaribus solitariis unifloris. ♀ in Carthagenâ Indorum. (v. s. in h. Banks.) †.
61. *C. CYNOPHALLOPHORA* (Lin. sp. 721. Jacq. am. t. 98.*) foliis glabris coriaceis oblongis breviter petiolatis, glandulâ axillari ovato-oblongâ, pedunculis paucifloris folio brevioribus. ♀ in ins. Caribæis et Amer. calidiore. *Capparis flexuosa* Lin! sp. 722. Species polymorpha forsân stirpes heterogeneas colligens. (v. s.)
62. *C. EUSTACHIANA* (Jacq. am. ed. pict. t. 146.) foliis glabris subcoriaceis ovatis acuminatis breviter petiolatis; glandulâ axillari ovato-oblongâ, pedunculis paucifloris folio brevioribus. ♀ in ins. Sti.-Eustachii. Flos purpureus.
63. *C. SALIGNA* (Vahl symb. 3. p. 66.*) foliis glabris coriaceis breviter petiolatis linearibus basi subdilatis cordatisque, glandulâ axillari ovato-oblongâ. ♀ in ins. Sauctæ-Crucis. (v. s.)
64. *C. HASTATA* (Lin. sp. 722. Jacq. am. t. 174. f. 56.) foliis glabris coriaceis breviter petiolatis basi hastatis subcordatisve hinc longè linearibus, axillis eglandulosis, racemis terminalibus. ♀ in sylvis Carthagenæ. (v. fol. in h. Banks.)
65. *C. LINEARIS* (Jacq! am. t. 102.*) foliis glabris coriaceis brevissimè petiolatis linearibus, axillis glandulosis, racemis subpaniculatis. ♀ in sylvaticis Carthagenæ et Cumanæ. (v. s. in h. Banks.)
66. *C. LANCEOLATA* (Ruiz. et Pav. fl. per. t. 423. f. a. ined.) foliis glabris lanceolatis mucronulatis, pedunculis paniculatis, bracteis basi glanduliferis. ♀ in Guayaquil ad Daule ripas. Siliqua 2 palmas longa.
67. *C. GUAYAQUILENSIS* (Humb. B. et Kunth! nov. sp. 5. p. 89.*) foliis glabris oblongis obtusis subretusis breviter petiolatis, glandulâ axillari truncatâ, racemis terminalibus. ♀ in Guayaquil. Siliqua 3-4 pollices longa torulosa. (v. s. in h. Humb.)
68. *C. LÆTA* (Humb. B. et K! nov. sp. 5. p. 88.*) foliis glabris oblongis utrinquè

quæ obtusis subemarginatis brevissimè petiolatis, glandulâ axillari truncatâ, racemis infernè ramosis 5 circa Turbaco. (v. s. in h. Humb.)

69. *C. SUBBILOBA* (Humb. B. et K! nov. sp. 5. p. 90.*) foliis glabris ellipticis apice emarginato-bilobis breviter petiolatis glandulâ axillari obtusis, racemis paniculatis abbreviatis. 5 circa Cumanam. (v. s. in h. Humb.) †.
70. *C. PAUCIFLORA* (Humb. B. et K! nov. sp. 5 p. 89.*) foliis glabris coriaceis ellipticis utrinquè obtusis subemarginatis, glandulâ axillari truncatâ, pedunculis 3-floris. 5 prope Cumanam in umbrosis. Flor. ignoti. Siliqua 2-3-pollices longa. (v. s. in h. Humb.)
71. *C. VERRUCOSA* (Jacq. am. t. 99.*) foliis glabris oblongis basi obtusis apice subacutis in axillâ eglandulosis, racemis terminalibus paucifloris. Siliquâ verrucosâ. 5 in sylvis Carthagenæ et Sancti-Thomasii. Siliqua sesquipollicaris. (v. s. in h. Deless.)
72. *C. MOLLIS* (Humb. B. et K! nov. sp. 5. p. 88.) foliis oblongis breviter petiolatis basi subovatis apice attenuatis supernè glabris, subtus ramulisque velutinis, glandulâ axillari obtusâ, racemis terminalibus 7-8-floris. 5 in Peruvîâ ad ripas Maceræ. Fruct. ignot.
73. *C. HETEROPHYLLA* (Ruiz. et Pav. ined. ex litt.) foliis glabris oblongo-obovatis mucronatis emarginatisve, glandulâ axillari pyxidiformi, racemis terminalibus subcorymbosis, fructibus clavatis. 5 in Guayaquil sylvis. Flores parvi. Fructus pollicares.
74. *C. AMPLISSIMA* (Lam. dict. 1. p. 607.) foliis glabris ovalibus breviter petiolatis, glandulâ ovatâ, pedicellis paucis solitariisve ad apices ramorum, fructibus ovatis. 5 in Hispaniolâ et Mexico. Plum. ed. Burm. t. 73. f. 2. excl. syn. (v. s. in h. Lamb.)
75. *C. SAXATILIS* (Humb. B. et K! ined.) foliis glabris ovali-oblongis retusis breviter petiolatis, pedicellis laxè racemosis, fructibus ovatis. 5 in fissuris rupium Angusturæ.
76. *C. MACROCARPÂ* (Ruiz. et Pav. ined. t. 540. ex litt.) foliis utrinquè glabris ovalibus mucronatis, glandulis calycinis maximis, baccâ subrotundâ. 5 in Guayaquil sylvaticis et prope Loxa. An sectio propria? (v. s. in h. Lamb.)

Sect. IV. CALANTHEA.

Sepala sublinearia acuta reflexa basi distantia. Fructus subrotundus.—Species Americanæ.

77. *C. PULCHERRIMA* (Jacq. am. t. 106.) foliis glabris oblongis, obtusis brevissimè petiolatis in axillâ eglandulosis, racemo terminali simplici, fructu stylo subacuminato. 5 in declivibus Carthagenæ. *C. arborescens* Mill. dict. n. 3. Petala e flavo-albentia tomentosa oblonga. (v. s. sine fl. in h. Banks.)
78. *C. NEMOROSA* (Jacq! am. ed. pict. t. 156.) foliis ovato-acuminatis supernè glabris subtus tomento deciduo vestitis, fructu ovato-oblongo. 5 in Carthagenæ sylvis. Flor. ign. (v. fol. in h. Banks.)

Sect. V. BREYNIATRUM. Breyniæ sp. Plum. gen. t. 16. non Forst.

Sepala parva petalis multò minora triangularia acuta, Bacca oblonga. Thecaphorum breviusculum. Species Americanæ inermes, pedunculis axillaribus multifloris.

* *Floribus polyandris.*

79. *C. AMYGDALINA* (Lam. dict. 1. p. 608 excl. syn. Jacq. et Lin.) foliis elliptico-oblongis utrinquè attenuatis calloso-mucronatis supernè glabris subtus ramulisque lepidotis, pedunculis axillaribus compressis corymbiferis. 5 in insulis Caribæis et Amer. continente.—Breyn. ic. rar. p. 13. ic. Plum. gen. 40. t. 16. Siliqua pedem longa. (v. s.)

β. *unbellata* (H. B. et K!), differt foliis paulò angustioribus.

80. *C. BARCELONENSIS* (Humb. B. et K! nov. sp. 5. p. 97.*) foliis elliptico-

oblongis acuminatis, supernè junioribus pubescentibus, subtùs ramulisque lepidotis, pedunculis axillaribus racemiferis. ζ in fruticetis Novæ-Barcellona. (v. s. in h. Humb.)

81. *C. COMOSA* (Jacq. *am.* 160.*) foliis lanceolatis petiolatis supernè glabris subtùs cinereo-tomentosis, pedunculis multifloris, staminibus corollà brevioribus. ζ in inundatis maritimis Sancti-Domingi.
82. *C. ODORATISSIMA* (Jacq. h. *schœn.* t. 110.*) foliis oblongis acuminatis longiusculè petiolatis supernè glabris subtùs piloso-lepidotis, pedunculis apice racemiferis, staminibus 28-32 corollæ longitudine. ζ ad Caracas. (v. s.)
83. *C. PACHACA* (Humb. B. et K! *nov. sp.* 5. p. 93.*) foliis ovato-ellipticis submarginatis coriaceis glabris brevissimè petiolatis, racemis terminalibus paucifloris, fructu subgloboso pendulo. ζ in sylvis Cumanæ. (v. s. in h. Humb.)
84. *C. SCABRIDA* (Humb. B. et K! *nov. sp.* 5. p. 95.*) foliis oblongis basi obtusis apice acutiusculis supernè scabridis subtùs subvelutinis longè petiolatis, ramis racemisque tomentosis, staminibus 24 basi villosis. ζ in America calidiore. Fructus ovato-oblongus sesquipollicaris. Sepala ovali-oblonga in alabastrum ovatum conniventia. An hujus sectionis? (v. s. in h. Humb.)
85. *C. TENUSILYQUA* (Jacq! *am. t.* 105.) foliis lanceolatis ovatisve glabris petiolatis racemis terminalibus multifloris, staminibus 16-24 siliquisque longissimis. ζ Carthagenæ frequens. (v. s.)
86. *C. OBOVATIFOLIA* (Humb. B. et K! *nov. sp.* 5. p. 92.*) foliis elliptico-obovatis subcordatis glabris longè petiolatis, racemis subaxillaribus multifloris, siliquâ elongatâ pendulâ. ζ propè Cumanam. (v. s. in h. Humb.)
87. *C. BREVISILYQUA* (fl. *mex. ic. ined.*) foliis ellipticis obtusis emarginatis glabris breviter petiolatis, pedicellia terminalibus 2-3 umbellatis, siliquâ torulosâ thecaphoro duplò longiore. ζ in Novâ-Hispaniâ. Petala oblongo-spathulata ex albo-virescentia.

** *Floribus octandris.*

88. *C. ANGUSTIFOLIA* (Humb. B. et K! *nov. sp.* 5. p. 96.*) foliis linearibus basi subcordatis brevissimè petiolatis, supernè glabris subtùs subpubescentibus, pedunculis corymbosis ramulisque velutinis. ζ inter Mexico et Acapulco. Ramuli angulato-sulcati. Alabastra ovata tomentosa. (v. s. in h. Humb.)
89. *C. FERRUGINEA* (Lin. *amœn.* 5. p. 398.*) foliis lanceolatis acuminatis petiolatis supernè glabris subtùs piloso-lepidotis, ramulis tomentosis, pedunculis axillaribus apice corymboso-racemosis. ζ in fruticetis maritimis Jamaicæ et Sancti-Domingi. Brown *jam. t.* 28. f. 1. *C. octandra* Jacq. *am. t.* 100. Flores albi, inodori, fragrantès, aut fœtidi à diversis dieti! (v. s.)
90. *C. INCANA* (Humb. B. et K! *nov. sp.* 5. p. 94.*) foliis ovato-oblongis acuminatissimis petiolatis ramulisque adpressè cano-velutinis, racemis paucifloris folio subbrevioribus. ζ inter Mescala et Estola in Amer. calid. Petala extùs tomentosa. Stam. filamenta brevissima. Fructus ovatus cano-tomentosus. (v. s. in h. Humb.)
91. *C. CORDATA* (Ruiz. et Pav. *t.* 432. *ined. ex litt.*) foliis cordato-subrotundis utrinquè pube stellatâ tomentosis, pedunculis racemosis paucifloris, baccâ subglobosâ oligospermâ. ζ in Guayaquil tractibus.

Sect. VI. QUADRELLA.

Alabastrum subquadratum, sepalis per æstivationem valvatis marginibus in suturas prominulas subrevolutis. Siliquæ dehiscentes. Thecaphorum elongatum. — Species Americanæ inermes, velutinæ seu lepidotæ, pedunculis axillaribus multifloris.

92. *C. CAOTONOIDES* (Humb. B. et K! *nov. sp.* 5. p. 95.* t. 437.) foliis cordato-subrotundis petiolatis supernè villosis subtùs ramisque tomentosis, racemis

- angulatis terminalibus paucifloris, staminibus 8, fructu globoso. § 5 in calidis ad flumen Magdalenæ. (v. s. in h. Humb.)
93. *C. FURFURACEA* (Ruiz. et Pav! in h. Lamb.) foliis elliptico-oblongis obtusis petiolatis supernè glabris, subtùs ramis petiolisque velutinis, racemis terminalibus angulatis multifloris, staminibus 12 basi hispidis. § in Mexico. (v. s. in h. h. Lamb.)
94. *C. SIDAIFOLIA* (Ruiz. et Pav! in h. Lamb.) foliis ovatis basi cordatis petiolatis supernè adultis glabris, subtùs ramisque pube-stellatâ velutinis, racemis terminalibus folio brevioribus subcorymbosis. § in Peruvîâ. Stam. et fruct. ignot. (v. s. in h. Lamb.)
95. *C. BREYNIA* (Lin! sp. 721. Jacq. am. t. 103.) foliis ellipticis coriaceis petiolatis supernè glabris subtùs ramulisque lepidotis, pedunculis angulatis racemoso-corymbosis, staminibus 16, siliquâ longissimâ. § in fruticetis ins. Caribæarum et continentis Amer. Breynia Indica et *C. cynophallophora* Lin. sp. ed. 1. p. 503 et 504. (v. s.)
- β. *uniflora*, *C. siliquosa* Lin. spec. 721. excl. syn. Hæc var. habet thecaphorum brevius et folia acutissima. (v. s.)
96. *C. JAMAÏCENSIS* (Jacq! am. t. 101.) foliis oblongis emarginatis coriaceis petiolatis supernè glabris subtùs ramisque cinereo-tomentosis, pedunculis paucifloris, staminibus 20. § § in Jamaicâ. Fruct. prioris? cujus forsân var? (v. s. in h. Balbis.)
97. *C. TORULOSA* (Sw. fl. ind. occ. 2. p. 932.*) foliis ovato-lanceolatis coriaceis petiolatis supernè glabris subtùs ramulisque lepidotis, pedunculis paucifloris racemoso-corymbosis, staminibus 28, siliquâ longissimâ. § in fruticetis montosis Jamaicæ et Barbados. (v. s.)
98. *C. INTERMEDIA* (Humb. B. K! nov. sp. 5. p. 98.*) foliis ovato-lanceolatis coriaceis petiolatis supernè glabris subtùs ramulisque lepidotis, pedunculis paucifloris apice racemoso-corymbosis, siliquis terebinis, thecaphora brevî crasso. § in Cumanâ ubi dicitur Olivo (v. s. in h. Humb.) †.

†. *Capparides quoad sectionem incertæ.*

* *Veteris orbis.*

99. *C. DIVARIGATA* (Lam. dict. 1. p. 606.*) stipulis spinosis uncinatis divaricatis, foliis glabris linearibus acutis brevissimè petiolatis. § in Coromandelîâ. Certè ad Eucapparidem pertinet sed flores, fructusque ignoti. (v. s. in h. Juss.)
100. *C. MITRIDATICA* (Forsk. descr. 99.*) foliis lineari lanceolatis obtusis glabris pendulis petiolatis. § in Arabiâ circa Surdud. An Capparis?
- N. B.* Capparidem Dahim ejusdem auctoris omisi, videtur *Zygophyllum* aut *Cæsalpinia*!

** *Americanæ* (omnes inermes).

101. *C. NITIDA* (Ruiz. et Pav. ined. ex litt.) foliis oblongis acuminatis nitidis glabris petiolatis, pedunculis axillaribus 1-floris, fructu ovali. § in Peruvîâ Andium nemoribus. Flores lutei.
102. *C. LINEATA* (Pers! ench. 2. p. 60.) foliis ovatis acuminatis breviter petiolatis supernè demùm glabris subtùs ramisque tomentoso-velutinis, pedunculis axillaribus 1-floris. § in Brasiliâ ad Rio-Janeiro. Etiamsi inermis videatur ad Eucapparides referenda, et ideò patria mihi dubia videtur. (v. s. in h. Juss.)
103. *C. FRUTICOSA* (Mill. dict. n. 7.) foliis lanceolatis acutis coriaceis breviter petiolatis, floribus axillaribus breviter pedicellatis. § in Amer. calid. propè Toln. †.
104. *C. AVICENNIAEFOLIA* (Humb. B. et K! nov. sp. 5. p. 94.*) foliis ellipticis utrinquè obtusis brevissimè petiolatis supernè glabris subtùs junioribus canescentibus, racemis folio brevioribus, calyce campanulato 4-lobo, sta-

minibus vix petala superantibus. Fructus globosus. 5 in arenosis Amer. calid. ad littora maris Pacifici et circa Guayaquil.

105. *C. LONGIFOLIA* (Sw. prod. 81.) foliis linearibus acuminatis breviter petiolatis supernè glabris subtùs ramulisque lepidotis. 5 in Jamaicà et Barbados. — Pluk. t. 327. f. 6. Verosimiliter *Quadrellæ* species? (v. s. sine fl.)
106. *C. DOMINGENSIS* (Spreng! in h. Balb.) foliis oblongis breviter petiolatis supernè glabris subtùs ramulisque tomentoso-velutinis, pedunculis corymbosis paucifloris, fructu-ovato velutino, thecaphoro velutino. 5 in Santo-Domingo. Bertero. Flores gnoti. An *Breyniastri* aut *Quadrellæ* species? (v. s.)
107. *C. LAURINA* (Humb. B. et K! nov. sp. 5. p. 93.*) foliis oblongis utrinquè obtusis petiolatis coriaceis glabris, racemis terminalibus, fructu globoso pendulo. 5 in calidis Americæ prope Guancabamba. Flores violacei. (v. s. sine fl. in h. Humb.)
108. *C. ANGULATA* (Ruiz. et Pav. fl. per. t. 431. ined. ex litt.) foliis ovato-oblongis mucronatis petiolatis supernè nitidis subtùs ramisque pube 10-radiata velutinis, pedunculis terminalibus corymboso-racemosis, fructu ovato angulato. 5 in Guayaquil montibus. Stamina filamenta basi submonadelpha.
109. *C. VIRIDIFLORA* (Humb. B. et K! nov. sp. 5. p. 92.*) foliis ovali-oblongis mucronulatis longè petiolatis supernè glabris subtùs ramulisque pubescentibus, racemis terminalibus angulatis, sepalis obovatis brevissimis. 5 in umbrosis prope Cumanam. (v. s. in h. Humb.)
110. *C. CONFERTA* (Mill! dict. n. 8. excl. syn.) foliis ellipticis utrinquè acuminatis glabris longè petiolatis, racemis terminalibus velutinis, pedicellis compressis thecaphoro brevioribus. 5 in Carthagena. fl. fructus que ignoti. (v. s. in h. Banks.)
111. *C. DIDYMOBOTRYS* (Ruiz. et Pav. fl. per. t. 423. f. b. ined. ex litt.) foliis ovalibus emarginatis subcarinatis supernè nitidis, pedunculis axillaribus geminis racemosis paucifloris distichis. 5 in Guayaquil sylvis.
112. *C. COMMUTATA* (Spreng. neu. entd. 3. p. 57.) foliis oblongo-lanceolatis utrinquè glabris, pedunculis racemosis elongatis nudis, staminibus corollà brevioribus. 5 in Guadalupà et Martinicà. Habitus *C. Breyniæ*. †.

†† *Species quoad genus dubiæ.*

113. *C? OVALIFOLIA* (Ruiz. et Pav. fl. per. t. 432. ined. ex litt.) foliis ovalibus mucronatis emarginatisve subtùs incano-tomentosis, calyce 4-crenato, glandulis calycinis squamiformibus ciliatis. 5 in Peruvà ad Ica et in Guayaquil. Petala unguiculata lutescentia margine purpurea. Stamina declinata 18, 12 exteriora basi submonadelpha. Fructus ovalis.
114. *C? RACEMOSA* (Mill. dict. n. 5.) foliis oppositis longè petiolatis ovatis, pedunculis 2-3-floris. 5 in Novà Hispanià prope Carthagenam. Flores albi. †.
115. *C? CANTONENSIS* (Lour. fl. coch. ed. Willd. 1. p. 404.) stipulis spinosis, foliis ovatis acuminatis glabris rugosis, pedunculis racemosis, floribus 5-petalis. 5 prope Cantonem Sinarum. Flores albi, fructus ovatus. †.
116. *C? BREVIS* (Spreng. ex Fisch. cat. h. Gor. 1808. p. 85.) ovario sessili. 5 Hab...

XIII. *STEPHANIA*. Willd. sp. 2. p. 239. non Lour.

Calyx campanulatus bilobus. Petala 4. Torus parvus. Stamina 6. Ovarium stipitatum oblongum. — Frutices inermes foliis longè petiolatis.

1. *S. CLEOMOIDES* (Willd. l. c.) foliis oblongo-lanceolatis valdè acuminatis vix petiolo longioribus. 5 ad Caracas in Amer. calid. *Capparis paradoxa* Jacq. Schœnb. t. 111. (v. s. sine flore)
2. *S. ELLIPTICA*, foliis ellipticis vix acuminatis petiolo duplò longioribus. 5 in insulà Trinitatis. Fructus utriusque speciei ignotus. (v. s. in herb. Banks.)

XIV. MORISONIA. *Plum. gen. t. 23.*

Calyx obovatus bifidus. Petala 4. Torus parvus. Stam. 20, basi submonadelpha? Bacca stipitata globosa. — Arbor foliis glabris petiolatis.

1. *M. AMERICANA* (Lin. sp. 719. Jacq. amer. t. 97.) 5 in insul. Caribæis. Capparis Morisonia Sw. obs. 272.

XV. THYLACHIUM. *Lour. coch. ed. Willd. 1. p. 417.*

Calyx follicularis demum circumscissus. Petala 0. Stamina ∞ toro brevi inserta. Bacca stipitata oblonga. — Frutices inermes foliis variis.

1. *T. LUCIDUM*, foliis integris ellipticis acutiusculis glabris, ramulis petiolisque velutinis, baccâ ovato-globosâ. 5 in insulâ Booby inter Nov.-Holl. et Nov.-Guineam. Capparis lucida Banks! herb. (v. s. in h. Banks.)
2. *T. AFRICANUM* (Lour! l. c. p. 418.) foliis integris ovalibus mucronulatis ramisque glabris, baccâ oblongâ. 5 in Africâ orientali. Thyl. ovalifolium Juss. ann. 12. p. 71. (v. s. in h. Mus. Par.)
3. *T. HETEROPHYLLUM* (Juss! ann. 12. p. 71.) foliis glabris, aliis simplicibus ovali-lanceolatis, aliis 3-foliolatis. 5 in Madagascar. Capparis panduri formis Pet. Th. obs. afr. austr. p. 26. (v. s. in h. Mus. Par.)
4. *T. PANDURIFORME* (Juss. ann. 12. p. 71.) foliis glabris, aliis simplicibus panduriformibus, aliis 3-foliolatis. 5 in ins. Mauritiî. Capparis panduriformis Lam. dict. 1. p. 609. (v. s. in h. Mus. Par.)
5. *T? OPPOSITIFLORUM*, foliis glabris integris ovato-lanceolatis basi subcordatis, pedicellis 1-floris oppositifoliis. 5 in insul. Caribæis? (v. s. sine fr. pet. et stam.)

XVI. HERMUPOA. *Læfl. itin. 307. †.*

Calyx duplex, exterior tubulatus, interior 4-sepalus minimus. Petala 4-linearia Stamina 6 longissima. Bacca oblongo-cylindracea. — Affinis Breyniæ ex Læfl. An calyx interior potius nectarium?

1. *H. LÆFLINGIANA*. 5 in Amer. calidior. Flos coccineus. †.

XVII. MÆRUA. *Forsk. æg. 104.*

Calyx tubulatus 4-partitus, fauce squamis petaloïdeis coronatâ. Petala 0. Torus elongatus. Stam. ∞ ad apicem tori submonadelpha. Siliqua carnosâ stipitata. — Frutices inermes, foliis simplicibus. Genus medium inter Capparideas et Passifloreas.

1. *M. UNIFLORA* (Vahl symb. 1. p. 36.) pedicellis axillaribus solitariis unifloris, coronâ petaloïdeâ filamentoso-laciniatâ, foliis aveniis. 5 in Yemen.
2. *M. ANGOLENSIS*, pedicellis axillaribus solitariis unifloris, coronâ petaloïdeâ 4-loba, lobis inciso-laceris, foliis mediis 1-nerviis. 5 ad Angolam. An satius à priore differt? (v. s. h. Mus. Par.)
3. *M. RACEMOSA* (Vahl symb. 1. p. 36.) racemis terminalibus, coronâ petaloïdeâ integrâ, 5 in Arabiâ.

Ordo XIII. FLACOURTIANÆ. *Rich. in mem. mus. 1. p. 366.*

Sepala 4-7 numero definita, basi inter se subcoalescentia. Petala sepalis alterna, tot quot sepala, rarius nulla. Stamina thalamo inserta, petalis numero æqualia, dupla aut multiplicia, interdum in squammulas nectariferas mutata. Ovarium ovato-globosum, liberum, sessile aut substipitatum. Stylus o aut filiformis. Stigmata tot quot ovarii valvulæ, plus minus distincta. Fructus 1-locularis, nunc indehiscens carnosus, nunc capsularis 4-5-valvis, intus pulpâ tenui fartus. Semina pauca, crassa, sæpè pelliculâ è pulpâ arescente factâ involuta, capsulæ valvis adfixa, nec margini (ut in Capparideis), nec lineæ longitudinali (ut in Violaceis Passifloreisve), sed, placentis nempè ramosis, in disco ipso subirregulariter et quasi areolatim adhærentia. Albumen carnosum suboleosum. Embryo rectus, axilis, radiculâ ad hilum versâ et idèd sæpius superâ, cotyledonibus planis ovalibus foliaceis.

Frutices aut arbusculæ æquatoriales. Stipulæ nullæ. Folia alterna, breviter petiolata, simplicia, penninervia, sæpius integra, coriacea. Pedunculi axillares, multiflori. Flores interdum sexu distincti. — Ordo affinis hinc Capparideis, illinc præsertim Passifloreis, ab omnibus Dicotyledoneis seminibus areolatim parietalibus seu placentâ parietibus adnexâ ramosâque distinctus. An huc revocari deberent Maunea et Vareca?

Trib. I. *PATRISIEÆ.*

Flores hermaphroditi apetali. Sepala 5 intus colorata persistentia. Stamina indefinita. Fructus capsularis baccatusve. — An potiùs genera ad Passifloreas propè Smeathmanniam referenda? sed semina lævia, nec ut in Passifloreis scrobiculata, disco toto nec lineæ valvularum longitudinali adfixa.

I. *RYANAEA.* — *Ryania*. *Vahl ecl. 1. p. 51. t. 9.* — *Patrisia* *Rich. act. soc. par. p. III.*

Urceolus petaloideus inter stamina et pistillum. Fructus baccatus indehiscens.

1. *R. SPECIOSA* (Vahl l. c.) foliis utrinquè viridibus subtus in costis pilos stellatim fasciculatos gerentibus, pedunculis unifloris. ♂ in Insulâ Trinitatis et Cayennæ. *Patrisia pyrifera* *Rich. l. c. Pers. ench. 2. p. 69.* Nomen tantillum mutavi ne confundatur cum Rianâ.

II. *PATRISIA.* *H. B. et K. nov. gen. 5. p. 356.*

Urceolus inter stamina et ovarium nullus. Fructus capsularis dehis-
cens suberosus 3-5-valvis. — In honorem cl. Patris qui duas species
piores detexit.

1. *P. BICOLOR*, foliis subtus tenuissimo tomento albidis, pedunculis multifloris. 5 in Cayennâ. (v. s.)
2. *P. PARVIFLORA*, foliis subtus incostis villosis, pedunculis unifloris. 5 in Cayennâ. (v. s.)
3. *P. DENTATA* (H. B. et K. nov. sp. 5. p. 357.) foliis glabris subtus in nervo et venis puberulis, floribus axillaribus 1-2 breviter pedicellatis, ramulis subflexuosis. 5 in umbris arenosis inter Aturas et Maypures mission. orinocensium.
4. *P. AFFINIS* (H. B. et K. l. c.) foliis glabris nervo venisque subtus pilosis, aculis ovatis acuminatis obsolete denticulatis, floribus 1-2 axillaribus breviter pedunculatis, ramulis subrectis. 5 in scopulis propè Carichana ad ripam Orinoci.

Trib. II. *FLACOURTIEÆ*.

Flores abortu dioici, apetalî. Stamina indefinita. Fructus baccatus indehiscens.

III. *FLACOURTIA*. *L'her. stirp.* 59. t. 30.

♂ Stamina suprâ torum hemisphæricum dense congesta, basi eglandulosa. ♀ Calyx deciduus. Stigmata 4-9 longitudinaliter superne 1-sulca. Semina ossea. — Frutices sæpius spinosi.

1. *F. RAMONTCHII* (L'her! l. c. Lam. ill. t. 826.) foliis subrotundo-ovatis acutis crenatis. 3 in Madagascar. Alamotou Flac. mad. 124. (v. s.)
2. *F. SAPIDA* (Roxb. cor. 1. p. 49. t. 69.) foliis ellipticis repando-serratis basi obtusis apice obtusiusculis. 5 in montibus Indiæ orientalis.
3. *F. INERMIS* (Roxb. cor. 222.) arborea inermis, foliis ellipticis crenato-serratis nitidis, racemis axillaribus brevibus, floribus hermaphroditis, stylo 5-fido. 5 in Mollucis. — Mal. misc. 1. n. 1. p. 25. — Stamina 20-30. Bacca rubro-purpurea nimis acida.
4. *F. SEPIARIA* (Roxb. cor. 1. p. 48. t. 68.) foliis obovato-oblongis repando-serratis. 3 in Indiâ orientali. — Rheed. mal. 2. t. 39. Synonyma plurima forsân cum *Elæodendron Argan* confusa. (v. s. in h. Juss.)
5. *F. CATAPHRACTA* (Roxb! in Willd. sp. 4. p. 830.) foliis ovali-oblongis acuminatis serratis. 5 in Indiâ orientali. (v. s. ♂.)
6. *F. FLAVESCENS* (Willd. sp. 4. p. 830.) foliis oblongis obtusiusculis serratis basi attenuatis. 5 in Guineâ.
7. *F. RHAMNOIDES* (Burch! cat. afr. austr. n. 4012.) foliis ellipticis subserratis acutiusculis. 3 in sylvaticis ad cap. Bonæ-Spei. Flores axillares, parvi. Pedicelli 1-flori. Baccæ ovatæ, rubræ. (v. s.)

IV. *ROUMEA*. *Poit. mem. mus.* 1. p. 62. t. 4.

♂ Stamina toro non dilatato inserta, glandulâ crenatâ basi cincta. ♀ Calyx persistens. Stigmata apice depresso-capitata. Semina cartilaginea. — Frutex spinosus.

1. *R. CORIACEA* (Poit! l. c.) spinis ramosum adultorum nudè magnâ ramosis, nuuc parvis simplicibus, foliis serratis. 3 in Sancti-Domingi siccis et apricis. *Kœlera laurifolia* W. (excl. descr. cum *Drypete* conf.) *Bessera spinosa* Spreng. pug. 2. p. 91. *Limacia laurifolia* Dietr. (v. s.)
- 2? *R. INERMIS*, spinis nullis, foliis integerrimis. 5 in Bengalo. *Bessera inermis* Spreng. pug. 2. p. 90. Rami nervis 3 à quoque folio decurrentibus angulati. Flores ad axillas 15-20 adgregati, tenuiter pedicellati minimi. ♂ antheræ pauce sessiles. Stirps non satis nota à genere, forsân ab ordine removenda. (v. s. in h. Balb.)

V. STIGMAROTA. *Lour. fl. coch. 2. p. 778.*

♂ ut in Roumea? ♀ Calyx deciduus. Stylus cylindricus brevis. Stigmata 6 radiantia. — An genus cum Flacourtiâ aut Roumeâ conjungendum?

1. S. JANGOMAS (*Lour. l. c.*) spinis fœminæ simplicibus, masculæ ramosis, foliis ovatis acuminatis serratis. 5 in Cochinchinâ ex *Lour. in Javâ et Baleyâ* ex *Rumph. app. p. 36. t. 19. f. 1. 2.*

Trib. III. KIGGELARIÆ.

Flores abortu dioici. Petala? 5 sepalis alterna. Stamina definita. Fructus subbaccatus demùm dehiscens.

VI. KIGGELARIA. *Lin. gen. n. 1128.*

Petala 5 ad unguem glandulas 3 gerentia. ♂ stam. 10-20, filam. brevibus, antheris apice birimulosis. ♀ styli 2-5. — Frutices foliis saliciformibus subtus subvelutinis.

1. K. AFRICANA (*Lin. sp. 1466.*) foliis serratis supernè demùm glabratis, staminibus 10, stylis 5. 3 in Africâ australi. *Lam. ill. t. 82j. (v. v.)*
2. K. INTEGRIFOLIA (*Jacq. coll. 2. p. 296. ic. rar. t. 628.*) foliis integris utrinquè subvelutinis, staminibus 20, stylis 2. 5 ad cap. *Bonzæ-Spei.*

VII. MELICYTUS. *Forst. gen. t. 62.*

Petala 5 eglandulosa. ♂ stam. 5, filamentis clavatis, antheris adnatis. ♀ stylus brevissimus. Stigmata 4-5 radiantia.

1. M. RAMIFLORUS (*Forst. l. c. Lam. ill. t. 812. f. 1.*) foliis dentatis oblongo-cuneatis utrinquè glabris. 5 in Novâ-Zeelandiâ.
2? M? UMBELLATUS (*Gaertn. fr. 1. p. 206. t. 44. f. 3.*) An eadem ac prior? An Erythrospermi cujusdam fructus?

VIII. HYDNOCARPUS. *Gaertn. fr. 1. p. 238. t. 60. f. 3.*

♂ Sepala 5, duo ext. ovata. Petala 5 margine villosa, intus squamâ sucta Stam. 5. ♀ Flos. Bacca sphaerica tuberculis 4 reflexis terminata, placentis 4 polyspermis.

1. H. INEBRIANS (*Vahl! symb. 3. p. 100.*) 5 in Ceylonâ. H. venenata *Gaertn. l. c.*

Trib. IV. ERYTHROSPERMEÆ.

Flores hermaphroditi. Petala et Stamina 5-7. Fructus indehiscens subbaccatus.

IX. ERYTHROSPERMUM. *Lam. ill. t. 274.*

Cal. 4-sepalus deciduus. Pet. 4-7 vix calyce longiora. Stam. 4-7 filam. brevissimis. Ovarium subrotundum. Stigmata 5-5. Fructus ∞-spermus. — Frutices glaberrimi inermes.

1. E. MACROPHYLLUM (*Poir. suppl. 2. p. 585.*) foliis sparsis lanceolatis obtusis mucronatis, racemis folio brevioribus. 5 in ins. Mauritiî.
2. E. PANICULATUM (*Poir. l. c. p. 585.*) foliis sparsis ovatis utrinquè obtusis, racemis paniculatis folio longioribus. 5 in ins. Mauritiî.

ℓ? pauciflorum, racemis 2-4-floris.

3. *E. ELLIPTICUM* (Poir. l. c. p. 585.) foliis sparsis elliptico-subrotundis utrinquē obtusis, racemis paucifloris folio longioribus. 5 in ins. Mauritii.
β. *mucronatum*, foliis ovatis mucronatis. — in ins. Java.
4. *E. AMPLEXICAULE*, foliis sparsis apice ramorum confertis ovalibus basi cordatis, pedunculis folio brevioribus subumbellatis. 5 in ins. Mauritii. (v. s. in h. Mus. Par.)
5. *E. PYRIFOLIUM* (Lam. ill. t. 274. f. 1.) foliis sparsis ovatis brevissimè petiolatis utrinquē obtusis, racemis subterminalibus folio brevioribus. 5 ins. Mauritii. (v. s.)
6. *E. VERTICILLATUM* (Lam. ill. t. 274. f. 2.) foliis ternatim verticillatis subsessilibus subrotundis, pedunculis apice corymboso-umbellatis folii longitudine. 5 in ins. Mauritii et Borboniæ. hujus speciei fructus omninò Kiggelariam refert. (v. s.)
-

Ordo XIV. BIXINÆ. *Kunth malv. p. 17. Nov. gen. am. 5. p. 331.*

Calyx 4-7-sepalus, sepalis æstivatione imbricatis, nunc omninò distinctis, nunc basi subcoalitis. Petala 5, sepalis sat similia aut nulla. Stamina numero indefinita, thalamo aut disco fundo calycis imposito inserta, filamentis liberis, antheris bilocularibus. Ovarium superum, sessile, 1-loculare. Ovula placentis 1-7 parietalibus inserta. Stylus 1 indivisus aut 2-4-fidus. Fructus capsularis baccatusve, 1-locularis, polyspermus. Semina placentis parietalibus adfixa, membranâ carnosâ forsâ involuta. Albumen carnosum aut tenuissimum. Embryo inclusus, subrectus aut curvatus, cotyledonibus foliaceis, radiculâ hilum spectante (Kunth.) — Arbores aut frutices sæpiùs glabri. Folia alterna simplicia integra interdum pellucido-punctata. Stipulæ caduæ. Pedunculi axillares, 1-∞-flori, bracteolati. — Ordo nuperrimè stabilitus, generibus plerisque non satis notis constans, ex habitu Malvaceis, Flacourtianeis, Rosaceisve affinis, ex characteribus carpologicis Flacourtianeis Cistineisve accedens.

I. BIXA. *Lin. gen. n. 654. H. B. et K. nov. gen. am, 5. p. 353.*

Cal. 5-sepalus, sepalis orbiculatis basi tuberculatis deciduis per æstivationem imbricatis. Pet. 5 obovata. Stam. ∞ hypogyna libera, filam. filiformibus, antheris ovatis. Ovarium 1. Stylus 1 longus apice fere ligulato-compressus. Capsula 1-locul. bivalvis, valvis extus setoso-hispidis medio intus placentam linearem gerentibus. Sem. 8-10 ad quamque placentam pulpâ farinaceâ coloratâ induta. Albumen carnosum. Embryo erectus cotyled. foliaceis planis parallelis medio transverse plicatis (Gaertn. fr. 1. t. 61.)

1. B. ORELLANA (Lind. sp. 730.) foliis utrinque glabris. ♂ in Amer. calid. ad fossas et fontes, culta in ins. Caribæis et Indicis. Sloan. hist. 2. p. 52. t. 181. f. 1. Comm. horst. amst. 1. t. 33 B. Americana Poir. dict. 6. p. 229. Pulpa semina involvens tinctoria sub nomine gallico *Rocou.* (v. s.)
2. B. URUCURANA (Willd. enum. 565.) foliis subtus lepidotis albis. ♂ in Brasiliâ.

II. BANARA. *Aubl. guian. 1. p. 547.*

Cal. persistens 6-partitus, lobis subrotundis. Pet. 6 subrotunda, infra discum inserta. Stam. 15 et ampliùs ad ambitum disci inserta. Ovarium disco insidens, orbiculatum. Styl. 1. Stigma capitatum. Bacca vix succulenta, stylo terminata. Semina ∞ striata.

1. B. GUIANENSIS (Aubl. l. c. t. 217.) ♂ in Cayennæ sylvis. Folia ovato-oblonga acuta denticulata subtus subtomentosa. Flores flavi.

III. LAETIA. *Lin. gen. n. 651.*

Cal. 5-partitus coloratus marcescens. Pet. 5 aut o. Stam. numero indefinita hypogyna, antheris subrotundis. Stylus 1. Stigma capitatum. Capsula carnosa 3-5-valvis parva globosa stylo acuminata. Semina plurima pelliculâ pulposâ arillata.

* *Floribus apetalis.*

1. L. APETALA (Jacq. am. 167. t. 108.) floribus apetalis, pedunculis axillaribus subtrifloris, foliis oblongis obtusis serrulatis nitidis. ♀ in Carthagenæ sylvaticis, et ad flum. Magdalenæ. H. B. et K. nov. gen. am. 5. p. 354.
2. L. GUIDONIA (Sw. prod. 83.) floribus apetalis, pedunculis 1-floris terminalibus, foliis oblongis acuminatis pubescentibus. ♀ in Jamaicâ. Guidonia Brown. jam. 249. t. 29. f. 4? Samyda icosandra Sw. fl. ind. 3. p. 1962. Stylus brevissimus. Capsula carnosa 3-valvis. An revera Samyda species?
3. L. THAMNIA (Sw! fl. ind. occ. 2. p. 950.) floribus apetalis, pedunculis axillaribus multifloris subdivisis, foliis oblongis acutis subcrenatis nitidis. ♀ in maritimis fruticetis Jamaicæ australis. Thamnina Brown jam. 245. t. 25. f. 2. Helwingia Adans. (v. s.)

** *Floribus completis.*

4. L. COMPLETA (Jacq. am. 167. t. 183. f. 60.) floribus pentapetalis, foliis ovatis obtusis serratis rugosis glabris. ♀ in Carthagenæ sylvis.

† *Species quoad flores non satis notæ.*

5. L. HIRTELLA (H. B. et K. nov. gen. am. 5. p. 355.) ramulis foliisque subtus in nervo venisque hirtellis oblongis obtusis argutè denticulatis, pedunculis 1-carpis. ♀ prope Cumana.
6. L. GUAZUMÆFOLIA (H. B. et K. l. c.) ramulis ferrugineo-tomentosis, foliis oblongis subobtusis obsolete denticulatis subtus in nervo et venis hirtellis, pedunculis dichotomis ∞-floris? ♀ in Novâ-Andalusîâ prope Bordonas.

IV. PROCKIA. *Browne in Lin. gen. n. 674. Lam. dict. 5. p. 625. ill. t. 465.*

Cal. persistens 3-5-partitus, lobis subrotundis inæqualibus. Pet. nulla. Stam. ∞ disco inserta, antheris subrotundis. Ovarium 1 subrotundum. Stigma integrum. Bacca subexsucca 4-6-sperma subrotunda. — Frutices americani aut Mauritanii, glabri. Folia alterna, integra, aut dentata. Flores interdum abortu 1-sexuales.

Sect. I. PROCKIARIA.

Stylus filiformis. Stigma obtusum seu acutiusculum.

1. P. CRUCIS (Lin. sp. 745.) foliis ovatis acuminatis serratis, stipulis subfalcatibus, racemis paucifloris terminalibus. ♀ in ins. Sanctæ-Crucis. Calyx 3-4-sepalus Vahl symb. 3. p. 69. t. 64. — P. acuta fl. mex. ic. ined? sed folia angustiora.
 - β. cordata, foliis cordatis, calyce 4-5-sepalo, stipulis linearibus. ♀ in ins. Sanctæ-Crucis. Lam. ill. t. 465. f. 1. — P. subcordata fl. mex. ic. ined. huc pertinere videtur.
2. P. DELTOIDES (Lam. ill. t. 465. f. 3.) foliis subrotundo-deltoidibus, crenato-subsinuatis, pedicellis axillaribus subgeminis. ♀ in ins. Mauritiî et Madagascar. (v. s.)

Sect. II. *APHLOIA*. — *Lightfootia*, Sw. prod. 83. non *L'her*.

Stylus nullus seu vix conspicuus. Stigma latum suborbiculatum planiusculum.

3. *P. serrata* (Willd. sp. 2. p. 1213.) foliis oblongo-ovatis serratis acuminatis, pedunculis lateralibus aggregatis unifloris. ♂ in ins. Montserrat. *Lightfootia serrata* Sw. fl. ind. occ. 2. p. 947.*

4. *P. integrifolia* (Willd. sp. 2. p. 1214.) foliis subcoriaceis ovalibus obovatisve integris aut subdentatis obtusis subemarginatisve, pedunculis 1-4-axillaribus 1-floris. ♂ in ins. Mauritii. *Lightfootia integrifolia* Vahl symb. 3. p. 70. (v. s.)

♀ *ovata* (Poir. dict. 5. p. 626.) pedicellis in umbellulam axillarem ferè dispositis. ♂ Lam. ill. t. 465. f. 2. (v. s.)

5. *P. thæformis* (Willd. sp. 2. p. 1214.) foliis submembranaceis lanceolato-ellipticis obtusè serratis, basi ad marginem subrevolutis et in petiolum decurrentibus, pedicellis 1-3 axillaribus 1-floris. ♂ in ins. Borboniæ. *Lightfootia thæformis* Vahl symb. 3. p. 69. *Ludia heterophylla* Bory voy. 2. p. 115. t. 24. fig. inf. (v. s.)

♀? *laciniata* (Poir. dict. 5. p. 627.) foliis pinnatifidis, lobis mucronatis integris dentatisve. ♂ *Ludia heterophylla* Bory voy. 2. t. 24. fig. sup. Potius rami steriles ejusdem arboris quàm varietas. (v. s.)

† *Species non satis notæ.*

6. P? *lobata* (Poir. dict. 5. p. 627.) foliis ovato-acuminatis serratis, pedunculis axillaribus racemosis 3-5-floris, stigmate subsessili 5-lobato. ♂ Patr. ign. *Litsea* Lam. ill. t. 834.

7. P? *macrostachya* (fl. mex. ic. ined.) foliis ovato-lanceolatis acuminatis integerrimis, racemis terminalibus elongatis nudis, calycibus 4-partitis obtusis, fructibus 1-spermis. ♂ in Mexico. A genere, forsan ab ordine removenda! Stam. 12 hypogyna. Stylus brevissimus. Stigma orbiculatum.

V. *LUDIA*. Lam. dict. 3. p. 612. ill. t. 466.

Cal. persistens 5-7-partitus, lobis ovalibus. Pet. nulla. Stam. ∞ disco inserta, antheris subrotundis. Ovarium 1 ovatum. Stylus filiformis. Stigma trifidum, rariùs 4-fidum. Bacca globosa stylo acuminata 6-8-sperma. — Frutices Mauritanii, floribus lateralibus subsessilibus, foliis sæpè in iisdem individuis formâ variis (1), undè nomen.

1. *L. heterophylla* (Lam. l. c. t. 466. f. 1. 2.) foliis obovatis nitidis venosis, junioris fruticis minimis dentato-spinosis, adulti majoribus integerrimis, pedicellis axillaribus solitariis. ♂ in ins. Mauritii. *L. Mauritiana* Rœusch. ex Steud. nom. (v. s.)

2. *L. myrtifolia* (Lam. l. c. t. 466. f. 3.) foliis ovatis integerrimis venosis, pedicellis axillaribus solitariis, stylo incurvo. ♂ in ins. Borboniæ.

3. *L. sessiliflora* (Lam. l. c.) foliis ovato-oblongis acutiusculis, floribus axillaribus subsessilibus, stylo recto. ♂ in ins. Mauritii. *L. tuberculata* Jacq. h. schœnb. 1. t. 112. Ovarium 1-loculare, ovulis 6 per paria placentis 3 parietalibus adnatis. (v. s.)

-(1) Hoc phenomenon frequens est in stirpibus Mauritanis, ex. gr. *Quivisiâ*, *Acaciâ heterophyllâ*, etc. etc.

VI. AZARA. *Ruiz et Pav. fl. per. prod. 79. t. 36. syst. p. 137.*

Cal. 4-6-partitus. Pet. o. Stam. ∞ , filamentis filiformibus liberis et fila plurima capillaria è toro orta. Antheræ 2-loc. Stylus subulatus. Stigma obtusum. Bacca stylo apiculata, 1-loc. ∞ -sperma. Placentæ 3 parietales. — Frutices Chilenses; Folia geminata inæqualia. Flores fragrantés.

1. *A. SERRATA* (Ruiz et Pav. l. c.) foliis geminis serratis, majori lanceolato, minori subrotundo, corymbis multifloris. ♂ in nemoribus Chilensibus ubi Corcolen dicitur. (v. s.)
 2. *A. INTEGRIFOLIA* (Ruiz et Pav. syst. p. 138.) foliis geminis integerrimis, majori obovato, minori subrotundo, floribus spicatis dependentibus. ♂ in nemoribus Conceptionis Chile. (v. s.)
 3. *A. DENTATA* (Ruiz et Pav. l. c.) foliis geminis dentatis, majori elliptico, minori subrotundo, floribus umbellatis. ♂ in nemoribus Conceptionis Chile.
-

Ordo XV. CISTINEÆ. — Cisti *Juss. gen.* 294. et Cistoidæ *Vent. tabl.* 3. p. 219 *excl. gen. affin.* — Cisti *DC. fl. fr.* 4. p. 811. (1)

Calycis sepalâ 5 cum pedicello continua persistentia sæpius inæqualia, 2 exteriora ceteris sæpius minora interdum evanida, 3 interiora per æstivationem contorta. Petalâ 5 hypogyna, caduca, æqualia, per æstivationem contorta, sed directione sepalis contraria. Stamina numero indefinita, sæpius ∞ , hypogyna, erecta. Filamenta libera. Antheræ ovatæ, biloculares, birimosæ, basi insertæ. Ovarium liberum. Stylus 1 filiformis. Stigma simplex. Capsula constans valvis 3-5, rarius 10, nunc medio nervum longitudinalem placentarium gerentibus et tunc capsula 1-locularis, nunc nervo medio in septum plus minusve completum abeunte et tunc capsula completè aut incompletè multilocularis. Semina idè semper verè parietalia, sed nunc placentæ parieti contiguæ nunc septi angulo interno adfixa plurima, parvula. Albumen farinosum. Embryo spiralis aut curvus intrâ albumen. — Suffrutices aut Herbæ. Folia simplicia, penninervia, integra aut subdentata, primordialia semper opposita, cetera sæpius opposita, interdum alterna, basi nunc nuda, nunc stipulis binis foliaceis instructa. Racemi sæpè floribus unilateralibus pedicellatis successivè evolutis et idè ferè scorpioidei. Flores rosacei, petalis fugacissimis, sæpius ephemeri, semper diurni, flavi albi aut purpurei, ungue sæpè discolore. Rami succo glutinoso sæpè obducti. — Ordo affinis hinc Violarieis quibuscum olim confusus et à quibus differt præcipuè staminibus indefinitis, illinc Bixineis à quibus non differt nisi petalis semper præsentibus nec interdum nullis, sepalorum æstivatione subdiversâ, albumine farinaceo, foliis nunquam pellucidis, etc.

I. CISTUS. *Tourn. inst.* 259. t. 136. *Gaertn. fr.* t. 76. *DC. fl. fr.* 4. p. 811. Cisti sp. *Linn.* *

Calyx 5-sepalus, sepalis duplici serie dispositis, 2 externis inæqualibus, interdum nullis. Petalâ 5, æqualia, subcuneata, caduca. Stamina ∞ , sæpè è disco glanduloso exserta. Stylus filiformis. Stigma capitatum. Capsula calyce oblecta, 10-5-locularis, valvis 10-5, medio septiferis. Semina ovato-angulata. Embryo filiformis spiralis. — Frutices suffruticesve. Folia opposita exstipulata integra vel subdenticu-

(1) Hujus ordinis hanc synopsis nobis communicavit cl. et amiciss. F. DUNAL, ex monographiâ inchoatâ excerptam; ad hunc laborem adminicula momentosa præbuerunt et Herbarium Thibaudianum ex quo cl. Persoon species plurimas instituerat, et collectio Cistinearum Hispanicarum, notis manuscriptis sagacissimis illustrata, à cl. Lagasca humanissimè nobiscum communicata.

lata. Pedunculi axillares, uni aut multiflori. Semina ex solo C. monspeliensi descripta.

Sect. I. ERYTHROCISTUS.

Sepala externa angustiora, sæpè minora, interna basi concava margine serrata. Petala rosea, rubra aut purpurea. Capsulæ 5-loculares.

* *Pedunculis unifloris, axillaribus vel terminalibus, solitariis umbellatisve; stylo cylindrico sæpè staminibus longiore; stigmatibus capitato 5-sulcato.*

1. C. PURPUREUS (Lam. dict. 2. p. 14.) foliis oblongo-lanceolatis utrinquè acuminatis rugosis, petiolis brevibus pilosis vaginantibus, pedunculis brevibus 1-2-ternisve. ζ in Oriente. Petala rubro purpurea, basi maculâ atro-purpureâ notata Alabastrum magnum. Ker. bot. reg. t. 408. (v. s.)
2. C. HETEROPHYLLUS (Desf. fl. atl. 1. p. 411. t. 104.) foliis ovato-lanceolatis breviter petiolatis basi vaginantibus margine revolutis, pedunculis hirsutis foliosis unifloris 1-2-ternisve. ζ in collibus incultis Algeriæ. Corolla magna, rosea. (v. s.)
3. C. PARVIFLORUS (Lam. dict. 2. p. 14.) foliis ovatis acutis subtomentosis in petiolum attenuatis subconnatis, pedunculis 1-ternisve subterminalibus. ζ in Cretâ. Corolla parva, rosea. (v. s.)
4. C. COMPLICATUS (Lam. dict. 2. p. 14.) foliis subrotundo-ovatis obtusiusculis albo-tomentosis approximatis subtus reticulatis petiolatis, petiolis basi dilatatis margine pilosis subcavis vaginantibus, pedunculis brevibus unifloris 3-4-nisve subterminalibus. ζ in Oriente et in Hispaniæ montibus Valentiniis. Flores parvi rosei. (v. s.)
5. C. VILLOSUS (Lam. dict. 2. p. 12.) foliis subrotundo-ovatis rugosis tomentoso hirtis petiolatis, petiolis sulcatis connatis, pedunculis unifloris, 1-3-nisve, sepalis villosis. ζ in Italiâ, Hispaniâ et Barbariâ. C. pilosus Lin. spec. 737. Duh. arb. 1. p. 67. t. 64. Petala rubro-purpurea magna. (v. s.)
 β . *virescens*, ramis foliisque viridioribus, pedunculis unibracteatis. ζ hab. cum var α . Plurimæ varietates aut species forsan adhuc sub hoc titulo latent. Syn. Lin. ex errore typographico provenit.
6. C. CRETICUS (Linn. spec. 1. p. 738.) foliis spathulato-ovatis tomentoso-hirtis in petiolum brevem attenuatis margine undulatis, pedunculis brevibus unifloris, sepalis villosis. ζ in Cretâ, Syriâ. Jacq. ic. rar. 1. t. 95.
 β . *crispatus*, foliis undulatis et crispis. (v. s.)
 γ . *Tauricus*, foliis planis subtus sepalisque villosissimis. ζ in Tauriâ. C. Creticus Bieb. (v. s.)
7. C. INCANUS (Linn. spec. 737.) foliis spathulatis tomentosis rugosis subtrinerviis sessilibus subconnatis, superioribus angustioribus, pedunculis unifloris. ζ in Hispaniâ, Galliâ Narbonensi. Petala rubro-purpurea emarginata. (v. v.)
 β . foliis linearibus longis. — C. mas 2. Clus. hist. 1. p. 69. ic.
8. C. UNDULATUS (Dunal ined.) foliis sessilibus lineari-oblongis acutis margine undulatis basi trinerviis, pedunculis solitariis unibracteatis, sepalis longè acuminatis villosis. ζ Hab. Fortè var. culta C. crispis. (v. s.)
9. C. CRISPUS (Linn. spec. 738.) foliis sessilibus lineari-lanceolatis undulato-crispis trinerviis rugosis pubescentibus, floribus subsessilibus 3-4-nis umbellatis. ζ in Galliâ australi, Hispaniâ, Lusitaniâ. Cav! ic. 2. p. 57. t. 174.* Petala purpurea. (v. v.)
10. C. ALBIDUS (Linn. spec. 737.) foliis sessilibus oblongo-ellipticis incanotomentosis subtrinerviis, floribus 3-8 terminalibus subumbellatis, sepalis externis majoribus. ζ in Galliâ Narbonensi, Hispaniâ, Lusitaniâ. C. mas 1. Clus. hist. 1. p. 68. ic. Petala diluè purpurea, maculâ parvâ luteâ basi notata. (v. v.)
11. C. CANDIDISSIMUS (Dun. ined.) foliis ovato-ellipticis acutis densè tomentoso-incanis trinerviis, breviter petiolatis, petiolis basi dilatatis margine

pilosis vaginantibus, pedunculis solitariis unifloris folio brevioribus, sepalis externis dimidio brevioribus. ζ in insulæ magnæ Canariæ pinetis elevatis. Chr. Smith. Flos pallide roseus. (v. s.)

12. *C. VAGINATUS* (Ait. hort. Kew. 3. p. 304.) foliis lanceolatis acutis trinerviis hirsutis subtus reticulatis petiolatis, petiolis basi dilatatis margine pilosis sulcatis vaginantibus, pedunculis 1-3 axillaribus vel terminalibus longis basi bracteatis. ζ in ins. Teneriffâ. Jacq. hort. Schœnb. 3. p. 17. t. 282. *C. symphytifolius* Lam. dict. 2. p. 15. Corolla basi lutescens, dilute purpurea. (v. v.)
- 13? *C. SERICEUS* (Vahl symb. 1. p. 37.) foliis ovatis tomentosis trinerviis, inferioribus petiolatis, supremis sessilibus, pedunculis hirtis. ζ in Hispaniâ. *C. latifolius* magno flore Barrel. ic. 1315. Pedunculorum pili longi, purpurascens. Petala filamentaque purpurea.
- 14? *C. HYBRIDUS* (Vahl symb. 1. p. 37.) foliis ovatis incanis petiolatis, ramis squamosis, pedunculis elongatis hirtis. ζ in Hispaniâ. Petala purpurea.

* *Pedunculis cymosis, stylo subnullo capitato staminibus brevior.*

15. *C. CYMOSUS* (Dunal ined.) foliis lato-ovatis apice contortis acutiusculis subtus rugosis incanis petiolatis, petiolis basi dilatatis sulcatis subvaginantibus, pedunculis cymosis 5-10-floris incanis axillaribus terminalibusve. ζ Hab. olim cultus in hort. Cels. sub nomine *C. incani*. (v. s.)

SECT. II. LEDONIA.

Sepala 5, 2 externa majora valdè acuminata vel nulla; petala alba aut albida; stamina numerosa pistillo longiora; stigma sessile magnum capitatum; capsulæ 5-10-loculares. — Frutices aut suffrutices; folia sæpè glutinosa.

\S . 1. *Pedunculis unifloris, aut multifloris cymosis, sepalis 5, externis sæpius cordatis acuminatis; capsulis 5-locularibus.*

* *Pedunculis basi nudis, sæpè infrà medium folia opposita gerentibus.*

16. *C. SALVIFOLIUS* (Linn. spec. 732.) foliis petiolatis ovatis obtusis rugosis subtus tomentosis, pedunculis longis tomentoso-albicantibus unifloris superne articulatis solitariis. ζ In Helvetiâ, Galliâ australi Italiâ, Hispaniâ, Lusitaniâ. *C. femina* Clus. hist. 1. p. 70. ic. *
- a. *erectiusculus*, caule suberecto.
- ß. *humifusus*, caule humifuso.
- γ. *ochroleucus*, floribus ochroleucis.
17. *C. CORBARIENSIS* (Poirr! in herb. DC.) foliis petiolatis subcordatis ovatis acuminatis margine limbriatis utrinquè rugosis leviter glutinosis, pedunculis longis 1-5-floris. ζ in Galliæ australis montibus dictis *les Corbières*, et in Hispaniâ. *C. hybridus* Poirr. chlor. narb. p. 36. nec Vahl. *C. salvifolius* ß. DC! fl. fr. 4. p. 813. (v. s.)
18. *C. FLORENTINUS* (Lam. dict. 2. p. 17*.) foliis angusto-lanceolatis rugosis subtus reticulatis subsessilibus, pedunculis villosis subtrifloris. ζ in Italiâ.
19. *C. MONSPELIENSIS* (Linn. spec. 737.) foliis linearis-lanceolatis sessilibus trinerviis utrinquè villosis, pedunculis cymosis subsecundis. ζ in Galliâ australi, Hispaniâ, Lusitaniâ. Lam. ill. t. 477. f. 4. Cav! ic. 2. t. 137. Jacq. coll. 2. t. 8. (v. v.)
20. *C. LEDON* (Lam. dict. 2. p. 17*.) foliis connatis oblongo-lanceolatis nervosis suprâ glabris nitidis subtus villosis cymosis subsecundis, floribus corymboso-cymosis, pedunculis calycibusque villosis-sericeis. ζ in Galliâ australi. Duh. arb. 1. p. 168. t. 66. (v. v.)
21. *C. HIRSUTUS* (Lam. dict. 2. p. 17*.) foliis sessilibus oblongis obtusis hirsutis, pedunculis brevibus unifloris aut cymoso-multifloris, capsulis parvis calyce maximo hirsuto et pyramidalis tectis. ζ in Armoraciâ, propè Landerneau et montosis Hispaniæ. Ledon. iv. Clus. hist. 1. p. 78* (v. v. et s.)
22. *C. LAXUS* (Ait. hort. Kew. 3. p. 305. exclus. syn.) foliis breviter petiolatis

ovato-lanceolatis acuminatis margine undulatis subdenticulatis subglabris, summis hirtis, floribus cymosis, pedunculis calycibusque hirsutis. ζ in Hispaniâ, Lusitaniâ. C. capensis Linn. sp. 736? ex herb. Burmann. (v. s.)

** *Pedunculis bracteatis, basi bracteolis caducis parvulis concavis coriaceis sublateis decussatis, infra mediûm 2 oppositis majoribus.*

23. C. LONGIFOLIUS (Lam. dict. 2. p. 16.*) foliis breviter petiolatis oblongo-lanceolatis margine pubescentibus et undulatis subtus venosis, pedunculis cymosis. ζ in Hispaniâ et Galliâ Narbonensi. C. nigricans Pourr. act. toul. 3. p. 311. (v. s.)

24. C. POPULIFOLIUS (Linn. spec. 736.) foliis petiolatis cordatis acuminatis lævibus, floribus cymosis, pedunculis bracteatis, bracteis oblongis. ζ

α . *major* (Pourr! in h. DC.) pedunculis calycibusque hirsutis, pilis longis albescentibus. — in Mauritiâ. (v. s.)

β . *minor*, pedunculis calycibusque subglabris nitidis viscosis. — in Galliâ australi, Hispaniâ, Lusitaniâ. Ledon latifolium 2. Clus. hist. 1. p. 78. C. populifolius Cav. ic. 3. t. 215. (v. s.)

\S . 2. *Pedunculis bracteatis, bracteis caducis decussatis, inferioribus minoribus unifloris, axillaribus solitariis vel terminalibus umbellatis; calycibus 3-sepalis; capsulis 5-10 locularibus.*

* *Stigmate magno sessili.*

25. C. LAURIFOLIUS (Linn. spec. 736.) foliis petiolatis ovato-lanceolatis trinerviis supra glabris subtus tomentosis, petiolis basi dilatatis connatis, capsulis 5-locularibus. ζ in Galliâ australi, Hispaniâ. Clus. hist. 1. p. 78. f. 1. (v. v.)

26. C. CYPRIUS (Lam. dict. 2. p. 16.) foliis petiolatis oblongo-lanceolatis supra glabris subtus tomentoso-incanis, pedunculis subtrifloris, petalis guttatis, capsulis 5-locularibus. ζ in ins. Cypri. An C. stenophyllus Link enum. 2. p. 74? (v. s. sine flore et fructu.)

27. C. LABANIFERUS (Linn. spec. 737.) foliis subsessilibus basi connatis lineari-lanceolatis supra glabris subtus tomentosis, capsulis 10-locularibus. ζ in Hispaniâ, Lusitaniâ collibus.

α . *albiflorus*, petalis omnino albis. Ledon 1 Clus. hist. 1. p. 78. ic. (v. s.)

β . *maculatus*, petalis albis, basi maculâ atro-sanguineâ notatis. (v. s.)

** *Stigmate capitato parvo, stylo cylindrico staminibus subæquali.*

28. C. CLUSII (Dunal! ined.) caule fruticoso erecto, foliis subtrinerviis linearibus margine revolutis subtus canescentibus, floribus subcapitatis, calyce 3-sepalo piloso, sepalis ovatis acutis, capsulis 5-locularibus. ζ in Hispaniâ, Barbariâ. C. Libanotis β . Lam. dict. 2. p. 18. Desf. fl. atl. 1. p. 412. excl. syn. Ledon VII. Clus. hist. 1. p. 80. ic. Bracteæ pilosæ lato-ovatæ acuminatæ ciliatæ involventes caducæ pedunculis sublongiores. Petala alba. Habitus Helianthemî Libanotis, characteres Ledoniæ. (v. s. comm. à cl. Lag.)

II. HELIANTHEMUM. Tourn. inst. 248. t. 128. Gaertn. fr. 1. p. 371. t. 76. DC. fl. fr. 4. p. 815. — Cisti sp. Linn.

Calyx 3-sepalus, sepalis æqualibus, vel 5-sepalus, sepalis duplici serie dispositis, 2 externis sæpè minoribus, rarè majoribus. Petala 5 sæpè apice irregulariter denticulata. Stigma capitatum. Stylus nunc subnullus, nunc rectus, nunc obliquus, nunc basi flexus. Ovarium triquetrum. Capsula 3-valvis, valvis medio septi aut seminiferis. Semina angulata, glabra. Albumen farinosum. Embryo uncinato-inflexus (in Hel. vulgare Gaertn. 1. p. 371. tab. 76. fig. 11.) — Herbæ, suffrutices, frutices. Folia opposita alternaque, exstipulata vel stipulata, trinervia vel penninervia. Pedicelli sæpissimè basi bracteati, interdum oppositi-

bractei seu oppositifolii, nunc solitarii, nunc umbellati, nunc racemosi, racemis secundis, nunc corymbosi, nunc paniculati.

Genus divisum in series tres nempè :

- 1^a. Stylo recto erecto subnullo aut staminibus brevior, stigma capitato. — Sect. 1-3. HALIMIUM. LECHEOIDES. TUBERARIA.
- 2^a. Stylo recto erecto staminibus æquali aut longiora. Sect. 4-5. MACULARIA. BRACHYPETALUM.
- 3^a. Stylo basi inflexo. Sect. 6-9. ERIOCARPUM. FUMANA. PSEUDECISTUS. EUHELIANTHEMUM.

Sect. I. HALIMIUM.

Calyx 3-sepalus, sepalis æqualibus, rarò 5-sepalus, sepalis 2 externis minutis. Petala rarò alba, sæpissimè lutea, cuneata, truncata, sæpè basi maculâ atro-sanguineâ vel intensè luteâ notata. Stylus rectus brevis vel subnullus. Stigma capitatum subtrilobum. Semina nigrescentia, minutè muricata, pauca, subangulosa. — Suffrutices vel frutices. Folia opposita trinervia exstipulata pilosa vel tomentosa. Pedunc. 1-3-flori axill. solit. vel umbellati, rarò paniculati.

* *Stylo brevi recto.*

1. H. LIBANOTIS (Willd. enum 570.) caule fruticoso subglabro ramoso, foliis sessilibus linearibus margine revolutis suprâ fusco-viridibus subtùs subcanescentibus, bracteis oblongo-linearibus pedunculo brevioribus, pedunculis solitariis unifloris, calycibus trisepalis, glabris nitidis ovato-acuminatis. 5 in Lusitaniâ, Beticâ et Mauritaniâ. C. Libanotis Linn. spec. 739. Brot. fl. lus. 2. p. 261, non Lam. Desf. H. rosmarinifolium Lag. in litt. nec Pursh. — Barr. ic. 294. Ledon viii. Clus. hist. 1. p. 80. ic.* Petala luteola; stylus staminibus dimidiò brevior; stigma parvum. (v. s.)
2. H. UMBELLATUM (Mill. dict. n. 5.) caule suffruticoso ramoso, ramis junioribus tomentoso-pilosis viscosis, foliis sessilibus lineari-oblongis margine revolutis viscosis subtùs tomentosis, pedunculis unifloris racemoso-verticillatis terminalibus umbellatis, calyce trisepalo villosa. 5 in Galliâ, Hispaniâ, Lusitaniâ. C. umbellatus Linn. spec. 739. Petala alba aut albo-luteola. Stylus staminibus dimidiò brevior. Stigma parvum. (v. s.)
 - a. *erectus*, caule erecto, foliis subtùs viridi-tomentosis. C. umbellatus. 6. Lam. dict. 2. p. 18.* C. verticillatus Brot. lus. 2. p. 262.* Ledon x. Clus. hist. 1. p. 80.* (v. s.)
 - β. *subdecumbens*, caule subdecumbente, foliis subovatis subtùs albo-tomentosis. C. umbellatus a. Lam. dict. 2. p. 18.* Brot. fl. lus. 2. p. 262.*
3. H. OCYMOIDES (Pers. ench. 2. p. 76.) caule suffruticoso ramoso, ramis incanis, foliis caulinis obovatis vel ovato-oblongis trinerviis subsessilibus viridibus, ramulorum petiolatis dorso carinatis apice reflexis utrinquè incanis, pedunculis longis ramoso-paniculatis, pedicellis oppositis subumbellatis, calycibus trisepalis valdè acuminatis. 5 in Hispaniâ, Lusitaniâ. C. ocymoides Lam. dict. 2. p. 18.* C. sampucifolius Cav. ic. 1. p. 65. t. 96.* — Clus. hist. 1. p. 72. ic.* Petala lutea, rarò alba, immaculata aut maculâ atropurpureâ notata. Stamina apice atropurpurea, basi lutea. (v. s.)
 - a. Pedunculis glabris.
 - β. ramis pedunculisque laxè hirsutis. Lam. l. c.

** *Stylo subnullo, stigmate magno.*

4. H. ALYSSEOIDES (Vent. choix t. 20.) caule suffruticoso ramosissimo diffusopatulo, ramis apice incanis tomentoso-hirsutis, foliis sessilibus basi attenuatis oblongo-ovatis obtusiusculis breviter hirsutis, junioribus subincanis, adultis viridibus, pedunculis terminalibus solitariis seu umbellatis 1-2-floris folio longioribus, calycibus 3-sepalis acuminatis hirsutis. 5 in Hispaniâ, Galliâ occid. C. alyssoides a. Lam. dict. 2. p. 20.* DC. fl. fr. 4. p. 818. Petala lutea. Alabastra apice intensè purpurea. (v. s.)

5. *H. RUGOSUM* (Dunal ined.) ramis subhirsutis tomentoso-leprosis scabris fusco cinereis, foliis sessilibus in petiolum attenuatis obovato-oblongis obtusiusculis subobliquis margine subdenticulatis subcrispis utrinquē tomentoso rugosis subtus incano-cinereis, pedunculis terminalibus axillaribus seu umbellatis 1-2-floris hirsutis folio brevioribus, calycibus 3-sepalis hirsutis. ζ in Lusitaniā. (v. s.)
 β . *microphyllum*, foliis minoribus. ζ in Galliā occid. circa Bardigalam. C. alyssoides β Lam. 2. p. 20. *H. alyssoides microphyllum* DC! fl. fr. suppl. p. 622. (v. s.)
6. *H. SCABROSUM* (Pers. ench. 2. p. 76.) caule suffruticoso erectiusculo, ramis piloso-tomentosis scabris canescentibus, foliis sessilibus basi attenuatis oblongo-ovatis acutiusculis scabriusculis supra viridibus subtus tomentoso-cinereis, pedunculis terminalibus 1-2-floris folio brevioribus, calycibus 3-sepalis hirsutis. ζ in Lusitaniā boreali. C. scabrosus Ait. Kew. 2. p. 236. Brot. fl. lus. 2. p. 265.* (v. s. sine flore.)
7. *H. ALGARVENSE* (Dunal ined.) caule assurgente, foliis sessilibus, caulinis subtus subincanis ovato-lanceolatis, superioribus viridibus piloso-scabris ovato-oblongis, pedunculis subpaniculatis pilosis, calycibus 3-sepalis acutis hirsutis. ζ in Lusitaniā. C. Algarvensis Sims bot. mag. 627. Hel. Algarvense Tourn. inst. 250? Rami folia pedunculi sepalaque pilis longis laxis obsiti. (v. s.)
8. *H. FORMOSUM* (Dunal ined.) caule fruticoso, ramis tomentoso-villosis canescentibus, foliis subpetiolatis obovato-lanceolatis tomentoso-villosis, junioribus incanis, pedunculis calycibusque villosis, calycibus trisepalis. ζ in Lusitaniā. C. formosus Curt. mag. 264. Petala magna, lutea, basi maculā nigrā notata. (v. s. sine fl.)
9. *H. ATRIPLEXIFOLIUM* (Willd. enum. 569.) caule fruticoso erecto, ramis leproso-candidis, foliis petiolatis lato-ovatis obtusiusculis basi undulatis utrinquē leproso-argenteis, pedunculis racemosis hirsutis, calycibus hirsutis 3-sepalis, raro 5-sepalis, externis minutissimis. ζ in Hispaniā. C. Atriplexifolius Lam. dict. 2. p. 19.* — Barr. ic. t. 292. Caulis 4-6-pedalis et ultrā. Folia super. sessilia. Pili longi, hirsuti, rufescentes. Petala lutea, magna. (v. s.)
10. *H. LASIANTHUM* (Pers. ench. 2. p. 76.) caule suffruticoso ramossissimo, ramis nigro-cinerascentibus apice tomentoso-hirsutis, foliis subsessilibus ovato-oblongis sæpè obtusis carinatis obscurè cinerascentibus tomentosis, pedunculis 1-2-floris hirsutis brevissimis, calycibus sæpè 3-sepalis hirsutissimis. ζ in Lusitaniā. C. lasianthus Lam. dict. 2. p. 19.* — Barr. ic. t. 289? Pedunculi calyces foliaque juniora hirsuta pilis longis albidis. Calyx interdum 5-sepalis, sepalis 2 externis angustissimis apice glabriusculis. Petala lutea immaculata aut ad unguem maculata. (v. s. comm. à cl. Lag.)
11. *H. INVOLUCRATUM* (Pers. ench. 2. p. 76.) caule suffruticoso ramoso erecto, ramis tomentosiusculis cineraceis, foliis inferioribus petiolatis subovatis parvis tomentoso-incanis, superioribus oblongo-lanceolatis sessilibus subviridibus scabriusculis, pedunculis brevissimis foliis circumvallatis, calycibus 5-sepalis, sepalis internis incano-tomentosis, externis linearibus subglabris virescentibus. ζ in Hispaniā, Lusitaniā. C. involucratum Lam. dict. 2. p. 20. Ramuli foliacei axillares. (v. s.)
12. *H. CHEIRANTHOIDES* (Pers. ench. 2. p. 76.) caule fruticoso erecto ramoso, ramis junioribus villosio-tomentosis incanis, foliis tomentosis cinereo-incanis oblongo-lanceolatis in petiolum attenuatis, pedunculis brevibus subbifloris, calycibus subvillosis 5-sepalis, sepalis externis minutissimis. ζ in Lusitaniā. C. cheiranthoides Lam. dict. 2. p. 19.* C. elongatus Vahl symb. 1. p. 38*? C. halimi folio 11. Clos. hist. 1. p. 71.* Folia floralia minorā alterna. (v. s.)
13. *H. HALIMIFOLIUM* (Willd. enum. 569.) caule fruticoso erecto ramoso, ramis supernè leproso-albidis, foliis utrinquē leproso-candidis basi attenuatis subpetiolatis ovato-oblongis, pedunculis longis ramosis subpaniculatis leproso-albidis, calycibus leprosis 5-sepalis, sepalis 2 externis angustissimis linearibus

bus. ♀ in Hispaniâ, Lusitanîæ maritimis. C. halimifolius Linn. spec. 738. C. folio halimi r. Clus. hist. 1. p. 71. Folia floralia miuora alterna. Petala lutea immaculata, sæpius maculâ atro-sanguineâ notata. Capsula 2-3-valvis. (v. s.)

♂. *obtusifolium*, foliis obtusis. ♀ hab. cum var. a.

Sect. II. LECHEOIDES.

Calyx 5-sepalus, sepalis externis linearibus angustis, internis acutis margine scariosis. Petala lutea. Stylus subnullus vel brevissimus erectus. Stigma magnum capitatum. Ovarium triangulare. Capsula glabra, nitida, trivalvis, unilocularis. Semina rufescentia, parva. — Caules perennes herbacei vel suffruticosi, ascendentes seu erecti, sæpè dichotomi. Folia inferiora opposita; caulina alterna, penninervia, breviter petiolata, sessilia, exstipulata.

* *Pedunculis multifloris axillaribus seu terminalibus; floribus parvulis confertis.*

14. H. CORYMBOSUM (Michx. fl. bor. am. 1. p. 307.) caule suffruticoso ramoso erecto, ramis dichotomis subpubescentibus apice subtomentoso-cinereis, foliis caulinis alternis lanceolato-oblongis obtusiusculis subtùs cano-tomentosis, superioribus margine revolutis, corymbis fastigiatis confertis, calycibus tomentoso-hirsutis canescentibus, sepalis externis linearibus obtusis, internis ovatis acutis capsulâ subbrevioribus. ♀ in New-Jersey et Georgiâ. Stylus brevissimus. (v. s.)

15. H. ROSMARINIFOLIUM (Pursh. fl. bor. am. 2. p. 364.) caule erecto ramoso dichotomo, ramis valdè erectis pubescentibus, foliis oblongo-linearibus margine sæpè revolutis subtùs cano-tomentosis, ramulis axillaribus brevissimè pedunculatis 1-3-floris folio brevioribus, sepalis internis ovatis acutis petalis triplò minoribus. — in Georgiâ (Pursh.) Canadâ (Spreng. in litt.) circa Bostoniam (Bigel.) Lechea major Big. bost. p. 29. Flores minuti, conferti. Capsula nitida, triquetra, rufa. Petala dilatè lutea. (v. s.)

16. H. GLOMERATUM (Lag! in litt.) caule suffruticoso subdichotomo, ramis subtomentoso-cinereis, foliis lanceolato-oblongis basi attenuatis subtùs præcipuè incanis, racemis axillaribus terminalibusve multifloris folio minoribus, floribus glomeratis. ♀ prope Acapulco et Cimapan in Novâ-Hispaniâ. C. glomeratus Lag! gen. et spec. p. 16. (v. s.)

** *Pedunculis ramealibus unifloris ebracteatis.*

17. H. RAMULIFLORUM (Michx. fl. am. 1. p. 307.) caule erecto piloso-pulverulento apice ramoso subdichotomo, ramulis virgatis florigeris, foliis caulinis lanceolato-ellipticis seu oblongis acutis margine vix revolutis subtùs tomentoso-canescens, floribus pedunculatis solitariis, sepalis internis lato-ovatis acuminatis, capsulis globosis longitudine calycis. 2♂ in Carolinâ. C. virgatus Thibaud! ined. Pedunculi calycesque pilosi. Stylus brevissimus, erectus. (v. s.)

18. H. CANADENSE (Michx. fl. am. 1. p. 308.) caule ascendente, ramis erectis hirsutis, foliis caulinis oblongo-lanceolatis acutis hirsutis subtùs pallidioribus, pedunculis hirsutis unifloris solitariis, sepalis internis lato-ovatis acuminatis, capsulis calyce brevioribus. 2♂ in Canadâ et Carolinâ. C. Canadensis Willd. spec. 2. p. 199. Calyces hirsuti. (v. s.)

19. H. BRASILIENSE (Pers. ench. 2. p. 77.) caule subsimplici hirsuto, foliis ovato-oblongis acutis sessilibus hirsutis, pedunculis calycibusque hirsuto-canescens, pedunculis solitariis unifloris, sepalis internis ovato-acuminatis. 2♂ ♀ ? in Brasiliâ. C. Brasiliensis Lam. dict. 2. p. 22. C. alternifolius Vahl symb. 1. p. 38. (v. s. comm. à cl. Lag.)

20. H. CAROLINIANUM (Michx! fl. bor. am. 1. p. 307.) caulibus herbaceis hirsutis erectis, foliis tomentoso-hirsutis subdenticulatis breviter petiolatis obtusis, inferioribus oppositis obovalibus, ceteris alternis oblongo-ovatis, pedunculis solitariis unifloris hirsuto-candidis, calycibus hirsutis, sepalis internis oblongis acutis. 2♂ in Carolinâ et Georgiâ. Vent. Cata. 74. 1c. (v. s.)

Sect. III. TUBERARIA.

Calyx 5-sepalus, sepalis externis minoribus vel majoribus sæpè patulis. Petala lutea, sæpè basi maculâ atro-purpureâ notata, integra vel denticulata, serrata. Stamina numerosa pistillo multò longiora. Stylus rectus subnullus. Stigma capitatum. Capsula trivalvis. Semina minuta flavescentia. — Radices perennes lignosæ seu herbacæ annuæ. Caules herbacei erecti seu adscendentes. Folia trinervia opposita exstipulata, superiora interdum alterna, sæpius stipulata, stipulis longis linearis-acutis. Flores subpaniculati vel racemosi secundi bracteati vel ebracteati.

* *Perennia, foliis omnibus exstipulatis, caulibus basi pilosis supernè glabris, floribus paucis bracteatis subpaniculatis.*

21. *H. GLOBULARIÆFOLIUM* (Pers. ench. 2. p. 77.) perenne, caulibus adscendentibus simplicibus apice subnudis, foliis radicalibus longè petiolatis subspathulatis obtusis, caulinis sessilibus acutis, omnibus hirsutis, pedicellis basi bracteatis subcymosis paucis, calycibus glabris. \mathcal{L} in Lusitaniâ boreali. *C. globulariæfolius*. Lam. dict. 2. p. 22.* Petala maculata aut immaculata. Stamina violacea. (v. s. in h. Banks.)

22. *H. TUBERARIA* (Mill. dict. n. 10.) perenne, caulibus adscendentibus subsimplicibus, foliis radicalibus in petiolum desinentibus ovato-oblongis trinerviis tomentoso-hirsutis canescentibus subtus nervosis suprâ sulcatis, caulinis sessilibus subglabris lanceolatis, summis alternis, pedicellis basi bracteatis paucis subpaniculatis, calycibus glabris nitidis. \mathcal{L} in Gallo-provincia, Italiâ, Hispaniâ, Lusitaniâ. *C. Tuberaria* Lin. spec. 741. Cav. icon. 1. p. 65. t. 67.* (v. s.)

** *Annua? foliis superioribus sæpè stipulatis, racemis secundis terminalibus.*

23. *H. BUPLEVRIFOLIUM* (Dunal ined.) perenne?? caule herbaceo erecto basi pubescenti-canescente, supernè glabro subnitido, foliis oblongis acutis levibus longis in petiolum desinentibus, caulinis oppositis, summis alternis stipulatis, pedunculis longis piloso-pubescentibus, pedicellis calycibusque piloso-hirsutis. \mathcal{L} ??? in Hispaniâ, Lusitaniâ. *C. huplevrifolius*. Lam. dict. 2. p. 22. Pedicelli ebracteati sepala externa ovata obtusa; internis dimidiò breviora, sepala interna acuta; stipulæ sublineares, longæ. (v. s.)

24. *H. HETEROBOXUM* (Dunal ined. in h. Banks.) caule erecto herbaceo hirsuto, pilis longis albidis, foliis sessilibus oblongo-lanceolatis tomentoso-scarbis suprâ nervis hirsutis, inferioribus oppositis, summis alternis stipulatis, racemis secundis hirsuto-hirtis ebracteatis, floribus brevissimè pedicellatis approximatis subimbricatis, sepalis externis majoribus interna tegentibus bractæas simulantibus! (1) ? in Africâ propè Vallè (Masson in h. Banks.) in Hispaniâ. *H. imbricatum* Lag! in litt. cum descr. Sepala externa utrinque hirsuta, interna intus glabra, nitida margine membranacea. Capsula acutiuscula. Semina numerosa, subglobosa, obscure flavescentia, glauca, minutè muricata. (v. s.)

25. *H. PLANTAGINEUM* (Pers. ench. 2. p. 77.) caule herbaceo erecto hirsuto, foliis elliptico-lanceolatis oppositis sessilibus trinerviis, subtus villosotomentosiusculis in nervis hirsutis, suprâ hirsutis pilis simplicibus adpressis, summis oblongo-linearibus stipulatis subalternis, racemis brevibus ebracteatis, sepalis externis subglabris angusto-linearibus internos villososcanos subæquantibus, petalis denticulatis. (1) in Cretâ, Corsicâ, Hispaniâ, Africâ boreali. *C. plantagineus* Willd. spec. 2. p. 1197. *C. serratus* Desf. fl. atl. 1. p. 416. excl. Cav. syn. *C. guttatus* γ Lam. dict. 2. p. 23. Planta hirsuta, pilis albidis. Petala flava immaculata. (v. s.)

26. *H. GUTTATUM* (Mill. dict. n. 18.) caule herbaceo annuo subhirsuto, foliis oppositis sessilibus oblongo-linearibus trinerviis villosis-hirsutis, extremis alternis, racemis laxis ebracteatis, pedicellis filiformibus subnudis, sepalis externis internis dimidiò brevioribus. (1) in Angliæ, Galliæ, Italiæ, Hispaniæ.

- nizæ, Lusitanizæ, Byzantii siccis. DC! fl. fr. 4. p. 819, 5. p. 623. C. guttatus Linn. spec. 2. p. 742. C. acuminatus Viv. fragm. 13. t. 14. f. 1? (v. s. s.)
- a. *Columnæ*, petalis integris basi maculatis. Hel. flore maculoso, Col. Ephr. 2. p. 78. t. 77.
 - ß. *Cavanillesii*, petalis basi maculatis apice laciniatis. C. serratus Cav. ic. 2. p. 57. t. 57. f. 1. nec Desf.
 - γ. *Lamarckii*, petalis maculâ parvâ notatis seu subimmaculatis. C. guttatus ß. punctatus Lam. dict. 2. p. 23. An H. punctatum. n. 29?
 - δ. *exstipulatum*, caule subsimplici, foliis exstipulatis.
 - ε. *fulcraceum*, caule ramoso 2-3-chotomo, foliis superioribus stipulatis, stipulis longis.
27. H. *KRIOCAYLON* (Dunal ined.) caule herbaceo ramoso di-trichotomo hirsutissimo, foliis oblongo-linearibus angustis hirsutis oppositis, superioribus stipulatis, extremis alternis, racemis simplicibus ebracteatis, pedicellis filiformibus longis pilosis, sepalis externis angustis. ① in Hispaniâ, Galliâ. C. serratiflorus herb. Lamb! H. semistipulatum Lag! in litt. Pili caulini albidii, patentes, tennes; pili foliorum adpressi. Petala lutea, vix maculata? serrata? (v. s.)
28. H. *INCONSPICUUM* (Thib! ined. in Pers. ench. 2. p. 77.) caule herbaceo ramoso 2-3-chotomo parcè hirsuto gracili, foliis oblongo-linearibus angustis hirsutis oppositis, superioribus stipulatis, extremis alternis, racemis longis filiformibus, pedicellis brevibus secundis, floribus minutis, petalis oblongo-linearibus calyce minoribus altero margine pilosis. In Hispaniâ, Corsicâ. H. præcox Saltzm. exsic. (v. s.)
29. H. *PUNCTATUM* (Willd. enum. 570.) caule herbaceo ramoso dichotomo tomentosiusculo-pubescente subcinereo, foliis oblongis 3-5-nerviis viridicineris breviter piloso-scabriusculis, pilis stellatis minimis, inferioribus oppositis obtusis?, superioribus alternis acutiusculis stipulatis, racemis longis pubescentibus cinereis paucifloris. ① in Galliâ occid. C. punctatum Willd. spec. 2. p. 1199. * Pedunculi rarò medio bracteam magnam seu folium parvum gerentes. Pedicelli filiformes, longi. Petala serrulata. (v. s.)

Sect. IV. MACULARIA.

Calyx 5-sepalus, sepalis externis angustis, internis striatis. Petala lutea, basi maculata. Stylus rectus erectusque ovario duplò longior, staminibus subæqualis. Stigma parvum, subtrilobum. Capsula lævis. — Suffrutescens aut herbæ? Folia petiolata, penninervia, angusta, exstipulata. Flores terminales, solitarii seu racemosi, racemis paucifloris, pedicelli secundi, basi bracteati, bracteis subulatis parvulis.

30. H. *LUNULATUM* (DC! fl. fr. 4. p. 816.) caule suffruticoso tortuoso ramoso, ramulis pubescentibus filiformibus, foliis planis oblongis basi acuminatis margine sæpè ciliatis, floribus solitariis seu racemoso-subumbellatis 2-4 terminalibus breviter pedicellatis. § in Alpibus Pedemontanis. C. lunulatus All! auct. p. 30. t. 2. f. 3. * Calyx per anthesin reflexus. Petala lutea, subintegra, lævissimè crenulata, lunulâ crocèâ versus unguem notata. (v. s.)
- a. foliis suprâ viridibus, subtùs subalbescentibus.
 - ß. foliis minoribus utrinquè incanis. Requien! herb.

31. H. *PETIOLATUM* (Thib! in Pers. ench. 2. p. 77.) caule herbaceo? basi tortuoso, ramis subdichotomis adscendentibus gracilibus tomentoso-leprosis cinereis, foliis oblongo-linearibus acutis in petiolum longum attenuatis utrinquè leproso-tomentosis, subtùs incanis suprâ viridi-glaucis racemis parvis paucifloris, pedicellis calycibusque pubescentibus. ①? in Hispaniâ. Petala stamina pistillumque ignota. (v. s.)

Sect. V. BRACHYPETALUM.

Calyx 5-sepalus, sepalis externis minutis, internis 3-nerviis acuminatis.

Petala lutea, interdum basi maculata, parva, sæpè calyce breviora. Stamina pauca 10-20, ovarium cingentia. Stylus rectus et erectus apice incrassatus. Stigma simplex. Ovarium triquetrum, angulis sæpè pilosis. Capsula triquetra, subuitida. Semina numerosa, minuta, pallida, angulosa. — *Herbæ annuæ.* Folia stipulata, petiolata, penninervia, subdenticulata, opposita, floralia alterna. Stipulæ oblongo-lineares, superiores longiores. Pedunculi uniflori, breves, solitarii, rarè axillares, sæpius suboppositifolii vel oppositibractei, suberecti, horizontales vel retrofracti.

* *Pedunculis suberectis foliis brevioribus, sepalis internis 3-nerviis.*

32. *H. VILLOSUM* (Thib! in Pers. ench. 2. p. 78.) caule tomentoso hirsuto subcinereo, foliis petiolatis oblongo-lanceolatis subdenticulatis utrinquè præcipuè subtus villosis stipulatis, racemis longis secundis, antè anthesin apice revolutis, axillaribus terminalibusque, pedunculis erectis villosis-cinereis suboppositibracteis, calycibus oblongis acuminatis villosis. (1) in Hispaniâ. *C. villosus* Thib! herb. Lag! in litt. nec Lam. *C. annuus* Lamb! herb. Bracteæ subsessiles ovato-oblongæ, interdum parè dentatæ. Petala lanceolata angusta sepalis breviora, plerumque denticulata. (v. s.)

a. caule simplicissimo erecto.

β. caule basi ramoso, ramis longis simplicibus adscendentibus.

33. *H. NILOTICUM* (Pers. ench. 2. p. 78.) caule herbaceo, ramis erectis vel adscendentibus tomentosiusculis seu villosis cinereis, foliis breviter petiolatis oblongo-ellipticis tomentoso-villosis oppositis, superioribus alternis floribus oppositis, omnibus stipulatis, pedunculis erectis et calycibus acuminatis tomentoso-hirsutis. (1) in AEGypto, Barbariâ, Hispaniâ, Galliâ austr. *C. Niloticus* Liu. mant. 246.

a. *erectum*, caule erecto simplici seu ramoso, ramis adscendentibus, pedunculis calycibusque villosis-hirsutis. (v. s.)

β. *majus*, caule erecto ramoso, ramis adscendentibus, ramis foliis pedunculisque tomentosiusculis. Var. culta ex omni parte major. (v. s.)

γ. *procumbens*, caulibus procumbentibus adscendentibus tomentoso-hirsutis subincanis, foliis utrinquè præcipuè subtus tomentosis. *C. ledifolius* Gouan fl. monsp. p. 264? *C. Ger. gall. prov. p. 392. n. 2.* *Au species diversa?* (v. s.)

34. *H. LEDIFOLIUM* (Willd. enum. 571.) caule erecto subglabro, foliis petiolatis oblongo-ellipticis subdenticulatis glabriusculis oppositis, superioribus alternis, floribus oppositis stipulatis, pedunculis erectis glabriusculis calyce brevioribus, calycibus acuminatis subpubescentibus. (2) in Galliâ australi? Lusitaniâ. *C. ledifolius* Liu. spec. 742. *C. annuus folio ledi* Lob. ic. 2. p. 118. Valdè affinis *H. Nilotico*. (v. s.)

** *Pedunculis horizontalibus foliis seu bracteis longioribus, floribus erectis; sepalis internis 3-nerviis.*

35. *H. INTERMEDIUM* (Thib! ined.) caule erecto ramoso, ramis gracilibus erectis vel patulis adscendentibus villosis-subcinereis, foliis petiolatis obovato-oblongis obtusiusculis subdenticulatis venosis leviter tomentosis stipulatis, stipulis linearibus oblongis, superioribus petiolo vix duplò longioribus, pedunculis calycibusque villosis-cinereis suboppositifolii, calycibus oblongis. (1) in Hispaniâ propè Aranjuez. *H. salicifolium intermedium* Pers. ench. 2. p. 78. *C. salicifolius* Cav! ic. n. 156. t. 144. Planta spithamea, subcinerea. Folia floralia seu bracteæ alterna, lineari-oblonga sæpè petiolata, 1-2-stipulata, interdum solitaria, integra vel incisa, sæpissimè pedunculis breviora. Flores exiles. (v. s.)

36. *H. DENTICULATUM* (Thib! in Pers. ench. 2. p. 78.) caule ramoso erecto vel patulo, ramis erectis vel adscendentibus minutè tomentoso-pubescentibus apice subcinereis, foliis breviter petiolatis obovato-oblongis acutiusculis subdenticulatis tomentosis supra virescentibus subtus incano-cinereis,

atipulis linearibus, superioribus folio dimidiò brevioribus, pedunculis calycibusque oppositis bracteis, bracteis alternis subincisis. ① in Galliâ austr. C. salicifolius Gouan. herb. p. 34? Bracteæ subovatæ, sæpè incisæ, sessiles, exstipulatæ. Calyces antè anthesin extûs incani. (v. s.)

37. H. SALICIFOLIUM (Pers. ench. 2. p. 78.) caule ramoso, ramis erectis vel adscendentibus subhirsutis, foliis breviter petiolatis obovato-oblongis acutiusculis denticulatis subtomentosis suprâ virescentibus, stipulis lineari-oblongis, superioribus folio dimidiò brevioribus, pedunculis calycibusque hirsutis suboppositibracteis, bracteis ovatis acuminatis sessilibus integris. ① in Hispaniâ, Italiâ. C. salicifolius. Linn. spec. 742. H. annuum, etc. Seg. ver. 3. p. 297. t. 6. f. 3? Rami pedunculi calycesque hirsuti, pilis tenuibus albidis. (v. s.)

β. latifolium, tota planta maxima, foliis bracteisque læviter tomentosis virescentibus. ① in Oriente. (v. s.)

*** Pedunculis ante anthesin cernuis, per anthesin erectis, post anthesin retroflexis; sepalis interioribus 4-nerviis.

38. H. SANGUINEUM (Lag! in litt.) caule herbaceo pumilo pubescenti-viscido sanguineo, foliis petiolatis oppositis ovatis obtusis scabriusculis, inferioribus exstipulatis subtus sanguineis, superioribus stipulatis, stipulis oblongo-linearibus obtusis petiolatis folio vix brevioribus, pedunculis piloso-viscidis axillaribus oppositifoliisque, fructiferis retroflexis. ① in Hispaniâ propè Chamartin. C. sanguineus Lag! gen. et spec. nov. p. 17. varied. ann. 2. n. 19. p. 40. H. retrofractum Pers! ench. 2. p. 78. C. pusillus herb. Lamb! Folia omnia opposita. Pedunculi semper axillares et oppositifolii. Sepala interna striata. (v. s.)

39. H. AEGYPTIACUM (Mill. dict. n. 23.) caule herbaceo pubescente erecto vel adscendente, foliis breviter petiolatis lineari-oblongis angustis margine revolutis obtusiusculis subtus pallidè cinereis oppositis, summis alternis stipulatis, stipulis lineari subulatis, pedunculis filiformibus pubescentibus, foliis apremis oppositis, calycibus ovato-oblongis inflatis petala includentibus. ① in AEGypto, Barbariâ, Hispaniâ. C. Aegyptiacus Linn. spec. 742. Jacq. obs. 3. p. 17. t. 68.* Pedunculi apice incrassati, interdum bracteis brevibus linearibus oppositi. Sepala externa angusta brevia, interna 4-nervia, nervis ciliatis. Petala lanceolata, brevissima. (v. s.)

Sect. VI. ERIOCARPUM.

Calyx 5-sepalus, sepalis extûs piloso-sericeis seu tomentosisculis intûs nitidis, 2 externis minutis linearibus, 3 internis ovatis 4-5-striatis. Petala calyce paulò longiora. Stylus subrectus basi flexus. Ovarium pilosum seu villosum. Capsula pilosa. Semina numerosa, rufescentia, minima. — Suffrutices; rami teretes, juniores cinereo-pubescentes. Folia opposita alternaque, subtuscano-cinerea obtusiuscula. Stipulæ petiolo breviores lineares. Racemi secundi parvi oppositifolii, floribus confertis parvulis sessilibus seu majoribus breviter pedicellatis.

40. H. LIPPII (Pers. ench. 2. p. 78.) caule suffruticoso erecto pubescente albicante subbifido subdichotomo, foliis oppositis alternisque breviter petiolatis elliptico-lanceolatis seu lineari-oblongis obtusis subscabris glaucescentibus subtus canescentibus, stipulis angustis erectis longitudine petiolorum, racemis brevibus, floribus sessilibus confertis basi bracteatis, bracteis minutissimis. ③ in AEGypto. C. Lippii Linn. mant. 245. Vahl symb. 1. p. 39. Sepala pubescentia, interna 4-5-costata obtusa. Petala ovata, flava, calyce vix longiora. Stamina sæpius 10 petalis breviora. (v. s.)

41. H. SÆSSILIFLORUM (Pers. ench. 2. p. 78.) caule suffruticoso erecto ramossimo, ramis pubescentibus, foliis oppositis alternisque linearibus brevissimè tomentosis cinereis margine revolutis, stipulis linearibus parvulis, racemis brevibus, floribus sessilibus minutè bracteatis. ③ in Africâ bor.

- collibus aridis. *C. sessiliflorus* Desf! fl. atl. 1. p. 427. t. 106.* Sepala pubescentia, interna obtusa. Petala flava calyce paulò longiora. (v. s.)
42. *H. ELLIPTICUM* (Pers. l. c.) caule suffruticoso ramoso erecto cinereo, foliis oppositis ellipticis leproso-tomentosis cinereo-candidis obtusis margine revolutis, stipulis linearibus parvis, racemis paucifloris, floribus sessilibus bracteatis subalternis. § in Barbariâ, Ægypto. *C. ellipticus* Desf. fl. atl. 1. p. 418. t. 107.* *H. Lippii* Delil. ægypt. 93. *C. stipulatus* α Forsk. ægypt. p. 101. Sepala villosa-tomentosa, interna ovata obtusiuscula. Petala flava calyce paulò longiora. Folia superiora subalterna. (v. s.)
43. *H. KAHIRICUM* (Delil. fl. æg. 93. t. 31. f. 2.) caule ramosissimo basi tortuoso, ramis adscendentibus, foliis inferioribus oppositis, ceteris alternis obovatis margine revolutis incanis stipulatis subtus nervosis, racemis secundis, floribus breviter pedicellatis, pedicellis calycibusque villosis, sepalis acutis, capsulis oblongis villosis. § in Ægypto. *C. stipulatus* β. Forsk. ægypt. 101. Petala apice coalita. (v. s.)
44. *H. CONFERTUM* (Dunal. ined.) caule suffruticoso basi nudo ramoso, ramis brevissimè tomentosis cinereis, foliis lanceolato-ellipticis obtusiusculis utrinquè minutissimè tomentosis suprâ virescentibus subtus canescentibus, stipulis linearibus parvis, racemis parvis secundis oppositifoliis terminalibusque, floribus confertis subsessilibus, calycibus pilosis, sepalis internis acutis. § in insulâ Teneriffæ. Broussonet. (v. s.)
45. *H. CANARIENSE* (Willd. enum. 571.) caule suffruticoso procumbente, ramis subtomentosis incano-cinereis, foliis petiolatis oppositis alternisque subovato-ellipticis obtusis utrinquè brevissimè tomentosis glaucis, subtus incano-cinereis, stipulis subulatis petiolo brevioribus, racemis terminalibus erectis minutè bracteatis, floribus brevè pedicellatis, pedicellis tomentoso-incanis. § in Canariâ magnæ et Lancerottæ aridis. *C. Canariensis* Jacq. ic. 1. p. 97. misc. 2. p. 339. *H. canescens* Moench. Calyx glaucus; sejala interna ovata, obtusiuscula. (v. s.)
46. *H. MUCRONATUM* (Dunal. ined.) caule suberecto patulo, ramis tomentoso-hirsutis incanis, foliis petiolatis ovato-ellipticis ventricosis mucronatis, suprâ viridibus scabriusculis pilis stellatis, subtus tomentoso-incanis, stipulis subulatis pilosis petiolo brevioribus, racemis subterminalibus, floribus brevè pedicellatis, pedicellis tomentoso-hirsutis. § in Teneriffâ. Broussonet. Calyx piloso-sericeus, sepalis lato-ovatis obtusiusculis. (v. s.)

Sect. VII. FUMANA.

Calyx antè anthesin apice tortus 5-sepalus, sepalis externis angustis parvulis, internis 3 ovato-acuminatis 4-5-venosis margine scariosis. Petala lutea parva calyce subduplò longiora. Stamina pauca. Stylus rectus staminibus sublongior, per anthesin obliquus, post anthesin suberectus. Stigma capitatum, fimbriatum sub-3-lobum. Capsula 3-valvis, aperta patula; semina nigrescentia vel rufescentia angulosa pauca. — Caules suffruticosi. Folia linearia sessilia aut subsessilia, angusta. Pedicelli uniflori, ante anthesin cernui, per anthesin erecti, post anthesin reflexi.

* *Foliis alternis exstipulatis.*

47. *H. XERICOIDES* (Dunal. ined.) caule fruticoso erecto, foliis alternis imbricatis semi-cylindricis brevibus glabriusculis, pedunculis solitariis unifloris suboppositifoliis terminalibusve folio longioribus, capsulis apertis nudis? § *C. ericoides* Cav. ic. 2. p. 56. t. 172.* *C. calycinus* plurimorum auctorum non ut videtur Linn. mant. 565? An sequentis varietas? Folia margine subrevoluta.
- α. *glabrum*, ramis pedunculisque glabris. § in Hispaniâ (v. s.)
β. *pubescens*, ramorum summitatibus pedunculisque pubescentibus. § in regno Neapolitano. (v. s.)
48. *H. FUMANA* (Mill. dict. n. 6.) caule suffruticoso ramoso tortuoso subdif-

fuso erectiusculo, ramis inferioribus procumbentibus, foliis alternis linearibus margine piloso scabrisculis subinvolutis, inferioribus brevibus confertis, superioribus sparsis longioribus, pedunculis solitariis unifloris raro ramealibus sæpius suboppositifoliis terminalibusve folio longioribus, capsulis apertis nudis. ζ in Galliâ austr. Hispaniâ, Lusitaniâ, Helvetiâ, Gotlandiâ, Italiâ. C. Fumana Linn. spec. 740. C. humilis seu chamæcistus ericæ folio luteus erectior Bauh. pin. 466. Magn. bot. 69. Pedicelli calycesque sparsè pilosi.

α . *majus*, foliis majoribus. C. Fumana A. Desf. fl. atl. 1. p. 414. t. 105. exclus. syn. Linn. et Barr. (v. s.)

β . *minus*, foliis subfiliformibus etc. Barr. ic. 286 et 446. (v. s.)

γ . *virgatum*, ramis virgatis. H. fumanaoides, olim in horto Parisiensi. (v. s.)

49. H. *PROCUMBENS* (Dunal. ined.) caule suffruticoso procumbente ramoso, ramis elongatis junioribus canescentibus, foliis alternis linearibus subulatis margine et subtus pilosis, pilis strigosis. pedunculis subaxillaribus folio brevioribus, capsulis apertis semina gerentibus. ζ in Galliâ austr. Italiâ, Tauriâ. — Barr. ic. t. 445. C. humilis aive chamæcistus ericæ folio humilior Magn. bot. p. 69. Capsula major. Nervi calycini strigosi. « Hos rectè separamus (inquit Magnol) quia non tantum magnitudine differunt sed et modo conservandi seminis, siquidem erectioris (fumane) capsula dum aperitur, decidit statim semen, sicut aliis accidit; at in humiliori (procumbenti) quamvis capsula aperta sit et prona versus terram, manet tamen semen quasi capsulæ adherens. » (v. s.)

** *Foliis alternis stipulatis.*

60. H. *ARABICUM* (Pers. ench. 2. p. 80.) caule suffruticoso hirsuto adacendente, ramis virgatis, foliis alternis lineari-oblongis hirsutis subsessilibus, pedunculis solitariis unifloris suboppositifoliis ramealibus terminalibusve, calycibus hirsutis. ζ in Arabiâ, Italiâ, Hispaniâ. C. Arabicus Linn. spec. 745. C. ferrugineus Lam. dict. 2. p. 25.* C. Savi Bertol! H. viscidulum Stev! — Folia superiora majora. Stipulæ ovato-acuminatæ. (v. s.)

61. H. *LÆVIPS* (Willd. enum. 570.) caule fruticoso ad-cendente, foliis stipulatis setaceis glaucis subglabris, gemmis foliaceis axillaribus, stipulis filiformibus longis, pedunculis longis racemosis secundis, pedicellis glabris basi bracteatis, calycibus hirsutis. ζ in Gallo-Provinciæ austr. Hispaniæ, Dalmatiæ rupibus apricis. C. lævipes Linn. spec. 739. Ger. gallo-prov. p. 394. n. 6. t. 14. (v. s.)

α . pedunculis pubescentibus.

β . pedunculis subglabris.

*** *Foliis oppositis alternisque stipulatis.*

62. H. *LÆVIS* (Pers. ench. 2. p. 78.) caule suffruticoso suberecto glabro ramoso, ramis erectis, foliis linearibus sessilibus glabris margine revolutis carinatis oppositis, summis alternis stipulatis, stipulis subulatis longis, pedunculis solitariis unifloris subterminalibus, calycibus lævibus. ζ in Hispaniæ collibus. C. lævis Cav! ic. 2. p. 35. t. 145. f. 1.* exclus. syn. Barr. (v. s.)

63. H. *VIRIDE* (Tenor. prod. fl. neap. p. 31.) caule suffruticoso adscendente glabro, foliis oppositis linearibus margine revolutis glabris viridibus submucronatis stipulatis, stipulis subulatis folio multò minoribus, pedunculis racemosis calycibusque villosis-glutinosiis. ζ in Siciiliâ. Folia lætè viridia. (v. s.)

54. H. *JUNIPERINUM* (Lag! in litt.) caule suffruticoso adscendente ramoso, foliis lineari-subulatis ciliatis mucronatis planis margine subrevolutis oppositis, superioribus alternis, stipulis subulatis, superioribus longioribus, peduncu-

- lis racemosis calycibusque villosis-glutinosus. \S in Galliâ australi, Italiâ, regno Tunetano. C. lævipes Durand! Gouan. fl. monsp. 263? C. mauritanicus Thib! ined. — Barr. ic. t. 443. Bracteæ solitariae, lineares. (v. s.)
55. H. *BARRERIERI* (Tenor! prodr. fl. neap. p. 31.) caule suffruticoso erecto, ramis villosis-pubescentibus, foliis lineari-oblongis basi attenuatis angustis pubescentibus margine revolutis et ciliatis oppositis, summis alternis, stipulis lineari-subulatis mucronatis erectis, pedunculis racemosis paucifloris calycibusque villosis-glutinosus. \S in Italiâ, Hispaniâ. — Barr. rar. ic. 416. (v. s.)
56. H. *THYMIFOLIUM* (Pers. ench. 2. p. 79.) caule suffruticoso procumbente, ramis pubescentibus, foliis sublinearibus brevissimis pubescentibus oppositis, summis alternis, stipulis mucronatis erectis, pedunculis villosis-glutinosus paucifloris. \S in Hispaniâ. C. thymifolius Linn. spec. 743. — Barr. rar. ic. t. 444. H. glutinosum β . fl. fr. 4. p. 821. (v. s.)
57. H. *GLUTINOSUM* (Pers. ench. 2. p. 79.) caule suffruticoso ascendente, ramis villosis-glutinosus subcinereis, foliis sublinearibus margine revolutis villosis-glutinosus subcinereis oppositis, summis alternis, stipulis inferioribus minutis, ceteris longis laxiusculis, pedunculis calycibusque villosis-glutinosus. \S in Galliâ austr. Hispaniâ. C. glutinosus Linn. mant. 246. — Barr. rar. 512. t. 415? (v. s.)

SECT. VIII. *PSYDROCISTUS.*

Calyx 5-sepalus, sepalis externis angustis minutis, internis 4-venis. Petala lutea, parva, calyce vix duplò longiora. Stylus basi contortus retroflexus apice inflexus, staminibus sæpè brevior, rarò longior. Stigma capitatum, trilobum. Capsulae parvae. Semina pauca, subrufescentia. — Herbae perennes aut suffrutices. Folia petiolata, penninervia, opposita, exstipulata, rarò stipulata in summitate ramorum. Flores secundi, racemosi seu paniculati. Pedicelli basi bracteati, bracteis sessilibus lineari-lanceolatis, antè anthesin recurvi, per anthesin erecti, post anthesin reflexi.

58. H. *MOLLE* (Pers. ench. 2. p. 76.) caule suffruticoso, ramis subsimplicibus piloso-hirsutis, foliis subrotundo-ovatis obtusis petiolatis utrinque hirsutotomentosis mollibus, racemis simplicibus calycibusque hirsutotomentosis cinereis. \S in Hispaniâ. C. mollis Cav. ic. 3. p. 32. t. 262. f. 2. Bracteae subulatae minutae, piloso-tomentosae, cinerae (v. s.)
59. H. *ORIGANIFOLIUM* (Pers. ench. 2. p. 76.) caule suffruticoso 2-3-chotomo, foliis petiolatis ovatis utrinque pilosis, racemis brevibus terminalibus, petalis calyce vix majoribus. \S in Hispaniâ. C. origanifolius Lam. dict. 2. p. 20.* Cav. ic. 3. p. 31. t. 262. f. 1.* Calyx oblongus. Petala duplò minora quam in H. molli. (v. s.)
60. H. *DICHOTOMUM* (Dunal. ined.) caule suffruticoso, ramis dichotomis subglabris, foliis minutis ovatis acutis glabris margine revolutis breviter petiolatis, racemis tenuibus paucifloris. \S in Hispaniâ. C. dichotomus Cav. ic. 3. p. 32. t. 263. f. 1.* Flores parvi, saturatè flavi, vix magnitudine floris Spergulae nodosae. Folia parva facie Thymi piperellae (v. s. sine fl.)
61. H. *OELANDICUM* (DC! fl. fr. 4. p. 817.) caule suffruticoso procumbente ramoso, foliis lanceolato-ellipticis obtusiusculis utrinque viridibus sæpè glabris interdum ciliatis petiolatis, summis sessilibus, racemis simplicibus paucifloris, calycibus subgloboso-ovatis. \S in Alpibus Europae praesertim borealis. C. Oelandicus Linn. sp. 741. Chamæcistus 2. Clus. hist. p. 73. β . Bracteae minutae sessiles, lineari-oblongae. Flores pauci, approximati. Petala calyce vix longiora. (v. s.)
62. H. *ALPESTRIS* (Dunal. ined.) caule suffruticoso procumbente ramoso, ramis piloso-hirsutis, foliis utrinque subviridibus oblongo-ellipticis subglabris vel fasciculatim piloso-hirsutis petiolatis, summis subsessilibus, pedicellis calycibusque piloso-hirsutis, pilis albido-cinereis. \S in petrosis Germaniae, Helvetiae, Italiae, Galliae. C. alpestris Crantz austr. p. 103. t. 6. f. 1. Wahlenb.

helv. p. 103. C. Oelandicus Jacq. austr. t. 399. Petala calyce duplò longiora. (v. v. et s.)

α. *glabratum*, foliis subglabris oblongo-ellipticis obtusiusculis, petiolis pilosis.

β. *elongatum*, ramis elongatis, foliis acutis utrinquè pilosis. C. Segneri Poarr. ined.

γ. *canescens*, foliis junioribus pedunculis calycibusque pilosis cinereo-canescentibus C. Oelandicus Gouan fl. monsp. p. 263?

63. H. *PERICILLATUM* (Thib! ined.) caule suffruticoso, ramis procumbentibus longis hispidis, foliis viridibus utrinquè nervis et margine hispidis, inferioribus petiolatis ovatis minoribus, superioribus lineari-oblongis subsessilibus, racemis simplicibus calycibusque hispidis, floribus minutis. ♂ in Hispaniâ et Galliâ australiori. C. echioides Lam. dict. 2. p. 21? C. Anglicus Linn. mant. 245? habitus *Myosotis Lappulæ*. (v. s.)

64. H. *OBOVATUM* (Dunal. ined.) caule suffruticoso, ramis patentibus subdichotomis apice tomentoso-cinereis, foliis obovatis vel oblongis obtusis utrinquè viridibus ciliatis læviter piloso-strigosis, inferioribus minutis, racemis simplicibus paucifloris, bracteis viridibus, calycibus piloso-cinereis. ♂ in Hispaniâ propè Aranjuez. C. Italicus Linn. spec. 740. exclus. syn? Folia in brevem petiolum desinentia. (v. s.)

65. H. *ITALICUM* (Pers. ench. 2. p. 76.) caule suffruticoso, ramis simplicibus erectis longis piloso-tomentosis, foliis piloso-hispidis, pilis strigosis adpressis, inferioribus ovatis minoribus, superioribus lanceolatis oblongis vel oblongo-linearibus, racemis simplicibus calycibusque piloso-hispidis canescentibus. ♂ in siccis regionis Mediterraneæ. C. Italicus Linn. spec. 740. C. marifolius Bieb. fl. taur. cauc. 2. p. 8. — Barr. rar. 510. t. 366. (v. s.)

α. *strigosum*, virescens, foliis utrinquè subviridibus piloso-strigosis, pedunculis calycibusque piloso-tomentosis incanis. H. strigosum Fisch! in litt.

β. *candissimum*, foliis pedunculis calycibusque tomentoso-candidis.

γ. *albidum*, foliis subtus tomentosiusculis albidis.

66. H. *VINEALE* (Pers. ench. 2. p. 77.) caule suffruticoso procumbente, ramis adscendentibus piloso-tomentosis canescentibus, foliis ovato-oblongis supra viridibus piloso strigosis, subtus tomentoso-incanis, racemis simplicibus paucifloris calycibusque piloso-tomentosis canescentibus. ♂ in Germaniâ, Helvetiâ, Galliâ, Hispaniâ. C. vinealis Willd. spec. 2. p. 1195. An H. cani varietas? (v. s.)

67. H. *CANUM* (Dunal. ined.) caule suffruticoso procumbente ramoso, ramis adscendentibus piloso-tomentosis incanis, foliis obovatis ovatis ovato-oblongis vel ellipticis pilosis supra virescentibus subtus subtomentoso-canis, racemis simplicibus, pedicellis calycibusque piloso-canescantibus. ♂ in Hispaniâ, Galliâ austr. Germaniâ. C. canus Linn. spec. 740. Chamæcistus 3. Clus. hist. p. 74.* Folia variabilia, pilis strigosis adpressis subtus margine, nervo medio et supra ubiquè oblecta. Ovarium triquetrum, angulis marginatis pilosis. (v. s.)

68. H. *MARIFOLIUM* (DC! fl. fr. 4. p. 817.) caule suffruticoso procumbente, foliis extipulatis petiolatis ovato-cordatis vel ovatis acutiusculis supra viridibus pilosis, subtus incanis, racemis solitariis simplicibus paucifloris terminalibus. ♂ in Italiâ, Hispaniâ, Galliâ austr. C. marifolius Linn. spec. 741. nec Bieb. flor. taur. — Barr. rar. 521. t. 441. Calyces ovato-oblongi. (v. s.)

69. H. *ROTUNDIFOLIUM* (Dunal. ined.) caule suffruticoso basi ramoso, ramis simplicibus tomentoso-incanis basi plerumque decumbentibus, foliis breviter petiolatis supra viridi-glaucescentibus subtus tomentoso-candidis, inferioribus suborbiculatis, ceteris ovatis, summis stipulatis, stipulis parvis oblongis deciduis, racemis solitariis bi-ternisve subpaniculatis confertis terminalibus, calycibus hirsutis. ♂ in Hispaniâ, Barbariâ. C. nummularia

Cav! ic. 2. p. 34. t. 142.* Desf! fl. atl. 1. p. 423. exclus. syn. Linn. et Magn. (v. s.)

70. *H. CRASSIFOLIUM* (Pers. ench. 2. p. 77.) caule suffruticoso erecto subglabro, foliis subcarnosis breviter petiolatis inferioribus ovatis acutis exstipulatis, superioribus oblongo-linearibus stipulatis, racemis brevibus subumbellatis, calycibus basi pilosis. ζ in Barbariâ et Hispaniâ. *C. glaucus* Desf. fl. atl. 1. p. 418.* nec Cav. *H. Setze* Lag! in litt. *Setze* vulgò in regno Valentino. Folia suprâ ad margines, subtusque nervo medio pilosiuscula; petioli pilosi, pilis raris longis albidis. (v. s.)

71. *H. PANICULATUM* (Dunal. ined.) caule suffruticoso procumbente, ramis adscendentibus erectisque floriferis longis supernè stipulatis, foliis petiolatis ovatis obtusiusculis rarò subrotundatis suprâ viridibus subtus incanis, racemis oppositis ternisque paniculatis. ζ in Hispaniâ et Siciliâ montosis. *C. marifolius* herb. Thib! *C. nummularius* var. Lag! in litt. *H. sp. nov.* Schouw! in litt. Stipulæ minutæ, lineari-acutæ. Flores parvi. (v. s.)

72. *H. POLYANTHOS* (Pers. ench. 2. p. 78.) caule suffruticoso erecto hirsuto, foliis petiolatis, inferioribus ovatis obtusis minoribus subtus tomentoso-incanis, caulinis ovato-oblongis seu lanceolatis utrinquè virescentibus margine ciliatis stipulatis, stipulis petiolo longioribus, racemis hirsutis paniculatis, pedicellis filiformibus calycibusque hirsutis. ζ in Africâ boreali. *C. polyanthos* Desf! fl. atl. 1. p. 420. t. 108.* (v. s.)

73. *H. CINEREUM* (Pers. ench. 2. p. 76.) caule suffruticoso erecto ramoso, ramis oppositis incanis, foliis ovatis acutis in petiolum attenuatis dense tomentosis incano-cinereis exstipulatis, summis stipulatis, racemis paniculatis axillaribus oppositis, terminalibus ternis, calycibus hispidis. ζ in Hispaniâ *C. cinereus* Cav! ic. 2. p. 33. t. 141.* Flores parvi. Petala integra. Cav. (v. s.)

β . *Lagascanum*, caulibus tenuibus, calycibus minùs hispidis. Lag. in litt.

74. *H. SQUAMMATUM* (Pers. ench. 2. p. 78.) caule suffruticoso, ramis longis erectis sublignosis leproso-argenteis, foliis petiolatis oblongo-obtusis leproso-argenteis stipulatis, stipulis parvis sessilibus acutis marcescentibus, racemis axillaribus solitariis ternisve terminalibus, pedicellis approximatis secundis basi bracteatis, bracteis marcescentibus, calycibus leprosis. ζ in Hispaniâ et Barbariâ. *C. squammatus* Linn. spec 743. Cav. ic. t. 139. Desf! fl. atl. 1. p. 416. — Barr. rar. ic. t. 328. male. Rami basi tetragoni. Planta leprosa, squamis orbiculatis centro depressis. Stylus basi contortus flexus staminibus longior. Rami foliacei axillares. (v. s.)

SECT. IX. EUHELIANTHEMUM.

Calyx ante anthesin apice subtortus, 5-sepalus, sepalis externis sæpè patulis multò minoribus, internis 2 sæpiùs 4-costatis sulcatis margine scariosis intùs nitidis, angulis sæpè pilosis. Petala calycibus 2-3-4-plo longiora. Stamina numerosa. Stylus basi flexus apice subclavatus. Stigma simplex. Capsula calyce oblecta trivalvis unilocularis apice dehiscens. Semina pauca extùs convexa, intùs angulosa. — Suffrutices; caules basi ramosi, ramis numerosis erectis vel procumbentibus, sæpiùs adscendentibus. Folia opposita, breviter petiolata, inferiora minorâ, sæpè margine revoluta stipulata, stipulis lineari-lanceolatis. Racemi terminales secundi simplices, antè anthesin incurvi, post anthesin erecti elongati. Pedicelli basi lateraliter bracteati, antè anthesin cernui, per anthesin erecti, post anthesin recurvi reflexi.

* *Petalis luteis.*

75. *H. LAVANDULIFOLIUM* (DC! fl. fr. 4. p. 820.) caule suffruticoso erecto ramoso, ramis longis teretibus canescentibus, foliis oblongo-linearibus margine revolutis subtus tomentoso-incanis, junioribus utrinquè canescentibus, stipulis bracteisque lineari-acutis ciliatis, racemis 1-3 terminalibus,

floribus confertis, calycibus glaucis, sepalis ciliatis externis minutis post anthesin reflexis, internis 2-nerviis obliquis acutis. ζ in Galliæ austr., Barbariæ, Hispaniæ, Syriæ locis siccis. C. lavandulæfolius Lam! dict. 2. p. 25.*—Barr. ic. t. 288. Rami foliacei axillares. (v. s.)

β . *Syriacum*, foliis subplanis suprâ viridi-griseis. C. Syriacus Jacq. ic. rar. t. 96. (v. s.)

γ . *Thibaudi* (Pers! ench. 2. p. 79.) racemis longis erectis. C. racemosus Cav. ic. 2. p. 33. t. 140.* eadem planta post anthesin. (v. s.)

76. H. *Broussonetii* (Dunal. ined.) caule fruticoso ramoso, ramis oppositis, foliis breviter petiolatis planis oblongo-lanceolatis obtusiusculis utrinquè tomentosis, subtus incanis, suprâ viridi-cinereis, stipulis bracteisque caducis linearibus tomentosisculis, racemis brevibus ramosis, floribus secundis, calycibus oblongis acutis, sepalis internis 4-nerviis tomentosisculis lutescentibus. ζ in insulâ Teneriffæ. Stylus staminibus duplò longior, subrectus. Stipulæ subfalcatæ. (v. s.)

77. H. *stæchadiifolium* (Pers. ench. 2. p. 79.) caule erecto, ramis tomentoso-canescensculis, foliis oblongo-linearibus obtusiusculis utrinquè subtomentosis, suprâ viridi-cinereis, subtus incanis margine revolutis, stipulis subvillosis lineari-lanceolatis, racemis antè anthesin revolutis, floribus confertis, calycibus villosis, sepalis externis ciliatis viridibus, internis acuminatis incanis. ζ in Lusitaniâ ex Brot. in Corsicâ ex Salzmann. C. stæchadiifolius Brot. fl. lusit. 2. p. 270. (v. s. ex Corsicâ.)

78. H. *croceum* (Pers. ench. 2. p. 79.) caule fruticoso subprocumbente ramoso, ramis simplicibus erectis tomentoso-canescensculis, foliis tomentosisculis subtus canescensculis suprâ glaucis margine revolutis, inferioribus suborbiculatis, mediis ellipticis obtusis, superioribus lanceolatis acutiusculis, stipulis bracteisque erectis lineari-oblongis villosis subvirescentibus, calycibus glauco-flavescentibus minutè pubescentibus. ζ in Hispaniâ, Barbariâ. C. croceus Desf. fl. atl. 1. p. 422. t. 110.* Stipulæ inferiores minutæ. Petala crocea. (v. s.)

α . stipulis petiolo longioribus.

β . stipulis setaceis petiolo brevioribus.

γ . ramis procumbentibus, foliis minoribus, racemis paucifloris.

79. H. *nudicaule* (Dunal. ined.) caule fruticoso ramoso, ramis infernè glabris supernè villosis-canescensculis, foliis oblongo-lanceolatis margine revolutis utrinquè tomentosis subtus incanis suprâ viridi-flavescentibus, stipulis linearibus petiolo longioribus, calycibus profundè sulcatis vix pubescentibus, nervis elevatis pilosis. ζ in montibus regni Valentini, Hispaniæ. ex Lagasca. Petala lutea. An H. crocei varietas? (v. s.)

80. H. *glaucum* (Pers. ench. 2. p. 78.) caule suffruticoso ramoso, ramis adscendentibus tomentosis-incanis apice hispidis, foliis margine vix revolutis ciliatis utrinquè tomentosis subtus incanis suprâ viridi-glaucensculis, inferioribus orbiculatis, ceteris ellipticis lanceolato-oblongis, stipulis bracteisque pubescentibus viridibus, pedicellis calycibusque hirsutis, pilis albidis. ζ in Hispaniâ, Italiâ. C. glaucus. Cav. ic. 3. p. 31. t. 261.* nec Desf.

α . *acutiusculum*, foliis superioribus oblongis subacuminatis suprâ glaucescentibus, calycibus molliter hirsutis.

β . *obtusiusculum*, foliis superioribus oblongo-ellipticis obtusiusculis suprâ scabriusculis viridibus, calycibus subhispidis.

81. H. *tomentosum* (Dunal. ined.) caule suffruticoso ramoso, ramis elongatis adscendentibus subcanescensculis, foliis lanceolato-oblongis margine sæpè revolutis subtus incano-tomentosis suprâ glabriusculis viridibus, calycibus sulcatis, nervis elevatis pilosis. ζ in Hispaniâ, Galliâ, Angliâ. C. tomentosus Smith engl. bot. t. 2208. — Scop. carn. t. 24? Pedicelli incani piloso-tomentosi. Bractæ glabræ. Calyx hinc indè violaceus. Petala lutea. (v. s.)

82. H. *leptophyllum* (Dunal. ined.) caule suffruticoso lignoso subprocumbente ramoso, ramis adscendentibus tomentosisculis subcinereis, foliis

angustis oblongo-linearibus in petiolum brevem attenuatis margine revolutis subtus breviter tomentoso-cinereis supra glabriusculis viridibus, stipulis subulatis pilosis petiolo vix longioribus, racemis longis, calycibus piloso-hirsutis. \S in Hispaniâ. C. angustifolius Lag! in litt. nec Jacq. C. stæchadifolius hortul! Calyces sulcati. Petala lutea. Bracteæ minutæ. Racemi laxi. (v. s.)

83. *H. ACUMINATUM* (Pers. ench. 2. p. 79.) ramis erectis basi et apice pilosis medio nudis, foliis longè petiolatis oblongis margine revolutis utrinque viridibus pilosis subtus tomentosiusculis, stipulis subglabris linearibus petiolo longioribus. racemis pilosiusculis paucifloris laxis, calycibus glabris nitidis hyalinis. \S in agro Nicæensi. C. serpyllifolius Balb! ined. (v. s.) C. acuminatus Viv. fragm. 13. t. 14. f. 1. ab hoc omnino differt et videtur varietas *H. guttati* aut species affinis.

84. *H. SERPYLLIFOLIUM* (Mill. dict. n. 8.) caule suffruticoso, ramis adscendentibus basi glabris apice pilosiusculis, foliis oblongo-ellipticis margine subrevolutis subtus tomentoso-incanis, supra intensè viridibus nitidis primum pilosiusculis dein glabris, stipulis bracteisque viridibus ciliatis, calycibus canescentibus pube subinconspicuâ, nervis parçè pilosis. \S in Alpibus Austriæ, Styriæ? ex Lin.; in montibus Hispaniæ ex Lagasca. C. serpyllifolius ex Lîp. spec. 743? ut in preced. et sequent. folia inferiora minora orbiculata ovataque. (v. s.)

85. *H. VULGARE* (Gaertn. fruct. 1. p. 371. t. 76.) caule suffruticoso procumbente ramoso, ramis elongatis, foliis margine vix revolutis subtus incanocinereis, supra viridibus pilosis subciliatis, inferioribus suborbiculatis, mediis ovato-ellipticis superioribus oblongis, stipulis oblongo-linearibus ciliatis petiolo longioribus, racemis laxis, pedicellis calycibusque pilosis. \S in Europæ pascuis siccis. C. Helianthemum Linn. spec. 1. p. 744. fl. dan. t. 101. Petala lutea, basi fulva, integra.

α. ramis tomentosiusculis pubescentibus, stipulis petiolo vix longioribus. (v. s.)

β. ramis basi glabris supernè pubescentibus, stipulis petiolo 2-3-plò longioribus. (v. s.)

86. *H. SURREJANUM* (Mill. dict. n. 15.) caule suffruticoso procumbente, foliis obovato oblongis subpilosis, racemis paucifloris terminalibus, petalis angustis lanceolatis. \S in Angliæ comitatu Surrejano propè Croydon. C. Surrejanus Linn. spec. 743. — Dill. elth. 177. t. 145. f. 174.* Stipulæ longitudine petiolorum, lineari-lanceolatæ. Calyces pilosi. (v. s.)

87. *H. OVATUM* (Dunal. ined.) caule suffruticoso procumbente ramosissimo, ramis villosis, foliis ellipticis lanceolatis in petiolum attenuatis obtusiusculis utrinque villosis-sericeis ciliatis, stipulis petiolo sublongioribus villosis-ciliatis, pedunculis 1-3-floris terminalibus, calycibus subvillosis. \S in montosis inter Viterbum et Ronciglione et in Alpibus Genevensibus. C. ovatus Viv! fragm. 1. p. 6. t. 8. f. 2. Pili sericei albicantes. (v. s.)

88. *H. GRANDIFLORUM* (DC! fl. fr. 4. p. 821.) caule suffruticoso adscendente, ramis piloso-hirsutis, foliis superioribus subplanis oblongis pilosiusculis supra viridibus subtus nunc viridibus nunc dilutè cinereis, stipulis ciliatis petiolo sublongioribus, floribus magnis, calycibus subhirsutis. \S in Alpibus, Pyrenæis, Tauricis. C. grandiflorus Scop. carn. ed. 2. n. 648. t. 25.* Differt ab *H. vulgari* habitu omnium partium majori, foliis subplanis, pedunculis calycibusque pilis patentibus tectis. (v. s.)

89. *H. OBSCURUM* (Pers. ench. 2. p. 79.) caule suffruticoso adscendente ramosissimo, ramis hirsutis, foliis utrinque hirsutis subviridibus, superioribus ellipticis, stipulis ciliato-hirsutis petiolo longioribus, racemis longis, calycibus hirsutis. \S in sylvis Europæ. C. barbatus Lam. dict. 2. p. 24? *H. obscurum* α DC! fl. fr. 6. p. 624. Ut in preced. et seq. speciebus, folia inferiora subrotunda ovataque (v. s.)

90. *H. NUMMULARIUM* (Mill. dict. n. 11.) caule suffruticoso, ramis procumben-

tibus hirsutis, foliis inferioribus orbiculatis, superioribus oblongo-linearibus hirsutis subtus viridi-cinereis, stipulis lineari-oblongis petiolo duplè longioribus, racemis calycibusque hirsutis. ζ in Galliâ australi, Italiâ. C. nummularius. α Lin. spec. 743 nec Desf. et Cav. H. obscurum β . nummularium DCI fl. fr. 6. p. 624.* H. angustifolium in plur. hortis botanicis. (v. s.)

91. H. ANGUSTIFOLIUM (Pers. ench. 2. p. 79.) caule suffruticoso diffuso, ramis tomentosisculis cinereis, foliis breviter petiolatis superioribus lineari-oblongis margine revolutis acutiusculis subtus tomentoso-canescens, supra subhispidis, stipulis pilosis petiolo longioribus, racemis laxis, calycibus pubescentibus subhirsutis pilis deciduis. ζ Patria dubia, verosim. eadem ac prioris. C. angustifolius Jacq. vind. 3. p. 53. Petala basi angusta, subunguiculata, margine non imbricata. (v. s.)

92. H. OBTUSIFOLIUM (Dunal. ined.) caule suffruticoso ramoso, ramis tomentoso-incanis, foliis parvis petiolatis lineari-oblongis obtusis margine revolutis, subtus tomentoso-incani, supra viridibus pilis albidis sparsis obtectis, stipulis viridibus oblongo-linearibus planis obtusis vix ciliatis longitudine petiolorum, calycibus hispidis. ζ in ins. Cypro. C. ciliatus Cas. Rostung! in litt. nec Desf. Petala lutea? (v. s.)

93. H. HIRTUM (Pers. ench. 2. p. 79.) caule suffruticoso ramoso, ramis adscendentibus numerosis tomentoso-hirtis cinereis, foliis ovatis seu oblongis margine revolutis tomentoso hirtis subtus canescentibus supra viridi-cinereis, stipulis angustis petiolo sublongioribus, calycibus hirsutissimis albidis. ζ in Hispaniâ, Galliâ australi. C. hirtus. Lin. spec. 744. exclus. Barr. syn. Cav! ic. 2. p. 37. t. 146.* Calyces parvi.

α . Bæticum, foliis ovato-oblongis supra viridibus. H. Bæticum hort. matr. olim. (v. s.)

β . aureum, foliis lineari-oblongis margine revolutis utrinquè canescentibus. H. aureum Thib! ined. Pers. ench. 2. p. 78. (v. s.)

γ . teretifolium, ramis tomentoso-incanis, foliis subtus revolutis subteretibus crassiusculis obtusis incano-cinereis. C. teretifolius Thib! ined. H. aureum β teretifolium Pers! ench. 2. p. 78. (v. s.)

94. H. LAGASCE (Dunal. ined.) ramis adscendentibus tomentoso-hirsutis canescentibus, foliis linearibus obtusis margine valdè revolutis subteretibus subhirtis virescentibus, stipulis planis vix pilosis petiolo duplè longioribus, pedicellis hispidis albicantibus, calycibus nitidis nervoso-sulcatis, nervis ciliatis, pilis albidis. ζ in Hispaniâ. H. hirti var.? Lag. in litt. Calyces parvi. Folia brevia. (v. s.)

** Petalis albis, roseis, rubris vel dilutè sulphureis.

95. H. VIOLACEUM (Pers. ench. 2. p. 78.) caule erecto seu adscendente ramosissimo, ramis oppositis, ramulis gracilibus tomentoso-hirsutis incanis, foliis parvis sublinearibus obtusis margine revolutis utrinquè subtomentosis subtus canescentibus, stipulis minutis pilosis, racemis paucifloris laxis, calycibus levibus violaceis nervoso-sulcatis. ζ in Hispaniâ. C. violaceus Cav. ic. 2. p. 38. t. 147.* Petala alba. (v. s.)

96. H. RACEMOSUM (Dunal. ined.) caule fruticoso ramoso, ramis erectis teretibus tomentoso-incanis, foliis breviter petiolatis angustis linearibus vel linearilanceolatis margine revolutis subtus incanis, stipulis subulatis petiolo longioribus, pedicellis incanis, calycibus nervoso-sulcatis violaceo-rufescentibus. ζ in Hispaniâ, Barbariâ, Teneriffâ. C. racemosus Lin. mant. 76? Lam. dict. 2. p. 25.* Vahl symb. 1. p. 39.* Willd. spec. 2. p. 1208.* exclus. syn. Cav. et Barrel. Petala alba.

α . violascens, foliis supra viridi-glaucis, calycibus glabris violaceo-rufis. (v. s.)

β . farinosum, foliis utrinquè canescentibus pulverulentis, calycibus farinoso-pulverulentis, pilis brevissimis. H. violacei var. Lag! in litt. (v. s.)

97. H. STRICTUM (Pers. ench. 2. p. 79.) caule suffruticoso erecto ramosissimò,

ramis strictis cano-tomentosis, foliis subsessilibus angustissimis lineari-subulatis margine revolutis canescentibus, stipulis linearibus setosis, calycibus nervoso-striatis lutescentibus glabriusculis. ζ in Hispaniâ. C. strictus Cav. ic. 3. p. 32. t. 263. f. 2.* Petala alba. (v. s.)

98. H. PILOSUM (Pers. ench. 2. p. 79.) caule suffruticoso ramoso, ramis erectiusculis, foliis linearibus seu lineari-oblongis utrinquè incanis apice setosis, stipulis subulatis, calycibus pilosiusculis nervoso-striatis. ζ in Hispaniâ et Galliâ australi. C. pilosus Lin. spec. 744. α ? Chamæcistus 4. Clus. hist. 1. p. 74.* Petala alba. (v. s.)

α . foliis linearibus canis, calycibus pilosis glaucescentibus.

β . foliis lineari-oblongis, calycibus nitidis subglabris acutiusculis, nervis pilosis.

99. H. LINEARE (Pers. ench. 2. p. 78.) caule suffruticoso, ramis elongatis adscendentibus subtomentoso-incanis, foliis linearibus cano-virescentibus margine revolutis, stipulis lineari-subulatis, racemis laxis virgatis paucifloris, calycibus striatis glabriusculis, nervis subviolaceis, sepalis acutis. ζ in Hispaniâ, Galliâ australi. C. linearis Cav. ic. 3. p. 8. t. 216.* C. pilosus DC. fl. fr. 5. p. 823? Calyx major quàm in præced. spec. Petala alba. (v. s.)

β . calycum angulis pilosis.

100. H. VIRGATUM (Pers. ench. 2. p. 79.) caule suffruticoso, ramis virgatis incanis adscendentibus seu erectis, foliis linearibus subtus canescentibus margine revolutis, stipulis lineari-subulatis, calycibus cano-pulverulentis pubescentibus. ζ

α . albiflorum, foliis suprâ viridibus, petalis roseis. — In Barbariâ. C. virgatus Desf. fl. atl. 1. p. 432.*

β . roseum, foliis utrinquè canescentibus, petalis albis. — Patr. ign.

101. H. APENNINUM (DC! fl. fr. 4. p. 824.) caule suffruticoso ramoso, ramis patulis cano-tomentosiusculis, foliis petiolatis oblongo-linearibus margine vix revolutis, subtus tomentosis, suprâ glaucescentibus demùm glabris, stipulis subulatis petiolo longioribus, calycibus brevissimè villosis striatis glauco-cinereis obtusiusculis. ζ in Hispaniâ, Galliâ, Italiâ, Germaniâ, collibus apricis. C. Apenninus Lin. spec. 744? — Dill. elth. 170. C. hispidus β . Lam. dict. 2. p. 26.* Petala alba. (v. s.)

α . foliis subplanis.

β . foliis linearibus angustis. C. pilosus Thib! herb. Gouan fl. monsp. p. 265?

102. H. HISPIDUM (Dunal. ined.) caule suffruticoso ramoso, ramis adscendentibus tomentoso-canescensculis, foliis petiolatis oblongis obtusiusculis submucronatis margine revolutis, subtus incanis, suprâ subscabris viridi-glaucis, calycibus piloso-hirsutis. ζ in Galliâ australi. C. hispidus Lam. dict. 2. p. 26.* Brot. fl. lus. 2. p. 271.* C. pilosus β Gouan fl. monsp. p. 265. H. majoranæfolium β DC. fl. fr. suppl. p. 625.* Petala alba. (v. s.)

103. H. PULVERULENTUM (DC! fl. fr. 4. p. 823.) caule suffruticoso ramosissimo prostrato, ramis incano-tomentosiusculis, foliis oblongo-linearibus margine revolutis obtusis subtus incanis suprâ glaucis, stipulis subulatis petiolo longioribus, calycibus canescentibus minutè tomentoso-pubescentibus. ζ in collibus sterilibus Galliâ. C. pulverulentus Pourr. act. tonl. 3. p. 311. C. polifolius Lam. dict. 2. p. 26.* nec Lin. Petala alba. (v. s.)

104. H. RHODANTHUM (Dunal. ined.) caule suffruticoso procumbente, ramis tomentosiusculis subincanis, foliis oblongis margine revolutis, subtus tomentoso-incanis, suprâ viridi-glaucescentibus, stipulis subulatis pilosis apice setosis, calycibus breviter tomentosis albidis. ζ in Hispaniâ. C. roseus Jacq. hort. vind. 3. p. 65? C. angustifolius olim in hort. Reg. Paris. C. piluliferus Thib! ined. H. pulverulento valdè affine. (v. s.)

- α. oblongifolium*, ramis tomentoso-canescens, foliis oblongis, floribus roseis vel rubris.
- β. subhirsutum*, ramis tomentoso-canescens, foliis calycibusque subhirsutis, floribus roseis vel rubris.
- γ. carneum* (Lag! in litt.) ramis subglabris, foliis sublinearibus, pedicellis tomentoso-incanis, floribus carneis. An species diversa?
105. *H. POLIFOLIUM* (DC! fl. fr. 4. p. 823.) caule suffruticoso, ramis procumbentibus glabriusculis apice subtomentosis, foliis oblongis ovatis obtusiusculis subplanis, subtus tomentoso-incanis, supra glabris viridibus, stipulis bracteisque linearibus viridibus ciliatis, calycibus striatis glabriusculis subnitidis. *♂* in Angliâ, Galliâ, Hispaniâ. *C. polifolius* Lin. spec. 745. *C. splendens* Lam. dict. 2. p. 26.* — Dill. elth. 175. t. 145. f. 172.* Petala alba. (v. s.)
106. *H. MUTABILE* (Pers. ench. 2. p. 79.) caule suffruticoso, ramis procumbentibus tomentosiusculis, foliis planis ovato-oblongis acutiusculis supra glabris subtus levissimè tomentosis pallidè cinereis, stipulis pilosiusculis petiolo subæqualibus vel longioribus, calycibus striatis glabriusculis. *♂* in Hispaniâ. *C. mutabilis* Jacq. ic. 1. t. 99. misc. 2. p. 340.* *H. polifolio* proximum. (v. s.)
- α. floribus albis.*
- β. floribus roseo-rubris minoribus.*
- γ. floribus roseo-rubris plenius.*
107. *H. SULPHUREUM* (Willd. enum. suppl. 39.*) caulibus ramosis procumbentibus, foliis angusto-lanceolatis planis, supra viridibus, subtus pallidioribus utrinquè stellato-pubescentibus, racemo terminali paucifloro. *♂* in Hispaniâ. Petala crenulata sulphurea.
108. *H. ROSEUM* (DC! fl. fr. 4. p. 822.) caule suffruticoso subprocumbente tomentosiusculo, foliis ovato-lanceolatis utrinquè tomentosiusculis, subtus pallidè cinereis, stipulis linearibus, pedicellis calycibusque piloso subhirsutis. *♂* in Europâ australi. *C. roseus* All. fl. ped. 2. p. 105. t. 45. f. 4. nec Jacq. H. vulgari proximum, an hujus varietas? (v. s.)
109. *H. FOETIDUM* (Pers. ench. 2. p. 79.) caule suffruticoso procumbente piloso-subhirsuto, foliis oblongis utrinquè virentibus hirsutis scabriusculis, stipulis hirsutis linearibus petiolo longioribus, pedicellis calycibusque subhirsutis. *♂* Patr. ign. *C. foetidus* Jacq. ic. rar. 1. p. 98. misc. 2. p. 341.* Odor. Bryoniæ. Petala *H. vulgaris* alba. (v. s.)
110. *H. CILIATUM* (Pers. ench. 2. p. 79.) caule suffruticoso decumbente, ramis tomentoso-incanis, foliis ovato-lanceolatis seu lanceolato-oblongis margine vix revolutis subtus tomentoso-incanis supra hirsutis, stipulis subviridibus petiolo longioribus, calycibus membranaceis, sepalis internis nervoso-sulcatis, nervis elevatis glanduloso-pilosis. *♂* in Hispaniâ, Africâ bor., Italiâ. *C. ciliatus* Desf. fl. atl. 1. p. 421. t. 109.* Petala rosea. (v. s.)
111. *H. ASPERUM* (Lag! ined.) caule suffruticoso ramoso, ramis longis adscendentibus erectisque subtomentoso-scabriusculis cinereis, foliis petiolatis oblongo-acuminatis margine revolutis subtus tomentoso-incanis supra viridibus subtomentosis scabriusculis apice setosis, stipulis subulatis setosis, calycem angulis hirsuto-hispidis, pilis longis numerosis. *♂* in Hispaniâ. *C. hirtus* herb. Thib! Petala alba. (v. s.)
- β. Roussæi*, caule foliis calycibusque hirsutissimis, pilis albidis. *♂* in Oriente. Rousseau. (v. s.)
112. *H. MAJORANÆFOLIUM* (DC! fl. fr. 6. p. 625. var. *α.*) suffruticoso erecto ramosissimo, ramis hirsuto-tomentosis canescentibus, foliis petiolatis ovato-oblongis acutiusculis margine revolutis subtus incano-tomentosis supra viridiglaucescentibus tomentoso-hirtis, stipulis subulatis setosis, calycibus hirsu-

tissimis, pilis albidis. ♂ in Galliâ australi. C. majoranæfolius Gouan herb. 26? (v. s.)

†. *Species non satis notæ.*

113. *H. ASTYLUM* (Moq. et Sessé fl. mex. ic. ined.) diffusum subherbaceum humile, foliis subalternis stipulatis ovali-oblongis, sepalis quinque; 2 exterioribus linearibus minimis, stylo nullo, stigmatibus subtrilobis. — In Novâ Hispaniâ.
114. *H. TRIPETALUM* (Moq. et Sessé fl. mex. ic. ined.) erectum multicaule gracile, foliis alternis linearibus exstipulatis, sepalis quinque, exterioribus 2 linearibus parvis, petalis 3. — In Mexico.
115. *H. OBOECORDATUM* (Moq. et Sessé fl. mex. ic. ined.) erectum suffruticosum, foliis alternis oblongis stipulatis subfasciculatis, sepalis 3, petalibus 5 oboecordatis. ♂ in Mexico.
116. *H. OLIGOPHYLLUM* (Clark. in Spreng. neu. entd. 3. p. 163. sub Cisto.) fruticosum stipulatum, foliis petiolatis ovato-lanceolatis nerviis integerrimis scabris margine revolutis, pedunculis unifloris. ♂ propè Jaffa. Petala flava.
117. *H. DISTACHIUM* (Roth in ann. bot. 2. p. 34. sub Cisto.) suffruticosum stipulatum erectum, foliis oppositis ovali-oblongis incanis, racemis terminalibus foliosis bipartitis, floribus oppositifoliis. ♂ in Lusitaniâ.
118. *H. HYSSOPIFOLIUM* (Ten. cat. h. neap. 1819. p. 48.) suffruticosum stipulatum, foliis inferioribus ovalibus, superioribus lanceolato-linearibus, utrinque viridibus planis hirsutis, calycibus hirsutis capsulâ duplò longioribus. ♂ in collibus Lucaniæ. Medium, ex cl. auct., inter *H. obscurum* et *grandiflorum*.
119. *H. LUCIDUM* (Horn. cat. h. hafn. 498.) suffruticosum procumbens stipulatum, foliis ovalibus lucidis margine revolutis. ♂ patr. ign. Affine, ex cl. auct. *H. vulgari*.
120. *H. HIRSUTUM*, suffruticosum stipulatum hirsutum, foliis petiolatis subtus canis, imis rotundatis, superioribus lanceolato-acutis, floribus terminalibus racemosis subsecundis. ♂ in Pyrenæorum orientalium rupibus. *Cistus hirsutus* Lapeyr. abr. 303. non Lam. Flores albi magni.
121. *H. PILOSELLOIDES*, suffruticosus exstipulatus, foliis elongato-ellipticis obtusis longè petiolatis suprà viridibus infrà tomentoso-incanis infrà supràque longioribus pilis hirsutis, floribus paniculatis. ♂ in Pyrenæorum rupibus apricis. *Cistus piloselloides* Lapeyr. abr. 301. Flores flavi.
122. *H. FUGAX*, caule herbaceo, foliis subovatis pilosis, flore fugaci. ① in monte Baldo. *H. fagacium*. Mill. dict. n. 19. An idem ac *H. guttatum*?
123. *H. CISTIFOLIUM* (Mill. dict. n. 9.) caulibus procumbentibus suffruticosis glabris, foliis ovato-lanceolatis oppositis, pedunculis longioribus. ♂ in Germaniâ. Flores lutei.
124. *H. FASCICULATUM* (Mill. dict. n. 22.) foliis angustis fasciculatis, pedicellis elongatis lateralibus et terminalibus. ②? ad cap. B.-Spei. Flores fugacissimi pallide straminei.

III. HUDSONIA. *Lin. mant.* 11. *Nutt. gen. amer.* 2. p. 4.

Petala 5? Stamina 15-30; filamenta filiformia; antheræ parvulæ, longitudinaliter dehiscentes. Stylus rectus erectus? simplex staminibus æqualis. Stigma simplex. Capsula 1-locularis, 3-valvis, 1-3 spermata, oblonga vel obovata, coriacea, lævis vel pubescens. Semina granulata. Embryo in albumine corneo immersus. — Suffrutices parvi, sæpius

cæspitosi. Folia alterna, parvula, subulata, acerosa, imbricata, exstipulata. Flores subsessiles vel pedunculati; pedunculi 1-flori, solitarii terminalesque vel lateraliter aggregati.

1. *H. ERICOIDES* (Lin. mant. 74.) pubescens, caule suffruticoso suberecto, ramis elongatis, foliis filiformibus subulatis subimbricatis, pedunculis lateraliter à gemmis foliaceis solitariis, calycibus cylindricis obtusis, capsulis pubescentibus semper monospermis, valvulis oblongis. ♂ in pinetis Novæ-Cæsareæ et Virginie. Willd. hort. ber. t. 15. Folia persistentia. Stamina circiter 15. Pedunculi 5-8 lin. longi. (v. s.) — An stirps Nuttaliana floribus aggregatis ex descriptione donata, omnino eadem?
2. *H. MONTANA* (Nutt. gen. 2. p. 5.) sublævis, caulibus cæspitosis decumbentibus, foliis longis subulato-filiformibus subimbricatis, pedunculis terminalibus, calycibus campanulatis lanuginosis, partitionibus acuminatis subulatis, capsulis villosis sæpius 3-spermis, valvulis ovatis. ♂ in borealis Carolinæ montium summitatibus. Stamina 20-25. Semina subangulata.
3. *H. TOMENTOSA* (Nutt. gen. 2. p. 5.) cæspitosa et incano-tomentosa, caulibus intricatis densis, foliis minutis dense imbricatis ovato-acutis, floribus aggregatis subsessilibus, calycibus subcylindricis partitionibus obtusis, capsulis monospermis, valvis ovatis lævibus. ♂ in sabulosis maritimis New-Jersey, Delaware, Marylandiæ, etc. Stamina 14-18.

IV. LECHEA. *Lin. gen. n. 142. Gaertn. fruct. 2. t. 129.*

Calyx 3-sepalus bracteis sepalisve exter. duobus exterioribus stipatus. Petala 3, lanceolata. Stamina 3-12, sæpius numero ternaria. Ovarium 1 subtrigonum. Stigmata 3 vix distincta. Capsula 3-valvis, valvis medio septi aut nerviferis; semina septo nervove adfixa paucissima sæpius 8; albumen carnosum. Embryo dorsalis rectiusculus, radicula infera, cotyledonibus ovato-oblongis. — Herbæ boreali-Americanæ incomptæ, floribus numerosis parvis, ramis infimis à floriferis sæpè diversis.

1. *L. VILLOSA* (Ell. sketch. 184.) undiquè hirsuta, foliis oblongo-lanceolatis mucronatis, paniculâ foliosâ pyramidatâ, ramis apice floriferis, floribus fasciculato-racemosis secundis brevissimè pedicellatis. ♀ à Canadâ ad Floridam. *L. major* Pursh fl. bor. am. 1. p. 90. non Lin. — *L. minor* Lin. ex Smith. Lam. ill. t. 52. f. 2. pessimè. (v. s.)

♂? *mucronata* (Raf. prec. 37.) pilosa, caule recto simplici, racemo composito, floribus bracteolatis. ♀ in sylvis Novæ-Cæsareæ.

2. *L. MINOR* (Pursh. fl. bor. am. 1. p. 91.) glabriuscula, foliis lineari-lanceolatis acutis, paniculâ foliosâ, ramis elongatis undiquè floriferis, floribus breviter pedicellatis, caule assurgente. ♀ in collibus à Canadâ ad Pensylvaniam. — Lam. ill. t. 52. f. 1. ex Pursh.
3. *L. RACEMULOSA* (Michx. fl. bor. am. 1. p. 77.) undiquè adpresso-pubescens, foliis linearibus acutis ciliatis, paniculâ gracili ramosissimâ pyramidatâ, racemis nudiusculis, floribus parvis alternis pedicellatis, caule erecto. ♀ in campis arenosis à Novâ-Cæsareâ ad Carolinam. Pursh 1. p. 91. Poir. suppl. 3. p. 340. — *Gaura* Lam. ill. t. 281. f. 3. opt. (v. s.)
4. *L. THYMIFOLIA* (Michx. fl. bor. am. 1. p. 77.) undiquè adpresso-cano-villosa, foliis linearibus acutis, paniculâ foliosâ elongatâ, ramis brevibus, floribus fasciculatis lateralibus terminalibusque, pedicellis brevissimis, floribus parvis cano-tomentosis, caule erecto. ♀ in siccis sylvis à Virginiâ ad Carolinam. Rami inferiores repentes *Thymo Serpyllo* similes Pursh. fl. bor. amer. 1. p. 91. *L. minor* Walt. car. 83 ex Ell. sketch. p. 185.

5. *L. tenuifolia* (Michx. fl. bor. am. r. p. 77.) sparsè pilosa, foliis angustissimis, paniculâ divaricatâ nudiusculâ, ramulis alternis, pedicellis elongatis divaricatis, caule erecto. \mathcal{U} in collibus sabulosis à Virginiâ ad Georgiam. *L. juncifolia* Walt. car. 83^p ex Ell. sketch. 185. Rami inferi foliis linearibus donati. Pursh l. c. p. 91.

6? *L. verticillata* (Willd. spec. r. p. 495.) caule hispido, foliis oblongo-ovatis serrulatis, floribus verticillatis. \mathcal{U} in Indiâ orientali Habitus *Spermaceæ*. — Species à genere imò ex ordine verosimiliter removenda.

N. B. L. Chinensis Lour. videtur *Commelinæ* quædam species ex DC.

(DUNAL.)

Ordo XVI. VIOLARIEÆ. DC. fl. fr. 4. p. 801. Juss. ann. mus.
18. p. 476. (1)

Sepala 5, persistentia, æqualia aut inæqualia, margine sæpius membranacea, libera aut basi inter se subcoalita, sæpius deorsum producta, æstivatione imbricata et ideò duplici serie disposita : nempè (in Violeis) 3 exteriora et 2 interiora, interdum (in Sauvageis) 2 exteriora et 3 interiora. Petala 5 cum sepalis alterna, hypogyna, thalamo inserta, sæpè marcescentia, æstivatione (sæpius) obliquè convolutivâ; nunc inter se æqualia, nunc inæqualia et tunc inferius ab aliis discrepans labelliforme aut basi calcaratum cucullatumve. Interdum inter petala et stamina urceolus staminifer vel appendices filiformes. Stamina 5 thalamo inserta, cum petalis (in Violeis) alterna, aut rarissimè (in Sauvageis) petalis opposita. Antheræ biloculares, introrsum rimâ longitudinali duplici dehiscentes, ovario adpressæ, sæpius liberæ, interdum basi plus minusve in discum monadelphum solutæ. Filamenta sæpissimè dilatata, nunc à basi in ligulam infernè latiore, antheras ad basin dimissiùs gerentem, nunc infernè unguiculata, hinc in ligulam dilatata, antheras ad basin hujus ligulæ et ideò altiùs gerentia, in utroque casu apice ultrà antheras in membranam aridam producta, stylum plus minusve imbricatum cingentem, rarissimè (in Sauvageis) subulata antheras non superantia; 2 in floribus irregularibus sæpius deorsum in appendicem seu glandulam nectaream (Nectaria Schk.) producta. Ovarium uniloculare polyspermum aut rarè abortu (?) monospermum. Placentæ 3 parietales, sepalis 3 exterioribus suprâ thalamum symetricè dispositis oppositæ. Stylus 1 persistens, sæpius declinatus perforatusque, canali pistillari sæpius apice recurvo et ideò stigma paulum laterale. Capsula 3-valvis, valvis sæpissimè ab apice usquè ad basin dehiscentibus, sæpius elasticis. Seminis integumentum 3-plex; exterius (Epidermis Gærtn. DC.) membranaceum tenuissimum adhærens, versùs hilum plus minus in carunculam (Arillum incompletum Gærtn.) incrassatus; vascula umbilicalia cum epidermide conferruminata, ab hilo ad verticem seminis producta, lineam vix pro-

(1) Hanc ordinis monographiam partim ex herbario et manuscriptis nostris ad syst. univ. paratis, partim et præcipuè ex observationibus propriis elaboratam nobiscum communicavit cl. DE GINGINS qui jam in Mem. soc. h. nat. gen. 2. p. 1. Violaricarum characteres sagacissimè enucleavit.

minentem (Raphe Gærtn.) formantia, in areolam rugosam (umbilicam intimum?) expansa, intrā testam penetrantia. Testa crustacea, fragilis, sæpius lævis, interdum striata (?) rarius scrobiculata; membrana interna (Endoplevra DC.) tenuissima adhærens, in vertice (Chalaza Gærtn.) orbiculari sæpè rufo-notata. Albumen carnosum, sæpius læve. Embryo rectus, in axi albuminis, ejusque ferè longitudine; radicula geometricam seminis basin (non hilum) spectante; plumula inconspicua; cotyledones sæpissimè planiusculæ.

Herbæ, suffrutices aut frutices mediocres. Folia sæpius alterna, rarius opposita, stipulacea, simplicia, sæpius per vernationem involutiva. Flores erecti aut sæpè cernui, pedunculati aut ramosi, axillares; pedunculi nunc solitarii aut plurimi, uniflori et bibracteolati, nunc ramosi et tunc pedicelli uniflori bibracteolati ex axillâ bracteæ solitariæ.

Violariæ sunt valdè affines Polygaleis, Droseraceis et imprimis Passifloreis. A Polygaleis differunt fructû uniloculari nec biloculari, foliis stipulaceis nec exstipulatis, antheris bilocularibus, nec unilocularibus. A Droseraceis stylo solitario, embryone elongato et foliis per vernationem involutivis nec circinalibus, stipulatis nec exstipulatis. A Passifloreis tandem fructû capsulari, nec verè baccato; albumine nitido, nec foveolato; staminibus hypogynis nec perigynis; antheris medifixis, nec adnatis; stigmatibus 3, nec 1. Genus Calyptrion affine Passifloreis ob caules volubiles; genus Hymenanthera Polygaleis ob pericarpium monospermum seminibus solitariis pendulis aut (secundum R. Brown Cong. p. 21) 2-loculare, loculis monospermis.

Trib. I. *VIOLÆ*. DC. mss.

Petala inter se inæqualia; sepala 3 exteriora, sæpius latiora, et 2 interiora. Pericarpia dehiscentiâ loculicidâ, nempè suturæ valvarum cum placentâ alternantes et ideò valvæ medio placenteræ dictæ. Stamina petalis alterna; filamenta dilatata, ultrâ antheras producta, libera (approximata aut coarctata) vel rarò coalita; loculi antherarum tardiùs bivalves.

I. CALYPTRION. Ging. mss. — Corynostylis. Martius ined.

Calyx minimus, basi in pedicellum decurrens; sepala subæqualia, 2 inferiora subinæquilatera; petalum inferius maximum posticè productum in tubum cuculliformem, lateribus compressum, medio coarctatum, contortum, multinervium, limbo æstivatione involuto, seriùs patulo, sæpius undulato, eroso vel lobato. Petala 2 lateralia vix breviora, patula; reliqua minima brevissimaque sursùm flexa, omnin

sæpius villosa. Stamina libera, coarctata; filamenta è basi dilatata oblonga, antheras demissius gerentia; antheræ ovario duplò longiores; lobis parallelis; 4? anteriores extus productæ in caudas (nectaria) totidem, (è quibusque 2 brevissimæ aut abortivæ), barbata, sursum recurvas, intrà cucullum intrantes. Capsula sæpius 3-gona? 1-3-sperma? Semina (in *C. Aubletii*) maxima subrotunda, abortu solitaria? — Arbusculæ vel frutices scandentes sæpius dextrorsum volubiles, inermes, cortice cinereo, striato, sæpius albo punctato vestitæ. Folia alterna, firma, reticulato-venosa. Flores sæpius magni, racemosi, racemi axillares aut subterminales, interdum in axillis fasciculato-congesti; pedicelli uniflori, basi bracteati, versùs medium bibracteolati, hinc articulati.

Obs. sub *V. Hybantho* Willd. quatuor stirpes inter se valdè diversæ et verosimiliter ad genera tria distincta pertinentes conjunguntur. (conf. May. prim. fl. essequéb. p. 123.)

§. 1. *Floribus maximis, racemosis aut in ramulis axillaribus solitariis.*

1. *C. AUBLETTI* (Ging. mss.) caule striato albo-punctato, foliis elliptico-oblongis acuminatis basi rotundatis serratis, sepalis ovatis obtusis, cucullo fauce vix inflato. § in Guyanâ, Cayennâ, insul. Trinitatis, Brasiliâ. *Viola Hybanthus* Aubl. guyan. 2. p. 811.* t. 319. May. fl. essequéb. p. 123.* non Willd. nec Pers. *Jonidium Aubletii* Roem. et Schult. syst. 5. p. 397. *Viola laurifolia* Sm. in Rees cyclop. *Corynostilis Hybanthus* Martius! ined. Semina maxima subrotunda? castanea. (v. s. s.)

2. *C. BERTERII* (Ging. mss.) caule striato albo punctato, foliis ovatis acuminatis integris, sepalis ovatis acutis, cucullo fauce inflato. § in Amer. æquinoctiali.

a. *Magdalenense*, racemis axillaribus. — In Baranquillâ ad ripas fl. Magdalenæ, locis inundatis. *Viola scandens* Bert! ined. (v. s. s.)

ß. *Mexicanum*, racemis (ad apicem ramulorum) terminalibus, flore ampliore? in Mexico Moc. et Sess. *Viola cucullata* fl. mexic. ic. ined. An species dist.? (v. s. comm. à cl. Lamb.)

3. ? *C. CITRIFOLIUM* (Ging. mss.) caule, foliis oblongis obtusis integerrimis, sepalis ovatis acutis, floribus solitariis, capsulis obsolete 3-gonis, seminibus oblongis compressis. § in Cumanâ. *Viola arborescens* Lœfl. itin. p. 282.

4. *C. ? ORINOCENSE* (H. B. et Kunth nov. gen. am. 5. p. 384.) caule anguloso puberulo, foliis ovato-oblongis acutis, basi obtusis serrulatis, suprâ glabris subtus pubescentibus, floribus solitariis?. § in locis humidis propè St. Fernando de Atabapo, Miss. del. Orinoco. *Simillimum C. Aubletii* ex auct. *Viola excelsa* Willd. herb. ex Roem. et Schult. syst. 5. p. 391.

§. 2. *Floribus minimis, in axillis foliorum fasciculato-congestis.*

5. *C. ? FRANGULÆFOLIUM* (Ging. mss.) caule dextrorsum scandente, foliis oblongo-ellipticis acutis basi rotundatis obsolete serrulatis utrinquè puberulis, floribus fasciculato-congestis petiolos superantibus, sepalis angustis acuminatis, petalo inferius apice emarginato, calcare (cucullo) obtuso lateribus compresso, staminibus (omnibus effætis) appendiculatis, appendicibus glabris longissimis. § in Andibus Popayanensibus, alt. 1004 hex. *Noisetia frangulæfolia* H. B. et Kunth nov. gen. am. 5. p. 384.* t. 499. a et b fig. 1. Flos ferè Noisetiæ longifoliæ, sed habitus ferè *Calyptrionis*. An inde rectius huc collocanda?

†. *Species dubia.*

6. *C. ? DIANDRUM* (Ging. mss.) caule herbaceo serpente, foliis oblongis, pedun-

culis solitariis, petalo infimo trifido *Viola diandra* Allamand in
Lin. syst. 669.* *Jonidium diandrum* Rœm. et Schult. syst. 5. p. 399.* †.

II. NOISETTIA. *H. B. et Kunth. nov. gen. am. 5. p. 382.* exclus. syn. Aubl. —*
Bigelovia. DC. mss. non Spreng.

Calyx mediocris basi in pedicellum decurrens, sepalis inæqualibus, 2 inferioribus sæpius inæquilateris; petalum inferius magnum, limbo æstivatione involuto, posticè in tubum cylindraceum, rectum longè productum, 2 lateralia breviora adscendentia, reliqua minora, omnia glabra?. Stamina libera; filamenta basi dilatata oblonga, antheras demissius gerentia; antheræ ovario vix longiores, lobis subsagittatis, 2 anteriores dorso productæ in appendices subulatas nectaríferas intrâ tubum intrantes. Ovarium superum; ovula crebra (12-30? subpyriformia (*H. B. et K. l. c.*) Capsula ovata, obsolete trigona, venosa, placentis linearibus, seminibus paucis. — Frutices erecti; folia alternata, simplicia, penninervia; stipulæ geminæ, sæpius in caule decurrentes; flores pedicellati; pedicelli plurimi aut (abortu?) solitarii, in axillis fasciculato-congesti, versùs medium hibracteolati, inarticulati.

1. *N. LONGIFOLIA* (*H. B. et K. l. c. p. 382. t. 499. b. f. II.*) caule fruticoso striato, foliis lanceolatis serratis acutis, basi in petiolum breviusculum attenuatis, calcare subulato longitudine pedicellorum, sepalis lanceolatis acuminatis, 2 inferioribus inæquilateris extùs basi auriculatis, petalo infero obcordato, lateralibus oblongis obtusis, ceteris ovatis, stigmatè uncinato. ‡ in *Guyenâ. Viola longifolia* Poir. dict. 8. p. 649.* *Jonidium longifolium* Rœm. et Schult. syst. 5. p. 398.

2. *N. ORCHIDIFLORA* (*Ging. mss.*) caule gracili lævi, foliis lanceolatis serrulatis longè petiolatis, floribus 4-6 fasciculatis, calcare obtuso pedicello paulò breviorè. ‡? in *Guyanâ Viola orchidiflora* Rudge plant. guyan. rar. 1. p. 111. t. 10. *Jonidium orchidiflorum* Rœm. et Schult. syst. 5. p. 400.

ℓ. foliis lanceolatis acuminatis serratis basi in petiolum longiusculum attenuatis glabris, floribus 3-4 lineas longis 3-4 fasciculatis, capsulis 3-gonis stramineis, valvis ovatis venosis glabris 2-3-spermis. ‡ in *Guyanâ. (v. s. s.)*

3. *N. ACUMINATA* (*DC. mss.*) foliis lanceolatis breviter petiolatis ciliatis hinc indè dentatis, apice in acumen integrum longè productis, pedicellis solitariis folio paulò brevioribus, capsulis ovatis. — In *Amer. bor. Violæoides Michx?* ined. An prioris var? (*DC. v. s. in h. Mus. Par.*)

III. GLOSSARRHEN. *Martius ined.*

Calycis sejala valdè inæqualia, basi in pedunculum decurrentia; 3 exteriora maxima, petalis majora sæpius cordata acuminata basi hastatim biauriculata; inferiora calcare interposito inæquilatera, auriculis exterioribus porrectis sæpius rotundatis; 2 interiora minima angustissima. Petala inæqualia, æstivatione. , unguibus 3-nerviis, 2 superiora breviora, 2 lateralia superiore longiora, infimum omnium magnum, limbo æstivatione involutivâ, posticè in calcar cavum productum. Stamina filamenta basi discreta, hinc dilatata, oblonga, ovario adpressa, antheras demissius gerentia; lobis antherarum apice productis? divergentibus, basi approximatis; filam. 2 anteriora dorso appendices subulatas nectaríferas in calcar intrantes gerentia. Stigma sæpius deorsum in appendicem subspathulatam productum adscendens, apice subinflexum. — Arbusculæ? cortice sæpius rubente; folia al-

terna, decidua? penninervia, stipulis minimis; flores ad apicem ramulorum? axillares; pedunculi solitarii, uniflori, bibracteati, articulati.

Genus medium inter Noisettiam et Violam; ab utrâque differt ob calycis formam.

1. *G. FLORIBUNDUS* (Martius! in litt.) foliis lanceolatis acuminatis. \S in Brasiliâ. (v. s.)

2. *G. PARVIFLORUS* (Mart! in litt.) foliis obovatis obtusis. \S in Brasiliâ. (v. s.)

IV. VIOLA. *Tourn. inst.* 419. t. 236. — *Violæ* sp. *Lin.*

Calycis sepala inæqualia, omnia plus minus deorsum in appendices auriculæformes (è dilatationibus nervorum prodeuntes) producta, post anthesin erecta. Petala inæqualia, æstivatione convolutivâ, ungue trinervi; inferius deorsum in calcar cavum plus minus productum. Stamina approximata aut coarctata (nec coalita) ad apicem dentium tori pentagoni 5-dentati inserta. Filamenta basi dilatata oblonga vel triangularia, antheras demissius gerentia; lobi antherarum basi divergentes; stamina 2 anteriora dorso appendices varias nectariferas in calcar intrantes gerentia. Ovarium nunc superum, nunc basi toro concavo cinctum et ideò semi-inferum. Valvulæ capsulæ elasticæ post maturationem contractæ semina ejicientes; semina horizontalia manifestè carunculata, plus minus ovoidea et nitida. Embryo oblongus, radícula teretiuscula, cotyledones sæpius oblongo-orbiculatæ planiusculæ radiculâ vix longiores. — Herbæ plerumque perennes, rarissimè annuæ, nunc caule brevissimo vel subterraneo, unde *acaules* dictæ, nunc caulescentes vel suffrutices parvi; folia alterna marcescentia; pedunculi solitarii axillares uniflori, 2-bracteolati non articulati, apice reflexi; flores cernui; folia seminalia oblonga ovatave, petiolata; primordialia opposita, sed rarò coëtanea.

Sect. I. NOMIMIUM. *Ging. mss.*

Stigma rostratum, foraminulo ad extremitatem rostri sito; rostrum plus minus recurvum, nunc deorsum marginatum et tunc planiusculum et obliquum, nunc absque margine et tunc deorsum convexiusculum. Stylus ab apice ad basin attenuatus. Stamina oblonga approximata. Torus planiusculus. Capsula sæpius trigona. Semina 15-27. Folia seminalia sæpè obovata vel oblonga.

\S . 1. *Stigmatibus apice depressis, aut deorsum in orbem marginatis; caulibus subnullis.*

*. *Truncis radiciformibus (Rhizomatibus?) dentatis plus minusve carnosis, sæpius obliquis.*

1. *V. PEDATA* (Lin. sp. 1323.) stigmatè incrassato hinc obliquè truncato, rostro brevissimo, foliis pellucido-punctatis palmatisectis, stipulis pectinatis laceris longè adhærentibus, petalis omnibus glabris. \mathcal{V} in Amer. boreal.

a. *lineariloba*, foliis pedatim 5-7-partitis, partitionibus multifidis, lobis lineari-lanceolatis integris ciliatis. *Curt. bot. mag.* t. 89. *V. digitata* Pursh. fl. am. p. 171. ex *Lewis am. journ.* 5. n. 1. p. 48. (v. s. sp.)

β . *atropurpurea*, foliis 5-7-partitis, partitionibus cuneatis incisissimis, pistillo pubescente. *V. atropurpurea* Rafin. in litt. *V. pedata* var. *bicolor* Pursh sec. Rafin. (v. s. sp. et c.)

γ . *ranunculifolia* (Ging. mss.), foliis inciso-lobatis. *V. ranunculifolia* Poir! dict. 8. p. 626? (v. s.)

2. *V. PALMATA* (Lin. sp. 1323.) pubescens, stigmatè triangulârè marginato,

rhizomate carnoso crasso, foliis hastato-cordatis palmato-lobatis, lobis polymorphis (raro indivisis), seminibus turgido-ovoideis cinereo-punctatis, petalo infero imberbi, omnium unguibus carinatis. $\frac{1}{2}$ in Amer. boreali.

a. vulgaris (Elliot sket. 1. p. 300.*) foliorum lobis extimis ad basin segmentis parvis acutis. — Communis in solo lævi. (v. s. sp. et c.)

β. fragrans (Elliot l. c.*) foliis magis dissectis, floribus fragrantibus. — prope Savannah.

γ. dilatata (Elliot l. c.) tota pubescens, foliis profundè dissectis *Vulgaris* in Carolinâ et Georgiâ super. Hæc est typus speciei ex Lewis. (v. s. sp.)

δ. heterophylla (Elliot l. c.), foliis vernis cordatis, ceteris hastatis divisis rugosis, (esculentis et mucilagineis). — secus rivulos uliginosos prope Ogeacher. An spec.?

ε. triloba (Ging. mss.) foliis biformibus, alteris cordato-reniformibus, alteris trilobis, lobis lateralibus minoribus subdivisis glabris, nectariis minoribus (ex Lewis l. c.) — in sylvis fertilibus et pratis Carolinæ septentrion. (Lewis) *V. triloba* Lewis amer. journ. 5. n. 1. sp. 5.* An species? (v. s. sp.)

3. *V. ASARIFOLIA* (Pursh fl. bor. am. 2. p. 732. non Muhl.) stigmate globoso immarginato?, rhizomate crassissimo corallino, foliis reniformi-cordatis indivisis raro lobatis pubescentibus, petalo infimo barbato. $\frac{1}{2}$ in oleraceis et pratis Carolinæ septent. Ambigit inter *V. palmatam* et *cucullatam*, sed ab utrâque differt stigmate non triangulari marginato nec depresso et petalo infimo barbato.

4. *V. PAPILIONACEA* (Pursh fl. bor. am. 1. p. 173.) stigmate triangulari marginato? rhizomate carnoso crassiusculo, foliis cordato-reniformibus serratis glabris, petalo infimo ad cavitatem pistillarem barbato. $\frac{1}{2}$ in humidis Carolinæ sept. Rœm. et Schult. syst. 5. p. 356. *V. barbata* Willd. mss? *V. cucullata* Lewis l. c. spec. 7.* non Ait.

a. sepalis acuminatis, foliis triangulari-cordatis acuminatis. (v. s. s.)

β. sepalis ovatis, foliis cordatis subacutis. (v. s. s.)

5. *V. CUCULLATA* (Elliot sket. 1. p. 298.* non Lewis.) stigmate triangulari marginato, rhizomate carnoso crasso, foliis cordatis acutis serratis glabriusculis, petali infimi limbo angusto imberbi, unguibus omnium carinatis. $\frac{1}{2}$ in pratis humidis Americæ borealis. (Elliot.)

a. obliqua, foliis glabriusculis acuminatis, sepalis lanceolato-acuminatis, seminibus — in Carolinâ sept. Deile; in pratis udis Lewis. — *V. obliqua* Lewis am. journ. sc. et art. 5. n. 1.* Pursh fl. am. sept. 1. p. 172. *V. cordata* Walt. carol. p. 219? (v. s. sp.)

β. glaberrima, foliis glaberrimis, serioribus acuminatis, sepalis lanceolatis, seminibus oblongo-ovoideis canescentibus rufo-punctatis. — in hortis Europæis culta. *V. cucullata* Ait. kew. 3. p. 288. Pursh fl. am. sept. 1. p. 173.* Sims bot. mag. t. 1795. *V. obliqua* Pio diss. p. 12. t. 3. f. 1. In var. β cultâ, folia verna cordata, dein amplifiata et acuminata, seriùs maximè dilatata et reniformia Flos *V. palmatæ* L. cultæ simillimus etiam quoad organa generationis, sed differt præcipuè ex caracteribus seminum. A *V. papilionaceâ* Pursh differt imprimis petalo infimo imberbi. (v. v. c.)

γ. hispidula, foliis supernè hispidulis breviter acuminatis. (v. s. sp.)

δ. cordiformis, foliis cordatis vix acuminatis supernè hispidulis, rhizomate perpendiculis nigricante. (v. s. sp.)

*** *Radixibus rugosis plus minus lignosis et ramosis.*

6. *V. PINNATA* (Lin. sp. 1323.) stigmate triangulari marginato, radice divisâ sublignosâ, foliis pinnatis multipartitis, partitionibus multilobatis, sepalis ovatis, seminibus turbinatis basi foveolatis rubicundis $\frac{1}{2}$.

- α. Sibirica*, foliis pinnatipartitis, partitionibus pinnatifidis dense ciliatis, capsulis (immaturis?) acutioribus, seminibus minoribus. — in Sibiria. Gmel. sibir. 4. p. 101. t. 48. f. 4. (v. s. sp.)
- β. Europæa*, foliis pinnatifidis, fissuris multilobatis remotè ciliatis, capsulis obtusis, seminibus maximis. — in Alpibus Europæ. All! miscell. taur. 3. p. 181. t. 5. f. 2. Hall. hist. n. 561. (v. v.)
7. *V. DACTYLOIDES* (Roem. et Schult. syst. 5. p. 349.) stigmate marginato, radice divisâ duriusculâ, foliis palmatim 5-7-sectis, foliolis oblongo-lanceolatis grosse serratis subtus villosis, sepalis ovatis, seminibus. . . . ♀ in betuletis circa Ircutiam et Nertschinsk-Sawod. V. palmata Patrini! herb. — Gmel. fl. sibir. 4. p. 100. t. 49. f. 3. (v. s. sp.)
8. *V. VARIEGATA* (Fisch! in litt.) stylo marginato, radice subdivisâ duriusculâ, foliis cordato-ovatis subrotundisve, capsulis obtusis, seminibus ovoideis rubicundis. ♀ in Dahuriâ. V. maculata Patrini! herb. non Cav. Stipulæ lanceolatæ adhærentes denticulatæ; folia subtus violacea, superne obscurè viridia ad venas albida hispidula; calcar cylindraceum rectum sepalorum longitudine. In plantâ fructiferâ folia ampliata rotundata subglabra. Flos V. pinnatæ proximus. Capsulæ obovato-trigonæ. (v. s.)
9. *V. CALTHÆFOLIA* (Poir. dict. 8. p. 627.) foliis ovato-cordatis obtusis subcrenatis subhirsutis, pedunculis folio brevioribus. — Patr. ign. An eadem ac V. asarifolia? Capsula ovato-triquetra. Semina globosa. †
10. *V. PRIMULÆFOLIA* (Lin. sp. 1324. excl. syn. Gmel.) stigmate marginato, radice fibrosâ subramosâ, foliis ovatis basi subcordatis, petiolis marginatis, stipulis liberis, sepalis lanceolatis subobtusis, petalis 2 barbatis infimo acuminato. ♀ in humidis Carolinæ et New-Jersey. Flores odorati ex Lewis
- α. cordata*, glabriuscula, petiolis folio brevioribus, scapis folio duplò longioribus. — in Carolinâ sept. V. primulifolia Ait. h. Kew. ed. 2. v. 2. p. 45? Lewis am. journ. 5. n. 1. p. 48.* V. cordata Walt. ? ex Bosc. (v. s. s.)
- β. Boscii*, glabriuscula, petiolis subnullis, scapis folia vix superantibus. — in Carolinâ. Bosc. Planta 6-pollicaris (v. s. s.)
- γ? Nuttallii*, pilosa, petiolis foliorum longitudine. — circa Nov.-Eboracum Torrey! V. primulifolia Nutt. gen. 1. p. 149.* (v. s. sp.)
- δ? ovata*, glabriuscula, petiolis foliorum longitudine, capsulis oblongo-3-gonis, valvis apice subacuminatis — in Pennsylvania. V. ovata Rafini! in litt. (v. s. sp.)
11. *V. LANCEOLATA* (Forst. in tr. lin. soc. 6. p. 310.) stolonifera, stigmate marginato, foliis lanceolatis utrinquè attenuatis, stipulis liberis, sepalis lanceolatis, petalis omnibus glabris? ex Nutt. 2 barbatis ex Elliot. ♀ in humidis Americæ boreal. V. lanceolata Lin! spec. excl. var. sib. Lodd. cab. t. 211. (v. s. sp. et c.)
- β. Foliis ovatis basi truncatis.* — in Georgiâ et Carolinâ Beauv. An var. V. primulifoliæ. (v. s. sp.)
12. *V. PATRIKI* (DC! mss.) stigmate triangulari marginato, radice truncata duriusculo, foliis ovato-oblongis basi truncatis, sepalis lanceolatis, petalis 2 barbatis, stipulis dimidiatis adhærentibus. ♀ in Sibiria. V. primulifolia Lin. sp. excl. var. Virginiana. V. prunellæfolia Fisch. in litt. (v. s.)
- β. Chinensis*, foliis glabris, petalis intus omnibus imberbibus. — in Chinæ prov. Kianang. Georg. Staunton. V. primulæfolia β Lour. fl. coch. 2. p. 628? (DC. v. s. in h. Bauks.)
- γ. Napaulensis*, foliis glabris, petiolis folio dimidiò brevioribus (nec duplò vel triplò longioribus ut in α et β.) — in Napauliâ circa Chittong. Buchanan. V. primulæfolia Buchan! ined. (DC. v. s. in h. Lamb.)
13. *V. OVATA* (Nutt. gen. 1. p. 148.) villosa, stigmate marginato, radice truncata crasso subcarnoso, foliis ovatis obtusis basi in petiolum abruptè atte-

nuatis, sepalis lanceolatis, stipulis lanceolatis longis, petalis 2 dense barbatis. \mathcal{L} in collibus siccis Amer. bor. à Canadâ ad Virginiam. V. primulifolia Pursh fl. bor. am. 1. p. 172.* Bigel! fl. bost. 59.

β . *Belvisiana*, hirsutissima, petiolis vix nullis, pedunculis folio brevioribus, floribus minoribus. in mont. editoribus Virginie. Beauvois. An spec? (v. s.)

14. V. *SAGITTATA* (Ait! hort. kew. 3. p. 287.*) stigmatē marginato, trunco radicis fibroso, foliis oblongis basi sagittato-cordatis pubescentibus, stipulis liberis, sepalis lanceolatis. \mathcal{L} in Amer. bor. Variat foliis basi truncatis abrupte attenuatis plus minus pubescentibus. V. sagittæfolia Salisb. prodr. 130. Confer cum V. betonicæfoliâ. (v. s.)

β . *emarginata* (Nutt. gen. 1. p. 147.*) foliis triangulari-cordatis hastatisve subtus glabris. — in arvis arenosis New-Jersey.

γ . *dentata* (Lewis am. journ. n. 2.) glabra, foliis oblongis subhastato-grandidentatis. — in herbosis et sylvis humidis Pensylvaniæ. V. dentata Pursh fl. bor. am. 1. p. 172*.

15. V. *BETONICÆFOLIA* (Sm. in Rees cycl. n. 7.) stigmatē triangulari marginato, trunco subsimplici nigricante, foliis glabriusculis sagittato-lanceolatis cordatis, petiolis abrupte attenuatis, stipulis liberis, sepalis lanceolatis, capsulâ oblongâ trigonâ, valvis subobtusis, seminibus turbinatis fusco-rubris in raphe albis. \mathcal{L} in Nov. Holl. ad Bot. Bay et Port Jackson. V. longifolia Brown! ined. Flores interdum incompleti. (v. s. sp.)

β . *longiscapa*, foliis angustis lanceolatis basi abrupte decurrentibus (nec cordatis). — in orâ occidentali Novæ Hollandiæ (v. s. s.)

*** *Radicibus subsufiformibus.*

16. V. *PHILIPPICA* (Cav. ic. rar. 6. p. 19. t. 529. f. 2.) stigmatē marginato, radicibus fusiformibus simpliciusculis, foliis ovato-oblongis basi in petiolum breviorē abrupte attenuatis, stipulis adnatis, sepalis ovato-lanceolatis, seminibus fusco-rubentibus subglobosis. \mathcal{L} in insulæ Manillæ et Luçon lapidibus. Differt à V. betonicæfoliâ radicis formâ, à V. Gmelinianâ tantum foliorum formâ et calcare recto nec recurvo. (DC. v. s. in h. Mousp. ex Née.)

17. V. *GMELINIANA* (Rœm. et Schult. syst. 5. p. 354.) stigmatē marginato, radicibus fusiformibus simpliciusculis, foliis obovato-oblongis basi in petiolum attenuatis, stipulis adhærentibus, sepalis ovato-oblongis subobtusis. \mathcal{L} in siccioribus locis Sibirie et Dahuriæ. Gmel. fl. sibir. 4. p. 99. t. 49. f. 2. (v. s.)

β . *scorpiurifolia* (DC. mss.) hispidula, sepalis acuminatis longioribus, foliis obovato-oblongis, capsulis ovato-trigonis subobtusis, seminibus atro-rubentibus basi umbonatis, carunculâ albâ. — in Sibiria ad Nerschinsky Sawod. Fisch. (v. s. sp.)

γ . *elongata* (Ging. mss.) foliis cordato-ellipticis. — In locis humidis prope Irentiam. — Gmel. fl. sib. 4. p. 99.

δ ? *cordifolia* (Ging. mss.) foliis cordatis, sepalis acuminatis. — in mont. Uralensibus. V. suavis? Fisch! in litt. Radices subdivisæ; stigma depressum. An spec? An V. suavis Bieb. vera? (v. s. sp.)

**** *Radicibus (rhizomatibusve) articulatis*

18. V. *PALUSTRIS* (Lin. sp. 1324.) stigmatē marginato, radicibus articulatis squamosis, foliis reniformi-cordatis, stipulis lato-ovatis acuminatis, sepalis ovatis obtusis, capsulâ oblongo-trigonâ, seminibus ovoideis atro-viridescentibus. \mathcal{L} in pratis humidis et uliginosis montanis Europæ.

α . *vulgaris*, floribus dilute cærulescentibus, petalis obovatis. Variat nunc floribus majoribus, nunc floribus apetalis, et tunc foliis amplificatis. — Hall. hist. n. 560.

β . *Pensylvanica*, floribus purpurascens, petalis suborbiculatis. — in Amer. bor. V. cucullata Bigelow! in litt. (v. s. sp.)

19. *V. BLANDA* (Nutt. l. c.) stigmatē marginato, radicibus articulatis, foliis reniformi-cordatis, stipulis ovato-lanceolatis acutis, sepalis ovatis acuminatis, petalis (omnibus?) glabris. \mathcal{U} in pascuis humidis Americæ borealis. *V. blanda* Willd. hort. herol. 1. t. 24*? (v. s. sp.)

20. *V. ROTUNDIFOLIA* (Michx! fl. bor. amer. 2. p. 150.) stigmatē marginato, radice articulata squamosa, foliis orbiculato-cordatis sinu (demum) clauso subtus glabris, stipulis subulato-lanceolatis, sepalis oblongis angustis obtusis, petalis lateralibus barbatis. \mathcal{U} in excelsis montanis Carolinæ. *V. clandestina* Pursh fl. am. bor. 1. p. 173. Flores primum majusculos luteos, adhuc tenella profert, dein foliis amplificatis, flores exiguos viridescentes vix conspicuos, stolonēque reptantes producit. (v. s. c. à cl. Bigelow comm.)

β. pallens (Banks! herb.) glabra, sepalis acuminatis. — in Labrador et Kamtschatka. An eadem ac α ? An species? (DC. v. s. in h. Banks.)

21? *V. VILLOSA* (Walter fl. carol. p. 219.) villosa, stigmatē marginato vix rostrato, trunco radicis obliquo rugoso, foliis orbiculato-cordatis utrinque villosis, sepalis oblongis, stipulis lanceolatis dentatis, petalo infero imberbi, capsulis glabriusculis. \mathcal{U} in collinis umbrosis Amer. bor. Lewis amer. journ. 5. n. 1. spec. 9. (v. s. sp.)

22. *V. CORDIFOLIA* (Lewis l. c.*) stigmatē depresso valdè rostrato, foliis orbiculato-cordatis subtus glabris, petalo infimo barbato. \mathcal{U} in Amer. bor. *V. Sororia* Willd. hort. ber. t. 72. An verè spec. distincta? Affinis *V. papilionacæ*. (v. s. sp.)

23? *V. ASPERA* (Ging. mss.) hispidula, stigmatē recurvo rostro breviusculo apice subdepresso nudo? foliis profundè cordatis, sepalis oblongis acutiusculis, stipulis setoso-laceris, petalis 2 lateralibus valdè barbatis, nectariis conicis falcatis. \mathcal{U} in Napauliâ. Wallich. Aff. *V. hirtæ*, sed stigmatē differt. (v. s.)

§. 2. *Stigmatibus convexis, deorsum immarginatis.*

* *Capsulis turgido-subrotundis, caulibus subnullis.*

24. *V. JAPONICA* (Langsdorff ex Fisch. in litt.) glabra, foliis cordatis, stipulis lineari-lanceolatis subulatis, sepalis lanceolatis acutis calcare grosso obtuso recto petalis dimidio breviorē. \mathcal{U} propè Nagasaki Japoniæ. *V. odorata* Thunb. fl. Jap. 326? Flores cærulei. (v. s.)

25. *V. HIRTA* (Lin. sp. 1324.) villosa, stigmatē uncinato acuto nudo, foliis cordatis, sepalis ovatis obtusis margine ciliatis, stipulis glanduloso-dentatis, petalis 2 mediis subbarbatis, calcare subconico, nectariis glabris, capsulis turgidis hirtis, seminibus turgidis fuscis. \mathcal{U} in nemorosis et pratis Europæ ferè totius. Smith fl. brit. p. 244. engl. bot. t. 394. *V. hirta* in herb. Lin! est alia ex Americâ.

α. vulgaris, foliis lanceolato-cordatis. — in nemorosis et campis. Sturm deut. fl. ic. opt. Variat in stationibus variis, petiolis plus minus longis, corollâ plus minus lilacinâ vel pallidâ, floribus sæpè apetalis, interdum multicalcaratis. (v. v. sp. et c.)

β? fragrans, foliis basi subcordatis, floribus odoratis. — in Tauriâ. nullo modo differt à *V. hirtâ* sec. cl. Treviranum in beitr. nat. ges. berl. VII. p. 148. *V. campestris* Bieb. et *V. ambigua* Waldst. et Kit. sunt eadem sec. Bieb. suppl. p. 162. at *V. ambigua* semina fusca habet? An species? (v. s. sp.)

γ. Alpina, foliis dilatato-cordatis breviter acuminatis infrâ petiolos violaceis. — in Alp. Sabaudia ad Sancti-Bernardi. Variat sinu plus minus inciso, petiolo apice plus minus dilatato. An eadem ac *V. Pyrenaica*? An *V. collina* Bess. cat. hort. crem. anno 1816. p. 151. (v. s. sp.)

δ. alba, foliis cordatis minutissimis, floribus albis, calcare lilacino. — in pratis siccis. Semina turgida fusca. Folia seriùs majora. (v. s. sp.)

ε? scabra, foliis acuminatis, sepalis acutis ex Braun. — circa Salisburgum. *V. scabra* Braun in florâ 1820. p. 469.*

26. *V. PYRENAÏCA* (Raml in DC. fl. fr. 4. p. 803.) stigmatē uncinato nudo, foliis latē subcordatis, petiolis apice dilatatis, sepalis ovatis obtusis. \mathcal{U} in petrosis Pyrenæorum. Flores odorati ex Ram. Non *V. palustris* nec *V. caninæ* varietas, verosimiliter eadem ac *V. hirta* var. *Alpina*? Sem. ign. (v. s. sp.)
- 27? *V. HUMILIS* (H. B. et Kunth am. nov. gen. 5. p. 369.* t. 492. f. 1.) glaberrima, foliis subrotundo - ovatis obtusis basi rotundatis leviter cordatis, stipulis apice dentato-ciliatis, calycibus acutiusculis, stigmatē uncinato acuto, ovario glabro, calcare brevi rotundato patente, appendicibus apice rotundatis, loculis antherarum dimidio brevioribus. \mathcal{U} in subalpinis regni Mexicani, propē Real del monte. alt. 1426 hex. Flores albi, venis luteis.
- 28? *V. HOOKERIANA* (H. B. et Kunth am. nov. gen. 5. p. 369.* t. 492. f. 2.) glaberrima, foliis reniformibus profundē cordatis, stipulis dentato-ciliatis, calycibus acutis, stigmatē uncinato apice truncato, ovario glabro, calcare brevi rotundato, appendicibus apice rotundatis, loculis antherarum dimidio brevioribus. \mathcal{U} crescit cum præcedente.
29. *V. ODORATA* (Lin. sp. 1324.) stigmatē uncinato nudo, foliis rotundato-cordatis, sepalis ovatis obtusis, calcare obtusissimo, capsulā turgidā hirtā, seminibus turpinatis albidis, stolonibus flagelliformibus. \mathcal{U} in nemorosis et sepibus Europæ totius et in Sibiria, China, Japonia? Sm. fl. brit. p. 245. Sturm deutsch. fl. ic. opt. V. suavis Bieb. suppl. p. 164.
- a. *vulgaris*, floribus violaceis. Variat flor. corollatis purpureis, cinereis, rubris, pallidis, albis; calcare lilacino, flore simplici, semipleno, pleno, sæpē floribus apetalis et tunc plantis amplificatis; foliis longē petiolatis et pedunculos multò superantibus; tempore antheses verno, vel superiore, floribus multicalcaratis, sepalis interdū acutiusculis. (v. v.)
- β. *alba*, floribus albis, calcare albo obtusissimo, stolonibus longissimis. \mathcal{U} in locis argillosis cum *V. hirta* vulgò mixta. *V. alba* Bess. fl. gall. 1. p. 177. (v. v. c.)
- γ. *variegata*, floribus variegatis, Tourn. inst. 419.
30. *V. SERPENS* (Wall! in litt.) glabriuscula, stigmatē marginato? caulibus gracilibus, foliis profundē cordatis, sinu lato auriculis rotundatis, sepalis lanceolatis acutis, stipulis lanceolatis dentatis, floribus minutis, petalis a mediis barbatis, calcare brevi obtusissimo. \mathcal{U} in Napauliā. Wallich. An eadem ac *V. repens* Buchan! in herb. Lamb? (v. s. sp.)
- ** *Capsulis oblongis trigonis.*
- † *Caulibus herbaceis.*
31. *V. KROCKERI* (Gmel. syst. 412.) stigmatē uncinato, caule procumbente, foliis reniformi-cordatis obtusiusculis, sepalis lanceolatis acutis, capsulā trigonā pubescente. \mathcal{U} in Bohemiā. *V. purpurascens* Schm. fl. boh. 1. cent. 3. p. 49. t. 311.
32. *V. ULIGINOSA* (Bess. prim. fl. gall. p. 169.) stigmatē subreflexo nudo, caulibus abbreviatis, petiolisque glabris, foliis ovatis obtusis basi subcordatis, stipulis integris, sepalis ovatis obtusis, calcare brevi conico. \mathcal{U} in Ducat. Parmæ? in paludosis Carniolæ, Carinthiæ, Lusatiz. *V. uliginosa* Schrad. in Roem. et Schult. syst. 5. p. 357? An *V. scaturiginosa* Wallr. sched. 1. p. 97? (v. s.)
33. *V. LANGSDORFII* (Fisch. ined.) stigmatē marginato? subreflexo obtuso, caulibus demū elongatis basi obliquā, foliis rotundato-cordatis glabriusculis, stipulis ovatis setaceo-acuminatis, imis fuscis setoso-dentatis, sepalis ovatis subacutis, petali inferi limbo subrhomboidali, calcare latē saccato obtusissimo. \mathcal{U} in insulā Unalaschkā.
- a. subacaulis, capsulis oblongo-3-gonis obtusis, seminibus magnis subpyriformibus fuscis. *V. odorata* variet. in herb. Banks! (DC. v. s.)

- β. caulescens*, fructibus..... *V. peduncularis* Langsdorff ex Fisch. in litt. (v. s.)
34. *V. MIRABILIS* (Lin! sp. 13:6.) stigmatē subreflexo nudo, caulibus rigidis petiolisque villosis, pilis secundis, foliis reniformi-cordatis acuminatis, sepalis 3 maximis oblongis acuminatis, calcare cylindrico obtuso longo, capsulæ valvis peracuminatis, seminibus pyriformibus fuscis. \mathcal{L} in sylvis et dumetis montium Europæ temperatæ et frigidioris ferè totius. (DC. v. s. in h. Lin.)
- a. caulescens*, floribus radicalibus corollatis sæpius abortientibus, caulibus apetalis seminiferis. Jacq. fl. austr. t. 19. (v. v. sp.)
- β. acaulis*, pedunculis omnibus radicalibus, Gmel. fl. bad. 3. p. 519. in obs. (v. v. sp.)
- 35? *V. MAGELLANICA* (Forst! in comm. gotting. 9. p. 41. t. 8.) stigmatē, caule brevissimo, foliis reniformibus, stipulis lanceolatis acutis integris, sepalis ovato-lanceolatis acuminatis, calcare saccato. \mathcal{L} ad fretum Magellanicum. Petala intus glabra? Fl. lutei. (DC. v. s. in h. Mus. Par.)
36. *V. MACULATA* (Cav. ic. 6. p. 20. t. 539.) stigmatē apice subplano, rostro brevissimo, caule abbreviato, foliis ovatis, stipulis ovatis apice fimbriatis, sepalis lanceolatis acuminatis, staminibus apice emarginatis, calcare brevi obtuso. \mathcal{L} in Chili et ad fretum Magellanicum. Petala intus valdè barbata, pili clavati. Flores lutei.
- a. megaphylla* (DC. mss.) trunco radices longo tereti, foliis elliptico vel subrotundo-ovatis. — in sylvis bahiæ Duclos ad fretum Magellanicum *V. pyrolæfolia* var. *a.* Poir! dict. 8. p. 636. excl. patriæ. *V. glandulosa* Dombey! herb. *V. lutea megaphyllos* Commers. (DC. v. s. sp. in h. Mus. Par.)
- β. microphylla* (DC. mss.) trunco radices brevissimo, foliis ovato-lanceolatis. — in collibus bahiæ Boucault Commers. *V. pyrolæfolia* var. *β.* Poir. l. c. *V. lutea microphyllous* Commers! herb. (v. s. s.)
37. *V. COMMERSONII* (DC. mss.) stigmatē caule brevissimo, foliis ovatis, stipulis lato-ovatis integris?, sepalis oblongo-lanceolatis obtusiusculis. \mathcal{L} in summis montibus freti Magell. suprâ Fort.-Gallant. Commerson. Flos albus. (DC. v. s. in h. Mus. Par.)
38. *V. RADICANS* (DC. mss.) rhizomate horizontali fibrilloso, stigmatē brevirostrato marginato, foliis ovato-lanceolatis, basi abruptè attenuatis subcordatisve serratis, stipulis linearibus setaceo-subulatis setoso-serratis, sepalis linearibus acutis, floribus minutis, petalis mediis obovatis imberbibus? infimo minori, calcare subnullo. \mathcal{L} in Carolinâ austr. Michx. fil. (v. s.)
39. *V. STRIATA* (Ait. hort. Kew. ed. 1. v. 3. p. 291?) stigmatē papilloso-subreflexo, rostro obtusiusculo, caulibus ramosis flexuosis, foliis ovato-cordatis acuminatis pubescentibus, stipulis ovato-lanceolatis dentato-laceris, sepalis ovato-lanceolatis, capsulâ breviusculâ, valvis subobtusis 3-7-spernis, seminibus subrotundis rufis. \mathcal{L} in sylvis Amer. bor. Poir. dict. enc. 8. p. 637. Elliot sket. 1. n. 111. p. 301.* *V. debilis* Michx. non Pursh. Species plurimæ confusæ sec. Lewis l. c. Flores flavidi. (v. v.)
40. *V. DICHOTOMA* (fl. mex. ic. ined.) stigmatē caule dichotomo, ramis patentibus, foliis cordatis, stipulis oblongis, apice tantum setoso-laceris. \mathcal{L} in Mexico. *V. striatæ* valdè proxima. Flores violacei. Fruct. ign. An eadem ac *V. scandens* Ruiz et Pav. in herb. Lamb?
41. *V. OCHROLEUCA* (Lewis l. c.*) caulibus assurgentibus, foliis cordatis, stipulis oblongis maximis remotè dentatis, sepalis angustissimè lanceolatis acuminatis, floribus magnis, petalis lateralibus densè barbatis, calcare longiusculè producto. \mathcal{L} ? secus flumen Dam-River, in montibus Saura Amer. bor. Flores ochroleuci. *V. striata* Pursh fl. bor. am. 1. p. 174. Nutt. 1. p. 150. Au var. v. striatæ? (v. s.)
42. *V. MÜHLENBERGIANA* (Ging. mss.) caulibus flexuosis assurgentibus, foliis

cordatis acuminatis, calcare brevissimo, stipulis lanceolatis ciliatis, radice fasciculatâ. — in Pensylvaniâ. Flores cœrulescentes. V. uliginosa. ex Muhl. cat. n. 18 non Schrad. nec Besser: Lewis l. c.* An V. dichotomæ var?

43. V. LEWISIANA (Ging. mss.) caulibus decumbentibus stoloniferis, foliis reniformibus cordatisque, stipulis magnis ovatis longissimè deusèque ciliatis, sepalis latè lanceolatis acuminatis ciliatis, floribus minoribus, petalo infimo minutissimo aequali lateralibus, barbâ albâ, calcare breviusculo. ½ in rupibus montium Sauræ Amer. bor. V. repens Lewis l. c.*

44. V. CANINA (Lin! sp. 1324.) stigmatè papilloso subreflexo, caule adultiore adscendente ramoso glabro, foliis cordatis, stipulis acuminatis pectinatim tenuiter laceris, sepalis subulatis, pedunculis glabris, capsulâ elongatâ, valvis acuminatis, seminibus pyriformibus fascis. ½ in sylvis dumetis sepiusque totius ferè Europæ, Japoniæ, Persiæ, oræ occid. Amer. sept., et in insulis Canariensibus. Smith engl. bot. t. 620. V. sylvestris Lam. fl. fr. 2. p. 680. V. cordata Willd. sp. 164. non Walt. V. neglecta Schm. fl. boh. n. 1. p. 51. t. 313. non Bieb. Multum variat altitudine et foliorum formâ. Sæpius excrecit et tum flores apetalos producit.

β. minor (DC! fl. fr. suppl. p. 617.) caule brevissimo, foliis reniformi-cordatis glabriusculis. — in montibus Pyrenæicis et Helveticis. (v. v. et sp.)

γ. alba, floribus albis, stipulis uno latere ciliato-laceris. — in Tauriâ Stev. V. neglecta Bieb. fl. taur. 1. p. 172. sec. Stev. V. canina fl. albo Dill. in Ray. synops. 364? (v. s. sp.)

δ. macrantha, floribus duplò triplòve majoribus. (v. s. sp. in herb. Balb.)
? Japonica, Foliis basi vix cordatis. — in Japoniâ propè Nagasaki. V. non odorata foliis cordatis Lamb! herb. (DC. v. s. in h. Lamb.)

45. V. PALMARIS (Buchan! ined.) stigmatè caule ramoso, ramis debilibus, foliis cordatis acuminatis, stipulis lineari-lanceolatis subserratis, sepalis oblongis acutiusculis. ½ in Napauliâ prope Bimpedi. An var. V. caninæ? (DC. v. s. in h. Lamb.)

46. V. DIFFUSA (Ging. mss.) stigmatè subcapitato, rostro brevissimo obtuso nudo, caulibus simplicibus decumbentibus, foliis ovatis crenatis in petiolum decurrentibus, nervis marginibusque pilosis, stipulis lanceolatis dentatis ciliatis, sepalis lanceolatis acutis, nectariis infernè latis, extus suprà emarginatis hinc attenuatis brevissimis angustis. ½ in Napauliâ. Wallîch. Aff. V. Nuttali Pursh. (v. s.)

47. V. ROSTRATA (Pursh fl. am. bor. 1. p. 174.) stigmatè rostrato, rostro suberecto acuto nudo, caulibus simplicibus decumbentibus, foliis cordatis glabris, sepalis subulato-lanceolatis glabris, calcare reflexo rostrato. ½ in rupibus umbrosis Pensylvaniæ et in montibus Alleghany. Petalâ omnia? glabra. (v. s. sp.)

48. V. DEBILIS (Pursh! fl. bor. am. 1. p. 174. non Michx.) stigmatè papilloso subreflexo, caulibus subsimplicibus decumbentibus, foliis reniformi-cordatis acutiusculis glabriusculis, sepalis lanceolatis glabris, calcare longè porrecto. ½ in Amer. bor. V. canina Walt. fl. carol. ex Elliot. An var. V. caninæ? Flores cœrulescentes. (v. s. sp.)

49. V. ARENARIA (DC! fl. fr. 4. p. 806.) pruinoso-pubescentis, stigmatè papilloso subreflexo, caulibus simplicibus erectiusculis, foliis cordato-subrotundis, stipulis erectis inciso-dentatis, sepalis lanceolato-oblongis, capsulis turbinato-ellipticis obtusis pubescentibus, seminibus ovatis castaneis. ½ in locis arenosis Vallesiæ, in Alpibus Pedemontii, Gallo-provinciæ, Bohemiæ. circa Mosquam, in Caucaso. V. Allionii Pio diss. p. 20. t. 1. f. 2. V. glauca Bieb. suppl. p. 165. V. rupestris Schm. fl. boh. cent. 3. t. 1. p. 50. t. 312. V. pusilla Schleich. in litt.

β. aprica, perpusilla, subacaulis. ½ in collibus ad Lobenheim et Bendstadt V. hirta aprica Spreng! nov. prov. 43. n. 97.

γ? sarmentosa, caulibus (ob chlorosim) luxuriantibus elongatis præ-

cumbentibus, pedunculis foliis brevioribus. — in *Alpium Caucasicarum cryptis* Bieb. *V. sarmentosa* Bieb. fl. taur. cauc. i. p. 172. et suppl. p. 166.

50. *V. PUMILA* (Vill! Dauph. 2. p. 266. cat. strasb. p. 288. t. 5.) glabriuscula, stigmate papilloso subreflexo, caulibus ramosis diffusis procumbentibus, foliis ovato-lanceolatis, petiolis marginatis, stipulis ovato-lanceolatis dentatis incisive, calcare cylindrico apice subrecurvo obtuso flavescente sepalorum vix longitudine, valvis subrectis glabris apice truncatis, seminibus ovatis nigro-castaneis. ♀ in Delphinatû Monte Corps, in dunis Vasceniæ, in ericetis siccioribus Germaniæ, Helvetiæ, Pedemontii. (v. s.)

α. *lancifolia*, foliis lanceolatis basi in petiolum gradatim attenuatis. *V. lancifolia* Thore! chlor. land. 357. (v. s. sp.)

β. *ericetorum*, foliis basi truncatis vel subcordatis in petiolum abrupte attenuatis. — In ericetis turfosis. *V. ericetorum* Schrad! ined. (v. s.)

γ. *littoralis*, caulibus elongatis, floribus apetalis. — Ad oras maris Baltici propè Warnemunde. *V. littoralis* Spreng! nov. prov. p. 43. (v. s.)

51. *V. MONTANA* (Lin! sp. 1325.) glabriuscula, stigmate papilloso subreflexo, caulibus simplicibus erectis, foliis imis cordatis, superioribus ovatis acutis, petiolis marginatis, stipulis oblongis dentatis incisive, calcare conico truncato recto viridescente sepalis brevioribus, valvis arcuatis subacutis, seminibus ovatis castaneis. ♀ In Europâ temperatâ et Sibirîâ. Flores albi: demum cærulescentes. (v. v.)

α. *stricta*, caulibus strictis, foliis imis cordatis. *V. stricta* Horn. hafn. *V. montana*. α. fl. fr. 4. p. 807.* in pratis montanis dumetisque. (v. v.)

β. *pubescens*, caulibus strictis, foliis imis cordatis pubescentibus. *V. erecta* Gerard! in litt. (v. s.)

γ. *lactea*, caulibus basi geniculatis, foliis omnibus lanceolatis. — in ericetis udis Angliæ, Helvetiæ, Galliæ. *V. lactea* Smith fl. brit. i. p. 247. engl. bot. t. 445. opt. *V. Lusitanica* Brot. phytog. p. 39. t. 17.

δ? *Ruppîi*, caulibus procumbentibus, foliis cordatis lanceolatisque. — In Pedemontio. *V. Ruppîi* All. fl. ped. 2. n. 1646. t. 26.* Lodd. bot. cab. t. 686. (v. v.)

†† *Caulibus fruticulosis.*

52. *V. DECUMBENS* (Lin! fil. suppl. 397.*) stigmate uncinato nudo, caulibus ramosissimis procumbentibus, foliis linearibus angustissimis integris confertis, stipulis subulato-linearibus adnatis, calcare tubuloso obtuso recto sepalorum vix longitudine. ♀ Lin. f. 5 ad Cap. Bon.-Spei in montibus Hottentot. holl. (DC. v. s. sp. in herb. Lin. et Lamb.)

53. *V. SCROTIFORMIS* (DC. mss.) stigmate.... caulibus ramosis, foliis linearibus obtusiusculis integris, imis remotis, stipulis subulatis adnatis, calcare saccato, auriculis sepalis vix superantibus. 5 in promont. B. Spei. Masson. (DC. v. s. sp. in herb. Lamb.)

54. *V. ARBORESCENS* (Lin. sp. 1325.) stigmate uncinato nudo, caulibus ramosis erectis teretiusculis pubescentibus, foliis lanceolatis basi attenuatis, stipulis oblongis adhärentibus, calcare cucullato obtuso auriculas vix superante, valvis naviculatis obtusis oligospermis, seminibus ovatis antè dehisceniam purpureo-nigris dein piceis, carunculis albis. 5 in saxosis et inter fissuras rupium sabuletaque Europæ merid. et Africæ bor. DC. fl. fr. suppl. p. 618.*

α. *linearifolia*, pruinosa, foliis elongato-linearibus integris. — in Corsicâ. (v. s.)

β. *integrifolia*, foliis lanceolatis integris. — in sabuletis oræ Mediterraneæ, in rupibus Hispaniæ. *V. Hispanica fruticans* Barr. ic. t. 568. (v. s.)

γ. *serratifolia*, foliis lanceolatis remotè dentatis. — In Barbariâ. *V. suberosa* Desf! fl. atl. 2. p. 313.* et emend. (v. s.)

δ. *latifolia*, glabriuscula, foliis rhombeo-spathulatis remotè dentatis. (v. s. ex Brouss.)

55. *V. ARBORÆA* (Forsk! fl. yem. cxx.) stigmatē caule basi tereti, apice angulato erecto simplici?, foliis oblongis utrinquē acuminatis remotē dentatis, stipulis subulatis integris, calcare obtusissimo parvo. $\frac{3}{4}$ in Yemen. An eadem ac *V. arborescens* var. *V. serratifolia*? (DC. v. s. in h. Banks.)

Sect. II. DISCHIDIUM. *Ging. mss.*

Stigma erostratum, apice plus minùs bilobum, foraminulo inter lobos sito. Stylus ab apice ad basin attenuatus. Stamina oblonga approximata Torus planiusculus. Capsula sæpè 3-gona oligosperma. Folia seminalia sæpè subrotunda,

56. *V. PYGMÆA* (Poir. dict. 8. p. 630.*) stigmatē bilobo?, lobis divergentibus, trunco crasso perpendiculari, foliis confertissimis linearibus acutis margine scabris sessilibus, sepalis linearibus, calcare breviusculo, petalis apiculatis. $\frac{2}{4}$ in locis frigidissimis Andium Peruviae. *V. Alpina* Ruiz et Pav! in herb. Deless. ined. (v. s.)

57. *V?* *COTYLEDON* (Ging. mss.) stigmatē bifido, lobis reflexis, trunco crasso tortuoso, foliis confertissimis rhombeo-spathulatis in petiolum longē attenuatis integris margine cartilagineis glabris, sepalis linearibus, calcare brevissimo. $\frac{2}{4}$ in montibus excelsis Chili. Née. Capsula 3-gona oblonga in cacumine pedunculi spiraler contorti; semina ovata 9-10 atrata. (v. s. in herb. Roux, nunc Mus. Genev.)

58. *V. TRIDENTATA* (Menzies! ined.) stigmatē caulibus ramosissimis procumbentibus, foliis confertis obovato cuneatis apice tridentatis coriaceis glabris, sepalis ovatis, calcare brevissimo. $\frac{5}{4}$ in Statenland vertice montium inter nives. (DC. v. s. in h. Lamb. et Banks.)

59. *V. WALLICHIANA* (Ging. mss.) stigmatē bifido?, lobis lamellatis divaricatis, foliis reniformibus cordatisve, stipulis lanceolatis denticulatis, sepalis subulatis, calcare subulato petalorum longitudine. $\frac{2}{4}$ in Napauliā. Calcar geniculatum descendens; nectaria elongata, capillaria; petalum inferum obovato-cuneatum, apiculatum, flos flavus? Habitus *V. bifloræ*. Flos ferè *V. ros-tratæ*; stigma ab omnibus discrepat. (v. s. comm. à cl. Wallich.)

60. *V. BIFLOA* (Lin! sp. 132.) stigmatē bifido, lobis crassis divergentibus, foliis reniformibus, stipulis ovatis, sepalis linearibus, petalis glabris, calcare brevissimo, seminibus rotundo-ovoideis fuscis obscure punctatis. $\frac{2}{4}$ in humidis montanis Europæ ferè totius, Sibiriae et Americæ bor. ad oram occidentalem? Variat caule unifloro (J. Bauh. hist. 3. p. 545. f. r.) et bifloro (Pluk. t. 233. f. 7. et 234. f. 1.). (v. s. sp.)

Sect. III. CHAMÆMELANIUM. *Ging. mss.*

Stigma sphæroïdeo-capitatum, utrinquē fasciculis pilorum onustum, foraminulo minuto sublaterali. Stylus compresso-clavatus. Stamina oblonga approximata. Torus planiusculus. Capsula sæpè trigona. Folia seminalia sæpiùs subrotunda. Petala 2 ungue barbata.

61. *V. TRIPARTITA* (Ell. sket. i. p. 302.*) foliis 3-5-partitis, lobis lanceolatis dentatis pubescentibus, stipulis lanceolatis, sepalis acutis. $\frac{2}{4}$ in Georgia propè Athens. Flores lutei. An var. *V. hastatæ*?

62. *V. NUTTALLII* (Pursh! fl. bor. amer. p. 174.*) caule simplici erectiusculo, foliis lanceolato-ovatis integris in petiolum attenuatis, nervis marginibusque pubescentibus, stipulis lanceolatis integris, sepalis lanceolatis acutis. $\frac{2}{4}$ in Americâ bor. inter flumen Roch River et Missouri. Nutt. gen. am. 1. p. 151.* Flores parvi, flavi, subtùs purpurei.

63. *V. HASTATA* (Michx! fl. bor. amer. 2. p. 149.) foliis cordato-lanceolatis hastatisve sinuato-serratis, stipulis ovatis acutis, sepalis lanceolatis acutis, calcare brevissimo. $\frac{2}{4}$ in Americæ borealis Georgia propè Athens. Elliot Sket. i. p. 302.* Flores lutei. (v. s.)

♀? glaberrima, foliis rhomboïdeo-lanceolatis. — In sylvis et collibus Carolinæ septentrion. *V. hirta* Lewis l. c. non auctorum. An spec.?

64. *V. CANADENSIS* (Linl. sp. 1326.) foliis latè cordatis serratis acuminatis, nervis pubescentibus, stipulis lanceolatis, sepalis subulatis lanceolatis, calcare brevissimo subsaccato, capsulâ oblongo-3-gonâ obtusissimâ, seminibus subrotundo-ovatis castaneis. 2 $\frac{1}{2}$ in Americâ bor. Petala albida dorso violacea. (v. s.)

α . stipulis angusto-lanceolatis. *V. Canadensis* Ell. Sket. p. 301.*

β . stipulis ovato-lanceolatis. *V. albiflora* Link. enum. h. berol. 1. p. 141.

65. *V. PUBESCENS* (Ait! hort. Kew. ed. 1. v. 3. p. 290.) villosa, caulibus simplicibus erectis, foliis lato-cordatis acuminatis serratis, stipulis ovatis, sepalis oblongo-lanceolatis, calcare brevissimo subsaccato. 2 $\frac{1}{2}$ in Americâ bor. sylvis rupestribus. Nutt. gen. 1. p. 150.* (v. s. sp.)

α . ovario glabriusculo, stipulis ovatis latis.

β . ovario villosa, stipulis oblongo-ovatis. *V. Pensylvanicâ* Michx! fl. bor. amer. 2. p. 149.

66. *V. KRIOCARPA* (Lewis am. journ. 5. n. 1.) scabriuscula, caulibus assurgentibus ramosis, foliis cordatis acutis, interdum reniformibus, stipulis ovatis, calcare brevissimo subsaccato, fructibus dense villosis. 2 $\frac{1}{2}$ in Carolinâ sept. Non prioris var. ex Nutt. (v. s. sp.)

67. *V. UNIFLORA* (Linl. sp. 1327.) foliis radicalibus reniformibus, caulinis ovatis acuminatis inciso-dentatis pubescentibus, stipulis lanceolato-subulatis dentatis, sepalis ovatis, calcare brevi lato saccato. 2 $\frac{1}{2}$ in Sibirîâ, solo turfoso Gmel. sib. 4. p. 101. t. 48. f. 5. (v. s.)

Sect. IV. MELANIUM. DC. mss. — Jacea DC. fl. fr. — *Violæ tricolores veter. auct.*

Stigma urceolatum utrinquè fasciculis pilorum onustum, aperturâ magnâ infernè labello munitâ. Stylus ab apice ad basin attenuatus. Stamina triangularia connexa. Torus concavus et idèò ovarium paulò inferum. Capsula obsoletè hexagona. Semina numerosissima 40.-60. Folia seminalia sæpius ovata. Petala 3 inferiora, ungue barbata.

68. *V. NUMMULARIFOLIA* (All! pedem. n. 1640. t. 9. f. 4.) glaberrima, caulibus ramosis decumbentibus, foliis subrotundis integris, stipulis lanceolatis setoso-dentatis, sepalis oblongis, calcare breviusculo. 2 $\frac{1}{2}$ in saxosis Alpium Pedemontii et Delphinatûs. DC. fl. fr. 4. p. 804.* (v. v.)

β . minima — in Alpibus Corsicis. — Bocc. mus. 2. p. 163. t. 127. (v. s.)

69. *V. ALPINA* (Jacq! obs. 1. p. 21. t. 11.*) caule brevissimo, foliis cæspitosis ovato subrotundis subcordatis crenatis longè petiolatis, stipulis lanceolatis vix dentatis adherentibus, sepalis oblongis obtusiusculis, calcare tubuloso obtuso sepalis breviorè. 2 $\frac{1}{2}$ in Alpibus Austriæ et Carpathis. (v. s.)

70. *V. CORNUTA* (Lin. spec. 1325.) radice fibrosâ diffusâ, caulibus ascendentibus, foliis cordato-ovatis crenatis ciliatis, stipulis obliquè cordatis inciso-dentatis ciliatis, sepalis subulatis, calcare subulato calyce longiorè. 2 $\frac{1}{2}$ in Pyrenæis et monte Atlante. DC! fl. fr. 4. p. 810.* Sims bot. mag. t. 791. (v. v.)

71. *V. CENISIA* (All! pedem. n. 1641.) radice trunco subfusiformi lignoso, caulibus diffusis procumbentibus indivisis, foliis spathulato-ovatis oblongisve integris, stipulis integris vel palmatisectis segmentis obovatis folio simillimis integris, calycibus pubescentibus, calcare tubuloso acuto calycis longitudine. 2 $\frac{1}{2}$ in saxosis Alpium Helvetiæ Pedemontii et Galloprovinciæ. (v. s.)

α . ovatifolia, foliis omnibus ovatis vix pubescentibus. *V. Cenisia* All! fl. pedem. t. 22. f. 6. DC! fl. fr. 4. p. 805.* (v. s.)

β . diversifolia, foliis imis ovatis, superioribus oblongis tomentoso-pubescentibus. — in Pyrenæis. *V. Cenisia* Lapeyr. abr. Aff. V. cheirantifoliæ. (v. s.)

- γ. *Valderia*, foliis omnibus oblongis hispidulis. V. *Valderia* All! fl. pedem.
2. p. 98.* t. 24. f. 3. (v. s.)
72. V. *MINUTA* (Bieb. fl. taur. i. p. 173.) caulibus simplicibus unifloris flac-
cidis, foliis subrotundis crenatis, stipulis ovatis integris ciliatis, calcare vix
petalorum longitudine. ℥ in Caucaso Iberico. An var. V. *Alpinæ*?
73. V. *CHEIRANTHIFOLIA* (Humb. et Bonpl. pl. equin. i. p. 111.* t. 32.) velu-
tina, radice crassâ lignosâ, caulibus adscendentibus, foliis lanceolatis inte-
gerrimis in petiolum attenuatis, stipulis linearibus, calcare tubuloso subacutâ
sepalorum vix longitudine. ℥ in insulis Canariis in summo monte Pico-
Teyde. (v. s.)
74. V. *CALCARATA* (Lin! spec. 1325.) radice fibrosâ diffusâ, caulibus simpli-
ciscusculis cæspitosis, foliis spathulato-subrotundis elongatisve crenatis, stipu-
lis palmatifidis integrisve, sepalis oblongis glanduloso-denticulatis, calcare
subulato calyce longiore, nectario staminibus vix duplo longiore, seminibus
ovatis. ℥ in pascuis montium editiorum ab Austria ad Galloprovinciam. DC.
fl. fr. 4. p. 810*. Species minè varians, nec intra limites suos certa.
- α. *Zoysii*, caule brevissimo, stipulis glabriusculis bidentatis, flore luteo
amplissimo. — In Alpibus Austriae et Helvetiae. V. *Zoysii* Wulf. in Jacq.
coll. 4. p. 297. t. 11. f. 1.* — Barr. icon. t. 691. (v. v.)
- β. *Halleri*, caule brevi, stipulis incisus vix ciliatis, flore caruleo amplo. —
Hall. hist. n. 566. var. α p. 243.* t. 17. In iisdem locis frequens. (v. v.)
- γ. *albiflora*, caule brevi, stipulis incisus vix ciliatis, flore albo amplo. —
In Alpibus Sabaudiae. Hall. hist. n. 566. var. β. 111.* (v. v.)
- δ. *Bertolonii*, caulibus elongatis, foliis imis ovatis, extimis elongatis sti-
pulisque palmatifidis glabriusculis. — In Apennini Genuensis summo
monte Scaggia. V. Bertoloni Pio diss. 34. t. 3. f. 2. — Cupan. pamph. ed.
bonon. t. 99. (v. v.)
- ε. *Æthnensis*, caulibus elongatis, foliis stipulisque velutino-pubescentibus,
calycibus glabriusculis. — In monte Ætnâ. V. gracilis Riv. mant. 4.
p. 25. ex cl. Moricaud. — Cupan. pamph. ed. bonon. t. 138. (v. s.)
- ϑ. *decepiens*, caulibus elongatis, foliis stipulisque hispidis, sepalis lanceo-
latis hispidulis. — In Pyrenæis orient. V. hispida Lapeyr. abr. 123.
excl. syn. (v. s.)
75. V. *OREADES* (Bieb. fl. taur. suppl. 167.) caule abbreviato, foliis oblongis,
stipulis pinnatifidis integrisve, laciniis obtusis, sepalis acutis denticulatis,
calcare tubuloso obtuso calyce brevior. ℥ in cacumine Tauriae montium.
An fortè varietas V. *amænæ*? (v. s.)
76. V. *AMÆNA* (Sym. synop. ex Smith engl. bot. t. 1287.) caule abbreviato,
foliis rotundis, stipulis pinnatifidis, lobis obtusis, sepalis ovato-lanceolatis
acutis integris, calcare tubuloso obtuso calyce brevior. ℥ in montibus
Scoticis. Flores amplissimus purpureus. Petala rotundata undulata. (v. s.)
77. V. *ALTAICA* (Ker. bot. reg. t. 54.*) caule abbreviato, foliis ovalibus, sti-
pulis oblongo-lanceolatis incisus, dentibus acutis, sepalis acutis denticu-
latis, calcare brevissimo appendiculis sepalorum vix longiore. ℥ in Alpi-
bus Altaicis ad fines Sinenses. Sims bot. mag. t. 1776. V. *grandiflora* Lin!
herb. et mant? V. *Pallasii* Fisch. cat. h. gor? Flores magni lutei. (v. s.)
- β. *purpurea*. V. *Altaica* purpurea Fisch. in litt.
78. V. *SUDETICA* (Willd. enum. suppl. 12.) radice fibrosâ diffusâ, caulibus
simpliciscusculis, foliis ovato-oblongis, stipulis palmatifidis, lobis integris,
sepalis lanceolatis, petalis cuneatis longè unguiculatis discretis, calcare subu-
lato acutiusculis porrecto nectariis staminum vix longiore, seminibus ovatis.
℥ in pratis Alpibus et montanis Europæ. V. *grandiflora* DC! fl. fr. 5. p. 620.*
et omnium ferè auctorum.
- α. *lutea*, caulibus brevioribus confertè foliosis, foliis ovato-oblongis,
floribus luteis. — in pascuis Alpium Helvetiae et Britanniae. — Cupan.

- pamph. t. 133. V. lutea Smith fl. brit. 1. p. 248.* engl. bot. t. 721. V. grandiflora var. *a*. DC! fl. fr. 5. p. 620. excl. syn. (v. v. c. et s.)
- β. *calaminaria*, caulibus adscendentibus, foliis subremotis ovatis, floribus luteis. — in pascuis siccis et calaminariis prope aquas Grani. Lejeune. V. lutea DC! fl. fr. 5. p. 619.* excl. syn. (v. s.)
- γ² *media*, caulibus elongatis erectiusculis, foliis remotis, floribus purpureis. — in montibus Jurassi, Vogesorum et Arvernæ. (v. s.)
- δ. *ramosior*, caulibus ramosis, foliis subremotis ciliatis, floribus tricoloribus. — Circa Verviers. Lejeune. V. Rothomagensis var. DC! fl. fr. 5. p. 619. An V. tricolor. var? An sint V. tricoloris aut V. calcaratæ varietates, aut species nescio!
79. V. *PROSTRATA* (Roem. et Schult. syst. 5. p. 582.*) radice simpliciusculâ, caulibus decumbentibus, foliis imis subrotundis, supremis ovalibus, omnibus subcordatis grossè dentatis, dentibus cartilagineis, stipulis pinnatifidis, lobo medio oblongo integro, calcare vix producto obtuso. 2/ in cacamine montium Teneriffæ et in Appenninis. V. Ludovicea Jan in litt. Petala ochroleuca nigro-striata vix incumbientia. (v. s.)
80. V. *ROTHOMAGENSIS* (Desf. cat. 153.) hispida, radice subfusiformi, caulibus ramosis diffusis, foliis ovalibus, imis subcordatis, stipulis pinnatifidis, calcare tubuloso obtuso sepalis brevioribus, nectariis staminum brevioribus, seminibus oblongo-obovatis. 2/ in rupibus et aggeribus cretaceis à Rothomago ad Meldam. DC! fl. fr. 4. p. 809.* Pio diss. t. 2. V. hispida Lam. fl. fr. 2. p. 679. Vix ac ne vix differt à V. tricolore. (v. v.)
81. V. *TRICOLOR* (Lin! spec. 1326.) radice subfusiformi, caulibus ramosis diffusis, foliis imis ovato-cordatis, stipulis runcinato-pinnatifidis, lobo medio crenato, petalis breviter unguiculatis incumbentibus, calcare crasso obtuso non porrecto, nectariis brevibus, seminibus oblongo-ovatis. ① ② in arvis Europæ, Sibiriz et Americæ bor. DC! fl. fr. 4. p. 808.* Species mirè varians.
- a*. *hortensis* (DC. mss.) petalis intensè velutinis calyce multò majoribus. V. tricolor OED. fl. dan. t. 623. (v. v.)
- β. *degener* (DC. mss.) subramosa, floribus tricoloribus subvelutinis calyce paulò majoribus, stipulis maximis. — in oleraceis (v. v.) *
- γ. *alpestris* (DC. mss.) elongata, foliis remotis, floribus sulfureis purpureo maculatis calyce majoribus. — in pratis Alpium. V. lutea Tratt. tab. 43? (v. s.)
- δ. *crassifolia* (DC. mss.) foliis maximis crassiusculis internodiis longioribus, floribus luteis calyce vix majoribus. — in Neustriâ maritimâ. (v. v. c.)
- e*. *calycina* (DC. mss.) foliis maximis ovatis subconfertis, calycibus maximis. — in montibus Teneriffæ. Brouss. (v. s.)
- ζ. *appendiculata* (DC. mss.) foliis utrinquè attenuatis, calycibus valdè appendiculatis corollâ majoribus. — inter Bagdad et Kermancha. Olivier et Brng. An eadem ac calycina? (v. s. s.)
- η*. *purpurea* (DC. mss.) foliis ovatis, sepalis breviusculis latis valdè appendiculatis, floribus purpureis calyce majoribus. — in Armoricæ inferioris campis. (v. s. s.)
- θ. *declinata* (Ging. mss.) foliis lanceolatis vix appendiculatis, sepalis elongatis, floribus purpureis calyce majoribus. — in graminosis Alpibus Transylvaniæ et Hungariæ; in præruptis rupium montis Corno-di-Canzo in Italiâ. Daby. V. declinata Waldst et Kit. pl. hung. 3. p. 248.* t. 223. (v. s. s.)
- ι. *ovatifolia* (DC. mss.) caulibus diffusis ramosis, foliis ovatis, floribus tricoloribus sepalis lanceolatis vix longioribus. — in (v. s.)
- κ*. *arvensis* (DC! mss.) caulibus ramosis assurgentibus, petalis calyce vix longioribus flavescentibus cæruleis vel purpureo-maculatis. — in agris et arenosis frequens. V. arvensis DC. fl. fr. 4. p. 808.* An V. tricolor Pursh fl. amer. sept. 1. p. 175. n. 22? V. tenella Lewis am. journ. scienc. 5. n. 1. spec. 29. (v. s. s.) — * Toro fructibusque ex Uredine

carbone? deformatis. — in campis prope Lazaretum Cafsæ D'Urville. (v. s. s.)

- λ *sabulosa* (DC. mss.) caulibus plurimis diffusis, foliis remotis ovatis elongatisve, sepalis angusto-lanceolatis corollâ vix brevioribus. — in arenosis maritimis Belgii et Galliæ. (v. s. s.)
- μ. *gracilescens* (DC. mss.) caulibus simpliciusculis elongatis erectis, sepalis angustis corollâ bicolore sæpius longioribus. — in agris prope Bernam. (v. s. s.)
- ν. *trimestris* (DC. mss.) caule erecto tenuissimo, stipulis minimis, sepalis linearibus. — circa Olyssiponam. V. tricolor trimestris flore variorum colorum elegans Grisl. vir. lus. (v. s.)
- ξ. *hirta* (Ging. mss.) tota velutino-hirta, ac præcocior (aprili floret!). in Branson Vallisæ. An V. Kitaibeliana Rœm. et Schult. syst. 5. p. 383? (v. s. s.)
- ο. *bellioides* (DC. mss.) hispidula, caulibus brevissimis, foliis subrotundis confertis, petalis calyce brevioribus. — in arenosis circa Montpelium et Castelnau, in Nebrodibus Siciliæ. V. parvula Tineol pug. sic. 5. n. 3. (v. s.)
- π. *nana* (DC. mss.) glabriuscula, caule brevissimo, cotyledonibus per anthesim persistentibus. in arenosis maritimis Neustriæ, Syriæ, Asiæ minoris. V. tenella Poir. dict. 8. p. 644. (v. s. s.)

Sect. v. LEPTIDIUM. *Ging. mss.*

Stigma proboscideum foramine minuto; stylus subulatus, flexuosus. Stamina oblonga approximata 2 sæpius appendicibus terminalibus longioribus. Torus planiusculus? Capsulæ subrotundæ? oligospermæ.

- 82. V. *RUBELLA* (Gav. ic. rar. 6. p. 20.* t. 351. f. 1.) caulibus erectis?, foliis ovatis oblongisve acutis, stipulis lanceolatis setaceo-dentatis, calcare conico brevi. ξ in Chili circa Santo-Carlos. Cav.
- 83. V. *CONCHORIFOLIA* (Domb! herb.) pubescens, caulibus erectis, foliis ovatis acutis basi inæqualibus, serraturis exsertis apiculatis, stipulis lanceolatis setaceo-laceris, calcare latissimo saccato, capsulâ ovato-trigonâ. ξ circa Huanaco in Peruvia. V. punicea Ruiz et Pav! in herb. Lamb. (DC. v. s. in h. Mus. Par.)
- 84. V. *CAPILLARIS* (Pers. ench. 1. p. 256.) fruticosa scandens glabriuscula, ramis angulatis, foliis ovatis basi inæqualibus subcordatis argutè serratis, serraturis exsertis glabris, stipulis oblongis acutis apice trifido-arietatis hinc setaceo-laceris, pedunculis folio brevioribus, sepalis acutis, petalo infero obovato subrotundo apiculato, calcare saccato-brevissimo, staminibus 2 inferioribus membranaceis terminalibus elongatis subulatis, appendicibus nectariferis dorsalibus latis basi rotundatis apice truncatis loculis brevioribus, seminibus fusco-punctatis. ξ in Regno Novo-Granatensi? V. stipularis. Cav. ic. 6. p. 21.* t. 531. f. 2. H. B. et Kunth nov. gen. am. 5. p. 372 non Swartz. (v. s. s.)
- 85. V. *SCANDENS* (H. B. et Kunth. nov. gen. am. 5. p. 371.* t. 493.) fruticosa scandens glabra, ramis subangulatis striatis, foliis cordato-ovatis crenatis glabris, stipulis oblongis dentato-ciliatis, pedunculis folii longitudine, sepalis acuminatis, petalo infero obovato oblongo, calcare saccato brevi, staminibus 2 inferioribus membranaceis terminalibus elongatis subulatis, appendicibus nectariferis dorsalibus oblongis obtusis loculis brevioribus, stigmate obtusiusculo. ξ in Peruvia prope Loxam, alt. 1060 hex.
- 86. V. *ARGUTA* (H. B. et Kunth. nov. gen. am. 5. p. 373.) fruticosa volubilis, ramis teretibus hirtellis, foliis profundè cordatis argutè serratis supernè puberulis subtus canis, petiolis villosis, stipulis lanceolatis setaceo-laceris, pedunculis folio brevioribus, sepalis acutis, petalo infero obovato mucronato, calcare lato obtusissimo laminâ vix dimidio breviorè, staminum 2 membranaceis longioribus obtusis reflexis, appendicibus nectariferis ovatis truncatis an-

therâ parùm brevioribus, seminibus glanduloso-punctatis fuscis. ζ in sylvis frigidis inter Gonzanam et Loxam, alt. 1060. hex.

87. *V. STIPULARIS* (Swartz fl. ind. occid. 3. p. 1956*) glabra, caulibus reptantibus, foliis ovato-lanceolatis utrinquè attenuatis, serraturis adpressis obtusis, stipulis oblongis acuminatis longè ciliato-laceris, calcare brevissimo, membranarum staminum a longioribus subulato-recurvis. ζ in Guadaloupâ et insulâ Sancti-Christophori. *V. persicariæfolia* Poir. dict. 8. p. 628. (v. s. sp.)

88. *V. DOMBEYANA* (DC. mss.) glabra, caulibus reptantibus, foliis ovalibus basi cuneatim attenuatis serratis, serraturis exsertis remotis, stipulis lanceolato-linearibus setaceo-dentatis, calcare brevissimo. ζ in Americâ meridionali. *V. repens* Domb. herb. An var. *V. stipularis* Swartz? (DC. v. s. sp. in herb. Mus. Par.)

89. *V. TRIFIDA* (Spreng. pug. 1. p. 22.) caule decumbente, foliis subtrifidis, lacinis lanceolatis basi attenuatis integerrimis, calycibus posticè vix appendiculatis, calcare calyce multò ampliore. \mathcal{L} ? hab. . . . †

90. *V?* *REDERACEA* (Labill! spec. Nov.-Holl. 1. p. 66. t. 91.*) caulis trunco abbreviato stolonifero, foliis subfasciculato-confertis reniformibus subrotundisve sinuato-dentatis, stipulis lanceolatis setoso-acuminatis, sepalis vix productis petalo infero subulato gibboso ampliato. \mathcal{L} in Novâ-Hollandiâ. An genus proprium? (v. s. sp.)

a. foliis basi subcuneatis. — In terrâ Van Diemen Labill. (v. s.)

β. *elatines* foliis reniformibus. — In Bot. - Bay. Brown *V. reniformis* Brown ined. An spec? Species distincta ex DC. (v. s.)

γ. *petiolaris* foliis reniformibus, petiolis pedunculisque longissimis nempè 6-9 poll. longis. — In Novâ-Hollandiâ et insulâ Van Diemen. *V. gracilis* Brown! ined. Species distincta ex DC. qui v. s. in h. Banks.

† *Species non satis notæ.*

91. *V. XIPHSILA* (Ledebour in Link enum. 1. p. 241.) caule repente, foliis cordato-subrotundis vix acuminatis glabris, lacinis calycinis basi productis barbatis. \mathcal{L} in Sibirîâ. Corolla pallidè cærulea, petalis 3 infimis lineis obscuris notatis.

92. *V. GIBBOSA* (Rafin. dec. pl. nov. am. sept. in litt. 1819.) caulescens glabra, foliis longè petiolatis cordato-deltoides crenatis obtusis subulatis pallidis, stipulis ovato-lanceolatis obtusis integris, petalis glabris, calcare gibboso brevissimo. — In mont. Alleghanis. Flores lutei.

93. *V. SERPYLLIFOLIA* (Humb. et Bonpl. in Willd. herb. ex Roem. et Schult. syst. 5. p. 391.) caule procumbente, foliis lineari-spathulatis integerrimis. \mathcal{L} in Americâ merid. — Cl. Kunthio ignota.

94. *V. TEUCRIIFOLIA* (Humb. et Bonpl. ex Willd. herb. in Roem. et Schult. syst. 5. p. 391.) foliis alternis ovatis serratis, caule fruticoso procumbente, pedunculis folio longioribus. ζ in Americâ merid. — Cl. Kunthio ignota.

95. *V. SCABRA* (Brown in florâ 1820. p. 469.) subcaulis, foliis cordatis acuminatis scabris, sepalis acutis, radice repente. \mathcal{L} ? circa Salisburgum.

96. *V. CRASSIUSCULA* (Bory ann. gen. 1820. v. 3. p. 16.) caulibus decumbentibus ebracteatis?, foliis alternis petiolatis ovatis oblongis integerrimis crassiusculis, floribus nudis longè pedunculatis. — In Sierrâ - Nevadâ Hispaniæ.

97. *V. SELKIRKII* (Pursh mss. ex Goldie edimb. phil. journ. 1822. p. 319.) foliis cordatis crenato-serratis pilosiusculis, petalis imberbibus, calcare longo crasso obtusissimo. — In montibus prope Montreal Americæ bor.

98. *V. PUNCTATA* (Lewis amer. journ. 5. n. 1. p. 48. non Willd. reliq.) caule breviusculo stolonifero simplici paucifolioso, foliis cordatis crassis glanduloso-punctatis, calcare longo in rostrum refracto. \mathcal{L} in Labrador Americæ bor.

99. *V. SPATHULATA* (Willd. rel. ex Rœm. et Schult. syst. 5. p. 353.) foliis lanceolâto-spathulatis fasciculatis subintegerrimis pubescentibus. $\frac{1}{2}$ in prov. Guilan ad mare Caspium.
100. *V. ALLEGHANENSIS* (Rœm. et Schult. syst. 5. p. 560.*) hirsuta, foliis ovatis et oblongo-cordatis subintegerrimis, petiolis latis membranaceis, petalis 2 lateralibus barbatis, calcare brevissimo saccato. $\frac{1}{2}$ in Alleghanis Pal Beauv. An var. *V. ovatae*?
101. *V. LABRADORICA* (Schrank denk. 1. bot. gesell. regensb. 11. p. 12.) caule erecto ramoso, foliis orbiculato-cordatis acuminatis, stipulis lanceolatis. — In Labradoriâ.
102. *V. ELONGATA* (Poir. dict. 8. p. 644.*) glabra, foliis ellipticis integerrimis petiolatis, floribus solitariis longè pedunculatis, caule debili, capsulis glaberrimis, seminibus rufescentibus globosis. — In Americâ ex herb. Desf.
103. *V. PRUNELLEIFOLIA* (H. B. et Kunth nov. gen. am. 5. p. 370.*) glaberrima, foliis ovato-oblongis acutis basi rotundatis obsolete cordatis crenatis, stipulis leviter ciliatis, calycibus acutis, petalis retusis, calcare subsaccato brevi rotundato, stylo supernè clavato incrassato, stigmatè laterali truncato. $\frac{1}{2}$ in altâ planitie Bogotensi, alt. 1370 hex. Flores violacei? *V. hirtæ* affinis?
104. *V. UMBRATICOLA* (H. B. et Kunth nov. gen. am. 5. p. 370.*) caulescens procumbens, foliis ovatis obtusiusculis basi in petiolum decurrentibus denticulatis suprâ glabris subtus petiolis pedunculisque tenuissimè hirtellis, stipulis dentato-ciliatis, calycibus linearibus acutis, petalo infero rotundato basi saccato-calcarato, calcare brevi rotundato, ceteris obovato-spathulatis, appendicibus staminum dorsalibus oblongis obtusis compressis loculis duplò brevioribus, ovario ovato glabro, ovulis 22 biseriatis, stylo cultriformi? stigmatè obtuso. $\frac{1}{2}$ in sylvis prope Real del monte Mexicanorum alt. 1426. hex. Flores violacei magn. *V. palustris*.
105. *V. CHAMÆDRIFOLIA* (Ruiz et Pav. in h. Lamb.) caulibus....., foliis ovatis obiter dentatis, petalis 2 lateralibus barbatis. — In Peruviâ. An *V. tenuifolia* Rœm. et Schult. syst. 5. p. 391? An var. *V. rubellæ* Cav?

IV. *SOLEA*. *Ging. mss.* — *Soleæ* spec. *Spreng. pug. rar.* 1. p. 22.

- Calycis sepala vix æqualia carinata? posticè non producta basi in pedicellum decurrentia (Nutt.) post anthesin reflexa. Petala subinæqualia, inferiùs ceteris paulò majus basi paulò gibbosum, cetera subæqualia, æstivatione convoluta? Stamina approximata 2 extùs ad basin glandulam nectaream gerentia? filamenta basi brevius unguiculata, unguibus latiusculis ovarium vix adæquantibus, antheram paulò altiùs gerentia. Stigma hamosum. — Herbæ pilosæ; caules virgati; folia alterna; pedunculi axillares gemini (uno sæpiùs abortivo) uniflori breves bibracteati.

1. *S. CONCOLOR* (*Ging. mss.*) $\frac{1}{2}$ in uliginosis Pennsylvaniae. *Viola concolor* Forster in lin. soc. trans. 6. p. 309.* t. 28. An *Solea stricta* Spreng. l. c? (*v. s. sp.*)

V. *POMBALIA*. *Vand. fasc. 7. t. 1.*

Calycis sepala maxima, margine aculeis callosis echinata, basi in pedicellum decurrentia. Petala inæqualia, 4 superiora breviora calycem vix excedentia, 5^{um} 2-3-plò longius basi subgibbum hinc stipitato-unguiculatum, limbo transversim latiore, marginibus involutis. Staminum filamenta basi longè unguiculata, unguibus linearibus ovarium adæquantibus hinc oblongo-dilatata, antheras altissimè gerentia; filamenta 2 dorso ad basin glandulam nectaream gerentia? lobi antherarum apice obtusi. Stylus rectus; stigma infundibuliforme. Capsula *Violæ*, sæpiùs

villosa. Cotyledones sæpius orbiculatæ, radiculæ teretiusculæ longitudine. — Herbæ perennes sæpius villosissimæ, radices crassiusculæ, duræ, tortuosæ; folia alterna; pedunculi axillares solitarii inarticulati 2-bracteolati; flores cernui, magni.

1. P. ITUBU (Ging. mss.) ♀ *Viola Itubu*, Aubl! guj. 2. p. 808. t. 318.* mala. *Ionidium Itubu* H. B. et K. nov. gen. am. 5. t. 496. (v. s. sp.)

α. caulibus hirsutissimis, floribus albis, seminibus albescentibus?, capsulis villosis. — in Guianâ.

β. caulibus hirsutissimis, flore cæruleo. — In Cayennâ. *Viola Itubu* var. flore cæruleo. Aubl. l. c.

γ. caulibus glabriusculis, flore albo, ovario villosa, seminibus nigris. In Brasiliâ. *Pombalia Ipecacuanha* Vand. fasc. 7. t. 1. mala. *Calceolar Lœffl. itin.* p. 184. n. 2. ? An spec ?? Au V. *Ipecacuanha*. Lin. mant. 484 ?

δ. caulibus villosis, flore albo?, seminibus.... capsulis glabriusculis. — In Brasiliâ. *Jonidium calceolaria* Mart. in litt. — An species? (v. s.)

VI. PIGEA. DC. mss.

*Calycis sepala inæqualia basi in pedicellum decurrentia. Petala inæqualia; inferius (labellum) ceteris quadruplò-quintuplòve majus, basi gibbosum, hinc unguiculatum et in limbum obovatum planum dilatatum, æstivatione convolutum? cetera calyce paulò longiora. Stamina filamenta à basi dilatata, antheras demissius gerentia; lobi antherarum sæpius apice in setas producti. Capsula sæpius trigona, 3-valvis; seminibus sæpius hinc angulatis. — Herbæ aut suffrutices, foliis sæpius alternis; flores erectiusculi, pedunculi solitarii, sæpius ad apicem ramulorum racemoso-congesti, bibracteolati, inarticulati? An rectius ad *Ionidii* sect. 3^{am} adnumerandæ?

1. P. FILIFORMIS (DC. mss.) caulibus indivisis, foliis alternis linearibus integerrimis, labello obovato integro, sepalis lanceolatis acutis. ☉? in Novâ-Hollandiâ circa Port-Jackson. (v. s. sp.)

2. PP BANKSIANA (Ging. mss.) caule erecto puberulo, foliis alternis oblongo-linearibus integerrimis marginibus revolutis supernè scabriusculis, stipulisque subulatis, labello calyce multò longiore, seminibus ellipsoideis albidis striatis? ♀ in Novâ-Cambriâ austr. *Viola angustifolia* Banks herb. (DC. v. s. in h. Banks.)

3. PP CALYCINA (DC. mss.) caulibus ramosis, foliis linearibus integerrimis, labello lanceolato acuto, sepalis ovatis acuminatis. ☉? in Novâ-Hollandiâ orâ occidentali. (DC. v. s. sp. in h. Mus. Par.)

4. PP MONOPETALA (Ging. mss.) caulibus ramosis, foliis linearibus integerrimis, labello cuneato-spathulato emarginato, sepalis ovatis. — *Ionidium monopetalum*. Rœm. et Schult. syst. 5. p. 400. †.

VII. IONIDIUM. — *Ionidii spec. Vent. malm. p. 27. — Soleæ spec. Spreng. pug. rar. p. 23.*

Calycis sepala parva inæqualia, basi in pedunculum decurrentia, non appendiculata, margine membranacea. Petala inæqualia; inferius ceteris 2-3-plò longius, basi carinato-concavum aut paulò gibbum, hinc unguiculatum, indè in limbum dilatatum, per æstivationem marginibus sæpius involutis. Stamina approximata; filamenta vix à basi oblongo-dilatata, antheram demissius gerentia, 2 anteriora sæpius ad basin anteriorem glandulâ nectariferâ basilari instructa. Capsula Violæ, non elastica? post maturitatem cum parte superiore pedunculi articulati de-

cidua? 1-6-rius g-sperma. Cotyledones sæpius reniformes; radícula breviuscula. — Herbæ aut suffrutices; folia nunc alterna, nunc opposita, aut inferiora opposita, superiora alterna; pedunculi solitarii, uniflori, suprâ medium 2-bracteolati, hinc articulati. Flores erectiusculi.

§. I. *Labello calyce duplò triplòve longiore stipitato.*

1. I? *ANOMALUM* (H. B. et Kunth nov. gen. an. 5. p. 381. t. 500.) puberulum, caule ramoso foliis ramescibus, alternis lanceolato-oblongis acuminatis serratis, suprâ glabris, subtùs canescentibus, stipulis latè ovatis acutis, sepalis ovatis acutis sericeo-pubescentibus, labello lanceolato longè attenuato calyce 4-quintuplòve longiore, petalis ceteris ovatis acutis. § in sylvis juxtâ Turbaco Novo-Granatensium. V. prunifolia Willd. rel. in Rœm. et Schult. syst. 5. p. 391. Flores antè folia, albi. Anthæræ lineari-oblongæ, 2 appendiculatæ, appendice uncinata descendentes apice villosæ in calcar geniculatum concavum intrantes. An genus?
2. I. *PARIETARIÆFOLIUM* (DC. mss.) caule ramoso pubescente, foliis alternis elliptico-ovatoe-lanceolatis acuminatis dentatis subpubescentibus discoloribus, stipulis subulatis ciliatis, sepalis acuminatis ciliatis, labelli limbo subrhomboideo ☉? in Americâ equinox.
 - a. *Houstoni*, foliis argutè serratis, caule hirsuto. — In Verà-Cruz., Peruvîa. *Viola parietariæfolia* herb. Banks. V. frutescens Ruiz et Pav. ined. (DC. v. s. sp. in herb. Banks et Lamb.)
 - β. *Berterii*, foliis subserratis, caule pubescente. — In Sanctâ-Marthâ. Bert. V. melanosperma Bertero! ined. Semina lenticulari-ovata atra nitida. Cotyledones reniformes. (v. s.)
3. I. *LEPTORRHIZUM* (DC. mss.) caule simplici aut parcè ramoso glabriusculo, foliis alternis glabris subglaucis ovatis acutis in petiolum breviter attenuatis dentatis, stipulis lineari-subulatis, sepalis acutissimis. (I)? in arenosis Malabarîæ, Tranquebarîæ. Rheed. h. mal. g. p. 109. t. 61. * Pluk. almag. t. 120. f. 8. A priore vix differt. (v. s. in h. Banks.)
4. I. *CAPENSE* (Rœm. et Schult. syst. 5. p. 393.) caule suffruticoso erecto, foliis alternis obovatis obsolete dentatis pubescentibus, stipulis subulatis ciliatis, sepalis acutis ciliatis. § ad cap. Bonæ-Spei. *Viola Capensis* Thunb. prod. 40. V. Massoni in h. Banks! (DC. v. s. sp. in h. Lin. ex Thunb.)
 - β. *Ovariense* foliis superioribus lanceolatis infernè glabriusculis subconfertis, sepalis pubescentibus, labello basi valdè gibboso. — in Oware Beauvois. (v. s.)
 - γ? *Burmanni* foliis superioribus oblongo-lanceolatis supernè glabriusculis, labello obovato basi gibboso extus pubescente. — in Indiâ orient. Burmann. — Ab I. heterophyllo tantùm differt sepalis ciliatis nec glabris, ab I. angustifolio seminibus ellipsoideis striatis. (v. s.)
5. I. *HETEROPHYLLUM* (Vent! malm. n. 27. in adn.) caule suffruticoso basi ramoso, foliis inferioribus obovatis, superioribus lineari-lanceolatis obsolete dentatis pubescentibus, stipulis subulatis rigidis, sepalis acuminatis glabris. § in Chinâ, Zeylonâ. *Polygala frutescens* Burmann fl. zeyl. 195. t. 35? (ex DC. adn. v. s. in h. Vent. et Burm.)
6. I. *BUXIFOLIUM* (Vent! malm. p. 27. in adn.) caule herbaceo diffuso, foliis alternis obovatis integris margine revolutis, stipulis subulatis rigidis, sepalis acuminatis glabris, labello basi gibboso apice subrotundo-truncato. ¶ in ipsâ Madagascariâ. Commerson. *Viola buxifolia* Poir. dict. 8. p. 646. * (v. s.)
7. I. *ENNEASPERMUM* (Vent. malm. p. 27.) caule suffruticoso basi ramosissimo, foliis alternis lanceolatis glabriusculis, stipulis subulatis rigidis, sepalis acuminatis glabris, labello....., seminibus nitidis. § in Indiâ orient. (Vent.)

a. Malabaricum, caulibus decumbentibus, foliis parcè serratis. — In Malabar. Rheed. h. malab. g. p. 117. t. 60.* (DC. v. s. sp. in h. Banks et h. Mus. Par.)

β. Zeylanicum, caulibus adscendentibus, foliis integriusculis. — In Zeylonà. *Viola enneasperma* Lin. sp. 1327. fl. zeyl. n. 317.* (DC. v. s. sp. in h. Banks.)

3. *I. LINARIFOLIUM* (Vahl! ecl. am. 2. p. 18.*) caule erecto, foliis caulinis alternis oblongis utrinquè attenuatis, stipulis subulatis rigidis. ☉? in insulis Caribæis. *Viola linariæfolia* Poir. dict. 8. p. 648*? An var. *I. stricti*? (v. s.)

9. *I. STRICTUM* (Vent! malm. n. 27. in adnot.) caulibus erectis imà basi ramosis puberulis, foliis inferioribus oppositis oblongis lanceolatisve basi attenuatis apice obtusioribus, stipulis subulatis rigidis, sepalis acuminatis glabris, labello subrotundo, gibbo nullo, capsulà subrotundo-3-gonà, seminibus ovato-subrotundis nigricantibus? 2/ in Santo-Domingo, Novà-Hispanià. *V. stricta* Poir! dict. 8. p. 648.* (DC. v. s.)

β. ramis erectis, foliis elongatis, imis obtusis, basi longè attenuatis glabriusculis. — in Guadaloupà. Bertero. (v. s.)

10. *I. ANGUSTIFOLIUM* (H. B. et Kunth nov. gen. am. 5. p. 377.*) caule lignoso elongato debili supernè ramoso glabro, foliis oppositis lineari-lanceolatis apice angustatis remotè serrulatis glabris, stipulis minutis, pedunculis in ramis racemoso-confertis glabris, fructiferis folio 4-quintuplòve brevioribus, sepalis lanceolatis acuminatis, capsulis subrotundo-ovatis trigonis 3-4-spermis. 3 locis humidis propè Carichana in sylvis Orinocensium.

§. 2. *Labello calyce vix duplò longiore subsessili.*

11. *I. THESIIFOLIUM* (DC. mss.) caule erecto glabro, foliis alternis angustis longissimis glabris integerrimis, stipulis sepalisque subulatis, petalis calyce vix longioribus. 2/ in Senegal. Adanson. *Viola thesiifolia* Poir! dict. 8. p. 649.* (DC. v. s. sp. in herb. Juss.)

12. *I. LINIFOLIUM* (DC. mss.) caule erecto puberulo, foliis alternis lineari-angustis integris glabris, stipulis subulatis. ☉? in Madagascar. Commerson. *Viola linifolia* Poir! dict. 8. p. 647.* (DC. v. s. sp. in h. Mus. Par.)

13. *I. POLYGALÆFOLIUM* (Vent! malm. p. et t. 27.*) caulibus ramosis diffusis procumbentibus, ramis puberulis, folio oppositis lanceolatis integriusculis, stipulis lanceolatis folio dimidiò brevioribus, sepalis ovato-oblongis acutis pubescentibus, labello spathulato apice rotundato calycem superante, squamulis nectareis carnosis, capsulis subrotundis trigonis 1-5-spermis, seminibus subglobosis lateribus compressiusculis nitidis nigris, cotyledonibus reniformibus. 3 in Novà-Hispanià. Mocq. et Sessé. prope Mexico. H. B. et Kunth nov. gen. am. 5. p. 376.* t. 496. Solea verticillata Spreng. in Schrad. journ. bot. 1800. v. 2. p. 190. t. 6.* *Viola verticillata* Orteg. dec. 4. p. 50. (v. s. a.)

14. *I. GRACILE* (Mocq. et Sessé fl. mex. ined. ic.) caulibus suberectis simpliciusculis, foliis alternis oblongis integris, stipulis ovato-lanceolatis folio dimidiò brevioribus, petalis calyce vix longioribus, infimo vix majore, seminibus ovatis nigris. 2/ in Mexico. An var. *I. polygalæfolii*?

15. *I. VISCIDULUM* (H. B. et Kunth nov. gen. am. 5. p. 377.*) caule lignoso erecto ramoso viscoso, foliis oppositis lanceolatis remotè serrulatis glabris, stipulis subulatis minutis, sepalis ovato-lanceolatis glabris, labello longè unguiculato latissimè ovato acuto basi dilatato concavo, petalorum lateraliùm ungue petali superioris longitudine, staminibus 2 anterioribus basi glandulà instructis, membranis terminalibus acutis. 3 in arenosis humidis propè Angosturam Guyanensium, ad ripam Orinoci. Flores suaveolentes albi, labello basi flavo maculato.

16. *I. RIPARIUM* (H. B. et Kunth nov. gen. am. 5. p. 378.*) caule herbaceo erecto subramoso bifariam pubescente, foliis inferioribus oppositis lanceo-

latis crenato-serratis glabris subciliatis, stipulis subulatis minutis, sepalis lanceolatis acuminatis subulatis ciliatis, labello unguiculato ovato subrotundo basi gaccato concavo, petalis lateralibus dimidio brevioribus, omnibus versùs basim ciliatis, staminum filamentis subnullis 2 anterioribus glandulâ squamulâformi instructis, membranâ terminalibus acutis, capsulis ovatis trigonis glabris, 3-6-spermis valvis navicularibus acutis, seminibus ovato-globosis subcompressis nitidis fuscis. ① in regno Novo-Granatensi propè Angostura de Carare ad ripam fluminis Magdalene alt. 120 hex. I. attenuatum. Willd. herb. ex Rœm. et Schult. syst. 5. p. 402. Flores cœrulei.

17. *I. VERBENACEUM* (H. B. et Kunth nov. gen. am. 5. p. 379. t. 497.) caule herbaceo erecto subranoso pubescente, foliis alternis ovato-ellipticis acutis basi in petiolum decurrentibus crenato-serratis utrinquè hirtellis, stipulis lineari-subulatis puberulis petiolo sublongioribus, sepalis lanceolatis acuminatis ciliatis, labello longè unguiculato subrotundo elliptico basi 2 gibbis aucto calyce duplò triplòve longiore, petalis lateralibus calycem parùm longioribus, staminum filamentis brevibus 2 inferioribus suprà basim glandulâ gibbis, membranâ terminalibus emarginatis. ② Colitur in horto Mexicano. Flores pallidè cœrulei.

18. *I. CIRCEOIDES* (H. B. et Kunth nov. gen. am. 5. p. 379. t. 498.) caule herbaceo erecto simplici bifariam puberulo, foliis oppositis ovatis acuminatis basi rotundatis serratis glabris, stipulis linearibus falcatis petiolum longè quantibus glabris, sepalis ovato-lanceolatis acuminatis glabris, labello longè unguiculato subrotundo - ovato obtuso calyce quintuplò longiore, petalis lateralibus inferiore triplò brevioribus oblongis falcatis medio angustatis, hinc infernè lobulo obtuso acutis, staminum filamentis brevibus, 2 inferioribus ad basim glandulâ notatis, membranâ terminalibus emarginatis, capsulis subrotundo-ovatis trigonis calyce triplò longioribus glabris 6-spermis, valvis oblongis, seminibus (immaturis?) globoso-ovatis compressiusculis carunculatis. ① propè Goyaquil locis umbrosis, ad littora maris Pacifici. Willd. herb. ex Rœm. et Schult. syst. 5. p. 402. Flores violacei. (v. s.)

19. *I. GLUTINOSUM* (Vent! malm. n. 27. in adnot.) caulibus suberectis pubescentibus viscosis, foliis ovato-lanceolatis in petiolum longè attenuatis argutè dentatis, stipulis subulatis petiolo brevioribus, labello obcordato. ② in Buenos-Aires. V. glutinosa Poir! dict. 8. p. 647.* Calyx in sicco viridis nec ut in sequente, cui valdè affine, violaceus. (DC. v. s.)

20. *I. PARVIFLORUM* (Vent! malm. p. 27. in adn.) fruticosum ramosum diffusum, ramis elongatis bifariam puberulis, foliis alternis ovatis serratis in petiolum breviter attenuatis, stipulis subulatis petioli vix longitudine, pedunculis glabris folium superantibus, sepalis ovato-lanceolatis acutis, labello calycem triplò superante ovato bilobo, lobis rotundatis patentibus, squamulis nectareis subclavatis, capsulis subrotundo-trigonis calyce triplò longioribus 3-5-spermis. ③ in Americæ merid. regionibus calidioribus propè Santa-Fe-de-Bogota. H. B. et Kunth nov. gen. am. 5. p. 375. Viola parviflora Lin. f. suppl. 396*. (DC. v. s.)

④? ramis longissimis. Viola filiformis Ruiz et Pav. ined. (v. s. s.)

21. *I. MICROPHYLLUM* (H. B. et Kunth nov. gen. am. 5. p. 374. t. 495.) fruticosum ramosum diffusum, ramis bifariam pubescentibus, foliis oppositis elliptico-oblongis acutis serratis pilosiusculis ciliatis, pedunculis folio triplò longioribus, hinc lineâ petiulâ notatis, stipulis lanceolatis acuminatis petiolo triplò longioribus, sepalis ovato-oblongis acutis, labello calycem duplò superante subpanduræformi, squamulis nectareis subovatis, capsulis subrotundo-ellipticis trigonis 3-6-spermis. ⑤ inter saxa propè Lactacunga Quitenium, alt. 1480. hex. Viola microcephala Bonpl! ined. V. microphylla Willd. herb. V. parvifolia Rœm. et Schult. syst. 5. p. 391. (v. s.)

†. *Species non satis notæ.*

22. *IP CLAYTONIOIDES* (Rœm. et Schult. syst. 5. p. 402.) caulibus monophyllis perfoliatis. Patria ign.
23. *IP ERÆCTUM* (Ging. mss.) caule stricto filiformi scabriusculo, foliis linearibus mucronatis remotè denticulatis strictis, stipulis lanceolatis ciliatis adpressis, floribus axillaribus solitariis posticè æqualibus nutantibus. — in Indiâ orient. *Viola erecta* Roth nov. sp. 165. Flores exigui.
24. *IP SUFFRUTICOSUM* (Ging. mss.) caule procumbente scabriusculo, foliis elliptico-lanceolatis serratis subpubescentibus, stipulis lanceolatis pilosis demùm patulis aculeiformibus, floribus axillaribus posticè æqualibus nutantibus. 3? in Indiâ orient. *Viola suffruticosa* Roth nov. spec. 165.*
25. *IP FRUTESCENS* (Ging. mss.) caule adscendente scabriusculo basi fruticoso, foliis oblongo-lanceolatis mucronatis glabris subciliatis serratis, stipulis setaceis erectis pilosis, floribus axillaribus solitariis posticè æqualibus nutantibus. 3? in Indiâ orient. *Viola frutescens* Roth nov. spec. 167.
26. *IP BREVICAULE* (Mart! in litt.) caule abbreviato adscendente, foliis confertissimis subsessilibus alternis ovato-lanceolatis acutis serratis pubescentibus. — In Brasiliâ. (v. s. sine fl.)
27. *I. LINEATUM* (Ging. mss.) ramis procumbentibus, foliis oppositis ovatis lineatis petiolatis subtus pubescentibus, stipulis subulatis. 3? in Cubâ. *Viola lineata* Orteg. dec. 4. p. 49.* Petalum inferum seu labellum violaceum ungue albedo, lateralìa violacea, superiora alba.
- 3? foliis imis obovatis glabrisculis. — in Cubâ. *V. lineata* herb. hort. monsp.
28. *IP CALCEOLARIUM* (Ging. mss.) caule ramoso bifariam piloso, foliis oppositis elliptico-lanceolatis in petiolum attenuatis subamplexicaulibus, stipulis subulatis, sepalis aristato-acuminatis glabris. ① in Mexico. Moc. et Sessé. *Viola calceolaria* Moc. et Sessé fl. mex. ined. non Poir. An *V. calcaria* labello obovato retuso Lessl. itin. p. 183. n. 1?
29. *IP LONGIFOLIUM* (Moc. et Sessé fl. mex. ic. ined. non Rœm. et Schult.) caulibus simpliciusculis, foliis oppositis lanceolato-linearibus rariter serratis acutissimis, stipulis lanceolatis petiolo sublongioribus, pedunculis folio triplò longioribus. 2? in Mexico.

VIII. HYBANTHUS. Jacq. am. 77. H. B. et Kunth nov. gen. am. 5. p. 385.*

Calycis sejala inæqualia basi in pedicellum decurrentia non appendiculata. Petala inæqualia, inferius ceteris longius basi saccatum, medium canaliculatum apice in limbum bilobum dilatatum, reliquia breviora 3-nervia. Stamina oblonga basi inter se in disco connata; antheræ demissiùs insertæ, in duobus inferioribus loculi contigui oblitterati; glandula nectarea magna conchæformis ad basin staminum duorum inferiorum in gibbo petali quinti locata. Capsula obovata oligosperma. — Frutices inelegantes, facie Randiæ, sæpiùs aculeati. Folia sparsa secus ramos alterna aut secus caules subfasciculata. Pedunculi alares vel laterales solitarii aut plures congesti apice bifidi. Flores pedicellati subramosi albidì, pedicellis basi bracteatis suprâ basim articulatis.

1. *H. HAVANENSIS* (H. B. et Kunth l. c. p. 385.* t. 494.) caule erecto aculeato ramoso, foliis oblongis remotè serratis, floribus albidis, petalo infimo apice bilobo, pedunculis bifidis subracemosis. 3? prope Havanam ins. Cubæ.
- α. *Jacquinianus*, foliis apice emarginatis basi cuneato-attenuatis, sepalis ovatis 3 inferioribus basi gibbis, floribus minoribus vix oculis nudis examinandis, petalis 4-ovatis obtusis, quinto oblongo medio attenuato paulò longiore apice bifido, stylo subulato. — In montibus sylvaticis ex Jacq. l. c. *Ionidium Jacquinianum* Rœm. et Schult. syst. 5. p. 397.*

- β. *Humboldtianus*, foliis apice integris vix petiolatis, stipulis ovatis obtusis, floribus racemoso-fasciculatis majoribus (magnitudine floris *Ionidi polygalæfolii*) sepalis oblongis obtusis glabris, petalo infimo calyce triplò longiore basi saccato concavo hinc canaliculato, limbo dilatato bilobo, lateralibus oblongis obtusis obsolete trilobis calyce duplò longioribus, superioribus vix brevioribus obtusis, antheris subcordatis, ovario subrotundo ovato glabro g-spermo, stylo geniculato apice subuncinato. (H. B. et K. l. c.)
2. H? *MEXICANUS* (Ging. mss.) caule aculeato, foliis oblongis, petalo infimo apice acuminato, pedunculis unifloris fasciculatis? ♀? in Mexico. Moc. et Sessé fl. mex. ined. icon.

Trib. II. *ALSODINEÆ*. Brown cong. p. 21.

Petala inter se æqualia. Stamina sæpiùs aut basi coalita, aut basi internæ urceoli inter petala et stamina elevati (tori expansi?) adnata.

IX. *CONOHORIA*. DC. mss. Conohoria, Passoura et Riana. Aubl. *Guian.* 1. p. 237. 239. t. 94 et 95. et app. p. 21. t. 380.

Calycis sepala imbricata. Petala æstivatione convolutivâ? Stamina libera approximata, filamenta breviter unguiculata, hinc in ligulam dilatata, extus sæpiùs lamellâ erectâ appendiculatâ, antheras paululò altius gerentia; lobi antheræ ad apicem setosi. Stylus flexuoso-subulatus. Stigma obtusum. Ovarium villosum. Capsula 3-valvis, valvis oligospermis. Semina, ex Aublet, globosa. — Frutices. Folia opposita. Flores racemosi. Racemi axillares aut terminales, pedicelli brevissimi erecti uniflori ad basin 1-bracteati, bracteolæ 2 secùs pedicellum; flores parvi; stipulæ deciduæ.

1. C? *RIANA* (DC. mss.) foliis ovatis serratis apice longè acuminatis, squamis staminum acuminatis, racemis spiciformibus. ♀ in *Guianæ* sylvis *Araurensibus*. Riana *Guianensis*. Aubl! *guian.* 1. p. 237. t. 94.* (v. s. sp.)
2. C. *ULMIFOLIA* (H. B. et Kunth nov. gen. am. 5. p. 387. t. 491.) foliis geminis vel ternis in apice ramulorum suboppositis oblongis acuminatis dentatis glabris nervo venisque pinnatis supra nitidis, racemis terminalibus solitariis ramosis folio 2-triplòve brevioribus. ♀ in regno *Novo-Granatensi*, ad ripam fluminis *Magdalenæ*. Sepala lanceolata ciliata; petala calyce 2-triplòve longiora oblonga obtusa venosa, venis subparallelis planis albidis, membranis terminalibus staminum apice rotundatis, squamis staminum inferiorum nullis? Ovarium triquetrum pilosum 3-spermum. Capsulæ obovato-turbinate triquetre apice pilosiusculæ aut glabræ fuscæ, seminibus subglobosis lateribus vix compressiusculis lævibus glabris hinc ab hilo ad foveolam oppositam sulco longitudinali notatis.
3. C. *PASSOURA* (DC. mss.) foliis ovatis integriusculis apice breviter acuminatis, sepalis lanceolatis acutis, squamis staminum acutis. ♀ in *Guianæ* sylvis. Conohoria flavescens Aubl! *guian.* 1. p. 239.* t. 93. Passoura Aubl. l. c. suppl. p. 21.* t. 380. (v. s. sp.)

X. *RINOREA*. Aubl. *guian.* 1. p. 235. t. 93.

Omnia *Conohoriæ*, sed staminum filamenta à basi in ligulam acuminatam dilatata, nec unguiculata, antheras dimissiùs gerentia approximata. — Arbor mediocris. Folia alterna stipulata, stipulis deciduis; flores racemosi, racemi axillares terminalesve laxè paniculati, pedicelli cernui uniflori ad basin 1-bracteati, versus medium 2-bracteolati, hinc *articulati*. Flores parvi. Fructus ignotus.

1. *R. GUIANENSIS* (Aubl! guian. 1. p. 235. t. 93.*) floribus racemosis, foliis dentatis. § in Guianæ locis cultis territorii Caux. (DC. v. s. sp. in h. Banks.)
2. *R? INTEGRIFOLIA* (Ging. mss.) pedunculis solitariis geminive unifloris, foliis integris. — in Brasiliâ. Conohoria alternifolia Spreng. neu. entd. 2. p. 151.

XI. *ALSOIDEIA*. Pet. Th. hist. veg. afr. 2. p. 55. t. 17 et 18. nov. gen. Madagasc. n. 65.

Calycis sepala acuta imbricata; 3 exteriora, 2. interiora. Petala alternâ calyce longiora, æstivatione contorta. Stamina petalis alterna, filamenta basi in urceolum ovarium cingentem soluta, hinc in ligulas non unguiculatas dilatata, antheras ad basin cujusque ligulæ gerentia, ovarium superantia, lobi antherarum sæpius apice in appendices setosas producti. Urceolus extus simplex aut cingulatus, vel appendicibus variis auctus. Ovarium simplex. Stylus clavatus. Capsula calyce et corollâ persistentibus induviata obversè turbinata obtusè 3-gona. Semina 1-2 ad quamque valvulam. Cotyl. orbiculatæ (r. r.)—Frutices vel arbores Madagascarienses; folia sæpius alterna penninervia; stipulæ parvæ deciduæ. Flores racemosi, parvi; racemi axillares et terminales; pedicelli bracteati articulati.

§. 1. *Urceolis simplicibus.*

1. *A. PAUCIFLORA* (Pet. Th. l. c. p. 57. f. 17.*) urceolo staminum simplici, foliis cuneatis breviter petiolatis, floribus paucis subcorymbosis, pedicellis reflexis. § in umbrosis insulæ Madagascar.
2. *A. ARBOREA* (Pet. Th. l. c. p. 57.) urceolo staminum simplici, foliis longè petiolatis deflexis, floribus paniculatis. § in Madagascar.

§. 2. *Urceolis cingulatis.*

3. *A. ANGUSTIFOLIA* (Pet. Th. l. c. p. 57. t. 17. f. 1.*) urceolo staminum cingulato, foliis oblongo-lanceolatis dentatis, racemis spicatis. § in umbrosis Madagascariensibus.
4. *A. LATIFOLIA* (Pet. Th. l. c. p. 57. t. 18. f. 2.) urceolo staminum cingulato, foliis ovatis obtusè acuminatis, racemis glabris densis. § in Madagascar.
5. *A. PUBESCENS* (Pet. Th. l. c. p. 57. t. 18. f. 3.) urceolo staminum cingulato, foliis ovalibus basi cuneatis serratis, racemis laxis pubescentibus. § in Madagascar.
6. *A? RACEMOSA* (Mart! in litt.) urceolo staminum cingulato fimbriato, foliis oppositis elliptico-lanceolatis acuminatis integris. § in Brasiliâ. Habitus Conohoriæ et imprimis *C. Rianæ*, sed stamina Alsodeiæ. (v. s.)

XII. *CERANTHERA*. Beauv. fl. ow. 2. p. 11. t. 65. — *Passalia Banks herb. ex Brown cong. p. 21.*

Calycis sepala acuta. Cor. . . petala lanceolata calyce duplò longiora. Urceolus dentatus basi cum petalis coalescens inter petala et stamina positus. Stam: filamenta basi unguiculata, hinc in squamam petaliformem dilatata, antheras ad basin squamæ altius gerentia, lobi antherarum setosi; ungues filamentorum basi cum urceolo coaliti, singuli apice liberi. Ovarium ovatum. Capsula ign. . . Frutices. Folia alterna Flores racemosi spicati, pedicelli bracteati. (Car. ex Beauv. l. c.)

Obs. Ab icone Beauv. l. c. videtur staminum structura eadem ac in Conohoriâ; sed stamina ad parietem internam urceoli affixa nec libera; differt tamen vix ac ne vix ab Alsodeiâ (ut Rinorea à Conohoriâ) filamentis unguiculatis nec à basi dilatatis, antheris altius (nec demissiùs) insertis.

1. *C. DENTATA* (Beauv! fl. ow. et ben. 2. p. 11. t. 65. sub nom. *C. dentulæ*.) foliis lanceolato-ovatis dentatis, petalis lanceolato-ovatis. ζ in Buonopozo. regni Owariensis. Passalia Soland. in herb. Banks et in Sierra-Leone detecta fortè non differt, ex R. Brown. l. c. (DC. v. s. in h. Deless.)
2. *C. SUBINTEGRIFOLIA* (Beauv! l. c. t. 66.) foliis lanceolato-oblongis rarè margine sinuosis, petalis ovatis. ζ circa Owariensem urbem (DC. v. s. in h. Deless.)

XIII. PENTALOBA. *Lour. coch. 1. p. 191.*

Sepala 5 lanceolata erecta pilosa. Petala 5 lanceolata apice subreflexa in campanulam conniventia. Nectarium 5-dentatum erectum. Filamenta 5 filiformia, planiuscula, incisuris nectarii insistentia, corollæ subæqualia. Ovarium pilosum, stylus brevis pilosus, stigma simplex. Bacca subrotunda 5-loba 1-locularis 5 sperma, semina 5 ovata (Lour. l. c.) Omnia Alsodeiæ, etiam quoad placentarum parietalium numerum ex cl. Brown cong. p. 22. — Arbor mediocris, folia alterna, flores pallidi sessiles congesti (Lour.) An Alsodeiæ congener?

1. *P. sessilis* (Lour. fl. coch. 1. p. 192.) ζ in montibus Cochinchinæ.

XIV. LAVRADIA. *Vellozo et Vand. in Ræm. script. lus. et br. p. 88. t. 6. f. 6.*

Calycis sepala 5 ovato-lanceolata persistentia. Petala 5 ovata acuminata calyce majora. Filamenta 0 vel brevissima. Urceolus conicus truncatus suprâ stamina et pistillum cucullatus (Vand. l. c.) Aut potius stamina Violæ coalita, ut in genere *Hymenantha* cuius membranæ approximatae pro cucullo proprio habitæ fuerunt. Ovarium oblongum. Stylus filiformis urceolo vix longior. Stigma obtusum. Capsula oblonga 1-locularis 3-valvis ∞ sperma. (Vell.) Cl. Brown dubitanter tamen ad *Alsodeiâ* refert.

1. *L. VELLOZIANA* (Stend. nom. phan. 466.) sepalis integris, stylo deciduo (ex icon) ζ in Brasiliâ. Folia lanceolata serrata. Caulis teres. Racemi terminales.
2. *L. MONTANA* (Mart! ined. DC. mss.) sepalis glanduloso-serratis, stylo ad apicem valvulæ cuiusdam in fructu persistente. ζ in Brasiliâ. Rami teretes. Stipulæ setoso-pinnatipartitæ erectæ. Folia oblonga subcuneata obtusa cum mucrone cartilagineo-serrata. Racemus terminalis compositus. Flores parvuli. (v. s. comm. à cl. Mart.)

XV. PHYSIPHORA. *Sol. in Banks herb. ex R. Brown cong. p. 21.*

Ab *Alsodeiâ* aut *Gerantherâ* differt tantum filamentis basi læviter connexis, capsulâque membranaceâ inflatâ, ex Brown. l. c.

Frutex ramosus; folia alterna ad apicem ramulorum subconferta; stipulæ deciduæ; pedunculi communes (laterales?) 1-5 ex iisdem punctis, (mixti cum ramis junioribus) laxè paniculati; pedicelli ex axillis bracteolarum infrâ medium 2-bracteolati hinc articulati.

1. *P. LÆVIGATA* (Sol! in Banks herb.) ζ in Brasiliâ. *Alsodeia Physiphora* Brown! in Banks herb. (DC. v. s. in Banks herb.)

XVI. HYMENANTHERA. *Banks herb. ex Brown cong. p. 23.*

Calycis sepala 5 imbricata. Petala 5 alterna ovato-acuminata demùm reflexa calyce longiora, æstivatione obliquè imbricatâ (Brown.). Stamina structurâ ad *Violam* accedentia, sed basi coalita in discum *monadelphum*; squamis totidem iis dorso oppositis. Stylus brevissi-

mus. Stigmata 2 acuta. Capsula subbaccata (in sicco rugosa aut venosoreticulata) tenuis ovata (unilocularis monosperma?) 2-locularis, loculis 1-spermis (sec. Brown.), calyce petalis staminibusque induviata. Semina capsulæ conformia illamque omnino replentia, ad ejus apicem ē placenta nerviformi (ut in Violâ) pendula. Seminis structura inter Violaceas et Polygaleas, ex Brown, media.

Frutices ramosi; folia nunc solitaria et alterna, nunc subfasciculata coriacea; flores axillares parvi; pedunculi solitarii, 1-flori basi 2-bracteati.

1. *H. ANGUSTIFOLIA* (Brown! in h. Banks.) foliis linearibus integerrimis. 5 in insul. Van Diemen ad portum Dalrymple. (DC. v. s. sp. in h. Banks.)
2. *H. DENTATA* (Brown! in herb. Banks.) foliis oblongis denticulatis. 5 in Novâ-Hollandiâ prope Port-Jakson. (DC. v. s. sp. in herb. Banks.)

Trib. III. *SAUVAGEÆ*. Ging. mss.

Capsulæ dehiscencia septicida; nempè suturæ in axi placentarum sitæ et ideo valvæ margine seminferæ dictæ. Stamina 5 fertilia petalis opposita, libera, filamenta non dilatata ultrâ antheras non producta. Squamæ 5 petaliformes cum staminibus alternantes. — Tribus media inter Violaceas et Frankeniaceas.

XVII. *SAUVAGESIA*. Jacq. pl. Amer. p. 77. — *Sauvagea*. Neck. elem. n. 1188.

Calycis sepala 5 concava persistentia, 2 exteriora, 3 interiora, æstivatione imbricata. Petala 5 sepalis alterna hypogyna decidua æstivatione sæpius contorta; corona filamentosa antè petala sita et his basi adnata, filamentis ∞ capillaribus, apice clavæformibus. Stamina 5 fertilia petalis opposita cum squamis 5 alternantia, stamina sterilia secund. Gaertn. fil., petala interna sec. Saint-Hil.) scariosa, seriùs decidua. Staminum filamenta subulata. Antheræ adnatæ, oblongæ (extrorsæ secundum Saint-Hil.) rimâ laterali duplici versùs apicem dehiscentes. Ovarium substipitatum; Stylo stigmatæque simplicibus. Capsula ovato-triquetra 1-locularis, 3-valvis; valvulis basi intus marginibus involutis, apice valvatim adplicatis dehiscentibusque; placentæ 3 duplices bipartitæ, parietales, sepalis 3 (interioribus) suprâ torum symmetricè dispositis oppositæ. Semina minutula, ovata, scrobiculata, prope basin capsulæ ope funiculi brevissimi, parti valvularum involutæ adfixa. Embryo subcylindricus ex Gaertn. fil.

Herbæ annuæ tenellæ; folia alterna; stipulæ membranaceæ sæpiùs pectinatim ciliatæ; pedunculi axillares uniflori solitarii aut bini, aut racemi multiflori axillares; pedunculi aut pedicelli, sæpiùs imo articulati, sæpiùs nudi, interdum bracteati.

§. 1. *Pedunculis 1-2 axillaribus unifloris.*

1. *S. GEMINIFLORA* (fl. mex. ic. ined.) caule ramoso erecto, foliis lanceolatis serratis, sepalis acuminatis, pedicellis geminis, petalis obtusis. ☉? in Mexico.
2. *S. ERECTA* (Lin. sp. ed. 2. p. 294. non Spreng.) caule ramoso procumbente erectoque, foliis lanceolatis serratis, pedicellis 1-2, sepalis aristato-acuminatis, petalis obovatis apiculatis. ☉ in Amer. æquinoctiali. *S. erecta* H. B. et Kunth nov. gen. am. 5. p. 389.* (v. s. s.)

a. foliis distantibus pollicaribus. *S. Adyma* Aubl. guian. t. 100. f. A. — in Cayenâ et Guyanâ.

ℓ. Foliis approximatis sesqui-pollicaribus. S. erecta Aubl. guian. t. 100. f. B.

γ. Foliis confertis vix semi-pollicaribus.

3. S. *nutans* (Pers. ench. 1. p. 253.) caule simplici foliis stipulisque omnino tecto, foliis lanceolatis serratis, pedicellis calyce duplò longioribus incurvolutantibus, sepalis acuminatis. ☉ in Madagascar. An S. erecta Jacq. amer. t. 77. f. 3? An var. S. erectæ? (v. s. sp.)

4. S. *tenella* (Lam. ill. 2. n. 2769.*†) caule ramoso, foliis lineari-lanceolatis subserulatis, imis confertis, summis distantibus, pedicellis calycis vix longitudine erectis, sepalis obtusis. ☉ in Cayennâ Patris. Poir. dict. 8. p. 669.* (v. s. sp.)

§. 2. *Floribus racemosis, racemis axillaribus.*

5. S. *ericoïdes* (Ging. mss.) caule simplici, foliis congestis lineari-lanceolatis margine revolutis integerrimis, stipulis setoso-subulatis, racemis dichotomis, sepalis ovatis acuminatis petalis triplò brevioribus. ☉ 2? in Brasiliâ. S. pendula Mart. ex Steven. in litt. (v. s. sp.)

† *Species dubiæ.*

6. S? *fruticosa* (Mart! in litt.) caule ramoso stipulis lanceolatis setoso-pennatis persistentibus obtecto, foliis ad apicem ramulorum confertis linearibus acutis marginibus revolutis argutè serrulatis. 5? in Brasiliâ. An S. erecta Spreng. nen. entd. 1. p. 296. excl. syn? (v. s. sine fl.)

7. S?? *salicifolia* (Ging. mss.) caulibus ramosis adscendentibus, foliis lanceolatis marginibus revolutis integerrimis, stipulis aristato-subulatis integris, floribus solitariis subsessilibus. 5? in Brasiliâ. (v. s. in herb. Moricand.) — An genus?

GENERA VIOLARIIS AFFINIA.

XVIII PIPAREA. Aubl. guian. 2. suppl. p. 31. t. 386. Gaertn. fil. carp. 3. p. 231. t. 224.

Sepala 5 æqualia persistentia demùm reflexa. Petala Filamenta? 10-15? subulata persistentia erecta calyce breviora glabra. Appendices totidem cum iis alternantes oblongæ hirsutissimæ omnes inter se basi coalitæ, capsulam cingentes. Capsula ovato-triquetra 1-locularis 3-valvis. Stylus filiformis, stigmatè tripartito (Gaertn. fil.) Capsula 3-valvis, 1-locularis, ab apice lateraliter dehiscens, intus pube densâ brevi fuscâ velutinâ tecta, valvis mediò semiuiferis. Semina ad quamque valvulam solitaria, globosa velutina (2 plerumque abortiva.) (car ex spec. sicco.) ad ventrem cicatriculâ lineari deorsum deliquescente umbilicali albâ notata. (Gaertn. l. c.)

Frutices arboresve. Folia alterna, vix petiolata, rigida, penminervia; stipulæ 2 caducæ. Flores axillares, brevissimi, pedicellati; pedicellis 1-floris, solitariis aut numerosis, articulatis?

Obs. An hoc genus verè pentandrum? an forsan ad Tiliaceas pertinet? An P. dentata Aubl. cum P. multiflorâ Gaertn. fil. congener?

1. P. *dentata* (Aubl. guian. 2. p. 31. t. 386.*) floribus solitariis geminisve, foliis ellipticis acuminatis subtus rufo-velutinis, capsulâ acuminatâ glabriusculâ 5. in sylvis Cayennæ prope Montem Serpent. (v. s. sp.)
2. P? *multiflora* (Gaertn. f. carp. 3. p. 231. t. 224. f. 1.*) floribus plurimis pedicellatis, foliis oblongis acuminatis subtus glabris, capsulâ subobtusâ velutinâ 5. in Cayennâ Patris. (v. s. sp.)

(DE GINGINS.)

Ordo XVII. DROSERACEÆ. DC. *theor.* 214. — Drosereæ.
Sal. parad. n. 95.

Sepala 5 persistentia, æqualia, per æstivationem imbricata. Petala 5, distincta, aut in Romanzowiâ corollam gamopetalam constituentia, hypogyna, sepalis alterna, sæpè marescentia. Stam. marescenti-persistentia, nunc petalis alterna et iis numero æqualia, nunc petalis dupla, tripla quadrupla, libera. Antheræ 2-loculares birimosæ. Ovarium 1 sessile. Styli 3-5, nunc basi coaliti, nunc distincti, bifidi ramosive. Capsula 1-3-locularis, 3-5-valvis, valvulis margine plus minus introflexis, ab apice dehiscentibus, intus nunc in nervo medio, nunc ad basin seminiferis. Semina secus nervum medium biseriata aut ad basin capsulæ congesta, ovata, nitida, nuda aut arillo tenui folliculari involuta. Albumen cartilagineum carnosumve. Embryo rectus, axilis, dicotyledoneus, radicula ad hilum versâ obtusâ, cotyledonibus crassiusculis. — Herbæ teneræ, pilos apice glandulosos sæpius gerentes aut glaberrimæ. Folia alterna, juniora saltem in D. legitimis, circinatim Filicum more ab apice ad basin voluta! Petioli basi ciliis stipularibus sæpius instructi. Pedunculi juniores circinatim voluti.

I. DROSERA. *Lin. gen.* 391. *Lam. ill. t.* 220.

Sepala petalæque 5 inappendiculata. Stam. 5. Styli 3-5, bipartiti. — Herbæ in uliginosis sphagnosis crescentes. Folia ciliis glandulosis rubidis irritabilibus ornata.

Sect. I. ROSELLA. — Ros-Solis. *Tourn. inst. t.* 127.

Styli simplices aut 2-3-partiti, lobis integris apice subcapitatis.

§. 1. *Acaules, foliis radicalibus sæpius rosulatis.*

1. D. ACAULIS (Thunb. prod. 57.) foliis oblongis obtusis basi angustatis, scapo brevissimo 1-floro. — Ad cap. B.-Spei. Flos albus.
2. D. UNIFLORA (Willd. enum. 340.) foliis subrotundis breviter petiolatis, scapo brevi 1-floro. — Ad fretum Magellanicum. (v. s.)
3. D. PYGMÆA, foliis subrotundis peltatis longè petiolatis, stipulis scariosis, scapo 1-floro. (†) in insulâ ad introitum Jervis'bay. Caley. D. pusilla Brown! ined. non Humb. (v. s.)
4. D. PAUCIFLORA (Banks! herb.) foliis obovato-oblongis basi attenuatis, scapo ciliato glanduloso 1-2-floro, petalis calyce triplò majoribus. 2 ad cap. B.-Spei. (v. s. in h. Banks.)
5. D. PUSILLA (Humb. B. et Kunth. nov. gen. am. 5. p. 390. t. 490. f. 1.) foliis spathulatis glandulosis limbo obovato suprâ margineque ciliato, scapis 2-3-floris calycibusque glabris, seminibus subglobosis. — in humidis arenosis ad ripam flum. Orinoci. D. biflora Willd. in Schult. syst. 6. p. 763.
6. D. TENELLA (H. B. et Kunth. l. c. p. 391. t. 490. f. 2. Willd. in Schult. syst. 6. p. 763.) foliis spathulatis limbo obovato-subrotundo suprâ margineque ciliato-glandulifero, scapis capillaceis elongatis 2-3-floris calycibusque glabris, seminibus oblongis. — In Alpibus Novæ-Andalusie. Affinis D. capillari.
7. D. UMBELLATA (Lour. ed. Willd. 1. p. 232.) foliis ovalibus longè petiolatis, scapo nudo apice umbellato 5-floro. — In Chinâ.
8. D. BREVIFOLIA (Pursh! fl. bor. am. 1. p. 211.) foliis cuneatis obtusissimis

- vix petiolatis brevibus, stipulis scariosis 3-4-fidis, scapo 3-4-floro. ① à Carolinà ad Georgiam. (v. s. in h. Lamb.)
9. *D. PALEACEA*, foliis oblongis petiolatis, stipulis scariosis acutè ad apicem incis, scapo erecto glabro foliis duplò longiore apice racemoso. ① in Novâ-Hollandiâ ad portum regis Georgii. (v. s. in h. Mus. Par.)
10. *D. TRINERVIA* (Spreng! anleit. i. p. 298.) foliis spathulato-cuneiformibus sessilibus trinerviis, scapo paucifloro calycibusque pubescentibus. — ad Cap. B.-Spei. (v. s.)
11. *D. CUNEIFOLIA* (Thunb. prod. 57.) foliis obovato-cuneiformibus sessilibus reticulato-subvenosis, scapo paucifloro calycibusque pubescentibus. — ad Cap. B.-Spei. Burch. trav. i. p. 57. cat. n. 599. (v. s.)
12. *D. BURMANNI* (Vahl symb. 3. p. 50.) foliis cuneiformi-spathulatis sessilibus reticulato-subvenosis, scapo erecto paucifloro calycibusque glabris. — In Zeylonâ, Coromandeliâ, forsan in Chinâ. Burm. zeyl. t. 94. f. 2.
13. *D. SPATHULATA* (Labill! nov. holl. t. 106. f. 1.) foliis oblongo-spathulatis in petiolum subangustatis, scapo apice calycibusque glandulosi, floribus breviter racemosis subsessilibus. — in terrâ Van-Diemen et Port-Jackson. (v. s.)
14. *D. PETIOLARIS* (Brown! ined.) foliis orbiculatis peltatis longè petiolatis, petiolorum calycis scapique pilis confertis non glandulosi. ½ in Novâ-Hollandiâ prope Endeavour-river. (v. s. in h. Banks.)
15. *CAPILLARIS* (Poir! dict. 6. p. 299.) foliis obovato-spathulatis breviter petiolatis, scapo erecto calycibusque glabris. ① in Carolinâ et ad Philadelphiam. *D. rotundifolia* Michx! fl. bor. am. i. p. 186. An satis ab Europæâ differt? (v. s.)
16. *D. ROTUNDFOLIA* (Lin. sp. 402.) foliis orbiculatis, petioli pilosis limbo longioribus, scapis rectis folia triplò superantibus, seminibus arillatis. ① in Europâ. Drev. et Hayn. pl. eur. 3. p. 40. t. 74. opt. (v. v.)
β. distachya, scapo bifido apice distachyo. — in Europâ et in Americâ boreali. (v. s.)
17. *D. INTERMEDIA* (Drev. et Hayn. pl. eur. 3. p. 43. t. 75. B.) foliis obovatis, petioli glabris limbo longioribus, scapis adscendentibus folia paulò superantibus, seminibus exarillatis. ①? in Europâ. *D. longifolia* Lin. sp. 403. engl. bot. t. 868. (v. v.)
β. corymbosa, scapo bifido, ramis divergentibus cymoso-corymbosis. — circa Haguenau. (v. s.)
γ. Americana, scapo simplici, foliis oblongo-obovatis. — in Amer. bor. *D. americana* Willd. en. 340. *D. longifolia* Michx! fl. i. p. 186. (v. s.)
18. *D. ANGLICA* (Huds. angl. 135.) foliis oblongis obtusis basi attenuatis, petioli glabris limbo vix longioribus, scapis erectis folia duplò superantibus, seminibus arillatis. ½ in Europâ mediâ et boreali. *D. longifolia*. Drev. et Hayn. 3. p. 42. t. 75. A. opt. (v. v.)
β. subuniflora, scapo 1-2-floro. — ad montem S.-Gothardi. (v. s.)
19. *D. LINEARIS* (Gold. in Edimb. phil. journ. 1822. p. 325.) foliis linearibus obtusis, petioli longissimis nudis, scapis radicatis, floribus paucis, calycibus glabris. ½ in Canadâ super. ad lacum Simcoe.
20. *D. FILIFORMIS* (Raf. in Desv. journ. 1808. i. p. 227.) foliis filiformibus longissimis, petioli limbo multò brevioribus basi lanatis, scapis erectis folia vix æquantibus. ½ in Novâ-Cæsareâ, circa Tuckerton. *D. tenuifolia* Willd. enum. 340. (v. s.)
21. *D. CAPENSIS* (Lin. sp. 403.) foliis subradicalibus oblongo-linearibus obtusis basi attenuatis, petiolo glabro limbo ferè brevior, scapo subascendente pilosiusculo foliis longiore. ½ ad Cap. B.-Spei. Burm! afr. t. 75. f. 1. (v. s.)
- §. 2. *Caulescentes, foliis caulinis.*
22. *D. RAMENTACEA* (Burch! cat. n. 7692.) caule erecto foliis veteribus deflexis oblecto, foliis ad apicem subrosulatis obovatis, petiolo ciliato limbo longiore, stipulis incis. ½ ½ ad Cap. B.-Spei. (v. s. in h. Banks et Burch.)
23. *D. MADAGASCARIENSIS*, caule adscendente, foliis sparsis obovatis, petiolo glabro limbo longiore, stipulis ciliato-laceris, pedunculo florifero foliis sextuplò longiore. ½ in Madagascar. (v. s.)

24. *D. FOLIOSA* (Ell. sketch. 1. p. 376.) caulescens, foliis confertis ovalibus basi cuneatis, petiolis glabris elongatis, stipulis subulatis. — in Carolinā meridionali. Species mihi dubia. Sub hoc nomine in herb. Balbisii vidi specimen ē Novā-Cæsareā ortum, staturā pollicari à *D. intermediā* americanā tantum diversum, et stipulis non subulatis sed ciliato laceris donatum. An vera *D. foliosa*?
25. *D. CISTIFLORA* (Lin. amœn. 6. p. 85.) caule erecto simplici, foliis oblongo-linearibus sessilibus, floribus paucis pedicellatis. ? ad Cap. B.-Spei. Burm! afr. t. 75. f. 2. Flos amplus purpureus; variat, ex Thunb., flore albo rubrove basi maculato.

℘. *violacea*, foliis angustioribus, caule bifloro. — *D. violacea* Willd. enum. 1. p. 340.

26. *D. INDICA* (Lin. sp. 403.) caule ramoso, foliis linearibus, petiolo glabro vix limbo angustiore, pedicellis calycibusque puberulis. — in Ceylonā et Malabariā. Burm. zeyl. t. 94. f. 1.

Sect. II. ERGALEIUM.

Styli apice capillaceo-multifidi penicilliformes.

§. 1. *Caulescentes, foliis caulinis peltatis.*

27. *D. LUNATA* (Buchan! ined.) caule erecto glabro, foliis sparsis petiolatis peltatis lunato-triangularibus, racemo laxo terminali, calycibus glabris. ① in Indiā Orientali circa Sembu. (v. s.)
28. *D. PELTATA* (Sm! in Rees. cycl. n. 5.) caule erecto glabro, foliis sparsis petiolatis peltatis subtriangularibus, calycibus ciliato-glandulosis. ① in Novā-Hollandiā. — Labill. n. holl. t. 106. f. 2. (v. s.)
29. *D. BANKSII* (Brown! ined.) caule erectiusculo glabro apice inter flores piloso, foliis sparsis petiolatis peltatis orbiculatis, calycibus hirtis. ① in Novā-Hollandiā prope Endeavour-river. (v. s. in h. Banks.)
30. *D. MENZESII* (Brown! ined.) caule erectiusculo flexuoso glabro in axillis ramuloso, foliis junioribus subfasciculatis petiolatis peltatis orbiculatis, racemis bifloris glabris, calycibus ciliato-villosis. ① in Novā-Hollandiā. (v. s.)

§. 2. *Acaules, foliis radicalibus partitis.*

31. *D. BINATA* (Labill! n. holl. 1. t. 105.) foliis longè petiolatis profundè bipartitis, lobis linearibus. ? in terrā Van-Diemen. (v. s.)
32. *D. PEDATA* (Pers. ench. 1. p. 357.) foliis longè petiolatis pedatim dichotomis, lobis linearibus. ? in Novā-Hollandiā. *D. dichotoma* Sm! in Rees cycl. n. 6. (v. s.)

II. *ALDROVANDA*. Monti act. bon. 2. p. 3. p. 404. t. 12. Lin. gen. 390. Lam. ill. t. 220.

Sepala petalaeque 5 inappendiculata. Stam. 5. Styli 5 filiformes breves. Stigmata obtusa. Caps. globosa 5-valvis, 1 locularis, 10-sperma.

1. *A. VESICULOSA* (Lin. sp. 402.) ? All. ① Sav. — in Europæ australis aquis stagnantibus natans. Folia verticillata, petiolo in cilia 6 setacea desinente, limbo complicato ferè vesiculari. DC. fl. fr. 5. n. 4294. (v. v.)

III? *ROMANZOWIA*. Cham. in hor. ber. 71. t. 14.

Sepala 5, basi coalita. Pet. 5 in corollam 5-fidam deciduam coalita. Stam. 5 imo tubo inserta. Capsula 2-valvis, 2-locularis, ∞-sperma. — An potiùs *Saxifragea*?

1. *R. UNALASCHENSIS* (Cham! l. c.) ? in vallibus insulæ Unalaschkæ. Herba habitus *Saxifragæ* aut *Adoxæ*. Folia petiolata subrotundo-reniformia grossè dentata. Flores albidì ebracteati. (v. s.)

IV. BYBLIS. Salisb. parad. t. 95.

Sepala et petala 5 inappendiculata. Stam. 5. Antheræ apice biporosæ. Stylus 1 filiformis. Stigma bilobum. Capsula 2-valvis, 2-locularis, ∞ sperma.

1. *B. LINIFLORA* (Sal. l. c.) ①? in Novā-Hollandiā. Herbula simplex; folia linearia more *Droseræ* piloso-glandulosa; flores cærulei.

V. RORIDULA. *Lin. syst. veg.* 244.

Sepala et petala 5 inappendiculata. Stamina 5, antheræ apice biporosæ basi in appendicem callosam productæ. Styl. 1. Stigma 3-lobum. Caps. 3-locularis, 3-valvis. Semina in quoque loculo subsolitaria.

1. *R. DENTATA* (Lin. l. c.) 5 ad Cap. B. Spei. Suffrutex 1-3-pedalis, foliis confertis linearibus ciliato-glandulosi. Bractæe foliaceæ. Lam. ill. t. 141. (v. s.)
β. muscicapa (Gaertn. fr. 1. t. 62. f. 8.) Hab. cum var. *α*. — Petala angustiora.

VI. DROSOPHYLLUM. *Link in Schrad. journ.* 1806. 1. p. 53.

Sepala et petala 5 unguibus approximata. Stam. 10. Styli 5 filiformes. Capsula 5-valvis, 1-locularis, ad medium valvulis introflexis ferè 5-locularis.

1. *D. LUSITANICUM* (Link l. c.) 2/5 in collibus arenisque Lusitaniæ. Caulis fruticosus, folia linearia integra glandulis stipitatis obsita. Pauicula corymbosa. Flores ampli sulphurei. — Pluk. alm. t. 117. f. 2. Saint-Hil. mem. mus. 2. t. 4. f. 13. *Drosera Lusitanica* Lin. sp. 403. *Spergula droseroides*. Brot. fl. lus. 2. p. 215. (v. s.)

VII. DIONÆA. *Ellis. nov. act. ups.* 1. p. 98. t. 8. *Lin. mant.* 151.

Sepala et petala 5. Stam. 10-20, antheris lateraliter dehiscentibus. Stylus 1. Stigma orbiculatum. Capsula 5-valvis, 1-locularis. Semina eo in substantiâ cellulosa ad basim capsulæ semi-immersa.

1. *D. MUSCIPULA* (Lin. mant. 238.) 2/ in Amer. boreali. — Herba glabra; folia radicalia petiolo apice dilatato, limbo bilobo ciliato irritabili complicato. Vest. malm. t. 29. Ker. bot. reg. t. 785. Delaun. herb. amat. t. 349. (v. v.)

VIII? PARNASSIA. *Tourn. inst. t.* 127. *Lin. gen.* 384.

Sepala 5. Petala 5. Squamæ 5 (an stamina abortiva?) unguibus petalorum oppositæ, in setas apice glandulosas desinentes. Stam. 5. Antheræ posticæ. Stigmata 4 sessilia. Capsula 4-valvis, 1-locularis, valvis medio septiferis. Semina arillata. — Herbæ glaberrimæ, habitu et forsan characterè hinc ad *Swertiam*, illinc ad *Violam* accedentes.

1. *P. PALUSTRIS* (Lin. sp. 391.) appendicibus 9-13-setis, petalis subsessilibus, foliis radicalibus cordatis, caulino amplexicauli. 2/ in paludibus submontanis Europæ. — Fl. Dan. t. 584. (v. v.)
2. *P. PARVIFLORA*, appendicibus 5-7-setis, petalis sessilibus, foliis radicalibus ovatis, caulino sessili. 2/ in Amer. bor. An *P. palustris* var? an *P. tenuis* Wahl fl. lapp. n. 137 huc potius quàm ad palustrem referenda? (v. s. in h. Banks.)
3. *P. OVATA* (Ledeb. act. petr. 1815. p. 514.) appendicibus 3-setis, foliis radicalibus ovatis, caulino subcordato-amplexicauli. 2/ in Sibiria orientali. *β? Belvisii*, foliis radicalibus 5-7-nerviis, caulino ovato. 2/ in Amer. bor. *P. ovata*. Beauv! ined. (v. s. sine fl. in h. Beauv.)
4. *P. CAROLINIANA* (Michx. fl. bor. am. 1. p. 184.) appendicibus 3-setis, petalis subsessilibus, foliis radicalibus suborbiculatis, caulino ovato sessili. 2/ à Carolinâ ad Canadâ. Sims. bot. mag. t. 1459. (v. s.)
5. *P. ASARIFOLIA* (Vent! malm. t. 39.) appendicibus 3-setis, petalis unguiculatis, foliis radicalibus reniformibus, caulino subcordato orbiculato. 2/ in Virginiâ et Carolinâ. (v. s.)
6. *P. GRANDIFOLIA*, appendicibus 3-setis, petalis oblongis sessilibus, foliis radicalibus ovatis subcordatis 7-nerviis, caulino subcordato-orbiculato. 2/ in Amer. septentr. apud Cherokees. Beauvois. (v. s.)
7. *P. FIMBRIATA* (Banks! in kœn. ann. 1. p. 391.) appendicibus 3-fidis, petalis obovatis unguiculatis basi ciliatis, foliis radicalibus reniformibus multi-nerviis, caulino cordato. 2/ in Americæ borealis orâ occidentali. (v. s. in h. Banks.)

Ordo XVIII. POLYGALEÆ *Juss. ann. mus. 14. p. 386. mem. mus. 1. p. 385.*

Calyx 5-sepalus, sepalis per æstivationem imbricatis, 2 interioribus sæpius petaliformibus, 3 externis minoribus, quorum 2 anteriora interdum coalita et tertium posticum. Petala 3-4 hypogyna, mediante tubo stamineo plùs minùsve connexa, rariùs distincta. Staminum filamenta cum petalis coalita, monadelpha, apice in phalanges duas oppositas æquales divisa. Antheræ 8 uniloculares, basi insertæ, apice poris dehiscentes. Ovarium 1 liberum 2-loculare, rariùs 1-3 loculare. Stylus 1 incurvus. Stigma infundibuliforme aut bilobum. Pericarpium capsulare aut drupaceum, biloculare aut (abortù) uniloculare, valvis medio septigeris. Semina in loculis solitaria, pendula, sæpè basi carunculato-arillata, interdum pilosa aut comosa. Embryo rectus, planus, nunc in albumine carnosio axilis, nunc rariùs exalbuminosus endoplevrà tunc tumidiore. — Herbæ aut suffrutices. Folia in plerisque alterna, super caulem articulata, integra. Flores racemosi. Succus lacteus præsertim ad radicem. Cortex radixque amara.

Ordo naturalissimus, sed affinitates dubiæ. Habitù florum refert et Leguminosas et Fumariaceas. Staminum sitù, numero et dispositione ad Fumariaceas forsàn magis accedit, pluribus etiam notis ad Tremandreas vergit. An potiùs admittendæ sunt bracteæ tres sub flore, sepala 2, petala 4 quorum duo, ut in Leguminosis, in carinam coalita, stamina 4 seu 8? tunc affinitas cum Fumariaceis multò clariùs appareret. An potiùs sepala 5, petala 5 quorum 3 in carinam coalita; tunc affinitas cum Leguminosis propinquior!

I. POLYGALA. *Tourn. inst. t. 79.*

Calycis sepala persistentia, 2 interiora alæformia. Petala 3-5 tubo stamineo connexa, inferiore carinæformi (forsàn e duobus coalitis constante). Capsula compressa elliptica obovata aut obcordata. Semina pubescentia, hylo carunculata, coinà destituta.

Sect. I. PSYCHANTHUS. *Raf. specch. 1. p. 116.*

Carina amplè cristata. Capsula glabra, marginata. Bracteæ tres sæpius persistentes ad basin pedicellorum. — Frutices Capenses (unicus Arabicus.)

* *Foliis oppositis.*

1. P. OPPOSITIFOLIA (Lin. *mant. 259.*) foliis oppositis ovatis acutis. ♂ in montosis cap. B. Spei. Valdè sequenti affinis. *Ker. bot. reg t. 636. (v. s.)*
2. P. CORDIFOLIA (Thunb. *prod. 120.*) foliis oppositis cordatis acuminatis, ramulis teretibus glabris. ♂ ad cap. B.-Spei. P. fruticosa Berg. *cap. 188.* (v. v.)*

3. *P. TETRAGONA* (Burch! cat. n. 4639.) foliis oppositis cordatis acuminatis, ramulis tetragonis glabris. ♂ ad cap. B.-Spei. (v. s.)
4. *P. NUMMULARIA* (Burch! cat. n. 3709.) foliis oppositis basi cordatis orbiculato-reniformibus mucronatis, ramulis teretiusculis glabris. ♂ ad cap. B.-Spei. (v. s.)
5. *P. BORONIEFOLIA* (Burch! cat. n. 6861.) foliis oppositis cordatis acuminatis, ramulis teretibus hispidis. ♂ ad cap. B.-Spei. (v. v.) — An hæc quinque Stirpes sint meræ varietates dubito?

** *Foliis alternis.*

6. *P. CLUTTHOIDES* (Burch! cat. n. 3326.) foliis ellipticis submucronatis, ramulis glaberrimis, petalis anticis bilobis, lobo acuto longissimo. ♀ ad cap. B.-Spei. (v. s.)
7. *P. MYRTIFOLIA* (Lin. am. 2. p. 138.) foliis obovatis oblongisve submucronatis, ramulis adpressè pubescentibus, bracteis persistentibus æqualibus, pedicellis flore brevioribus. ♀ ad cap. B.-Spei.
 - a. *vera*, foliis obovato-oblongis planis. — Burm. afr. t. 73. f. 1. — Bot. reg. t. 669. (v. s.)
 - β. *angustifolia*, foliis oblongis. — An *P. tenuifolia* Link. enum. 2. p. 220? (v. s.)
 - γ. *rosmarinifolia*, foliis oblongo-linearibus margine revolutis. — An potius sequentis var.? sed rami glabriusculi. (v. s.)
8. *P. LIGULARIS* (Ker. bot. reg. t. 637.) foliis lingulato-linearibus subobtusis confertis patulis glabris, ramis villosis, bracteis persistentibus æqualibus rachique villosis, racemo brevi conferto. ♂ ad cap. B.-Spei.
9. *P. INTERMEDIA*, foliis oblongo-linearibus mucronatis margine revolutis, ramulis glabris, bracteis persistentibus æqualibus, pedicellis flore sublongioribus, alis subcuspidatis. ♀ ad cap. B.-Spei. (v. s.)
10. *P. BRACTEOLATA* (Lin. am. 2. p. 137. var. β.) foliis lineari-lanceolatis planis, ramis glabris hispidisve, bracteis persistentibus inferiore longiore, pedicellis flore sublongioribus, alis cuspidatis. ♀ ad cap. B.-Spei.
 - a. *stricta*, foliis erectis, ramisque glabris. — Pluk. t. 53. f. 2. (v. v.)
 - β. *patula*, foliis patulis, ramisque glabris (v. v.)
 - γ. *hispida*, foliis erectis ramisque hispidis (v. s.)
 - δ. *umbellata* (Thunb. prod. 120.) foliis erectis ramisque hispidis, floribus paucis. ① Burm! afr. t. 73. f. 5. An forsitan cum var. γ speciem propriam constituens? (v. s.)
11. *P. TINCTORIA* (Vahl. Symb. 1. p. 50.) foliis obovatis ramisque villosovelutinis, calycibus pubescentibus. ♀ in montibus Arabiæ felicis. *P. bracteolata*. Forsk. descr. 213. (v. s. in h. Banks.)
12. *P. PUBIFLORA* (Burch! cat. n. 6205.) foliis lanceolatis acuminatis ciliatis, ramulis hirsutis, calycibus pubescentibus. ♀ ad cap. B.-Spei. (v. s. in h. Burch.)
13. *P. BURMANNI*, foliis linearibus obtusiusculis, ramulis pubescenti-velutinis, racemis supra-axillaribus, pedicellis flore brevioribus, bracteis deciduis. ♀ ad cap. B.-Spei. *P. bracteolata* δ Lin. — Burm! afr. t. 73. f. 4. Burch! cat. n. 6437. — An hujus sectionis? (v. s.)
14. *P. AFFINIS*, foliis linearibus obtusis, junioribus subvillosis, ramulis pubescentibus, racemis oppositifoliis, bracteis minimis persistentibus. ♀ ad cap. B.-Spei. (v. s. in h. Deless.)
15. *P. MICROLOPHA* (Burch! cat. n. 3416.) foliis linearibus mucronatis ramulisque glabris, racemis paucifloris, pedicellis brevissimis, alis obovatis. ♀ ad cap. B.-Spei. Crista inter *Psycanthos* parva. (v. s. in h. Burch.)
16. *P. SIMPLEX*. (Burch. cat. n. 1933.) foliis oblongo-cuneatis obtusis submucronatis cauleque simplici glabris, bracteis deciduis, pedicellis patulis,

alis ovalibus. $\frac{1}{2}$ ad cap. B.-Spei. Flores dimidiò minores quàm in *P. speciosa*, cui affinis. (v. v. in hort. Burch.)

17. *P. SPECIOSA* (Simms. bot. mag. t. 1780.) foliis inferioribus oblongo-cuneatis obtusis, superioribus linearibus, ramisque virgatis glabris, bracteis deciduis, pedicellis patulis, alis subrotundis. $\frac{1}{2}$ ad cap. B.-Spei. Bot. reg. t. 150. Herb. amat. t. 193. Bot. mag. t. 621. (v. v.)

18. *P. PEDUNCULARIS* (Burch! cat. n. 5163.) foliis linearibus margine revolutis, supernè scabris, pedunculis rigidulis patulis apice paucifloris, ramulisque angulatis. $\frac{1}{2}$ ad cap. B.-Spei. Flores ferè *P. myrtifoliæ*. (v. s. in h. Burch.)

19. *P. TERETIFOLIA* (Thunb. prod. 120.) foliis linearibus margine revolutis subteretibus subtus sulcatis patulis glabris, ramulis albo-velutinis, racemis brevibus paucifloris. $\frac{1}{2}$ ad cap. B.-Spei. *P. empetrifolia*. Houtt. hist. nat. 11. D. v. t. 28. f. 1. (v. s.)

20. *P. PINIFOLIA*. (Lam. ill. t. 598. f. 2.) foliis linearibus margine revolutis subteretibus subtus sulcatis patulis petiolulatis, ramisque glabris. $\frac{1}{2}$ ad cap. B.-Spei. An à priore satis differt? folia duplò longiora.

21. *P. REFRACTA* (Burch! cat. n. 4891.) foliis linearibus acuminatis patentibus, ramisque glabris, pedunculis lateralibus paucifloris, pedicellisque refractis, bracteis deciduis, alis ovalibus. $\frac{1}{2}$ ad cap. B.-Spei.

22. *P. MACRA*, foliis minimis sublinearibus acutis erectis ramisque glabris, racemis paucifloris, pedicellis flore brevioribus, alis ovatis. $\frac{3}{4}$ in aridis cap. B.-Spei ad Breda-River. (v. s. comm. à cl. Lamb.)

23. *P. GENISTOIDES* (Poir! dict. 5. p. 492.) foliis linearibus acutis distantibus paucis ramisque virgatis glabris, racemis elongatis, bracteis deciduis. pedicellis demùm patulis flore brevioribus, alis ovalibus obtusis. $\frac{1}{2}$ ad cap. B.-Spei. (v. s.)

℞. *ephedroides*. (Burch! cat. n. 1793. itin. 1. p. 485.) ramis vix foliosis, racemo laxo. (v. s.)

γ. *spartioides*. (Burch. cat. n. 4449.) ramulis foliisque junioribus vix pubescentibus, alis subrotundis. (v. s.)

24. *P. GARCINI*, foliis linearibus ferè subulatis erectis subdistantibus ramisque virgatis glabris, racemis elongatis, pedicellis bracteis vix duplò superantibus, alis ovalibus obtusis. $\frac{1}{2}$ ad cap. B.-Spei. Flores purpurascens. — *P. bracteolata*. γ Lin. am. 2. p. 137. Burm. afr. t. 73. f. 3. Burch! trav. 1. p. 19. 37. (v. s.)

25. *P. ERICEFOLIA*, foliis linearibus erectis confertis acutis glabris, superioribus ciliatis, pedicellis subumbellatis flore longioribus, alis ovalibus mucronatis. $\frac{1}{2}$ ad cap. B.-Spei. Burch! cat. n. 5514. Forsan à plurimis cum umbellatâ confusa, sed toto cœlo diversa. (v. s.)

26. *P. PUNGENS* (Burch. cat. n. 1598. trav. 1. p. 304.) foliis linearibus acutiusculis angustis parvis, ramulis divaricatis glaucis rigidis apice spinescentibus, racemis 2-4-floris. $\frac{3}{4}$ ad cap. B.-Spei. Crista purpurascens. Alæ albidæ, venis viridibus. Capsula obcordata.

27. *P. RIGENS* (Burch! cat. n. 1821. trav. 1. p. 465.) foliis lineari-oblongis canaliculatis obtusis apice mucrone subrecurvo ramisque junioribus pubescentibus, racemis paucifloris, alis oblongis subciliatis. $\frac{3}{4}$ ad cap. B.-Spei. (v. s.)

28. *P. LEPTOPHYLLA* (Burch! cat. n. 2380. trav. 1. p. 400.) foliis linearibus erectis acutiusculis ramulisque junioribus subpubescentibus, racemis laxifloris, bracteis minimis deciduis, alis ellipticis. $\frac{1}{2}$ ad cap. B.-Spei. (v. s.)

29. *P. HISPIDA* (Burch! cat. n. 4473.) foliis inferioribus ellipticis, summis sublinearibus ramulisque molliter hispidis, racemis elongatis pubescentibus, bracteis deciduis infimâ pedicello subæquali, alis ellipticis. $\frac{3}{4}$? ad cap. B.-Spei. (v. s.)

30. *P. ASBESTINA* (Burch! cat. n. 2030. trav. 1. p. 543.) foliis oblongis basi at-

tenuatis, supremis mucronatis, glaucescentibus ramulisque subpuberulis, racemis subunifloris lateralibus, alis oblongo-linearibus. ζ ad cap. B.-Spei ultrâ flumen Gariiep. Flores albi. Capsula ovalis apice bifido. An hujus sectionis? (v. s.)

31. *P. POLYPHYLLA* foliis oblongis utrinquè acutis confertis dorso et margine scabris, ramulis subpuberulis, pedicellis axillaribus. 1-floris, alis suborbiculatis. ζ ad cap. B.-Spei. An hujus sectionis? habitus ferè Mundiaë. (v. s.)

Sect. II. POLYGALON.

Carina cristata. Capsula glabra. Bractea 3 mox decidua ad basin pedicellorum. — Herbae aut suffrutices Europæ, aut Asiæ temperatæ incolæ.

32. *P. TENUIFOLIA* (Willd. sp. 3. p. 879.) foliis linearibus mucronatis cauleque erecto glabris, racemis elongatis laxis subebracteatis, alis ovali-oblongis subacutis. \mathcal{Z} in Sibiria. Var. flore carneo et cæruleo. (v. s.)
33. *P. SIBIRICA* (Lin. sp. 987.) foliis lanceolatis inferioribus ovatis glabris, junioribus subpuberulis, racemis laxis paucifloris demum ebracteatis, alis ellipticis mucronatis. \mathcal{Z} in Sibiria. (v. s.)
34. *P. JAPONICA* (Houtt. syst. 8. t. 62. f. 1.) foliis glabris inferioribus subrotundis, superioribus ovatis acutis, caule erecto, racemis lateralibus subcorymbosis, calycibus patulis, alis oblongis. \mathcal{Z} in Japonia. *P. vulgaris*. Thunb! fl. jap. 277. (v. s.)
35. *P. SAXATILIS* (Desf. fl. atl. t. 175.) foliis oblongo-lanceolatis mucronatis, caule basi fruticoso decumbente, racemis extrâ-axillaribus paucifloris, alis ellipticis. ζ in rupestribus Mauritaniae, Hispaniae, Galliae austr. *P. rupestris*. Pourr. (v. s.)
36. *P. OXYCOCOÏDES* (Desf. atl. t. 174.) foliis ellipticis crassiusculis, caule fruticoso decumbente, racemis lateralibus brevibus paucifloris, alis ovatis ζ in Atlantis fissuris. (v. s.)
37. *P. SUPINA* (Schreb. dec. t. 10.) foliis obovatis subretusis, caule fruticuloso supino, racemis subterminalibus 7-10-floris, alis ovalibus subtrinerviis corollâ longioribus. ζ in glareosis Armeniae, Tauriae. *P. andrachnoides* Willd. — Buxb. cent. t. 70. f. 2. — Var. fl. cæruleo et purpureo. (v. s.)
38. *P. ADSCENDENS* (Clark! in Spreng. nen. entd. 3. p. 165.) foliis oblongis basi attenuatis, infimis obovatis subvillosis, caule basi fruticoso adscendente racemis paucifloris, alis oblongis obtusis corollâ dimidiò brevioribus, ovario sessili. ζ in insulâ Cos. (v. s. in h. Lamb.)
39. *P. VENULOSA* (Sibth. et Sm. prod. 2. p. 52.) foliis elliptico-lanceolatis, caule adscendente, racemis subterminalibus, alis oblongis trinerviis transversè venosis corollâ dimidiò brevioribus. \mathcal{Z} in insulis Archipelagi et in Græciâ.
40. *P. ROSEA* (Desf. atl. t. 176.) foliis inferioribus oblongis obtusis, superioribus lanceolatis, caule erecto basi suffruticoso, alis ovalibus multinerviis corollâ paulò brevioribus, ovario brevissimè stipitato. ζ in Atlante prope Tlemsen. (v. s.)
41. *P. MAJOR* (Jacq. austr. t. 413.) foliis lineari-lanceolatis acutis, caulibus erectis, alis ovatis multinerviis corollâ paulò brevioribus, ovario longè stipitato. \mathcal{Z} in montosis pratis Italiae, Austriae, Græciæ, Orientis. Flores roseo-purpurei. (v. s.)
42. *P. FLAVESCENS* (DC. cat. h. monsp. 134.) foliis linearibus sublanceolatis acutis, caulibus adscendentibus, alis ellipticis utrinquè acutis corollâ capsulâque longioribus, ovario subsessili. \mathcal{Z} in Appenninis Romanis. — Seb. pl. rom. fasc. 1. t. 1. (v. s.)
43. *P. VULGARIS* (Lin. sp. 986.) foliis lineari-lanceolatis obtusiusculis, caulibus adscendentibus, alis ovatis capsulâ parùm longioribus corollâ subbrevioribus, ovario subsessili. \mathcal{Z} in pratis Europæ.

- a. vera*, caulibus erectiusculis, foliis inferioribus obovatis obtusis, superioribus linearibus acutis — OEd. fl. dan. t. 516. (v. v.)
- β. pubescens*, caule decumbente foliisque pubescentibus. Rohd! Journ. bot. 2. p. 359. (v. s.)
- γ. elata*, caulibus erectis foliis ex ovali oblongis. P. comosa Schkuhr. handb. t. 194. (v. s.)
- δ. Verviana* (Lej. fl. spa. 2. p. 92.) caulibus adscendentibus, foliis lanceolato-linearibus, floribus albis.
- ε. acutifolia*, caulibus erectis, foliis linearibus, floribus mediocribus cæruleis. (v. s.)
- ζ. angustifolia*, caulibus erectis, foliis linearibus, floribus mediocribus roseis. P. Monspeliaca Vill. dauph. 3. p. 388? (v. s.)
- η. grandiflora*, caulibus erectis, foliis linearibus, floribus magnis roseis. P. Monspeliaca All. ped. n. 1089. (v. s.)
44. P. AMARA (Lin. sp. 987.) foliis radicalibus obovatis obtusissimis, caulibus linearibus, caulibus erectiusculis, alis ellipticis corollam æquantibus, capsulâ suborbiculatâ. \mathcal{L} in pratis montanis Europæ. — Wahl. carp. n. 701.*
- a. Austriaca* (Crantz. austr. t. 2. f. 4.) amarissima, floribus minoribus albicantibus, alis post anthesin viridescentibus. (v. v.)
- β. alpestris*, insipida, alis fructibus coloratis, radice magis cespitosa. Jacq. austr. t. 412. P. Vaillantii. Bess. cont. 2. p. 73? Vaill. par. t. 32. f. 3? (v. v.)
- γ? cespitosa*, floribus cæruleis, caulibus cespitoso-procumbentibus, foliis linearibus summis ad flores approximatis. P. repens. Merat? (v. s.)
- δ. Alpina* (Poir. dict. 5. p. 488.) floribus cæruleis paucis, caulibus cespitoso-procumbentibus brevissimis, foliis inferioribus obovatis. (v. s.)
- ε. obtusifolia*, floribus cæruleis, caulibus cespitoso-procumbentibus, foliis inferioribus obovatis obtusissimis. (v. s.)
- ζ? decipiens* (Bess. cont. 2. p. 73.) foliis obovato lanceolatis, radicalibus maximis obtusissimis, alis trinerviis, caulibus adscendentibus. \mathcal{L} in Volhyniâ. P. amara fl. lith.
45. P. HYBRIDA, glabra erectiuscula simplex, foliis inferioribus oblongis, ceteris linearibus, alis oblongis corollâ longioribus. \mathcal{L} in desertis Uralensibus. Media inter duas priores. (v. s.)
46. P. PODOLICA, glabra erectiuscula simplex, foliis linearibus acutis, bracteis subulatis duplò flore longioribus, alis oblongis uninerviis corollæ æqualibus (albis cum fasciâ mediâ viridi.) \mathcal{L} in Podoliâ australi (v. s.)
47. P. MONSPELIACA (Lin. sp. 987.) glabra erecta subsimplex, foliis linearibus acuminatis, alis oblongis utrinquè acutis trinerviis capsulâ multò longioribus. $\textcircled{1}$ in sterilibus circuitibus maris Mediterranei. DC! ic. rar. 1. p. 3. t. 9. (v. v.)
48. P. EXILIS (DC! cat. h. monsp. 133.) erecta glabra ramosissima, foliis linearibus crassiusculis, alis ovalibus obtusis 1-nerviis capsulâ longitudine, racemo laxiusculo. $\textcircled{1}$ in Galliâ australi, Hispaniâ. — P. linearis Lag. P. nova. Boiss. fl. eur. 1. t. 474. f. 1. (v. v.)
49. P. GLUMACEA (Sibth. et Sm. prod. 2. p. 52.) suberecta, foliis lanceolato-linearibus acuminatis, alis elliptico-lanceolatis corollâ duplò longioribus trinerviis obsolete venosis. \mathcal{L} in insulâ Cypro et ad Gibraltar.
50. P. LEPTALEA, erecta glabra simplex, foliis inferioribus oblongis obtusis, superioribus linearibus acutis, alis oblongis subtrinerviis, corollâ capsulâque longioribus. $\textcircled{1}$ in Napauliâ. Wallich. Herbula vix digitalis. Flores parvi penduli. (v. s.)
51. P. OLIGOPHYLLA, glabra suberecta basi subramosa, foliis linearibus distantibus, alis oblongis subtrinerviis corollâ capsulâque longioribus. $\textcircled{1}$ in Napauliâ. Wallich. Priori valdè affinis, duplò major. (v. s.)

Sect. III. BLEPHARIDIUM.

Corollæ carina cristata. Capsula apice emarginata margine ciliata. Racemi breves laterales. Bracteæ 3 parvæ ad basin pedicellorum. — Herbæ parvæ sæpius ramosissimæ Indiæ aut Ægyptiæ et Guineenses. — Flores pallidi aut virescentes.

52. *P. ERIOPIERA*, caulibus adscendentibus ramosis apice pubescenti-velutinis, foliis glabris linearibus acutiusculis, racemis 4-5-floris, alis ellipticis pubescentibus, capsulâ oblongâ velutinâ longioribus. 2℥ in Senegaliâ et Ægypto. *P. paniculata* Forsk. fl. arab. n. 429? (v. s.)

53. *P. OBTUSATA*, caulibus diffusis virgatis glabris, foliis linearibus obtusissimis, racemis 2-6-floris, alis oblongis obtusis glabris subciliatis, capsulâ subpubescente longioribus. 2℥ in Ægypto. (v. s. in h. Balb.)

54. *P. LINEARIFOLIA* (Willd. sp. 3. p. 877.) caulibus erectis virgatis ramosis pubescentibus, foliis linearibus strictis, racemis 6-8-floris, alis ovali-oblongis acutis, capsulâ obcordatâ ciliatâ. 2℥ 5 in insulâ Mindanao et Samboagang. Flores reflexi. (v. s. in herb. Deless.)

55. *P. PROCUMBENS* (Roth. nov. sp. p. 329.) caule procumbente piloso, foliis oblongo-ovalibus obtusis submucronatis ciliatis glabris, racemis axillaribus brevibus paucifloris, alis acuminatis, capsulâ emarginatâ longè ciliatâ. — in Indiâ orient. — Aff. *P. linearifoliæ* ex cl. Roth.

℘. angustifolia (Roth. l. c.) foliis longioribus angustioribus.

56. *P. GLOMERATA* (Lour! coch. ed. W. 2. p. 518.) caule suberecto ramoso pubescente, foliis obovato-oblongis, racemis densis 7-8-floris, alis ovalibus acuminatis, corollâ subripetalâ, capsulâ orbiculatâ. 2℥ 5 in Chinâ prope Canton. (v. s. in h. Mus. Par.)

57. *P. GLAUCOIDES* (Lin. sp. 991.) caulibus diffusis subpubescentibus, foliis ovali-oblongis subpetiolatis glabris glaucis, racemis 7-8-floris, alis ovalibus acutis, capsulâ obovatâ emarginatâ subciliatâ. 2℥ in Ceylonâ. (v. s. in h. Banks.)

58. *P. ARENARIA* (Willd. sp. 3. p. 880.) caulibus procumbentibus apice pubescentibus, foliis oblongis petiolulatis, racemis ovato-globosis breviter pedunculatis, alis obovatis pubescentibus, capsulâ ovali emarginatâ ciliatâ. ① in Guineâ. (v. s. in h. Juss.)

59. *P. SERPYLLIFOLIA* (Poir. dict. 5. p. 499.) caulibus à basi ramosis pubescentibus, lateralibus procumbentibus, foliis oblongis obtusis subpetiolatis, racemis 4-5-floris, alis ovalibus acutis, capsulâ ovatâ immarginatâ ciliatâ. — in Indiâ orient. (v. s. in h. Juss.)

60. *P. ARVENSIS* (Willd. sp. 3. p. 876.) caulibus à basi ramosis procumbentibus pubescentibus, foliis obovatis mucronatis, racemis 7-8-floris, bracteis persistentibus ferè pedicelli longitudine, alis oblongis vix capsulâ ovatâ emarginatâ ciliatâ longioribus. ① in Indiâ orientali. (v. s. ex Roxb.)

61. *P. VAHLIANA*, caule erecto pubescente, foliis obovatis mucronatis subvillosis, superioribus oblongis, racemis 3-5-floris, alis ellipticis capsulâ ovali emarginatâ ciliatâ vix longioribus. ① in Indiâ orient. *P. tomentosa* Vahl! non Thunb. (v. s. in h. Deless.)

62. *P. BRACHYSTACHYA*, caule diffuso subpubescente, foliis linearibus mucronatis glabris, racemis 3-5-floris, alis ovalibus acuminatis subfalcatis capsulâ ovali emarginatâ ciliatâ paulò longioribus. — In Indiâ orientali. (v. s. in h. Deless.)

63. *P. LEPTORHIZA*, caule à basi ramoso diffuso subpubescente, foliis linearibus pubescentibus, racemis 7-8-floris, alis ellipticis obtusis capsulâ ovali emarginatâ ciliatâ paulò longioribus. ① patr. ign. (v. s. in h. Mus. Par.)

64. *P. PERSICARIEFOLIA*, caule erecto apice bifido subpiloso, foliis oblongis utrinquè acuminatis, racemis 10-15-floris è dichotomiâ ortis, alis obovatis capsulâ obcordatâ ciliatâ longioribus. ① in Napauliâ. Wallich. (v. s.)

65. *P. CROTALARIOIDES* (Buchl ined.) caule à basi ramoso suffruticulosos decumbente pubescente, foliis obovatis basi cuneato-petiolatis, racemis 8-10 floris, alis obovato rotundis capsulæ suborbiculatæ ciliatæ æqualibus. 4? in Napauliâ. Bractææ persistentes acutæ. (v. s.)
66. *P. JAVANA*, caule à basi ramoso suffruticoso diffuso, foliis obovatis cuneato-petiolatis. racemis 7 8 floris folii longitudine, alia ovato-rotundis villosis capsulâ suborbiculatâ pubescente paulò longioribus. 5 in Javâ. (v. s. in h. Deless.) An hujus sectionis?

Sect. IV. CLINCLINIA.

Carina cristata. Alæ oblongæ. Capsula glabra marginata apice emarginata utrinquè ad apicem subdentata. — Herbæ suffruticulosæ Americanæ.

67. *P. THASIOIDES* (Willd. sp. 3. p. 877.) caulibus plurimis erectis, foliis oblongo linearibus confertis, racemis 8-12 floris, capsulâ apice subbicorni. 4 5 in regnis Chilensi et Peruviano. Clinclin Feuille. obs. 2. t. 13. (v. s.)
68. *P. GNIDIOIDES* (Willd. sp. 3 p. 78.) caulibus diffusis ramosis, foliis linearibus obtusiusculis, racemis 6-10 floris, capsulâ subemarginatâ. 4 5 in regnis Chilensi et Peruviano. (v. s.)
69. *P. CAPILLIFOLIA* (Desf! herb.) caulibus erectis vix ramosis, foliis capillaceis, racemis gracilibus, alis obovatis, capsulis bifidis. ① in..... (v. s. in h. Desf.) An hujus sectionis?

Sect. V. TIMUTUA.

Carina cristata (crista interdum minutissima). Capsula glabra, ovali-oblonga, vix emarginata. Racemi nunc elongati floribus parvis, nunc conferti spiciformes, floribus inferioribus mox deciduis. Folia alterna aut verticillata. — Herbæ Americanæ.

70. *P. TRICHOSPERMA* (Lin. mant. 259.) racemis spiciformibus ovatis, foliis alternis oblongo-linearibus erectis, caulibus erectis virgatis. 4 in Novâ-Granadâ. Jacq! obs. 3. t. 67. (v. s. in h. Banks.)
71. *P. ASPALATHA* (Lin. mant. 99.) racemis spiciformibus subcapitatis, foliis alternis lineari-setaceis patulis, caulibus erectis virgatis simplicissimis. — in Brasiliâ. (v. s.)
72. *P. STELLERA*, racemis spiciformibus subcapitatis, foliis sparsis paucis, imis obovatis, ceteris linearibus, caulibus erectis ramosis virgatis. ① in Brasiliâ, Portorico, Sancto-Domingo. (v. s.)
73. *P. BYGROPHILA* (H. B. et Kunth. nov. gen. am. 5. p. 395. t. 508.) racemis spiciformibus oblongis, foliis sparsis linearibus glabris glanduloso-punctatis, caule erecto apice subramoso. ① in humidis prope Esmeralda Orinocensium. Flores albidis? Semina retrorsum pubescentia.
74. *P. INCARNATA* (Lin. sp. 986 excl. Pluk. ic.) racemis spiciformibus oblongis glandulosis, foliis sparsis raris subulatis, caulibus erectis vix ramosis. ① in arenosis Virginiæ, Carolinæ. (v. s.)
75. *P. LONGICAULIS* (H. B. et K. l. c. p. 396.) racemis spiciformibus subrotundis, corollis glandulosis, foliis sparsis remotis linearibus glanduloso-punctatis, caule elongato subramoso. ① in Novâ-Audalusiâ et Novâ-Granadâ. Glabra sesquipedalis. Alæ roseæ. Semina conica retrorsum sericeo-pilosa.
76. *P. ADENOPHORA*, racemis spiciformibus oblongis, alis capsulisque glandulosis, foliis sparsis raris subulatis, caulibus erectis subramosis. ① in Guyanæ pratis. *P. incarnata* Aubl. Guian. 2. p. 739. excl. syn. (v. s.)
77. *P. BICOLOR* (H. B. et Kunth. nov. gen. am. 5. p. 394. t. 507.) racemo spiciformi cylindraceo densifloro, sepalis 3 brevioribus nervos 2 parallelos crassos gerentibus, foliis pellucido-punctatis, infer. 3-5^{um} verticillatis, summis sparsis lineari-lanceolatis. ① prope Santa-Rosa Mexicanorum. Glabra simplex monostachya. Flores magni. *P. austriacæ*, albidis, inferiores rubescenti-crocei.

78. *P. TIMOUTOU* (Aubl! guian. 2. t. 295.) racemis spiciformibus cylindraceis imbricatis, floribus demum reflexis, foliis oblongis submucronatis, infimis verticillatis, superioribus alternis, caule erecto ramoso alato. ① in Cayennæ uliginosis. *P. lupulina* Wild. *P. uliginosa* Pers. (v. s.)
79. *P. CUSPIDATA*, racemis spiciformibus cylindraceis imbricatis, floribus demum reflexis, foliis oblongis cuspidato-mucronatis, superioribus alternis, caule adscendente alato. 2? in Brasiliâ. An à priore satis diversa? (v. s.)
80. *P. VARIABILIS* (H. B. et Kunth l. c. p. 397. t. 509.) racemis spiciformibus laxiusculis, foliis sparsis remotis angustè linearibus glabris pellucido-punctatis, caulibus adscendentibus supernè subfastigiato-ramosis. ① in Amer. calid. Semina conica retrorsum pilosa.
- a. floribus roseis. — ad ripam fl. Orinoci.
- β floribus albis. — prope Maypures.
- γ. caule simplici. — juxta Esmeralda.
81. *P. BERTERIANA*, racemis spiciformibus oblongis imbricatis, floribus demum reflexis, alis obtusis, foliis linearibus acutis, infimis verticillatis, ceteris sparsis, caule erecto tenui subangulato. ① in Sauto-Domingo. Bertero. (v. s.)
82. *P. CRUCIATA* (Lin. am. 2. p. 138.) racemis spiciformibus ovatis imbricatis, alis junioribus longè cuspidatis, foliis linearibus 4^{im}-verticillatis, caule erecto ramoso angulato. ① ad ripas rivulorum Americæ borealis. (v. s.)
83. *P. BREVIFOLIA* (Nutt. gen. am. 2. p. 89.) racemis spiciformibus partim capitatis, alis cordato-ovatis acutis, foliis oblongo-linearibus resinoso-punctatis 4^{im}-verticillatis, caule erecto ramoso alato. ① in arenosis paludosis New-Jersey et Ohio.
84. *P. FASTIGIATA* (Nutt. gen. am. 2. p. 89.) racemis spiciformibus subcapitatis, alis patentibus acutis, foliis alternis linearibus acutis, caule gracili fastigiatim ramoso. ① in New-Jersey.
85. *P. LUTEA* (Lin. sp. 990.) racemis spiciformibus densis ovatis, alis ovatis mucronatis, foliis oblongo-lanceolatis acutis, radicalibus subpetiolatis, caulibus simplicibus elongatis. ① à New-Jersey ad Floridam. Pluk. t. 438. f. 6. (v. s.)
86. *P. NANA*, racemis spiciformibus densis cylindraceis, alis ovatis cuspidato-acuminatis, foliis oblongo-spatulatis petiolatis, caule simplici vix foliis longiore. ① in udibus Carolinæ et Georgiæ. *P. viridescens*. Nutt. 2. p. 88. non alior. *P. lutea* β Mich. — (v. s.)
87. *P. ANGULATA*, racemis spiciformibus brevibus 8-10-floris, alis oblongis, foliis ovalibus cuspidato-mucronatis, caulibus ramosis adscendentibus angulato-alatis fruticosis. 2 5 in Brasiliâ.
88. *P. PURPUREA* (Nutt. gen. 2. p. 88.) racemis spiciformibus ovatis imbricatis, alis ovatis capsulâ duplò longioribus, foliis alternis oblongo-linearibus, caule ramoso-fastigiato erecto. ① in Amer. bor. *P. sanguinea*. Mich. fl. 2. p. 52. (v. s.)
89. *P. SANGUINEA* (Lin. sp. 991.) racemis spiciformibus oblongis, alis obovatis longitudine capsulæ, foliis alternis angustè linearibus, caule erecto fastigiato ramoso. ① in Carolinâ. *P. viridescens*. Poir! 5. p. 502. — in hac et in priore carina subimberbis. (v. s.)
90. *P. SETACEA* (Mich! fl. 2. p. 52.) racemis spiciformibus densis oblongis, foliis squamæformibus subnullis, caule erecto setaceo subsimplici. ① in Carolinâ et Georgiâ. Flores parvi subsessiles. (v. s.)
91. *P. GRACILIS* (H. B. et Kunth nov. gen. am. 5. p. 401.) racemis spiciformibus gracilibus laxiusculis, foliis sparsis linearibus, caule erecto elongato vix ramoso. ① in humidis prope Maypures Orinocensium. Glabra. Flores albi. Semina obliquè oblonga retrorsum pubescentia.
92. *P. SUBTILIS* (H. B. et Kunth. nov. gen. am. 5. p. 393. t. 506.) racemis spici-

- formibus densifloris, floribus dorso glandulosis, foliis squamæformibus remotis lineari-subulatis, caule erecto apice ramoso. ① in humidis ad ripam fl. Orinoci. Flores minimi albidi. Semina elliptica glabriuscula.
93. *P. tenuis*, racemis spiciformibus oblongo-cylindraceis, foliis sparsis lineari-setaceis, caule tenui erecto apice parçe ramoso. ① in Brasiliâ. Flores parvi albi. (v. s.)
94. *P. glochidata* (H. B. Kunth nov. gen. am. 5. p. 400.) racemis gracilibus spiciformibus laxiusculis, foliis angustè linearibus oppositis verticillatisve, rameis sparsis, caule erecto paniculato-ramoso. ① in humidis juxtâ Oriuocum et Magdalenam. Glabra. Flores pallidè carnei. Semina oblonga glochidato-pilosa.
95. *P. ambigua* (Nutt. gen. am. 2. p. 89.) racemis spiciformibus acutis longè pedunculatis, alis rotundis capsulæ æqualibus, foliis linearibus, infimis verticillatis, ceteris sparsis, caule erecto virgatim ramoso. ① in New-Jersey et Virginiâ.
96. *P. verticillata* (Lin. am. 2. p. 159.) racemis spiciformibus acutis pedunculatis, alis ovatis, foliis linearibus verticillatis, caule erecto ramoso. ① in Amer. bor. Pluk. t. 438. f. 4. (v. s.)
97. *P. galioides* (Poir! dict. 5. p. 503.) racemis spiciformibus gracillimis laxiusculis pedunculatis, alis ovatis, foliis oblongis 4im-verticillatis, caule erecto gracili parçe ramoso. ① in Cayennâ et insulis Caribæis. (v. s.)
98. *P. crucianelloides*, racemis spiciformibus gracilibus laxiusculis pedunculatis, alis oblongis, foliis rhombo-ovatis basi cuneatis verticillatis, summis alternis, caule erecto angulato. ① in Santo-Domingo. Nectoux. (v. s. in h. Mus. Par.)
99. *P. asperuloides* (H. B. et Kunth nov. gen. am. 5. p. 403.) racemis spiciformibus gracilibus laxiusculis, corollis dorso glandulosis, foliis oblongis pellucido-punctatis glabris, oppositis aut 3-6im verticillatis, caulibus diffusis fasciculato-cespitosis. ½ in Amer. calid. Semina oblonga retrorsum sericeo-pubescentia.
100. *P. paniculata* (Lin. am. 5. p. 402.) racemis subspiciformibus laxiusculis elongatis pedunculatis, alis ovali-oblongis vix capsulâ longioribus, foliis sparsis linearibus acutis, caule erecto ramosissimo. ① in insulis Caribæis et Amer. calid. — Sw! obs. 272, t. 6. f. 2.
 β. *Brasiliæna*, floribus minoribus, foliis latioribus. (v. s.)
 γ. *Africana*, foliis lineari-subulatis. — In Sierrâ Leonâ. (v. s.)
101. *P. scoparia* (H. B. et Kunth nov. gen. 5. p. 399.) racemis spiciformibus laxiusculis, foliis sparsis linearibus, caule erecto subfastigiato-ramoso. radice vermiculari tortuosa. ½ prope Mexico. Glabra. Flores albidi. Semina oblongo-cylindracea retrorsum sericeo-pilosa.
102. *P. corymbosa* (Michx. fl. 2. p. 54.) racemis plurimis in corymbum terminalem dispositis, alis oblongis cuspidatis, foliis radicalibus ovatis, caulibus linearibus, summis subulatis. ½ à Carolinâ ad Floridam in sphagnosis. *P. cymosa*. Walt. fl. car. 179. (v. s.)
103. *P. graminifolia* (Poir. dict. 5. p. 500.) racemis plurimis in racemum terminalem dispositis, alis oblongis acutis, foliis radicalibus lineari-spatulatis, caulibus lineari-subulatis. ½ in herbidis à Carolinâ ad Floridam. Priori nimis affinis. *P. attenuata*. Nutt. 2. p. 90. (v. s.)
104. *P. baldwinii* (Nutt. gen. am. 2. p. 90.) racemis capituliformibus corymbosis pedunculatis, alis lanceolatis cuspidatis, foliis radicalibus spatulatis, caulibus alternis lanceolatis, caule erecto angulato apice subramoso. ½ in Floridâ ad Sanctam-Mariam. (v. s. in h. Lamb.)
105. *P. Neesii*, racemis pluribus brevibus corymboso-confertis, alis oblongis acutis, foliis subulatis, caulibus è radice plurimis apice tantum ramosis. ①? in Peruvîâ aut Chili. Née. (v. s.)

106. *P. LIMOIDES* (Poir! dict. 5. p. 449.) racemis subspicatis elongatis, alis ovatis obtusis vix capsulâ longioribus, foliis linearibus acuminatis, caulibus adscendentibus vix ramosis. \mathcal{L} in Brasiliâ. circa Montevideo et Buenos-Aires. Flores certè cristati. Habitus ferè *Resedæ* glaucæ. (v. s.)
107. *P. ALBA* (Nutt. gen. 2. p. 87.) racemis spiciformibus pedunculatis, alis rotundatis corollæ longitudine, foliis alternis linearibus margine revolutis, caule simplici elongato. \mathcal{L} in Louisianâ superiore. (v. s. in h. Lamb.)
108. *P. RUBELLA* (Porsch! fl. 2. p. 464.) racemis subspicatis laxiusculis elongatis, alis ovalibus obtusissimis, foliis lanceolato-linearibus mucronatis caule erecto sulcato apice subramoso. \mathcal{L} à Pensylvaniâ ad Georgiam. (v. s.)

Sect. VI. SENECA.

Carina imberbis. Calycis sepala 3 exteriora subæqualia. Folia alterna. — Herbar Americanæ.

109. *P. SENECA* (Lin. sp. 990.) caulibus plurimis erectiusculis simplicibus teretibus, foliis ovato-lanceolatis, summis acuminatis, racemis subspiciformibus, alis orbiculatis, capsulâ ellipticâ emarginatâ \mathcal{L} in Amer. bor. Woodv. med. bot. 3. t. 93 Radix angulo exserto hinc longitudinaliter aucta. Var. floribus albidis et roseis. (v. s.)
110. *P. POLYGAMA* (Walt. fl. car. 179.) caulibus plurimis simplicibus erectis et procumbentibus foliis oblongis acutis, racemis inferioribus apetalis. \mathcal{L} in Carolinæ pinetis.
111. *P. MONTICOLA* (H. B. et Kunth nov. gen. am. 5. p. 405.) caulibus subsimplicibus folisque birto-pubescentibus remotis lanceolatis acutis, racemis terminalibus laxifloris, alis ellipticis basi unguiculatis, capsulâ emarginatâ. \mathcal{L} in aridis montis Tumiriquiri Novæ Andalusie.
112. *P. ANGUSTIFOLIA* (H. B. et K. l. c. p. 405. t. 511.) caule supernè ramoso puberulo, foliis remotis lineari-lanceolatis acutis margine revolutis glabris nervo venisque subtis puberulis, racemis spiciformibus laxiusculis. \mathcal{L} ad flum. Orinocum prope Carichanam.
113. *P. MOLLIS* (H. B. et K. l. c.) caulibus villosis simplicibus vel ramosis, foliis lanceolato-oblongis molliter caescentibus, racemis spiciformibus laxiusculis. \mathcal{L} in arenosis ad Carichanam. Semina retrorsum sericea. Alæ orbiculato-obovatae.
114. *P. NUTKANA* (Moc. ic. ined.) caulibus basi suffruticulosis adscendentibus, foliis ovalibus utrinquè acuminatis petiolatis, racemis laxis 4-5-floris, alis orbiculatis, capsulâ emarginatâ. \mathcal{L} in Amer. bor. orâ occidentali prope Nutka.
115. *P. HERECARPA*, caulibus simplicibus, foliis ovatis reticulato-venosis, racemis 10-12-floris, alis ovali oblongis, capsulâ ovali pubescente. \mathcal{L} \mathcal{L} in Sante-Domingo. An *P. ovata*. Poir. dict. 5. p. 498? (v. s.)
116. *P. VIOLACEA* (Vahl symb. 2. p. 79.) caule erecto ramis infimis subprocumbentibus, foliis ovalibus breviter petiolatis subvillosis, racemis 7-15-floris, alis obovatis capsulæ ovali glabræ aequalibus. \mathcal{L} in Cayennâ. (v. s.)
117. *P. CINEREA* (Willd. sp. 3. p. 880.) caule erecto ramoso hirsuto, foliis lanceolatis acuminatis subtomentosis, racemis 8-10-floris, carinâ margine fimbriatâ, capsulâ oblongâ glabrâ. $\textcircled{1}$ in pratis Guianæ et Cayennæ. *P. violacea*. Aubl. Guian. t. 294. (v. s. in h. Banks.)
118. *P. AMERICANA* (Mill! dict. n. 7.) caule erecto ramoso, juniore pubescente, foliis oblongis mucronatis velutino-pubescentibus, racemis multifloris, alis obovatis, capsulâ orbiculatâ velutinâ. \mathcal{L} \mathcal{L} in Americâ prope Vera-Cruz. (v. s. in h. Banks.)
119. *P. PUBESCENS* (Nutt. gen. am. 2. p. 87.) caule erecto pubescente virgatim ramoso, foliis ovato-lanceolatis ciliatis, carinâ apice glandulosâ. \mathcal{L} in Georghiâ circa Savannah. \dagger .

120. *P. CARACASANA* (H. B. et Kunth nov. gen. am. 5. p. 407.) caulibus procumbentibus et erectis, ramis foliisque hirsuto-pubescentibus ellipticis acutis, racemis spiciformibus terminalibus paucifloris, capsulâ ellipticâ emarginatâ pubescente. ζ in siccis prope Caracas.
121. *P. MONNINOIDES* (H. B. et K. l. c. p. 408.) ramis villosa-pubescentibus, foliis ovato-oblongis subacuminatis hirsuto-pubescentibus, racemis subterminalibus. ζ in montibus Novo-Granatensibus. Fruct. ign.
122. *P. RIVINÆFOLIA* (H. B. et K. l. c. p. 409. t. 512.) caulibus herbaceis subsimplicibus pubescentibus, foliis ovatis valdè acuminatis utrinquè puberulis ciliatis, racemis spiciformibus terminalibus multifloris. \mathcal{Z} in regno Mexicano prope Ario. Capsula *Thlaspeos arvensis* elliptico-orbiculata emarginata glabra ciliata.
123. *P. OVALIFOLIA*, caulibus teretibus subvelutinis foliis ovalibus breviter petiolatis, racemis 7-8-floris, alis obovatis, capsulâ ovali pubescente. ζ \mathcal{Z} in Novâ-Hispaniâ. (v. s. in h. Deless.)
124. *P. HEBECLADA*, caulibus teretibus villosis, foliis linearibus pubescentibus, racemis elongatis, alis obovatis, capsulâ oblongâ glabrâ. ζ in Brasiliâ (v. s. in h. Mus. Par.)
125. *P. BUXIFOLIA* (H. B. et Kunth l. c. p. 407.) ramosa diffusa, ramis foliisque hirsuto-pubescentibus oblongo-ellipticis apice rotundatis subemarginatis margine revolutis, racemis paucifloris terminalibus et lateralibus. ζ prope Sancta Rosa Mexicanorum.
126. *P. GLANDULOSA* (H. B. et Kunth l. c. p. 404. t. 510.) caulibus ramissimis procumbentibus punctato-glandulosis, foliis obovato-cuneatis mucronatis enerviis subpuberulis pellucido-punctatis, floribus pedicellatis subsolitariis extra-axillaribus aut oppositifoliis. ζ in Novâ-Hispaniâ. Semina retrorsum pubescentia. *Viola punctata* Willd. ex Schlect. in Roem. et Schult. syst. 5. p. 391.

Sect. VII. *CHAMÆBUXUS*. *Dill. nov. gen. t. 9.*

Calyceis sepalum inferius concavo-cucullatum intus ad basin glandulam fovens. Corollæ carina subcristata aut apice callosa. Flores pauci, magui. Herbæ aut suffrutescentes.

127. *P. PAUCIFOLIA* (Willd. sp. 3. p. 880.) caulibus simplicissimis erectis infernè nudis, foliis ovatis, floribus ternis terminalibus, carina cristatâ. \mathcal{Z} à Pensylvaniâ ad Virginiam. — *P. purpurea*. Ait. Kew. ed. 2. v. 4. p. 244. non Nutt. *Triclisperma grandiflora*. Raf. specch. 1. p. 117. (v. s.)
128. *P. UNIFLORA* (Michx. fl. 2. p. 58.) caule adscendente subsimplici, foliis ovalibus acutis, floribus solitariis pedicellatis sparsis, carinâ imberbi. \mathcal{Z} in finibus Canadæ ad lacum Ontario. (v. s. sine fl.)
129. *P. CHAMÆBUXUS* (Lin. sp. 989.) caulibus suffruticosis ramosis procumbentibus, foliis oblongo-lanceolatis mucronatis, racemis 1-2-floris, carinâ cristatâ. \mathcal{Z} ζ in montanis sylvaticis Europæ. Jacq. fl. austr. t. 233. Flores flavidi apice purpureo notati. (v. v.)
130. *P. VENENOSA* (Juss. in Poir. dict. 5. p. 493.) caule fruticoso, foliis petiolatis obovatis acuminatis maximis, floribus racemosis, carinâ cristatâ. ζ in Javâ. (v. s.)
131. *P. OXYPHYLLA*, caule fruticoso, foliis oblongis acuminatissimis basi attenuatis, floribus breviter racemosis, carinâ imberbi, alis ovalibus. ζ in Brasiliâ. (v. s. in h. Mus. Par.)
132. *P. SPECTABILIS*, caule fruticoso, foliis ovali-oblongis acuminatis basi attenuatis, summis suboppositis, floribus longè racemosis, carinâ imberbi, alis orbiculatis. ζ in Parâ Amer. Austr. (v. s. in h. Mus. Par.)
133. *P. CHINENSIS* (Lin. sp. 989.) caule suffruticoso prostrato, foliis ovalibus obtusis, floribus breviter racemosis, carinâ imberbi, alis obovatis. ζ in Indiâ. (v. s. in h. Juss.) An hujus sect.?

Sect. VIII. BRACHYTROPIS.

Carina imberbis alis corollinis calycinisque multò brevior. — An genus proprium Comespermæ affine?

134. *P. MICROPHYLLA* (Lin. sp. 989) caulibus fruticulosus virgatis, foliis minutissimis ellipticis, racemis axillaribus sessilibus 8-10-floris. ζ in dumetis Lusitaniz. Flores cærulei. Fructus ignotus. (v. s.) — *P. juniperina* Cav. anal. cienc. nat. 1801. 4. p. 53. ab eâ non videtur diversa sed descr. non sufficit.

† *Polygalæ non satis notæ.*

§. 1. *Carinâ cristatâ.*

135. *P. LONGIFOLIA* (Poir. dict. 5. p. 501.) floribus cristatis, racemis spiciformibus, alis oblongis, caule subsimplici, foliis longissimis graminiformibus, superioribus filiformibus. $\textcircled{1}$ in Javâ. Nulli sectioni affinis videtur. (v. s. in Mus. Par.)
136. *P. ELONGATA* (Klein. in Willd. sp. 3. p. 879.) floribus cristatis, racemis axillaribus elongatis, caule basi ramofo, foliis linearibus obtusis mucronatis margine subciliatis. $\textcircled{1}$ in Indiâ orientali.
137. *P. TELEPHIOIDES* (Willd. sp. 3. p. 876.) floribus cristatis, racemis axillaribus folio duplò longioribus, caule simplici erecto, foliis oblongis obtusis basi attenuatis. $\textcircled{1}$ in Indiâ orientali.
138. *P. TRANQUEBARICA* (Mart. ex Goett. anz. 1817. p. 159.) floribus cristatis, racemis paucifloris lateralibus, caule herbaceo ramofo, foliis linearibus mucronatis. — In littore Coromandeliano.
139. *P. MARTII*, floribus cristatis, racemis lateralibus, foliis lanceolato-linearibus obtusiusculis, caule procumbente herbaceo. — *P. pubescens*. Mart. ex Goett. anz. 1817. p. 159. non Nutt.
140. *P. VARIANS* (Mart. l. c.) floribus cristatis, racemis axillaribus, foliis inferioribus obcordatis ovatisve, superioribus lanceolatis, caule herbaceo ramofo procumbente, pedunculis hirtis. — In Bengalo.
141. *P. UMBROSA* (Mart. l. c.) floribus cristatis, racemis axillaribus, foliis oblongis acutiusculis basi attenuatis. — In Bengalo.
142. *P. MACROPTERA*, floribus cristatis, alis cordato-orbiculatis, racemis axillaribus, foliis obovatis apice emarginatis. ζ in Madagascar. — Vouné Flac. mad. p. 136. n. 94. ic. ? (v. s. in h. Mus. Par.)
143. *P. RARIFOLIA*, floribus cristatis, alis obovatis capsulâ ellipticâ longioribus, foliis linearibus acutis, caule suffruticoso erecto sulcato virgato. \mathcal{L} \mathcal{S} in Sierrâ Leonâ. (v. s. in h. Lamb.) An sect. propria?
144. *P. GUINEENSIS* (Willd. sp. 3. p. 882.) floribus cristatis nutantibus, racemis axillaribus secundis, foliis filiformibus, caule ramofo-herbaceo. — In Guineâ.
145. *P. CERNUA* (Thunb. prod. 120.) floribus cristatis racemosis, foliis lanceolatis glabris. — Ad cap. B.-Spei.
146. *P. VIRGATA* (Thunb. prod. 120.) floribus cristatis racemosis, foliis obovato-oblongis. — ad cap. B.-Spei. An Mundiz species?
147. *P. AMENA* (Thunb. prod. 120.) floribus cristatis lateralibus, foliis obovatis obtusis glabris. — Ad cap. B.-Spei.
148. *P. TOMENTOSA* (Thunb. prod. 120.) floribus cristatis verticillatis, foliis cordatis subtus tomentosis. — Ad cap. B.-Spei.
149. *P. SULCATA* (Willd. sp. 3. p. 878.) floribus cristatis, racemis terminalibus pedunculatis, canibus erectis sulcatis superne ramosis, foliis linearibus adpressis. $\textcircled{1}$ in Montevideo Brasiliz. Affinis *P. paniculatæ*. Flores violacei.
150. *P. TENELLA* (Willd. sp. 3. p. 878.) floribus cristatis, racemis terminalibus

longissimè pedunculatis, caulibus erectis supèrne ramosis, foliis linearibus ovatis mucronatis. ① in Panamâ. Affinis *P. paniculatæ*. Flores incarnati.

151. *P. BRASILIENSIS* (Lin. mant. 99.) floribus cristatis subspicatis, caulibus prostratis simplicissimis, foliis lanceolatis. — In Brasiliâ. — Videtur ad *Timutuas* referenda.

152. *P. MULTICAULIS* (Tausch. in Flor. 1821. p. 563.) floribus cristatis, racemis laxis paucifloris, alis calycinis ellipticis acutis corollâ brevioribus grossè trinerviis margine venosis, caulibus filiformibus ramosissimis humifusis, foliis lineari-lanceolatis acutis. \mathcal{N} in pratis montium Sudetorum Iseræ majoris. — Ad *Polygali* sect. referenda videtur.

§. 2. *Carinâ imberbi.*

153. *P. TRIFLORA* (Lin. fl. zeyl. 269.) floribus imberbibus, pedunculis subtrifloris folio brevioribus, caule herbaceo erecto, foliis linearibus alternis. ① in Ceylonâ.

154. *P. PROSTRATA* (Willd. sp. 3. p. 896.) floribus imberbibus, pedunculis multifloris, caulibus diffusis herbaceis, foliis lanceolatis obtusis. ① in Indiâ orientali. — Affinis *P. glaucoïdi*.

155. *P. MULTIFLORA* (Poir. dict. 5. p. 497.) floribus imberbibus, racemis longis terminalibus? caulibus herbaceis ramosis virgatis, foliis linearibus raris. — In Sierrâ-Leonâ.

156. *P. THUNBERGII*, floribus imberbibus racemosis, foliis ovatis inermibus. 5 ad cap. B.-Spei. *P. microphylla*. Thunb. prod. 121. non Lin.

157. *P. MUCRONATA* (Willd. sp. 3. p. 888.) floribus imberbibus, racemis terminalibus? caule fruticoso, ramis pilosis, foliis lanceolatis mucronatis subtus pubescentibus. 5 in Amer. calid.

158. *P. MYRTILLOIDES* (Willd. sp. 3. p. 889.) floribus imberbibus, racemis oppositifoliis, caule fruticoso, ramis procumbentibus, foliis subrotundo-ovatis reticulato-venosis. 5 in Americâ australi.

159. *P. MEXICANA* (fl. mex. ic. ined.) floribus imberbibus demùm cernuis, racemis terminalibus? subspicatis, caulibus simplicibus erectiusculis foliis linearibus acutis. \mathcal{N} 5 in Mexico.

160. *P. LINEARIFOLIA* (Poir. dict. 5. p. 495.) floribus subimberbibus? racemis terminalibus? densis ovatis, caule herbaceo tereti, foliis linearibus summis verticillato-congestis. — Hab. An *Muraltix* species?

161. *P. UNGUICULATA* (Poir. dict. 5. p. 494.) floribus imberbibus congestis axillaribus, petalis 2 unguiculatis, calycibus 4-sepalis, caule fruticoso, foliis ovatis mucronatis. 5 Hab. An genus proprium?

162. *P. RAMOSISSIMA* (Cav. annal. cienc. nat. 1801. 4. p. 53.) caule herbaceo ramosissimo, foliis linearibus, spicis terminalibus, floribus minutis. — In ins. Teneriffæ ex Broussonet.

163? *P. THEEZANS* (Lin. mant. 260. excl. Burm. syn. 2.) floribus imberbibus solitariis, pedunculis unifloris axillaribus, foliis alternis lineari-lanceolatis. 5 in Javâ et Japoniâ. — Icon. Burm. fl. zeylan. p. 195. t. 85. est certò Ionidii species; *P. thea* Burm. fl. ind. p. 154. est *Leptospermi* species; ut videtur ex specimenibus à Burmanno collectis, et in herb. Delessert. servatis. Quid sit ergo Linnæi *P. theezans*?

N. B. *P. triphylla* et *P. pinnata*. Burm. prod. fl. cap. p. 20. videntur *Leguminosarum* species.

II. SALOMONIA. *Lour. coch. ed. W. p. 18.*

Calyx 5-sepalis, sepalis subæqualibus. Corollæ tubus longitudinaliter fissus, limbus 3-fidus, carina cucullata. Stam. monadelphæ, Antheræ 4. Capsula biloba compressa sæpius ciliato-serrata. — Herbæ minimæ Asiaticæ.

1. *S. CANTONIENSIS* (Lour! loc. cit.) glabra erecta ramosa, foliis subpetiolatis cordatis, summis subovatis, capsulis margine cristato-dentatis. ① circa Cantonem Sinarum. *Salmonia Cantoniensis*. Vahl enum. 1. p. 8. (v. s. in h. Mus. Par.)
2. *S. EDENTULA*, glabra erecta ramosa, foliis subpetiolatis latè ovatis mucronatis, capsulis margine subedentulis. ① in Napauliâ. Wallich. (v. s.)
3. *S. OBLONGIFOLIA*, glabra erecta simplicissima, foliis oblongis ellipticisve, capsulis margine cristato-dentatis. ① in Napauliâ. Wallich. (v. s.)
4. *S. CILIATA*, caule erecto ramoso pedunculisque hirtis, foliis lanceolatis obtusis, capsulis ciliato-dentatis. ① in Indiâ orientali. *P. ciliata*. Lin. sp. 991.

III. COMESPERMA. Labill. specim. nov.-holl. 2. p. 21.

Cal. 5 sepalus deciduus, sepalis 2 interioribus alæformibus. Petala coalita in corollam 3-partitam, lobo medio imberbi integro aut emarginato. Stam. 8 ferè monadelpha. Capsula bilocularis spathulata basi longè attenuata. Semina apice longè comosa, carunculo lineari.

1. *C. RETUSA* (Labill! loc. cit. t. 160.) glabra erecta, foliis oblongis obtusis, racemis contractis, corollæ lobo medio integro. 5 ad cap. Van-Diemen. (v. s.)
2. *C. SECUNDA* (Banks! herb.) pubescens erecta, foliis ovatis mucronatis, racemis secundifloris. 5 in Nov.-Holl. prope flumen Endeavour. (v. s. in h. Banks.)
3. *C. CONFERTA* (Labill! l. c. t. 161.) glabra erecta, foliis linearibus confertis erectis margine revolutis, racemis contractis, corollæ lobo medio emarginato. 5 in terrâ Van Leuwin. folia 7-8. lin. longa. (v. s.)
4. *C. ERICINA*, glabra erecta, foliis linearibus, margine subrevolutis obliquè erectis, racemis laxiusculis, corollæ lobo medio integro. 5 in Novæ-Hollandiæ orâ orientali. folia 3-4. lin. longa. (v. s.)
5. *C. FLAVA*, glabra erecta, foliis linearibus planis erectis, racemis contractis obtusis, corollæ lobo medio integro. 5 in Novæ-Hollandiæ orâ orientali. Lechenault. Flores flavi nec ut in aliis purpurascens. (v. s.)
6. *C. CALYMEGA* (Labill! l. c. t. 162.) glabra herbacea erecta, foliis lanceolatis, alis vix cetera sepala superantibus, petali lobo medio integro. 4 in promont. Van Diemen.
7. *C. NUDUSCULA*, glabra erecta virgata, foliis linearibus subsetaceis acutis, summis minimis, racemis laxiusculis, alis vix cetera sepala superantibus. 5 in Novâ-Hollandiâ ad portum regis Georgii. Lechenault. (v. s.)
8. *C. VIRGATA* (Labill! l. c. t. 159.) glabra erecta virgata, foliis lanceolato linearibus obtusè acuminatis, racemis elongatis, corollæ lobo medio emarginato. 5 in Novâ-Hollandiâ. (v. s.)
9. *C. VOLUBILIS* (Labill! l. c. t. 163.) glabra herbacea volubilis, foliis paucissimis lanceolatis, racemis contractis, corollæ lobo medio dentato. 4 in insulâ Van-Diemen. (v. s.)

IV. BADIERA. — Penæa. Plum. gen. p. 22. t. 25. non Lin.

Cal. 5-sepalus, deciduus, sepalis subæqualibus. Petala 3 imâ basi subcoalita, medio concavo imberbi. Stam. 8-monadelpha. Capsula compressa, obcordata, bilocularis, margine sulcata. Semina glabra arillo maximo oleoso loculum supernè implente. — Suffrutices Americani.

1. *B. DIVERSIFOLIA*, racemis brevibus axillaribus, foliis ovali-oblongis utrinquè acuminatis. 5 in Jamaicæ sylvis. *Polygala diversifolia*. Lin. am. 2. p. 140. — Brown! jam. t. 5. f. 3-4. (v. s.)

2. *B. DOMINGENSIS*, racemis brevibus axillaribus, foliis obovatis obtusis. ♂ in montosis Santi-Domingi. Pol. Domingensis Jacq. am. ed. min. 252. (v. s.)
3. *B. PENSA*, floribus solitariis axillaribus pedunculatis, foliis obovatis obtusis. ♂ in Amer. calid. Pol. Pensa. Lin. am. 2. p. 140. Plum. ed. Burm. t. 214. f. 1.
4. *B? ACUMINATA*, racemis lateralibus oppositifoliis folio longioribus, foliis oblongis acuminatis. ♀ in Novâ-Hispaniâ. Pol. acuminata. Willd. sp. 2. p. 887. Similis *B. diversifoliæ* ex W.
5. *B? DIVARICATA* (racemis axillaribus paniculatis velutinis, foliis ovali-oblongis glabris, alis obovatis. ♀ in Parâ Amer. merid. An hujus generis? (v. s. in h. Mus. Par.)

V. SOULAMEA. Lam. dict. 1. p. 449.

Cal. 5-sepalus, sepalis 3-exterioribus minimis, 2 interioribus majoribus concavis. Petalum 1 concavum. Stam. 6? Capsula (samara) non dehiscens, compressa, suberosa, orbiculata, emarginata, bilocularis. Semina exalbuminosa.

1. *S. AMARA* (Lam. l. c.) ♀ in insulis Moluccanis. Rex amaroris. Rumph. amb. 2. p. 129. t. 41. Folia ampla ovali-oblonga ferè Polygalæ venenosæ. Flos non satis notus. (v. s.)

VI. MURALTIA. Neck. elem. n. 1382. — Heisteria. Berg. cap. 185. non Lin.

Calyx 5-sepalus glumaceus, sepalis subæqualibus. Petala 3 coalita, medio bifido, lobis obtusis. Ovarium cornubus tuberculisve 4 coronatum. Capsula 4-cornuta aut 4-tuberculata, bivalvis, bilocularis. — Suffrutices omnes ad cap. B.-Spei nascentes.

* *Foliis apice mucronato-pungentibus.*

1. *M. HEISTERIA*, foliis triquetris mucronato-spinosis rigidis fasciculatis, ramulis puberulis, floribus axillaribus sessilibus ♂ Heisteria pungens Berg. cap. 185. Pol. Heisteria. Lin. sp. 989. curt. bot. mag. t. 340. Cornua capsulâ longiora. (v. s.)
- β. *pilosa*, foliis planiusculis pilosis ciliatisque. — An species propria? (v. s.)
2. *M. CONFERTA*, foliis oblongo-lanceolatis fasciculatis confertis mucronato-subspinosis margine scabris glabris, ramulis hirsutis, floribus axillaribus sessilibus. ♂ an Pol. thymifolia. Thunb. prod. 121? Cornua capsulâ longiora. (v. s.)
3. *M. BREVIFOLIA*, foliis elliptico-oblongis mucrone recurvo spinulosis sparsis, junioribus ramulisque pubescentibus, floribus axillaribus sessilibus. ♀ Capsula pubescens 4-cornuta. (v. s. comm. à cl. Lambert.)
4. *M. SERPYLLOIDES*, foliis ovatis mucronato-spinosis margine subrevolutis in nervo subtus pilosis ciliatisque, ramis pilosis, floribus axillaribus sessilibus. ♀ Cornua capsulâ breviora. (v. s. in h. Deless.)
5. *M. ALOPECUROIDES*, foliis floralibus ovatis mucronato-subspinosis confertissimis glabris ciliatis, inferioribus sublinearibus, ramis pilosiusculis, floribus axillaribus sessilibus. ♂ Polygala alopecuroides. Lin. mant. 260. Sims. bot. mag. t. 1006. Cornua capsulæ longitudine. (v. s.)
6. *M. TRINERVIA*, foliis ovato-cordatis cuspidato-pungentibus trinerviis glabris, ramis angulatis glabris, floribus axillaribus pedicellatis. ♂ Pol. trinervia Thunb. prod. 121. Capsula videtur bicornis. (v. s. in h. Banks.)
7. *M. SQUARROSA*, foliis lanceolatis mucronato-spinosis fasciculatis rigidulis, superioribus patulis, inferioribus reflexis, ramulis villosis, floribus axillaribus sessilibus. ♂ Pol. squarrosa. Thunb. prod. 121. Muraltia n. 3493. Burch! cat. videtur hujus varietas. (v. s. in h. Deless.)

8. *M. STIPULACEA* (Burch! trav. 1. p. 19. 29.) foliis linearibus subulatis mucronatis subtus punctatis mucronatis glabris subfasciculatis distantibus, ramulis subpubescentibus, floribus axillaribus sessilibus. § Pol. stipulacea Lin. mant. 260. excl. Seb. syn. Andr. bot. rep. t. 263. capsula 4-cornuta. (v. s.)
9. *M. JUNIPERIFOLIA*, foliis triquetris subulatis mucronatis rectis ramisque glabris, floribus axillaribus subpedicellatis, capsulis subinermibus 4-tuberculatis. § Pol. juniperifolia. Poir. dict. 5. p. 496. (v. s. in h. Deless.)
10. *M. ERICÆFOLIA*, foliis trigonis subulatis mucronatis rectis fasciculatis glabris, ramulis subpubescentibus, floribus axillaribus sessilibus, capsula 4-cornuta. § An Pol. mixta Thunb. prod. 121? non Lin. (v. s.)
11. *M. SPRENGELIOIDES* (Burch! cat. n. 4957.) foliis subtrigonis mucronato-pungentibus sparsis demum patulis ramisque glabris, floribus axillaribus subpedicellatis. § (v. s. comm. à cl. Busch.)
12. *M. SATURNEOIDES* (Burch! cat. n. 5617.) foliis subtrigonis mucronato-pungentibus fasciculatis margine subpuberulis, ramis pubescentibus, floribus axillaribus sessilibus. § Folia vix 2 lin. longa. Capsula 4-rarius casu 3-cornuta. (v. s.)
13. *M. TENUIFOLIA*, foliis subulatis apice recurvis mucronatis patulis fasciculatis tenuissimis, ramis pubescentibus, floribus axillaribus subsessilibus. § Pol. tenuifolia. Poir. dict. 5. p. 497. non Willd. Capsulae cornua 4 divergentia (v. s. in h. Desf.)
14. *M. ASPALATHA*, foliis superioribus linearibus subtrigonis acuminato-pungentibus longe pilosis, inferioribus subulatis glabris, ramis pilosis, floribus axillaribus sessilibus, capsula pubescente 4-cornuta. § (v. s.)
15. *M. DIFFUSA* (Burch! cat. n. 916 et 354.) foliis linearibus acuminato-pungentibus subsparis distantibus, ramis tenuissimis diffusis, floribus axillaribus sessilibus. § Variat ramulis foliisque junioribus glabris aut pubescentibus. Pol. micrantha Willd? (v. s.)
16. *M. VIRGATA* (Burch! cat. n. 7074 et 7104.) foliis lineari-subulatis pungentibus sparsis rectis ramisque tenuissimis glabris, floribus axillaribus sessilibus, sepalis oblongis acuminatissimis. § Pol. micrantha Thunb. prod. 121? Caules plurimi à radice. (v. s.)
17. *M. LINOPHYLLA* (Burch! cat. n. 510.) foliis linearibus ferè subulatis subpungentibus sparsis ramisque tenuissimis subpuberulis glabrisve, floribus axillaribus sessilibus, sepalis ovato-oblongis mucronatis. § Priori valde affinis, sed calyce differt. An Pol. filiformis Thunb. prod. 121? (v. s.)
18. *M. MACROCRAS*, foliis lineari-subulatis mucronatis fasciculato-confertis rectis ramisque tenuissimis glabris, floribus axillaribus sessilibus, cornubus capsula longioribus. § Burch! cat. n. 3985. Multicaulis. (v. s. comm. à cl. Lambert.)
19. *M. MIXTA* (Lin. f. suppl. 316.) foliis lineari-subulatis mucronulatis fasciculato-confertis rectis, ramisque tenuissimis glabris, floribus axillaribus sessilibus, cornubus capsula brevioribus. § Pol. mixta Andr. bot. rep. t. 445. (v. v.)
20. *M. CILIARIS*, foliis lineari-subulatis mucronatis fasciculatis patentibus ramisque hispido-ciliatis, floribus axillaribus sessilibus, capsulis subinermibus. § Burch! cat. n. 5566? (v. s. comm. à cl. Lambert.)
 β. *laxiuscula*, foliis brevioribus magis distantibus. — An species propria? (v. s.)

§. 2. Foliis apice obtusiusculis.

21. *M. PUBESCENS*, foliis linearibus fasciculatis patulis, ramisque velutino-pubescentibus, floribus axillaribus solitariis. § Species distinctissima. (v. s. comm. à cl. Lamb.)
22. *M. DEPRESSA* (Burch! cat. n. 6264.) foliis subtrigonis callosis glabris fasci-

chlato-rosulatis, ramulis junioribus pubescentibus, floribus axillaribus sessilibus. ♂ Fruticulus humillimus ramoissimus demissus. (v. s. comm. à cl. Burch.)

23. *M. BREVIORNIS*, foliis linearibus obtusis glabris extus basi tuberculatis sparsis, ramis virgatis glabris, floribus axillaribus sessilibus, capsulæ cornubus brevissimis. ♂ species distinctissima. (v. s. comm. à cl. Lamb.)
24. *M. OBOVATA*, foliis obovatis obtusis subfasciculatis ramulisque glabris, floribus axillaribus sessilibus. ♂ habitus ferè Mundiaë, sed inermis. (v. s. comm. à cl. Lamb.)

† *Muraltiæ non satis notæ* (1).

25. *M. PHYLIROIDES*, foliis lineari-lanceolatis mucronatis margine revolutis? subtus ramulisque pubescentibus. ♂ Polyg. phylcoïdes Thunb. prod. 121.
26. *M. THYMIFOLIA*, foliis oblongo-lanceolatis carinatis mucronatis denticulatis ramisque divaricatis glabris. ♂ Polyg. thymifolia Thunb. prod. 121. an Willd?
27. *M. PILOSA*, foliis lanceolatis mucronatis rigidis pilosis, ramulis velutinis. ♂ Polyg. pilosa. Poir. dict. 5. p. 497.
28. *M. FILIFORMIS*, foliis trigonis mucronatis solitariis. ♂ Polyg. filiformis Thunb. l. c.
29. *M. PAUCIFLORA*, foliis linearibus mucronatis, floribus axillaribus pedunculatis. ♂ Polyg. pauciflora Thunb. l. c.
30. *M?* *BURMANNI*, foliis ovato-linearibus adpressis strictis. ♂ Polyg. ericoïdes Burm. fl. cap. prod. p. 20.
31. *M?* *POIRETI*, foliis subulatis ramisque virgatis glabris, floribus axillaribus sessilibus congestis. ♂ Polyg. ericoïdes Poir. dict. 5. p. 497. An eadem ac *M. virgata*?
32. *M. DUMOSA*, foliis sublinearibus obtusis apice reflexis, floribus sessilibus solitariis. ♂ Polyg. dumosa Poir. dict. 5. p. 495.
33. *M?* *LAXA*, foliis lanceolatis solitariis, floribus racemosis. ♂ Polyg. laxa Thunb. prod. p. 121.
34. *M?* *FASCICULATA*, foliis tenuissimis subciliatis, ramis fasciculatis, floribus sessilibus solitariis aut geminis. ♂ Polyg. fasciculata Poir. dict. 5. p. 502. An eadem ac *M. ciliaris*?
35. *M. STRIATA*, foliis subteretibus inermibus. ♂ Pol. striata Thunb. prod. 121.
36. *M?* *PARVIFLORA*, foliis setaceis glabriusculis minutis stipulatis? ramis gracilibus, floribus axillaribus solitariis sessilibus (1) Polyg. parviflora Poir. dict. 5. p. 504.
37. *M. HUMILIS*, foliis lanceolato-linearibus erectis sparsis, caule humili ramoississimo, floribus axillaribus sessilibus solitariis. ♂ Polyg. humilis Lodd. bot. cab. 5. t. 420.

VII. MUNDIA. Kunth nov. gen. am. 5. p. 392. in adn.

Calyx glumaceus persistens 5-sepalus, sepalis 2 interioribus alaeformibus. Petala 3 basi vix coalita, medio cucullato imberbi. Stam. 7-8 subvillosa basi monadelpha tubo anticè fisso. Drupa bi-et abortu unilocularis 1-sperma. Alhumen carnosum. — Frutices ramosissimi, ramis apice spinescentibus, foliis coriaceis integerrimis. — Genus calyce ferè Muraltiæ et fructu Monniæ donatum, jam à decennio à nobis admissum et sub nomine Vascoe amicis communicatum.

(1) Pleræque forsàn jam inter priores enumeratæ, sed ob descriptiones valdè incompletas non dignoscendæ.

1. *M. spinosa*, ♂ ad Cap. B.-Spei. *Polygala spinosa* Lin. amœn. 2. p. 141. *Ulex Capensis* Lin. sp. 1046. ex Burm! herb.

a. latifolia, foliis obovatis ovalibusve. ramis brevibus spinosis. (v. s.)

β. angustifolia, foliis oblongis lanceolatisve, ramis virgatis. *Polyg. viminea* Houtt! in herb. Deless. (v. s.)

- VIII. *MONNINA*. Ruiz et Pav. *fl. per. syst.* 1. p. 169. — *Kunth. nov. gen. am.* 5. p. 409. — *Hebeandra*. *Bonpl. ges. berl.* 1808. p. 40.

Flores resupinati. Calyx 5-sepalus deciduus, sepalis 2 internis alæformibus, 3 exterioribus ovatis, 2 quorum sæpè coalitis. Petala basi connata, medio concavo sæpè 3-dentata. Stam. 8, filam. subdiadelphis pilosiusculis. Drupa 2-loc. 2 sperma aut abortu 1-loc. 1-sperma, interdum margine membranaceo cincta. Semen ferè ex apice loculi pendulum. Albumen nullum. — Suffrutices aut herbæ in Amer. calid. spontaneæ. Species pleræque non satis notæ!

Sect. 1. HEBEANDRA.

Drupæ apteræ nec margine cinctæ.

* *Racemis compositis.*

1. *M. POLYSTACHYA* (Ruiz et Pav! l. c.) fruticosa, foliis ovatis acutiusculis, paniculis ramulisque villosis. ♂ in Peruvix Andium nemoribus ubi vulgò dicitur Yalhoi. Ruiz diss. madr. 1805. ic. (v. s.)
2. *M. LIGUSTRIFOLIA* (Kunth nov. gen. 5. p. 417.) foliis oblongo-lanceolatis acutis basi attenuatis reticulato-venosis membranaceis, ramulis paniculisque pubescentibus, racemis subternis. ♂ in regni Quitensis locis calidis prope Ayavacam.
3. *M. LATIFOLIA* (Bonpl. l. c. sub Heb.) herbacea, foliis obovato-oblongis acuminatis subtus pubescentibus, ramis sulcatis, racemis paniculatis. ♀ in monte Quindiu.
4. *M. PARVIFLORA* (Kunth. nov. gen. am. 5. p. 419.) foliis ovato-oblongis basi angustatis apice attenuatis subtus pubescentibus reticulato-venosis membranaceis, ramis teretibus molliter hirtellis, racemis paniculatis subcorymbosis. ♀ in monte Quindiu regni Novo-Gran. Heb. parviflora Bonpl. l. c.
5. *M. FASTIGIATA* (Bonpl. l. c. sub Heb.) foliis oblongis utrinquè acuminatis subtus molliter pubescentibus, racemis compositis. ♀ in monte Quindiu. An à priore satis differt?
6. *M. PUBESCENS* (Kunth l. c. 5. p. 418.) foliis oblongis acutis submucronatis basi attenuatis reticulato-venosis membranaceis supra puberulis subtus ramulisque hirtis pubescentibus, racemis solitariis et ternis. ♂ ad Caracas. Heb. pubescens Bonpl. l. c.
7. *M. PILOSA* (Kunth. nov. gen. am. 5. p. 419.) foliis oblongis acuminatis reticulato-venosis membranaceis supra pilosiusculis subtus præsertim nervo venisque pilosis, racemis compluribus fasciculato-corymbosis. ♀ prope Ayavacam Quitensium. Arborea.
8. *M. ÆSTUANS*, fruticosa, foliis lanceolatis petiolatis ramulisque subpubescentibus, racemis ramosis puberulis. ♀ in Novâ-Granadâ. *Polygala æstuaus* Lin! suppl. 315. (v. s. in h. Lin. et Banks.)
9. *M. CESTRIFOLIA* (Kunth nov. gen. am. 5. p. 413. t. 502.) foliis oblongo-lanceolatis acutis basi angustatis margine revolutis venosis coriaceis, ramulisque glabris, racemis compluribus corymboso-congestis. ♂ in Amer. calid. prope urbem Quito, si saltem huc referenda sit. Heb. cestrifolia Bonpl. l. c.?

10. *M. PHYTOLACCEIFOLIA* (Kunth l. c. t. 503.) foliis elliptico-oblongis utrinquè acutis vel oblongis utrinquè angustato-acutis reticulato-venosis crassiusculis glabris, ramulis pilosiusculis, racemis subternis. ♂ *Hebeandra padifolia* Bonpl. l. c. ?

a. ramulis pilosiusculis, foliis elliptico-oblongis utrinquè acutis. — Prope Mariquitam Novo-Granatensium.

β. ramulis villosio-hirtis, foliis oblongis utrinquè angustato-acutis. — In monte Avilæ prov. Caracasana. H. padifolia β. Bonpl. l. c.

** *Racemis simplicibus aut solitariis.*

11. *M. KALAPENSIS* (Kunth nov. gen. am. 5. p. 414.) foliis oblongis acuminatis basi angustatis reticulato-venosis membranaceis glabriusculis, racemis solitariis geminisve. ♂ prope Kalapam Mexic. Heb. evonymoides Bonpl. l. c.

12. *M. MEMOROSA* (Kunth l. c. p. 416. t. 504.) foliis elliptico-oblongis acutis mucronatis basi angustatis et revolutis obsolete denticulatis subvenosis crassiusculis glabris, junioribus ramulisque pubescentibus, racemis bifidis simplicibusve. ♂ in nemoribus regni Quitensis. Heb. mucronata Bonpl. l. c.

13. *M. MYRTILLOIDES* (Bonpl. l. c. sub *Hebeandra*.) foliis ellipticis utrinquè acutis subcoriaceis, racemis simplicibus. ♂ in Peruvæ nemoribus.

14. *M. OBTUSIFOLIA* (Kunth nov. gen. am. 5. p. 411.) foliis oblongis obtusis basi cuneatis obsolete venosis subcoriaceis ramulisque glabris, racemis solitariis. ♂ in regno Novo-Granatensi prope Almaguer. Heb. phylliræoides Bonpl. l. c. ?

15. *M. CRASSIFOLIA* (Kunth l. c. p. 411.) foliis lineari-oblongis apice rotundatis basi obtusis aveniis carnosio-coriaceis glabris, ramulis pubescentibus foliosis, racemis solitariis. ♂ in montibus Quitensibus. Heb. crassifolia Bonpl. l. c.

16. *M. REVOLUTA* (Kunth l. c. p. 412. t. 501.) foliis oblongo-lanceolatis acutis basi obtusiusculis margine revolutis aveniis coriaceis leviter hirtellis, ramulis pubescentibus, racemis solitariis. ♂ in Andibus prov. Pastoensis. Heb. revoluta Bonpl. l. c.

17. *M. RUTIFOLIA* (Kunth l. c. p. 415.) foliis oblongis subacuminatis mucronatis basi angustatis irregulariter undulato-denticulatis reticulato-venosis membranaceis glabriusculis, ramulis pilosiusculis, racemis solitariis. ♂ in monte Saragura Novo-Granatensium. An *M. celtastroides* aut *M. ligustrina* Bonpl. l. c. ?

18. *M. CONFERTA* (fl. per. syst. 1. p. 173.) fruticosa, foliis confertis oblongis acumine minimo, racemis brevibus corymbosis. ♂ in Andium nemoribus. Drupa ovata glabra. (v. s.)

19. *M. CROTALAROIDES*, fruticosa, foliis ellipticis utrinquè acuminatis, inferioribus ovatis, ramulis adpressè pubescentibus, racemis elongatis. ♂ in Amer. calid. Baccæ ovatæ nigræ. (v. s.)

20. *M. LANCEOLATA*, suffruticosa?, foliis lanceolatis acuminatis glabris, ramis divaricatis, racemis elongatis pedunculatis. ♂ ? in Peruvâ. *Polygala lanceolata* Poir! dict. 5. p. 498. (v. s. in h. Mus. Par.)

21. *M. SALICIFOLIA* (fl. per. syst. 1. p. 172.) fruticosa, foliis lanceolatis acumine minimo, ramis hirtis, racemis brevibus ferè pyramidalis. ♂ in Peruvâ præruptis. (v. s.)

22. *M. LONGIFOLIA*, suffruticosa, foliis angusto-lanceolatis longissimis, ramis subvelutinis, racemis brevibus laxiusculis pyramidalis, ♂ in Brasiliâ. *Polygala salicifolia* Poir! dict. 5. p. 498. (v. s. in h. Mus. Par.)

23. *M. BIFURCATA* (fl. mex. ic. ined.) herbacea, apice bifurca, foliis lanceolatis breviter petiolatis, summis suboppositis glabris, racemis terminalibus geminis. — In Mexico.

340 · DICOTYL. SEU EXOGENÆ.

24. *M. CILIOLATA* (fl. mex. ic. ined.) suffruticosa dichotoma, foliis oblongis acutis basi attenuatis, petiolo brevi ciliolato, racemis brevibus. ♂ in Mexico.
25. *M. LINEARIFOLIA* (fl. per. syst. i. p. 173.) suffruticulosa, foliis linearibus subsessilibus glabris, racemis longis infernè nudis. ♂ in Chili subalpinis. (v. s. in h. Mus. Par.)
26. *M. HERBACEA*, herbacea, apice subramosa glabra, foliis lanceolato-rhombis, racemis subelongatis, drupis pubescentibus apteris. ① prope Lima. (v. s.)

Sect. II. PTEROCARYA.

Fructus in alam membranaceam margine expansi.

27. *M. MACROSTACHYA* (fl. per. syst. i. p. 173.) herbacea, foliis lanceolato-rhombis, racemis longissimis, drupis glabris membranâ nervosâ brevi dentatâ cinctis. — In Peruvîæ collibus.
28. *M. PTEROCARPA* (fl. per. syst. i. p. 174.) herbacea glabra, foliis lanceolatis, inferioribus subovatis, racemis longissimis, drupis membranâ utrinquè emarginatâ cinctis. ♀ in Peruvîæ versuris. (v. s. in h. Mus. Par.)
29. *M. ANGUSTIFOLIA*, herbacea glabra; foliis linearibus, inferioribus sublan- ceolatis, racemis longissimis, drupis membranâ cinctis. ♀ ? circa Lima. (v. s. comm. à cl. Lagasca.)

IX. BREDEMEYERA. Willd. nov. act. nat. scrut. berol. 3. p. 411. t. 6.

Calyx 5-sepalus, sepalis 2 interioribus petaliformibus. Petala 3, medio carinæformi. Stam. 8 basi monadelphæ. Drupa ovata nuce biloculari. — Genus dubium.

1. *B. FLORIBUNDA* (Willd. l. c.) ♂ ad Caracas. Folia alterna oblongo-lanceolata integerrima glabra breviter petiolata. Bractæe subulatæ ad basin pedicello- rum. Flores flavi paniculati. †.

X. SECURIDACA. Lin. gen. n. 852. Lam. ill. t. 599. non Tourn.

Cal. 5-sepalus, sepalis 2 interioribus petaliformibus. Pet. 5 basi imâ subcoalita, 3 in carinam sæpè trilobam coalita, 2 oblonga. Stam. 8 diadelphæ. Capsula ovata indehiscens 1-locul. 1-sperma apice desinens in alam foliaceam lingulatam hinc crassiorem, illinc mem- branaceam. Semen ex apice loculi pendulum. Albumen nullum. — Suffrutices. Folia alterna ovata aut oblouga. Flores racemosi. Interdum appendix ex apice loculi ad latus alæ crassius adest, et alterius loculi abortivi signum fert.

* *Caule volubili aut scandente.*

1. *S. VIRGATA* (Sw! fl. ind. occ. 3. p. 1231.) glabra, caule scandente, ramis virgatis, foliis subrotundis obtusissimis, racemis terminalibus, capsulæ alâ rotundatâ. ♀ in frutetis Jamaïcæ, Hispaniolæ, Porto-Ricci. — Plum. ed. Burm. t. 248. f. 1. Folia decidua vix semuncialia. Flores fragrantés variegati (v. s.)
2. *S. VOLUBILIS* (Lin. sp. 992.) caule scandente, ramulis subpubescentibus, foliis ovali-oblongis acutis, racemis lateralibus, capsulæ alâ obovatâ dorso ad basin appendiculatâ. ♀ in Americâ meridionali prope Sanctam Mar- tham et Carthagenam. Jacq amer. 197. t. 183. f. 38. — Plum. ed. Burm. t. 247. f. 1. (v. s.)
3. *S. MOLLIS* (H. B. et Kunth nov. gen. am. 5. p. 421.) caule scandente, ramulis pubescentibus, foliis subrotundo-ovatis basi et apice obtusis suprâ puberu- lis, subtus molliter pubescentibus, racemis terminalibus. ♀ in ripâ flum.

Amazonum. Petala inferiora spathulata, superiora breviora. Legumen ferè S. volubilis.

4. *S. PUBESCENS*, caule scandente, foliis ovalibus ovatis obovatisve subtùs ramulisque subpubescentibus, racemis terminalibus. ζ in Cayennâ aut Guyanâ. An duæ species hic confusæ?

a. obovata, foliis ovatis obtusis. (v. s. sine fl. et fr. ex Mus. Par.)

β. ovata, foliis ovatis acutiusculis. (v. s. comm. à cl. Perrotet.)

5. *S. HEBEGLADA*, caule subsicante, ramis nervisque foliorum subtùs pedicellisque velutino-hirtis, foliis ovalibus mucronatis supernè lucidis, racemis terminalibus subfoliosis. ζ in Brasiiliâ. Flores magnitudine *S. virgatæ*, glaberrimi. Fruct. ign. (v. s.)

6. *S. COMPLICATA* (H. B. et Kunth nov. gen. am. 5. p. 423.) caule scandente, foliis ovato-ellipticis obtusis basi rotundatis subtùs hirtellis complicatis, ramulis molliter hirtis. ζ ad ripam fluminis Orinoci. Glandulæ duæ ad basin petioli loco stipularum. Petala inferiora lineari-oblonga superiùs æquantia.

**** Caule erecto.**

7. *S. ERRECTA* (Lin. sp. 992.) caule erecto, foliis oblongis utrinquè acutis, ramulis pedicellisque subpubescentibus, racemis terminalibus simplicibus. ζ in saxosis Hispaniolæ et Sancti-Thomasii. Jacq. amer. p. 197. t. 183. f. 39. Sw. obs. 274. (v. s.)

8. *S. PANICULATA* (Lam. ill. t. 599. f. 2. Poir. dict. 7. p. 52.) caule erecto, foliis ovali-oblongis subacutis, ramis glabriusculis, racemis terminalibus subcompositis subterminalibus. ζ in Cayennâ. Fructùs alæ hinc dorso ad basin appendiculatæ. (v. s.)

XI. *KRAMERIA*. *Læfl. itin.* 195. *Ruiz. et Pav. prod. fl. per.* t. 3. *Juss. mem. mus.* 1. p. 390.

Sepala 4, rariùs 5, extùs sericea, intùs colorata. Petala 3, 2 orbiculata, tertium è 2-3 petalis unguiculatis basi coalitis constans. Stam. 3-4 basi submonadelpha, antheris duplici poro dehiscens. Fructus globosus indehiscens setoso-aculeatus 1-locularis 1-spermus. Embryo rectus in centro albuminis carnosus. — Frutices multicaules. Folia alterna. Flores supremis ramulorum foliis axillares subsolitarii sessiles. Genus aliquantisper affine *Polygalæis* discrepat numero et structurâ partium, et forsàn ab ordine removendum?

*** Foliis simplicibus.**

1. *K. IXINA* (Lin. sp. 177.) foliis ovato-lanceolatis mucronato-spinosis villosopubescentibus, pedicellis axillaribus in racemum dispositis bibracteatis. ζ in Cumanâ et Santo-Domingo. (v. s.)

2. *K. SECUNDIFLORA* (fl. mex. ic. ined.) foliis linearibus acutis villosis, pedicellis folio longioribus bibracteatis in racemum unilateralem dispositis. \mathcal{Z} in Mexico.

3. *K. PAUCIFLORA* (fl. mex. ic. ined.) foliis oblongo-linearibus villosis, pedicellis paucis folio longioribus medio bibracteatis, calycis lobo inferiore gibbo. \mathcal{Z} in Mexico.

4. *K. TRIANDRA* (Ruiz et Pav! fl. per. 1. t. 93.) foliis oblongis acutiusculis villosis-sericeis, pedicellis folio sublongioribus bibracteatis racemum brevem constituentibus. ζ in Peruvix declivibus. — Radix vulgò *Ratanhia* dicta; præstans stypticum. (v. s.)

5. *K. LINEARIS* (Ruiz et Pav. fl. per. 1. t. 94. f. a.) prostrata, foliis linearibus, villosis, pedicellis tomentosis bibracteatis. ♂ in Peruviae collibus. *K. pentapetala* Syst. fl. per. p. 35.
6. *K. GLABRA* (Spreng. neu. entd. 2. p. 157.) foliis oblongo-lanceolatis nitidis, fructibus glabriusculis. — in Brasiliâ.

** *Folius trifoliolatus.*

7. *K. CYTISOIDES* (Cav! ic. 4. t. 590.) foliis petiolatis 3-foliolatis, foliolis oblongis villosis, pedicellis petiolo duplò longioribus. ♂ in Novâ-Hispaniâ. (v. s.)

Ordo XIX. TREMANDREÆ. *R. Brown gen. rem. p. 12.*

Cal. sepala 4-5 inæqualia æstivatione valvata, imâ basi subcoalescentia, decidua. Petala totidem, æqualia, sepalis alterna, æstivatione involuta, stamina includentia, calyce multò majora, decidua. Stam. hypogyna, distincta, 2 antè petalum quodque nempè 8-10, filamentis erectis, antheris basi insertis 2-4-locularibus poro tubulove apice dehiscentibus. Ovarium ovatum compressum biloculare, loculis ovula 1-3 pendula foventibus. Stylus 1. Stigmata 1-2. Capsula ovata, compressa, bilocularis, bivalvis, valvis medio septiferis. Semina apice septi inserta, pendula, ovata, umbilico nudo, appendice carunculæformi terminata. Embryo cylindræus rectus in axi albuminis carnosi, huic longior, radiculâ umbilicum spectante nempè superâ.

Fruticuli Ericoidei graciles sæpiùs erecti, pilis sæpè glanduloso-capitatis instructi, omnes ex Australasiâ orti. Folia alterna aut verticillata, exstipulata, integra aut dentata. Pedicelli axillares, solitarii, 1-flori.

Ordo affinis Polygaleis et paululùm etiam Droseraceis, generibus nondum satis forsàn cognitis constans.

I. TETRATHECA. *Smith nov.-holl. 1. t. 2. Labill. nov. holl. specim. 1. p. 95. t. 122. 123.*

Cal. 4-sepalus subæqualis. Petala 4. Stam. 8. antheris 4-locularibus. Semina subsolitaria.

1. T. JUNCEA (Smith. loc. cit.) foliis alternis paucis lanceolatis ramisque glabris. \mathcal{L} 5 in Novâ-Holl. Variat flore albo et sæpiùs purpureo. Caules ancipites subnudi. (v. s.)
2. T. GLANDULOSA (Labill! nov. holl. 1. p. 95. t. 123. non Sm.) foliis alternis ovato-oblongis dentatis ramisque piloso glandulosis, calycibus ovato-lanceolatis. \mathcal{L} 5 in terrâ Van-Diemen. Sequenti valdè affinis. Flores majores. (v. s.)
3. T. PILOSA (Labill! nov. holl. 1. p. 95. t. 122.) foliis sparsis subverticillatisve oblongis linearibusve integris margine revolutis ramisque piloso-glandulosis, calycibus suborbiculatis. \mathcal{L} 5 in terrâ Van-Diemen. T. glandulosa. Smith exot. bot. t. 21. Flores axillares solitarii. Pedunculi basi 1-bracteati. Sepala suborbiculata acuta ciliato-glandulosa. Petala obovata. (v. s.)
4. T. ERICIFOLIA (Smith. exot. bot. p. 37. t. 20.) foliis verticillatis linearibus margine revolutis scabris, ramis glabris, calycibus ovatis acutiusculis. \mathcal{L} 5 in Novâ-Hollandiâ. Rudg. trans soc. lin. 8. t. 11. Pedunculi et Calyces læves. Flores subnutantes. Petala obovato-rosea, aut rarius alba ex Sm. (v. s.)
5. T. THYMIFOLIA (Sm. exot. bot. t. 22.) foliis verticillatis lanceolatis acutis ciliatis, ramis pilosiusculis, calycibus lanceolatis acutis ciliatis. \mathcal{L} 5 in Novâ-Hollandiâ. Pedunculi calycisque scabri. Sepala acutiora quàm in priore, cui affinis.

II. TREMANDRA. *Brown. ined.*

Calyx 5-sepalus. Petala 5. Stam. 10, antheris 2-ocularibus.

1. *T. STELLIGERA* (Brown! ined.) erecta, foliis ovatis integris subtus canis supernè piloso-scabris, pilis stellatis. ♀ in Novâ-Hollandiâ orientali. Habitus ferè Hermannia aut Cisti alyssoidis. (v. s.)
 2. *T. DIFFUSA* (Brown! ined.) ramosissima diffusa, foliis ovatis grossè paucidentatis supernè glabris subtus pilis stellatis sparsis scabris. ♀ in Novâ-Hollandiâ. (v. s.)
-

Ordo XX. PITTOSPOREÆ. *Brown gen. rem. ter. austr. p. 10.*

Calyx 5-sepalus deciduus, sepalis nunc liberis nunc ad medium coalitis, æstivatione imbricatâ. Petala 5 hypogyna, unguibus conniventibus interdum cohærentibus, laminis patulis æstivatione imbricatis. Stam. 5 hypogyna, distincta, cum petalis alternantia. Ovarium 1 liberum, loculis placentisve 2-5 polyspermis. Stylus 1. Stigmata numero placentarum. Pericarpium capsulare vel baccatum, loculis polyspermis quandoque incompletis. Semina sæpè pulpâ glutinosâ obducta. Embryo minutus, prope umbilicum, in albumine carnoso inclusus, dicotyledoneus, radicula longiusculâ, cotyledonibus brevissimis.

Frutices vel Arbusculæ Africanæ, Asiaticæ aut pleræque Australasicæ. Folia simplicia, alterna, exstipulata, penninervia, sæpius integra. Flores terminales vel axillares, interdum polygami.

I. BILLARDIERA. *Smith exot. bot. t. 1. non Vahl. nec Mærch.*

Calyx 5-sepalus, sepalis acuminatis. Petala 5 unguibus margine subconvolutis approximatis. Bacca ellipsoïdea, stylo terminata. — Suffrutices sæpius scandentes, australasici, pedicellis axillaribus 1-2-floris, fructibus edulibus.

* *Foliis integris.*

1. *B. SCANDENS* (Smith. exot. bot. t. 1.) ramis scandentibus junioribus villosis, foliis lineari-oblongis integris, pedicellis 1-floris villosis flore brevioribus, baccis velutinis. ζ in Novæ-Hollandiæ orâ orientali. Rami adulti glabri. Folia sesquipollice longa, 2 lin. lata. Flores ochroleuci. *B. Canariensis* Wendl. her. 3. t. 15. (v. v.)
2. *B. MUTABILIS* (Sal! par. lond. t. 48.) ramis scandentibus, junioribus subvillosis, foliis lanceolato-linearibus integris, pedicellis 1 floris glabris floribus longitudine, baccis glabris. ζ in Novâ-Hollandiâ. Flores demum purpurascens. Sims bot. mag. t. 1313. (v. v.)
3. *B. FUSIFORMIS* (Labill! nov. holl. t. 90.) ramis vix scandentibus junioribus subvillosis, foliis ovali-oblongis integris, paniculis paucifloris, baccis fusiformibus villosis. ζ in insulâ Van-Diemen. (v. s.)
4. *B. LONGIFLORA* (Labill! nov. holl. t. 89.) ramis scandentibus junioribus vix pubescentibus, foliis oblongis linearibusve integris, pedicellis 1-floris glabris flore dimidiò brevioribus, baccis subglobosis torosis glabris. ζ in insulâ Van-Diemen Sims. bot. mag. t. 1507. (v. s.)
5. *B. ANGUSTIFOLIA*, ramis scandentibus, junioribus pubescentibus, foliis linearibus integris planis glabris, pedicellis 1-floris baccisque oblongis glabris. ζ in Novâ-Hollandiâ. Folia acuta. Pedicelli glabri. Flores ochroleuci. Petala acuta. (v. s. comm. à cl. Lambert.)
6. *B. ROSMARINIFOLIA*, ramis scandentibus glabris, foliis linearibus margine

revolutis nervoso-subreticulatis glabris. ζ in Novæ-Hollandiæ orâ orientali. (v. s. sine fl. et fr. et ideo dubia.)

**** *Foliis variis, aliis integris, aliis dentatis incisive.***

7. *B. PARVIFLORA*, ramis scandentibus, junioribus subvillosis, foliis oblongis glabris integris aut apice paucidentatis, pedicellis 1-2-floris bracteolatis hispidulis. ζ in Novæ-Hollandiæ orâ orientali. Lechenault. Sepala villosa hispida. Flores 4-5 lin. longi, ex sicco, subærulescentes. Petala acuta. (v. s.)
8. *B. VARIFOLIA*, ramis filiformibus villosis, foliis oblongis subvillosis integris aut dentato-subpinnatifidis, pedicellis villosis vix calyce longioribus. ζ in Novâ-Hollandiâ ad portum Regis Georgii. Lechenault. Sepala villosa. Flores 4 lin. longi, ad apices ramorum subcorymbosi. (v. s.)

II. PITTOSPORUM. *Ranks in Gaertn. fr. 1. p. 286. t. 59.*

Calyx 5-sepalus. Petala 5 unguibus in tubum conniventibus. Capsula 2-3-valvis, 1-locularis, valvis medio septiferis. Semina pulpâ resinosa obducta. — Frutices foliis integris persistentibus.

1. *P. CORIACEUM* (Ait. Kew. ed. 1. v. 3. p. 488.) foliis obovatis obtusis coriaceis glaberrimis, pedunculis umbellatim ramosis multifloris calycibusque villosis. ζ in Maderâ. Andr. bot. rep. t. 151. Lodd. bot. cab. t. 569. Vahl symb. 2. p. 43. (v. s.)
2. *P. VIRIDIFLORUM* (Sims bot. mag. t. 1684.) foliis obovatis retusis basi cuneatis nitidis subtus reticulatis, paniculâ subglobosâ terminali glabrâ. ζ ad Cap. B.-Spei. Burchell. P. Capense et *P. arbutifolium* Hortul. P. Sinense Desf. cat. 231. Capsulæ bivalves. ad Senaciam Napaulensem formâ valdè accedentes. Flores flavo-virentes, Jasminum spirantes. (v. s. sine fl. ex. Burch.)
3. *P. TOBIRA* (Ait! Kew. ed. 2. v. 2. p. 27.) foliis obovatis obtusis coriaceis glaberrimis, pedunculis 1-floris adgregato-umbellatis pubescentibus ζ in Japoniâ. Sims bot. mag. t. 1396. Tobira Kœmpf. amœn. t. 797. Evonymus Tobira Thunb. jap. 99. P. Chinense Donn. cent 48. Capsulæ sæpius 3-valves, nunc 2-5-valves, valvis crasso-coriaceis, ferè lignosis. (v. v.)
4. *P. UNDULATUM* (Andr. bot. rep. t. 393.) foliis ovali-lanceolatis utrinquè attenuatis undulatis petiolisque glabris, pedunculis terminalibus aggregatis pubescentibus ramoso-multifloris. ζ in Novâ Hollandiâ; nec in Canariis. Vent. h. Cels. t. 76. Delaun. herb. amat. t. 36. Ker. bot. reg. t. 16. Schrad. gen. em. t. 4. (v. v.)
5. *P. REVOLUTUM* (Ait. Kew. ed. 2. v. 2. p. 27.) foliis elliptico-oblongis obtusiusculis subtus pubescentibus margine revolutis, pedunculis terminalibus adgregato-umbellatis villosis. ζ in Novâ-Hollandiâ ad Port-Jakson. Ker. bot. reg. t. 186. bot. cab. t. 506. (v. s.)
6. *P. TOMENTOSUM* (Bonpl. nav. t. 21.) foliis obovali-oblongis utrinquè acutiusculis glabris, subtus ramisque pubescenti tomentosis, margine subrecurvis, pedunculis adgregatis terminalibus. ζ in Novâ-Hollandiâ. (v. s.)
7. *P. FULVUM* (Rudg. tr. soc. lin. 10. p. 298. t. 20.) foliis latè lanceolatis obtusis in petiolis et per nervos tomentosis, ramis tomentosis, pedunculis adgregatis, calycibus patulis. ζ in Novâ-Hollandiâ. An à priori satis distinctum? (v. s. ex h. Mus. Par?)
8. *P. FERRUGINEUM* (Ait. Kew. ed. 2. v. 2. p. 27.) foliis ellipticis utrinquè acuminatis superne glabris subtus in nervis et petiolo ferrugineo-tomentosis,

pedunculis terminalibus adgregato-umbellatis ramosis. 5 in Guinæa. Sims bot. mag. t. 2075.

ℓ? *flarium*, cortex flarius. Rumph. amb. 7. p. 13. t. 7. in insul. Moluccanis. An omnino eadem conf. Sm. in Rees. cycl.

9. *P. HIRTUM* (Willd. enum. 261.) foliis ovato-oblongis subtus pubescentibus, adultis calvescentibus, petiolis ramulisque hirtis. 5 in Canariis. *P. hirsutum* Link. enum. 233.
10. *P. LAURIFOLIUM* (Willd. rel. in Rœm. et Schult. syst. 5. p. 432.) foliis oblongis obtusis coriaceis, pedunculis 1-floris lateralibus adgregatis. 5 in insulâ Teneriffæ.
11. P? *PHYLLIRHOÏDAS*, foliis oblongis mucronatis coriaceis planis glaberrimis subaveniis, pedicellis lateralibus 1-floris solitariis. 5 in Novâ-Hollandiâ. Lechenault. Capsula ovata subcompressa bivalvis extus lævis subcarnosa. (v. s. sine fl.)

† *Species vix notæ.*

12. *P. TENUIFOLIUM* (Gaertn. fr. 1. p. 286. t. 59.) capsulis trivalvibus subglobosis rugosis. 5 in Australasiâ.
13. *P. UMBELLATUM* (Gaertn. l. c.) capsulis bivalvibus compresso-globosis extus punctis elevatis scabris. 5 in Australasiâ.
14. *P. EXPENSE*. Nois. hort. ex Steud. nom. 628.
15. *P. TUBERCULATUM* Zeyh. ex Steud. nom. 628.

III. BURSARIA. Cav. ic. 4. t. 350.

Calyx 5-dentatus. Petala 5 à basi distincta. Capsula compressa, substipitata, obcordata, bilocularis, bivalvis (ferè Polygalæ). Semina (ex Cav.) resinâ induta. — Frutex ramosissimus, foliis oblongo-cuneatis nervo medio donatis cœterum aveniis integerrimis, floribus albis paniculatis.

1. *B. SPINOSA* (Cav. l. c.) 5 in Novâ-Hollandiâ. Variat ramis spinosis et inermibus. *Itea spinosa* Andr. bot. rep. t. 314. *Cyrrilla spinosa* Spreng. nov. prov. 15? (v. v.)
- IV. *SENACIA*. Comm. ined. *Du Pet. th. obs. pl. isl. d. fr. p. 27.* — *Senaciæ* sp. Lam. ill. 2. p. p. 95.

Calyx 5-dentatus minimus. Petala 5 à basi distincta lanceolata. Stam. 5 hypogyna. Capsula junior bacciformis demum bivalvis semibilocularis. Semina 4-8 ad basin aut medio septorum adnata, arillata, albumine corneo. Embryo minimus ad hujus basin. — Frutices ramosi glabri, foliis integris penninerviis, floribus corymbosis, habitu Celastrorum quibuscum diu confusi.

1. *S. UNDULATA* (Lam. ill. n. 2709.) foliis lanceolatis undulatis, floribus terminalibus umbellato-corymbosis, fructibus breviter pedicellatis 4-spermis. 5 in insulis Mauritanis ubi gallicè *Bois de joli cœur* dicitur. *Celastrus undulatus* Lam. dict. 1. p. 662 Ovarium substipitatum. Stylus brevis. (v. s.)
ℓ. Foliis ovali-lanceolatis. (v. s.)
2. *S. NAPAULENSIS*, foliis lanceolatis planiusculis, floribus terminalibus umbellato-corymbosis, fructibus sessilibus 4-spermis. 5 in Napauliâ. *Celastrus verticillatus* Roxb! cat. calc. 18. non fl. per. Folia non verticillata sed circa

florum corymbos conferta. Capsulæ subcompressæ longitudine suâ latiores, valvis post dehiscentiam apice mucronatis. (v. s.)

3. *S. LANCEOLATA* (Lam. ill. n. 2710.) foliis lanceolatis planiusculis, floribus umbellato-corymbosis, seminibus 8, 4 in fundo, 4 in medio loculorum. 5 in insulis Mauritianis, conf. Pet. th. loc. cit.
 4. *S. ELLIPTICA* (Lam. ill. n. 2711.) foliis ellipticis obtusis, floribus axillaribus subfasciculatis. 5 in insulis Antillis. Similis *Rhamno sarcomphalo*. Species valdè dubia, forsân potiùs ad *Celastrum* aut *Maytenum* referenda.
-

Ordo XXI. FRANKENIACEÆ. *St.-Hil. mem. plac. cent. p. 39.*

Cal. Sepala 4-5 erecta basi coalita in tubum sulcatum, persistentia, æqualia, linearia, acuta, concavo-canaliculata. Petala tot quot sepala, iis alterna, hypogyna, unguiculata, ungue longitudine calycis, limbo patente, fauce squamulis subcoronata. Stamina hypogyna, nunc petalis æqualia et iis alterna, nunc 1-2 addita petalis opposita et ideò stamina verosimiliter habenda ut petalis numero dupla. Filamenta filiformia. Antheræ subrotundæ. Ovarium 1 liberum. Stylus filiformis 2-3-fidus. Capsula calyce persistente stipata, ovato-oblonga, subtrigona, bi aut sæpius tri-imò quadrivalvis, 1-ocularis; valvulis ad marginem utrumque placentiferis, polyspermis. Semina minuta. Embryo rectus in medio albuminis (ex Gaertn. fil.) vaginalis in laminas duas divisi, radiculâ brevi ad umbilicum spectante, cotyledonibus planis ellipticis.

Herbæ aut suffrutices. Caules teretes ramosissimi. Folia opposita aut verticillata, exstipulata, basi producta in membranulam amplexicaulem, sæpè glandulosa, oblonga, integra, marginibus revoluta. Flores ad apices et in dichotomiis ramorum sessiles foliis floralibus stipati, sæpius roseo-violacei.

Ordo distinctus à Caryophylleis seminibus ad margines capsulæ adnatis, à Lineis fructu uniloculari et plurimis notis aliis, à Gentianeis floribus polypetalis et seminis structurâ paulò diversâ, à Violaceis seminibus margini nec centro valvularum adnatis.

I. FRANKENIA. *Lin. gen. n. 445.*

Stylus 3-fidus, lobis oblongis intus stigmatosis. Capsula 3-4-valvis polysperma.

1. *F. PULVERULENTA* (Lin. sp. 474.) foliis obovatis retusis glabris subtus pulverulentis in petiolo ciliatis, radice exili. ① in arenosis maritimis Europæ, Sibiriae, Tauriæ, forsan Novæ-Hollandiæ. — Clns. hist. 2. p. 186. f. 2. (v. v.)
2. *F. NODIFLORA* (Lam. ill. t. 262. f. 4.) foliis ovatis glabris in petiolo non ciliatis, caulibus prostratis. calycibusque glabris. 5 2 ad Cap. B.-Spei. F. Burch! cat. n. 513. (v. s.)
3. *F. NOTHRIA* (Thunb. prod. 58. fl. cap. ed. Schult. 1. p. 295.) foliis linearibus margine revolutis glabris basi ciliatis, ramulis pubescentibus, caulibus prostratis calycibusque glabris. 5 2 ad Cap. B.-Spei. Nothria repens Berg. cap. 171. t. 1. f. 2. An à sequente diversa?
4. *F. LEVIS* (Lin. sp. 473.) foliis linearibus margine revolutis glabris basi ciliatis, caulibus prostratis calycibusque glabris. 5 2 in maritimis Europæ, ins. Canariensium. — Mich. gen. t. 22. f. 1. Flos roseo-violaceus interdum albus. (v. v.)
5. *F. INTERMEDIA*, foliis linearibus margine revolutis glabris basi ciliatis, caulibus prostratis velutinis, calycibus hispidis. 5 2 in maritimis Europæ australis secus Mediterraneum et in littore Algeriensi. *F. hirsuta* var. *Calabrica* Lin. *F. hirsuta* fl. fr. 4. p. 766. — Mich. gen. t. 22. f. 2. (v. v.)
6. *F. HISPIDA*, foliis linearibus margine revolutis glabris basi subciliatis, caulibus diffusis calycibusque hispidis. 5 2 in salsuginosis et rupestribus à Cypro ad Sibiriam. *F. hirsuta* var. *Cretica*. Lin. *F. hirsuta* Sibth! fl. gr. t. 313. (v. s.)

7. *F. CORYMBOSA* (Desf. fl. atl. 1. p. 315. t. 93.) foliis linearibus margine revolutis glabris basi subciliatis, caulibus velutinis erectis, calycibus glabris, floribus corymbosis. ζ in maritimis Barbariæ prope Arzean. (v. s.)
8. *F. ERIGIFOLIA* (C. Smith. in Buch. can. p. 30.) foliis linearibus petiolatis margine revolutis basi subciliatis supernè glabris subtus caulibus calycibusque pruinoso-velutinis. ζ in maritimis insularum Canariensium. Species inquirenda nunc ad Frank. corymbosam, nunc ferè ad Frank. intermediam habita accedens. (v. s.)
9. *F. VELUTINA*, foliis lineari-oblongis obtusis basi non ciliatis utrinquè nec non in caulibus calycibusque molliter velutinis. ζ in agro Mogadoriensi. Broussonet. (v. s.)
10. *F. MOLLIS* (Bieb. suppl. 276.) foliis ovato-lanceolatis acutis villosis, ramis floriferis alternis paniculatis villosis, floribus pedicellatis. ζ in Tauriâ? Tota pilis mollibus patulis vestita subcanescit.
11. *F. THYMIFOLIA* (Desf. fl. atl. 1. p. 316.) foliis oblongis obtusis parvis rigidulis glabris cinereis basi ciliatis, caulibus erectis, junioribus vix puberulis, calycibus glabris. ζ in arenis deserti Barbariæ circa Tozzer, in Hispaniâ circa Aranjuez. (v. s.)
12. *T. MICROPHYLLA* (Cav. ic. 6. t. 597. f. 1.) foliis ovalibus minimis quadrifariam imbricatis glabris, caulibus prostratis calycibusque glabris. ζ in Americâ meridionali.
13. *F. FRUTIGULOSA*, foliis linearibus acutiusculis margine revolutis glabris basi non ciliatis, caulibus erectis glabris, calycibus acutis glabris ζ in arenis Novæ-Hollandiæ ad insulam Sancti-Francisci. (v. s. in h. Mus. Par.)
14. *F. PAUCIFLORA*, foliis linearibus margine revolutis ferè filiformibus glabris basi ciliatis, caulibus erectiusculis canescentibus calycibusque glabris acutis. ζ in Novæ-Hollandiæ orâ occidentali. (v. s. in h. Mus. Par.)
15. *F. TETRAPETALA* (Labill. nov. holl. 1. t. 114.) foliis linearibus margine revolutis ferè filiformibus subsolutis, floribus 4-petalis, calycis lobis 4 margine tomentosis. ζ in Novæ-Hollandiæ insulâ Van Diemen.
16. *F. REVOLUTA* (Forsk. æg. 75.) foliis ovatis margine revolutis, caulibus dichotomis, capsulis 4-valvibus. ζ in Alexandriæ desertis. — Flores 5-sepalos vidi in specim. hujus speciei à cl. Siebero Alexandriæ lecto. Habitus *F. corymbosa* sed flores sparsi solitarii. (v. s. in h. Balb.)

II. BEATSONIA. Roxb. fl. St.-Hel. in ~~h.~~ Beats. p. 300.

Stylus bifidus, lobis in stigmata globulosa desinentibus. Capsula bivalvis oligosperma. — An satis à Frankeniâ differt?

1. *B. PORTULACIFOLIA* (Beats. l. c.) foliis subrotundis carnosis glabris, caule fruticoso dumoso. ζ in rupibus maritimis ad littus australe insulæ Sanctæ-Helenæ. (v. s.)

III. LUXEMBURGIA. St.-Hil. mem. mus. 9. p. 351.

Cal. sepalâ 5 subinæqualia caduca. Pet. 5 inæqualia. Antheræ hypogynæ subsessiles lineares tetragonæ apice biporosæ in massulam concavam secundam coalitæ. Stylus subulatus incurvus. Ovarium 3-gonum. Capsula 3-valv. 1-loc. ∞ -sperma, marginibus introflexis seminiferis. Semina marginata. — Frutices ramosi glaberrimi. Folia alterna dentata cuspidata, nervis lateralibus parallelis numerosis, petiolo basi bi-stipulato, stipulis ciliatis caducis. Flores terminales, racemosi, lutei. — Genus videtur medium inter Violaceas et Frankeniacæ, prioribus flore irregulari, posterioribus capsulæ structurâ accedens.

1. *L. OCTANDRA* (St.-Hil. l. c.) foliis subsessilibus lineari-lanceolatis angustis, floribus octandris. ζ in Brasilæ prov. Minas-Novas.
2. *L. POLYANDRA* (St.-Hil. l. c.) foliis petiolatis lanceolato-ellipticis oblongis, floribus polyandris. ζ Hab. cum priore

Ordo XXII. CARYOPHYLLÆ. (1) *Juss. gen.* 299. — Caryophyllæ et Arenariæ. *Lam.*

Calyx sepalis 4 aut sæpius 5 cum pedicello continuis, nunc liberis, nunc inter se in tubum 4-5-dentatum coalitis, æstivatione imbricatis constans, sæpius persistens. Petala totidem hypogyna toro plùs minùsve elevato (Anthophoro dicto) (2) inserta, sepalis alterna unguiculata, limbo integro aut bifido, ad faucem interdum squamis petaloideis aucta, rarissimè abortu nulla. Stamina petalorum numero dupla toro inserta, alia petalis alterna libera præcociora, alia petalis opposita et cum iis basi coalita seriora interdum abortiva, filamenta subulata interdum basi submonadelpha; antheræ biloculares birimosæ, basi sæpius insertæ. Ovarium tori apice insertum simplex ovatum oblongumve 2-5-valve, stylis totidem superatum filiformibus clavatisve à basi ipsâ distinctissimis, intùs stigmatosis papillois. Capsula valvis 2-5 basi coalitis apice dehiscentibus et tunc dentiformibus, nunc integris nunc bifidis constans, nunc sæpius 1-locularis, nunc 2-5-locularis, valvis proferentibus è medio septa, nunc incompleta, nunc ad axim continua. Placenta semper centralis, intùs fila viridia seu flavida nutritientia fovens, in capsulis unilocularibus libera subconica cum stylorum basi continua ope filamentorum quorundam alborum exteriorum ex apice placentæ ad capsulæ apicem liberorum, post anthesin dessicatorum, in capsulis multilocularibus cum septis coalita filamentis auræ conductoribus etiam coalitis nec oculo nudo visibilibus. Semina ∞ (rarissimè pauca et definita) secùs placentam centram in tot series geminatas quot sunt fila conductoria et styli sæpius disposita, seriebus seminum cum stylis alternantibus. Albumen farinosum sæpius centrale. Embryo nunc periphericus plùs minùsve incurvus, rarissimè rectus centralis, radiculâ ad hilum versâ. DC. mss.

Herbæ aut suffrutices. Caules nodosi. Folia constantissimè opposita, sæpè connata, integra. Flores terminales.

Trib. I. *SILENÆÆ*. DC. mss. — Caryophyllæ *Lam.*

Sepala coalita in tubum cylindraceum apice 4-5-dentatum.

I. *GYPSOPHILA*. *Linn. gen. n.* 768.

Calyx campanulatus angulatus sub-5-lobus, laciniis margine membranaceis. Petala 5 non unguiculata. Stamina 10. Styli 2. Capsula 1-locularis.

(1) Ordinem elaboravit cl. *SERINGE*.

(2) Dicitur nullum ubi non manifestum.

Sect. I. STRUTHIUM. Ser. mss.

Calyces squamis destituti.

1. *G. OCELLATA* (Sibth. et Smith fl. græc. t. 387. ex prod. 1. p. 281.) floribus capitatis, petalorum laminis ovatis zonulâ notatis, caulibus diffusis, foliis spathulatis utrinquè tomentosis. \mathcal{L} in Delphi monte Eubæa. Cucubalus polygonoides Willd. spec. 2. p. 690?
2. *G. THYMIFOLIA* (Sibth. et Smith fl. græc. t. 388. ex prod. 1. p. 282.) floribus subcapitatis, caulibus diffusis subdichotomis, foliis spathulatis utrinquè tomentosis, petalorum laminis obovatis rotundatis immaculatis. \mathcal{L} in Monte Parnasso. Habitus *G. ocellatæ*. Caules magis elongati magis ramosi. Calyx pilosus.
3. *G. STRUTHIUM* (Lin. spec. 582.) floribus confertis, caulibus simplicibus scabriusculis, foliis linearibus carnosissimis. \mathcal{L} in Hispaniâ. — Barr. ic. 64. t. 119. Caulis infernè fruticosus.
4. *G. FASTIGIATA* (Lin. amœn. 3. p. 23.) floribus corymboso-fastigiatis, caule ascendente, foliis lanceolato-linearibus obsolete triquetris lævibus obtusis secundis, staminibus exsertis. \mathcal{L} in Galliæ, Helvetiæ, Hærcyniæ rupibus. — Gmel. fl. sib. 4. p. 144. t. 61. f. 1. (v. s.)
5. *G. DICHOTOMA* (Besser prim. fl. galiz. 1. p. 372.) floribus corymboso-fastigiatis, caule erecto dichotomo compresso glabro, foliis linearibus subcarnosis triquetris acutis, petalis oblongis patentibus, stylis stamina æquantibus. \mathcal{L} in saxosis circa Jaslo.
6. *G. COLLINA* (Stevl in litt.) caule ramoso lævi, floribus fastigiatis, foliis linearibus subcarnosis acutis scabriusculis, lobis calycinis obtusiusculis, staminibus exsertis. \mathcal{L} ? circa Odessam. à sequente differt paniculæ glabritie. Aff. *G. fastigiata*. (v. s.)
7. *G. ARENARIA* (Waldst. et Kit. pl. hung. 40. t. 41.) floribus corymboso-fastigiatis, petalis rarissimè submarginatis, foliis linearibus subcarnosis glabris planis. \mathcal{L} in arenosis Hungariæ, Volhyniæ. Capsula globosa. (v. s.)
8. *G. VISCOSA* (Murr. in comm. gott. 1783. p. 9. t. 3.) floribus corymboso-fastigiatis, ramis divaricatis, foliis ovato lanceolatis lævibus basi cordatis amplexicaulibus, internodiis ramorum medio viscosis, petalis retusis, laciniis calycinis latius obtusis. \textcircled{I} in Oriente. (v. s.)
9. *G. TENELLA* (Poir. suppl. 2. p. 874.) floribus paniculato-umbellatis longè pedicellatis, caule tenerissimo. \textcircled{I} Hab. Bractæ breves ovato-lanceolatæ. Corolla alba.
10. *G. GRANDIFLORA* (Poir. suppl. 2. p. 874.) caule stricto, floribus paniculatis, ramulis subnudis paucifloris, petalis emarginatis, foliis linearibus subulatis hirsutis. — Hab. Panicula pauciflora. Flores maximi.
11. *G. ALTISSIMA* (Lin. spec. 582.) ramis patulis, floribus paniculatis minimis, paniculâ ramosissimâ, pediculis viscosis, caulibus rectis, foliis lanceolatis subtrineriis. \mathcal{L} in Sibirîâ. — Gmel. fl. sib. 4. p. 143. t. 60. Capsula minima subrotunda. Lacinie calycinæ obtusæ. (v. s. ex h. Patr. et comm. à cl. Steven.)
12. *G. PERFOLIATA* (Linn. spec. 583.) floribus paniculatis, paniculâ ramosissimâ lævigatâ, foliis ovato-lanceolatis semi-amplexicaulibus. \mathcal{L} in Hispaniâ et Oriente. — Dill. helth. 368. t. 276. (v. s.)
 β . tomentosa, foliis tomentosis. *G. tomentosa* Linn. spec. 582? — In Tauriâ. — Barr. ic. t. 1002. (v. s. comm. à cl. Beauprè.)
13. *G. SCORZONERIFOLIA* (hort. Mus. Par.) floribus paniculatis, paniculâ viscosâ, foliis lanceolatis semi-amplexicaulibus acutis 3-5-nerviis lævibus \mathcal{L} prope Kislar. *G. sabulosa* Stevl in litt. *G. perfoliata* Bieb. fl. taur. ex Steven! Differt à *G. perfoliata* paniculâ viscosâ, floribus majoribus et calycibus acutioribus. Steven. (v. s.)

14. *G. ACUTIFOLIA* (Fisch. cat. hort. gor. 1812. p. 59.) floribus trichotomo-paniculatis, pedunculis villosis-viscosis, petalis emarginatis calyce duplò longioribus genitalia superantibus, foliis lineari-lanceolatis acuminatis planis obscure 3-nerviis. \mathcal{L} in collibus lapidosis Cancasi. *G. altissima* Bieb. fl. taur. 1. p. 280. non Linn. (v. s. comm. à cl. Fisch. et Stev.)
15. *G. PANICULATA* (Linn. amœn. 3. p. 23.) floribus paniculatis minutissimis dioicis, pedunculis lævibus filiformibus divaricatis, laciniis calycinis obtusis, foliis lineari-lanceolatis scabris acutis. \mathcal{L} in sabulosis Siciliæ, Sibiria. Jacq. fl. austr. 5. t. 1. Aff. *G. altissimæ*. (v. s.)
16. *G. ASCENDENS* (Jacq. hort. vind. 2. t. 138.) floribus paniculatis parvis, pedunculis lævibus filiformibus divaricatis, laciniis calycinis obtusis corollâ staminibus pistillisque ejusdem longitudinis, caulibus prostratis, foliis lanceolato-linearibus. \mathcal{L} in Oriente. (v. s.)
17. *G. GLAUCA* (Stev! in litt.) floribus paniculatis, paniculâ divaricatâ, ramis paucifloris pubescentibus viscosis, foliis lineari-lanceolatis subcarnosis obtusis. \mathcal{L} ? in Caucaso. *G. repens* Bieb. fl. taur. 318. ex Stev. in litt. (v. s.)
18. *G. ELEGANS* (Bieb. fl. taur. 1. p. 319. suppl. 296.) floribus dichotomo-paniculatis glabris, petalis emarginatis calyce genitalibusque duplò longioribus, foliis lanceolatis subcarnosis. ① in glareosis Tauriæ. Schrank pl. mon. t. 21. Aff. *G. viscosæ*, sed foliis utrinquè acutis angustioribus. (v. s. comm. à cl. Stev. et Fisch.)
19. *G. OCHROLEUCA* (Sibth. et Smith fl. græc. t. 385. ex prod. 1. p. 281.) caule dichotomo divaricato, petalorum laminis spathulatis basi punctatis. foliis subulatis infimis linearibus flaccidis. ①? in monte Hymetto prope Athenas. Flores similes *G. Creticæ*, petala magis spathulata, pallida straminei coloris eleganter striata.
20. *G. STEVENI* (Fisch. in litt. Schrank pl. h. mon. t. 32.*) floribus paniculatis, caule diffuso, foliis lineari-lanceolatis gramineis carinatis cæsiis, petalis dilatato-linearibus obtusis integris. \mathcal{L} in Iberiæ lapidosis circa Tiflin. *G. repens* Bieb. fl. taur. 1. p. 318. excl. syn. (ex Fisch. in litt.)
21. *G. GRAMINEA* (Sibth. et Smith prod. 1. p. 279.) floribus paniculatis, caule nudiusculo, foliis radicalibus linearibus cespitosiis, petalis emarginatis, calyce pubescente. \mathcal{L} in Peloponesi montibus. Folia radicalia numerosa graminea, læte viridia, margine spinuloso-scabra.
22. *G. ARMERIOIDES* (Ser. mss.) caulibus cespitosiis rigidis, floribus dichotomo-paniculatis, pedunculis calycibusque glanduloso-pilosis, calyce striato, laciniis acutiusculis margine scariosis, petalis spathulatis integriusculis, foliis linearibus obtusiusculis firmis internodiorum longitudine. \mathcal{L} in Troadâ. Habitus Dianthi Armeriæ. (v. s. ex herb. Olivier.)
23. *G. CRETICA* (Sibth. et Smith fl. græc. t. 384. ex prod. 1. p. 280.) caule subdichotomo, petalorum laminis lanceolatis immaculatis, foliis subulatis 3-nerviis. \mathcal{L} in Cretæ rupestribus. *Saponaria Cretica* Lin. spec. 584. Radix cespitosa.
24. *G. TENUIFOLIA* (Bieb. fl. taur. 1. p. 319. suppl. 296.) caulibus subpaniculatis paucifloris, petalis emarginatis calyce genitalibusque duplò longioribus, foliis subulato-filiformibus glabris. \mathcal{L} circa Portas Caucasicas. Radix tortuosa crassissima foliaque similia Plantagini subulatæ. Petala rosea inflexo-emarginata. (v. s.)
25. *G. REPENS* (Lind. amœn. 3. p. 23. spec. 581.) caulibus subpaniculatis paucifloris, staminibus petalis emarginatis brevioribus, foliis linearibus glabris. \mathcal{L} in petrosis ad vias et secus torrentes Alpium et Pyrenæorum. Lam. ill. t. 375. f. 2. Jacq. fl. austr. 5. p. 4. t. 407. (v. v.)
26. *G. PATRINI* (Ser. mss.) caulibus subpaniculatis paucifloris, calycibus obtusiusculis, petalis stamina æquantibus, foliis angustissimis crassis internodiis

- triplò longioribus. \mathcal{Z} in campestribus ad Irtim circa Oustka-Menogorik. (v. s. ex h. Patr. comm. à cl. Deless.)
27. *G. DUBIA* (Willd. enum. suppl. p. 23.) petalis obovatis emarginatis campanulatis, staminibus corollà brevioribus, foliis linearibus subcarnosis. \mathcal{Z} Hab. An *G. repens* Lin? ex cl. Trevirano.
28. *G. PROSTRATA* (Lin. spec. 1. p. 581.) caulibus subpaniculatis, pistillis petalis emarginatis longioribus, foliis lineari-lanceolatis levibus. \mathcal{Z} in Europâ? Sims bot. mag. t. 1281. — Pluk. alm. 22. t. 75. f. 2. Flores rosei. Aff. *G. repenti*. (v. s.)
29. *G. ROKEJEKA* (Delil. fl. egypt. 87. t. 29. f. 1.) caule erecto, ramis exilibus paniculatis dichotomis, pedunculis capillaribus, corollà majusculà lineatopictâ, foliis radicalibus ovatis oblongis, superioribus linearibus. (I) \mathcal{Z} in AEGYPTO prope Soueys. Rokejeka Forsk. descr. 90. n. 77. Capsula globosa, Semen hemisphaericum nigrum rugosum. (v. s. in h. Ser. comm. à cl. Sieb.)
30. *G. THESIFOLIA* (Ser. mss.) caulibus numerosis subpaniculatis paucifloris strictis, petalis subintegris stamina æquantibus, foliis linearibus angustissimis glabris. \mathcal{Z} circa fodinam Schamanaikhæ. Patr. (v. s. comm. à cl. Deless.)
31. *G. MURALIS* (Lin. amœn. 3. p. 24. spec. 583.) caule dichotomo-paniculato ramosissimo, floribus axillaribus solitariis, petalis crenatis, foliis linearibus planis longitudine pedicellorum. (I) in Galliâ, Helvetiâ, Germaniâ, Sueciâ, Tauriâ, ad vias et inter segetes. Lam. ill. 375. *G. serotina* Hayne. (v. v.)
32. *G. COMPRESSA* (Desf. fl. atl. 1. p. 343. t. 97.) caule erecto paniculato hinc compresso, pedicellis pubescentibus, floribus terminalibus, calycibus elongatis striatis lobis acutis, foliis subulato-lanceolatis striatis. (O) in Barbariæ arvis arenosis. Petala subtus venis violaceis variegata. Similis *G. saxifragæ*, at caret squamis calycinis. (v. s.)

Sect. II. PETRORHAGIA. Ser. mss.

Calyces squamis scariosis 2-4 oppositis basi instructi.

33. *G. GLOMERATA* (Pall. ind. taur. ex Bieb. fl. taur. 1. p. 321. suppl. 297.) floribus capitato-glomeratis, caulibus basi ramosis herbaceis patulis, foliis linearibus carinato-triquetris scabriusculis, staminibus petala obovata retroflexa subæquantibus. \mathcal{Z} in Tauriæ et Caucasi lapidosis. (v. s.)
- β . *capitata* (Bieb. fl. taur. 1. p. 321.) foliis longioribus minùs carnosius subglaucescentibus, caulibus suffruticosis.
34. *G. SAXIFRAGA* (Lin. spec. 584.) caulibus numerosis erectis rigidis, floribus paniculatis terminalibus, calycibus squamis quaternis involucreatis, foliis linearibus rigidis. \mathcal{Z} in saxosis Europæ austr. *Dianthus saxifraga* Lin. spec. ed. 1. p. 413. — Barr. ic. t. 998. Flores rosei. Dentes calycis obtusissimi. (v. v.)
- β . *capillacea* (Ser. mss.) foliis caulibusque capillaceis elongatis.
- γ . *hispidula* (Ser. mss.) foliis caulibusque hispidulis. — In Vallesiâ. (v. s. in h. Ser.)
- δ . *rigida* (DC! fl. fr. suppl. p. 600.) floribus agglomeratis. *G. rigida* Lin. amœn. 3. p. 24. spec. 583. (excl. syn. Sauv. et Dalech. ad var. α refer.) in Galliâ austr.
- 35? *G. DIANTHOIDES* (Sibth. et Smith fl. græc. t. 383. ex prod. 1. p. 280.) foliis linearibus, floribus capitatis, bracteis confertis dilatatis scariosis, petalis obtusis integerrimis. \mathcal{Z} in Cretâ. *Lychnis orient.* petalis albis, lineis cæruleis subtus variegatis Tourn. cor. 24. Dentes calycis obtusiusculi. (v. s. in herb. Balb.)
36. *G. MULTICAULIS* (Poir. suppl. 2. p. 875.) caulibus numerosis strictis simplicibus, foliis adpressis basi vaginantibus subulatis nervosis, flore terminali solitario sessili. \mathcal{Z} in Helvetiâ. An var. *G. saxifragæ*?

II. BANFFYA. *Baumg. fl. trans. ex Spreng. neue entd.* 1. p. 300.

Calyx tubulosus profundè 5-partitus persistens. Petala 5 indivisa. Stamina 5 fertilia, 5 abortientia. Pistilla 2. Capsula 1-locularis oligosperma. — Vix à Gypsophila differt, ni calicè et numero seminum.

1. B. PETRÆA (Baumg. l. c.) foliis linearibus obtusis, caulibus cespitosis. 2/2? in Alpibus Dinariis.

III. DIANTHUS. *Lin. gen. n.* 770.

Calyx tubulosus 5-dentatus basi squamis 2 4 oppositis imbricatis. Petala 5 longè unguiculata. Stamina 10. Styli 2. Capsula 1-locularis. Semina compressa hinc convexa indè concava, peltata. Embryo vix curvatus.

Sect. I. ARMERIASTRUM. *Ser. mss.*

Flores capitati vel corymbosi, sessiles vel pedunculati.

§. I. *Bracteis ovatis muticis.*

1. D. PROLIFER (Lin. spec. 587.) floribus aggregatis capitatis, squamis calycinis ovatis muticis tubum superantibus, foliis serrulatis. ① ad marginem sylvarum et agrorum Europæ. — OEd. fl. dan. t. 221. Tunica prolifera Scop. fl. carn. n. 503. Flores pallidè rubri, semina ovata compressa. (v. v. et s.)

℘. *diminutus* (DC! fl. fr. 5. p. 741.) floribus subsolitariis. D. diminutus Lin. spec. 587.

2. D. SPINOSUS (Desf. in ann. mus. 1. p. 198. t. 16. f. 1.) fruticosus procumbens, floribus capitatis sessilibus, foliis rigidis subulatis spinulentibus. 3 in Persiâ. Petala linearia.

§. 2. *Bracteis lanceolatis acutis, calycibus striatis villosis.*

3. D. ARMERIA (Lin. spec. 586.) floribus laxè fasciculatis, squamis calycinis lanceolato-subulatis tubum æquantibus, foliis subulatis calycibusque hirsutis. ① in sterilibus Europæ. — OEd. fl. dan. t. 230. D. hirtus Lam. fl. fr. 2. p. 533, non Vill. (v. v.)

℘. Floribus solitariis. (v. s.)

4. D. PSEUDO-ARMERIA (Bieb. fl. taur. 1. p. 323, suppl. 297.) floribus densè aggregato-fasciculatis, squamis calycinis ovato-subulatis tubum æquantibus, petalis barbatis, foliis subulatis pubescenti-scabris strictis. 2/2 in Tauriæ siccis lapidosis. Sims bot. mag. t. 2288. D. barbatus Pall. ind. taur. Aff. D. Armeriæ, sed undiquè pube exili tantum canescens, nec villis insigniter hirsutus.

℘. Bracteis divaricatis, calycibus abbreviatis. — Monstruositas ex Uredine flori adnatâ.

5. D. PUBESCENS (Sibth et Smith. fl. græc. t. 397. ex prod. 1. p. 286, et D'Urv. ! enum. 47.) floribus aggregatis solitariisque, squamis calycinis ovato-subulatis tubo dimidio brevioribus, calycibus striatis villosis glutinosis, dentibus brevibus, foliis subulatis glabriusculis. ②? circa Tenedos. (v. s. comm. à cl. D'Urville.)

- 6.? D. ARMERIOIDES (Rafin. in Desv. journ. bot. 1814. v. 2. p. 269.) floribus aggregatis, squamis calyce longioribus striatis scabris, foliis linearibus internodiis brevioribus scabris, caule simplici superne aspero. ②? in pratis New-Jersey. An D. discolor Sims bot. mag. t. 1162?

§. 3. *Bracteis ovatis vel lanceolatis, calycibus vix striatis glabris.*

7. D. BARBATUS (Lin. spec. 586.) floribus aggregatis fasciculatis, squamis calycinis ovato-subulatis tubum æquantibus, foliis lanceolatis. 2/2 in sterilibus Galliæ austr. et Germaniæ (v. v.)

- β. *latifolius* (Ser. mss.) foliis ovato-lanceolatis, floribus aggregatis.
- γ. *paniculatus* (Ser. mss.) foliis ovato-lanceolatis, caule paniculato. (v. s. in h. DC. et Ser.)
- δ. *pedunculatus* (Ser. mss.) foliis lanceolatis, floribus pedunculatis laxis. (v. s. in h. Ser.)
8. D. *LATIFOLIUS* (Willd. enum. p. 466.) floribus aggregatis racemoso-corymbosis, squamis calycinis ovato-lanceolatis demum calycem superantibus, foliis oblongo-lanceolatis. 2ℓ habitu refert ferè D. barbatus, sed folia latiora, flores racemoso-corymbosi duplò majores. An var. γ D. barbati?
9. D. *SILENCIOSUS* (Poir. dict. 4. p. 514.) caule. . . , squamis calycinis ovato-lanceolatis, foliis subaggregatis subovatis glanduloso-punctatis ciliatis. — Hab.
10. D. *AGGREGATUS* (Poir. suppl. 4. p. 124.) floribus aggregatis sessilibus, squamis calycinis latis margine membranaceis mucronatis tubo longioribus, foliis lanceolatis multinerviis. 2ℓ Hab. . . . An D. barbati var.?
11. D. *JAPONICUS* (Thunb. fl. jap. 183. t. 23.) floribus aggregatis fasciculatis, squamis calycinis acutis ciliatis tubo dimidiò brevioribus, foliis ovatis brevibus. 2ℓ in Japoniâ. Habitus D. barbati var. pedunculosi, sed folia brevissima.
12. D. *PIXIFOLIUS* (Sibth. et Smith prod. 1. p. 284.*) floribus aggregatis capitatis, squamis calycinis obcordatis aristatis tubo brevioribus, foliis setaceis. 2ℓ in Lemno insulâ et circa Byzantium. — Tabern. ic. 668. f. 13. Aff. D. juniperino act. soc. Lin. 2. p. 303.
13. D. *CEPHALOTES* (Ser. mss.) floribus subsessilibus capitatis, squamis calycinis imbricatis apice mucronulatis patulis tubo paulò minoribus, foliis elongatis angustis, caule angulato subpiloso. 2ℓ Hab. . . . D. capitatus Poir. suppl. 4. p. 124. non DC.
14. D. *CAPITATUS* (DC! cat. h. monsp. 1813. p. 103.) glaucescens, floribus sessilibus capitatis, squamis calycinis latis ovatis longè aristatis capitulo longioribus, foliis superioribus basi dilatatis. 2ℓ in Tauriâ Caucasique herbidis. D. atro-rubens Bieb. fl. taur. 1. p. 324 non All. (v. s.)
15. D. *POLYMORPHUS* (Bieb. fl. taur. 1. p. 324. suppl. 298.) atro-virens, floribus sessilibus capitatis, squamis calycinis ovatis brevissimis submuticis foliis angustioribus scabris. 2ℓ in Tauriâ, Caucaso. D. atratus Beaupré in litt. (v. s. comm. à cl. Fisch. et Stev.)
- β. *diutinus* (Ser. mss.) floribus paniculato-fastigiatis solitariisque pedunculatis. D. diutinus Kit. in Link enum. 1. p. 419 et Schrad! in litt. (v. s.)
16. D. *BALBISII* (Ser. mss.) glaucescens, floribus subsessilibus capitato-fasciculatis, squamis calycinis lanceolatis patulis tubo paulò minoribus, foliis lanceolato-linearibus, caule angulato. 2ℓ prope Gennam. D. glaucophyllus Horn. ex herb. Balb. Folia lanceolato-linearibus basi non dilatata. (v. s.)
- β. *paniculatus* (Ser. mss.) floribus paniculato-fasciculatis.
17. D. *GIGANTEUS* (D'Urv! enum. pl. archip. p. 45.) viridis, floribus sessilibus capitato-hemisphæricis numerosis basi suffultis, bracteis foliaceis, squamis calycinis ovato acuminatis adpressis dimidium tubi æquantibus, foliis linearibus longissimis, basi longè connatis, caule tereti. 2ℓ ad littora Bulgarie in rupibus maritimis. Caules 2-3 ped. Flores purpureo-ferruginei. (v. s.)
18. D. *FERRUGINEUS* (Linn. mant. 561.) floribus aggregatis, involucris squamisque scariosis rufis oblongis aristatis calyce paulò brevioribus, foliis linearibus longè connatis. 2ℓ in Pyreneis Narbonensibus. — Barr. ic. t. 497. An D. Carthusianorum var.? (v. s.)
19. D. *CARTHUSIANORUM* (Lin. spec. 586.) floribus aggregatis sessilibus pedunculatisque, squamis calycinis ovatis aristatis tubo brevioribus, involucri oblongo aristato capitulo brevioribus, foliis linearibus trinerviis. 2ℓ in incultis et sterilibus totius Europæ. (v. v.)

- Æ. atrorubens* (Ser. mss.) floribus aggregato-capitatis sessilibus 3-8 — in Vallesiâ. *D. atrorubens* All. fl. pedem. n. 1545. Jacq. ic. rar. 3. t. 467. non Bieb. (v. v.)
γ. nanus (Ser. mss.) subacaulis, floribus aggregatis. — in Sancto-Go-thardo. (v. s.)
δ. anisopodus (Ser. mss.) floribus aggregatis pedunculatis sessilibusque. *D. atrorubens* prolifer Schleich! exs. — in Vallesiâ et circa Parisios. (v. s.)
 20. *D. ARBOREUS* (Lin. spec. 590.) caule fruticoso, floribus aggregatis, unguibus petalorum longissimis, squamis calycinis mucronatis arcuè imbricatis brevissimis, foliis subulatis subcarnosis, caule fruticoso. ζ in Cretâ. Lodd. bot. cab. 5. p. 459. ic. (v. s.)
 21. *D. JUNIPERINUS* (Smith in act. linn. soc. 2. p. 303.) caule fruticoso multifloro, floribus squamis calycis subquaternis obovatis mucronatis tubo duplò brevioribus, foliis subulatis pungentibus. ζ in Græciâ. An *D. arborei* var? †.
 22. *D. FRUTICOSUS* (Linn. spec. 591.) caule fruticoso, floribus aggregatis, unguibus petalorum calycem æquantibus, squamis calycinis mucronatis arcuè imbricatis brevissimis, foliis lanceolatis obtusis. ζ in Cretâ. — Tourn. itin. r. p. 183. t. 9. (v. s. comm. à cl. Sibth.)
 23. *D. SUFFRUTICOSUS* (Willd. enum. 466.) caule suffruticoso, floribus subaggregatis, squamis calycinis ovatis subulatis tubo triplò brevioribus, foliis lineari-lanceolatis utrinquè attenuatis. ζ Hab. Bractæ lineares reflexæ, squamæ calycinæ nervo exserto reflexo, corolla magna. (Link enum. 1. p. 419.)
 24. *D. AURICOLA* (Biv. Bern. sicul. cent. 1. p. 31.) floribus aggregatis fasciculatis, squamis calycinis interioribus brevibus margine ciliatis, exterioribus setaceo-subulatis longissimis, foliis crassiusculis linearibus glaucis glabris integerrimis. \mathcal{L} in fissuris rupium Siculæ. Flores rosei fragrantés. *D. Bisigniani* Tenor. cat. giard. S. Bisig. 1809. p. 18. fl. napol. t. 39. *D. involucratus* Poir. suppl. 4. p. 132. (v. s. s.)
 25. *D. CAROLINIANUS* (Walt. fl. car. 140.) floribus aggregatis longè pedunculatis, squamis calycinis tubo dimidiò brevioribus. — in Carolinâ. †.
 26. *D. CARYOPHYLLOIDES* (Schult. obs. 78.) floribus aggregatis fasciculatis, squamis calycinis ovatis mucronatis brevibus, petalis dentatis imberbibus, foliis latis canaliculatis enerviis connatis basi ciliatis. \mathcal{L} Flores eleganter rubri odore *D. caryophylli*. †.

Sect. II. CARYOPHYLLUM. Ser. mss.

Flores pauculati vel solitarii.

§. 1. *Petalis dentatis.*

27. *D. ASPER* (Willd. enum. 466.) floribus fasciculatis, squamis calycinis ovato-lanceolatis tubo brevioribus, petalis acutè dentatis, foliis lineari-lanceolatis 3-5-nerviis cauleque scabris apice serrulatis. \mathcal{L} .
a. angustifolius (Ser. mss.) pauciflorus, foliis linearibus acutis. — in Vallesiâ. *D. scaber* Schleich. ex! Sut. fl. helv. 1. p. 259. *D. serratus* δ . DC! fl. fr. 5. p. 601. (v. s.)
β. collinus (Ser. mss.) multiflorus, foliis lineari-lanceolatis. — in Hungariâ. *D. collinus* Waldst. et Kit! pl. rar. 1. p. 36. t. 38. *D. umbellatus* DC! cat. monsp. 104. (v. s.)
γ. serratus (Ser. mss.) canlibus 1-3-floris, floribus majoribus, foliis linearibus. — in Pyrenæis orient. *D. serratus* Lapeyr. abr. 241. et DC! fl. fr. 5. p. 601. var. α . (v. s. comm. ex herb. Xatard.)
 28. *D. CAMPESTRIS* (Bieb. fl. taur. 1. p. 326.*) caule paniculato subhirsuto, floribus solitariis, squamis calycinis ovatis acutis calyce dimidiò brevioribus,

- foliis subulatis. \mathcal{Z} in Tauriæ campestribus. D. bicolor Adams. Aff. D. aspero. (v. s. ex herb. Stev.) †.
29. D. CORYMBOSUS (Sibth. et Smith fl. græc. 1. p. 395. ex prod. 1. p. 285.) floribus subaggregatis, squamis lanceolatis villosis tubo brevioribus, caule divaricato multifloro pubescente. $\textcircled{1}$ in Asiâ minore.
30. D. ALPESTRIS (Balb! act. acad. taur. 7. p. 11. t. 1.) floribus subbinis, squamis calycinis 4, duabus interioribus latis mucronatis, petalis emarginatis. \mathcal{Z} in pascuis Alpinis. An D. asperi vel nitidi var.? † (v. s. comm. à cl. Balbis.)
31. D. NITIDUS (Waldst. et Kit! pl. rar. 2. p. 209. t. 191.*) floribus subfasciculatis binis, squamis calycinis aristatis calycem subæquantibus, petalis crenatis, caule decumbente flexuoso, foliis sublancoelatis obtusis. — \mathcal{Z} in Carpathis ad terminum Fagi. Ad D. Carthusianorum accedere videtur ob calycem aristatum, sed folia enervia potius D. alpinum L. referunt. (v. s. à cl. Kit. comm.)
32. D. RAMOSISSIMUS (Poir. suppl. 4. p. 130.) floribus solitariis, squamis calycinis 4 ovatis mucronatis tubo cylindraceo brevioribus, foliis planis angustis, caule debili ramosissimo. \mathcal{Z} ? in Tataria. Corolla alba.
33. D. DIFFUSUS (Sibth. et Smith fl. græc. 396. ex prod. 285.) floribus subcorymbosis, squamis sulcatis mucronatis tubo duplò brevioribus, caulibus diffusis glabresculis. \mathcal{Z} in insulâ Cypro.
34. D. ATTENUATUS (Smith act. soc. Lin. 2. p. 301.) floribus solitariis, squamis calycinis subsenis brevibus lanceolatis acuminatis margine membranaceis, calyce longo apice attenuato, petalis crenatis unguibus calycem superantibus, foliis subulatis. \mathcal{Z} in maritimis Gallia merid. Caules diffusi basi tortuosi ramosissimi. D. Lusitanicus Brot. fl. lus. 2. p. 173. t. 73. D. longiflorus Lam. dict. 4. p. 522. (v. s.)
35. D. HIRTUS (Vill. delph. 3. p. 593. t. 46.) floribus subsolitariis, squamis calycinis senis ovatis mucronatis calyce multò brevioribus, petalis crenatis, foliis subulatis margine scabris. \mathcal{Z} in collibus Delphinatùs et Galloprov., in Tauriâ. D. scabei Chaix non Thunb. nec Schleich. An potius D. attenuati var.? (v. s.) †
36. D. ACIPHYLLUS (Sieb! exs.) fruticosus, floribus paniculatis paucis, bracteis obovatis obtusissimis subobcordatis acutissimèque mucronatis calyce dimidiò brevioribus, petalis integris vel bilobis, foliis subdistichis linearibus pungentibus margine vix serrulatis non striatis numerosis. 5 in Creta. (v. s. in herb. Ser. comm. à cl. Sieber.)
37. D. GUTTATUS (Bieb. fl. taur. 1. p. 328. suppl. 300.*) caule paniculato lævi, floribus solitariis, squamis calycinis ovato-aristatis tubum subæquantibus, foliis subulatis nervoso-striatis. \mathcal{Z} circa Odessam. Facies D. montani sed arista multò longior. Petala purpurea punctis plurimis albis irrorata. (v. s. comm. à cl. Steven.)
38. D. PALLIDIFLORUS (Ser. mss.) caule lævi laxè paniculato, floribus solitariis, squamis calycinis ovato-lanceolatis aristatis tubo duplò brevioribus, foliis subulatis rectis margine scabris. \mathcal{Z} in herbidis ad Volgam. D. pallens Bieb. fl. taur. 1. p. 325. suppl. 298. non Sibth. (v. s. comm. à cl. Stev. et Goldb.) †
39. D. VERSICOLOR (Fisch. in Link enum. 1. p. 420.) caule multifloro lævi, squamis calycinis cuspidatis patulis tubo brevioribus, petalis fauce pubescentibus, foliis linearibus scabriusculis. \mathcal{Z} in imperio Ruthenico. (v. s. comm. à cl. Fisch.)
40. D. REPENS (Willd. spec. 2. p. 681.) caule unifloro, squamis calycinis binis ovatis acuminatis patulis ferè longitudine tubi, petalis dentatis, radice repente. \mathcal{Z} in Sibiria. †
41. D. PRATENSIS (Bieb! fl. taur. suppl. 300.*) caule paniculato, floribus solitariis, squamis calycinis foliaceis acuminatis subadpressis, exterioribus lanceolatis tubum æquantibus, interioribus basi ovatis brevioribus, petalis acutè

- dentatis, foliis subulato-lanceolatis. \mathcal{L} ? in Tauriâ. D. chlorolencus Fisch! in Horn. hort. hafn. suppl. 137. D. ochroleucus Link. enum. 1. p. 420. non Pers. D. Tataricus Fisch! in litt. (v. s. in h. Balb. comm. à cl. Bieberstein.)
42. D. CHINENSIS (Lin. spec. 588.) caule ramoso, floribus solitariis, squamis calycinis linearibus foliaceis patulis tubum æquantibus, petalis dentatis purpureis vel albis rubro maculatis, foliis lineari-lanceolatis. ① ② in Chinâ. — Mill. ic. 81. f. 2. (v. v.)
43. D. MONTANUS (Bieb. fl. taur. 1. p. 328.*) caule supernè ramoso confertim dichotomo, floribus solitariis, bracteis senis ovato-acuminatis tubo brevioribus, acumine subfoliaceo patulo, foliis lineari-subulatis trinerviis hirtis. \mathcal{L} in Caucaso. D. discolor Sims bot. mag. t. 1162. Flos magnus purpureus, petala subtus olivacea. (v. v. et s.)
44. D. OCHROLEUCUS (Pers. ench. 1. p. 494. non Link.) caule elongato ramoso, floribus solitariis, squamis calycinis 2 ovato-lanceolatis tubi dimidium superantibus, laciniis calycinis angustis longis, petalis obovato-linearibus subintegris, foliis brevibus lineari-subulatis. \mathcal{L} ? in Oriente.
45. D. CARYOPHYLLUS (Lin. spec. 537.) caule ramoso, floribus solitariis, squamis calycinis quaternis brevissimis ovatis submucronatis, petalis latissimis imberbibus, foliis lineari-subulatis canaliculatis glaucis. \mathcal{L} in Galliâ merid. Smith engl. bot. t. 214. Curt. bot. mag. t. 39. Variat. fl. semi-pleno et pleno, albo, purpureo, variegato etc.
- β . *imbricatus* (Sims bot. mag. t. 1622.) ramis abbreviatis, squamis calycinis numerosis imbricatis. (v. s.)
- γ . *carduinus* (Ser. mss.) foliis calycibus petalisque spinis fistulosis obsitis. D. caryophyllus var. Tratt. in florâ 1821. p. 717. ic.
46. D. SYLVESTRIS (Jacq. coll. 1. p. 237. ic. rar. t. 82.) caule ramoso vel simplici, floribus solitariis, squamis calycinis brevissimis quaternis ovatis, exterioribus acutis, interioribus obtusissimis, petalis latissimis imberbibus dentatis. \mathcal{L} in saxosis Alpium et Jurassi. D. virgineus Sims bot. mag. t. 1740. non Lin. An typus D. caryophylli? Variat caule brevissimo, 1- vel oo-floro. (v. v.)
47. D. LONGICAULIS (Tenor. cat. 1819. p. 76.) caulibus paniculatis multifloris cespitosis, floribus solitariis, squamis calycinis ovatis omnibus emarginato-cuspidatis, petalis crenatis imberbibus, foliis lineari-lanceolatis crassiusculis glaucis. \mathcal{L} prope Neapolim.
48. D. GRANDIFLORUS (Poir. dict. 4. p. 514.*) caule....., floribus subaggregatis, bracteis ovatis acutis tubo dimidiò brevioribus. \mathcal{L} in Hispaniâ. Υ .
49. D. MONADELPHUS (Vent. hort. cels. t. 39.*) caule dichotomo paniculato multifloro glauco, floribus solitariis, squamis calycinis quaternis lanceolatis subpungentibus patulis tubo brevioribus. \mathcal{L} in Oriente. D. procumbens Pers. ench. 1. p. 494. Calyx attenuatus; ovarium stipitatum. (v. s.)
50. D. BURCHELLII (Ser. mss.) caule ramoso, floribus solitariis, squamis calycinis 4 ovato-lanceolatis brevissimis submucronatis, petalis....., foliis linearibus subulatis striatis, inferioribus numerosissimis longissimis, caulinis rameisque squamiformibus brevissimis. \mathcal{L} in Africâ austr. (v. s. comm. à cl. Burch. cat. plant. Africæ austr. n. 2456.)
51. D. MICROPETALUS (Ser. mss.) caulibus cespitosis unifloris, squamis calycinis 4 ovato-lanceolatis mucronulatis calyce striato dimidiò brevioribus, dentibus ciliatis, petalorum lamina brevissimis dentatis, foliis parvis radicalibus subulatis dentato-ciliatis, caulinis squamiformibus. \mathcal{L} in Africâ austr. Folia Plantaginis subulata; calyx D. Caryophylli. (v. s. comm. à cl. Burch. cat. pl. Africæ austr. n. 1851.)
52. D. EMARGINATUS (Ser. mss.) caulibus cespitosis? 1-2-floris, squamis calycinis obtusissimis latissimis, calyce striato abbreviato, lobis obtusiusculis, petalis emarginatis minimis intus albescentibus, extus purpurascen-

tibus, foliis inferioribus numerosis linearibus planis serratis. $\frac{1}{2}$? circa Astypaleam. (v. s. comm. à cl. D'Urville.)

53. D. *SUAVEOLENS* (Spreng! nov. prov. 16.) caule....., floribus subgeminatis pedunculatis, squamis calycinis lanceolatis erecto-adpressis calyce brevioribus, petalis inciso-dentatis barbatis maculatis, foliis linearilanceolatis glaucis enerviis margine scabris. Ex Herb. Paris. Accepi sub nomine D. Chinensis. — Hab. . . . Proximus D. Caucasicus (Spreng. l. c.) An var. D. pratensis, Bieb? (Ser.) (v. s. in herb. Balb.)
54. D. *POIRETIANUS* (Ser. mss.) floribus terminalibus corymbosis, inferioribus solitariis, squamis calycinis apice subulatis tubo multò brevioribus, foliis oblongo-lanceolatis margine asperis. $\frac{1}{2}$ Hab. D. corymbosus, fl. purpureo. Poir. suppl. 4. p. 124. non Sibth. et Smith. Variat flore pleno. (v. s. in herb. Ser.)
55. D. *LIMOSCHITZIANUS* (Ser. mss.) caule subunifloro glabro, squamis calycinis longè ovato-acuminatis tubo paulò brevioribus, foliis subulatis strictis margine scabris. $\frac{1}{2}$ in Armeniæ, Iberiæ petrosis. D. petraeus Bieb! fl. taur. 1. p. 328. suppl. p. 300. non Waldst. et Kit. D. bracteatus herb. Willd. ex Stev. in litt. Squamis calycinis tubo duplò ferè brevioribus variat. (v. s. in herb. Balb. comm. à cl. Bieb.)
56. D. *SYLVATICUS* (Hoppe in Willd. enum. 467.) floribus solitariis subcorymbosis, squamis calycinis ovato-lanceolatis tubo brevioribus, foliis linearilanceolatis obsolete subtrinerviis glabris, petalis duplicato-dentatis. $\frac{1}{2}$ ad Ratisbonam. †
57. D. *CINNAMOMEUS* (Sibth. et Smith fl. græc. t. 400. ex prod. 1. p. 287.) caule....., floribus solitariis, squamis rhombeis obtusissimis brevissimisque, petalis emarginatis dentatis, foliis laxis obtusiusculis. $\frac{1}{2}$ in ericetis Laconiz et Asiæ minoris.
58. D. *POMERIDIANUS* (Lin. spec. 1673.) floribus solitariis, squamis calycinis ovatis acutis brevissimis, tubo apice tantum striato, petalis emarginatis subintegerrimis. $\frac{1}{2}$ in Palestinâ. Smith act. soc. Lin. 2. p. 300. D. tricolor Adam. sec. Fisch. (Steud.) Petala flava lateribus revoluta. (v. s. in herb. Balb. comm. à cl. Bieb.)
59. D. *LEPTOPETALUS* (Willd. enum. 468.) caule ramoso, floribus solitariis, squamis calycinis ovatis acutis brevissimis subaristatis, petalis lanceolatis angustis lateribus revolutis subintegerrimis, foliis subulatis scabriusculis. $\frac{1}{2}$ in Caucaso. Sims bot. mag. t. 1739. D. pomeridianus Bieb. fl. taur. 1. p. 329. non Lin. Petala supernè alba, subtus flavescenti-viridi tincta. An var. D. pomeridiani? (v. s.)
60. D. *PUNGENS* (Lin. mant. p. 240.) caule paucifloro, floribus solitariis, squamis calycinis brevissimis mucronatis patentibus, tubo gibbo, petalis integris, foliis cespitosius subulatis. $\frac{1}{2}$ in Hispaniæ maritimis.
 β . *Hispanicus* (Ser. mss.) squamis calycinis ovatis, petalis linearibus integerrimis. — Asso. syn. n. 371. t. 3. Dufour ann. gen. 7. p. 309. (v. s.)
61. D. *DIVARICATUS* (D'Urv. enum. pl. archip. p. 46.) caule ramoso divaricato, ramis longissimis strictis unifloris, squamis calycinis ovato-lanceolatis acutissimis margine membranaceis calycem subæquantibus, calycibus conicis multistriatis, striis granulato-punctatis, petalis acutè dentatis, foliis angustissimis. ② in insulâ Samo. D. tripunctatus Sibth. et Smith prod. ex D'Urville l. c. Flores dilutè purpurei extus virides. (v. s. comm. à cl. D'Urville.)
62. D. *BICOLOR* (Bieb. fl. taur. 1. p. 329.*) caule paniculato, floribus solitariis, squamis calycinis brevissimis suborbiculatis mucronulatis, petalis dilatatis, foliis subulatis, inferioribus tomentosis. $\frac{1}{2}$ in Tauriæ campes-tribus aridis. D. saxatilis Pall. ind. taur. Statura D. caryophylli. (v. s.) †

β. minor (Ser. mss.) foliis brevioribus, floribus minoribus, calycibus brevioribus basi turgidioribus. In aridis lapideis Tauriæ merid.

63. *D. FURCATUS* (Balb! act. acad. taur. 7. p. 12. f. 2. non Horn.) caule dichotomo 2-4-floro, floribus solitariis, pedunculis distantibus, squamis calycinis oppositis subbinis tubo multo brevioribus \mathcal{Z} in Pedemontio prope Tendam. *D. geminiflorus* Loisel. fl. gall. p. 725? Flores carnei. (v. s.)

64. *D. VIRGINEUS* (Lin. spec. 590.) caule subunifloro, squamis calycinis brevissimis obtusissimis binis distantibus, petalis crenatis, foliis cespitosis linearibus rigidis serrulatis. \mathcal{Z} circa Monspelium. *D. rupestris* Lin. f. suppl. p. 240. *D. pungens* Poir. dict. 4. p. 526. — Dill. elth. 401. p. 298. t. 385? (v. s.)

β. subacaulis (Ser. mss.) caulibus cespitosis brevissimis 1-floris, foliis floribus squamis calycinis abbreviatis. — In rupibus montis Ventosi.

D. subacaulis Vill. Delph. 3. p. 597. Loisel. not. p. 66. t. 6. f. 1. (v. s.)

65. *D. DELTOIDES* (Lin. spec. 588.) caulibus decumbentibus ramosis, floribus solitariis, squamis calycinis ovato-lanceolatis acutis subbinis, foliis superioribus angustis acutis, inferioribus oblongis obtusis. \mathcal{Z} in arenosis sylvaticis Europæ. *D. supinus* Lam. fl. fr. 2. p. 534. *D. Pyrenaicus* Pourr. act. toul. 3. p. 318. *D. Volgensis* hort. erf. ex herb. Balb. — Dill. elth. 400. t. 298. f. 384. (v. s. et v.)

β. glaucus (Ser. mss.) floribus albis, foliis cauleque glaucis. *D. glaucus* Lin. spec. 588. (v. s.)

66. *D. PALLENS* (Sibth. et Smith fl. græc. t. 399. ex prod. 1. p. 286.) caule paniculato multifloro, floribus solitariis, squamis ovatis acuminatis brevissimis, foliis laxis acutis. \mathcal{Z} in Asiâ minore prope Smyrnam.

67. *D. NAZARÆUS* (Clarke itin. vol. 3. ex Spreng. neue entd. 3. p. 161.) floribus solitariis, caulibus subramosis, squamis calycinis tubo dimidiò brevioribus adpressis ovatis acutis, petalis sexdentatis, foliis elongatis lineari-subulatis trinerviis margine scabris. — In Troadis monte Gargaro.

68. *D. MARGINATUS* (Poir. suppl. 4. p. 131.) floribus solitariis, squamis calycinis acutis inæqualibus calyce brevioribus, foliis linearibus marginatis. \mathcal{Z} ? in Jurasso. Aff. *D. nitido*. Squamæ calycis marginatæ; corolla alba.

69. *D. CRENATUS* (Thunb. prod. 81.) caule ramoso, floribus solitariis, squamis calycinis senis lanceolatis adpressis, petalis glabris cuneato-obovatis, unguibus calyce longioribus, foliis lineari-acuminatis subtetragono-canaliculatis. \mathcal{Z} ad Cap. B. Spei. Flos carneus. Calyx longè tubulatus.

70. *D. RIGIDUS* (Bieb. fl. taur. 1. p. 325. suppl. p. 298.) caulibus cespitosis suffrutescentibus paucifloris, floribus solitariis, squamis calycinis ovatis acutis brevibus, foliis subulatis patulis pubescenti-scabris. \mathcal{Z} in Tauriâ. Caulis lignescens distortus. Flores pallidi. (v. s. comm. à cl. Fisch. Stev. et Beaupré.)

71. *D. SERRATIFOLIUS* (Sibth. et Smith fl. græc. t. 402. ex prod. 1. p. 287.) caulibus unifloris, radice multicipite lignosa, squamis quaternis, petalis dentatis imbebibus, foliis serratis pungentibus. \mathcal{Z} in monte Hymetto prope Athenas.

72. *D. CLAVATUS* (Spreng. neue entd. 2. p. 169) caule unifloro, squamis binis ovatis acutiusculis brevissimis patulis, calyce medio constricto, petalis crenatis fauce nudis, foliis linearibus canaliculatis margine scabriusculis. — Hab. An *D. cæsius*? (Spreng.) An *D. plumarius*? (Ser.)

73. *D. GRACILIS* (Sibth. et Smith fl. græc. t. 404. ex prod. 1. p. 288.) caulibus 1-floris, squamis mucronatis subsenis, petalis crenatis barbatis, foliis acuminatis margine scabris. \mathcal{Z} in monte Atho.

74. *D. SUAVIS* (Willd. enum. suppl. p. 24. Spreng. neu. entd. 2. p. 168.) caule subunifloro, squamis calycinis quaternis acutis abbreviatis, petalis barbatis duplicato-serratis, foliis linearibus patentibus. — Hab. Hic videtur *D. cæsius* Smith engl. bot. t. 62. (Spreng. l. c.) Potius *D. plumariæ* var. (Ser.)

75. *D. strictus* (Sibth. et Smith fl. græc. t. 403. ex prod. 1. p. 288.) caulibus unifloris, squamis quaternis, petalis crenatis imberbibus, foliis planis obtusiusculis margine scabris. $\frac{1}{2}$ in monte Atho.
76. *D. cæsius* (Smith act. soc. Linn. 2. p. 302. engl. bot. t. 62.) caulibus cespitosus subunifloris, squamis calycinis subrotundis brevibus, petalis crenatis pubescentibus, foliis obtusiusculis margine scabris. $\frac{1}{2}$ in saxosis montanis Jurassi, etc. *D. pulchellus* Pers! ench. 1. p. 495. *D. cespitosus* Lam. dict. 4. p. 525. — Dill. elth. p. 401. t. 298. f. 385. opt. An huc referend. *D. nitid.* Poir. suppl. 4. p. 131? (v. v. et s.)
77. *D. alpinus* (Linn. spec. 590.) caule unifloro, squamis calycinis exterioribus tubum æquantibus, interioribus dimidio brevioribus, petalis crenatis, foliis oblongis obtusis. $\frac{1}{2}$ in Styriâ et Austriâ. — Clus. hist. 1. p. 283. f. 1.
78. *D. glacialis* (Hænk. in Jacq. coll. 2. p. 84.) caulibus erectis cespitosus brevibus subunifloris, squamis calycinis elongatis tubum æquantibus vel superantibus, calyce striato, petalis serratis, foliis linearibus acutis serrulatis. $\frac{1}{2}$ in montanis Gallo-Provinciæ et Delphinatûs. *D. alpinus*. DC! fl. fr. 4. p. 746. *D. neglectus* Lois. not. 65. (v. s.)
- β. acaulis* (Ser. mss.) caule brevissimo. — In Monte-Cenisio.
- γ. latifolius* (Ser. mss.), foliis oblongo-linearibus, flore magno. *D. alpinus* herb. Balb. ex *Alpibus austriacis* à cl. Frœlich. miss.
79. *D. tener* (Balb! act. acad. taur. 7. p. 13. t. 3.) caule unifloro, petalis fimbriatis glabris, squamis calycinis quaternis lanceolato-linearibus tubo paulò brevioribus. $\frac{1}{2}$ ad margines agrorum Tendæ. Flores rubri. (v. s.)
80. *D. elegans* (D'Urv. enum. pl. archip. p. 46. n. 875.) caulibus erectis strictis unifloris, squamis calycinis ovatis mucronatis medium tubi æquantibus, calyce striato, lobis obtusis submembranaceis, petalis dentatis, foliis linearibus acutis longissimis striatis integerrimis. $\frac{1}{2}$ in ins. Cos. An *D. crinitus* Sibth. et Smith ex D'Urv. l. c. (v. s. comm. à cl. D'Urville.)
81. *D. multipunctatus* (Ser. mss.) caulibus erectis strictis subunifloris elongatis, squamis calycinis ovatis longè mucronatis tubum æquantibus margine membranaceis, calyce verrucoso-multipunctato, laciniiis nervosis, petalis maculatis barbatis 3-4-dentatis, foliis linearibus acutis integris longissimis. $\frac{1}{2}$ ad vias in ins. Cos. (v. s. comm. à cl. D'Urville.)
82. *D. pumilus* (Vahl symb. 1. p. 32.) cespitosus acaulis, foliis linearibus. $\frac{1}{2}$ in Arabiâ felici. *D. uniflorus* Forsk. cat. pl. arab. p. 111. n. 284. Calyx glaber striatus, squamæ 4 oblongæ acuminatæ flore breviores, exteriores minores.
83. *D. hornemanni* (Ser. mss.) pedunculis bifidis terminalibus, squamis calycinis lanceolato-cuspidatis erectis tubo brevioribus, petalis incisis, foliis linearibus nervosis margine serrulato-scabris. $\frac{1}{2}$ in Italiâ et ad Tergestum *D. furcatus* Horn. h. hafn. suppl. p. 47. non Balb.
84. *D. leucophæus* (Sibth. et Smith fl. græc. t. 405. ex prod. 1. p. 288.) caulibus unifloris, squamis 4, petalis 3-dentatis imberbibus, foliis ovato-lanceolatis basi 5-nerviis. $\frac{1}{2}$ in montis Olympi Bithyni summitate.
85. *D. sternbergii* (Sibth. ex cat. h. taur. 1821. p. 24.) caulibus subbifloris, squamis calycinis 4 ovatis acutis adpressis tubo duplò brevioribus, petalis cuneiformibus serratis pubescentibus, foliis linearibus. $\frac{1}{2}$ Hab.
86. *D. purpureus* (Lam. dict. 4. p. 523. ill. t. 376. f. 2.) caule subunifloro, floribus solitariis, squamis calycinis latis tubo dimidiò brevioribus acutis, petalis integriusculis. $\frac{1}{2}$ Hab. Calyx amplus brevis. (v. s.)
87. *D. petræus* (Waldst. et Kit! pl. rar. 3. t. 222.*) caule subunifloro, squamis calycinis obovatis mucronatis, petalis imberbibus multifidis, foliis subulatis integerrimis glabris nervosis. $\frac{1}{2}$ in Hungariâ. Flores albi. An var. *D. plumaris*? (v. s. in herb. Balb. comm. à cl. Kit)

ℓ? floribus majoribus, petalis subdeflexis. D. petræus Sims bot. mag. t. 1204. An D. plumarii var?

88. D. SCABER (Thunb. prod. p. 81. non Schleich.) caulibus unifloris, squamis calycinis quaternis lanceolatis, petalis crenatis. — Hab. †.

89. D. CESPITOSUS (Thunb. prod. p. 81.) caulibus unifloris, squamis calycinis 4 lanceolatis, petalis integris, foliis triquetris. 2ℓ ad Cap. B.-Spei. †.

90. D. BIFLORUS (Sibth. et Smith fl. græc. t. 393. ex prod. 1. p. 285.) floribus geminatis, squamis calycinis cuneatis obtusissimis aristatis patulis calyce brevioribus, foliis linearibus trinerviis. 2ℓ in monte Delphi Eubæ.

91. D. IBERICUS (Willd. enum. suppl. p. 24.) caule lævi, foliis scabriusculis, squamis patulis acumine lanceolato, petalis pubescentibus. 2ℓ Hab. D. Willdenovii Link enum. 1. p. 420. Caulis pauciflorus. Bractæ calyce paulò breviores patulæ, coroll. purpurascens. Link. l. c. (v. s. in herb. Balb. comm. à cl. Panzer.)

92. D. TRIPUNCTATUS (Sibth. et Smith fl. græc. t. 398. ex prod. 1. p. 286.) caule patulo multifloro, floribus solitariis, squamis scariosis tenuiter aristatis tubo paulò brevioribus. ☉ in insulâ Cypro.

93? D. RUTHENICUS (Roem. in Poir. suppl. 4. p. 131.) floribus solitariis, squamis calycinis involucrentis lanceolatis acutis tubo brevioribus, foliis linearibus acutis. — Cult. in h. Mus. Par. †.

§. 2. *Petalis fimbriatis.*

94. D. GALLICUS (Pers. ench. 1. p. 495.) caulibus adscendentibus subunifloris, squamis calycinis brevibus ovatis submucronatis, petalis dentato-multifidis, foliis linearibus subciliatis. 2ℓ in Galliæ meridionalis arenâ mobili. D. arenarius DC. syn. n. 4325.* ic. gall. rar. p. 14. t. 41.* non Linn. (v. s.)

95. D. ALBENS (Ait. h. kew. ed. 1. vol. 2. p. 90. ed. 2. v. 3. p. 8.) caulibus...., floribus solitariis, squamis calycinis lanceolatis quaternis brevibus, petalis emarginatis. 2ℓ ad Cap. B.-Spei. Differt à D. deltoide petalis vix crenatis et absentia circuli purpurei (Ait. l. c.) Folia angustissima striata serrulata, calyx longissimus, petalorum lamina ovata apice fimbriato-dentata. (v. s. comm. à cl. Jan.)

96. D. SERRULATUS (Desf. fl. atl. 1. p. 346.) foliis lanceolatis serratis, pedunculis unifloris, squamis externis imbricatis acutis calyce interiore brevioribus, petalis fimbriatis. — 2ℓ in arenis apud Tunetanos. Corolla pallidè rosea duplò minor quam in D. plumario Linn. (Desf. l. c.)

97. D. PLUMARIUS (Linn. spec. 589. hort. upsal. 105. n. 4.*) glaucescens, caulibus 2-3-floris, dentibus obtusis, bracteis subovatis brevissimis mucronulatis, petalis laciniato-multifidis barbatis, foliis linearibus margine scabris. 2ℓ Hab. D. dubius Hornem. hort. hafn. 1. p. 408? D. moschatus hort. parl. Variat fl. simplici et pleno, petalis albis, purpureis, maculatis, plùs minùsve fimbriatis, floribusque suaveolentibus. (Vulgò *Mignardise*.) (v. v. et s.)

ℓ. *hortensis* (Ser. mss.) petalis fauce barbatis. D. hortensis Schrad. ex Willd. enum. p. 469. (v. v.)

γ. *Portensis* (Ser. mss.), foliis minoribus, petalis vix fimbriatis. D. portensis Libosch. ex herb. Balb!

98. D. CAUCASEUS (Sims bot. mag. t. 795.*) caule paniculato paucifloro, floribus solitariis, squamis calycinis ovato-acuminatis adpressis, petalis cuneiformibus equaliter inciso-serratis, foliis glaucis margine scabris. 2ℓ in pratis subalpinis Caucasi orient. Bieb. fl. taur. 1. p. 327. Aff. D. plumario, sed squamis calycinis elongatis (v. s. comm. à cl. Fisch.)

99. D. SAXATILIS (Pers. ench. 1. p. 494.) cespitosus subdecumbens, caule 2-3-floro, squamis ovatis mucrone longiusculo distinctis tubo brevioribus, petalis multifidis fauce glabris. 2ℓ? ad rupes prope Clermont. D. plumarius

quorumdam, sed folia linearia non glauca; corolla fauce non pubescens, nec squamæ obtusæ. (Pers. l. c.) An *D. plumarius* var. *hortensis*?

100. *D. squarrosus* (Bieb. fl. taur. 1. p. 331. suppl. p. 302. cent. 1. t. 33.) caulibus subunifloris, squamis calycinis ovatis acutis brevissimis, petalis multifidis, foliis subulatis canaliculatis rigidis brevibus recurvis. $\frac{1}{2}$ in Tauriâ *D. arenarius* Pall. it. 3. p. 600. Flores *D. plumario* similes, sed calyx longior. Petala albida tenuissimè lacinia. (v. s. comm. à cl. Fisch.)
101. *D. MUSENI* (Horn. h. hafn. 1. p. 408) caulibus subunifloris procumbentibus, squamis calycinis ovalibus mucronatis tubo triplo brevioribus, petalis multifidis imberbibus, foliis subulatis reflexis. $\frac{1}{2}$ in Caucaso? Var. videtur *D. squarrosi*. †.
102. *D. FRAGRANS* (Bieb. fl. taur. 1. p. 331. suppl. p. 301.) caulibus subunifloris, squamis calycinis senis ovato-lanceolatis acuminatis tubo brevioribus, petalis semi-multifidis imberbibus, foliis subulatis margine scabrisculis. $\frac{1}{2}$ in Caucaso subalpino. Sims bot. mag. t. 2067? *D. multifidus* Willd. herb. ex Stev. in litt. Folia angustissima longa; calyx elongatus; petala alba purpureo-maculata; laciniae calycinæ angustæ acutæ. (v. s.)
- 103? *D. PUNCTATUS* (Spreng. neue entd. 2. p. 169.) caule ramoso paucifloro, squamis calycinis quateruis adpressis obtusiusculis brevissimis, petalis imberbibus multifidis punctatis, foliis glaucis linearibus flaccidis. Hab. Nullo affinis nisi *D. fragrant* Adam. qui differt squamis calycinis acuminatis subaristatis, petalis immaculatis, foliis rigidis viridibus subulatis, etc. (Spreng. l. c.)
104. *D. SEROTINUS* (Waldest. et Kit. pl. rar. 2. p. 188. t. 172.) caulibus subunifloris, squamis calycinis subovatis obtusiusculis calyce quadruplo brevioribus, petalis multifidis subnudis, foliis subulatis glaucis ciliatis. $\frac{1}{2}$ in Hungariâ. *D. arenarius* Towns. it. Hung. 3. p. 488. t. 16. *D. plumarius* Æ. Wahl. fl. carp. p. 126. An potius *D. plumarii* var? (v. s.)
105. *D. ARENARIUS* (Linn. spec. 589) caulibus subunifloris, squamis calycinis ovatis obtusis, petalis multifidis, foliis linearibus. $\frac{1}{2}$ in Europæ frigidioris arenâ mobili. Sims bot. mag. t. 2038. Petala ultra medium disci divisa, lobis angustissimis, basi maculâ lividâ et pilis purpurascensibus adpressis. An var. *D. plumarii*? (v. s. comm. à cl. Schrad. et Fisch.)
106. *D. PROSTRATUS* (Jacq. hort. Schœn. 3. p. 11. t. 271.) caule fruticoso, floribus laxè paniculatis, squamis calycinis lanceolatis calyce elongato triplo brevioribus, petalis fimbriatis glabris, foliis linearibus in egerimis. $\frac{1}{2}$ ad promontorium R.-Spei. Aff. *D. fimbriato* et *superbo*.
107. *D. FIMBRIATUS* (Bieb. fl. taur. 1. p. 332.* suppl. p. 302.*) caule suffruticoso basi ramoso subbifloro, floribus solitariis, squamis calycinis senis lanceolatis. calyce brevioribus, petalis oblongis cuneatis multifido-dentatis imberbibus, foliis subulatis scabris. $\frac{1}{2}$ in Iberiæ saxosis. *D. orientalis* Sims bot. mag. t. 1069. *D. contortus* Smith in Rees. cycl. vol. XI. Flores *D. plumario* minores, calyce lineari squamis triplo longiore. (v. s.)
108. *D. CRINITUS* (Smith. act. soc. Lin. lond. 2. p. 300.*) floribus solitariis, squamis calycinis ovalibus mucronatis subdivergentibus tubo triplo brevioribus, petalis multifidis imberbibus... — in Oriente. Willd. spec. 2. p. 678. Petala angusta, limbo ad basin usque irregulariter multifido-capillaceo. (v. s.)
109. *D. PLUMOSUS* (Spreng! pugill. 2. p. 64.) floribus sparsis solitariis, squamis calycinis lanceolato-linearibus patentibus tubo paulo brevioribus, petalis imberbibus multifido-incisis, foliis linearibus nervosis flaccidis. $\frac{1}{2}$? in monte Baldo. Fortè *D. Monspeulanus* huc pertinet. Spreng. l. c. (v. s. in herb. Balb.)
110. *D. MONSPEULANUS* (Lin. amœn. 4. p. 313.* spec. p. 588.) caule paniculato paucifloro, floribus solitariis, squamis calycinis subulatis rectis tubo dimidio brevioribus, petalis digitato-multifidis fauce glabris, foliis linearibus

serrulatis. \mathcal{U} in Pyrenæis et Jurasso. D. Monspeliacus DC! fl. fr. n. 4324. D. Sternbergii Schleich! exl. (v. v. et s.)

β . *brevifolius* (Ser. mss.) foliis caulibusque abbreviatis. — in Pyrenæis prope St.-Jean de Luz. (v. s. comm. à cl. Bory St.-Vincent.)

111. D. *SUPERBUS* (Lin. amœn. 4. p. 272 * spec. p. 589.) caule paniculato multifloro, floribus subfastigiatis, squamis calycinis brevibus ovatis mucronatis, petalis ultra medium pinnato-multifidis fauce pilosis. \mathcal{U} in nemoribus et pratis umbrosis montanis Delaun. herb. amat. t. 21. Caryophyllus sylvestris vi Clus. hist. 1. p. 284.

β . *rubicundus* (Ser. mss.) petalis purpureis (v. v. et s. in h. Ser.)

112. D. *FISCHERI* (Spreng. cat. sem. h. hal. 1810. pl. min. cog. 2. p. 62.) caule. . . . floribus subaggregatis, squamis calycinis ovatis cuspidatis erectis tubo duplò brevioribus, petalis multifidis barbatis, foliis lanceolatis uninerviis scabriusculis. \mathcal{U} ? ad Moscoviam. Petala rosea barbata. Habitus D. collini, sed petalis multifidis. †

113. D. *LIBANOTIS* (Labill! pl. syr. 1. p. 14. t. 5.*) caule erecto, floribus subaggregatis, squamis calycinis senis acuminatis divaricatis tubo brevioribus, petalis multifidis barbatis, foliis lanceolatis. \mathcal{U} in altis montibus Libani. (v. s.)

IV. SAPONARIA. Lin. gen. n. 769.

Calyx tubulosus 5-dentatus basi nudus. Petala unguiculata, ungue calycem æquante. Stam. 10. Styli 2. Caps. 1-locularis.

Sect. I. VACCARIA. Dod. pempt. 104.

Flores paniculati, calyx angulosus glaber.

1. S. *VACCARIA* (Lin. spec. 585.) floribus paniculatis, calycibus pyramidalis quinqueangularibus glabris, bracteis membranaceis acutis, foliis ovato-lanceolatis sessilibus. ① inter segetes Europæ. — J. Bauh. hist. 3. p. 357. f. 2. (mala) Lychuis vaccaria Scop. fl. caru. n. 511. Petala rubra.

β . *grandiflora* (Fisch! in litt.) petalis latis. — In Iberiâ. (v. s.)

2. S. *PARFOLIATA* (Roxb. hort. bengal. p. 34. Willd. enum. p. 464.) floribus paniculatis, calycibus pyramidalis 5-angularibus glabris, bracteis foliaceis acutis, foliis oblongo-lanceolatis basi connatis. ② in Indiâ Orient. An S. vaccariæ var? †.

Sect. II. BOOTIA. Neck. delic. gallo-belg. 1. p. 193.

Flores fasciculato-paniculati, calyx teres sæpius pilosus.

3. S. *OFFICINALIS* (Lin. spec. 584.) floribus fasciculato-paniculatis, calycibus teretibus villosis luteolis, petalorum appendicibus linearibus, foliis ovato-lanceolatis acutis obtusisve. \mathcal{U} ad margines viarum Europæ. Smith engl. bot. t. 1060. Bootia vulgaris Neck. delic. gallo-belg. 1. p. 193. Petala rosea. Variat flore pleno. (v. v.)

β . *glaberrima* (Ser. mss.) foliis calycibusque glaberrimis. (v. s.)

γ . *hybrida* (Lin. spec. 584.) — moris. hist. 2. p. 548. s. 5. t. 22. f. 52.

4. S. *CORYMOINES* (Lin. spec. 585.) caulibus procumbentibus dichotomis, floribus fasciculatis, calycibus teretibus villosis purpureis glanduloso-pilosis, foliis ovato-lanceolatis subuninerviis. \mathcal{U} in Europæ rupibus calcareis. Jacq. fl. austr. 5. p. 59. append. t. 23. Cav. ic. 2. p. 29. t. 134. (v. v.)
5. S. *GLUTINOSA* (Bieb. fl. taur. 1. p. 322. cent. 2. t. 66.) caulibus erectis ramosis, floribus paniculatis corymbosisque, calycibus longè teretibus glanduloso-hispidis, foliis ovatis trinerviis. ② in Tauriæ montosis. Silene armeria Pall. ind. taur. Flores magnitudine Silenes conoidæ; petala minutissima sanguinea apice bidentata, ad faucem coronata. (v. s. comm. à cl. Stev. et Beaup.)

Sect. III. PROTEINIA. *Ser. mss.*

Flores solitarii axillares. Petala fauce nuda? Calyx pilosus raro glaber.

6. *S. ILLYRICA* (Lin. mant. 70.) caule erecto ramosissimo dichotomo viscido, floribus laxè spicatis alternis, calycibus subcylindricis semi-5-fidis margine membranaceis, petalis integris punctatis, foliis linearibus glabris. \mathcal{L} in Illyriâ. Ard. spec. 2. p. 24. t. 9. *Gypsophila Illyrica* Sibth. et Smith. fl. græc. t. 386 ex prodr. 1. p. 281. (v. s.)
7. *S. CRETICA* (Lin. spec. 584.) caule dichotomo ramoso villos-viscido, floribus axillaribus solitariis strictis, calycibus quinquangularibus, foliis linearisubulatis lævibus, petalis parvis integris. \mathcal{L} ? in Cretæ aridis. *Saxifraga altera* Alp. exot. 292. t. 291.
8. *S. PORRIGENS* (Lin. mant. 239. *) caule erecto, ramis divaricatis piloso-viscidis, floribus longè pedunculatis axillaribus, pedunculis filiformibus, calycibus teretibus, fructibus ovoideis pendulis, foliis lanceolato-linearibus. \odot in Oriente. Jacq. h. vind. 2. p. 49. t. 109. *Silene porrigens* Gouan. ill. 29. (v. s.)
9. *S. ORIENTALIS* (Lin. spec. 585. *) caule dichotomo, ramis divaricatis, floribus axillaribus, pedunculis rigidis florem æquantibus calycibusque hispidulis teretibus, laciniis acutis, foliis lineari-spathulatis. \odot in Oriente et Carniolâ. — Dill. h. elth. 205. t. 167. f. 204. (v. s.)

β . *lævis*, glaberrima, ramis divaricatissimis. (v. s. cult. in h. DC. et Ser.)

Sect. IV. BOLANTHUS. *Ser. mss.*

Caules foliaque cespitosi. Flores aggregati. Calyx teres villosus.

10. *S. PROSTRATA* (Willd. enum. 465. *) caule prostrato, ramis adscendentibus brevibus, floribus corymbosis confertissimis, calycibus cylindricis hirtis, foliis oblongo-spathulatis. \mathcal{L} in Galatiâ. Corolla \hat{S} . ocyroidis, sed parum minor.
11. *S. HIRUTA* (Labill. icon. pl. syr. dec. 4. t. 4. f. 2. *) caulibus adscendentibus paucifloris, floribus terminalibus subcapitatis sessilibus, calycibus 5-angularibus foliisque lanceolatis hirtis, petalis integerrimis. \mathcal{L} in Oriente. (v. s. comm. à cl. Labill.)
12. *S. DEPRESSA* (Biv. stirp. rar. manip. 2. *) cespitosa, floribus subumbellatis pedunculatis, calycibus longissimis 5-angularibus pubescenti-viscosis, petalis bifidis acutè coronatis, foliis radicalibus cespitosiis obovato-ellipticis depressis. \mathcal{L} in collibus sabulosis regionis apertæ M. Ætnæ. Flores terminales rosei magni. *S. cespitosa* Biv. in Rafin. stat. gan. di Sicil. p. 27. Bon. t. 163. f. 1. *S. Sicula* Rafin. specch. 2. p. 7. Cupani panphyt. t. 167. f. 1. (v. s.)
13. *S. CESPITOSA* (DC. rap. voy. 2. p. 78. 1808. fl. fr. 5. p. 601. * ic. gall. rar. fasc. 2. ined.) cespitosa; caulibus subnudis apice floriferis, floribus subumbellatis, calycibus cylindricis villosis profundè lobatis, lobis acutis, petalis apice emarginatis appendice bifido, lobis angustissimis, foliis linearibus glabris subradicalibus vix denticulatis. \mathcal{L} in sterilibus Pyrenæorum editionum. *S. elegans* Lapey. abr. pyr. p. 238. — *Habitus* *S. lateæ*, sed flores rosei majores. (v. s.)
14. *S. LUTEA* (Lin. spec. 585. excl. syn. Bocc.) cespitosa, caulibus diphyllis, floribus capitatis involucreatis, calycibus cylindricis lanatis, lobis obtusis brevibus, petalis obovatis integris nudis, foliis linearibus basi ciliatis subradicalibus. \mathcal{L} in Alpibus Vallesiacis et Pedemontanis. All. fl. ped. n. 1560. t. 23. f. 1. Corolla flava, Stam. violacea. *Habitus* *Lychnidis Alpinae*. (v. s.)
15. *S. BELLIDIFOLIA* (Smith spic. bot. 5.) cespitosa, caulibus tetraphyllis, floribus capitatis densis, calycibus teretibus hirtis, petalis linearibus crenatis, foliis ovato-lanceolatis margine undulatis. \mathcal{L} in Italiæ montibus altissimis. Portenschlag. pl. dalm. t. 7. f. 2? — Bocc. mus. p. 75. t. 62. f. 1. Folia radicalia Globulariæ simillima. Stam. flava.

† *Species non satis notæ.*

16. *S. SMITHII* (Ser. mss.) caulibus simplicibus unifloris nudiusculis, calycibus basi contractis glabris, foliis spathulatis cespitosis. \mathcal{L} in Delphi monte Eubææ. *S. cespitosa* Sibth et Smith fl. græc. t. 389. ex prodr. 1. p. 282. Smith. in Rees. cycl. vol. 31.

17. *S. SAXATILIS* (Bory ann. gen. 3. 1820. p. 13.) caulibus divaricatis dichotomis, foliis ovato-oblongis oppositis sessilibus, pedunculis longissimis, petalis emarginatis. \mathcal{L} ? in rupibus Sierræ-Nevadæ.

V. CUCUBALUS. Gaertn. fruct. 1. p. 376. t. 77. f. 7.

Calyx campanulatus 5-dentatus nudus. Petala 5 unguiculata, limbo bifido. Capsula carnosa 1-locularis.

1. *C. BACCIFERUS* (Lin. spec. 591.) ramis divaricatis, foliis ovatis, calycibus campanulatis, petalis distantibus. \mathcal{L} in umbrosis Europæ. — Mill. ic. t. 112. (v. v.)

VI. SILENE. Lin. gen. n. 772. Gaertn. fruct. 2. p. 233. t. 130 f. 8. (1).

Calyx tubulosus 5-dentatus nudus. Petala 5 unguiculata, fauce sæpissimè coronata, limbo bifido. Stamina 10. Styli 3. Capsulæ basi 3-loculares apice in 6 dentes dehiscentes.

Sect. I. NANOSILENE. Otth mss.

Cespitosæ. Caules ferè nulli. Calyx subinflatus. Pedunculi uniflori.

1. *S. ACAULIS* (Lin. spec. 603.) glabra, caulibus densis humilibus, foliis lineari-lanceolatis, floribus abortu dioicis, pedunculis brevibus unifloris, calycibus campanulatis, petalis obovatis seu obcordatis. \mathcal{L} in Alpibus. Sims bot. mag. t. 1081. Petala purpurea. (v. v.)

♂. floribus majoribus longius pedunculatis, staminibus exsertis, ovario stylisque abortivis. *S. acaulis* Lin. l. c. DC. fl. fr. 4. p. 749. (v. v.)

♀. floribus minoribus ferè sessilibus, staminibus abortivis, stylis longè exsertis. *S. exscapa* All. ped. n. 1584. t. 79. f. 2. *S. acaulis* β *exscapa* DC. fl. fr. 4. p. 749. (v. v.)

β. *elongata* (DC. fl. fr. 4. p. 749.) pedunculo elongato, flore masculino. (v. v.)

γ. *parviflora* (Otth mss.) pedunculo nullo, flore minimo feminino. (v. s.)

δ. *alba* (Otth mss.) flore albo. (v. v.)

ε. *plena*, floribus magnis polypetalis — in monte Joms Alpium Rhæticiæ. (v. s.)

2. *S. PUMILIO* (Sturm deutschl. fl. 1. fasc. 22. t. 11.) caulibus minus densis, foliis lineari-spathulatis subpubescentibus, pedunculis brevibus unifloris, calycibus inflatis hirsutis. \mathcal{L} in Alpibus Germaniæ. Petala purpurea. *Cucubalus pumilio* Lin. Mant. 71. (v. s.)

β. *alba* (Otth mss.) flore albo. Sturm. l. c. in descr.

Sect. II. BEHENANTHA. Otth mss.

Caulescentes. Flores solitarii aut paniculati. Calyx vesiculato-inflatus.

3. *S. FIMBRIATA* (Sims bot. mag. t. 980.) pubescens, foliis magnis ovato-lanceolatis, floribus amplo-paniculatis, calycibus amplissimè vesiculatis, dentibus latis, petalis fimbriatis planis post anthesin incurvis. \mathcal{L} in Cretæ et Sicilia. *Cucubalus fimbriatus* Bieb. fl. taur. 1. p. 333. suppl. 303. (v. v.)

4. *S. LACERA* (Sims bot. mag. t. 2255.) hispida, foliis ovato-lanceolatis longè petiolatis undulatis, calycibus amplissimè vesiculatis, petalis lacris appendice bipartito, staminibus alternis deflexis. \mathcal{L} ? in Alpibus Caucasi. *Cucubalus lacerus* Bieb. fl. taur. 3. p. 303. Petala alba. (v. s.)

(1) Hoc genus proprio Marte elaboravit ornat. Adolph. OTTH Bernensis.

5. *S. STELLATA* (Ait. h. Kew. ed. 2. v. 3. p. 84.) caulibus erectis ramosis pubescentibus, foliis quaternis verticillatis lanceolatis longè acuminatis glabris, floribus paniculatis, calycibus vesiculosis pubescentibus, petalis fimbriatis. \mathcal{L} in Virginiâ et Canadâ. *Cucubalus stellatus* Lin. spec. 592. Petala alba. (v. s.)
6. *S. COULTERIANA* (Otth mss.) glaberrima, caule ramoso, foliis parvis linearilanceolatis, floribus maximis paniculatis, flore longissimè pedunculo in dichotomiâ, calycibus subsphæricis amplissimè inflatis, petalis unguiculo cuneiformi, lamina ferè bipartita, lobis latis subtruncatis. \mathcal{L} ? in Iberniâ et circa Constantinopolim. (v. s.)
7. *S. INFLATA* (Smith fl. brit. 467.) caulibus ramosis, floribus paniculatis, calycibus vesiculato-ovatis, petalis bifidis nudis, unguiculis cuneiformibus, stylis longissimis. \mathcal{L} in Europâ frequens. *Cucubalus behen* Lin. spec. 591. Petala alba.
- a. vulgaris* (Otth mss.) glabra, foliis lanceolatis. — In agris. *Cucub.* behen Lin. spec. 591. Smith engl. bot. t. 164. (v. v.)
- ß. angustifolia* (DC! fl. fr. 4. p. 747.) glabra, foliis linearibus. *Cucubalus angustifolius* Schrank hort. monac. t. 83. Ten. fl. neap. t. 37. (v. v.)
- γ. rubra* (Ram. pyr. ined.) petalis purpureis. In Pyrenæis et alpinis Berrensibus. (v. v.)
- δ. maritima* (DC! fl. fr. 4. p. 747.) piloso-pubescent, foliis ovato-lanceolatis. — In maritimis Galliæ. (v. s.)
- ε. viridiflora* (DC! fl. fr. 4. p. 747.) calycibus foliaceis profundè 5-lobis, petalis semi-abortivis viridibus. *Cucubalus viridis* Lam. dict. 2. p. 221.
- ζ. castrata* (Lapeyr. abr. pyr. 247.) uno sexu petalisque abortivis. — in Pyrenæorum pratis humidis. (v. v.)
- η. uniflora* (Otth mss.) humilis, glabra, floribus solitariis rarò geminis aut ternis. — in Alpibus. *S. uniflora* Roth cat. 1. p. 52. (v. s.)
- θ. fabaria* (Otth mss.) repens. foliis minoribus ferè spathulatis. — in Alpibus Saxoniz. *Cucub. fabarius* Thor. chl. land. 172. *S. uniflora* γ. DC. fl. fr. 4. p. 747. (v. s.)
8. *S. OVATA* (Pursh fl. bor. amer. 1. p. 316.) caule simplici, foliis ovato-lanceolatis acuminatis, floribus paniculatis, calycibus ovatis, genitalibus exsertis. \mathcal{L} in Georgiâ et Carolinâ. Petala pallida aut alba. An *S. inflatæ* var.?
- * *flore pleno*. *Cucub. polypetalus* Walth. fl. carol. 141? (v. s.)
9. *S. BEHEN* (Lin. spec. 599.) glabra ramosa, foliis lanceolatis, inferioribus petiolatis, floribus paniculatis, calycibus ovatis striato-venosis, dentibus brevibus, petalis bilobis, lobis brevissimis obtusis, appendicibus 2 bicuspidatis. \mathcal{L} in Cretâ. *Lychnis vesicaria* etc. Dill. elth. 427. t. 317. f. 409. (v. s.)
10. *S. INDICA* (Roxb. cat. h. calc. p. 34.) pubescens, caulibus altissimis ramosis, foliis magnis lanceolatis, floribus paniculatis, calycibus ovatis 10-striatis reticulatis, dentibus longis obtusiusculis, petalis unguiculo lato basi piloso, lamina bifida basi margine bidentata. \mathcal{L} in Napauliâ. Petala purpurea. (v. s.)
11. *S. LANUGINOSA* (Bertol. journ. bot. 4. p. 76.) fruticoso-cespitosa, caulibus lanuginosis, foliis lanceolato-linearibus margine lanuginosis, inferioribus longissimis, calycibus ovatis pubescentibus, petalis bifidis (quadrifidis ex Bertol.) \mathcal{L} in Italiâ. Lych. Alpina, etc. Till. cat. 4. pil. 105. t. 41. f. 2. (v. s.)
12. *S. GRAMINIFOLIA* (Otth mss.) glabra, caule erecto simplici basi valdè foliosissimo suprâ ferè nudo, foliis linearibus vix ciliatis, floribus spicato-paniculatis, pedunculis oppositis nunc calyce longioribus nunc brevioribus, calycibus ovatis 10-striatis, petalis semibifidis, unguiculis ciliatis. \mathcal{L} ? in Alpibus Altaicis. (v. s.)
13. *S. VISCAGINOIDES* (Horn. h. hafn. suppl. 4. p. 49.) glabra, caule erecto simplici subfolioso, foliis linearibus vix ciliatis, floribus spicato-paniculatis, pedunculis oppositis longitudine calycis, calycibus ovatis 10-striatis, petalis semibifidis, unguiculis non ciliatis. \mathcal{L} ? in Dahuriâ. (v. s.)

14. *S. PROCUMBENS* (Murr. com. goett. 1784 et 1785. p. 83. t. 2.) glabra, caulibus procumbentibus ramosis foliosissimis, foliis lanceolatis, floribus axillaribus oppositis et terminalibus, calycibus ovatis, petalis subauriculatis bifidis, anthophoro longo. $\frac{1}{2}$ in Sibiria. *S. decumbens* Schreb. (v. s.)
15. *S. RUBELLA* (Lin. spec. 600. fere glabra, caulibus parum ramosis, foliis obovatis serrulato-ciliatis, floribus paniculatis, calycibus subovatis 10-striatis, petalis obcordatis, appendicibus bifidis obtusis in anulum coalitis. ① in Lusitania et Oriente. Delile fl. ægypt. t. 29. f. 3. (v. s.)
16. *S. OBLONGIFOLIA* (Otth mss.) vix pubescens, caulibus assurgentibus, foliis oblongis obtusis, floribus parvis paniculatis laxis, calycibus subinflatis vix 10-striatis, anthophoro longo. ①? Patr. ign. (v. s.)
17. *S. APETALA* (Willd. spec. 2. p. 307.) pubescenti-canescens, caule erecto ramoso, foliis lanceolatis, superioribus linearibus, floribus paucis terminalibus aut in dichotomia, calycibus obovatis 10-striatis, petalis nullis. ①? in Hispania. (v. s.)
18. *S. SPERGULIFOLIA* (Bieb. fl. taur. suppl. 305.) caulibus procumbentibus diffusis 2-3-chotomo-ramosis, foliis parvis linearibus, floribus paniculatis confertis, calycibus inflatis 10-striatis, petalis semibifidis subdeflexis, appendice obcordata. $\frac{1}{2}$ in Armenia. Desf. cor. Tournef. 73. t. 55. *S. polyphylla* Bieb. fl. taur. n. 835. excl. syn.
19. *S. GYPSOPHYLLA* (Desf. cat. hort. paris. 184.) fere glabra, caulibus flexuosis ramosissimis foliosis, foliis lineari-lanceolatis, floribus paniculatis confertis oppositis, calycibus vesiculato-clavatis 10-striatis, petalis bilobis. $\frac{1}{2}$ Hab. Petala alba.
20. *S. CARYOPHYLLOIDES* (Otth mss.) caulibus erectis gracilibus, foliis linearibus angustissimis, superioribus latioribus, floribus magnis terminalibus, calycibus inflatis subinfundibuliformibus basi angustatis, petalis bilobis. — in Oriente. *Cucubalus caryophylloides* Poir. dict. suppl. 2. p. 416. An hujus sect.?
21. *S. HISPANICA* (Otth mss.) pilosa, caulibus procumbentibus subtetragonis, foliis lineari-oblongis obtusis, floribus axillaribus fere solitariis, calycibus ovatis 10-striatis, petalis minimis, anthophoro nullo. $\frac{1}{2}$ in Hispania. *S. parviflora* Zea in Poir. dict. suppl. 5. p. 150. An hujus sect.?
22. *S. CARNOsa* (Moench. suppl. 206.) glabra, caule erecto, foliis acutis glaucis, floribus solitariis, calycibus lævibus anastomosantibus, petalis lanceolatis, appendice bipartita. ① Patr. ign. Petala purpurea albo marginata. An hujus sect.?
23. *S. ANGUSTIFOLIA* (Bieb. fl. taur. 1. p. 337.) caule ramoso, foliis lineari-subulatis glabris, floribus terminalibus, calycibus subcampanulatis subhirsutis, petalis bifidis, anthophoro vix capsulæ longitudine. $\frac{1}{2}$ circa portas Caucasicas. An hujus sect.?

Sect. III. OTITES. *Otth mss.*

Caulescentes. Flores verticillato-spicati.

24. *S. OTITES* (Pers. ench. 1. p. 497.) caulibus erectis raro ramosis vix pubescentibus parum foliosis, foliis inferioribus numerosis spathulatis subcarnosis, superioribus lanceolatis, floribus dioicis parvis, calycibus femininis sphaericis, masculinis subclavatis, petalis linearibus indivisis. $\frac{1}{2}$ In Silesia, Austria, Gallia, etc., in arenosis. Smith engl. bot. t. 85. (v. s.)
- β. umbellata* (Otth. mss.) radice crassa ramosa, caulibus aphyllis humilibus, foliis radicalibus spathulatis, floribus umbellatis. $\frac{1}{2}$ in Syria. (v. s.)
- γ. macrophylla* (Otth. mss) subpubescens, caule altissimo ramoso, foliis magnis spathulatis obtusis seu acutis, floribus numerosissimis subpaniculatis. $\frac{1}{2}$ in Galloprovinciæ monte Cousson. (v. s.)
- δ. densiflora* (Otth mss.) pilosa, caule altissimo, verticillis densis multifloris distantibus. — in Tauria. *S. densiflora* D'Urv. enum. 48.

25. *S. WOLGENSIS* (Othh mss.) caule pubescenti-ramoso, foliis inferioribus magnis lanceolato-spathulatis, superioribus linearibus longis, floribus paniculato-verticillatis pedunculatis, petalis linearibus. \mathcal{L} ad Wolgam. Cucub. Wolgensis Willd. enum. suppl. 24. (v. s.)
26. *S. PARVIFLORA* (Pers. ench. 1. p. 497.) canescenti-pubescent, caulibus assurgentibus ferè simplicibus, foliis spathulato-lanceolatis, verticillis inferioribus in paniculas elongatis, calycibus sphaericis 10-striatis. \mathcal{L} in Hungariâ. (v. s.)
27. *S. EFFUSA* (Othh mss.) caulibus erectis ferè simplicibus, foliis linearibus, inferioribus obtusiusculis, floribus numerosissimis minimis effuso-paniculatis, calycibus obovato-clavatis 10-striatis. \mathcal{L} ad Wolgam. Cucubalus effusus Fisch! in litt. \mathcal{H} spicato-paniculati. (v. s.)
28. *S. VERTICILLATA* (Othh mss.) glaberrima, caulibus ramosissimis foliosissimis, foliis linearibus acutis, floribus verticillato-spicatis, verticillis distantibus, calycibus ovato-clavatis, petalis bifidis. \mathcal{L} Cucub. cespitosus Poir. dict. suppl. 2. p. 416.*
29. *S. SIBIRICA* (Pers. ench. 1. p. 497.) suffruticosa glabra, caulibus ramosissimis, foliis lineari-lanceolatis brevissimè ciliatis numerosis, floribus interruptè spicatis, calycibus subinflato-clavatis 10-striatis, petalis indivisis (emarginatis) genitalibus longis. \mathcal{L} in desertis Sibiricæ. Cucub. Sibiricus Lin. spec. 592. (v. v.)
30. *S. DISTANS* (Othh mss.) caule longissimo virgato vix pubescente, foliis radicalibus latis lanceolato-spathulatis obtusis, caulinis linearibus apice rotundatis, verticillis longè distantibus paucifloris, calycibus longioribus clavatis. \mathcal{L} ? Hab....
31. *S. MULTIFLORA* (Pers. ench. 1. p. 497.) vix pubescens, caule simplici, foliis lineari-lanceolatis, inferioribus latoribus petiolatis, floribus interruptè spicatis, pedunculis brevibus, calycibus clavato-cylindricis 10-striatis, petalis bipartitis, lobis angustatis, genitalibus longissimis. (1) ? in Sibiria et Hungariâ. Cucub. multiflorus Waldst. et Kit. pl. hung. 1. p. 56. t. 56. (v. s.)
32. *S. ELATA* (Othh. mss.) glabra, caule longissimo virgato simplici, foliis caulinis paucis linearibus, radicalibus lanceolato-spathulatis, floribus interruptè spicatis, verticillis 2-6-floris, calycibus clavatis non striatis, petalis bipartitis. \mathcal{L} in Tauriâ. *S. chlorantha* Stev! in litt. non Willd. (v. s.)
33. *S. RUTHENICA* (Othh mss.) caulibus repentibus ramosissimis, ramis oppositis erectis, foliis lineari-lanceolatis serrulatis, florum spicâ longissimâ, pedunculis calycem æquantibus, calycibus clavatis 10-striatis, petalis bipartitis. \mathcal{L} in Russiâ? Affin. *S. Sibiricæ* sed distinctissima. An *S. Tatarica*? (v. s.)
 β . *pedunculata* (Othh mss.) pedunculis calycem multò superantibus inferioribus ramosis. \mathcal{L} in Russiâ et Volhiniâ. (v. s. comm. à cl. Stev. et Goldb.)
34. *S. STATICEFOLIA* (Sibth et Smith fl. græc. t. 435. ex prodr. 1. p. 301.) glabra, foliis oblongis linearibus, floribus subverticillato-spicatis, spicâ elongatâ, calycibus inflatis, petalis bifidis. \mathcal{L} ? in Oriente.
35. *S. TATARICA* (Pers. ench. 1. p. 497.) glabra, caulibus erectis simplicibus foliosissimis, foliis lanceolatis parvis, florum spicâ elongatâ densâ, verticillis 2-4-floris, calycibus clavatis 10-striatis reticulatis, petalis bipartitis, genitalibus longissimis. \mathcal{L} in Tatariâ. Cucub. Tataricus Linn. spec. 592. (v. v.)
36. *S. GIGANTEA* (Linn. spec. 598.) velutinâ, foliis radicalibus cochleariformibus carnosis, verticillis distantibus, calycibus subtubulatis 10-striatis, petalis bifidis, genitalibus longissimis. \mathcal{L} in Africâ. *Lychnis*, etc. Walth. hort. 32. t. 11. Folia magna, obovata. (v. v.)
37. *S. VISCOSA* (Pers. ench. 1. p. 497.) pubescens viscosissima, caule simplici crasso folioso, foliis magnis lineari-lanceolatis undulatis, floribus magnis

nutantibus, spicâ longâ, calycibus cylindrico-clavatis 10-striatis, petalis bipartitis, genitalibus longissimis. ② in Italiâ, Angliâ. Sueciâ et in Oriente. Cucub. viscosus Linn. spec. 592. Folia ferè Cynoglossi offic. (v. v.)

38. *S. INVOLUTA* (Forsk. fl. ægypt. arab. suppl. 210. n. 47.) caulibus crassis sublignosis villosis, foliis lineari-lanceolatis subtùs tomentosis, floribus oppositis sæpè supernè verticillatis, calycibus 10-angularibus, petalis bifidis. ② prope Hellespontem.

Sect. IV. CONOIMORPHA. *Outh mss.*

Caulescentes. Calyx conoideus, fundo umbilicato dentibus longissimis.

39. *S. CONICA* (Linn. spec. 598.) pubescens, foliis linearibus mollibus, floribus solitariis aut paniculatis, calycibus brevioribus conicis 30-striatis, petalis bipartitis, capsulâ ovatâ. ① in Galliâ, Hispaniâ et Orientis arenosis. Jacq. fl. austr. t. 253. Petala purpurea. (v. s.)
40. *S. CONOIDEA* (Linn. spec. 598.) caulibus pubescentibus, foliis lanceolato-linearibus ferè glabris, floribus solitariis aut paniculatis, calycibus longis conoideis 30-striatis, petalis integris obovatis, capsulâ lageniformi. ① in Galliâ, Germaniâ, etc. arenosis. — Clus. hist. 1. p. 288. f. 2. Petala purpurea. (v. v.)
41. *S. CONIFLORA* (Nees! in litt.) caulibus pubescentibus, foliis gramineis longis acutis vix pubescentibus, floribus paniculatis, calycibus cylindrico-conicis 30-striatis, petalis obcordatis. ① in Mesopotamiâ (v. s. ex b. Olivier.)
42. *S. CYLINDRIFLORA* (Outh mss.) pubescens, caulibus ramosis foliis lineari-lanceolatis, floribus paucis paniculatis, calycibus cylindrico-conicis 30-striatis, petalorum unguiculo calycem superante unguiculato laminâ bifidâ. ② in Oriente. (v. s. comm. à cl. Rousseau.)
43. *S. UNDULATA* (Ait. h. kew. 2. p. 96.) pubescens, foliis lanceolatis undulatis inferioribus petiolatis, floribus magnis laxè dichotomo-paniculatis, calycibus longissimis cylindrico-conicis 10-striatis, petalis bilobis subreticulatis. — Ad cap. Bonæ-Spei. Ungues petalorum longissimi. (v. s.)

Sect. V. STACHYMORPHA. *Outh mss.*

Caulescentes. Flores spicati axillares, non oppositi. Calyx 10-striatus.

§. 1. *Calycibus cylindricis.*

44. *S. ANGLICA* (Linn. sp. 594.) pilosa, caulibus ramosis, foliis lanceolatis acutis, calycibus ventricosocylindricis 10-striatis, dentibus longissimis acutis, petalis obcordatis parvis, appendicibus pyramidalibus. ① in Angliâ, Galliâ, etc. Smith engl. bot. t. 1178. Petala alba. (v. v.)
45. *S. LUSITANICA* (Linn. spec. 594.) pilosissima, caulibus ramosissimis, foliis inferioribus obovato-spathulatis, superioribus lanceolatis obtusis, spicâ distichâ, calycibus cylindrico-ventriculosis 10-striatis, dentibus longioribus, petalis indivisis, lamina triangulari margine exteriore denticulatâ. ① in Lusitaniâ. Dill. elth. t. 311. f. 401. Petala incarnata. (v. s.)
46. *S. TRIDENTATA* (Desf. fl. atl. 1. p. 349.) caulibus ramosis, foliis lineari-lanceolatis, spicâ secundâ, calycibus subcylindricis 10-costatis, dentibus longis, petalis tridentatis. ① in Algeriâ et Hispaniâ arvis. Petala rosea. (v. s.)
47. *S. GALLICA* (Linn. spec. 595.) hirsuta, caulibus ramosis, foliis inferioribus spathulatis, superioribus lanceolatis obtusis, spicâ secundâ, calycibus cylindrico-ventriculosis 10-striatis, dentibus brevibus acutis, petalis obovatis coronatis. ① in Galliâ. — Vaill. bot. par. t. 16. f. 12. Petala carnea. (v. s.)
48. *S. COARCTATA* (Lag. gen. et spec. 15) foliis inferioribus lanceolatis petiolatis, superioribus lanceolato-linearibus, floribus subsessilibus, calycibus hirsutis, fructiferis ovatis fance coarctatis, petalis bifidis. ① in Valenciâ et Murciâ.
- 49? *S. OCYMOIDES* (Desf. cat. hort. par. 184.) pilosa, caulibus ramosis, foliis

- spathulatis, spicâ secundâ pauciflorâ, calycibus cylindrico-ventriculosis pilosissimis, dentibus longioribus, petalis obovatis vix crenatis, appendicibus 2 longioribus. ① Hab. . . . S. pedicellata. Poir. suppl. 5. p. 150. Petala purpurea margine pallescentia. (v. s. in h. Ser.) †.
50. S. DISTICHA (Willd. enum. p. 476.) pilosa, caule ramosissimo, foliis lanceolatis cuspidatis, spicis geminatis densis, flore solitario in dichotomiâ, calycibus cylindricis 10-striatis, petalis parvis bifidis. ① Hab. . . . S. microphylla Roem. in Schrank pl. rar. t. 39. Petala rosea. (v. v.)
51. S. CERASTOIDES (Linn. spec. 596.) caulibus simplicibus villosis, foliis pubescentibus, inferioribus spathulatis, superioribus lanceolatis, spicâ distichâ pauciflorâ, calycibus cylindrico-ventriculosis 10-striatis pilosissimis, petalis obovatis retusis coronatis. ① in Europâ australi. S. rigidula Linn. amœn. 4. p. 313. (v. s.)
52. S. QUINQUEVULNERA (Linn. spec. 595.) pubescens, caulibus ramosis, foliis lanceolatis, inferioribus obtusis, spicâ secundâ, calycibus villosissimis rostriatis dentibus brevibus, petalorum laminâ subrotundâ, appendicibus bicauspidatis. ① in Hispaniâ, Galliâ, etc. Smith engl. bot. t. 86. Cucub. variegatus Lam. fl. fr. 3. p. 28. Petala purpurea margine pallescentia. (v. v.)
53. S. SCIOTICA (Otth mss.) pilosa, caule erecto simplici, foliis lucidis acuminatis, radicalibus latis spathulatis, floribus disticho-spicatis confertis, calycibus vesiculato-cylindricis, petalis obcordatis. ① in insulâ Scio. Petala prioris. (v. s. ex h. Olivier.)
54. S. NOCTURNA (Linn. spec. 595.) caule ramoso infrâ piloso, foliis pubescentibus basi longè ciliatis, inferioribus spathulatis, superioribus lineari-lanceolatis, spicis secundis densis, floribus adpressis, calycibus cylindricis ferè glabris reticulato-costatis, petalis bipartitis angustatis ① ②³ in Hispaniâ, Galliâ, etc. S. spicata. DC. fl. fr. 4. p. 759. (v. v.)
β. pauciflora (Otth mss.) floribus paucis distantibus, petalis minoribus. Cucub. reflexus. Linn. spec. 594. S. mutabilis Linn. spec. 596.
- 55? S. CINEREA (Desf. fl. atl. 1. p. 355.) sericeo-canescens, caulibus ramosis, foliis inferioribus ovatis, floribus subsessilibus solitariis binis aut ternis, calycibus cylindricis 10-striatis, petalis bifidis angustatis. ①² in arvis Algeriz. Petala alba.
56. S. BRACHYPETALA (Rob. et Cast! mem ined. in DC. fl. fr. suppl. 607.) caule simplici, foliis obovato-spathulatis obtusis hirsutis basi longè ciliatis, floribus secundis erectis, calycibus subcylindricis hirsutis, dentibus longis, petalis bifidis minimis. ① Massiliæ. (v. s.)
57. S. HIRSUTISSIMA (Otth mss.) pilis longis densis albis, caule ramoso, foliis lineari-lanceolatis, bracteis longis, floribus secundis erectis, calycibus cylindricis, anthophoro brevi. ① in Hispaniâ. S. hirsuta Lagasc. varied. de cienc. 1805. p. 212.
- 58? S. MICROPETALA (DC! cat. monsp. 146. non Lag.) pilosa, caule ramosissimo folioso, foliis lineari-lanceolatis, floribus terminalibus aut in dichotomiâ, calycibus cylindricis, petalis bifidis, anthophoro brevi. ① Hab. . .
59. S. MICRANTHA (Link Cav. herb.) hirsuta, floribus sessilibus secundis, calycibus cylindricis adpressis, petalis parvis profundè marginatis. ① in Lusitaniâ. S. micropetala Lagasc. gen. et spec. 15. non DC.
60. S. CANARIENSIS (Spreng. neue endt. t. 3. p. 60.) caule subramoso folioso, piloso, foliis magnis ovato-lanceolatis ciliatis, floribus secundis subnutantibus, calycibus cylindricis longis pilosis. ① in insulis Canariis. (v. s.)
61. S. SETACEA (Otth mss.) caulibus numerosis diffusis sæpius simplicibus, foliis obovatis obtusis setaceo-pilosissimis, floribus laxis secundis, calycibus cylindricis pilosis, petalis bifidis, nuguicula calycem superante. ①² in insulâ Melos. (v. s. comm. à el. D'Urville.)

§. 2. *Calycibus clavatis.*

62. *S. VILLOSA* (Forsk. descr. plant. cent. 3. p. 88. n. 71.) viscido-pubescent, caulibus numerosis procumbentibus foliosis, foliis oblongis obtusis, floribus laxis secundis, pedunculis post anthesin deflexis, calycibus cylindrico-clavatis, petalis longissimè unguiculatis bifidis. ♀ ? in Ægypto. (v. s.)
63. *S. HISPIDA* (Desf. fl. atl. 1. p. 348, non hortul.) hirsutissima pilis longis albis, caule ramosissimo, foliis latius lanceolatis obtusiusculis, bracteis brevibus, floribus secundis suberectis, spicis sæpè dichotomis, calycibus longè clavatis, petalis semibifidis. ① in Barbariâ. *S. hirsuta* Poir. dict. 7. p. 169. (v. s.)
64. *S. LAXIFLORA* (Brot. flor. lus. 2. p. 188.*) pilosa, caulibus erectis geniculatis ramosis, foliis inferioribus lanceolatis, superioribus sublinearibus, floribus secundis remotiusculis, calycibus subclavatis, petalis bifidis, capsulâ cylindræâ. ① in Lusitaniâ. An species?
65. *S. CANESCENS* (Tenor. prodr. 25.) hirsuto-canescens, caulibus prostratis ramosis, foliis obovato-spathulatis basi ciliatis, floribus secundis erectis, calycibus vesiculato-subcylindricis, petalis bifidis. ♀ prope Neapolim. (v. s.)
66. *S. DECUMBENS* (Bern sicul. cent. 1. p. 75.) pubescens, caulibus numerosis decumbentibus diffusis ramosis, foliis parvis spathulato-lanceolatis, floribus inclinatis, spicâ dichotomâ, calycibus clavatis angustis, petalis longè unguiculatis bipartitis. ① in Hispaniâ et circa Neapolim. (v. s.)
67. *S. LERCHENFELDIANA* (Baumg. stirp. transylv. 1. p. 398.) glabra, caulibus decumbentibus filiformibus, foliis margine vix crenulatis, radicalibus lineari-lanceolatis, caulinis ovato-lanceolatis, floribus secundis, pedunculis patentid-flexiusculis, calycibus clavatis, petalis emarginatis. — in Alpibus Hunyadensibus.
68. *S. OLIVERIANA* (Oth mss.) subpubescens, caule erecto subsimplici, foliis linearibus basi ciliatis, floribus paucis, calycibus longè clavatis, petalis semibifidis subangustis. ① ? circa Alep et Mossul. (v. s. ex h. Olivier.)
69. *S. DIANTHYFOLIA* (Oth mss.) ferè glabra, caule ramosissimo folioso, foliis linearibus acutis, floribus terminalibus seu axillaribus, calycibus clavatis, petalis bipartitis, genitalibus longissimis. — In Sibirîâ. Cucub. fruticosus Pall. (v. s. comm. à Cl. Deless.)
70. *S. DIFFUSA* (Oth mss.) pubescens, caulibus diffusis ramosis, foliis parvis spathulatis obtusis basi ciliatis, floribus suberectis, spicâ 2-5-florâ, calycibus clavatis subangustis, petalis bifidis longè unguiculatis. ♀ ? in littore Massensi. *S. sericea* Bertol! in litt. non All. (v. s.)
71. *S. IBERICA* (Bieb. fl. taur. 1. p. 335.) caule ramoso pubescente, foliis lanceolatis pubescentibus, spicis secundis dichotomis, calycibus brevibus ovato-clavatis glabris, petalis bifidis. — In Turiâ et Iberiâ. An *S. dichotoma*? (v. s.)
72. *S. DICHOTOMA* (Ehrh. beytr. 7. p. 144.) caulibus ramosis pubescentibus, foliis scabris basi ciliatis, inferioribus spathulatis, superioribus lanceolatis, spicis geminis secundis, floribus sessilibus subnutantibus, calycibus subventricosis 10-costatis villosis, petalis bipartitis. ① in Hungariâ. *S. membranacea* Poir. dict. 7. p. 165. (v. v.)
73. *S. NYCTANTHA* (Willd. enum. 472.) pubescens, foliis subcarnosis, inferioribus spathulatis, superioribus lanceolatis, floribus secundis, calycibus longè clavatis post anthesin subtrigonis, petalis bifidis. ① Hab.
74. *S. TRINERVIA* (Seb. et Maur! fl. rom. 152.) pilosa, pilis nodosis, caule gracili ramoso, foliis lineari-lanceolatis subtus trinerviis, floribus secundis, calycibus clavatis hispidis, pilis sursum imbricatis, petalis semibifidis. ① circa Romam. (v. s. comm. ex h. Mauri.)
- 75? *S. VELUTINA* (Pouff. in Desf. herb. ex Loisel. in journ. 2. p. 324.) velatina, caulibus erectis ramosis, foliis inferioribus obovato-lanceolatis, superioribus lanceolato-linearibus, floribus nunc in axillis solitariis, nunc agglomeratis, calycibus longè clavatis, petalis semibifidis. ♀ in Corsicâ.

76. *S. SABULIFLORUM* (Link in Spreng. 1. nov. prov. 39.) hirsuta, foliis lanceolatis, floribus spicatis et in dichotomiis ramorum, calycibus longè pilosis angustis cylindrico-clavatis, petalis emarginatis. ①? Petala purpureo-vioacea. (v. s.)
77. *S. BELLIDIFOLIA* (Jacq. hort. vind. 3 t. 81.) pilosa, caule erecto gracili ramoso, foliis lanceolatis, spicis geminatis secundis, calycibus cylindrico-clavatis, petalis bifidis, antophoro longo. ① Hab... (v. s.)
78. *S. CHEIRANTHIFOLIA* (Schott in Isis 1818, fasc. 5. p. 821.) floribus secundis, calycibus breviter clavatis, petalis bipartitis, capsulâ cylindroidæ, seminibus dorso vix revolutis canaliculatis. — ad San-Rocco — Petala intus rosea.
79. *S. VESPERTINA* (Retz obs 3. p. 31.) pubescens, caulibus ramosis, foliis spathulatis obtusis, spicis geminis secundis, calycibus vesiculato-clavatis, petalis bipartitis, lobis ovatis. ① in Mauritania et Lusitaniâ. S. bipartita Desf. atl. 1. p. 352. t. 100. S. distachya Brot. fl. lus. 2. p. 189.† (v. v.)
80. *S. OBTUSIFOLIA* (Willd. enum. p. 473.) caule hirsuto, foliis elliptico-spathulatis rotundatis pubescentibus, floribus secundis cernuis, calycibus clavatis pubescentibus, petalis bifidis. ①? Hab... Petala purpurea. An *S. colorata*?
81. *S. COLORATA* (Schousb. maroc.) caulibus ramosis hirsutissimis, foliis obovato-spathulatis numerosis pubescentibus basi ciliatis, floribus secundis, calycibus vesiculato-clavatis, petalis bipartitis coronatis. ① in Marocco et in insulâ Scio. Horn. h. hafn. 1. p. 412. (v. s.)
82. *S. CRASSIFOLIA* (Lin. spec. 597.) velutina, caule procumbente ramoso folioso, foliis ovato-spathulatis carnosius, bracteis minimis, floribus secundis, calycibus vesiculato-clavatis, petalis longè unguiculatis emarginatis. ① ad caput B.-Spei. Petala tristis coloris. (v. s.)
83. *S. BURCHELLII* (Otth mss.) pubescens, caulibus assurgentibus simplicibus, foliis parvis lanceolatis, floribus paucis subsessilibus, calycibus longissimè clavatis, antophoro longissimo. — ad cap. B.-Spei. Burchell cat. geogr. n. 271. (v. s.)
84. *S. GRACILIS* (DC! cat. monsp. 145.) glabra, caule erecto gracili ramoso, foliis linearibus vix ciliatis, inferioribus ovatis, floribus erectis alternis distantibus, pedunculis longis, calycibus vesiculato-clavatis, petalis bipartitis, lobis linearibus. ① Hab... Petala alba. (v. s.)
85. *S. IMBRICATA* (Desf. fl. atl. 1. p. 349. t. 98.) infernè pilosa, caulibus erectis ramosis, foliis lanceolatis, spicis secundis densis, floribus erectis, calycibus clavatis, petalis obcordatis, limbo dentibus 2 marginalibus coronatis. ① prope Mascar.
86. *S. LINEARIFOLIA* (Otth mss.) caule recto parum ramoso glabro, foliis angustissimis acutis, floribus erectis secundis, calycibus vesiculato-clavatis amplis basi bracteatis, petalis bifidis coronatis. — in Alpibus Caucasiciis. (v. s.)
87. *S. JENISENCIS* (Willd. enum. 473.) glabra, caulibus sæpius simplicibus, foliis subcarnosis linearilanceolatis, floribus secundis, calycibus ovatis ventricosius, petalis bifidis, appendicibus 4-lobis. 2½ in Sibiria. *S. Jenisea* Poir. dict. suppl. 5. p. 154.
88. *S. ALTAICA* (Pers. ench. 1. p. 497.) radice crassâ lignosâ subramosâ, caulibus simplicibus assurgentibus, foliis linearibus acutissimis, floribus in summis caulibus paucis alternis, calycibus cylindraceo-ventricosius, petalis revolutis bifidis linearibus. 2½ in Sibiria. Cucub. fruticosus Gmel. syst. 2. p. 713. An *S. ciliata*?
89. *S. CILIATA* (Pourr. act. toul. 3. p. 328.) pubescens, caulibus numerosis prostratis basi foliosissimis, foliis linearibus setaceo-ciliatis, floribus paucis secundis, calycibus inflato-clavatis, petalis bipartitis, sinibus calycis deflexis. 2½ in Creta et Pyrenæis. *S. stellata* Lapeyr. non Ait. *S. arvensis* Lag. in var. de cienc. 1805. p. 212. *S. Pourretii* Poir. Petala purpurea. (v. s.)

- β. geniculata* (DC! fl. fr. 4. p. 756.) vix pubescens, calycibus pallidioribus, petalis albis. *S. geniculata* Pourr. act. toul. 3. p. 323.
90. *S. LEGIONENSIS* (Lag. gen. et spec. p. 14.*) caule gracili simplici, foliis radicalibus lanceolatis acutis in orbem prostratis breviter ciliatis, caulinis paucis lineari-subulatis, bracteis ovato-acuminatis, floribus 5-6 secundis, calycibus clavatis membranaceis. \mathcal{L} in Hispaniæ provinciâ Legionensi.
91. *S. SESSILIFLORA* (Desf. in Poir. dict. suppl. 5. p. 154.) villosa, caulibus erectis vel procumbentibus subsimplicibus subquadrangularibus, foliis inferioribus oblongo-spathulatis subcarnosis, superioribus angusto-linearibus, floribus sessilibus dichotomo-spicatis, calycibus turgidis, petalis bipartitis. (1) in Syriâ.
92. *S. PENDULA* (Linn. spec. 599.) pubescens ramosa supina, foliis ovato-lanceolatis, floribus axillaribus pendulis, calycibus clavato-inflatis, petalis bifidis coronatis. (1) in Sicilia. Sinus bot. mag. t. 114. Petala incarnata.
- β. erectiflora* (Oth mss.) floribus minoribus erectis. In Lusitaniâ. *S. scabrisflora* Broter. fl. lus. 2. p. 184.*
93. *S. LONGICAULIS* (Pourr. elench. hort. reg. madr. an. 1803.) foliis inferioribus spathulatis pubescentibus, floribus spicatis, calycibus ovatis glabris, petalis bipartitis acutis. (1) in Hispaniâ.
94. *S. SECUNDIFLORA* (Oth mss.) glaherrima, foliis lanceolatis ciliatis, floribus secundis pedunculatis, calycibus floriferis cylindricis, frutiferis clavatis, petalis bipartitis. (1) Hab. ... *S. glauca* Pourr. elench. hort. reg. madr. 1803. Lag. gen. et spec. 15. non Zea.
95. *S. VISCOSISSIMA* (Ten. prodr. fl. nap. xxvi.) villosa-viscosa, pilis articulatis, caulibus erectis lævibus, foliis carnosis, radicalibus obovato-spathulatis, caulinis lanceolato-oblongis obtusis canaliculatis, floribus spicato-racemosis, petalis bipartitis. (1) prope Neapolim. An hujus sect. ?

Sect. VI. RUPIFRAGA. *Oth mss.*

Caulescentes. Canles stricti. Pedunculi filiformes. Calyces campanulati vel cylindrici.

96. *S. QUADRIDENTATA* (DC! fl. fr. 4. p. 748.) cespitosa, caulibus erectis gracilibus ramosis, foliis parvis linearibus angustissimis, floribus minimis paniculatis, calycibus campanulato-clavatis, petalis breviter quadridentatis. (1) \mathcal{L} in Alpibus. Cucub. quadrifidus Linn. spec. 414. *S. fontana* Ten? (v. v.)
- β. pusilla* (Oth mss.) ferè acaulis, foliis latioribus radicalibus, pedunculis longis erectis 1 raro 2-floris. \mathcal{L} in Alpibus Hungariæ. *S. pusilla* Wulst. et Kit. pl. hung. p. 235. t. 212.
97. *S. ALPESTRIS* (Jacq. fl. austr. 1. p. 60. t. 96.) radice ramosâ, caulibus simplicibus subfoliosis, foliis lanceolato-linearibus obtusiusculis, floribus majoribus, paniculatis calycibus campanulato-clavatis, petalis lamina lata 4-fida, appendice bipartito, seminibus ciliatis. \mathcal{L} in Austriâ.
98. *S. GRATA* (Henk. adumbr. plant. 28.) glabra, caulibus filiformibus diffusis ramosis, foliis retroflexis carnoso-canaliculatis mucronatis, floribus terminalibus, calycibus ventricoso-clavatis, petalis bipartitis reticulatis. \mathcal{L} ? Poir. dict. suppl. 5. p. 155.
99. *S. RUPESTRIS* (Lin. spec. 602.) glabra, caulibus erectis ramosis, foliis ovato-lanceolatis, floribus paniculatis minimis, calycibus campanulato-clavatis, petalis obcordatis. \mathcal{L} in Alpibus. Sturm deutschl. fl. 1. fasc. 22. t. 10. (v. v.)
100. *S. GLAUCIFOLIA* (Lag! in varied de cienc. 1805. p. 213.) caule procumbente ramoso folioso, foliis ovatis obtusis, inferioribus petiolatis, superioribus ferè subcordatis, floribus parvis paniculatis, calycibus clavatis, petalis obovatis. \mathcal{L} in Hispaniæ montibus Legionensibus. Aff. *S. rupestri* sed Anthophoro longiore. (v. s.)
101. *S. LYCENIDIFLORA* (Oth mss.) viscosa, caule erecto ramoso basi pubes-

- cente, foliis inferioribus lanceolato-spathulatis pubescentibus, superioribus lineari-lanceolatis glabris, floribus laxè paniculatis, calycibus subovatis, petalis lamina lata obcordata. ① in Cretâ? (v. s.)
102. *S. INAPERTA* (Lin. spec. 600.) glabra, caulibus erectis ramosis, foliis lanceolatis acutis, inferioribus obtusiusculis, floribus laxè paniculatis, calycibus clavatis, petalis inapertis obcordatis, staminibus sæpè 5. ① in Enropâ austr. Viscago lævis, etc. Dill. elth. t. 314. f. 407. *S. polyphylla*. DC. fl. fr. 4. p. 750. (v. s.)
103. *S. CLANDESTINA* (Jacq. coll. suppl. 111. t. 3. f. 3.) pubescens, caule erecto ramosissimo gracili, foliis inferioribus oblongis obtusis, superioribus lanceolatis subangustis, floribus laxè paniculatis, calycibus ventricosis, petalis brevibus erectis bifidis. ① ② in promont. Bon.-Spei. (v. v.)
β. angustifolia (Otth mss.) foliis linearibus, inferioribus petiolatis. ① in Lusitaniâ. *S. Portensis* Linn. spec. 704.
104. *S. ANTIRRHINA* (Linn. spec. 600.) ferè glabra, caule erecto ramoso subflososo, foliis lanceolatis acutis subciliatis, floribus parvis paniculatis, calycibus ovatis glabris, petalis obcordatis coronatis. ① in Virginiâ. Viscago amer. etc. Dill. hort. elth. p. 422. t. 313. (v. s.)
105. *S. GEMINIFLORA* (Willd. enum. 472.) pubescens, caulibus subramosis, foliis inferioribus elliptico-spathulatis, superioribus lanceolatis, floribus terminalibus geminatis, calycibus clavatis, petalis bifidis. ① Hab. Petala purpurea. (v. s.)
106. *S. FLAVESCENS* (Waldst. et Kit. pl. hung. 2. p. 131. t. 175.) pubescens, caulibus erectis ramosissimis strictis, foliis inferioribus lanceolato-spathulatis, superioribus linearibus, floribus laxè paniculatis, calycibus cylindricis, petalis bilobis. 2 in Hungariâ. *S. mollis* Horn. hort. hafn. 1. p. 418. Petala flavescentia. (v. s.)
107. *S. LINIFOLIA* (Willd. enum. 473.) caulibus ramosis, foliis lineari-spathulatis, floribus terminalibus, calycibus cylindraceo-clavatis, petalis bifidis. ① Hab. ... Petala viridi-flavescentia.
108. *S. CRETICA* (Linn. syst. veg. 421. n. 29.) glabra, caulibus erectis ramosis, foliis inferioribus ovatis petiolatis obtusis, superioribus linearibus acutis, floribus laxè paniculatis, calycibus ovato-clavatis, petalis bifidis. ① in Cretâ. Viscago, etc. Dill. hort. elth. 422. t. 314. f. 404.
109. *S. DIVARICATA* (Clem. elench. hort. reg. madr. an p. 1806. 105.) viscosa, caule erecto pubescente ramoso, foliis ciliatis inferioribus spathulatis obtusis, superioribus lanceolatis acutis, floribus terminalibus et in dichotomiâ, calycibus cylindrico-clavatis, petalis obovatis emarginatis coronatis. ① Hab. ... *S. Sicula* Cyrillo. Petala rosea. (v. v.)
110. *S. SEDOIDES* (Jacq. coll. suppl. p. 112. t. 14. f. 1.) viscoso-pubescentis, caulibus erectis ramosissimis, foliis subcarnosis inferioribus spathulatis, superioribus ovatis, floribus parvis terminalibus et in dichotomiâ, calycibus clavatis, petalis obcordatis coronatis. ① in Cretâ. *S. ramosissima* Sibth. et Smith fl. græc. t. 425? (v. s.)
111. *S. COSTATA* (Otth mss.) pubescens, caulibus ramosissimis erectis, foliis parvis linearibus angustissimis, floribus terminalibus solitariis rarè geminis, calycibus clavatis scabris 10-costatis, petalis semi-bifidis. 2 Hab. ... (v. s.)
112. *S. FILIFORMIS* (Otth. mss.) scabra, caulibus filiformibus ramosissimis, foliis minimis filiformibus, floribus parvis numerosis terminalibus, calycibus cylindrico-clavatis. Patr. ign. (v. s. in h. Otth.)
113. *S. PARVIFOLIA* (Otth mss.) pubescens, caulibus repentibus ramosis, foliis parvis lineari-lanceolatis acutis, floribus terminalibus et in dichotomiâ, calycibus clavatis pilosis, petalis bifidis. 2 Hab. ... (v. s.)
114. *S. FRUTICULOSA* (Siebl. pl. exsicc.) vix pubescens, radice lignosâ ramosâ

- simâ, caulibus simplicibus diffusis filiformibus, foliis parvis lanceolato-spathulatis, floribus terminalibus longè pedunculatis, calycibus clavatis. $\frac{1}{2}$ in Cretâ. (v. s. in h. Otth.)
115. *S. SAXIFRAGA* (Linn. spec. 602.) caespitosa, caulibus assurgentibus, foliis linearibus acutis, pedunculis longissimis, floribus terminalibus solitariis rarò axillaribus, calycibus longè clavatis, petalis bipartitis, unguiculo ciliato, appendice bicuspidatâ. $\frac{1}{2}$ in Alpibus. Waldst. et Kit. pl. hung. 2. t. 163. (v. s.)
116. *S. PETRÆA* (Waldst. et Kit. pl. hung. 2. t. 164.) caespitosa, breviter setosa, caulibus assurgentibus, foliis linearibus setaceo-denticulatis, floribus minoribus terminalibus solitariis, calycibus clavatis, petalis bifidis, appendice bifidâ. $\frac{1}{2}$ in Hungariâ. (v. s.)
117. *S. CAMPANULA* (Pers. ench. 1. p. 500.) glabra, caulibus erectis seu assurgentibus subramosis basi foliosis, foliis lanceolato-linearibus acutis imis spathulatis, floribus terminalibus solitariis seu geminis, pedunculis longissimis, calycibus campanulatis amplioribus, petalis 2-partitis nudis. $\frac{1}{2}$ in rupibus Pedemontanis. Cucub. alpestris All. auct. p. 28. t. 1. f. 3. (v. s.)
118. *S. VISCIDA* (Spreng. pl. min. cog. 2. p. 65.) viscidissima, pilis glanduliferis, caulibus diffusis ramosis, foliis lineari-lanceolatis, floribus solitariis terminalibus aut in axillis foliorum, petalis dentatis. $\frac{1}{2}$? in Carniolâ.
119. *S. URVILLII* (Schott mon. sil. ined.) rigida, ferè glabra, caule basi suffruticoso ramoso foliosissimo, foliis horizontalibus linearibus pungentibus, floribus terminalibus rarò axillaribus, calycibus clavatis, petalis semibifidis. $\frac{1}{2}$ in insulâ Cos. (v. s. miss. à cl. D'Urville.)
- 120? *S. NIVEA* (Mühlenb. catal.) caule supernè ramoso, foliis oblongo-lanceolatis pulverulento-pubescentibus, floribus solitariis, calycibus campanulato-inflatis subpilosis, petalis parvis reflexis bifidis longè unguiculatis. — in vicino Columbiae Nutt. gen. amer. 1. p. 287.

Sect. VII. SIPHONOMORPHA. *Oth mss.*

Canescentes. Flores paniculati rarò solitarii; pedicelli oppositi breves. Calyx tubulatus.

§. 1. *Floribus nutantibus, calycibus cylindricis.*

121. *S. LONGIPETALA* (Vent. hort. cels. p. 83. t. 83.) glabra viscosa, caulibus erectis, foliis subcarnosis lineari-lanceolatis setaceo-ciliatis, floribus nutantibus laxè paniculatis, calycibus vesiculato-clavatis, petalis laminâ bipartitâ reflexâ longissimâ. ① circa Aleppum.
122. *S. LONGICILIA* (Otth mss.) radice lignosâ, caulibus basi pubescentibus ramosis, foliis lanceolatis acutis, inferioribus petiolatis pubescentibus lanuginoso-ciliatis, floribus nutantibus paniculatis, calycibus clavatis, petalis bipartitis tuberculis 2 coronatis. $\frac{1}{2}$ in Lusitaniâ. Cucub. longicilius Broter. fl. lus. 2. p. 180.*
123. *S. NUTANS* (Linn. spec. 596.) pubescens, caulibus basi foliosissimis, foliis radicalibus spathulatis, superioribus lanceolato-linearibus, floribus paniculatis secundis pendulis, calycibus cylindrico-ventricosus, petalis bipartitis reflexis longè coronatis. $\frac{1}{2}$ in Europæ pratis. *S. infracta* Waldst. et Kit. pl. hung. 3. t. 213. *S. latifolia* Horn. h. hafa. suppl. 49.
- β. *viridella* (Otth mss.) pubescens ramosissima, petalis viridibus aut viridi-pubescentibus. (v. v.)
- γ. *incana* (Ser. herb.) caule foliis calycibusque incano-subtomentosis, calycibus pallidis vel purpurascens — In Vallesiâ. (comm. ex herb. Venetz.)
- δ. *glabra* (DC! fl. fr. 5. p. 605.) glabra minùs ramosa minùsque foliosa, foliis linearibus, petalis sæpè viridibus. — in Sibirîâ et circa Spa. *S. saxatilis* Sims bot. mag. 689. *S. Amblevana* Lejeun fl. spa. 1. p. 199. (v. s.)
- ε. *oligophylla* (Otth mss.) pubescens humilis, caule simplici ferè nudo

foliis lanceolatis obtusiusculis, floribus paniculatis, calycibus cylindrico-ventricosus, striis alternis venosis, petalis bipartitis, lobis latis denticulatis. — ad cap. B-Spei. Sims bot. mag. t. 382. (v. s.)

149. *S. CORSICA* (DC! fl. fr. 4. p. 756.) pubescens viscosissima, caulibus procumbentibus foliosis, foliis parvis obovatis, floribus terminalibus erectis, calycibus clavatis, petalis longissimè unguiculatis bipartitis coronatis. ♀ in Corsicâ. — Bocc. mus. t. 54. (v. s.)

150. *S. PUBESCENS* (Loisel. fl. gall. p. 727.) caule humili ramosissimo, foliis lanceolatis obtusis basi attenuatis ciliatis, floribus terminalibus erectis, calycibus clavatis hirsutis, petalis longissimè unguiculatis bipartitis coronatis. ①? in Corsicâ circa Ajaccio. S. hirta Willd. h. berol. t. 23. An *S. sericea*? (Ser.) (v. s.)

151. *S. ÆGYPTIACA* (Lin. f. suppl. 241.) subtomentosa, caulibus ramosis, foliis obovatis petiolatis, floribus terminalibus erectis, calycibus clavatis, petalis laminâ obcordatâ basi bidentatâ. ① in Ægypto.

♀. *retroflexa* (Pers. ench. 1. p. 500.) foliis linearibus, petalis retroflexis.

152. *S. SERICEA* (All. ped. n. 1573. t. 79. f. 3.) sericea, caulibus ramosis, foliis basi longè ciliatis inferioribus spathulatis, superioribus oblongis, floribus magnis terminalibus seu paucifloro-paniculatis, calycibus clavatis, petalis longè unguiculatis bipartitis coronatis. ① in Corsicâ. Petala rosea. (v. s.)

♀. *minor* (Otth mss.) foliis opacis subcarnosis, floribus habituque minoribus. ① ad Tagum. *S. littorea* Brot. fl. lus. 2. p. 186.*

153. *S. PICTA* (Pers. ench. 1. p. 498.) caulibus ramosissimis vix pubescentibus, foliis inferioribus obovato-spathulatis, superioribus linearibus acutis, floribus laxè paniculatis, calycibus clavatis rubro-striatis bipartitis reticulatis. ① Hab. . . *S. Reinwardtii* Roth. *S. reticulata* hort. *S. anastomosans* Lag. gen. et spec. 15. (v. v.)

154. *S. RUGOSA* (Pers. ench. 1. p. 498.) glabra, caule ramoso, foliis linearilanceolatis acutis, floribus terminalibus vel subpaniculatis, calycibus clavatis, striis crenato-rugosis subasperis. — Hab. . .

155. *S. NICOLOR* (Thori land. p. 174.) cespitosa glabra subviscosa, caulibus basi ramosis, foliis linearibus, floribus paniculatis, calycibus longè clavatis reticulatis, petalis bifidis, lobis lanceolatis. ① in Galliâ occidentali. DC. ic. pl. rar. gall. t. 42. *S. Portensis* Bonam. prodr. (v. s.)

156. *S. KAULFUSSII* (Spreng. pl. min. cog. 2. p. 64. n. 123.) glaberrima, radice fusiformi, caule subsimplici, foliis radicalibus oblongis, caulinis ovato-lanceolatis, floribus paniculatis, calycibus clavatis, petalis longè unguiculatis, laminâ dentatâ. ♀ ad Polnitz.

157. *S. LÆVIGATA* (Sibth. et Smith fl. græc. t. 418. ex prodr. 1. p. 295.) foliis elliptico-rotundatis, floribus laxè paniculatis, calycibus lævissimis aveniis, petalis semibifidis angustatis nudis. ① in Cypri montibus.

158. *S. RAMOSA* (Otth mss.) caule ramosissimo, foliis spathulatis recurvis, calycibus clavatis 10-angularibus, petalis bifidis, appendice 4-fidâ. ① in Cretâ. *S. ramosissima* Sibth. et Smith fl. græc. t. 425. ex prodr. 1. p. 297. non Desf.

§. 3. Floribus erectis, calycibus longè clavatis.

159. *S. RETICULATA* (Desf! fl. atl. 1. p. 350. t. 99.) glabra viscosa, caulibus ramosis, foliis lanceolato-linearibus, floribus paniculatis, calycibus longissimè clavatis reticulatis, petalis obcordatis basi utrinquè unidentatis. ①? in Algeriâ. (v. s.)

160. *S. ECHINATA* (Otth mss.) pubescens, caule gracili ramoso, foliis caulinis parvis linearibus, floribus paniculatis, calycibus longis cylindrico-clavatis, costis 10 setaceo-echinatis, petalis bifidis. ①? Hab. . . (v. s.)

161. *S. PENNSYLVANICA* (Michx. fl. bor. amer. 1. p. 272.) viscidè-pubescent,

caulibus procumbentibus, foliis linearibus longis, floribus paniculatis, calycibus longis tubulatis, petalis leviter emarginatis subcrenatis. — in Pensylvaniâ. Lodd. bot. cab. 1. t. 41.

162. S. SUCCULENTA (Forsk. descr. p. 89.) viscidè-pubescent, caulibus diffusis crassis ramosis foliosis, foliis obovatis carneis, floribus paniculatis, calycibus longis subventricosus costatis, petalis longissimè unguiculatis bipartitis coronatis. $\frac{1}{2}$? in Ægypto. Delil. fl. ægypt. t. 29. f. 2. (v. s. ex herb. Olivier.)
163. S. VALLESIA (Linn. spec. 603.) cespitosa viscidè-pubescent, radice lignosâ, caulibus humilibus assurgentibus parùm ramosis, foliis lanceolatis, inferioribus spathulatis, floribus terminalibus rarò geminis, calycibus longissimis reticulatis, petalis bifidis. $\frac{1}{2}$ in Alpibus. All. ped. n. 1574. t. 23. f. 2. (v. s.)
164. S. FRUTIGOSA (Linn. spec. 597.) suffruticosa, caulibus suberectis glabris basi ramosis, foliis lato-spathulatis nitidis ciliatis, floribus paniculatis, calycibus longissimis subcylindricis viscidè-villosis, petalis bilobis. $\frac{1}{2}$ in Sicilia. S. nitida Lagasc. gen. et spec. 15. (v. v.)
165. S. CASPICA (Pers. ench. 1. p. 497.) scabra, caulibus ramosis, foliis spathulato-lanceolatis, floribus terminalibus aut in dichotomiâ, calycibus longis cylindricis, petalis bipartitis basi utrinquè unidentatis. $\frac{1}{2}$ in Caucaso, etc. S. frutescent. Bieb. casp. 175. S. fruticulosa Dietr.
166. S. SALZMANNII (Oth mss.) tomentoso-pilosa, radice lignosâ ramosâ, caulibus simplicibus foliosis, foliis lanceolato-spathulatis basi numerosis, floribus densè paniculatis erectis, calycibus longis, petalis latis obcordatis nudis. $\frac{1}{2}$ in Corsicâ. (v. s.)
167. S. REQUIENII (Oth mss.) pubescens, radice crassâ lignosâ, caule erecto simplici folioso præcipuè basi, foliis oblongo-obovatis acuminatis, floribus paucis paniculatis, calycibus longissimis. $\frac{1}{2}$ in Corsicâ prope Bonifacio (v. s.)
168. S. AMOENA (Linn. spec. 596.) pubescens, radice lignosâ, caulibus diffusis ramosis, foliis mollibus infrâ numerosis lanceolato-spathulatis subacutis, floribus numerosis amplè-paniculatis, calycibus cylindrico-clavatis, petalis semi-bifidis. $\frac{1}{2}$ in Tataria. (v. s.)
169. S. SUPINA (Bieb. fl. taur. 1. p. 336.) cespitosa viscidè pubescens, caulibus lignosis supinis diffusis ramosis, foliis linearibus acutis, floribus paniculatis, calycibus longis cylindrico-clavatis, petalis longè unguiculatis bifidis, lobis augustis divergentibus. $\frac{1}{2}$ in Caucaso. Sims bot. mag. t. 1997. (v. v. et s.)
170. S. DEPRESSA (Bieb. fl. taur. 1. p. 336.) viscosa, radice lignosâ, caulibus numerosis basi ramosis foliosis pubescentibus, foliis lanceolatis parvis subciliatis, floribus terminalibus rarò geminis, calycibus longissimis cylindrico-clavatis, petalis longè unguiculatis bifidis. $\frac{1}{2}$ in Iberiâ. (v. s. comm. à cl. Fisch. et Stev.)
171. S. PARADOXA (Lin. spec. 1673.) caulibus erectis pubescentibus, foliis scabriusculis vix ciliatis, inferioribus lanceolato-spathulatis, superioribus linearibus, floribus magnis paniculatis, calycibus longis cylindrico-clavatis, petalis bilobis, lobis latis obovatis, appendice bipartitâ. $\frac{1}{2}$ in Delphinatu. Jacq. hort. vind. 3. t. 84. Cucubalus viscosus Huds. angl. 186. (v. s.)
- β . *tenuifolia* (Oth mss.) caule virgato simplici, foliis linearibus acutis — circa Genuam.
172. S. CHLORÆFOLIA (Smith ic. ined. 1. p. 13. t. 13.) glaberrima glauca, caulibus ramosis, foliis subrotundis acuminatis, superioribus subcordatis, floribus magnis terminalibus aut paniculatis, calycibus clavatis non striatis, petalis bifidis, lobis latis ovatis, appendice bicuspidato. $\frac{1}{2}$ in Armeniâ. Sims bot. mag. t. 807. (v. v.)
173. S. ITALICA (DC. fl. fr. 4. p. 753.) piloso-pubescent, caulibus ramosissimis, foliis inferioribus ovato-spathulatis, superioribus linearibus, floribus amplè-paniculatis, calycibus longis clavatis, petalis bilobis nudis. $\frac{1}{2}$ in Italiâ, etc.

Cucub. Italicus Lin. spec. 593. Cucub. silenoides Vill. Dauph. 3. p. 614. (v. v.)

ℓ. *cana* (Otth mss.) parum ramosa molliissima, panicula densa. Cucub. molliissimus Loiseleur not. 166. (v. s.)

γ. *rubriflora* (Otth mss.) molliissima, calycibus purpurascens, petalis lamina purpurea. (v. s.)

174. *S. PANICULATA* (Otth mss.) velutino-viscosa, caule erecto ramosissimo folioso, foliis lanceolatis subcrassis, floribus amplò-paniculatis, calycibus cylindrico-clavatis longis, petalis bifidis, genitalibus longis. 2ℓ? Hab... (v. s.)

175. *S. PAUCIFLORA* (Salzmann! exsic.) cespitosa pubescens, caulibus numerosis simplicibus gracilibus ferè aphyllis, foliis lineari-lanceolatis basi numerosis, floribus terminalibus vel paucifloro-paniculatis, calycibus angustis cylindricis, petalis bipartitis reticulatis. 2ℓ in Corsicà. (v. s.)

176. *S. PATULA* (Desf. fl. atl. 1. p. 356.) pubescens viscosa, caulibus erectis ramosis, ramis patentibus, foliis inferioribus ovato-spathulatis, superioribus lanceolatis, floribus paniculatis, calycibus longè clavatis angustis, petalis longè unguiculatis semi-bifidis coronatis. 2ℓ in Barbarià.

177. *S. POLYPHYLLA* (Lin. spec. 601.) pubescens, caulibus assurgentibus ramosissimis foliosissimis, foliis linearibus acutis, floribus erectis paniculatis, calycibus clavatis, petalis bifidis. 2ℓ in Austrià, etc. — Clus. hist. 1. p. 290. ic. (v. s.)

178. *S. NEMORALIS* (Waldst. et Kit. pl. hung. 3. p. 277. t. 249.) caule simplici pubescente, foliis pubescentibus, inferioribus magnis subrotundis petiolatis, superioribus lanceolatis, floribus paniculatis, calycibus longè clavatis, petalis bipartitis coronatis. ②? in Hungarià. Cucub. floccosus Fic. (v. s. comm. à cl. Jan.)

179. *S. LONGIFLORA* (Ehrh. beytr. 7. p. 144.) glabra, caule virgato, foliis lineari-lanceolatis, radicalibus longissimis, floribus paniculatis, pedunculis unifloris, calycibus longissimè clavatis, petalis bipartitis coronatis. 2ℓ in Hungarià et in Taurià. Waldst. et Kit. pl. hung. 1. p. 7. t. 8. (v. v.)

ℓ. *juncea* (Otth mss.) foliis omnibus linearibus, panicula parva. In Iberià. *S. juncea* Roth catalect. 1. p. 54. (v. s.)

180. *S. BUPLEUROIDES* (Lin. spec. 598.) glabra viscosa, caule assurgenti ramoso, foliis lineari-lanceolatis acutis, imis longissimis, bracteis superioribus margine lato membranaceo, floribus paniculatis, pedunculis subtrifloris, calycibus longius clavatis, petalis bilobis coronatis. 2ℓ in Persià, etc. (v. s.)

181. *S. CANA* (Otth mss.) pubescenti-canescens, caule virgato simplici, foliis inferioribus ovato-oblongis petiolatis basi ciliatis, superioribus linearibus, floribus paniculatis, pedunculis trifloris, calycibus longè clavatis purpurascens, petalis longè unguiculatis bifidis. 2ℓ Hab... (v. s.)

182. *S. MOLLISSIMA* (Pers. ench. 1. p. 498.) holosericea, caulibus erectis ramosis, foliis undulatis, inferioribus magnis spathulatis, superioribus lanceolatis, floribus amplò-paniculatis, calycibus clavato-cylindricis, petalis bipartitis coronatis. 2ℓ in Italià. Cucubalus molliissimus Lin. spec. 593. Waldst. et Kit. pl. hung. 3. t. 248. (v. s.)

183. *S. REGIA* (Sims bot. mag. t. 1724.) viscoso-pubescent, foliis lanceolatis, floribus magnis paniculatis, calycibus longè tubulatis, petalis lamina lanceolata, appendice bicuspidata, genitalibus longissimis. 2ℓ? in Armenià. *S. virginica* var. *illinoensis* Michx. fl. bor. am. 1. p. 272.

184. *S. BALDWINII* (Nutt. gen. amer. 1. p. 288.) pilosa, foliis sublanco-latis, floribus maximis trichotomis, petalis divaricatis laciniatis. — in Floridà. Petala rosea.

185. *S. MOENIANA* (DC. mss.) subvillosa, caule simplici erecto, foliis lanceolatis, floribus paniculatis, calycibus cylindrico-clavatis longissimis, petalis

sexfidis, appendicibus 2-bicuspidatis. \mathcal{Z} in Mexico. S. Moq. et Sess. pl. mex. icon. ined. Flores purpurei.

186. S. LACINIATA (Cav. icon. 6. p. 44. t. 564.) pubescens, caule erecto ramoso, foliis magnis lanceolatis acutis, floribus maximis terminalibus subumbellatis, calycibus longis cylindrico-ventricosis, petalis semi-4-fidis, appendicibus 2-ovatis, genitalibus inclusis. (I) in America meridionali.

187. S. ROTUNDIFOLIA (Nutt. gen. amer. 1. p. 288.) caulibus decumbentibus pilosis, foliis dilatato-ovalibus petiolatis ciliatis, floribus paucis paniculatis, calycibus cylindrico-clavatis, petalis laciniatis sub-4-fidis coronatis. — in Ohio et Tennessee. Petala coccinea.

188. S. ASCENDENS (Lag. gen. et spec. 15.) villosa-viscosa, foliis lineari-lanceolatis obtusis ciliatis, pedunculis unifloris fructiferis patentibus, calycibus oblongo-clavatis basi circulariter reflexis, petalis bifidis. (I) in Hispania.

189. S. DIANTHOIDES (Pers. ench. 1. p. 500.) foliis linearibus, floribus paniculatis, calycibus striatis inter 2 foliola involucri similia sessilibus, petalis bifidis. — in Oriente. Cucubalus saxifragus Schreb. dec. 9. t. 5.

Sect. VIII. ATOCION. Oth mss.

Caulscentes. Flores corymbosi. Calyx clavatus, 10-striatus.

190. S. CORDIFOLIA (All. ped. n. 1581. t. 23. f. 3.) viscosa, caulibus simplicibus pilosis, foliis ovatis cuspidatis pubescentibus, superioribus subcordatis, floribus 1-4 terminalibus fere sessilibus, calycibus clavatis ovato-cylindricis, petalis bipartitis coronatis. \mathcal{Z} in Pedemontii saxosis. (v. s.)

191. S. CESPITOSA (Stev! in mem. soc. mosq. 2. p. 262.) cespitosa scabriuscula, radice crassa lignosa ramosa, caulibus simplicibus gracilibus basi foliosisimis, foliis parvis linearibus acutis, floribus 2-3 terminalibus, calycibus clavatis, petalis bilobis. \mathcal{Z} in Caucaso. (v. s.)

192. S. TOMENTOSA (Oth mss.) mollissima, radice lignosa ramosa, caulibus simplicibus, basi foliosisimis supra fere aphyllis, foliis carnosis, inferioribus magnis cochleariformibus, superioribus linearibus, floribus in corymbum confertis, calycibus cylindrico-clavatis, petalis bipartitis. \mathcal{Z} in Gibraltar. (v. s.)

193. S. PLATYPETALA (Oth. mss.) piloso-pubescent, caule humili basi foliosisimo, foliis basi ciliatis, inferioribus lanceolato-spathulatis, superioribus lineari-lanceolatis, floribus paniculato-corymbosis paucis, calycibus clavatis, petalis longe unguiculatis, laminâ latâ obovatâ. \mathcal{Z} in America sept. S. Virginica Rafin! in litt. non Linn. (v. s.)

194. S. ATOCION (Murr. syst. ed. 13. p. 421.) viscosa, caule ramosissimo pubescente foliis rotundis subciliatis inferioribus longe petiolatis, floribus laxè corymbosis, calycibus longe clavatis, petalis laminâ latâ obcordatâ basi utrinque unidentatâ coronatâ. (I) in Oriente. S. orchidea. Lin. f. suppl. 241. (v. s.)

195. S. PSEUDO-ATOCION (Desf. fl. atl. 1. p. 353.) viscosa, caule ramosissimo, foliis glabris, inferioribus obovato-spathulatis, floribus laxè corymbosis, calycibus longe clavatis, petalis laminâ integerrimâ lineari coronatâ. (I) in Atlante.

196. S. POLYGONOIDES (Pers. ench. 1. p. 500.) pilosa, radice crassâ, caulibus procumbentibus ramosis, foliis parvis, inferioribus lanceolatis, superioribus ellipticis acutis, floribus corymbosis sessilibus, calycibus cylindricis, petalis emarginatis. \mathcal{Z} in insulâ Naxo. Cucub. polygonoides Willd. spec. 2. p. 690.

197. S. CRISPA (Poir. dict. 7. p. 162.) radice gracili, caule erecto subsimplici hirtio, foliis oblongis linearibus obtusis margine crispato-spinulosi, floribus solitariis densis fere fasciculatis, calycibus longe clavatis, petalis bifidis. (I) in Barbariâ.

198. S. ARMERIA (Lin. spec. 601.) glaberrima glauca viscosa, caule ramoso, foliis ovato-lanceolatis, floribus paniculato-corymbosis, calycibus longe cla-

vatis, petalis obcordatis coronatis. ① in Angliâ, Galliâ, etc. Smith engl. bot. t. 1398. Cucub. fasciculatus Lam. Calyx et petala purpurea.

℘. *alba*, calycibus et floribus albis. (v. v.)

199. S. SIEGERI (Baumg. stirp. transylv. 1. p. 400.) caule erecto simplici sulcato, foliis radicalibus numerosis oblongo-lanceolatis canaliculato-petiolatis, caulinis 2 linearibus, floribus terminalibus solitariis aut 2-3 umbellatis, calycibus clavato-campanulatis, petalis obcordatis. — in Alpibus Rodnensis.

200. S. CONGESTA (Sibth. et Smith prod. fl. græc. 1. p. 300.) pubescens, caule subramoso, foliis obovatis acuminatis petiolatis, floribus corymbosis congestis, calycibus longissimè clavatis, petalis bipartitis nudis. ☉ in Pyrenæis et Græciâ.

201. S. COMPACTA (Fischl in Horn. hort. hafn. 1. p. 417.) glabra, caulibus erectis ramosis, foliis superioribus lanceolatis inferioribus lanceolato-linearibus, floribus in umbellam densam congestis, calycibus longissimè clavatis, petalis obovatis. ① in Russiâ. S. armeria Bieb. fl. taur. n. 837. (v. s.)

202. S. COESIA (Sibth. et Smith fl. græc. t. 417. ex prodr. 1. p. 294.) radice longâ, caulibus erectis ramosis, foliis obovato-rotundatis, floribus erectis corymboso-paniculatis, calyce vix reticulato, petalis bipartitis, lobis linearibus. ℥ in Parnasso.

203. S. PERFOLIATA (Otth mss.) glaberrima, caulibus gracilibus filiformibus subsimplicibus, foliis rotundato-perfoliatis, floribus capitato-corymbosis, involucri monophyllo foliis majore concavo, calycibus subinflatis oblongis, petalis parvis. — in Oriente. Cucub. chloræfolius Poir. dict. 2. p. 416. An Silene?

+ *Species non satis notæ.*

204. S. RIGIDULA (Sibth. et Smith fl. græc. t. 480, ex prod. 1. p. 299.) caule alternè ramoso patente, foliis lanceolatis glabris, petalis bipartitis acutis, appendicibus 4-dentatis. ① prope Athenas.

205. S. SPINESCENS (Sibth. et Smith fl. græc. t. 431, ex prodr. 1. p. 299.) caule fruticuloso, ramis oppositis horizontalibus spinescentibus foliis spatulatis undique pubescentibus, petalis bipartitis. ℥ in Asiâ minore.

206. S. LINOÏDES (Otth mss.) caulibus supernè paniculato-ramosis, foliis lineari-lanceolatis scabris, petalis bipartitis. ℥ in Parnasso. S. linifolia Sibth. et Smith fl. græc. t. 433. ex prodr. 1. p. 301. non Willd.

207. S. LEUCOPHÆA (Sibth. et Smith fl. græc. t. 424, ex prodr. 1. p. 297.) foliis lineari-oblongis recurvis glutinosis, calycibus decangularibus, petalis angustatis, appendice bifidâ. ☉ in Cypro.

208. S. JUNCEA (Sibth. et Smith fl. græc. t. 421. ex prodr. 1. p. 295.) foliis spatulatis undiquè scabris, floribus erectis, petalis bipartitis angustatis, appendicibus tridentatis. ① in Asiâ minore.

209. S. RACEMOSA (Otth mss.) caule dichotomo divaricato racemoso, foliis lanceolatis angustis, petalis bipartitis rotundatis-appendiculatis. ② in agro Cariense. S. divaricata Sibth. et Smith fl. græc. t. 414, ex prodr. 1. p. 293. non Clem.

210. S. THYMIFOLIA (Sibth. et Smith fl. græc. t. 411. ex prodr. 1. p. 292.) caulibus procumbentibus lignosis ramosissimis, calycibus hirtis glutinosis, petalis bipartitis angustatis, appendicibus emarginatis. ℥ in Cypro.

211. S. AURICULATA (Sibth. fl. græc. t. 435. ex prodr. 1. p. 301.) caulibus unifloris, foliis lanceolatis fimbriatis, calycibus campanulatis pubescentibus, petalis semibifidis utrinquè auriculatis. ℥ in monte Delphi Eubææ.

212. S. FALCATA (Sibth. et Smith fl. græc. t. 436. ex prodr. 1. p. 301.) cau-

- libus unifloris, foliis subulatis falcatis pilosis, calycibus clavatis, petalis semibifidis, unguiculis cuneiformibus. \mathcal{L} in monte Olympe Bithyno.
213. *S. LATIFOLIA* (Poir. voy. barbar. 2. p. 165.) caulibus gracilibus fistulosis ramosis, foliis ovatis acuminatis lævibus, superioribus et bracteis ciliatis, floribus axillaribus solitariis aut 2-3 terminalibus, calycibus subclavatis, petalis bifidis. \mathcal{L} ? in Barbariâ.
214. *S. NISCOLOR* (Sibth. et Smith fl. græc. t. 410, ex prodr. 1. p. 292.) caulibus diffusis raro ramosis foliis obovatis, floribus racemosis, calycibus villosis, petalis bipartitis, lobis linearibus, appendicibus emarginatis. (I) in Cypro.
215. *S. INCLUSA* (Horn. h. hafn. 1. p. 413.) hirsuta, floribus pedunculatis aut sessilibus, calycibus setoso-pilosis, pedunculo communi flexuoso, fructibus erectis, petalis emarginatis. (I)
216. *S. PINGUIS* (Vahl ex Horn. hort. hafn. suppl. 49.) sericeo-pilosiuscula, foliis carnosius, radicalibus spathulatis, superioribus obovato-lanceolatis, calycibus erectis clavatis, petalis bifidis. (I) ad cap. Sparte.
217. *S. CERNUA* (Thunb. fl. cap. ed. Schult. 1. p. 394.) hirsuta, floribus racemosis secundis cernuis, calyce 10-striato fructifero clavato, foliis linearibus villosis scabris. — ad cap. B.-Spei. An ad sectionem Siphonomorpharum referenda?

VII. LYCHNIS. DC. fl. fr. 4. p. 761. — *Lychnis* et *Agrostemma* Lin.

Calyx tubulosus, 5-dentatus, nudus. Petala 5 unguiculata, fauce sæpius coronatâ Stamina 10. Styli 5. Capsula 1-5-locularis, anthophoro longo vel nullo.

Sect. I. VISCARIA. DC. fl. fr. 4. p. 761.

Calyx cylindrico-clavatus; capsula semiquinque-locularis; anthophorum longum.

1. *L. VISCARIA* (Lin. spec. 625.) caule subgeniculis viscoso, limbo petalorum subintegro, foliis linearibus subspathulatis. \mathcal{L} in pratis siccis Europæ. Smith engl. bot. t. 788. — Clus. hist. 1. p. 289. f. 2. *Viscaria vulgaris* Roelh. ex Steud. nom. (v. v.)

*. Flore pleno. (v. v. in hort. cult.)

Sect. II. EULYCHNIS. DC. fl. fr. 4. p. 761.

Calyx cylindrico-clavatus; capsula unilocularis; petala fauce appendiculata; anthophorum longum aut in quibusdam subabbreviatum.

2. *L. CHALCEDONICA* (Lin. spec. 625.) glabriuscula, floribus fasciculatis, calycibus cylindrico-clavatis costatis, petalis bilobis, anthophoro longo, foliis lanceolatis cordatis subpilosis amplexicaulibus. \mathcal{L} circa Barnaoul et in Japoniâ; in hortis culta sub nom. *Croix de Jérusalem*. — Clus. hist. 1. p. 292. f. 1. Variat fl. coccineo, roseo, albo, simplici et pleno. (v. v.)
3. *L. FLOS-JOVIS* (Lin. spec. 625.) tomentosa, floribus umbellato-capitatis, calycibus cylindrico-clavatis costatis, petalis bilobis, foliis lanceolato-linearibus sericeo-tomentosis, anthophoro brevi crasso, foliis lanceolato-linearibus amplexicaulibus sericeo-tomentosis. \mathcal{L} in saxosis apricis Vallisæ, Pedemontii. — Moris. hist. 2. p. 540. s. 5. t. 36. f. 2. Pedunculi breves subramosi.

β . *ramosa* (Ser. mss.) floribus paniculato-corymbosis. — in monte Bresson Sabaudie. (v. s. et v. in h. Ser.)

4. *L. GRANDIFLORA* (Jacq. coll. 1. p. 149, pl. rar. t. 84.) glabra, floribus terminalibus axillaribusque 1-3, calycibus tereti-clavatis costatis, petalis laciniato-dentatis, anthophoro longissimo, foliis ovatis subsessilibus. \mathcal{L} in Chinâ, Japoniâ. *L. coronata* Thunb. fl. jap. 187. Delaun. herb. amat. t. 25. Flores coccinei, pulchri. (v. v.)

ℓ. *tetrapetala* (Ser. mss.) calyce quadridentato; petala 4. (v. s.)

5. *L. FULGENS* (Fisch. in Sims bot. mag. t. 2104.) pilosa, floribus 2-3 fastigiatibus, calycibus tereti-clavatis lanatis, petalis quadrifidis, laciniis exterioribus subulatis, anthophoro brevi, foliis ovatis hispido-tomentosis. 2ℓ in Sibiria. Sims. bot. mag. t. 478.* Reichenb. ic. t. 5. Corolla magnitudine *L. grandifloræ* sed colore *L. Chalcodoniceæ*. (v. s. comm. à cl. Fisch.)

6. *L. CœLI-ROSA* (Desrouss. in Lam. dict. 3. p. 644.) glabra, caule dichotomo paniculato erecto, floribus solitariis terminalibus, calycibus clavatis decemcostatis, lobis acutissimis, petalis lobatis, foliis linearibus acutis, anthophoro longo. ① in Sicilia, Barbaria et Oriente. *Agrostema cœli-rosa* Lin. spec. 624: — Moris. s. 5. t. 2. f. 32. (v. v.)

α. *pusilla* (Poir. suppl. 3. p. 537.) caule basi ramosissimo, foliis linearibus acutis subglabris. — in littore Hispaniæ.

β. *lævis* (Poir. l. c.) caule angulato erecto, foliis lævibus.

γ. *aspera* (Poir. l. c.) caule subtereti, foliis angustis asperis, angulis calycinis serrulato-scabris.

δ. *diffusa* (DC! cat. h. monsp. p. 122.) caule flexuoso decumbente.

Sect. III. AGROSTEMA. DC. fl. fr. 4. p. 763.

Calyx ovoideus, dentibus brevibus; capsula 1-locularis (5-locularis?); anthophorum brevissimum vel nullum.

7. *L. APETALA* (Lin. spec. 626.) caule stricto, uniflora, flore nutante, calyce inflato vesiculoso striato hirsuto, petalis calyce brevioribus, anthophoro brevissimo, foliis linearibus, inferioribus spathulatis. 2ℓ in Alpibus Lapponicis, Sibiricis. Lin. fl. lap. t. 12. f. 1. (v. s.)

ℓ. *pauciflora*, caule 1-2-vel 3 floro, petalis calyce vix longioribus. — *L. uniflora* et *pauciflora* Fisch. in litt. *L. brachypetala* cat. hort. berol. ex Horn. hort. hafn. suppl. 51.

8. *L. MAGELLANICA* (Lam. dict. 3. p. 641.*) subvillosa, petalis apice bifidis calyce campanulato vix longioribus, foliis linearibus gramineis. 2ℓ? ad fretum Magellanicum.

9. *L. VARIEGATA* (Desf. cor. tourn. 74. t. 56.*) glabra, floribus terminalibus, calycibus inflatis, petalis emarginatis variegatis, genitalibus exsertis, foliis rotundatis carnosis glaucis, petiolis basi connatis. 2ℓ in monte Idæ.

10. *L. SYLVESTRIS* (Hop! cent. exs. 3. n. 33. DC! fl. fr. 4. p. 763.) floribus (rubellis) dichotomo-paniculatis subdioicis, petalis semibifidis, lobis angustis divergentibus, capsulis subrotundis, valvis recurvis, foliis ovatis vel lanceolatis. 2ℓ in umbrosis humidis Europæ. *L. diurna* Sibth. oxon. 145 ex Hofm. fl. germ. 1800. p. 211. *L. dioica* α. Lin. spec. 626. *L. dioica* fl. rubro. Smith. engl. bot. t. 1579. Flores inodori. subsessiles. Variat fl. purpureis vel roseis plenis. (v. v.)

β. *pygmæa* (Ser. mss.) parvula, foliis subimbricatis, caulibus 1-2-floris. in monte Margazolo.

11. *L. DIOICA* (Lin. spec. 626. var. β, DC! fl. fr. 4. p. 762.) floribus (albis) dichotomo-paniculatis dioicis, petalis semibifidis, lobis latis approximatis, capsulis conicis, apicibus rectis, foliis ovatis. ① ad vias et sepes Europæ. *L. vespertina* Sibth. oxon. 146. Hoffm. fl. germ. 1800. p. 211. *L. alba*. Mill. dict. n. 4. *L. dioica* fl. albo Smith engl. bot. t. 1580. Flores odorati pedunculati. (v. v.)

* *multiplex* floribus albis plenis. 2ℓ in hortis rarè culta.

β. *viridiflora* (Ser. mss.) flore pleno, petalis viridibus. (v. s.)

12. *L. DICLINA* (Lag. gen. et spec. 15.*) pilosa, caulibus cespitosus diffusis dichotomis foliosis, floribus dioicis longè pedicellatis terminalibus axillaribusque, calycibus teretibus striatis, fructiferis subglobosis, dentibus parvis, petalis emarginatis, genitalibus inclusis, foliis bracteisque ovato-lanceolatis,

anthophoro nullo. $\frac{1}{2}$ in regno Valentino. *Agrostema dioica*. L. Duf. in litt. (v. s.)

13. L. *LÆTA* (Ait. hort. kew. 2. p. 118. ed. 2. v. 3. p. 134.) floribus solitariis, calycibus decemcarinatis, petalis bifidis, foliis lineari-lanceolatis subciliatis. (I) in Lusitaniâ. L. palustris Brot. fl. lus. 2. p. 221. phyt. lus. fasc. 1. ex fl. lus.

14. L. *CORSICA* (Lois. not. 73.) caule erectiusculo ramoso subdichotomo, pedunculis elongatis unifloris, petalis oblongis subemarginatis, foliis lineari-lanceolatis glabris. $\frac{1}{2}$ in Corsicâ. An diversa à L. lætâ? (v. s. comm. à cl. Robert.)

15. L. *SIBIRICA* (Lin. spec. 626.) caulibus cespitosis, floribus dichotomo-fasciculatis solitariisque in dichotomiâ longè pedunculatis, calycibus campanulato-globosis, lobis brevissimis obtusis, petalis bifidis, anthophoro nullo, foliis linearibus caulibusque subhirsutis. $\frac{1}{2}$ in Sibiria. Statura *Silenes rupestris*. (v. s. ex h. Patr. et comm. à cl. Fisch.)

16. L. *PYRÆNICA* (Berg. fl. bass. pyren. 2. p. 264.) glabra, caulibus cespitosis diffusis, floribus dichotomo-fasciculatis, flore longè pedunculato in dichotomiâ, calycibus campanulatis, lobis brevibus, petalis subemarginatis appendiculatis, foliis coriaceis, radicalibus spatulatis longè petiolatis, caulinis cordatis sessilibus. $\frac{1}{2}$ ad rupes in Pyrenæis occid. DC! fl. fr. 5. p. 608. ic. gall. rar. fasc. 2. ined. L. nummularia Lapeyr. abr. p. 263. (v. s.)

17. L. *FLOS-CUCULI* (Lin. spec. 625.) caulibus ascendentibus glabresculis, floribus dichotomo-fasciculatis, calycibus campanulatis decemcostatis, petalis laciniatis appendiculatis, foliis lanceolato-linearibus. $\frac{1}{2}$ in pratis et humidis Europæ. Lam. ill. t. 391. Smilh engl. bot. t. 573. Variat flore albo et pleno. (v. v.)

18. L. *ALPINA* (Lin. spec. 626.) glabra, caulibus cespitosis strictis, floribus densè umbellato-capitatis, calycibus campanulatis, petalis bifidis, anthophoro nullo, fructibus globosis, foliis lineari-lanceolatis. $\frac{1}{2}$ in pratis editiorum Alpium et Pyrenæorum. Hall. helv. n. 922. t. 17. f. 4. (v. v. et s.)

19. L. *CORONARIA* (Lam. dict. 3. p. 643.) tomentosa, caule dichotomo, pedunculis elongatis 1-floris, calycibus subcampanulatis costatis, petalis emarginatis coronatis serratis, foliis lanceolatis latissimis coriaceis. $\frac{1}{2}$ ad margines sylvarum in Alpibus et in Tauria. Curt. bot. mag. t. 24. *Agrostema coronaria* Lin. spec. 625. Variat fl. rubro, albo et pleno. (v. v.)

Sect. IV. *GITHAGO*. Desf. cat. 159.

Calyx cylindrico-campanulatus coriaceus, laciniis longissimis; capsulæ 1-loculares; anthophorum nullum.

20. L. *GITHAGO* (Lam. dict. 3. p. 643. DC! fl. fr. 4. p. 764.) hirsuta, caule dichotomo, floribus longè pedunculatis, foliis linearibus. (I) inter segetes Europæ. *Agrostema Githago* Lin. spec. 624. *Githago segetum*. Desf. l. c. — Fuchs hist. 127. ic. (v. v.)

β. *Nicæensis* (Willd. sp. 2. p. 805.) floribus pallidis, laciniis calycinis longioribus.

† *Species non satis nota.*

21. L. *LUSITANICA* (Mill. dict. n. 8.) caule erecto, calycibus striatis acutis, petalis dissectis. — in Lusitaniâ. †

VIII. *VELEZIA*. Lin. gen. n. 448. Gaertn. fruct. 2. t. 129. f. 12.

Calyx tubulosus, 5-6-dentatus. Petala 5-6 breviter, unguibus filiformibus barbatis, limbo emarginato. Stamina 5-6. Styli 2. Capsula 1-locularis.

1. V. *RIGIDA* (Lin. spec. 474.) calycibus filiformibus pubescentibus, petalis

bifidis. ① in Galliâ austr. et in Cretâ. — Bocc. mus. 2. p. 50. t. 45. Flores decandri ex Sibth. et Smith prod. 1. p. 283. Corolla parva, limbo roseo. (v. s.)

2. V. QUADRIDENTATA (Sibth. et Smith fl. græc. t. 391. ex prod. 1. p. 283.) calycibus clavatis glabris, petalis quadridentatis. ① in Asiâ minori. V. clavata D'Urville enum. pl. archip. 28. n. 234?

IX. DRYPIS. *Lin. gen. n. 519. Gaertn. fruct. 2. t. 28. f. 12.*

Calyx tubulosus 5-dentatus. Petala 5 unguiculata, fauce bidentatâ, limbo bipartito. Stamina 5. Styli 3. Capsula 1-locularis circumcissa abortu monosperma.

1. D. SPINOSA (Lin. spec. 390.) ② in Istriâ, Italiâ, Mauritaniâ. Jacq. h. vind. t. 49. Rœm. et Schult. syst. 6. p. 680. — Moris. hist. 3. p. 161. s. 7. t. 32. f. 8. Folia subulata, bractææ utrinquè tridentatæ.

Trib. II. ALSINÆÆ. DC. fl. fr. 4. p. 766.

Calycis sepala 4-5 libera aut imâ basi vix subcoalita.

X? ORTEGIA. *Lin. gen. n. 33. Gaertn. fruct. 2. t. 129. f. 8.*

Calyx 5-partitus. Corolla 0. Stamina 3, antheræ cordatæ. Stylus unicus, stigma capitatum. Capsula 1-locularis 3-valvis. Semina imâ capsulæ adfixa. — Glandula nigra utrinquè ad basin foliorum.

1. O. HISPANICA (Lin. spec. 49.) caulibus dichotomis, pedunculis brevibus, floribus dichotomo-corymbosis. 2/ in Hispaniâ. (v. s.)
2. O. DICHOTOMA (Lin. mant. 174.) caulibus dichotomis pedunculisque elongatis, floribus dichotomo-paniculatis. 2/ in Pedemontio et in Hispaniâ? O. Hispanica Cav. ic. 1. t. 47. — All. ped. st. t. 4. f. 1. O. dichotoma DC. fl. fr. 4. n. 4376. An species distincta? (v. s.) †.

XI. GOUFFEIA. *Robill. et Cast. diss. ined. DC. fl. fr. 5. p. 609.*

Calyx 5-partitus patulus. Petala 5 integra. Stamina 10. Styli duo. Capsula globosa 1-locularis 2-valvis 1-2-sperma.

1. G. ARENARIODES (Rob. et Cast! ined. DC! l. c.) ramis divaricatis, floribus subcorymbosis, foliis linearibus brevibus deflexis. ① circa Massiliam. Habitus Arenariæ tenuifol. (v. s.)

XII. BUFFONIA. *Sauv. meth. fol. 141. Lin. gen. n. 225. Gaertn. fruct. 2. t. 129. f. 1.*

Calyx 4-sepalus. Petala 4 integra. Stamina 4. Styli 2. Capsula compressa, 1-locularis, 2-valvis, 2-sperma.

1. B. ANNUA (DC! fl. fr. 4. p. 768.) caule basi laxè paniculato, ramis divaricatis brevibus firmis, striis calycinis rectis parallelis, capsulis calycem vix æquantibus, foliis subulatis basi dilatatis. ① in Galliæ austr. rupestribus. B. tenuifolia Lam. ill. n. 1710. t. 87. f. 1. (v. s.)
2. B. OLIVERIANA (Ser. mss.) caule apice ramoso, ramis filiformibus elongatis, striis calycinis rectis parallelis, capsulis subsessilibus calyce brevioribus, foliis brevissimis squamiformibus. ①? circa Teheran Persiæ. (v. s. ex herb. Olivier.)
3. B. MACROCARPA (Ser. mss.) caule ramosissimo dichotomo, ramis filiformibus rigidis longissimis nudiusculis divaricatis, sepalis ovatis obtusis, striis rectis parallelis, capsulis longè pedicellatis magnis calyce multò longioribus, seminibus subgloboso-compressis margine echinatis. 2/ in Persiâ a Teheran ad Isphahan. (v. s. ex h. Olivier.)

4. *B. PERENNIS* (Pourr. act. toul. 3. p. 319.) caule apice ramoso, ramis filiformibus elongatis, striis calycinis arcuatis convergentibus. 2 in Galliâ et in Vallesiâ. Lam. ill. n. 1711. t. 87. f. 2. DC! fl. fr. 4. p. 768. (v. s.)

XIII. *SAGINA*. Lin. gen. n. 236. Gaertn. fruct. 2. t. 129. f. 10.

Calyx 4-5-partitus. Petala 4-5 aut 0. Stamina 4-5. Capsula 4-5-valvis, 1-locularis, oo-sperma.

1. *S. CERASTOIDES* (Smith in trans. lin. soc. 2. p. 343.) caule diffuso dichotomo, foliis spathulatis obovatisque recurvis, pedunculis fructiferis reflexis, petalis obcordatis, laciniis calycinis lanceolatis acutis. ① in arenosis maritimis Bodotriæ insulis. Smith engl. bot. t. 166. (caps. 8-valv.) Rœm. fl. europ. fasc. 6. C. tetrandrum Smith comp. fl. brit. ed. 3. p. 72. Habitus Alsines mediæ.
2. *S. PROCUMBENS* (Lin. spec. 185.) glabra, ramis procumbentibus, foliis linearibus mucronatis, pedunculis fructiferis adscendentibus, petalis brevibus obtusis, laciniis calycinis subrotundis. ① ad muros et in arvis sabulosis Europæ. Smith engl. bot. t. 380. (v. s.)
3. *S. FILIFORMIS* (Pourr. chlor. hisp. n. 593.) caulibus erectis ramosissimis geniculatis, foliis linearibus glabris, pedunculis axillaribus terminalibusque dichotomis. ① in Pyrenæis frequens promiscuè cum *S. procumbente*, quâcum confunditur. Rœm. et Schult. syst. 3. p. 499.
4. *S. MARITIMA* (Smith engl. bot. t. 2195.) ramis erectiusculis, foliis lanceolatis brevissimis, pedunculis fructiferis adscendentibus, petalis lanceolatis calycem æquantibus, laciniis calycinis ovatis obtusis. ① in Angliâ. (Descript. ex ic.)
5. *S. STRICTA* (Fries novit. fl. suec. 3. p. 122.) glaberrima, caule pedunculisque strictis, foliis subcylindricis muticis, lobis calycinis lanceolatis acutis (calyce obtuso Horn. h. hafn. suppl. p. 122.) ① in littoribus Daniæ et Sueciæ ad Cimbrishavn. (v. s. comm. à cl. Horn.)
6. *S. CILIATA* (Fries in Billb. svensk. bot. ex Spreng. neue entd. 3. p. 220.) caule diffuso, foliis subulatis patulis ciliatis, calycibus acuminatis. — in Sueciâ.
7. *S. APETALA* (Lin. mant. p. 559.) hispido-pubescent, ramis erectiusculis dichotomis, foliis linearibus mucronatis, pedunculis fructiferis adscendentibus, petalis brevissimis subrotundis, laciniis calycinis lanceolatis obtusiusculis. ① in arenosis sterilibus Angliæ, Italiæ et Germaniæ.
β. agglomerata, floribus sessilibus pedicellatisque agglomeratis. — Circa Agatham. Ph. Thom.
8. *S. QUINTENIS* (H. B. et Kunth. gen. et spec. am. 6. p. 19.*) caulibus filiformibus procumbentibus? foliis subfalcatis, pedunculis axillaribus unifloris supra medium quadribracteatis, fructiferis elongatis rectis, floribus apetalis subpentandris. ① in ripâ arenosâ Rio-Blanco, altit. 1580-1600. hex. Valvæ revolutæ.
9. *S. ERRECTA* (Lin. spec. 185.) caulibus erectis subunifloris glabris, foliis linearibus acutis, pedunculis strictis, petalis brevibus, sepalis lanceolatis acutis. ① in sterilibus glareosis Europæ. Smith engl. bot. t. 609. DC! fl. fr. 4. p. 769. — Vaill. bot. par. t. 3. f. 2. *Mœnchia glauca* Pers. ench. 1. p. 153. (v. s.)
10. *S. FASCICULATA* (Poir. dict. 6. p. 390.) caulibus repetitis radicanibus articulatis, ramis suberectis, foliis fasciculatis secundis, pedunculis axillaribus unifloris. 2 circa Algeriam. Aff. *S. procumbenti*. Calyx glaber, lobis brevibus ovalibus obtusis patentibus, à fecundatione reflexis.

XIV. *HYMENELLA*. fl. mex. ic. ined.

Calyx 4-partitus patens. Petala 4 oblonga integra calycis longitudine. Stamina 4, petalis alterna basi coronulâ petaloideâ octodentatâ juncta. Ovarium ovatum. Styli 3. Capsula 3-locularis.

1. *H. MOEHRINGIODES* (fl. mex. ic. ined.) ? in Mexici horto P. Carmelitarum Sancti Angeli. *Buffonia tenuifolia* Moc. et Sessé fl. mex. ined. ic. *Caules debiles, folia linearia acuta glabra, pedicelli solitarii 1-flori axillares, flores parvi albi.*

XV. *MOEHRINGIA*. *Lin. gen.* 264. *Gaertn. fruct.* 226. *t.* 129. *f.* 11.

Calyx 4-partitus. Petala 4. Stamina 8. Styli 2. Capsula 4-valvis, 1-locularis, ∞-sperma.

1. *M. MUSCOSA* (Lin. spec. p. 515.) foliis linearibus connatis, laciniis calycinis planis longitudine ferè internodiorum lanceolatis acutis. ? in montibus humidis Europæ ad oras sylvarum. Lam. ill. t. 314.
2. *M. SEDIFOLIA* (Willd. ber. mag. 1818. p. 101. t. 3. f. 23.) foliis oblongis obtusis utrinquè convexis subimbricatis carnosis, floribus terminalibus solitariis. ? in Alpibus Tendæ. *M. muscosa* variet. Balb. misc. bot. 20. t. 5. f. 2. *M. muscosa sedoides* Pers. ench. 1. p. 438. Flores ut in *M. muscosa*, sed parum minores.
3. *M. STRICTA* (Sibth. et Smith fl. græc. 362. ex prodr. 1. p. 264.) foliis strictis basi scariosis ciliatis geniculo caulino brevioribus. ? in insulâ Cretâ.

XVI? *ELATINE*. *Lin. gen. n.* 685. *Gaertn. fruct.* 2. p. 142. t. 102. f. 2.

Calyx 3-4-partitus. Petala 3-4 unguibus destituta. Stamina petalis numero æqualia? aut dupla, idè 3-4 aut 6-8. Styli 4 apice capitati. Capsula 4-valvis 4-locularis ∞-sperma. Semina cylindracea.

1. *HYDROPIPER* (Lin. spec. 527.) foliis oppositis, floribus alternis pedicellatis octandris tetrapetalis. ? in inundatis totius ferè Galliæ. *E. hydro Piper a.* Lin. spec. 527. DC! ic. pl. rar. 1. p. 13. t. 43. f. 2. — Vaill. bot. par. t. 2. f. 2. (v. s.)
2. *E. HEXANDRA* (DC! ic. pl. rar. 1. p. 14. t. 43. f. 1.) foliis oppositis, floribus alternis pedicellatis hexandris tripetalis. ? in inundatis circa Parisios. *E. hydro piper* var. *β.* Lin. spec. 527. — Vaill. bot. par. t. 2. f. 1. *Tillæa hexandra* Lapièrre journ. phys. fl. an xi. *Birolia paludosa* Bellard. mem. acad. tur. 1808. ic. (v. s.)
3. *E. TRIANDRA* (Schkuhr bot. handb. n. 1023. t. 109. b. f. 2.) foliis oppositis, floribus sessilibus oppositisque triandris tripetalis. ? in locis inundatis circa Ratisbonam. Flores seminaque rosea.
4. *E. ALSINASTRUM* (Lin. spec. 527.) foliis verticillatis, floribus sessilibus verticillatisque. ? in aquilegiis Lutetiæ, Fontisbleaudi. — Vaill. bot. par. 1. f. 6. *E. verticillata* Lam. fl. fr. 3. p. 11. DC! fl. fr. 4. p. 772. * *Habitus Hippur. vulgaris.* (v. s.)

XVII? *BERGIA*. *Lin. gen. n.* 791.

Calyx 5-partitus. Petala 5. Stamina 10. Styli 5 approximati. Caps. 5-valvis, 5 locularis.

1. *B. VERTICILLATA* (Willd. sp. 2. p. 770.) foliis lanceolatis apice denticulatis, floribus verticillatis subsessilibus confertissimis. (U)? *Rosettæ et Damiatæ* in oryzatis. In Indiâ orient., nec ad cap. B.-Spei? *B. Capensis*. Lin. ment. 241. *Elatine laxurians* Delil! fl. ægypt. p. 13. t. 26. f. 1. (v. s. comm. à cl. Delil.)
2. *B. GLOMERATA* (Lin. f. suppl. 243.) foliis ovatis crenulatis, floribus glomeratis. — ad cap. B.-Spei. Thunb. prodr. 82.
3. *B. AMMANNIOIDES* (Roth. nov. pl. spec. 219.) caule ramoso setis capitatis aspero, foliis ellipticis acutis argutè inciso-serratis subtus tomentoso-argenteis, floribus axillaribus pedunculatis glomeratis pentandris — in Indiâ orient. *Corolla albida.*

XVIII. MOLLUGO. *Ser. mss.*

Calyx 5-partitus. Petala 0. Stamina 3-5. Styli 3. Capsula 3-valvis 3-localaris, ∞-sperma. — Folia verticillata rarò opposita. Flores solitarii vel umbellati.

Sect. 1. MOLLUGO. *Lin. gen. n. 139. Gaertn. fruct. t. 130. f. 8.*

Pedunculi uniflori verticillati.

1. *M. OPPOSITIFOLIA* (Lin. spec. 131.) foliis oppositis lanceolatis, ramis dichotomis, pedunculis verticillatis (I) in Zeylonâ. Pluck. phyt. t. 75. f. 6. Pedunculi fere longitudine foliorum.
2. *M. SERPYLLIFOLIA* (Ser. mss.) caulibus ramosis dichotomis filiformibus, pedunculis 1-floris, foliis oppositis petiolatis internodiis multoties brevioribus. (I) ad cap. B.-Spei. Pharnaceum serpyllifolium. Lin. suppl. p. 186. Roem. et Schult. syst. 6. p. 687.*
3. *M. HIRTA* (Thunb. prod. p. 24. fl. cap. 1. p. 444.) decumbens, foliis quaternis petiolatis, floribus verticillatis brevissimè pedunculatis. (I) ad cap. B.-Spei. †.
4. *M. VERTICILLATA* (Lin. spec. 131.) foliis verticillatis inæqualibus obovatis cuneiformibus acutis, caule decumbente dichotomo, pedunculis unifloris verticillatis. (I) prope Cumana ad littora maris Antillarum. Roem. et Schult. syst. 2. p. 871.* — Pluk. phyt. 4. t. 332. f. 5. Folia carnosâ. (v. s.)
5. *M. SCHRANKII* (Ser. mss.) foliis verticillatis sublinearibus acutis, caule dichotomo diffuso, pedunculis unifloris paucis subverticillatis. (I) in Brasiliâ. *M. dichotoma* Schrank pl. rar. 64.* ic. excl. Lin. f. syn. Similis *M. verticillatæ* à quâ differt foliis angustioribus longioribus et floribus 1-3 ex quovis foliorum verticillo.
6. *M. SPERGULA* (Lin. spec. 131.) foliis verticillatis inæqualibus subpetiolatis subcarnosis elliptico-lanceolatis, caule procumbente dichotomo, pedunculis verticillorum simplicibus 1-2, terminalibus subramosis. (I) in Indiâ Orient. Burm. fl. ind. 31. t. 5. f. 4. zeyl. 13. t. 7. Pharnaceum mollugo Lin. mant. p. 561. An variet. pauciflora *M. verticillatæ*?
7. *M. BERTERIANA* (Ser. mss.) caule dichotomo-gracili, internodiis longissimis, foliis angustissimis obtusiusculis, pedunculis filiformibus verticillatis. (I) ? in Porto-Ricco secus torrentes. Pharnaceum lineare Bert! ined. (v. s.)

Sect. 2. PHARNACEUM. *Sp. Lin. gen. n. 517. Gaertn. fruct. t. 130. f. 4.*

Pedunculi bifidi racemosi vel umbelliferi.

8. *M. STRICTA* (Lin. spec. 131.) foliis subquaternis lanceolatis, ramis paniculatis nutantibus, caule erecto anguloso. (I) in Asiâ. — Pluk. alm. p. 21. t. 257. f. 2.
9. *M. BELLIDIFOLIA* (Ser. mss.) caulibus erectiusculis aphyllis, foliis radicalibus ovato-spatulatis basi in petiolum attenuatis, floribus paniculatis. — In Jamaicâ, Guianâ. Pharnaceum bellidifolium Poir. dict. 5. p. 262.* Pharnaceum spathulatum Swartz fl. ind. occid. 1. p. 568. — Plum. amer. t. 21. f. 1. (v. s. in herb. Balb.)
10. *M. PARVIFLORA* (Ser. mss.) foliis ovatis obtusis subtus strigosis, umbellis lateralibus, caulibus herbaceis adscendentibus. — in Indiâ orient. Pharnaceum parviflorum Roth nov. pl. spec. p. 186.*
11. *M. PENTAPHYLLA* (Lin. spec. 131.) foliis subquinis obovatis glabris, floribus paniculatis, caule decumbente. (I) ? in Zeylonâ. Burm. zeyl. 13. t. 8. f. 1. Folia 3-4-5, utrinquè saturatè viridia; pedunculi axillares longissimi. (v. s.)
12. *M. NUDICAULIS* (Lam. dict. 4. p. 234.*) foliis radicalibus oblongis numerosis obovatis confertis, scapis trichotomo-paniculatis decumbentibus nudis.

- ①? in Indiâ orient. Burm. zeyl. 14. t. 8. f. 2. Folia radicalia glauco-viridia magna. Flores ochroleuci. Semina nigerrima. (Burm. l. c.)
13. *M. RADIATA* (Ruiz et Pav. fl. peruv. 1. p. 48.) caulibus procumbentibus foliosis ramis paniculatis, foliis obovatis acuminatis. — in inundatis Chili Conceptionis. Rœm. et Schult. syst. 2. p. 872. — Pluk. phyt. t. 118. f. 1. (v. s.)
14. *M. DISTICHA* (Ser. mss.) foliis sublinearibus pubescentibus, racemis bipartitis flexuosis. ① in Indiâ orient. et ad cap. B.-Spei. Pharnaceum distichum Lin. mant. p. 211. Mollugo racemosa. Lam. ill. n. 1197. — Pluk. phyt. 3. p. 22. t. 130. f. 6.
15. *M. TRIPHYLLA* (Lour. fl. coch. ed. Willd. 1. p. 79. non Link.) caulibus diffusis, ramis ternatis, foliis ternis lanceolatis sessilibus, floribus dichotomis longè pedunculatis, pedunculis apice ramosis. ①? ad Cantonem Sinarum.
16. *M. CORDIFOLIA* (Ser. mss.) caulibus decumbentibus, foliis quaternis obcordatis floribus dichotomis, flore solitario in dichotomiâ. ① ad cap. Bonæ-Spei. Pharnaceum cordifolium Lin. amœn. 6. p. 85. Jacq. b. Schœn. t. 349.
17. *M. LXXII* (Ser. mss.) caule erectiusculo, foliis verticillatis tribus maximis, paniculis terminalibus lateralibusque. ① in Brasiliâ. *M. triphylla* Link enum. 1. p. 108. non Lour. Folia acuta glaberrima.
18. *M. MULTIFLORA* (Ser. mss.) caulibus diffusis dichotomis, foliis verticillatis lanceolatis inæqualibus, floribus disticho-paniculatis numerosis. ① in Chinâ. (v. s. comm. à cl. Lamb.)
19. *M. DICHOTOMA* (Ser. mss. non Schrank pl. rar.) foliis verticillatis linearibus, pedunculis axillaribus elongatis dichotomis. ① ad cap. B.-Spei. Pharnaceum dichotomum Lin. f. suppl. p. 186. Schult. syst. 6. p. 689.*
20. *M. GLOMERATA* (Ser. mss.) caulibus flexuosis, foliis verticillatis linearibus acutis glabris, floribus glomeratis subsessilibus. ① ad cap. B.-Spei. Pharnaceum glomeratum Lin. f. suppl. p. 185. — Pluk. phyt. 163. t. 331. f. 4. (v. s. in herb. Balb.)
21. *M?* *MARGINATA* (Ser. mss.) caulibus filiformibus decumbentibus, foliis verticillatis ovatis marginatis obtusis, floribus axillaribus sessilibus. ①? ad cap. B.-Spei. Pharnaceum marginatum Thunb. prod. p. 54. fl. cap. 2. p. 239.
22. *M. CERVIANA* (Ser. mss.) caulibus verticillatis rariùs dichotomis, floribus umbellatis, foliis verticillatis 3-5 angustissimis glaucis obtusis glabrinisculis. ① in Russiâ, Hispaniâ, Guineâ et Asiâ. Pharnaceum Cerviana Lin. spec. ed. 1. p. 272. ed. 2. p. 388. Poir. dict. 5. p. 259.* Lam. ill. t. 214. f. 1. — Pluk. phyt. t. 332. f. 11. (v. s.)
23. *M. ARENARIA* (H. B. et Kunth gen. et spec. am. 6. p. 21.*) glabra, caulibus dichotomis diffusis teretibus, foliis quinis lineari-spathulatis obsolete uninnerviis internodio quadruplò brevioribus, umbellis subsessilibus 5-6-floris. ① in insulâ arenosâ fluminis Apures prope El Diamante. Stamina tria calyce breviora.
24. *M. SPERGULOIDES* (Ser. mss.) caulibus verticillatis 3-4, ramis simplicibus, umbellis involucretis terminalibus paucifloris, foliis fasciculato-verticillatis numerosis angustissimis acutis. ①? in Indiâ. Pharnaceum sperguloides Poir. dict. 5. p. 260.* Lam. ill. t. 214. f. 2.
25. *M. LINEARIS* (Ser. mss.) caulibus verticillatis, ramis dichotomis, umbellis terminalibus lateralibusque, foliis linearibus obtusis. ①? ad cap. B.-Spei. Pharnaceum lineare Lin. f. suppl. p. 185. Poir. dict. 5. p. 261. Schult. syst. 6. p. 633. (v. s. in herb. Balb.)
26. *M?* *MICROPHYLLA* (Ser. mss.) tomentosa, pedunculis umbellatis, foliis ovatis teretiunculis obtusis lanâ intertextis. ①? ad cap. B.-Spei. Pharnaceum microphyllum Lin. f. suppl. 185. Schult. syst. 6. p. 686.* Caulis fruticosus rigidus, ramuli subverticillato-aggregati, folia è nodis fasciculata et per caulem sparsa. †

27. *M. TERETIFOLIA* (Ser. mss.) glabra, ramis oppositis parum ramulosis, foliis filiformibus mucronatis teretibus, caule erecto frutescente, umbellis lateralibus simplicibus pedunculatis. ♂ ad cap. B.-Spei. Pharnaceum teretifolium. Thunb prodr. p. 53.
28. *M. QUADRANGULARIS* (Ser. mss.) caulibus suffrutescentibus erectis, ramis alternis flexuosis aphyllis, foliis linearibus quadrifariam imbricatis, floribus umbellatis, pedunculis brevibus. ♂ ad cap. B.-Spei. Pharnaceum quadrangulare Lj. f. suppl. 185. Schult. syst. 6. p. 688.* Facies *Ericæ*.
29. *M. MUCRONATA* (Ser. mss.) caule herbaceo subnullo, foliis ovatis mucronatis integris, floribus verticillato-aggregatis subsessilibus (I)? ad cap. Bonæ-Spei. Pharnaceum mucronatum Thunb. in phyt. bl. p. 29., fl. cap. 2. p. 239.
30. *M. UMBELLATA* (Ser. mss.) foliis radicalibus stellatis, caulinis obovatis, pedunculis umbellatis involucreatis, involucri linearibus in campis Lohagæ argillosis. Pharnaceum umbellatum Forsk. fl. ægypt. p. 58.

† *Species non satis notæ.*

31. *M. MARITIMA* (Ser. mss.) floribus lateralibus solitariis sessilibus, foliis subteretibus obtusis carnosius — in Carolinâ. Pharnaceum maritimum Walt. fl. car. p. 117.
32. *M. LICHTENSTEINIANA* (Ser. mss.) foliis verticillatis lanceolato-linearibus acutis, pedunculis axillaribus — ad montem Witsemberg cap. B.-Spei. Pharnaceum Lichtensteinianum. Schult. syst. 6. p. 692. Ph. galioides Lichtenst. spic. fl. cap. mss. Simillima *Spergulæ*, sed monogyna. An genus proprium?
33. *M. HOFFMANNSEGGIANA* (Ser. mss.) foliis verticillatis linearibus acutis, pedunculis 1-floris elongatis. (I) in Brasiliâ. Ph. Hoffmannseggianum Schult. syst. 6. p. 692. Ph. galioides Willd. mss. ex Schult. l. c.

XIX. PHYSA. *Pet. Thouar. nov. gen. mad. p. 20.*

Calyx 5-sepalus. Petala 0. Stamina 10. Stigmata 3. Capsula trisulca, 3-locularis, 3-valvis; valvæ septiferæ cum receptaculo conniventes.

1. *P. MADAGASCARIENSIS* Plantula prostrata. Cauliculi articulati. Folia 4^{im} verticillata inæqualia. Pedunculi 1-flori. (P. Th. l. c.)

XX. HOLOSTEUM. *Lin. gen. n. 136. Gaertn. fruct. t. 130. f. 5. excl. sp.*

Calyx 5-sepalus. Petala 5 dentata. Stamina 5 aut abortu 3-4. Styli 3. Capsula 1-locularis apice in 6 dentes dehiscens. Embryo intra albumen replicatus.

1. *H. DIANDRUM* (Swartz prodr. p. 27. icon. t. 7.) caulibus procumbentibus rigidioribus, foliis subrotundis, floribus diandris, stipulis utrinque quaternis. (C) in petrosis Jamaicæ. Rœm. et Schult. syst. 2. p. 857.* Aff. *H. cordato* sed multo minus. An *Drymaria* spec? (v. s.)
2. *H. MUCRONATUM* (Moc. et Sessé. pl. mex. ined. ic.) pubescens, caulibus procumbentibus, foliis cordatis, stipulis pedunculis trifloris. — in Mexico. An var. *H. cordati*? an *Drymaria* sp? (ex ic stylus 1. stigm. 3.)
3. *H. SUCCULENTUM* (Lin. amœn. 3. p. 21.) foliis ellipticis carnosius glabris, petalis subtrifidis calyce minoribus. — in Noveboraco. Pursh fl. bor. am. 1. p. 90. Polycarpon uniflorum Walt. fl. car. p. 83.
4. *H. HIRSUTUM* (Lin. amœn. 3. p. 21.) foliis orbiculatis hirsutis, floribus sessilibus? — in Malabariâ.
5. *H. UMBELLATUM* (Lin. spec. 130.) foliis radicalibus ellipticis glaucis glabris, caulinis majoribus ovatis, floribus umbellatis, pedunculo-communi viscoso-pubescente, pedicellis post anthesin deflexis, valvis circinnatis. (I) in Europæ arvis murisque. Lam. illust. t. 51. f. 1. (v. v. et s.)

ℒ. floribus roseis. *H. umbellatum* Smith. engl. bot. t. 27.*

XXI. SPERGULA. *Lin. gen. n. 798. Gaertn. fruct. t. 130. f. 4.*

Calyx 5-partitus. Petala 5 integra. Stamina 5-10. Styli 5. Capsula 1-locularis, 6-valvis, ∞-sperma.

§. 1. *Foliis verticillatis basi stipulatis.*

1. *S. ARVENSIS* (Lin. spec. 630.) foliis verticillatis basi stipulatis, floribus decandris, seminibus sphæroideis hispidulis angustè marginatis nigris. ① in Europæ agris. Lam. ill. t. 392. f. 1, fl. dan. t. 1033. (v. s.)

β. *geniculata* (Poir. dict. 7. p. 303.) caule ultra pedali geniculato ad nodos quasi viviparo. — *S. geniculata* Pers. ench. 1. p. 522. n. 3.

2. *S. PENTANDRA* (Lin. spec. 630.) foliis verticillatis basi stipulatis, floribus subpentandris, seminibus lenticularibus margine punctatis alatisque. ① in sylvaticis Europæ vere. Lam. ill. t. 392. f. 2. (v. s.)

3. *S. VILLOSA* (Pers. ench. 1. p. 522.) pubescens, foliis verticillatis basi stipulatis, petalis calyce brevioribus. — in Monte Video. Poir. dict. 7. p. 304.* *Sepala linearia.*

4. *S. VISCOSA* (Lag in varied. de cienc. 1805. p. 213. gen. et spec. 15.) foliis verticillatis villosis-viscidis, floribus decandris, petalis calyce longioribus. ② in Hispaniâ juxta nives in summo monte vulgò de *Lospozos* prope Arvas.

5. *S. PALLIDA* (Salisb. prod. 298.) caule viscoso pubescente, stipulis magnis foliis semiteretibus mucronulatis, petalis ovalibus integerrimis obtusis. — in promontorio B. Spei. Semina imbricata paululum reniformia, margine scariosa dense striata erosula.

§. 2. *Foliis oppositis, junioribus in axillâ fasciculatis; stipulis nullis.*

6. *S. NODOSA* (Lin. spec. 630.) foliis oppositis subulatis lævibus obtusiusculis, infimis latè vaginantibus, caulibus cespitosus subsimplicibus paucifloris, petalis calyce duplò longioribus, seminibus subreniformibus rugosis. ② in paludosis Europæ. Smith engl. bot. t. 694. — OEd. fl. dan. t. 96. (v. v. et s.)

β. *brevifolia* (Pers. ench. 1. p. 522. Poir. dict. 7. p. 305.) caule simplicissimo, nodis approximatis, foliis caulinis brevissimis.

γ. *maritima* (Pers. ench. 1. p. 522.) foliis carnosulis incurvo-patulis. — ad maris littora.

7. *S. LARICINA* (Lin. spec. 631.) decumbens, foliis oppositis rigidis subulatis ciliatis, floribus decandris, petalis calyce duplò longioribus, seminibus disciformibus ciliatis. ② in Sibiria. Smith ic. ined. 1. p. 18. t. 18. Lam. ill. t. 392. f. 3. (v. s. comm. à cl. Fisch.)

8. *S. SAGINOIDES* (Lin. spec. 631.) caule repente, foliis oppositis linearibus lævibus, pedunculis solitariis longissimis, petalis obtusissimis sepala obtusa vix æquantibus, seminibus subreniformibus punctatis. ① in Europâ et Carolina merid. Smith engl. bot. t. 2105. Habitus *Saginx* procumbentis. (v. s.)

9. *S. SUBULATA* (Swartz act. holm. 1789. p. 45. t. 1. f. 3.) subpilosa, foliis oppositis subsecundis lineari-subulatis subaristatis, pedunculis solitariis longissimis, petalis calycis longitudine. ② ① in sabulosis humidis Europæ. Smith engl. bot. t. 1082. *S. saginoides* Lam. fl. fr. 3. p. 55 (v. s.)

10. *S. PILIFERA* (DC! fl. fr. 4. n. 291.*) foliis oppositis linearibus aristatis subrigidis glabris fasciculatis caulibus repentibus ramosis cespitosus, pedunculis longissimis, petalis calyce duplò majoribus, seminibus subovoideis. — in montibus editioribus Corsicæ. (v. s.)

11. *S. GLABRA* (Willd. spec. 1. p. 821.) decumbens, foliis oppositis filiformibus glabris subacutis, floribus decandris, petalis calyce majoribus. ② in pratis umbrosis Europæ. Poir. dict. 7. p. 306.* *S. saginoides*. All. pedem. n. 1735. t. 64. f. 1. (v. s.)

12. *S. GLANDULOSA* (Besser prim. flor. galic. p. 298.) pubescenti-glandulosa,

foliis oppositis subulatis, caulibus supernè ramosis, axillis fasciculiferis, petalis ellipticis sepalis obtusis duplò longioribus.—in pascuis humidiusculis Leopoli.

13. *S. ARENARIOIDES* (Ser. mss.) glabra, radice crasso, caule prostrato, ramis adscendentibus, foliis lanceolato-linearibus acutis exstipulatis, floribus pentandris, petalis oblongis calycis longitudine. — in Mexico. *Arenaria?* pentagyna Moc. et Sessé pl. mex. ined. ic. (descr. ex. icon.)
14. *S. APETALA* (Labill! nov. holl. 1. p. 112.* t. 142.) subcaulis, foliis oppositis connatis imbricatis longis, floribus apetalis pentandris, sepalis lanceolatis acutissimis. — in capite Van^dDiemen. (v. s.)

XXII. LARBREA. *St. Hil. in mem. mus. 2. p. 261.*

Calyx 5-fidus basi urceolatus. Petala 5 bipartita perigyna. Stamina 10 perigyna. Styli 5. Ovarium 1-loculare, ∞ -spermum, Capsula apice 6-valvis.

1. *L. AQUATICA* (St. Hil. 1. c.) caulibus dichotomis, foliis lanceolatis superioribus sessilibus, floribus axillaribus longè pedunculatis post anthesin deflexis. $\frac{1}{2}$ in aquosis Europæ. *Cerastium aquaticum*. Lin. spec. 629. Smith engl. bot. t. 538. *Stellaria aquatica* Scop. fl. carn. n. 546. (v. s.)

XXIII. DRYMARIA. *Willd. herb. ex Rœm. et Schult. syst. 5. p. 406. H. B. et Kunth gen. et spec. 6. p. 21.*

Calyx 5-partitus. Petala 5 bifida. Stamina 5. Styli 3. Capsula 4-sperma ad basim trivalvis 5- ∞ -sperma. Embryo periphericus subannularis. — Herbæ diffusæ ramosæ. Stipulæ petiolares geminæ vel plures.

1. *D. FRANKENIOIDES* (H. B. et Kunth gen. et spec. am. 6. p. 21. t. 515.) ramosissima glanduloso-hirtella viscidula canescens, foliis brevissimè petiolatis lanceolato-oblongis, pedunculis 1-floris, petalis calyce brevioribus 6-fidis, laciniis 4 intermediis capillaceis, ovariis sub 50 spermis. (I) in arenosis prope Pachuca, alt. 1270 hex! (regno Mexicano.) *D. arenarioides* Willd. herb. ex Rœm. et Schult. syst. 5. p. 406. Semina angulato-reniformia tuberculis minutissimis densissimè obsita.
2. *D. STELLARIOIDES* (Willd. herb. ex Rœm. et Schult. syst. 5. p. 406.) caulibus apice dichotomis paucifloris, foliis sessilibus glabris, calyce viscido pubescente petalis bifidis brevioribus, ovariis sub 50-spermis. (I) prope Hambato, alt. 1380. hex. (regno Quitensi.) H. B. et Kunth gen. am. et spec. 6. p. 22.* t. 5.6.
3. *D. OVATA* (Rœm. et Schult. syst. 5. p. 406.) caulibus supernè ramisque villosis, foliis petiolatis ovatis acuto-mucronatis basi rotundatis puberulis, pedunculis bifidis paucifloris, calyce glabro corollam æquante, petalis profundè bifidis, ovariis 13-17-spermis. (I) prope Quito, alt. 1460. hex. H. B. et Kunth gen. et spec. am. 6. p. 23.*
4. *D. CORDATA* (Willd. herb. ex Rœm. et Schult. syst. 5. p. 406.) caule foliisque glabris breviter petiolatis ovato-subrotundis acuto-mucronatis basi rotundatis vel obsolete cordatis, pedunculis dichotomis multifloris, calyce glabro petala superante, ovariis 7-10-spermis. (I) in Jamaica, Surinami et in Novâ-Andalusiâ. H. B. et Kunth gen. et spec. 6. p. 23.* *Holosteum cordatum* Lin. amœn. 3. p. 21. mant. 327. Lam. ill. t. 51. f. 2. (v. s.)
5. *D. DIVARICATA* (H. B. et Kunth gen. et spec. 6. p. 24.*) caule foliisque glabris petiolatis late subrotundo-ovatis acutis basi rotundatis, pedunculis irregulariter ramosis divaricatis multifloris, calyce glabro petalis parum brevioribus, ovariis subocto-spermis. (I) ad littora Oceani Pacifici prope Lima Peruvianorum. Differt à præcedente præsertim pedunculis irregulariter ramosis, pedicellis longioribus, petalis calycem superantibus et stylo longiore.

XXIV. STELLARIA. *Lin. gen. n. 773. Gaertn. fruct. 2. t. 130. f. 3. (1)*

Calyx 5-partitus. Petala 5 bifida. Stamina 10 aut abortu 3-8. Styli 3. Capsula 1-locularis apice 6 valvis oo-sperma.

1. *S. NEMORUM* (Lin. spec. 603.) foliis inferioribus cordatis petiolatis, superioribus lanceolatis subsessilibus, paniculâ dichotomâ, petalis calyce duplò longioribus, seminibus orbiculatis compressis margine tuberculatis. ♀ in nemoribus et umbrosis Europæ. Smith engl. bot. t. 92. *Alsine nemorum* Schreb. spic. 30. — OEd. fl. dan. t. 271. (v. v. et s.)
2. *S. CUSPIDATA* (Willd. herb. ex Schlecht. berl. mag. 1816. p. 196.) caule dichotomo glabro, foliis petiolatis ovato-oblongis acuminatis cordatis membranaceis glabris, pedunculis alaribus, calyce basi ciliato, ovariis sub 3o-spermis. ♀ in regni Quitensis nemoribus. H. B. et Kunth gen. et spec. am. 6. p. 27.* *Stellariz nemorum* simillima. (Kunth l. c.)
3. *S. SAXIFRAGA* (Bertol! pl. ital. rar. ed. 1. p. 55. n. 4.*) pubescens diffusa, foliis sessilibus ovato-acutis basi dilatatis rugosis, paniculâ dichotomâ, pedunculis terminalibus subternis lateralibus bracteatis sepalis angustè lanceolatis corollâ dimidiò brevioribus, seminibus subreniformibus rugosis. ♀ in vertice *Sagro*, *Alpium Appuanorum*. (v. s. in h. Balb. comm. à cl. Bertol.)
4. *S. CORDATA* (Willd. herb. ex Schlecht. berl. mag. 1813. p. 197.) foliis subrotundis cordatis, pedunculis axillaribus. — in vertice *Caripæ* in America. Humb. et Bonpl.
5. *S. VIRGATA* (Ser. mss.) glabra, caule striatò dichotomo, foliis ovato-cordatis trinerviis, floribus terminalibus 3-4 subumbellatis brevè pedunculatis, sepalis oblongis margine scariosis obtusiusculis corollâ multò longioribus, stylis brevissimis, capsulis. . . — Patr. ign. *Alsine virgata* Deless. herb. (v. s.)
6. *S. ARISTATA* (Ser. mss.) caulibus cespitosis, dichotomis, foliis ovatis apice aristatis basi ciliatis patentibus, floribus cymosis pedunculatis, sepalis lanceolatis corollâ brevioribus, petalis bidentatis, staminibus 5, ovariis ovatis, stylo 1, stigmatibus 3. ♀ in Mexico. *Alsine?* *aristata* Moç. et Sess. pl. mex. ined. (descript. ex ic.)
7. *S. RADIANS* (Lin. spec. 603.) caule dichotomo, foliis lanceolatis acutis suberrulatis sessilibus, petalis 5-partitis calyce duplò longioribus. (1)? in Sibiriz uliginosis Poir. dict. 7. p. 416.* *Wikstrøm* acad. handl. 1822*. (v. s.)
Æ. uniflora caule simplici uniflora. — *Amm. stirp. ruth.* t. 10.
8. *S. VILLOSA* (Poir. dict. 7. p. 416.* non Schlecht.) caule pedunculisque hirsutis, foliis sessilibus ovato-lanceolatis pubescentibus, floribus longè pedunculatis subpaniculatis terminalibus, corollis calyce longioribus. — in insulâ *Borboniz*. *Commerson*.
9. *S. LANCEOLATA* (Poir. dict. 7. p. 416.) foliis lanceolatis oblongis acutis, paniculâ pubescente, sepalis corollâ longioribus. — ad fretum *Magellanicum*.
10. *S. LATIFOLIA* (Pers. ench. 1. p. 501.) caulibus diffusis dichotomis basi subradicantibus, foliis ovatis, inferioribus petiolatis subcordatis, superioribus sessilibus, pedicellis axillaribus, junioribus umbellato-aggregatis, fructiferis reflexis, petalis calyce brevioribus, seminibus disciformibus rugosis. (1) in aquaticis circa *Monspelim* et in *Germaniâ* DC. fl. fr. suppl. 5. p. 614.* *S. Cerastium*. *Linn. syst. veg.* ed. 15. p. 452. (v. s.)
11. *S. MEDIA* (Smith engl. bot. t. 537. fl. brit. p. 473.) caulibus procumbentibus lineâ laterali pilosâ alternâ, foliis lanceolatis tenerrimis, fructibus deflexis, capsulis profundè 6-valvibus calyce vix longioribus, seminibus

(1) Genus quoad specierum diagnoses difficillimum et omnino de novo elaborandum.

- subreniformibus rugosis. ① ad sepes Europæ frequens. *Alsine media* Lin. spec. 389. Lam. ill. t. 214. DC! (v. v.)
12. *S. ROTUNDIFOLIA* (Poir. dict. 7. p. 416. *) foliis petiolatis rotundis crassiusculis 3-nervatis submucronatis, paniculâ terminali, ramis virgatis, corollâ sepalis longiore. — ad fretum Magellanicum Commerson. *Drymaria* species ex cl. Kunth.
13. *S. CILIATA* (Vahl in h. Juss. Pers. ench. 1. p. 503. non H. B. et Kunth nec Scop.) foliis cordatis acutis subpetiolatis parvis, petiolis ciliatis, floribus axillaribus solitariis, petalis calycem vix superantibus. — in Peruvîâ. Poir. dict. 7. p. 516.
14. *S. DICHOTOMA* (Lin. spec. 603. non Willd. herb.) pilosa, foliis cordato-ovatis subamplexicaulibus, caule dichotomo, floribus solitariis, pedunculis fructiferis reflexis, sepalis lanceolatis acutis petala superantibus. ①? in Alpibus Sibiricis. Smith ic. ined. 1. p. 14. t. 14. opt. (v. s.)
15. *S. PUBERA* (Michx. fl. bor. am. 1. p. 273.) pubens, foliis ovalibus ciliatis, pedicellis erectis breviusculis, sepalis ovatis, petalis calyce longioribus. — in montibus sylvaticis Carolinæ sept. (v. s. in herb. Balb.)
16. *S. SULBOSA* (Wulf. in Jacq. coll. 3. p. 21. * ic. rar. 3. t. 468.) foliis ovato-lanceolatis subtus aveniis, caule subramoso, pedunculo unifloro, sepalis lanceolatis acutis petalis duplò brevioribus, radice filiformi repente bulbiferâ. 2/ in Alpibus Carinthiæ. (v. s.)
17. *S. VISCIDA* (Bieb. fl. taur. 1. p. 342. *) villosa-viscosa, foliis linearilanceolatis, caulibus dichotomis diffusis, petalis calyce longioribus, capsulis subcylindricis sepalis lanceolatis duplò longioribus. ① in Tauriæ et Caucasi campestribus. Waldst. et Kit. pl. hung. 1. t. 22. *Cerastium anomalum* Willd. spec. 2. p. 812. (v. s.)
- ℓ. *glabriuscula* (Bieb. fl. taur. suppl. p. 307.)
18. *S. SABULOSA* (Fisch! in litt.) villosa-viscosa, foliis linearibus angustissimis, caulibus paniculato-dichotomis, petalis calyce vix longioribus, capsulis subcylindricis sepalis lanceolatis ferè duplò longioribus. ①? in Persiâ circa Leukheran. An var. *S. dubiæ* Bast? (v. s. comm. à cl. Fisch.)
19. *S. DUBIA* (Bast! suppl. p. 24. DC! fl. fr. suppl. p. 614. *) foliis linearibus glabris margine subciliatis, caule erecto, pedunculis erectis, sepalis trinerviis. ① in Galliâ. *Cerastium arvense trigynum* Bast. fl. Maine et Loire. p. 163. (v. s.)
20. *S. HOLOSTEA* (Lin. spec. 711.) foliis lanceolatis acuminatis serrulatis scabriusculis, superioribus latioribus brevioribusque, pedunculis filiformibus prælongis, petalis semibifidis calyce lanceolato acuto longioribus. 2/ ad sepes et in dumetis Europæ. Smith engl. bot. t. 511. (v. v. et s.)
21. *S. LAXMANNI* (Fisch! in litt.) caule erecto paucifloro, foliis linearibus acutis integris subglabris, pedunculis filiformibus prælongis, petalis bipartitis sepalis lanceolato-acutis subduplò longioribus. — in Sibirîâ orient. (v. s. comm. à cl. Fisch.)
22. *S. VELUTINA* (Ser. mss.) molliter pilosa, foliis oblongo-linearibus distantibus, pedunculis filiformibus prælongis, petalis bilobis sepalis oblongis longioribus *S. mollis* Fisch! in litt. non Schlecht. (v. s. comm. à cl. Fisch.)
23. *S. GRAMINEA* (Lin. spec. 604.) foliis linearibus margine lævibus, caulibus diffusis, floribus paniculatis divaricatis, petalis calycis longitudine. 2/ ad margines sylvarum Europæ. Smith engl. bot. t. 803. (v. v. et s.)
24. *S. FRIESIANA* (Ser. mss.) foliis linearibus canaliculatis margine carinâque scabris basi ciliatis, caule flaccido, pedunculis solitariis, calyce nervi petala æquante. — ad Frensjö in Smolandîâ. *S. longifolia* Fries in Billb. svensk. bot. ex Spreng. neue entd. 3. p. 217. non Muhl. in Willd. enum.
25. *S. GLAUCA* (With. bot. arrang. 1. p. 420.) glaucæ, foliis linearilanceolatis margine lævibus, floralibus scariosis, petalis calyce duplò longioribus, caule

pubescente, foliis oblongo-lanceolatis anacronulatis, pedunculis lanceolatis solitariis longissimis, floribus apetalis. — in Carolinâ et Georgiâ.

51. *S. LONGIFOLIA* (Muhlenb. in Willd. enum. p. 479. non Fries.) foliis linearibus acutis, panicula terminali, petalis acutis bipartitis calyce brevioribus — in Pensylvaniâ.

52. *S. MURALIS* (Link enum. 1. p. 429.) glanduloso-pubescentis viscosa, caule procumbente, foliis ovatis in petiolum attenuatis carnosissimis, petalis calyce vix longioribus exsertis. (1) in Cretâ. *Arenaria muralis* Sieber. cret. exsic. Sepala acuta nervosa. Species intermedia inter *Stellariam* et *Arenariam*. (v. s. in herb. Ser.)

53. *S. RADICANS* (Lapeyr. fl. pyr. t. 93. ex Abr. p. 250.) caulibus prostratis teretibus multifloris, foliis ellipticis obtusis lucidis, caulinis subsecundis, pedunculis divaricatis, sepalis linearibus obtusis corollâ duplò longioribus, capsulis pyriformibus. 2 in Pyrenæis.

54. *S. RUPESTRIS* (Scop. fl. carn. 1. p. 317. t. 18. f. 1.) foliis planis subtus 3-nerviis, caulibus villosulis paucifloris, petalis ovatis acuminatis calyce brevioribus. — in Alpibus Carniolæ. †.

55. *S. ELEGANS* (Ser. mss.) caulibus basi prostratis, foliis ellipticis parvis glabris, floribus dichotomo-paniculatis, pedunculis calyce duplò longioribus puberulis, sepalis lanceolatis acutis glabriusculis margine submembranaceis, petalis calyce duplò longioribus, capsulis. . . . — in Sibiria Altaicâ. *Cerastium elegans* Fisch! in litt. (v. s. comm. à cl. Fisch.)

56. *S. LONGIPES* (Goldie plant. canad. in edimb. phil. journ. april. 1822.) glaberrima, foliis lineari-lanceolatis, pedunculis terminalibus dichotomè ramosis bracteatis, pedicellis longis, petalis latè obovatis bipartitis calyce obtuso trinervi vix longioribus. — in sylvis prope lacum Ontario.

XXV. ARENARIA. *Lin. gen. n. 774. Gaertn. fruct. 2. t. 130. f. 9.*

Calyx 5-sepalus. Petala 5 integra. Stamina 10 aut abortu pauciora. Styli 3. Capsula 1-locularis apice 6-3 valvis, ∞-sperma.

Sect. I. SPERGULARIA. *Pers. ench. 1. p. 504. — Stipularia Haw. — Lepigonum Wahlénb fl. goth. ex Spreng. neue entd. 3. p. 231.*

Capsula trivalvis. Folia linearia, stipulis scariosis ad basin donata.

1. *A. SEGETALIS* (Lam. fl. fr. 3. p. 43.) glabra, caule erecto, foliis subulatis stipulatis subsecundis, sepalis scariosis medio lineâ viridescente longitudinalliter notatis, petalis calyce brevioribus, pedunculis defloratis deflexis, seminibus subpyriformibus rugosis. (1) inter segetes Gallie et Hispanie. *Alsine segetalis* Lin. spec. 390. — Vaill. bot. par. t. 3. f. 3. Capsula 3-valvis, valvis latis obtusis. (v. s.)

2. *A. MOLLUGINEA* (Ser. mss.) glaberrima, foliis linearibus crassis glaucis stipulatis apice subcallosis, caule patulo dichotomo, floribus axillaribus post anthesin deflexis, sepalis ovatis obtusis margine membranaceis, capsulis subrotundis 3-valvibus, valvis obtusissimis calycem æquantibus, seminibus reniformibus vix rugosis. (1) in Novâ-Hispaniâ. *Alsine molluginea* Lag. gen. et spec. 1815. p. 13. (v. s. comm. à cl. Lag.)

3. *A. PROSTRATA* (Ser. mss.) caule dichotomo prostrato glabro, foliis stipulatis linearibus subverticillatis, ramis extremis filiformibus, pedunculis axillaribus 1-4-floris, sepalis lanceolatis inæqualibus corollam æquantibus, stylo unico, stigmatibus 3, capsulis ovato subrotundis 3-valvibus, valvis ovatis obtusissimis. 2? in Ægypto. *Alsine prostrata* Forsk. descr. p. 207. Delil. fl. æg. p. 68.* t. 24. f. 4. Flores rosei.

4. *A. SUCCULENTA* (Ser. mss.) caule humifuso passillo glabro, ramis dichotomis, foliis spathulatis, floribus terminalibus subpaniculatis, sepalis oblongis obtusis margine membraceis corollæ longitudine, stylo 1, stigmatibus 3, cap-

arenaria 3-valvis, foliis lanceolatis post dehiscentiam margine involutis. ① in AEGYPTO. *A. succulenta* Delil. fl. ægypt. p. 68. t. 24. f. 3. Flores albi.

5. *A. PURPUREA* (Pers. ench. 1. p. 50. non Schlecht.) hispida, caule erecto, ramis divaricatis, foliis setaceis stipulatis internodio dimidio ferè brevioribus, sepalis lanceolatis obtusiusculis margine scariosis, petalis obtusissimis calyce brevioribus, pedunculis defloratis deflexis, seminibus subpyriformibus nigris. ① in Hispaniâ. Corolla purpurea, capsula 3-valvis valvis lanceolatis acutis. (v. s. ex herb. Thib.)

6. *A. RUBRA* (Lin. spec. 606.) caulibus prostratis pilosis, foliis filiformibus internodio brevioribus stipulatis, sepalis lanceolatis obtusiusculis margine scariosis, pedunculis defloratis deflexis, seminibus angulato-subpyriformibus rugosis immarginatis. ① in Europâ et Africâ. Flores rubri; Capsulæ 3-valves calyce vix longiores. (v. v.)

a. campestris (Lin. spec. 606.) pilosa. — in arvis sabulosis. Smith engl. bot. t. 852.

β. marina (Lin. spec. 606.) glabriuscula. — in maritimis salis. *A. marina* Roth, fl. germ. 1. p. 189. non Smith.

7. *A. SALINA* (Ser. mss.) foliis semi-cylindricis carnosis muticis internodio longioribus, stipulis membranaceis vaginantibus, capsulis oblongis calyce majoribus, seminibus compressis angulatis scabrisculis. ① in salsis pascuis Bohemiæ. *Spergularia salina* Presl. fl. cech. p. 93. *A. marinæ* simillima dicitur.

8. *A. GLUTINOSA* (Bieb. fl. taur. 1. p. 344.) villosa-viscosa, foliis oblongis obtusiusculis, caule erecto paucifloro, petalis obovatis calyce longioribus. ① prope Astracan. Planta pusilla facie Cerastii, flores magnitudine et formâ *A. rubræ*, petalis purpurascens. An species propria?

9. *A. SPERGULA* (Duf. ann. gen. 7. p. 293.) prostrata pubescens, foliis linearibus carnosus internodio longioribus stipulatis, pedunculis subsecundis tandem retroflexis, sepalis subacutis capsulâ brevioribus, petalis roseis calycis vix longitudine, seminibus limbo membranaceo cinctis. ①? in arvis humidis prope St.-Filipe. Media inter *A. rubram* et mediam.

10. *A. MEDIA* (Lin. spec. 606.*) caulibus prostratis, foliis semi-cylindricis carnosus internodia æquantibus stipulatis, sepalis lanceolatis obtusiusculis margine scariosis, pedunculis defloratis deflexis, seminibus alâ membranacâ cinctis. ① in pratis et sabulosis maritimis Europæ. *A. marina* Smith. fl. brit. 480* engl. bot. t. 958. *A. marginata*. DC. fl. fr. 4. p. 793. ic. rar. t. 48. *A. glandulosa* Jacq. hort. Schoenb. 3. t. 355. Affinis *A. rubræ marinæ* sed minus ramosa. (v. s.)

11. *A. GRANDIS* (H. B. et Kunth. gen. et spec. am. 6. p. 30. in ad not.) foliis verticillatis basi stipulatis internodio longioribus, pedunculis dichotomis, floribus secundis, seminibus alâ membranacâ albâ cinctis. — prope Monte-video. *Spergula grandis* Pers. ench. 1. p. 522. Poir. dict. 7. p. 305.*

12. *A. CANADENSIS* Pers. ench. 3. p. 504.) piloso-subhispida, foliis filiformibus internodio longioribus stipulatis, sepalis lanceolatis obtusiusculis margine lato scarioso, staminibus 5, pedunculis defloratis deflexis, seminibus subsemicordatis compressis, radiculâ prominulâ. ① in Canadâ. *A. rubra* ≠ Michx fl. bor. am. 1. p. 274. Capsula subglobosa calyce longior 3-valvis, valvis latè obtusis. (v. s. comm. à cl. Rafin.)

Sect. II. ARENARIUM. Ser. mss.

Folia graminea linearia lanceolata vel rotundata, exstipulata. Capsula 3-6-valvis.

§. 1. Foliis gramineis.

13. *A. GRAMINIFOLIA* (Schrader. hort. goett. t. 5. neu. journ. 1810. 2. p. 139.) caulibus erectis simplicibus, foliis subulato-filiformibus serrato-scabris, pau-

culâ trichotomâ pubescente laxâ, calycibus obtusissimis petalis obovatis subtriplo brevioribus. \mathcal{L} in campestribus Caucasi. Bieb. fl. taur. suppl. n. 852. A. filifolia Bieb. fl. taur. n. 852. p. 344. non Vahl. (v. s. comm. à cl. Desf.)

\mathcal{L} . *glaberrima*, paniculâ glabrâ minùs multiflorâ, floribus majoribus. A. graminifolia Willd. enum. p. 481. ex Bieb. l. c.

14. A. LONGIFOLIA (Bieb. fl. taur. i. p. 345. suppl. p. 308.) foliis subulato-filiformibus serrulato-scabris, caulibus erectis simplicibus, paniculâ trichotomâ glabrâ confertâ, sepalis ovatis obtusis petalis obovatis duplo brevioribus. \mathcal{L} ad Volgâ inferiorem. Gmel. fl. sib. 4. p. 157. n. 65. t. 63. f. 2. Facies A. graminifoliz quoad caulem et folia. Flores minores numerosiores, sepala angustiora carinata (Bieb.) An var. A. graminifoliz? (v. s. comm. à cl. Fisch. Stev. et Goldb.)

15. A. DAHURICA (Fisch! in litt.) foliis subulato-filiformibus serrulato-scabris, caulibus erectis simplicibus, paniculâ dichotomâ pauciflorâ, pedunculis calycibusque villosis glutinosis, sepalis lanceolato-linearibus nervosis petalis vix æquantibus. \mathcal{L} in Dahuriâ. (v. s. comm. à cl. Fisch.)

16. A. CUCUBALOIDES (Smith ic. ined. i. p. 17. t. 17.*) foliis subulato-filiformibus serrulato-scabris, caulibus erectis simplicibus, paniculâ dichotomâ laxiusculâ, sepalis lanceolatis acutis, petalis obovatis calyce duplo longioribus. \mathcal{L} in Armeniâ. Calyx viscosus.

17. A. OTITOIDES (Adams ex Fisch! in litt.) foliis subulato-filiformibus serrato-scabris, caulibus erectis simplicibus, paniculâ dichotomâ congestâ multiflorâ, sepalis lanceolatis carinatis, petalis obovatis vix longioribus, stylis longissimis exsertis. \mathcal{L} in Sibirîâ. Facies A. graminifoliz quoad caulem et folia, sed flores minimi numerosiores et densiores. A. androsacea Fisch. in litt. (v. s. comm. à cl. Fisch. Stev. et Deless.)

18. A. HOLOSTEA (Bieb. fl. taur. i. p. 345.*) foliis subulato-filiformibus margine scabriusculis, caulibus erectis simplicibus, paniculâ pauciflorâ, sepalis acuminatis nervosis, petalis obtusis brevioribus. \mathcal{L} in Caucaso Iberico. Flos Stellariz holostez similes, sepala dorso viridia, margine albicantia. Accedit ad A. dianthoidem, sed sepala lanceolata acuminata nec obtusa. (v. s. comm. à cl. Stev.)

19. A. RIGIDA (Bieb. fl. taur. i. p. 346.* suppl. p. 309.*) foliis lineari-setaceis ciliato-scabris, caulibus erectis simplicibus, paniculâ trichotomâ coarctatâ glabrâ, sepalis acutis subcristatis corollam vix superantibus. \mathcal{L} in Ucraniâ arenâ mobili. Bractez majores quàm in A. graminifol. et longifol; A. holostea Beaupré! in litt. (v. s. comm. à cl. Stev. Fisch. et Beaupré.)

20. A. FORMOSA (Fisch! in litt.) foliis lineari-setaceis ciliato-scabris, caulibus erectis simplicibus pedunculisque piloso-viscidis, paniculâ trichotomâ pauciflorâ, sepalis crassis obtusis, interioribus latissimis dorso glanduloso-pilosis petalis obovatis dimidio brevioribus. \mathcal{L} in Dahuriâ. Habitus et magnitudo Scheuchzeriz palustris. Flores ampli. (v. s. comm. à cl. Fischer.)

21. A. LYCHNIDEA (Bieb. fl. taur. i. p. 346. suppl. p. 309.) cespitosa, foliis angustissimis, radicalibus fasciculatis lineari-setaceis ciliato-scabris, caulibus erectis hispidulis, paniculâ 2-3-chotomâ pauciflorâ post anthesin divaricatâ, sepalis ovatis subacutis petalis obovatis duplo brevioribus. \mathcal{L} in Caucaso Iberico. A. airæfolia et A. setacea Fisch! in litt? Flores terni magnitudine Arenariæ Austriacæ. (v. s.)

\mathcal{L} . *glabra*, A. Laxmanni Fisch! in litt. (v. s.)

22. A. GMELINI (Fisch! in litt.) foliis linearibus obtusis, caulibus geniculatis, internodiis arcuatis folio duplo longioribus, paniculâ trichotomâ subcorymbosâ, sepalis lanceolatis viridibus margine scariosis, petalis obovatis calyce duplo longioribus. \mathcal{L} in monte Tagannai jugi Uralensis. — Gmel. fl. sib. 4. p. 144. t. 61. f. 1. Flores similes Gypsophylæ repentis. (v. s. comm. à cl. Fisch.)

23. A. CEPHALOTES (Bieb. fl. taur. i. p. 346.* suppl. p. 309.) foliis subulato-filiformibus ciliato-scabris, caulibus crassis erectis simplicibus, capitulo ter-

minali fasciculato hemisphærico, calycibus acuminatis corollam subæquantibus: \mathcal{U} in campestribus Tauriæ circa Tyraspolin. Bractææ striatæ. Species hujus generis elegantissima: (v. s. comm. à cl. Stev.)

24. *A. DIANTHOIDES* (Smith ic. ined. p. 16. t. 16.*) foliis subulato-filiformibus ciliato-scabris, caulibus erectis simplicibus, floribus capitatis, bracteis ventricosis, pedunculos superantibus, sepalis ovatis obtusis, petalis obovatis calyce triplò longioribus. \mathcal{U} in Armeniâ.

25? *A. GYPSOPHILOIDES* (Schreb. act. nov. acad. N. C. n. 6. 4. p. 139.* ex Willd. spec. 2. p. 723.) foliis linearibus, radicalibus setaceis, paniculâ subpubescente, sepalis ovatis mucronatis duriusculis, petalis lanceolatis calyce triplò longioribus \mathcal{U} in Oriente. Statura *A. saxatilis*.

26. *A. CAPILLARIS* (Poir. dict. 6. p. 380.*) foliis infimis cespitosus longissimis capillaribus, caule subnudo umbellato trifloro, pedunculis elongatis unifloris capillaribus, bracteis membranaceis ovatis acuminatis, sepalis ovatis obtusis margine membranaceis glabris, petalis ovatis subcrenatis calyce duplò longioribus. \mathcal{U} ? in Sibiria. *A. Sibirica* Pers. ench. 1. p. 504.

27. *A. PINIFOLIA* (Bieb. fl. taur. 1. p. 348.*) caulibus adscendentibus paucifloris pubescentibus, foliis setaceis rigidis, caulinis strictis, sepalis obtusis striatis villosis corollâ brevioribus. \mathcal{U} in Alpibus Caucasiciis. Calyx viridis striis pube confertâ brevi subviscidâ opliteratis.

28. *A. SUBULATA* (Ser. mss.) foliis setaceis rigidis mucronatis striatis, inferioribus fasciculatis, caulibus paniculatis paucifloris, sepalis lanceolatis corollâ multò brevioribus, capsulis ovatis calyce vix longioribus \mathcal{U}

a. *glandulifera*, caulibus pedunculis sepalisque piloso-glandulosis, paniculis 3-5-floris. — in regionibus Baicalensibus. *A. pungens* Steph. ex Stev! in litt. *A. viscosa* Fisch! in litt. *A. setacea* Adams ex Fisch! (v. s.)

ß. *glabrata*, glaberrima, floribus 1-3. — in Sibiria orient. (v. s.)

29. *A. PUNGENS* (Clem. in Lag. gen. et spec. 15. non Steph.) foliis subulatis calycibusque macronato-pungentibus, petalis calyce longioribus, stylis 3-4. \mathcal{U} in summis montibus regni Granatensis.

§. 2. Foliis subulatis linearibusve.

30. *A. SQUARROSA* (Michx. bor. am. 1. p. 273.) cespitosa, foliis imis squarrosos-imbricatis canaliculatis glabris, cauliculis simplicissimis oligophyllis, floribus paniculato-dichotomis erectis, sepalis ovato-subrotundis, petalis obovatis calyce triplò longioribus, capsulis ovalibus calycem superantibus trivalvibus, valvis obtusis. \mathcal{U} in sabulosis et pinetis Carolinæ (v. s. comm. à cl. Torrey.)

31. *A. IMBRICATA* (Bieb. fl. taur. 1. p. 344.* non Rafin.) foliis lineari-subulatis ciliatis confertis, caule repente hirsuto, pedunculis unifloris terminalibus petalis calyce obtusiusculo striato hirtò longioribus, capsulis trivalvibus. \mathcal{U} in Alpibus Caucas orient. Stev. in mem. soc. mosq. 3. p. 263. Folia brevia rigida patentia ciliata, in axillis fasciculata. Flores magnitudinis *A. laricifoliæ*. (v. s. comm. à cl. Fisch.)

32. *A. JUNIPERINA* (Lin. mant. 72.) foliis subulatis rigidis spinosis, inferioribus subfasciculatis, superioribus distantibus, caulibus erectis firmis, sepalis ovatis subuninerviis, petalis obovatis calyce subduplò longioribus, capsulis ovato-rotundis trivalvibus calycem vix superantibus. \mathcal{U} in Oriente, in Sibiria orientali. Smith ic. ined. p. 35. t. 35.* Poir. dict. 6. p. 378. *A. acicularis* Fisch! in litt. (v. s.)

33. *A. STRICTA* (Michx. fl. bor. amer. 1. p. 274.) erecta glabra multicaulis, foliis subulato-linearibus erectis, paniculâ pauciflorâ, sepalis ovali-lanceolatis, petalis conspicue striatis calyce multò longioribus. — in rupibus Novæ-Angliæ, Canadæ. Aff. *A. striatæ*, sed multò major.

34. *A. PROCERA* (Spreng. ex Horn. hort. hafn. p. 424.) caule erecto simplicel

paucifloro, calycibus obtusis petalis emarginatis triplò brevioribus. \mathcal{U}
Hab. †.

35. *A. LARICIFOLIA* (Lin. spec. 607.) foliis subulatis denticulato-ciliatis, caulibus adscendentibus 3-6-floris subscabris, calycibus cylindricis, sepalis oblongis obtusiusculis triplinerviis hirtis corollà duplò brevioribus, capsulis trivalvibus calycem superantibus. \mathcal{U} in Alpibus Helveticis, Gallicis, etc. (v. s.)

α. multiflora (Ser. mss.) caulibus debilibus multifloris, foliis junioribus fasciculatis arcuatis axillaribus, calycibus pedunculisque hirtis. *A. laricifolia* Vill. dauph. 4. p. 629. t. 47. f. 5. opt. *A. rostrata* Waldst. et Kit. ex Willd. enum. 481. *A. macrocarpa* Kit. ex hort. gott. (v. v. et s.)

β. striata (Ser. mss.) caulibus rigidis paucifloris, foliis rectis longis, pedunculis calycibusque hirsuto-viscosis. *A. striata* Vill! dauph. 4. p. 630. t. 47. f. 6. opt. *A. liniflora* Jacq. coll. 2. t. 3. f. 3. (v. v. et s.)

36. *A. LYCOPODIOIDES* (Willd. herb. ex Schlecht. berl. mag. 1816. p. 212.) ramosissima repens glabra, foliis subimbricatis mucronatis rigidis uninerviis, petalis calyce brevioribus, capsulà polyspermà. \mathcal{U} prope Moran Mexicanorum, alt. 1330 hex.

37. *A. ARCTICA* (Steud. in litt.) parvula cespitosa, foliis linearibus obtusis, caulibus 1-floris, pedunculis piloso glandulosis sepalis oblongis obtusis vix nervosis, petalis obovatis calyce duplò longioribus. \mathcal{U} in littore Sibirico maris glacialis. *A. bryoides* Fisch! in litt. *A. Altaica* Fisch! in litt. *A. serpens* Fisch! in litt. Habitus *Silenes acaulis* (v. s.)

38. *A. GLOBULOSA* (Labill. pl. syr. dec. 4. p. 6. t. 3. f. 1. *) ramosissima, foliis subulatis nervosis pilosis, floribus racemosis numerosis, sepalis inæqualibus petalis longioribus, capsulis profundè 3-valvibus, seminibus reniformibus ambitu rugosis. (I) juxta Tripolim Syriæ.

39. *A. AUSTRIACA* (Jacq. fl. austr. 3. p. 39. t. 270. *) foliis lineari-subulatis 3-nerviis, caule subpaniculato, pedunculis terminalibus longissimis binis puberulis, petalis obtusis emarginatis, sepalis 3-nerviis acutissimis patentibus. \mathcal{U} in Austria et Italia Alpibus. — All. pedem. n. 1700. t. 64. f. 2. *A. elongata* Schlecht. berl. mag. 1816. p. 200. *Stellaria biflora* Jacq. coll. 1. p. 251. t. 18. *A. stellarioides* Pers. ench. 1. p. 503. (v. s.)

β. glabra (DC! fl. fr. 4. p. 787. *) *A. Villarsii* Balb! misc. 21. excl. var. hirsut. *A. triflora* Vill. Dauph. 4. p. 623. t. 47. *A. mixta* Lapeyr. abr. 255. (v. s.)

40. *A. CAPILLACEA* (All. pedem. n. 1705. p. 365. t. 89. f. 2. *) foliis setaceis rigidis glabris, floribus erectis, petalis calyce majoribus. (I) in summo Pedemontii jugo Tendæ dicto. †.

41. *A. GRANDIFLORA* (Lin. spec. 608.) foliis subulatis latiusculis planis trinerviis ciliatis, radicalibus confertis, caulibus sub-unifloris, pedunculis longissimis pubescentibus, sepalis ovatis aristatis grossè 3-nerviis corollà duplò minoribus, capsulis ovatis 6-valvibus calycem vix superantibus. \mathcal{U} in petrosis Alpium et montium Occitanie. All. pedem. n. 1711. t. 10. f. 1. Cav. ic. 3. t. 249. f. 2. *A. mixta* Lapeyr. abr. p. 255.

β. multiflora, caulibus 3-4-floris, pedunculis longissimis, foliis vix ciliatis. *A. juniperina* Vill. dauph. 4. p. 624. — All. pedem. n. 1715. t. 26. f. 5.

γ. triflora, caulibus 2-4-floris, foliis angustioribus recurvis. *A. triflora* Lin. mant. 240. DC! fl. fr. 4. p. 788. — in montosis sabulosis Fontis-bleaudi, et monte Ventoso.

δ. stolonifera, caulibus ramosissimis, foliis inferioribus latiusculis brevibus densis, superioribus sepalisque angustissimis longissimis distantibus. — in Delphinatu et in rupibus ad Tschussowaja (Pall. herb.) *A. stolonifera* Vill! ex h. DC. *A. laxa* Fisch! in litt. (v. s.)

42. *A. HELMII* (Fisch! in litt.) glanduloso-pilosa, foliis linearibus obtusiusculis 3-nerviis, radicalibus confertis, caulibus 1-3-floris, pedunculis longissimis, sepalis ovatis submucronatis obsolete trinerviis, petalis obovatis

calyce duplò longioribus, capsulis ovatis calycem superantibus 3-valvibus. \mathcal{L} in montibus Uralensibus. Aff. *A. grandifloræ* sed distincta. (v. s. comm. à cl. Fisch.)

43. *A. MACROCARPA* (Pursh fl. bor. amer. i. p. 316. non Horn.) cespitosa, foliis congestis subulato-linearibus planis margine ciliatis, pedunculis terminalibus unifloris foliosis, sepalis linearibus, petalis ovatis calyce duplò longioribus, capsulis oblongis calyce triplò longioribus. \mathcal{L} in Americà bor.

44. *A. HIRTA* (Wormsk. in fl. dan. t. 1646.) foliis lineari-subulatis obtusis, bisuleis hirtis, caulibus hirtis 2-3-floris, calycibus 3-nerviis acutis capsulà brevioribus, petalis oblongis calyce subbrevioribus. \mathcal{L} ? in Groenlandià australi. Alsine rubella Wahlenb. fl. lapp. p. 128. t. 6. Alsine rubella Swartz. summ. veget. scand. p. 17. Aren. sulcata Schlecht. in berl. mag. 1816. p. 212. Planta minima, flores purpurei. (v. s. comm. à cl. Hornem.)

45. *A. VERNA* (Lin. mant. 72.) cespitosa multicaulis, foliis subulatis obtusiusculis nervosis, caulibus subpaniculatis elongatis, sepalis ovato-lanceolatis acuminatis striatis, petalis obovatis calycem vix æquantibus, capsulis cylindricis 3-valvibus. \mathcal{L} in submontanis Europæ. Smith engl. bot. t. 512. opt. — Vaill. bot. par. t. 2. f. 3. *A. divaricata* Adams ex Fisch! in litt. *A. flaccida* Schleich. (v. v. et s.)

β . *cespitosa*, caulibus valdè foliosis, calycibus pedunculisque glabriusculis. *A. cespitosa* Ehrh. herb. 55. DC! fl. fr. 5. p. 613. (v. s.)

γ . *minor* (Ser. mass.) caule nano unifloro.

46. *A. GERARDI* (Willd. spec. 2. p. 729.) erecta ramosa, foliis lineari-subulatis 3-nerviis, pedunculis geminis terminalibus unifloris, sepalis acuminatis margine membranaceis trinerviis. \mathcal{L} in pratis sterilibus Alpium, Gallopro. et Delphin. — *A. liniflora* Jacq. fl. austr. t. 445. ? — Ger. fl. Gallopr. 405. n. 7. t. 15. f. 1 ? An var. præcedentis ? †.

47. *A. RAMOSISSIMA* (Willd. enum. suppl. p. 24.) caulibus ramosissimis prostratis elongato paniculatis, foliis subulatis obtusiusculis nervosis recurvis, petalis ovatis calycem æquantibus, sepalis acuminatis striatis pedunculisque hispidis, capsulis 3-valvibus. \mathcal{L} in Hungarià. Hornem hort. hafn. p. 964. Link enum. i. p. 431. Aff. *A. vernæ* sed caulis ramosissimus procumbens, folia recurva uti calyx minus striata. (v. s. comm. ex hort. Gotting.)

48. *A. HISPIDA* (Lin. spec. 608.) cespitosa erecta multicaulis piloso-hispida, paniculà dichotomà, foliis subulatis sepalisque subtus hispidis vix striatis, corollà calycem superante, capsulis subglobosis trivalvibus calyce vix longioribus. \mathcal{L} circa Mospelium. Habitus *Spergula*; caules simplices pilis raris adpersi. DC! fl. fr. 4. p. 789.* (v. s. s.)

49. *A. ECHINATA* (Poir. dict. 6. p. 377.*) pumila, caulibus subramosis brevibus, foliis filiformibus subulatis vix pubescentibus, floribus subpaniculatis, pedunculis calycibusque hispido-glandulosis, corollà calycem æquante, capsulis longitudine calycis. — in Alpibus.

50. *A. FATULA* (Michx. fl. bor. amer. i. p. 273.) tota puberula, caulibus filiformibus paniculatis multifloris, foliis setaceo-subulatis patentibus, petalis submarginatis calyce acutissimo et striato paulò longioribus. — in rupibus Amer. bor. circa Knoxville. Habitus *A. tenuifoliæ*.

51. *A. SAXATILIS* (Lin. spec. 607.) foliis subulatis, caulibus pauculatis, sepalis ovatis. \mathcal{L} in Germanià, Helvetià, Gallià, Sibià. (Lin.) — Barr. ic. t. 580. †.

52. *A. PENDULA* (Waldst. et Kit. pl. rar. i. p. 90. t. 87.*) caulibus filiformibus radicanibus longissimis diffusis, ramis floriferis erectis paucifloris. foliis linearibus planis acutis subfasciculatis, sepalis lanceolatis acutis petalis brevioribus, capsulis depresso-globosis (sex-valvibus?) \mathcal{L} in rupibus calcareis Hungariæ. Filamenta staminum infernè villosa (v. s.)

53. *A. TENUIFOLIA* (Lin. spec. 607.) foliis subulatis setaceis, caule ramosissimo dichotomo, sepalis subulatis striatis, petalis multò longioribus, cap-

sulis 3-valvibus calycem vix superantibus. ① ad muros et in sabulosis. DC! fl. fr. 4. p. 789.* (v. v. et s.)

α. *Vaillantiana*, caulibus cespitosis strictis glabris supernè ramosis. — Vaill. bot. par. t. 3. f. 1.

β. *Barrelieri*, caulibus ramosissimis glabris subprocumbentibus. A. Barrelieri Vill. Dauph. 4. p. 634. — Barr. ic. t. 580.

γ. *simpliciuscula* (Ser. mss.) caule supernè ramoso glabro stricto, calycibus glabris, floribus 3-5-andris.

δ. *hybrida*, caule supernè ramoso glabro stricto, calycibus piloso-glandulosis viscidis. A. hybrida Vill. Dauph. 4. p. 634. t. 47. A. pentandra Duf. ann. gen. 7. p. 292. ex descript.

ε. *viscidula*, caule erecto supernè ramoso calycibusque hispido-viscidis. A. viscidula Thuill. fl. par. ed. 2. p. 219. A. dubia. Sut. fl. helv. 1. p. 266. A. viscosa Pers. ench. 1. p. 504.

54. A. *VILLOSA* (Ledeb. ex Steud. nom. 67. descript. ex Fisch. in litt.) foliis linearibus subulatis 3-nerviis ciliatis, caulibus pubescentibus villosis, sepalis ovato-lanceolatis acuminatis striatis 3-nerviis margine membranaceis corollâ longioribus, capsulis. . . . — in Sibiria Orient.

55. A. *CALYCINA* (Poir. voy. 2. p. 167. dict. 6. p. 370.*) glaberrima, foliis gramineis brevibus, pedunculis longissimis subunifloris, sepalis lanceolatis acutissimis margine membranaceis, petalis oblongis angustis calyce multò brevioribus, capsulis ovalibus calycem subæquantibus 5-valvibus. ① in Barbaria. Facies Stellarie gramineæ.

56. A. *SPINULIFLORA* (Ser. mss.) glaberrima, foliis filiformibus obtusis, floribus dichotomo-paniculatis numerosis minimis, sepalis lanceolatis membranaceo-pellucidis longè mucronatis, petalis brevissimis, stam. 5? capsulâ. . . . — in Oriente. (v. s. comm. à cl. Rousseau.)

57. A. *TRIANDRA* (Schrank. hort. monac. t. 30.*) caule dichotomo diffuso, foliis planis linearibus angustissimis, corollis calyce minoribus, floribus triandris, capsulis trivalvibus. ① hab. . . . Minuartia tenuifolia. Mart. hort. erlang. p. 44. ex Schrank. l. c. An var. A. tenuifolia? (v. s. in herb. Balb.)

58. A. *EMARGINATA* (Brot. fl. lus. 2. p. 202.) caulibus erectis, foliis linearibus obtusiusculis, sepalis lanceolatis, petalis emarginatis calyce brevioribus. ①? in collibus siccis in Transagana et Algarbiis. Caules racemosi viscosi; folia sessilia; petala rubella emarginata. An var. A. tenuifolia? Schlecht. in berl. mag. 1816. p. 212?

59. A. *MEDITERRANEA* (Ledeb. ex Link enum. 1. p. 431.) caule ramosissimo, foliis linearibus recurvis, sepalis longè acutatis striatis margine membranaceis corollâ multò longioribus, capsulâ. . . . calyce longiore. ① hab. . . . Habemus sub nomine A. calycinæ ex hort. hafn. et A. calycinæ. Pers. pedunculi longissimi sec. characterem. (Link. l. c.)

60. A. *RECURVA* (All. pedem. n. 1713. t. 89. f. 3.*) foliis radicalibus congestis recurvis subulatis subsecundis, caulibus cespitosis simplicibus subtrifloris, sepalis ovato-lanceolatis striatis pedunculisque piloso-glandulosis, petalis ovatis calycem æquantibus, capsulis ovatis 3-valvibus calyce brevioribus, seminibus subreniformibus vix punctato-rugosis. ② in pratis saxosis Alpium editiorum. Jacq. coll. 2. p. 244. t. 16. f. 1. sed fl. in icon. nimis magni. Variat caulibus 1-8-floris. (v. s. s.)

β. *hispidula* (Ser. mss.) caulibus foliis pedunculisque glanduloso-hispidulis. — in Valsesia. (comm. à D. Venetz.)

61. A. *HIRSUTA* (Bieb. fl. taur. 1. p. 349.) hirsuta, foliis subulatis trinerviis obtusiusculis, inferioribus imbricatis recurvis, caulibus declinatis, paniculâ multiflorâ dichotomâ, sepalis acutis subquinquennerviis hirsutis corollam subæquantibus, seminibus subdisciformibus margine echinatis. ② in Tauria montibus altis. An var. A. recurvæ? (v. s. comm. à cl. Fisch. Stev. et Beaupr.)

62. A. *CALYCVLATA* (Poir. suppl. 5. p. 7.*) subpubescens, foliis crassiusculis

subfasciculatis semi-cylindricis subulatis curvatis, sepalis lanceolatis elongatis acutiusculis 5-nerviis margine membranaceis, petalis integris, capsulis ovatis calyce longioribus. ♀ in Hungariâ?

63. *A. GLABRA* (Michx. fl. bor. am. 1. p. 274.) glaberrima erectiuscula multicaulis, foliis subulato-linearibus planis patulis, pedicellis unifloris elongatis divaricatis, sepalis ovalibus obtusiusculis lævibus corollâ brevioribus. — in rupibus Carolinæ sept.

64. *A. RUPESTRIS* (Labill. pl. syr. dec. 4. p. 8. t. 4. p. 1.*) caulibus adscendentibus, foliis setaceis obtusis subcurvatis congestis infra subciliatis, floribus paniculatis 2-4, sepalis ovato-oblongis subhirsutis trinerviis, petalis ovali-oblongis calyce duplò longioribus, disco sub ovario glanduloso, capsulis 3-valvibus calycem vix superantibus. ♀ in Libano.

65. *A. SETACEA* (Thuill. fl. par. ed. 2. p. 220.) caule ramosissimo, floribus paniculato-fastigiatis, foliis setaceis fasciculatis secundis basi ciliatis, calycibus ovato-acutis margine albo-membranaceis petala subæquantibus, capsulis 3-valvibus calycem superantibus. ♀ ad saxa in collibus circa Parisios et Fontisbleaudum.

α. pilosa (Ser. mss.) caule inferne pilosè. *A. heteromalla*. Pers. ench. 1. p. 504. *A. saxatilis* Loisel. fl. gall. p. 261. excl. syn. Lin. et Vaill. (v. s.)

β. rugosa (Ser. mss.) caule infernè rugoso. *A. heteromalla* Bieb. fl. taur. 1. p. 350.* (v. s. s. comm. à cl. Beaupr. Stev. et Fisch.)

66. *A. FASCICULATA* (Gouan. ill. 30.) foliis subulatis fasciculatis setaceis, caulibus erectis strictis subsimplicibus, floribus fasciculatis, pedicellis folio brevioribus, sepalis acuminatis submembranaceis bilineatis petalis subduplò longioribus, capsulis 3-valvibus calyce brevioribus, seminibus reniformibus rugosis margine subserratis. ① ② in sabulosis apricis Europæ. Jacq. fl. austr. t. 182? DC! fl. fr. 4. p. 731. *Alsine mucronata* Lam. dict. 4. p. 310. *A. fastigiata* Smith engl. bot. t. 1744. — Hall. hist. t. 17. (v. s.)

67. *A. FILIFOLIA* (Forsk. descr. p. 211. Vahl symb. 1. p. 33. t. 12.*) foliis setaceis subfasciculatis bistriatis, caulibus suffruticosis dichotomis, pedunculis terminalibus 1-2-floris, sepalis angustissimis corollam æquantibus, capsulis . . . ♀? in monte Boka arabie.

68. *A. MUCRONATA* (DC! fl. fr. 4. p. 791.*) foliis setaceis basi non ciliatis, caulibus cespitosis basi prostatis, paniculâ subdichotomâ, pedunculis folio longioribus, sepalis acuminatis submembranaceis bilineatis aristatis petala capsulasque 3-valves superantibus. ①? in saxosis Occitanie *A. fasciculata* ≠ *rostrata*. Pers. ench. 1. p. 504. *Alsine mucronata* Gouan. ill. 22. *A. mutabilis* Lapeyr. abr. 256. An var. *A. fasciculata*? (v. s.)

69. *A. GLOMERATA* (Bieb. fl. taur. 1. p. 350.* suppl. p. 311.) pubescenti-incana, foliis subulatis strictis, floribus capitato-glomeratis brevissimè pedicellatis, calycibus scariosis bistriatis acuminatissimis petalis multoties longioribus. ①? in Tauriæ saxosis calcareis. Aff. *A. fasciculata* (v. s. in herb. Balb.)

β. elatior (Ser. mss.) paniculâ paulò laxiore, pedicellis longioribus. — in saxosis circa Odessam.

70. *A. ULIGINOSA* (Schleich! cent. exs. 1. n. 47. DC! fl. fr. 4. p. 786. ic. pl. gall. p. 14. t. 46.) erecta basi ramosa glabra, foliis linearibus obtusiusculis, ramis supernè nudis, pedunculis geminis unifloris subterminalibus basi bibracteatis, sepalis lanceolatis enerviis petalis ovatis vix longioribus, capsulis ovatis calycem æquantibus trivalvibus, seminibus subreniformibus aurantiaciis. ♀ in paludosis turfosis Jurassi et Lapponiæ. *Spergula stricta* Swartz. act. Stockholm et in Schrad. journ. 1800. 2. p. 256. (v. s.)

β. purpurata (Ser. mss.) caulibus calycibusque purpureis. *A. muscorum* Adams ex Fisch! in litt. (v. s.)

71. *A. POLYGONOIDES* (Wulf. in Jacq. coll. 1. p. 241. t. 15.*) procumbens, foliis linearibus obtusis, pedunculis geminis ternisve unifloris basi bibracteatis,

sepalis oblongo-linearibus obtusis enerviis corollâ vix longioribus, capsulis ovatis calycem vix superantibus 3-valvibus, seminibus subdisciformibus atris. \mathcal{L} in pratis saxosis Helvetiæ. A. obtusa All! pedem. t. 64. f. 4. med. (v. s.)

β . *occulta*, pedunculis 1-floris brevissimis. — in Sibirîâ Altaicâ. A. *occulta* Fisch. in litt!

72. A. *BAVARICA* (Lin. amœn. 4. p. 315.*) foliis linearibus semi-cylindricis carnosius obtusis internodio longioribus, caulibus ramosis subdichotomis duris geniculatis, pedunculis terminalibus subbinis, sepalis... petalis... capsulis... \mathcal{L} in Baviâ et Sibirîâ. Saxifraga Bavarica. Rai hist. 1033. (v. s.)
73. A. *SAJANENSIS* (Willd. herb. ex Schlecht. berl. mag. 1816. p. 200.) foliis linearibus obtusiusculis, caulibus procumbentibus, pedunculis terminalibus brevissimis unifloris, calycibus obtusis uninerviis... — in montibus Orientalibus Altaicis. †.
74. A. *JUNCEA* (Bieb. fl. taur. suppl. p. 309.) caulibus erectis simplicibus, foliis lineari-setaceis ciliato-scabris, paniculâ dichotomâ pubescente, sepalis acutis villosis obscure trinerviis corollam subæquantibus. — in Tauriâ. †.
75. A. *VERTICILLATA* (Willd. spec. 2. p. 725.*) foliis subulatis rigidis spinosis floribusque verticillatis, pedunculis quadrifloris subcapitatis, sepalis lineari-subulatis pungentibus, petalis lanceolatis. \mathcal{L} in Armeniâ inter Ezeram et Tocat.
76. A. *FILIFORMIS* (Labill. pl. syr. dec. 4. p. 8. t. 3. f. 2.) caule simplici filiformi suprà nudo, foliis setaceis acuminatis, paniculâ subdichotomâ pauciflorâ, sepalis ellipticis glabris, petalis ovali-oblongis infrà attenuatis calyce duplò longioribus, capsulis 3-valvibus (calyce longioribus?) seminibus reniformibus. ① in insulâ Cypri.
77. A. *PHARNACEOIDES* (Ser. mss.) foliis filiformibus mucronulatis, subfasciculatis, caule ramoso, floribus subumbellatis, sepalis ovatis obtusis corollâ duplò brevioribus, capsulis ovatis 3-valvibus calyce vix longioribus. — in Oriente. Caules pedunculique partim piloso-glandulosi. (v. s. comm. à cl. Rousseau.)
- 78? A. *PICTA* (Sibth. et Smith fl. græc. t. 440, ex prodr. 1. p. 304.) caulibus aphyllis dichotomis supernè pilosis, foliis cespitosus subulatis, petalis emarginatis subtus venosis. ① in insulæ Cypri campestribus.
- 79? A. *MARGINATA* (Bieb. in hab. Willd. ex Schlecht. her. mag. 1816. p. 212. non DC.) caulibus ciliatis, foliis lineari-subulatis glabriusculis, calycibus acutis marginatis. \mathcal{L} in Sibirîâ orient. Parva, ferè lignosa, valdè ramosa.
- 80? A. *CANESCENS* (Vahl. herb. ex Horn. h. hafn. 1. p. 964. in add.) foliis subulatis carnosius, calycibus nervosis petalis duplò brevioribus, A. macrocarpa Horn. h. hafn. 1. p. 964. non Pursh.
- 81? A. *JURESSI* (Willd. herb. ex Schlecht. berl. mag. 1816. p. 212.) foliis lineari-subulatis subsecundis, sepalis lanceolatis. — in monte Gerez Lusitanîæ. Humilis, cespitosa.
82. A. *DECUSSATA* (Willd. herb. ex Schlecht. in berl. mag. 1816. p. 35.) ramosissima repens, ramis hirtellis, foliis lanceolato-linearibus acuto-mucronatis margine incrassatis uninerviis rigidis ciliatis, petalis calyce longioribus, ovario 4-5-spermo. \mathcal{L} prope Moran Mexicanorum, alt. 1330 hex. H. B. et Kunth. gen. et spec. 6. p. 34.* Capsulæ am. ign.
83. A. *NEMOROSA* (H. B. et Kunth. gen. et spec. am. 6. p. 35.*) ramosa procumbens puberula, foliis petiolatis oblongis acuto-mucronatis venosis membranaceis, petalis calyce brevioribus, capsulâ oligospermâ, seminibus lævibus nitidis. \mathcal{L} A. alsinoides Willd. herb. ex Schlecht. in berl. mag. 1816. p. 201. Stellaria pubescens Willd. herb. ex Kunth l. c. Semina lenticulari-reniformia fusco-nigra.

- α. *Quitensis*, internodiis longioribus, caulibus ramisque teretibus. — in nemoribus regni Quitensis.
- β. *Novogranatensis*, foliis oblongis semipollicaribus vel lanceolatis subpollicaribus, caulibus ramisque teretiusculis. — in locis temperatis Andium Novo-Granatensium, in Alto de Quilquase, alt. 1000 hex.
- 84? *A. FULCHRA* (Willd. herb. ex Schlecht. berl. mag. 1816. p. 212.) recta glabra, foliis lineari-subulatis glabris, floribus paniculatis, calycibus membranceis marginatis petalis brevioribus. — in monte Ararat. (Adams) Flores magni.
85. *A. ARMERINA* (Bory. ann. gen. sc. ph. 3. (1820.) p. 5.) caulibus diffusis lignosis, foliis connatis linearibus acutis rigidis vestustate recurvatis, floribus capitatis. ♂ in sepiibus prope Sierra Nevada Hispaniæ.
86. *A. EXTENSA* (Duf. ann. gen. 7. p. 291.) levissimè pubescens diffusa ramosa, caulibus elongatis, foliis linearibus, floribus laxè paniculatis, corollâ rosæa calycis longitudine, capsulâ obconicâ inclusâ. — in collibus aridis regni Valentini.
87. *A. RAPHINESQUIANA* (Ser. mss.) foliis imbricatis subulatis acutis, caulibus parçè ramosis suberectis, pedunculis solitariis terminalibus subnudis 1-3-floris, staminibus 8-10, sepalis. . . . petalis. . . . capsulis. . . . in New-Jersey. *A. imbricata* Rafin. journ. bot. 1. p. 229. Poir. suppl. 5. p. 8. non Bieb. (Rafin. l. c.) †.
88. *A. P. CHERLERIÆ* (Fisch! in litt.) cespitosa pygmæa, foliis lineari-subulatis apice curvatis? subimbricatis, floribus terminalibus vel subcorymbosis, sepalis oblongis obtusiusculis submembranaceis corollâ duplò longioribus, petalis angustis, capsula. . . . ①? †.
- α. *uniflora*, floribus solitariis terminalibus, caule subnullo, foliis apice curvatis. — in rupibus prope Grædina in Dahuriâ. Habitus Cherleriæ sedoidis. (v. s. comm. à cl. Fisch.)
- β. *fasciculata*, floribus subumbellatis, caule elongato, foliis non curvatis. — in Dahuriæ apricis frequentissima primo vere. (v. s. comm. à cl. Fisch. ex herb. Pall.)

§. 43. *Foliis lanceolatis ovalibus rotundatisve.*

89. *A. TETRAQUETRA* (Lin. spec. 605.) foliis ovatis carinatis recurvis marginatis quadrifariam imbricatis, caulibus strictis puberulis floribus subcapitatis, sepalis rigidis acutis carinatis ciliatis corollam subæquantibus, capsulis ovatis truncatis sexvalvibus, valvis apice callosis, seminibus reniformibus exasperatis. ♀ in sterilibus montosis Galliæ Mediterraneæ. DC! fl. fr. 4. p. 781.*
- α. *laxifolia*, foliis distantibus, capitulis multifloris, caulibus elongatis. All. fl. pedem. n. 1718. t. 89. f. 1. — Magn. monsp. p. 53. ic. (v. s.)
- β. *densifolia*, foliis dense imbricatis, capitulis paucifloris, caulibus abbreviatis. *A. imbricata* Lag. in litt. *Gypsophila aggregata* Lin. spec. 581. (v. s.)
- 90? *A. AMABILIS* (Bory. ann. gen. sc. phys. 3. (1820.) p. 5.) foliis triangularibus concavis acuto revolutis, caulibus ramoso-fastigiatis cespitosus internè denudatis, pedunculis subnullis foliosis, sepalis rigidis. . . . petalis. . . . capsulis. . . . — ad Sierra-Nevada Hispaniæ. An *A. imbricata*. Lag. aun. 5. p. 278?
91. *A. GIESEKII* (Hornem! fl. dan. t. 1518.) glanduloso-hispida, foliis lineari-lanceolatis rigidis trinerviis, pedunculis longissimis unifloris subnudis, sepalis lanceolatis acutis 3-nerviis corollâ longioribus, capsulis 3-rariter 6-valvibus calyce ferè duplò longioribus, seminibus reniformibus rugulosis. ♀ in Groenlandiâ. (v. s. comm. à cl. Hornem.)
92. *A. MUSCORUM* (Fisch! in litt. non Adams) cespitosa, foliis oblongis obtusis striatis ciliatis, caulibus brevibus 1-floris puberulis, sepalis ovatis obtusis substriatis

corollâ multò minoribus, capsulâ.... ♀ in Kamschatkâ. (v. s. ex herb. Pall. comm. à cl. Fisch.)

93. *A. modesta* (Duf! ann. gen. 7. p. 291.) pubescenti-viscidula erecta gracilis subdichotoma, foliis oblongo-linearibus, infimis ovato-ellipticis, pedunculis filiformibus demum divaricatis, sepalis lanceolatis acutis enerviis, petalis ovato-oblongis vix calycis longitudine, capsulis ovatis apice 6-valvibus calycem subæquantibus. (†) circa Valentiam (v. s. comm. à cl. Duf.)

94. *A. galioides* (Ser. mss.) gracilis piloso-hispida, foliis lanceolato-linearibus acutis tenuibus ciliatis fasciculato-subverticillatis, caulibus filiformibus, floribus dichotomo-paniculatis, pedunculis capillaribus elongatis, sepalis lanceolatis acutis enerviis piloso-hispidis margine submembranaceis corollâ vix longioribus, capsulis..... — Patr. ign. Habitus Galii Anglici, sed tenuior.

95. *A. lanceolata* (All! fl. pedem. n. 1715. t. 26. f. 5.*) cespitosa villosula, ramis adscendentibus, foliis lanceolatis angustis acutis rigidis nervosis, pedicellis foliis duplò longioribus, sepalis lanceolatis acutis nervosis corollâ vix longioribus, capsulis ovatis longitudine calycis trivalvibus, valvis apice callosis. ♀ in pratis lapideis Alpium Pedemontii. (v. s. comm. ex herb. All. à cl. Balb.)

♀. *cherlerioides* (DC! fl. fr. 4. p. 785.) parvula, caulibus reptantibus cespitosisque, foliis imbricatis. — *A. cherlerioides* Vill. Delph. 4. p. 626. t. 47. f. 1.

96. *A. densa* (Kit! ex Schlecht. Berl. mag. 1816. p. 200.) foliis oblongis lanceolatis scabris, caule cespitoso ramoso adscendente, ramis sterilibus, abbreviatis, floralibus subtrifloris, sepalis ovatis subscariosis petalis duplò minoribus. ♀ in Croatiâ. (v. s. in herb. Balb. comm. à cl. Kit.)

97. *A. purpurascens* (Ram! pyr. ined. DC! fl. fr. 4. p. 785.* et 5. p. 611. ic. pl. rar. p. 14. t. 45.) cespitoso-decumbens, ramis erectis 2-3-floris, foliis ovato-lanceolatis acuminatis glabris, pedicellis tomentosis folia vix superantibus, sepalis lanceolatis lævibus margine scariosis corollâ longioribus, capsulis ovato-cylindricis semi-sexvalvibus calycem superantibus, valvis acutis, seminibus reniformibus. ♀ in Pyrenæis editissimis. *A. cerastoides*. Pers. ench. 1. p. 502. Lapeyr. abr. p. 252. non Poir. (v. s.)

98. *A. violacea* (Ledeb. ex Fisch! in litt.) foliis lanceolatis glabrinaculis, floribus subpaniculatis erectis terminalibus, pedunculis piloso-viscosis calycem æquantibus, sepalis oblongis piloso-viscosis obtusis corollâ duplò brevioribus, petalis obovatis angustis, capsula..... — Hab. versùs Ochotek. (v. s.)

99. *A. montana* (Lin. amœn. 4. p. 272.* spec. 606.) pubescens, foliis lanceolato-linearibus, caulibus sterilibus longissimis procumbentibus, pedunculis terminalibus longis unifloris, fructiferis nutantibus, sepalis lanceolatis acuminatis corollâ multò brevioribus, capsulis ovato-globosis 6-valvibus calycem æquantibus, valvis obtusiusculis, seminibus reniformibus rugosis. ♀ in sterilibus montanis Galliæ occid. et Hispaniæ. DC! fl. fr. 4. p. 784.* Vent! hort. cels. t. 34. — *A. linearifolia*. Poir. dict. 6. p. 366.* (v. s.)

♀. *intricata*, foliis sublinearibus. *A. intricata*. Duf. in litt. — circa Valentiam.

100. *A. ruscifolia* (Poir. dict. 6. p. 365.*) foliis coriaceis ovatis spinulosi marginatis approximatis, floribus paniculato-dichotomis, sepalis ovatis pungentibus striatis corollâ ferè duplò brevioribus, capsulis ovatis calycem superantibus 6-valvibus, valvis angustis acutis. ♀? Hab. (v. s. comm. ex hort. matr. à cl. Vahl.)

101. *A. repens* (Ser. mss.) pilosa, foliis ovatis subsessilibus, caulibus repentibus, floribus solitariis axillaribus, pedunculis foliis brevioribus, sepalis lanceolatis acutiusculis margine membranaceis corollâ vix longioribus, cap-

solis ovatis 6-valvibus calycem æquantibus, seminibus reniformibus nitidis nigris. — *Alsine repens* herb. Deless. (v. s. comm. à cl. Deless.)

102. *A. WALLICHIANA* (Ser. mss.) glaberrima, foliis ovato-lanceolatis, caulibus valdè ramosis numerosissimis, sepalis lanceolatis acutis subpellucidis 1-nerviis corollà majoribus, capsulis ovatis profundè 6-valvibus calycem æquantibus, seminibus subpyriformibus rugosis. (1) ? in Napaniâ. Affinis. *A. serpyllifolia* sed distincta. (v. s. comm. à cl. Wallich.)

103. *A. BUXIFOLIA* (Poir. dict. 6. p. 362.*) pubescens, foliis ovato-oblongis sessilibus, caulibus repentibus, pedunculis dichotomis subbifloris, sepalis linearibus brevibus obtusis margine membranaceis, petalis calyce vix longioribus, capsulis ovatis obtusis calycem æquantibus (5-valvibus?) — in Canadâ. (Poir. l. v.)

104. *A. SERPYLLIFOLIA* (Lin. spec. 606.) foliis ovatis acutis sessilibus rugulosis ciliatis lævibusque, caulibus ramosissimis, sepalis lanceolatis acutis trinerviis viridibus opacis corollà ferè duplò majoribus, capsulis ovatis apice 6-valvibus calycem æquantibus, seminibus exactè reniformibus rugosis. (1) ad muros et in sabulosis arvis Europæ. Fuchs. hist. 23. ic. Smith engl. bot. t. 923. *Stellaria serpyllifolia* Scop. carn. n. 544. (v. v.)

β. *viscida* (Ser. mss.) caulibus nanis, foliis calycibusque nervosissimis viscidis. — in monte Pennino.

105. *A. CONIMBRICENSIS* (Brot. fl. lus. 2. p. 200. phyt. lus. p. 179. t. 73.) foliis carnosus lanceolatis enerviis, subtus glabris, suprâ pubescentibus, caule paniculato-dichotomo erecto, petalis calyce duplò longioribus, capsulâ ipso vix breviori apice sexvalvi, sepalis obtusiusculis enerviis corollâ brevioribus. (1) in solo sabuloso prope Conimbram.

106. *A. PUBESCENS* (d'Urville, enum. pl. Archipel. 50.) undiquè pubescens, foliis ovatis acutis subpetiolatis, caulibus patulis ramosis elongatis, sepalis acutis corollâ brevioribus, capsulis. 2℥ in summis montibus insul. Cois, altitud. 400. hexap. Aff. *A. ciliatæ* sed differt pube densâ in omnibus partibus sparsâ, sepalis non striatis et foliis angustioribus. (d'Urv. l. c.)

107. *A. CINEREA* (DC! fl. fr. 5. p. 811.*) piloso-cinerea, foliis ovato-lanceolatis acutis ciliatis, superioribus distantibus lanceolatis acutis, floribus paniculato-dichotomis longè pedunculatis, sepalis lanceolatis acutis subcarinatis corollâ ferè duplò minoribus, capsulis ovatis calycem æquantibus sexvalvibus, valvis callosis. 2℥ in Galliâ australi. (v. s. comm. à cl. Requier.)

108. *A. BREVICAULIS* (Sternb. in Spreng. pl. min. cog. 1. p. 31.) foliis oblongis acutiusculis trinerviis ciliatis subimbricatis, caule prostrato, sepalis lanceolatis acuminatis striatis petala superantibus margine membranaceis, capsulis. 5 ? in Alpibus Rhæticiis. *A. ciliatæ* similis, sed hujus caules erecti, folia uninervia obtusa, sepala minora, petala uninervia Spreng. l. c.

109. *A. SCABRA* (Poir. dict. 6. p. 377.*) foliis lanceolato-acutis patentibus duris scabris, caule simplici brevissimo, pedunculis paniculatis dichotomis, sepalis ovatis acuminatis striatis asperis margine membranaceis corollâ brevioribus, capsulis. . . — in Alpibus.

110. *A. CILIATA* (Lin. spec. 608.) foliis ovatis obovatisve bullato-rugosis raris pilosis plus minùsve nervosis ciliatisque, caulibus procumbentibus, sepalis lanceolatis acutis nervosis corollâ ferè duplò brevioribus, capsulis ovatis 6-valvibus calycem æquantibus. 2℥ in Alpibus et montanis excelsis Europæ. Smith engl. bot. t. 1745. Jacq. coll. 1. p. 245. t. 16. f. 1. — Hall. hist. t. 17. f. 3. (v. v.)

β. *multicaulis* (DC! fl. fr. 4. p. 783.*) foliis pulposo-subcoriaceis sepalisque vix nervosis. *A. multicaulis* Wulf. in Jacq. Coll. 1. p. 248. t. 17. f. 1. (v. s.)

111. *A. SCOPULORUM* (H. B. et Kunth gen. et spec. am. 6. p. 31.*) caulibus elongatis ramosissimis procumbentibus, foliis lanceolato-linearibus acutis uni-

- nerviis membranaceis margine dorsoque ciliatis, calyce glabro petalis brevioribus, capsulis 2-3-spermis, seminibus lævibus nitidissimis. \mathcal{L} in Andibus Peruvianis, alt. 1800 hex. A. digyna Willd. herb. ex Schlecht. berl. mag. 1816. p. 201.
112. A. SERPENS (H. B. et Kunth gen. et spec. am. 6. p. 32. *) ramosissima serpens glabra, foliis oblongo-subspathulatis obtusis subcarnosis aveniis basi ciliatis, petalis calyce vix longioribus, capsulis sub-15-spermis, seminibus lævibus nitidis. \mathcal{L} ad radicem montis Chimborazo, alt. 1640 hex.
113. A. RADDIANA (Ser. mss.) foliis ovato-lanceolatis basi attenuatis ciliatis, caulibus longissimis prostratis glabris quadrangulis, pedunculis axillaribus 1-3-floris, floribus apetalis. — in insula Maderâ. Sepala pellucida lanceolato-acuminata tribus nervis viridibus; bractæ duæ sepalis subsimiles. A. alsinoides Raddi brev. observ. p. 13. non Willd.
114. A. NORVEGICA (Gunn. fl. norv. 2. n. 1100. t. 9. f. 7-9, ex Hørnerm. fl. dan. t. 1269) glabra, caulibus subteretibus procumbentibus 1-2-floris, foliis spatulatis glabris, floribus terminalibus subglobosis, sepalis ovatis obtusis margine submembranaceis vix nervosis corollam subsequantibus, capsulis ovato-globosis calycem vix superantibus apice 6-valvibus. \mathcal{L} in Norvegiâ.
115. A. TRINERVIA (Lin. spec. 605.) caule gracili ramoso, foliis ovatis acutis petiolatis ciliatis nervosis, pedunculis longis post anthesin deflexis, sepalis lanceolatis acutis margine membranaceis corollâ multo longioribus, capsulis ovatis calycem æquantibus 6-valvibus, valvis apice circinnatis. (I) ? in nemorosis Europæ. — OEd fl. dan. t. 429. (v. v.)
116. A. LATERIFLORA (Lin. spec. 605.) caule gracili ramoso, foliis obtusis ciliatis nervosis, pedunculis lateralibus longis 1-2-floris, sepalis ovatis obtusis corollâ duplò brevioribus, capsulis ovatis obtusis calyce duplò longioribus (3-6-valvibus?) valvis obtusis. \mathcal{L} ? in Kamschatka. Aff. A. trinervia, sed distinctissima. (v. s. comm. à cl. Deless. ex herb. Patr. et Fisch.)
117. A. CERASTIIFOLIA (Ram! in DC! fl. fr. 4. p. 783. *) cespitosa subignosa ramosa, foliis ovato-lanceolatis 3-nerviis pubescentibus, pedunculis terminalibus unifloris piloso-glandulosis, sepalis lanceolatis obtusiusculis nervosis corollâ brevioribus, capsulis ovatis calycem vix superantibus 3-valvibus, valvis apice callosis. \mathcal{L} in rimis saxorum in Pyrenæis. A. Ramondi Poir. suppl. 5. p. 3. * (v. s.)
118. A. CALYCANtha (Ledeb. ex Fisch. in litt.) foliis oblongis acutis sessilibus basi ciliatis, floribus nutantibus, sepalis oblongis acutis margine membranaceis, petalis nullis (an semper), staminibus calyce longioribus, capsulis. (I) ? in Sibiria orient. (Fisch. l. c.)
119. A. ROTUNDIFOLIA (Bieb. fl. taur. 1. p. 343. *) foliis orbiculatis basi vix ciliatis, caulibus filiformibus procumbentibus, pedunculis lateralibus diphyllis subdichotomis, sepalis lanceolatis acutis margine membranaceis, petalis obovatis longitudine calycis. \mathcal{L} in Alpibus Caucasiciis.
120. A. BALEARICA (Lin. syst. nat. ed. 12. app. 230.) cespitosa repens, foliis ovatis lucidis subcarnosis ciliatis, pedunculis elongatis 1-floris, floribus cernuis, sepalis ovatis obtusiusculis 3-nerviis corollâ multo brevioribus, capsulâ ovatâ calycem vix superante 6-valvi, valvis apice subcircinnatis. \mathcal{L} in Corsicâ et ins. Balearicis. L'Her! stirp. 1. t. 15. A. muscosa Med. act. pal. 3. p. 202. t. 12. (v. s.)
121. A. BIFLORA (Lin. mant. 71.) foliis ovato-subrotundis glabris subimbricatis, caulibus procumbentibus ramosissimis, pedunculis bifloris lateralibus, alabastris subrotundis, sepalis ovatis obtusis corollâ brevioribus, capsulis ovatis longitudine calycis 3-valvibus, valvis apice callosis. \mathcal{L} in Alpibus editissimis prope nives. All. fl. pedem. n. 1699. t. 44. f. 1. et t. 64. f. 3. opt. (v. s. et v.)
- β . *apetala* (DC! fl. fr. 4. p. 782. *) foliis imbricatis ciliatis, floribus apetalis. — A. apetala Vill! Dauph. p. 622. t. 48. (v. s. comm. à cl. Vill.)

122. *A. PYSODES* (Fisch! in litt.) foliis acutis distantibus tenuibus, floribus terminalibus solitariis subnatis longe pedunculatis, sepalis lanceolatis acutiusculis corollam æquantibus, capsulis depresso-globosis (5-locularibus?) seminibus subpyriformibus 2? in Kamschatka. Merk. *Merkia physodes* Fisch? *Semina minor*. (v. s.)
123. *A. MUSCOIDES* (H. B. et Kunth gen. et spec. am. 6. p. 32. *) repens microphylla glabra, foliis subimbricatis lineari-lanceolatis acutiusculis enerviis parum carnosius, petalis vix longitudine calycis, capsulis oligospermis, seminibus lævisculis. 2? in summo monte Antisanâ, alt. 2300 hex. A. nana Willd. herb. ex Schlecht. berl. mag. 1816. p. 201. *Semina minutissima sublenticularia*.
124. *A. TETRAGYNA* (Willd. herb. ex Schlecht. in berl. mag. 1816. p. 201.) ramosissima repens glabra, foliis approximatis oblongo-lanceolatis acutiusculis obsolete uninerviis parum carnosius basim versus ciliatis, corollâ calycem superante, stylis plerumque 4, capsulis 4-valvibus 8-10-spermis, seminibus tuberculatis. ① in monte Antisanâ, alt. 2100-2200 hex. H. B. et K. gen. et spec. 6. am. p. 30. *
125. *A. BRYOIDES* (Willd. herb. ex Schlecht. berl. mag. 1816. p. 201.) caespitosa microphylla glabra, foliis quadrifariam imbricatis ovatis obtusius concavis enerviis crassis ciliatis, petalis calyce brevioribus, capsulis 1-2-spermis, seminibus lævibus nitidis. 2? in locis frigidissimis montis Tolucæ, alt. 1900 hex., in regno Mexicano. H. B. et K. nov. gen. et spec. am. 6. p. 33. *
126. *A. DICRANOIDES* (H. B. et K. gen. et spec. am. 6. p. 34. *) caespitosa glabra, foliis quadrifariam imbricatis ovato-lanceolatis acutis concavis uninerviis ciliatis, floribus terminalibus solitariis sessilibus digynis. 2? in monte Antisanâ, alt. 2100 hex. *Lobelia bryoides* Willd. herb. et mss. ex Schlecht in Roem. et Schult. syst. 5. p. 41. *Fructus ign.*
127. *A. PEPLIOIDES* (Lin. spec. 605.) foliis ovatis acutis carnosius approximatis, floribus solitariis breviter pedunculatis, sepalis oblongis acutiusculis corollam subæquantibus, capsulis crassissimis depressis (1-locularibus!) 3 valvibus calycem superantibus, seminibus maximis subpyriformibus. 2? in sabulosis maritimis Oceani. Smith engl. bot. t. 189. *Honkenya peploides* Ehrh. Beitr. 2. p. 181. — OEd. fl. dan. t. 624. (v. s.)
128. *A. SPATHULATA* (Desf! fl. atl. 1. p. 358. *) pubescens, foliis spathulatis basi ciliatis, caule erecto filiformi ramoso, floribus dichotomo-paniculatis, sepalis oblongis margine membranaceis puberulis corollâ multo brevioribus, capsulâ ovatâ calycem æquante (5 valvi?), valvis apice callosis. ① in arenosis Barbariæ. A. cerastoides Poir! dict. 6. p. 363. * non DC. nec Lapeyr. (v. s. comm. à cl. Desf. et Poir.)
129. *A. PROCUMBENS* (Vahl. symb. 1. p. 50. t. 33. *) tota pubescens, foliis lineari-lanceolatis, caulibus prostratis ramosissimis, sepalis lanceolatis obtusiusculis margine membranaceis corollâ paulo longioribus, capsulâ longitudine calycis (5-valvi?) semina minima. 2? in rudertis Toneti inque Ægypto. A. herniariæfolia. Desf. fl. atl. 1. p. 358. * A. geniculata Poir. dict. 6. p. 365. * Petala rubra.
- †. *Species non satis notæ.*
130. *A. ROSANI* (Ten. prod. p. 26. et cat. 1819. p. 43.) foliis lineari-lanceolatis mucronatis hirsutis striatis, caule erecto hirsuto subtrifloro petalis obovatis, calycibus striatis duplò longioribus. 2? in Lucaniâ. *Planta sesquipollicaris*. †.
131. *A. BARTOLOTTI* (Tineo pl. rar. sic. pug. 1. p. 10.) foliis lanceolatis subglabris, caule decumbente paniculato dichotomo acicpiti, paniculâ pubescente, petalis ovatis calycem subæquantibus, sepalis ovato-lanceolatis margine membranaceo. — in maritimis Siciliæ prope Gelam. (Tineo l. c.)
132. *A. MOLLUGINIFOLIA* (Pourr. ex Schlecht. berl. mag. 1816. p. 201.) foliis obovatis acutis, pedunculis bifidis cernuis. sepalis petalis capsulis — in Hispaniâ. †.

133. *A. OXYPTALA* (Sibth. et Smith fl. græc. t. 437. ex prodr. 1. p. 303.) foliis ovatis acutis petiolatis uninerviis, calycibus hirsutis obsolete 5-nerviis, petalis acuminatis. ① in agro Eiensis.
134. *A. THYMIFOLIA* (Sibth. et Smith fl. græc. t. 305, ex prodr. 1. p. 305. non Pursh.) foliis spathulato-linearibus, caule paniculato, petalis unguiculatis ovatis calyce trinervi longioribus..... capsulis..... ① in insulâ Cretâ.
135. *A. PURSHIANA* (Ser. mss.) foliis ovalibus subacutis, caule dichotomo diffuso, pedunculis alternis axillaribus solitariis elongatis, sepalis acutis..... corollâ longitudine calycis, capsulis..... ① in Labrador. *A. thymifolia* Pursh. fl. bor. am. 1. p. 317. non Sibth. et Smith. †.
136. *A. UMBELLATA* (Soland. in Russel Alep., ed. 2. p. 252.) foliis obovatis ciliatis, caule lævi, floribus umbellatis, petalis erosis, staminibus quinque cetratis, sepalis..... capsulis..... ① in Asiâ minore. Sibth. et Smith fl. græc. t. 439. Planta Holosteo umbellato aff.
137. *A. SERICEA* (Ser. mss.) foliis sericeis lanceolatis ciliatis, caule recto, ramis unifloris, corollâ calyce brevior, sepalis..... capsulis..... in Sibiria. (Bieb.) *A. purpurea* Willd. herb. ex Schlecht. berl. mag. 1816. p. 211. †.
138. *A. CLANDESTINA* (Portenschlag. pl. dalm. t. 1. ined.) foliis lineari-lanceolatis acutis, caule simplici 1-floro, flore magno nutante, pedunculo brevi, sepalis oblongis obtusis corollâ quintuplò brevioribus, petalis longissimis angustissimisque linearibus acutis, (staminibus o?) corpusculis ovoideis 5 inter ovaria et petala, ovario oblongo, stylis 3, stigmatibus furcatis. — in Dalmatiâ. An genus proprium? (descript. ex icon.) †.
- 139? *A. STELLARIOIDES* (Willd. herb. ex Schlecht. berl. mag. 1816. p. 209.) foliis..... ciliato-serrulatis, caulibus pubescentibus subpaniculatis, sepalis lanceolatis corollâ longioribus, capsulis.... — In Caucaso. †.
140. *A. AFFINIS* (Willd. herb. ex Schlecht. berl. mag. 1816. p. 198.) foliis oblongis acutis petiolatis, summis sessilibus, corollis calycæ brevioribus. †.

XXVI. CERASTIUM. Lin. gen. n. 797. Gærtn. fruct. 2. p. 231. t. 130. f. 6.

Calyx 5-partitus. Petala 5 bifida. Stamina 10. Styli 5. Capsula 1-locularis cylindrica aut globosa, apice dehiscens, dentibus decem circinnatis vel adscendentibus.

Sect. I. STREPHODON. Ser. mss.

Capsula cylindrica, dentibus circinnatis.

1. *C. PAUCIFLORUM* (Stev! in litt.) pilosum, foliis lanceolatis acutis, floribus paucis dichotomis longè pedunculatis nutantibus, flore solitario in dichotomiâ, petalis capsulisque calyce multò longioribus. — in Sibiria. (v. s. comm. à cl. Stev. et Fisch.)
2. *C. MEMORALE* (Bieb. fl. taur. suppl. p. 317.*) hirsuto-viscosum, caule erecto supernè dichotomo, foliis caulinis lanceolatis acutis, floribus axillaribus solitariis patentibus longè pedunculatis, petalis longitudine calycis, capsulis calyce vix longioribus. ① in nemoribus promontorii Caucasi. (v. s. comm. à cl. Fisch.)
3. *C. PERFOLIATUM* (Lin. spec. 627.) glabrum glaucescens, caule erecto ramoso vel simpliciter, foliis lanceolatis connatis obtusiusculis, floribus umbellatis, petalis calyce multò brevioribus. ① in Græciâ, Sibiria et Barbaria arenosa. — Dill. h. elth. 295. t. 217. f. 284. optima. (v. v. et s.)
4. *C. CAUCASICUM* (Fisch. in litt.) caule erecto dichotomo, ramis elongatis, foliis caulinis lanceolatis glabris margine scabris, petalis longitudine calycis glabri, capsulis pedicello pubescente brevioribus ①? ad Caucasum. *C. elongatum* Bieb. fl. taur. suppl. p. 316.* non Pursh. Flores magnitudine floris *Stellaris* holostear; capsula subcylindrica lata calyce paulò longior.

5. *C. FRIGIDUM* (Bieb. fl. taur. p. 362,* suppl. p. 320.*) villosum, caulibus basi ramosis adscendentibus dichotomo-subumbellatis, foliis lanceolatis acutis hirtis, sepalis lanceolatis acutis, petalis calyce multò longioribus, capsulis oblongis. ④ in Alpibus Caucasiacis. *C. purpurascens*. Adam. sp. Web. et Mohr. cat. 1. p. 60. Flos colore Lini sativi. (v. s. comm. à cl. Deless. ex h. Patrin.)
6. *C. DAHURICUM* (Fisch. in Spreng. plant. min. cog. 2. p. 65. Schrank. hort. (mon. t. 75, opt.) glabrescens, caule dichotomo, foliis cordato-ovatis amplexicaulibus, pedunculis fructiferis, longissimis deflexis, sepalis lanceolatis margine scariosis, petalis semibifidis calyce longioribus. ④ in Dahuriâ. *C. connatum*? Willd. ex Steud. nom. *C. amplexicaule*. Sims bot. mag. t. 1789.* opt. — Gmel. fl. sib. 4. p. 148. n. 49. t. 62. f. 1. (v. s. comm. à cl. Deless. ex h. Patrin.)
β. holosteuum, foliis lanceolato-linearibus amplexicaulibus subciliatis, caule pedunculisque subpilosis. *C. holosteuum*. Fisch! in litt. *C. nitens*. Stev! in litt. (v. s.)
- 7? *C. MAXIMUM* (Lin. spec. 629.) puberulum, caulibus diffusis, foliis lanceolato-linearibus acuminatis basi apiceque attenuatis, floribus maximis dichotomo-subumbellatis, petalis crenatis bilobisque, capsulis ovatis calycem subæquantibus. ④? in Sibiria. — Gmel. fl. sib. 4. p. 150. t. 62. f. 2. (v. s. comm. à cl. Patr. à cl. Deless. et à cl. Fisch.)
8. *C. STELLARIOIDES* (Moq. pl. nutk. ic. ined.) caule erecto dichotomo ramoso subtrifloro foliisque glabris oblongis acuminatis, pedicellis 1-floris terminalibus, sepalis lanceolatis, petalis semibifidis calyce duplò longioribus. ① ②. In America bor. circa Nutka.
- 9? *C. MOLLISSIMUM* (Poir. suppl. 2. p. 164.*) foliis amplexicaulibus lanceolatis acutis molliter pubescentibus, paniculâ subumbellatâ diffusâ. — in Peruvia. Aff. *C. perfoliato*. †.

Sect. II. ORTHODON. Ser. mss.

Capsulæ cylindricæ vel ovatæ, dentibus margine revolutis.

§. 1. *Petalis calycem æquantibus vel minoribus.*

* *Capsulis cylindricis calyce longioribus.*

10. *C. DICHOTOMUM* (Lin. spec. 628.) piloso-glutinosum, caule ramoso dichotomo, flore solitario in dichotomiâ, pedunculis petalisque calycem subæquantibus, laciniis calycinis lanceolatis acutis, capsulis longissimis erectis, foliis lanceolatis. ① Inter segetes Hispaniæ et Algeriæ. *Myosotis dichotoma* Mönch. meth. 225. *Alsine corniculata* Clus. hist. 2. p. 184. (v. v. et s.)
11. *C. RUDRALE* (Bieb. fl. taur. 1. p. 357,* suppl. p. 318.*) hirsutum, caule erecto dichotomo, foliis oblongo-lanceolatis obtusiusculis, floribus subumbellatis, pedunculis calyce multò longioribus, laciniis calycinis lanceolatis acutis, petalis longitudine calycis, capsulis pendulis calyce duplò longioribus. ① ad Caucasum. (v. s. comm. à cl. Fisch.)
12. *C. TAURICUM* (Spreng! ex herb. Balb.) hirsutum, caule erecto ramosissimo, foliis ovatis inferioribus basi attenuatis superioribus sessilibus, floribus dichotomo-paniculatis pedunculum æquantibus, petalis longitudine calycis, capsulis oblongis subattenuatis calyce duplò longioribus. ① culta in horti. Lugdunensi. Aff. *C. vulgato* sed caules ramosi, folia numerosa, flores multò minores. (v. s. in herb. Balb.)
13. *C. VULGATUM* (Lin. spec. 627.) hirsutum pallidè virens, caulibus erectis, foliis ovatis obtusissimis hirsutis, floribus dichotomis subumbellatis pedunculo longioribus, petalis calycem æquantibus, capsulis oblongis attenuatis calyce duplò longioribus. ① ② In ruderalis Europæ. Smith fl. brit. 2. p. 496.* engl. bot. t. 789. opt. DC! fl. fr. 4. n. 4395.* *C. viscosum* Bieb. fl. taur.

et suppl. n. 884,* cum syn. — Vaill. bot. par. t. 3o. f. 3. *C. barbulatum* Wahlenb. fl. carp. n. 446. *C. rotundifolium* Sternb. et Hopp. in mem. soc. ratisb. 1818. p. 113, ex Bieb. l. c. (v. v. et s.)

β. *glomeratum* (DC! fl. fr. 4. p. 776.) foliis obtusissimis, floribus umbellato glomeratis. C. ovale Pers. ench. 1. p. 521.

γ. *Americanum* (Ser. mss.) caule brevissimo paucifloro, foliis imbricatis. *C. pumilum* Rafin. in litt. non Curt. lond. *C. semidecandrum* Amer. auct. ex Raf.

δ. *tenellum* (Ser. mss.) minutissimum filiforme, caule 1-2 floro; in arenosis circa Genevam. *C. tenellum*. Gaud! fl. helv. mss. (v. s. comm. à cl. Gaud.)

14. *C. viscosum* (Lin. spec. 627.) hirsuto-viscosum saturatè virens, caulibus erectis, foliis lanceolato-oblongis, floribus dichotomo-subumbellatis, pedunculis petalisque calycem æquantibus, capsulis subpendulis teretibus calyce duplò longioribus. ① In ruderatis. Smith fl. brit. 2. p. 497.* engl. bot. t. 790. opt. DC! fl. fr. 4. p. 776.* *C. vulgatum* Bieb. fl. taur. et suppl. n. 883.* cum syn. C. sylvaticum Schleich. exs. — Vaill. bot. par. t. 3o. t. 1. C. obscurum Chaub. in St. Amans. fl. agen. p. 180. bouq. t. 4. f. 1. (v. v.)

15. *C. holosteoides* (Fries nov. fl. suec. III.) caule altero latere pubescente, pedunculis eglanduloso-pilosis, foliis oblongis glabris, petalis calyce margine scarioso brevioribus. ② In Blekingiâ Sueciæ. Hornem. hort. hafn. suppl. p. 138. (v. s. in herb. Balb.)

16. *C. pellucidum* (Chaubard in St.-Amans. fl. agen. p. 181. bouq. t. 4. f. 2.) villosio-viscosum pentandrum, caulibus erectis, foliis ovato-subrotundis, pedunculis 3-4-ve calyce longioribus, bracteis dichotomiæ universalis semimembranaceis pellucidis, petalis calyce longioribus. ① In arenosis versus Aginum. An var. *C. semidecandri*? †.

17. *C. SEMIDECANDRUM* (Lin. spec. 627.) hirsuto-viscidum, caulibus erectis, foliis ovato-lanceolatis, floribus pentandris dichotomo-subumbellatis pedunculis calyce longioribus, capsulis teretibus post florescentiam deflexis calyce duplò longioribus. ① In sabulosis et ad margines agrorum Europæ. Smith fl. brit. 2. p. 497, engl. bot. t. 1630. — Vaill. bot. par. t. 3o. f. 2. (v. s.)

β. *alsinoides* (DC. fl. fr. 4. p. 776.*) lobis calycinis margine apiceque scariosis. C. alsinoides Pers. ench. 1. p. 521. circa Montpelium. (v. s.)

18. *C. PENTANDRUM* (Lin. spec. 627.) procumbens subviscidum, foliis radicalibus spathulatis, caulinis ovali-orbiculatis, sepalis acuminatis petalis bidentatis longioribus, capsulis pedicello brevioribus. ① In Hispaniâ et in Tauriæ ruderatis. Bieb. fl. taur. 1. p. 359.* suppl. p. 319.*

18.*? *C. ANDROSACEUM* (Ser. mss.) pusillum pilosissimum, foliis ovatis caule dichotomo, floribus sub capitatis ternis, pedunculatis basi involucriatis, sepalis angustis acutissimis, petalis... capsulis... ① circa Constantinopolim: C. pilosum Castagne! in litt. non Horn. nec Ledeb. Habitus et ferè magnitudine *Androsac. villosæ*. (v. s. comm. à Castagne.)

19. *C. GRACILE* (Duf! in ann. gen. sc. ph. 7. p. 304.) erectum gracile dichotomum viscoso-pubescent, foliis inferioribus ovatis petiolatis, superioribus ovato-lanceolatis sessilibus, floribus solitariis distantibus, pedunculis flore vix longioribus, fructiferis deflexis, corollâ calycis longitudine, capsulâ exsertâ oblongâ, staminibus 5-10. ① In rupestribus Hispaniæ loco dicto la Sierra de Vernisa prope St.-Philippum. Aff. C. pentandro vel C. semidecandro. (v. s. comm. à cl. Dufour.)

20. *C. BRACHYPETALUM* (Desp! in Pers. ench. 1. p. 520.) caule erecto tomentoso dichotomo, foliis ovatis, floribus paniculatis, pedunculis flore longioribus, calyce villosio petalis longiore, capsulis calycem vix superantibus. ① In ruderatis circa Cænomanum et Argentoratum. DC! fl. fr. 4. p. 777. ic. pl. gall. t. 44.* *C. canescens*. Hornem ex Spreng. in herb. Balb. Caulis foliaque pilosissima. (v. s. s.)

21. *C. SPATHULATUM* (Pers. ench. 1. p. 520.) caule simplici subvillosio, foliis hirsutis, inferioribus obovato-spathulatis petiolatis, caulinis sessilibus subovatis, floribus glomeratis. ①? In Jamaicâ. Capsula calyce ferè longior. (v. s. in herb. Balb. comm. à cl. Bertero.)

22. *C. VULVUM* (Rafin. prec. p. 36, Journ. bot. 1814. p. 269.) fusco-villosum, caule erecto anguloso, foliis obtusis, floribus dichotomis, sepalis lanceolatis acutis, petalis calycem æquantibus, capsulâ nutante arcuatâ. ① In Pennsylvania et Virginia.
23. *C. MURALE* (Desp! in DC. fl. fr. 5. p. 609. *) hirsutum suberectum rigidum, foliis ovato-oblongis acutis numerosis, floribus fasciculatis longitudine pedicelli. petalis emarginatis calycem æquantibus, capsulis oblongis calycem vix superantibus ½? circa Cœnomanum. (v. s. comm. à cl. Desp.)
24. *C. DIFFUSUM* (Pers. ench. 1. p. 520.) caule ramosissimo villosulo opaco diffusum, foliis ovato-lanceolatis subhispidis, floribus dichotomo-paniculatis numerosis, pedicellis longitudine florum, petalis emarginatis calyce brevioribus, capsulâ obovatâ calycem æquante. ½? hab..... (v. s.) Habitus Stellariæ arenariæ.

* *Capsulis ovoideis calycem æquantibus vel brevioribus.*

25. *C. SERPYLLIFOLIUM* (Willd. enum. suppl. p. 26? Link. enum. 1. p. 433. *?) caule basi decumbente hirsuto-subviscoso, foliis lanceolatis basi attenuatis distantibus, floribus dichotomis breviter pedicellatis, flore solitario longèque pedicellato in dichotomiâ, capsulis ovatis calyce brevioribus. ½? in Sibiria *C. serpyllifolium*. Bieb. ex. Stev. in litt. 1817. †.
26. *C. LONGIFOLIUM* (Willd. spec. 2. p. 814, * non Poir.) caule erecto dichotomo piloso-viscoso, foliis lineari-lanceolatis, sepalis margine membranaceis, petalis calyce brevioribus, pedunculis fructiferis horizontalibus, capsulis longitudine calycis. ① in Armenia.
27. *C. COMMERSONIANUM* (Ser. mss.) caule dichotomo viscido angulato, foliis linearibus longissimis sessilibus viscidulis, floribus dichotomo-subumbellatis, sepalis lanceolatis acutis vix margine membranaceis, corollâ capsulâque longitudine calycis. . . — In Monte-Video. Commerson. *C. longifolium* Juss. ex Poir. supp. 2. p. 164. non Willd.

§. 2. *Petalis calycem superantibus.*

* *Capsulis calycem æquantibus.*

28. *C. MANTICUM* (Lin. spec. 629.) glaberrimum, foliis lanceolato-linearibus, caule stricto dichotomo, pedunculis longissimis, sepalis bracteisque lanceolatis acutis margine membranaceis corollâ brevioribus, capsulâ ovatâ calycem subæquante. ① in pratis montium Italiæ borealis et Hungariæ. Waldst. et Kit. pl. rar. hung. 1. p. 96. t. 96. *Stellaria Mantica*. DC! fl. fr. 4. p. 794. * (v. s.)
29. *C. RUPESTRE* (Fisch! in litt.) subpilosum, caulibus prostratis ramosis, foliis ovato-linearibus, floribus dichotomis, flore longè pedunculato in dichotomiâ, sepalis ovatis obtusis-margine membranaceis, petalis bilobis calycem multò superantibus, capsulâ ovoideâ calycem subæquante. ½? in locis uliginosis ad profluvia aquæ nivalis in Alpibus Sibiricis supra Tschala. (v. s.)
30. *C. ELONGATUM* (Pursh fl. bor. am. 1. p. 321, non Bieb.) hirsutum, foliis linearibus internodio longioribus divaricatis, pedunculis terminalibus elongatis 2-3 chotomis, bracteis ovatis, petalis emarginatis calyce acuto duplò longioribus, capsulis subglobosis. ½. In campis ad fl. Columbia.
31. *C. DEFLEXUM* (Ser. mss.) puberulum, caule elato dichotomo paniculato, foliis ovato-lanceolatis undulatis, floribus erectis parvis, petalis calycem superantibus, pedicellis capsuliferis deflexis, capsulis calycem vix æquantibus. ½? in Persia boreali. (v. s. comm. à cl. Fisch.)
32. *C. TENUIFOLIUM* (Pursh fl. bor. amer. 1. p. 321.) tenuissimè pubescens cespitosum, foliis angusto-linearibus internodio longioribus, floribus longè pedunculatis (deflexis?), petalis obovatis emarginatis calyce acuto subtriplo longioribus, capsulis ovoideis (calycem subæquantibus?) ½ in Pennsylvania. (v. s. comm. à cl. Rafin.)
33. *C. CAMPANULATUM* (Viv. annual. bot. 1. p. 2. p. 191. t. 1.) adscendens diffusum villosum, foliis radicalibus spathulatis, caulinis oblongis, pauci-

culâ dichotomâ, corollis campanulatis, petalis semi-bifidis calyce duplò longioribus, capsulâ ovoideâ calycem æquante. ① circa Romam. Sebast. rom. plant. fasc. 2. p. 12. t. 3. f. 1. *C. Ligusticum* Viv. cat. h. Diuagro. C. præcox. Tenor fl. nap. 1. p. 27. (v. s. comm. à cl. Viv.)

34. *C. INCANUM* (Ledeb. mem. acad. scienc. Potz. 5. p. 514.) erectum incanopubescent, foliis oblongo-linearibus acutis, pilis brevissimis adpressis, pedunculis 3-choromis, sepalis margine membranaceo cinctis, petalis calyce duplò longioribus spathulatis apice semi-bifidis, capsulâ globosâ inclusâ. ½ in Sibiariâ merid.

35. *C. GRANDIFLORUM* (Waldst. et Kit. pl. rar. hung. 2. t. 168.) caulibus repentibus, foliis linearibus acutis margine subrevolutis tomentoso-canis, floribus dichotomis, sepalis oblongis vix canis margine scariosis, petalis calyce duplò longioribus, capsulis oblongis. ½ in collibus siccis Hungariæ et Iberiæ. West in florâ 1820, p. 357. *C. argenteum*. Bieb. fl. taur. 1. p. 361.* suppl. p. 320. *C. incanum* Hoffm. h. mosq. ann. 1808, ex Bieb. l. c. Affine *C. tomentoso* sed differt superficie minùs canâ, foliis angustioribus acutis vix margine revolutis.

** *Capsulis calycem superantibus.*

36. *C. TOMENTOSUM* (Lin. spec. 629. var. β.) caulibus repentibus diffusis foliisque oblongis spathulatis tomentoso-canis pedunculis erectis dichotomis, sepalis oblongis tomentoso-canis margine scariosis, capsulâ subcylindricâ calyce longiore. ½ in montibus Gallo-provinciæ et Neapolitani agri. In hortis vulgarissimum ubi vulgò *oreille de souris* gallicè dictum. Col. phytob. ed. 1744. p. 115. t. 31. *C. Columnæ* Tenor. prod. p. 27. cat. app. p. 44. *C. tomentosum*. Lam. dict. 1. p. 680. (v. v.)

37. *C. WILLDENOWII* (H. B. et Kunth nov. gen. et spec. am. 6. p. 29.) caulibus erectis apice furcatis foliisque albido-lanatis ovato-lanceolatis acutis, calyce pubescente, capsulis oblongo-cylindraceis polyspermis subarcuatis calycem vix superantibus, seminibus fascis parvis. ½ in regno Quituensi. *Stellaria mollis* Willd. herb. ex Schlecht. berl. mag. 1816. p. 196. Flores magnitudinem *C. arvensis* æquantes.

38. *C. BIERBERSTEINII* (DC! in mem. soc. phys. gen. 1. p. ...) caulibus repentibus diffusis foliisque ovato-lanceolatis tomentoso-lanatis, pedunculis erectis dichotomis, sepalis oblongis tomentosis margine scariosis, capsulâ ovato-subcylindricâ calyce longiore. ½ in montium Tauriæ altiorum. *C. tomentosum* var. α. Lin. spec. 629? *S. repens*. Bieb. fl. taur. 1. p. 360. suppl. 320. non Lin. — Moris. ox. 2. s. 5. t. 22. f. 44? Differt à *C. tomentoso* foliis multò latioribus, flore fructuque majoribus. (v. s. comm. à cl. Stev. et v. v. in hort. gen.)

39. *C. FUSILLUM* (Ser. mss.) caulibus erectis subbifloris, foliis ovatis tenuiter tomentosis sessilibus, sepalis lanceolatis acutis pilosis, petalis. . . . , capsulis cylindricis calyce subtriplo longioribus, dentibus minutis. . . . — in Sibiariâ. (descrip. et ic. comm. à cl. Fisch.)

40. *C. LANATUM* (Lam. dict. 1. p. 680.*) caulibus prostratis cespitosis, foliis densè lanatis inferioribus subrotundis, superioribus ovatis, floribus dichotomis, sepalis lanceolatis margine scariosis, capsulâ ovato-cylindricâ calyce subduplò longiore. ½ in Pyrenæis et Alpibus. DC! fl. fr. 4. p. 778.* *Myosotis lanata* Moench. suppl. p. 308. (v. s.)

β. *Thomasianum* (Ser. mss.) caulibus foliis calycibusque glabris. — in valle Pyren. orient. dictâ Eynes. Ph. Thomas. (v. s.)

41. *C. IMBRICATUM* (H. B. et K. nov. gen. et spec. am. 6. p. 28.*) ramosissimum repens, foliis submembranaceis quadrifariam imbricatis subspathulato-oblongis obtusis utrinquè molliter villosis, capsulis oblongo-cylindraceis calyce piloso longioribus, seminibus subrotundis basi emarginatis fascis. ½ in summis montibus Catopaxi et Antisana, alt. 1900-2200 hex. Flores magnitudinè *C. vulgati*.

42. *C. FISCHERIANUM* (Ser. mss.) *hirsuto-anhviscosum*, *caulibus prostratis*, *foliis ovatis sessilibus distantibus*, *floribus dichotomo-subumbellatis*, *breviter pedunculatis*, *flore solitario longè pedunculato in dichotomiâ*, *sepalis lanceolatis obtusiusculis margine scariosis*, *petalis calyce duplò longioribus*, *capsulâ*. ? in Kamtschatkâ. *C. hirsutum* Fisch! in litt. non Tenor. *C. pilosum* Ledeb. ? OEd. fl. dan. t. 6. (v. s. comm. à cl. Fisch.)
43. *C. ALPINUM* (Lin. spec. 628.) *caulibus prostratis*, *foliis ovato-lanceolatis pilosis nudisve*, *floribus dichotomis paucis longè pedunculatis*, *sepalis oblongis obtusiusculis margine scariosis pilosis*, *petalis calyce duplò longioribus*, *capsulâ oblongâ calyce subduplò longiore*. ? in herbosis humidis Alpium, Pyrenæorum, etc. Smith engl. bot. t. 472. — Rat. syn. p. 349. t. 15: OEd. fl. dan. t. 6? Variat fl. magno minoreque. (v. s.)
44. *C. LITHOSPERMIFOLIUM* (Fisch! mem. soc. mosk. 3. p. 81.) *caule ramoso patulo*, *foliis lanceolatis acutis pubescentibus*, *floribus solitariis*, *petalis calyce duplò longioribus emarginatis*, *sepalis ellipticis*, *ovario globoso*? ? in Sibiria. (ic. comm. à cl. Fisch.)
45. *C. OVATUM* (Hoppe in Willd. enum. p. 493.) *subhirsutum*, *caulibus prostratis*, *foliis ovatis acutis glabris subciliatis*, *floribus terminalibus subcorymbosis*. ? in Alpibus Carinthiæ. *C. Carinthiacum* West? *C. Alpinum* Hoppe herb. viv? An *C. Alpini* variet.? †.
- β. filiforme* (Schleich. pl. exs.) *caulibus unifloris*, *pedunculis elongatis deflexis*. *C. pedunculatum*. Gaud. in litt. 1814. An spec. propria?
46. *C. LATIFOLIUM* (Lin. spec. 629.) *hirsuto-subviscosum*, *caulibus prostratis*, *foliis ovatis 1-rarò 3-floris*, *floribus terminalibus*, *pedunculo florem æquante*, *sepalis ovatis margine scariosis*, *petalis calyce duplò longioribus*, *capsulâ ovatâ turgidâ calycem superante*. ? in Alpibus Helveticis, Gallicis, Austriacis, in Angliâ, etc. Smith engl. bot. t. 473. Jacq. coll. 1. p. 156. t. 20. (v. v. et s. s.)
47. *C. GLACIALE* (Gaud. in litt. 1814.) *hirsuto-viscosissimum*, *caulibus cespitosus densis 1-floris*, *pedunculo florem æquante*, *foliis ellipticis ovatisve*, *sepalis ovatis margine subscariosis*, *petalis calyce duplò longioribus*, *capsulâ*. . . ? in Alpibus Helvetiæ editissimis prope nives. *C. uniflorum* Thom. exs. An var. *C. latifolii*? (v. v. et s.)
48. *C. SYLVATICUM* (Waldst. et Kit. pl. rar. hung. 1. p. 100. t. 97.) *diffusum repens*, *caule dichotomo-paniculato*, *foliis inferioribus ovatis*, *reliquis ovato-lanceolatis*, *floribus erectis longè pedunculatis*, *petalis subsemibifidis calyce duplò longioribus*, *capsulis calycem multò superantibus*. ? in vallibus sylvaticis Hungariæ et verosimiliter in Sibiria. *C. Sibiricum* Stev! in litt. (v. s. s.)
49. *C. ARVENSE* (Lin. spec. 628.) *caulibus declinatis*, *foliis lineari-lanceolatis obtusiusculis subpilosis*, *floribus dichotomo-paniculatis*, *pedunculis deflexo-pubescentibus*, *petalis calyce duplò majoribus*, *capsulâ oblongo-cylindricâ calyce duplò longiore*. ? in versuris agrorum et ad vias Europæ. Smith engl. bot. t. 93. — Vaill. bot. par. t. 30. f. 4. 5. *C. repens*. Lin. spec. 628? ex DC. fl. fr. 4. p. 778. (v. v.)
50. *C. STRICTUM* (Lin. spec. 629? DC! fl. fr. 5. p. 610.) *caulibus declinatis*, *foliis sublinearibus acuminatis glabris aut subhirsutis*, *pedunculis glanduloso-pubescentibus*, *petalis calyce duplò majoribus*, *capsulâ oblongâ*. ? in Alpibus. An *C. arvensis* var?
- a. suffruticosum*, *foliis angustissimis glabriusculis*. *C. suffruticosum* Lin. spec. 629. *C. laricifolium* Vill. delph. 4. p. 644. (v. s.)
- β. molle*, *foliis angustissimis hirsutis*. *C. molle* Vill! delph. 3. p. 644. (v. s.)
- γ. lineare*, *foliis lineari-lanceolatis elongatis acuminatis glabris*. *C. lineare* All. ped. 2. p. 365. t. 88. f. 4. (v. s.)
- δ. commune*, *foliis linearibus obtusiusculis*. *C. strictum* Lin. spec. 629? Centunculus angustifolius Scop. fl. carn. 1. t. 19. f. 1. *C. ambiguum* Fisch! in litt. (v. s.)

51. *C. PENNSYLVANICUM* (Hornem. h. hafn. p. 435.) caule prostrato foliisque lineari-lanceolatis pubescentibus, corollis calyce duplò longioribus, paniculâ longissimè pedunculatâ dichotomâ. \mathcal{P} in Pensylvaniâ Aff. *C. arvensi*, sed, ex cl. Horn., petala angustiora et capsula globosa? Au *C. stricti* var? (v. l. in herb. Balb.)
52. *C. NUTANS* (Rafin. prec. p. 36.) caule erecto viscido-pubescente, foliis lanceolato-linearibus, floribus subumbellatis longè pedunculatis, petalis calycem superantibus, capsulis calyce duplò longioribus nutantibus. ① in Pensylvaniâ. *C. longèpedunculatum* Muhl. cat. 1813. *C. glutinosum* Nutt. gen. am. 1. p. 291. non H. B. K. (v. s.)
53. *C. LEDEBOURIANUM* (Ser. mss.) caule erecto piloso, foliis oblongis obtusis subtus glaucis, floribus 1-3 erectis post anthesin nutantibus, petalis calyce triplò longioribus, sepalis obtusis margine membranaceis, capsulâ oblongâ calyce longiore. \mathcal{P} in Sibirîâ. *C. pilosum*. Ledeb. acad. scienc. petersb. 5. p. 514. n. 26.* non Hornem.
54. *C. FILIFORME* (Vest in florâ, 1820. p. 353.) caulibus cespitosis, foliis filiformibus trigonis carnosius, sepalis lanceolatis, petalis calyce duplò longioribus, capsulis oblongis calycem superantibus. \mathcal{P} in Styriâ superiore. Pedicelli paniculâ dichotomâ circiter adæquantes longitudinem calycis erecti etiam fructiferi. (Vest l. c.)
55. *C. RIGIDUM* (Ledeb. mem. acad. petersb. 5. p. 514. n. 25.) hirsutum, caule erecto infernè simplicissimo apice dichotomo, foliis oblongis acutis, pedunculis elongatis, sepalis lanceolatis acutis, petalis bifidis calyce longioribus, capsulis oblongis nitidis calyce duplò et ultrâ longioribus. \mathcal{P} in Sibirîâ. Tota planta pilis patentibus rigidis undiquè hirsuta; caulis ferè bipedalis strictus rigidus.
- 56? *C. GLUTINOSUM* (H. B. et Kunth gen. et spec. am. 6. p. 29.*) villosopilosum glutinosum fuscescens, caulibus adscendentibus apice dichotomo-ramosis, foliis lanceolatis angustato-acutis, capsulis cylindraceis subarcuatis calyce duplò longioribus, seminibus rugosis fuscis. \mathcal{P} in regno Novo-Granatensi. Petala biloba.

†. *Species non satis notæ.*

57. *C. BRACTEATUM* (Raf. prec. p. 36.) omninò pubescens, caule debili cylindrico, foliis oblongis ferè mucronatis, floribus erectis dichotomis bracteatis, bracteis ovatis acutis, petalis calyci ferè æqualibus, capsulis enerviis erectis. in Pensylvaniâ.
58. *C. PUBESCENS* (Goldie pl. canad. in edimb. phil. journ. april. 1822.) pubescenti-hirtum, caule deflexo piloso, foliis lineari-lanceolatis internodio longioribus, paniculâ terminali subquadriflorâ, petalis acutè emarginatis calyce duplò longioribus. — in Canadâ.
59. *C. MATRANSE* (Kit. in Spreng. pl. min. cogn. 1. p. 33.*) caule stricto trichotomo-paniculato pubescente, foliis lanceolato-linearibus pubescentibus revolutis basi subconnatis, sepalis lanceolatis obtusiusculis pubescentibus, petalis calyce longioribus. Hab. ad Matram Pannoniæ. Link. enum. 1. p. 434. Panicula terminalis, aphylla, dichotoma.
60. *C. DIOICUM* (Ait. hækew. ed. 1. v. 2. p. 120. ed. 2. v. 3. p. 137.) hirtum viscidum, foliis lanceolatis, floribus dioicis, petalis calyce triplò longioribus. \mathcal{P} in Hispaniâ. — Cult. anno 1766. in h. Oxford.
61. *C. ILLYRICUM* (Arduin. spec.) floribus subpentandris, petalis emarginatis, caulibus patentissimis, caule pilosissimo. ① in Peloponesi et insulæ Cyprî montibus. *C. pilosum*. Sibth. et Smith fl. græc. t. 454. ex prod. 1. p. 316. non Horn. Flores pentandri; apud Arduinum decandri.
62. *C. PILOSUM* (Horn. b. hafn. p. 965.) foliis lineari-lanceolatis obtusis piloso-lanatis margine reflexis, petalis calyce majoribus. \mathcal{P} ... Miss. ab illustriss. Schradero sub nomine. *C. lanati*. Pers. (Horn. l. c.) Link. enum. 1. p. 434.

63. *C. PALLASII* (Vest. in florâ, 1820. p. 356. *) foliis lanceolatis pubescentibus rigidis acutis, inferioribus internodia subæquantibus, summis longioribus, caule subuniiflora, petalis semibifidis. Hab. Flos majusculus.
64. *C. SPRENGELII* (Ser. mss.) foliis linearibus longissimis cauleque pubescentibus, pedunculis terminalibus umbellatis. (2) Hab. *C. tenuifolium* Spreng. in Horn. h. hafn. suppl. p. 133. non Pursh.
65. *C. FIMBRIATUM* (Ledeb. mem. acad. scienc. petersb. 5. p. 516. n. 27.) diffusum, caule angulato piloso, foliis lanceolatis acuminatis glabris ciliatis, sepalis oblongis æqualibus, petalis multifidis, capsulis globosis. ♀ in Sibiria.
66. *C. TENOREANUM* (Ser. mss.) diffusum pilosum, foliis ellipticis obtusis calycibusque corollâ longioribus ciliato-pilosis, pilis patentibus mollissimis, floribus paniculatis, capsulis oblongis. (1) in montibus Samnii et in Magellâ. *C. pilosum* Tenore cat. 1819. p. 44. non Hornem.
67. *C. SCARANI* (Tenor. prod. p. 27. cat. 1819. p. 44.) diffusum, foliis lanceolato-oblongis acutis utrinque tomentoso-viridibus, corollis calyce duplò longioribus, capsulis ovatis. pedunculis fructificantibus horizontalibus. ♀ in montibus Samnii.
68. *C. SAMNIANUM* (Ser. mss.) diffusum, ramis divaricatis, foliis lanceolato-linearibus tomentosis sublanato-virescentibus, paniculâ dichotomâ, corollis calyce duplò longioribus, capsulis oblongis. ♀ in montibus Samnii. *C. longifolium* Tenor. prod. fl. napol. p. 27. cat. hort nap. 1819. p. 451. non Willd. Juss. nec Poir.
69. *C. HIRSAUTUM* (Tenor. prod. p. 27. cat. 1819. p. 45. non Fisch.) diffusum hirsuto-viscosum, caulibus repentibus, foliis oblongis obtusis basi attenuatis hirsutis canescentibus, floribus paniculatis, corollis calyce duplò longioribus, capsulis oblongis subincurvis. ♀ in montibus Samnii.

XXVII. *CHERLERIA*. Hall. itin. helv. 1. Lin. gen. n. 775. Lam. ill. t. 379.

Calyx 5-sepalus. Petala 5 minima emarginata. Stamina 10. Styli 3. Capsula 3-locularis? 3-valvis, loculis 2-spermis? — Herbæ cespitosæ glabræ, foliis parvis confertis, floribus breviter pedicellatis pallidè virescenti-albidis.

1. *C. SEDOIDES* (Lin. spec. 608.) cespitosa parvula, foliis triquetris obtusiusculis nervosis patulis, capsulæ valvis obtusiusculis apice callosis calyce duplò longioribus. ♀ in pratis alpinis Europæ prope nives æternas. Smith engl. bot. t. 1212. DC! fl. fr. 4. p. 781. *C. cespitosa* Lam. fl. fr. 3. p. 46. (v. v.)
2. *C. IMBRICATA* (Ser. mss.) cespitosa parvula, foliis triquetris obtusissimis nervosis imbricatis, capsulæ valvis. ♀ ? in Carinthiæ et Tyrolæ Alpibus etiamque in Sibiria. Cherlerioides Hoppe? pl. sel. 2. (v. s. incompl.)

XXVIII. *SPERGULASTRUM*. Michx. fl. bor. amer. 1. p. 275. Micropetalon. Pers. ench. 1. p. 509.

Calyx 5-sepalus. Petala 5 minutissima integra aut nulla. Stam. 10. Stigmata 4 sessilia ligulato-setacea. Capsula ovata calyce longior 4-valvis — Herbæ boreali-Americane. Habitus Stellaris aut Spergulæ.

1. *S. LANUGINOSUM* (Michx. fl. bor. amer. 1. p. 275.) pubescens, foliis lanceolatis in petiolum angustatis, pedunculis subsolitariis longis demum reflexis, floribus apetalis. — in Americæ septentrionalis provinciis australibus. Micropetalon lanuginosum. Pers. ench. 1. p. 509. — Caulium lanugo tenuissima et densa.
2. *S. LANCEOLATUM* (Michx. fl. bor. amer. 1. p. 275.) glabrum, foliis lanceolatis utrinque attenuatis, floribus paniculatis, petalis ovalibus brevissimis. — in Americæ sept. provinciis borealibus. Micropetalon lanceolatum. Pers. ench. 1. p. 509. Interdum stigmata tria.

3. *S. GRAMINEUM* (Michx. fl. bor. amer. 1. p. 276.) glaberrimum, foliis linearibus erectis, floribus corollatis. — in Pensylvaniâ. Similis *Stellaris gramineæ*.

XXIX? HYDROPITYON. Gaertn. fruct. 3. p. 19. t. 183. f. 2.

Calyx 5-sepalus. Petala 5 ovato-rotundata. Stamina 10, filamentis crassis pilosisque, antheris cordatis. Ovarium oblongum. Stylus 1. Stigma orbiculatum. Capsula 1-sperma semen nudum mentiens. — *Herbæ aquaticæ Indicæ*, foliis verticillatis pinnatisectis, floribus axillaribus parvis. — Genus recognoscendum, forsân ab ordine removendum.

1. *H. ZYLANICUM* (Gaert. l. c.) foliis verticillatis pinnatisectis, caulibus arcuatis, floribus axillaribus sessilibus. — in Indiâ orient.
2. *H. PEDUNCULATUM*, foliis verticillatis pinnatisectis, caulibus strictis, floribus axillaribus longè pedunculatis. — in Indiâ orientali. *Hottonia Indica* Linn. sp. 208. ex syn. — Burm. fl. zeyl. p. 121. t. 55. Rheed, hort. Malab. 12. p. 71. t. 36.

(SEALING.)

Ordo XXIII. LINEÆ. DC. theor. ed. 1. p. 217.

Cal. sepala 3, 4 aut sæpius 5, basi vix imâ subcoherentia, per æstivationem imbricata cum pedunculo continua et ideò persistentia. Petala tot quot sepala, hypogyna, basi unguiculata, cum staminum annulo et interdum inter se subcoalita, sepalis alterna, per æstivationem contorta. Stam. petalorum numero æqualia, basi in annulum monadelpha, petalis alterna interjectis dentibus qui aut stamina abortiva aut filamentorum processus sunt habendi; Antheræ ovatæ, basi insertæ, biloculares, birimosæ. Ovarium subglobosum, loculis sæpissimè intus tot quot sunt sepala, rarò paucioribus. Styli tot quot loculi apice capitati. Capsula globosa, stylium basi persistente sæpius acuminata, constans carpellis marginibus induplicatis, apice bivalvi dehiscentibus, intus dissepimento incompleto è centro orto in concamerationes duas divisis. Semina in quoque carpello 2, aut in concamer. solitaria, ovata, compressa, nitida, inversa. Albumen rariùs parcum, sæpissimè nullum at hujus loco endoplevra tumida carnosâ. Embryo rectus planus carnosus-oleosus, radicula ad hylum sursum tendente, cotyl. ellipticis.

Herbæ aut suffrutices. Folia integra, exstipulata. Flores pedunculati. Petala caducissima. Ordo ferè medius inter Caryophyllas et Malvaceas Geraniaceasque.

I. LINUM. C. Bauh. Faill. Smith. fl. brit. 1. p. 342. Schult. syst. 6. p. 736.

Flores proportionè partium quinariâ. Sepala integra. Styli rarissimè 3, cum petalis staminibus sepalisque 5.

§. 1. *Floribus flavis.*

1. L. GALLICUM (Lin. sp. 401.) glabrum, sæpius multicaule, foliis alternis lineari-lanceolatis, floribus laxè paniculato-corymbosis, pedicellis calycis longitudine, sepalis basi ciliatis apice subulatis, petalis obtusis calyce duplò longioribus. ① in collibus à Galliâ ad Iberiam. — Ger. Gallo-prov. t. 16. f. 1. (v. v.)

β. *medium*, ramis subangulatis, foliis margine sub lente serrulato-scabris, circa Conimbram. Brot. fl. lus 1. p. 483. An potiùs ad seq. referendum?

2. L. AUREUM (Waldst. et Kit. pl. hung. 2. t. 177.) glabrum, erectum, foliis alternis lineari-lanceolatis, floribus laxè paniculato-corymbosis, pedicellis calyce duplò triplòve longioribus, calycibus basi subciliatis apice subulatis, petalis emarginatis calyce duplò longioribus. ① in Hungariæ et Croatiae graminosis. L. Liburnicum Scop. carn. ed 2. n. 385. Styli 3.

3. L. SETACEUM (Brot! phyt. n. 22. t. 6.) puberulum erectum, foliis alternis confertis lineari-subulatis subserrato-scabris, floribus paniculato-corymbosis, pedicellis calycis longitudine, sepalis subulatis basi ciliatis, petalis calyce duplò longioribus. ① in Lusitaniâ, Mauritaniâ. L. Broteri Willd. mss. in Schult. 6. p. 758. L. tenuifolium Schousb. (v. s.)

4. L. TENUE (Desf. atl. 1. p. 280. t. 81.) glabrum erectiusculum, foliis alternis

- linearibus acutis, floribus laxè paniculato-corymbosis, pedicellis vix calycis longitudine, sepalis mucronatis, petalis subretusis calyce quadruplò longioribus. ① in Algeriæ collibus. — *L. virgatum* Schousb. mar. r. p. 136. — *L. melianthum* Brot. fl. lus. r. p. 484. ex Link.
5. *L. LUTEOLUM* (Bieb. fl. taur. r. p. 256.) glabrum erectum, caule angulato, foliis alternis floralibusque oppositis linearibus acutis sepalisque basi biglandulosis serrulato-scabris, paniculæ ramis dichotomis, pedicellis brevissimis petalis calyce duplò longioribus. ① in Tauriæ et Iberiæ campis apricis. — Buxb. cent. 5. t. 59. (v. s.)
6. *L. NODIFLORUM* (Lin. sp. 401.) glabrum, foliis lanceolatis lævibus basi biglandulosis, inf. alternis, floralibus oppositis, paniculæ ramis subdichotomis, pedicellis brevissimis, calyce longitudine foliorum. ①? in Italiâ et Archipelagi ins. — Moris. ox. s. 5. t. 26. f. 11. Sibth. fl. græc. t. 307. ex prodr.
7. *L. STRICTUM* (Lin. sp. 400.) glabrum erectum, foliis lineari-lanceolatis strictis sepalisque acuminatissimis ciliato scabris, paniculâ corymbosâ confertâ, pedicellis brevissimis, petalis calycis longitudine. ① in Europæ australis et Africæ borealis agris et arenosis. — Lob. ic. t. 411 f. a. — *L. sessiliflorum* Lam. dict. 3. p. 523. (v. v.)
- β. *alternum*, (Pers! ench. r. p. 336) caule supernè evidentius corymboso. In Corsicâ. (v. s.)
- γ. *spicatum*, (Lam. l. c.) caule apice subspicato.
8. *L. RIGIDUM* (Pursh. fl. bor. am. r. p. 210.) foliis rigidè erectis linearibus brevibus, caule angulato sulcato, sepalis ovatis acuminatis trinerviis glanduloso, ciliatis, petalis oblongis angustissimis. — Ad flumen Missouri. Flores sulphurei.
9. *L. SELAGINOIDES* (Lam. dict. 3. p. 525.) foliis filiformibus alternis confertis, caulibus suffrutescis diffuso-procumbentibus, floribus terminalibus subsessilibus. ♂ in rupibus ad Monte-Video et in Chili. (v. s.)
10. *L. PROSTRATUM* (Lam. dict. 3. p. 525.) glabrum, foliis alternis ovali-oblongis, caulibus prostratis ramosissimis, floribus lateralibus solitariis brevissimè pedunculatis, sepalis obtusiusculis. — in Peruviz collibus siccis prope Limam. Folia femè Polygalæ vulgaris Flores ferè L. Gallici. (v. s.)
11. *L. VIRGINIANUM* (Lin. spec. 398.) glabrum erectum. foliis lineari-lanceolatis alternis, radicalibus ovatis, paniculâ laxâ corymbosâ, calycibus acutis, capsulis muticis. ① in agris et collibus à Noveboraco ad Virginiam. *L. Virginicum* Pursh. fl. am. r. p. 210. Flores vix L. Gallici majores. (v. s.)
12. *L. MEXICANUM* (H. B. et Kunth nov. sp. am. 6. p. 39.) glabrum erectum, foliis alternis, superioribus suboppositis, oblongis aut ovato-oblongis acutis basi rotundatis, ramis paniculatis, sepalis ovato-subrotundis acutis trinerviis ciliatis, capsulis acuto-mucronatis. ♀? in sylvis prope Santa-Rosa Mexicanorum.
13. *L. AQUILINUM* (Mol. chil. ed. germ. 126.) glabrum erectum, foliis alternis lanceolatis acutis, pedunculis bifidis, pedicellis calyce longioribus. ♀ in Chili montibus. — Feuill. per. 3. p. 32. t. 22. f. 2.
14. *L. CORYMBIFERUM* (Desf. atl. r. p. 279. t. 80.) glabrum erectum, foliis lanceolatis erectis alternis trinerviis retrorsum asperis, paniculâ laxâ subcorymbosâ, sepalis apice subulato-mucronatis, petalis calyce quadruplò longioribus. — in Atlante prope Mayane. Flores magni. L. usitatissimi.
15. *L. DAHURICUM* (Schult. syst. 6. p. 752.) glabrum erectum, foliis infer. lanceolato spatulatis confertis, caulibus lineari-lanceolatis remotis margine scabriusculis revolutis, paniculâ laxâ corymbosâ, pedicellis brevissimis, sepalis ovato lanceolatis, petalis calyce triplò longioribus. ♀ ♂ in Dahuriâ.
16. *L. MARITIMUM* (Lin. spec. 400.) glabrum erectum, foliis lanceolatis trinerviis alternis, infimis oppositis obtusis, paniculâ laxâ subcorymbosâ, sepalis ovatis breviter mucronatis, petalis calyce triplò longioribus. ♀ in Europæ

australis herbosis maritimis. *L. heterophyllum* Mœnch. suppl. 99. — Lob. ic. t. 412. f. 2. — *Trat. tab. t. 579.* (v. v.)

17. *LINUM GLANDULOSUM* (Mœnch. meth. 287.) glabrum erectum, basi suffruticosum, foliis basi biglandulosis lanceolatis alternis, infimis subspathulatis obtusis, sepalis acuminatis, petalis calyce triplò longioribus. \mathcal{L} \mathfrak{z} *L. campanulatum*. fl. fr. ed 3. v. 4. p. 797.

a. arboreum, (Lin. spec. 400.) fruticosum glaucum, floribus paucis subcapitatis, sepalis ovali-lanceolatis. \mathfrak{z} in Cretæ montibus. — Sibth! fl. gr. t. 305. ex prod. 1. p. 216. (v. s.)

ß. campanulatum, (Lin. sp. 400.) basi fruticosum glaucum, floribus corymbosis, sepalis lineari-lanceolatis. \mathfrak{z} in Europæ austr. præsertim Galliæ, et Tauriæ aridis (v. v.) Variat rarius in Tauriâ ramis foliisque pubescentibus.

γ. Tauricum (Willd. enum. 399.) basi fruticosum subvirens, floribus corymbosis, sepalis subserratis. \mathfrak{z} in Tauriâ. (v. s.)

δ. capitatum (Schult. syst. 751.) basi fruticosum, floribus capitatis, sepalis subserratis. \mathfrak{z} in Alpibus Croaticis.

ε. flavum, (Lin. sp. 399.) basi suffruticosum subvirens, floribus corymbosis, sepalis subserratis. \mathcal{L} \mathfrak{z} in Eur. austr. et Russiâ. *L. monopetalum* Steph. mosq. 214. Forsan hic pluræ species distinguendæ, sed characteres noti non sufficiunt. (v. v.)

18. *L. CESPITOSUM* (Sibth. fl. gr. t. 305 ex prodr. 216.) glabrum glaucum cespitosum basi suffruticosum, foliis obovatis acutis, sepalis ovatis obtusis. \mathcal{L} \mathfrak{z} in Cretæ montibus elatioribus. — *L. globulariæ-folium* Poir. suppl. 3. p. 445.

19. *L. TRIGYNUM* (Roxb. ex asiat. res. 6. p. 357.) glabrum fruticosum, foliis alternis ellipticis, utrinquè acuminatis, floribus bracteatis trigynis. \mathfrak{z} in Indiæ montibus. — Specim. sylvestre videtur vix palmare demissum, hortense erectum 2-3-pedale Sm. ex. bot. t. 17. Sims bot. mag. t. 1100. Delaun. herb. amat. t. 290. (v. v.)

20. *L. AFRICANUM* (Lin. mant. 360.) glabrum erectum basi fruticosum, foliis lineari-lanceolatis acutis alternis aut irregulariter oppositis verticillatisve, paniculâ corymbosâ erectâ, floribus secus ramos subsessilibus, sepalis subserrato-ciliatis. \mathfrak{z} ad cap. Bonæ-Spei. — Curt. bot. mag. t. 403. L'her! mss. (v. s.)

21. *L. ÆTHIOPICUM* (Thunb. fl. cap. 2. p. 143.) glabrum basi fruticosum erecto-patulum, foliis ovatis mucronatis oppositis. \mathcal{L} \mathfrak{z} ad cap. B.-Spei.

22. *L. QUADRIFOLIUM* (Lin. sp. 402.) glabrum erectum, foliis ovatis submucronatis quaternatim verticillatis, summis interdum oppositis. \mathcal{L} \mathfrak{z} ad cap. Bonæ-Spei., Curt. bot. mag. t. 431. An *L. quadrifolium* Thunb. fl. cap. 2. p. 244 et *L. tetraphyllum* Habenat. in herb. Panzer ab hoc diversa? (v. s.)

§. 2. *Floribus cœruleis roseisve, rarò variantibus albis.*

23. *L. VERTICILLATUM* (Lin. sp. 402.) foliis verticillatis, superioribus calycibusque villosis. ① in Italiâ circa Romam. — Barr. ic. t. 1226. Bocc. mus. t. 42. — Petala cœsia. Caulis pedalis. Folia lineari-lanceolata.

24. *L. STRIATUM* (Walt. car. 118.) foliis oppositis ovatis aut oblongis margine nervisque subdecurrentibus, floribus paniculatis terminalibus. — in Carolinâ. Petala parùm calyce majora. — Poir. suppl. 3. p. 443. (v. s. ex cl. Bosc.)

25. *L. VISCOSUM* (Lin. sp. 398.) foliis lanceolatis 3-5-nerviis alternis suboppositisque, cauleque hirsutis, mediis superioribus sepalisque margine pilos glanduloso-capitato gerentibus, stylis staminibus brevioribus aut subæqualibus. \mathcal{L} in Italiæ et Germaniæ australioris apricis. — Flores pallidè vinosi rarius cœrulescentes. — Bertol. am. it. p. 139. (v. s.)

ß. sylvestre (Scop. carn. n. 383. t. 11.) foliis ferè omnibus pilis glandulosis ciliatis et quasi serratis. — in Carnioliâ.

- γ. *Nestleri*, foliis infer. glabiusculis subbovatis. — in campis sabulosis Austriæ. (v. s.)
- δ. *hypericifolium*, (Salisb. parad. t. 79.) foliis ovato-oblongis distinctius 5-nerviis. — Sims bot. mag. t. 1048. — L. venustum Bot. rep. 477. — Flores ampli, ferè Malvæ.
26. L. *HIRSUTUM* (Lin. sp. 398.) foliis lanceolatis 3-5-nerviis alternis suboppositisque cauleque hirsutis, summis sepalisque margine pilos glanduloso-capitato gerentibus, stylis stamina duplò ferè superantibus. 2/ in Italiæ, Galliæ australis, Græciæ, Tauriæ, Caucasi, apricis. — Flores cærulescentes, rariùs subrosei aut albid. — Jacq. fl. austr. t. 31. — DC. fl. fr. ed. 3. v. 4. p. 800 et 5. p. 616. (v. v.)
- β. *ascyrifolium* (Sims bot. mag. t. 1087.) foliis basi subcordatis. — in Lusitaniâ. — Flores albi venis cæruleo-purpureis.
27. L. *NERVOSUM* (Waldst. et Kit. pl. hung. 2. t. 105.) caule basi piloso, foliis lanceolatis cuspidatis 3-5-nerviis glabris, paniculâ laxâ, sepalis subulato cuspidatis basi serratis. 2/ in Hungariæ, Ucraniæ, Tauriæ collibus. — Flores cærulei magni. (v. s.)
- β. *glabratum*, caule basi glabro. — ad Tanaïm. Goldbach. (v. s.)
28. L. *NARBONENSE* (Lin. sp. 398.) glabrum erectum, foliis lanceolato-linearibus acutissimis rigidiusculis, paniculâ subcorymbosâ, calycibus acuminatissimis basi ad marginem scariosis. 2/ in Hispaniæ, Galliæ australis, Italiæ apricis. — Flores ampli, pulchri cærulei, rarissimè albi. — Barr. ic. t. 1007. — (v. v.)
29. L. *USITATISSIMUM* (Lin. sp. 397.) glabrum erectum, foliis lanceolatis linearibusve, paniculâ corymbosâ, sepalis ovatis acutis margine membranaceis, petalis subcrenatis calyce triplò majoribus. ① in arvis Europæ. Tratt. tab. t. 744. L. arvense Neck. gallob. 159. L. sativum Blackw. herb. t. 160. (v. v.)
- β. *humile* (Mill. dict. n. 2.) petalis emarginatis, caule ramoso.
30. L. *MARGINATUM* (Poir. suppl. 3. p. 443.) glabrum erectum, foliis linearilanceolatis reflexis, calycibus albo-marginatis capsulisque mucronatis, petalis crenatis. 2/ Patr. ign. — L. angustifolium Willd. enum. 338. non Huds. L. affine Panz. mss. — L. usitatissimo simillimum, sed flores minores.
31. L. *REFLEXUM* (Ait. Kew. 1. p. 307.) foliis ovato-lanceolatis acuminatis reflexis levibus, calycibus acuminatis, filamentis connatis. 2/ in Europâ australi. An L. tenuifolio affinius? sed flores cærulei. †
32. L. *RUBRUM* (Rafin. car. 74.) glabrum, caule recto ramoso supernè angulato, foliis linearibus acutis 1-nerviis, pedunculis subcorymbosis striatis, sepalis ovatis acuminatis 3-nerviis margine scariosis. — in Siciliâ prope Trapanum, Agrigentum. Flores rosei. Affine L. usitatissimo.
33. L. *SQUAMULOSUM* (Rud. in Willd. enum. 338.) glabrum erectum, foliis linearilanceolatis erectis, inferioribus minoribus confertis, sepalis ovatis acutis 5-nerviis margine membranaceis, petalis calyce vix triplò majoribus. 2/ in Tauriâ et ad Tanaïm. Flores cærulei, interdum ex Rud. apetal. L. austriacum Bieb. fl. taur. 1. p. 254. non Lin.
34. L. *DIFFUSUM* (Schult. obs. 63.) glabrum diffuso-adscendens, caulibus ramosis, ramis patulis, foliis linearilanceolatis erectis acutis, sepalis ovato lanceolatis acutis 1-nerviis, petalis calyce duplò majoribus. 2/ Patr. ign. Cult. in hort bot. Flores pallidè cærulei. L. austriacum Lam. dict. 3. p. 521. (v. s.)
35. L. *ANGUSTIFOLIUM* (Huds. angl. 134.) glabrum multicaule suberectum, foliis linearilanceolatis acutis sub-3-nerviis, sepalis ellipticis sub-3-nerviis acutiusculis, petalis calyce vix duplò majoribus. 2/ in pascuis Angliæ, Galliæ, Italiæ; recepi etiam ex Asiâ et Novâ-Hollandiâ, at ex Europâ migratum? Flores pallidè cærulei. — Engl. bot. t. 381. (v. v.)
- β? *agreste* (Brot. fl. lus. 481.) capsulis ad dissepimenta lanuginosis, pe-

talibus albidis-purpureis. ② in Lusitania. — Huc referendum ex cl. Link.

36. *L. SIBIRICUM*, glabrum erectum elatum, foliis linearibus acutis patulis, sepalis ovalibus basi 5-nerviis, exterioribus acutiusculis, interioribus obtusissimis margine membranaceis, petalis integris calyce triplò quadruplò majoribus. ½ in Sibiria. L. perenne var. *Sibirica* Lin. sp. 397. Mill. ic. t. 166. f. 2. opt. L. *austriacum* Sims bot. mag. t. 1086. L. *Lewisi* Pursh. fl. bor. am. 1. p. 300 ex Nutt. — Flores ampli pulchrè cærulei. (v. s.)
37. *L. ANGLICUM* (Mill. dict. n. 5.) glabrum adscendens, foliis linearibus acutis erectis, sepalis ovalibus, ext. submucronatis, int. obtusis margine membranaceis, petalis emarginatis calyce duplò triplòve longioribus. ½ in Angliæ montosis cretaceis. — L. perenne var. *anglica* Lin. — L. perenne Sm. engl. bot. t. 40. Flores pallide cærulei.
- β. *procumbens* (Rai angl. 3. p. 362.) caulibus procumbentibus. ½.
38. *L. AUSTRIACUM* (Lin. mant. 359.) glabrum erectum, foliis linearibus sublanceolatisve acutis erectiusculis, ramis racemosis, pedicellis fructiferis deflexis, sepalis ovalibus obtusis basi 3-5-nerviis, petalis retusis triplò quadruplòve majoribus. ½ ? ② ? in Austriæ collibus. Jacq. fl. austr. t. 418. Petala pallide cæruleo-purpureascentia. (v. s.)
39. *L. MONTANUM* (Schleich! cat.) glabrum erectiusculum multicaule, foliis linearibus lanceolatisve acutis erectiusculis, floribus paniculato-corymbosis, pedicellis fructiferis erectis, sepalis ovalibus basi 3-nerviis, ext. acutis, int. obtusis margine membranaceis, petalis obtusis calyce triplò majoribus. ½ in Galliæ, Helvetiæ, Italiæ, Hungariæ, herbis montanis. L. *austriacum* DC. suppl. 615. L. *alpinum* fl. fr. 4. p. 615. L. perenne Lam! fl. fr. 3. p. 66. Bertoll. amen. it. 352. L. *læve* Scop. carn. ed. 2. n. 387. t. 11. L. *Narbonense* Sut. helv. 184. L. *Alpinum* β elatius. Wahl. carp. 299. Flores cærulei. (v. v.)
40. *L. ALPINUM* (Lin. sp. 1672.) glabrum multicaule decumbens, foliis linearisubulatis erectis, floribus paucis subcorymbosis, pedicellis fructiferis erectis, sepalis ovalibus basi 3-nerviis, ext. subacutis, int. obtusis margine membranaceis, petalis obtusis calyce triplò majoribus. ½ in Galliæ australis, Italiæ bor. et Austriæ siccis montanis. — Jacq. austr. t. 321. — Flores cærulei. (v. s.)
41. *L. DECUMBENS* (Desf! atl. 1. p. 278. t. 79.) glabrum multicaule decumbens, foliis linearisubulatis acutis erectis, floribus paucis subcorymbosis, pedicellis fructiferis erectis, sepalis basi ovatis membranaceis apice cuspidatis, petalis retusis calyce duplò longioribus. ½ in regni Tunetanis arvis. Cor. rosea. (v. s.)
42. *L. GRANDIFLORUM* (Desf! atl. 1. p. 278. t. 78.) glabrum basi ramosum erectiusculum, foliis linearilanceolatis acutis suberectis, floribus laxè paniculatis, sepalis lanceolatis acutis serrato-ciliatis capsulà sublongioribus. ½ ? in arvis prope Mascar. Petala magna rosea. (v. s.)
43. *L. TENUIFOLIUM* (Lin. spec. 398.) caulibus à basi ramosis erectis glabris, foliis linearisetaceis glabriusculis, sepalis lanceolatis acuminatis medio glanduloso-ciliatis capsulam superantibus, petalis calyce triplò longioribus. ½ in collibus aridis Europæ. Jacq. fl. austr. t. 215. Clus. hist. 1. p. 318. f. 2. — Petala sordide albo-carnea. (v. v.)
44. *L. SALSOLIOIDES* (Lam. dict. 3. p. 521.) caulibus basi fruticosis subtor-tuosis ramosis, ramis adscendentibus, sterilibus brevioribus, foliis linearisetaceis glabriusculis, sepalis ovatis acuminatis medio glanduloso-ciliatis capsulà subbrevioribus, petalis calyce triplò longioribus. ½ in Occitanis, Gallopr. et Vasconis aridis. — L. *suffruticosum* DC! fl. fr. 5. p. 616. non Cav. — Barr. ic. t. 795? Corolla præcedentis. (v. v.)
45. *L. SUFFRUTICOSUM* (Lin. spec. 400.) caulibus basi fruticosis ramosis foliisque linearibus acutis ex omni parte scabris, sepalis acuminatis medio glanduloso-ciliatis, petalis calyce quintuplò longioribus. ½ in aridis Kila-

paniæ. Cav! ic. 2. t. 108. *L. tenuifolium* Assol syn. arr. 41. *Linum* Bory. ann. gen. 3. p. 102 — A priore differt habitu canescente, petalis oblongis, etc. (v. s.)

§. 3. *Floribus albis, foliis oppositis.*

46. *L. CATWARTIGUM* (Lin. sp. 401.) glabrum erectum, foliis oppositis obovato-lanceolatis, caule supernè dichotomo. ① in Europæ pascuis. — Schkuhr. handb. 1. t. 87. Blackw. herb. t. 368. (v. v.)

†. *Species non satis notæ.*

47. *L. MONOGYNUM* (Forst. prod. n. 145.) caule suffruticoso, foliis alternis lineari-lanceolatis 3-nerviis. 5 in Novâ-Zeelandiâ.

48. *L. HISPANICUM* (Mill. dict. n. 7) caule paniculato procumbente, foliis sparsis lineari-lanceolatis, calycibus acutis. 24 in Hispaniâ. Flores ferè *L. usitatissimi*.

49. *L. BIKENNE* (Mill. dict. n. 8.) caule ramoso, foliis alternis linearibus, calycibus patulis acuminatis. ② in Istriâ. Flores *L. usitatissimi*.

50. *L. BICOLOR* (Schousb. mar. 135.) caule simplicissimo, foliis margine scabris, corollâ luteâ fundo cæruleâ striis purpureis. ① prope Tanger. — Nequit esse var. *L. tenuifolii* ob colorem floris et radicem annuam.

51. *L. PALLASIANUM* (Schult. syst. 6. p. 758.) pubescens, foliis linearibus acutis subcanis, sepalis glabriusculis acutis margine albo lacero. 24 in Chersoneso Heracleotico. — *L. pubescens* Willd. herb.

52. *L. PUBESCENS* (Russ. alep. ex Schult. syst. 6. p. 758) caule tereti, foliis floralibus oppositis lanceolatis pilosis, floribus alternis subsessilibus, sepalis longitudine foliorum pilosis ciliatis. — in Aleppo.

53. *L. OLIGOPHYLLUM* (Willd. mss. in Schult. l. c.) caulibus ramosis striato-scabriusculis, foliis linearibus acutis squamiformibus, floribus solitariis, sepalis ovatis acuminatis. — Patr. ign.

54. *L. TRINERVIUM* (Roth. nov. sp. 187.) glabrum, foliis alternis linearibus trinerviis, sepalis obtusis tricostatis. — in Indiâ orient. — Flores ign.

II. *RADIOLA*. *Gmel. syst. 1. p. 289.*

Flores proportionè partium quaternariâ. Sepala ad medium ferè coacta, apice trifida.

1. *R. LINOIDES* (Gmel. l. c.) ① in Europæ sabulosis humidis. *R. millegrana*. Sm. engl. bot. t. 895. *Linum radiola* Lin. spec. 402. — Vaill. par. t. 4. f. 6. Herba pusilla dichotoma floribus numerosis minimis. Petala calyci æqualia forsân ut squamæ stamineæ habenda. (v. v.)

Ordo XXIV. MALVACEÆ. *Brown cong. p. 8. Kunth diss. 1822. p. 1. — Malvaceæ §. 1. 2. 3. Juss. gen. 271.*

Calyx constans sepalis 5, rarissimè 3-4, basi plus minùs coactis, per æstivationem valvatis!, sæpè extùs gerens bracteas seu sepala exteriora involucrum calycemve exteriorem constituentes. Petala tot quot sepala et iis alterna, hypogyna, inter se æqualia, per æstivationem spiraliter contorta, nunc omninò distincta, sæpiùs imò staminum tubo adnata. Stamina numero petalorum æqualia aut sæpiùs multiplicia, definita aut sæpiùs indefinita, hypogyna; filamentis in tubum coalitis, inæqualibus, exterioribus brevioribus; antheris unilocularibus reniformibus rimâ transversali dehiscentibus. Ovarium carpellis plurimis sæpiùs circa axim verticillatis, interdum congestis, rariùs liberis, ferè semper coalitis constans; styli tot quot carpella nunc distincti, nunc in unum coaliti; stigmata tot quot carpella plus minùs distincta. Carpella nunc 1-2-sperma intùs rimâ dehiscentia, nunc polysperma loculicido-dehiscentia aut medio septifera, septo ad latus interius semina gerente; nunc sublibera, nunc in capsulam multilocularem, nunc in baccam anomalam connata. Semina ovata aut subtriquetra, intereà epidermide villosa gerente indusiata. Albumen nullum. Embryo rectus, dicotyledoneus, radiculâ tereti, cotyledonibus chrysa-loideo-contortuplicatis.

Herbæ, frutices aut arbores. Folia alterna sæpiùs petiolata, dentata aut lobata. Villi sæpiùs ramoso-stellati. Stipulæ 2 ad latera foliorum. Pedunculi axillares uni aut multiflori, interdum foliis subevanidis spicato-racemosi.

Div. I. *Calyx duplex seu involucri cinctus.*

I. MALOPE. *Lin. gen. n. 843. Lam. ill. t. 583.*

Calyx cinctus involucri 3-phylo, foliolis cordatis. Carpella plurima 1-sperma in capitulum adgregata. — Herbæ floribus purpuræscens aut albis.

1. MALACOIDES (Lin. sp. 974.) foliis ovatis crenatis, stipulis oblongo-linearibus, pedunculis axillaribus 1-floris. ① in Italiæ, Galloprovinciæ, Hispaniæ, Mauritanie, insulæ Scio pratis. — Cav. diss. t. 37. f. 1. (v. v.)

♀. *sinuata*, foliis nonnullis obtusè subtrifidis sinuatis pinnatifidisve. — in Mauritaniam. — Cav. diss. t. 27. litt. x. Media inter malacoidem et stipulaceam.

2. M. STIPULACEA (Cav. ann. cienc. nat. 3. p. 74.) foliis ovatis crenatis, stipulis cordatis ovato-acutis, pedunculis axillaribus 1-floris. ① circa Mogador. Broussonet! (v. s.)

3. M. TRIFIDA (Cav. diss. 2. p. 85. t. 27. f. 2.) foliis trinerviis trifidis dentatis

glabris, lobis acuminatis, pedunculis axillaribus. ① in pratis Brasilie prope Carmonam et in Mauritaniâ. (v. s.)

4. *M. MULTIFLORA* (Trig. in Cav. diss. 2. p. 85.) foliis subrotundis crenatis villosis, floribus 3-4 axillaribus. ① in Bæticâ. Flores albi. †.

II. *MALVA*. Lin. gen. n. 841. Lam. ill. t. 582.

Calyx cinctus involucro 3-phylo rariùs 5-6-phylo, bracteolis oblongis setaceisve. Carpella capsularia plurima in orbem disposita.

Sect. I. *MALVASTRUM*.

Carpella 1-locularia 1-sperma.

- §. 1. *Chrysanthæ*, floribus flavis, foliis indivisis. — Flores in axillis superioribus subsessiles et interdum foliis evanidis spiciformes.

1. *M. TRICUSPIDATA* (Ait. h. Kew. ed. 2. v. 4. p. 210.) foliis oblongis ovatisve acutis serratis, floribus axillaribus glomeratis, carpellis tricuspidatis. ② in Jamaicâ *M. Americana* Cav. diss. 2. t. 22. f. 2. *M. carpinifolia* Desrous. in dict. enc. 3. p. 754. *M. Coromandeliana*. Willd. Sw. *Sida Jamaicensis* Mill. (v. v.)

β. *subtriloba*, foliis subtrilobis. *M. Antillarum* Zucc. obs. n. 79.

2. *M. AMERICANA* (Lin. sp. 968.) foliis ovatis acutis crenato-serratis pilosiusculis, floribus axillaribus subsolitariis, terminalibus capitato-spicatis, carpellis muticis. ① in Sancto-Domingo. *M. ulmifolia* Balb. herb. *M. Curassavica* Desrous. in dict. enc. 3. p. 754. (v. v.)

3. *M. SCABRA* (Cav. diss. 5. t. 138. f. 1.) foliis ovato-lanceolatis duplicato-dentatis obsolete subtrilobis, subtus ramisque pilis stellatis scabris, pedunculis axillaribus subbifloris. ③ in Peruviae locis aridis. *M. scoparia* Jacq. ic. rar. t. 139. *M. corchorifolia* Desrous. in dict. enc. 3. p. 743. *M. Lagascae* cat. h. taur. 1821. p. 36? Variat floribus subsessilibus. (v. s.)

4. *M. SCOPARIA* (L'her. stirp. t. 27.) foliis ovatis crenato-serratis subtus ramisque pube stellatâ molliter subvelutinis, floribus axillaribus confertis. ③ in Peruvia. — Cav. diss. 2. t. 21. f. 4. (v. s.)

5. *M. BORBONICA* (Willd. enum. 728.) foliis ovatis acutis grossè inæqualiter dentatis stellato-pubescentibus, superioribus basi cuneatis, floribus axillaribus et ad apices ramorum spicatis. ⑤ in insulis Mauritiæ et Borboniæ. (v. s.)

6. *M. POLYSTACHYA* (Cav. diss. 5. t. 138. f. 3.) foliis ovato-acuminatis serratis scabris, floribus axillaribus et ad apices ramorum spicatis, capsulis 12 glabris muticis. ③ in Peruvia. *M. betulina* Desrous. in dict. enc. 3. p. 754. (v. s.)

7. *M. TIMORIENSIS*, foliis ovatis dentatis ramisque stellato-scabris, floribus in spicas longas cylindraceas digestis, capsulis 8 glabris muticis. ⑤ in insulâ Timor. (v. s.)

8. *M. SPICATA* (Lin. sp. 967.) foliis ovatis subcordatisve pube stellatâ supernè scabris subtus tomentosis, floribus in spicas ovatas digestis, capsulis 14 glabris muticis. ⑤.

α. Foliis ovatis. — in Brasiliâ. *M. ovata* Cav. diss. 2. t. 20. f. 2.

β. Foliis subcordatis. — in Jamaicâ. *M. spicata* Cav. diss. 2. t. 20. f. 4. *M. sublobata* Desrous. in dict. enc. (v. s.)

9. *M. SUBHASTATA* (Cav. diss. 2. p. 72. t. 21. f. 3.) foliis ovatis cordato-subhastatis obsolete subtrilobis grossè dentatis ramisque hirsutis, pedunculis solitariis axillaribus petioli longitudine. ⑤ in Brasiliâ et Peruvia. (v. s.)

10. *M. TOMENTOSA* (Lin. sp. 967.) foliis cordatis crenatis obtusis ramisque tomentosis, floribus lateralibus congestis. ⑤ in Indiâ. — Pluk. am. t. 356. f. 12 (v. s. in herb. Balb.)

11. *M. WATHERIFOLIA* (Link. enum. 2. p. 209.) foliis subcordatis acutis den-

tatis subtus tomentosis, floribus sessilibus, calycis lobis ovatis, involucri foliolis angustissimis. 2 $\frac{1}{2}$ in Javā. Folia subtus canescentia. Corollā calyce major.

12. *M. TRACHELIOLIA* (Link. l. c.) foliis cordatis acuminatis serratis scabris, inferioribus lobatis, pedunculis axillaribus, floribus capitatis. ① Patr. ign. Folia longè petiolata. Cal. hirtus. Cor. calyce major.
13. *M. GANGETICA* (Lin. sp. 967.) foliis cordatis obtusis glabris, floribus sessilibus glomeratis, carpellis 10 muticis crenulatis. ① in Indiā. Pluk. t. 74. f. 6. Aff. *M. tricuspidatæ*.

14. *M. DOMINGENSIS* (Spreng! in h. Balb.) nama, foliis ovatis dentatis, adultis subglabris, junioribus petiolis ramulisque pilosis, floribus axillaribus solitariis breviter pedicellatis, carpellis hispidis tricuspidatis. 3 $\frac{1}{2}$ in Sancto-Domingo. Bertero. (v. s.)

15. *M. SUBTRIFLORA* (Lag. nov. gen. et sp. 21.) foliis cordatis angulato-subhastatis crenatis, pedunculis axillaribus brevissimis tri-automultifloris, carpellis 8-9 glaberrimis muticis. 5 in Novā-Hispaniā. Aff. *M. spicatæ* et *scopariæ*.

§. 2. *Cymbalarix*, floribus purpurascens aut albis, pedicellis axillaribus 1-floris solitariis, calyce externo 3-phylo, foliis subrotundis, caule herbaceo.

16. *M. LEPROSA* (Ort. dec. 8. p. 95.) foliis reniformibus latè crenatis ramisque albido-leprosis, caulibus prostratis, pedunculis petiolo longioribus. 2 $\frac{1}{2}$ in insulā Cuba. (v. v.)

17. *M. SHERARDIANA* (Lin. sp. 1675.) foliis orbiculatis cordatis velutino-tomentosis crenatis, caulibus prostratis, pedicellis arcuatis petiolo longioribus. 2 $\frac{1}{2}$ in Bithyniā. — Cav. diss. 2. t. 26. f. 4. (v. s.)

18. *M. CYMBALARIEFOLIA* (Desrous. in dict. enc. 3. p. 753.) foliis cordato-subrotundis obsolete 5-lobis crenatis subglabris, caule decumbente? subviloso, pedicellis petiolo longioribus. — Patr. ign. †.

19. *M. CRETICA* (Cav. diss. 5. t. 138. f. 2.) foliis cordatis subrotundis quinquangularibus crenatis villosis, pedicellis petiolo longioribus, caule erecto. ① in Cretā.

§. 3. *Bibracteolatæ*, floribus purpurascens aut albis, calyce exteriore 2-phylo, pedicellis axillaribus 1-floris, caule herbaceo.

20. *M. HISPANICA* (Lin. sp. 970.) foliis semi-orbiculatis crenatis, summis subrhombeis, caule erecto hirtio. ① in Hispaniā et Mauritiā. — Desf. fl. atl. 2. t. 170. (v. s.)

♀. *spithamea* (Cav. diss. 2. t. 18. f. 3.) foliis minimis subrotundo-crenulatis.

21. *M. STIPULACEA* (Cav. diss. 2. t. 15. f. 2.) foliis inferioribus trilobis integerrimis, superioribus multifidis, laciniis trifidis apice dentatis, stipulis lanceolatis petiolo longioribus. ① in Hispaniā. Asso. arr. t. 5. f. 1.

22. *M. CUNEIFOLIA* (Cav. diss. 2. t. 20. f. 1.) foliis obovato-cuneiformibus apice truncatis repandis, pedicellis petiolo longioribus. — Patr. ign.

23. *M. PAPAVER* (Cav. diss. 2. t. 15. f. 3.) foliis 3-5-palmatisectis, segmentis integerrimis linearibus ciliatis. — in Lusitiā. Flores ferè *Papaveris Rhæadis*.

24. *M. ÆGYPTIA* (Lin. sp. 971.) foliis tripartitis, laciniis trifidis ciliatis apice dentatis, caulibus erectis, corollis calyce ferè minoribus. ① in Ægypto, Mauritiā, Hispaniā. — Cav. diss. 2. t. 17. f. 1. *M. diphylla* Mœnch. (v. v.)

25. *M. TRIFIDA* (Cav. diss. 5. t. 137. f. 2.) foliis tripartitis, laciniis trifidis linearibus obtusis, caulibus erectis, corollis calyce triplō majoribus. ① in Hispaniā. (v. s.)

§. 4. *Bismalvæ* (Medik.) *floribus purpurascentibus aut albis, pedicellis axillaribus solitariis unifloris, involucro 3-phylo, foliis multipartitis, caule herbaceo.*

26. *M. TOURNEFORTIANA* (Lin. sp. 971.) foliis multipartitis, lobis trifidis linearibus, caule decumbente, pilis subnullis fasciculato-stellatis. (1) in Gall provinciæ et Hispaniæ maritimis. — Cav. diss. 2. t. 17. f. 3. *M. maritima* Lam. fl. fr. 3. p. 140. (v. s.)

β. *ciliata* (Dufour! in litt.) caule erecto humillimo. (1) Tudelæ.

27. *M. ALCEA* (Lin. sp. 971.) foliis inferioribus angulatis, superioribus quinque-partitis incis, caulibus calycibusque pube stellatâ fasciculatâ subvelutinoscabris. 2 in Europâ. Cav. diss. 2. t. 17. f. 2. Sims bot. mag. t. 2197. — Blackw. t. 309. (v. v.)

β. *Moreni* (Poll! veron. 1816.) foliis superioribus palmato-5-fidis, lobis dentatis. — in Italiâ. *M. alceoides* Ten. cat. 1819. p. 55. (v. s. in h. Balb.)

28. *M. MOSCHATA* (Lin. sp. 971.) foliis inferioribus reniformibus incis, caulibus multipartitis, laciniis linearibus, caulibus calycibusque pilis simplicibus scabris. 2 in Europâ. — Cav. diss. 2. t. 18. f. 1. (v. v.)

β. *laciniata* (Desrous. in dict. enc. 3. p. 750.) foliis etiam inferioribus multifidis, laciniis linearibus. — Moris. ox. s. 3. t. 18. f. 4. *M. tenuifolia* Sav. cent. p. 122. *M. moschata* β undulata, Sims bot. mag. t. 2293. flore albo. — Variat etiam omninò glabra. (v. s.)

29. *M. ALTHÆOIDES* (Cav. diss. 2. t. 135.) foliis palmatifidis, lobis lanceolatis dentatis, pilis simplicibus, pedicellis folio longioribus, calycibus acuminatis. (1) in Hispaniâ. (v. s.)

β. *hirsuta* (Ten. prod. 40.) foliis infimis subrotundo-cordatis 7-lobatis, ceteris 5-partitis. — prope Agrigentum Siciliæ. (v. s.)

30. *M. FASTIGIATA* (Cav. diss. 2. t. 23. f. 2.) foliis cordatis 5-lobis acuminatis inæqualiter dentatis medio productiore cauleque erecto pube stellatâ tomentosis, pedicellis axillaribus et in corymbum terminalem approximatis. (2)? in Arverniâ. — DC. fl. fr. n. 4510. (v. v.)

β. *lobata* (Cav. diss. 2. t. 18. f. 4.) foliis crenatis subcordatis. — in Hispaniâ.

γ. *Albulensis* (Cav. diss. 2. t. 34. f. 3.) foliis superioribus basi cuneiformibus. — in Hispaniâ.

§. 5. *Fasciculatæ, floribus purpurascentibus aut albis, foliis angulatis cordatis 5-nerviis, pedicellis axillaribus plurimis 1-floris, calyce exteriori 3-phylo, caule herbaceo.*

31. *M. MAURITIANA* (Lin. sp. 970.) caule erecto, foliis 5-lobatis obtusis, pedicellis petiolisque glabriusculis aut ad latus superius tomentosis. (1) in Italiâ, Hispaniâ, Lusitaniâ. *M. obtusa* Mœnch. Cav. diss. 2. t. 25. f. 2. — Flores intensè purpurei. (v. s.)

β. *Sinensis* (Cav. diss. 2. t. 25. f. 4.) caule reclinato glaberrimo. — *M. glabra* Desrous. in dict. enc. 3. p. 752.

32. *M. SYLVESTRIS* (Lin. sp. 969.) caule erecto, foliis 5-7-lobatis acutis, pedicellis petiolisque pilosis. (2) in Europæ campestribus. Cav. diss. 2. t. 26. f. 2. *M. vulgaris* Ten! prod. fl. nap. (v. v.)

β. *albiflora*, lobis obtusioribus; floribus albis. *M. Veneta* hortul. (v. s.)

33. *M. HENNINGII* (Goldbl! act. mosc. 5. p. 133.) caule diffuso, foliis subreniformibus obtusissimè 7-lobatis, pedicellis petiolisque glabriusculis, petalis calycem non excedentibus, carpellis rugosis. 2 circa Mosquam frequens. (v. s.)

34. *M. ROTUNDIFOLIA* (Lin. sp. 969.) caule prostrato, foliis cordato-orbiculatis obtusissimè 5-lobatis, pedicellis fructiferis declinatis petiolisque pubescen-

tibus, corollis calyce duplò majoribus. $\frac{1}{2}$ in Europæ campestribus. Cav. diss. 2. t. 26. f. 3. Albo flore ludit. (v. v.)

β . *pusilla* (Smith engl. bot. t. 241.) petalis vix calyce majoribus. M. parviflora Huds. angl. 307. non Lin.

γ . *crenata* (Kit! in litt.) petalis calyce vix majoribus, foliis indivisis crenato-serratis. $\textcircled{1}$ in Hungariæ subsalsis. An species propria?

35. *M. BRASILIENSIS* (Desrous. in dict. enc. 3. p. 744.) caule diffuso, foliis cordato-orbiculatis 7-lobatis molliter villosis, lobis acutiusculis, floribus aggregatis pedicellatis, involucelli foliolis setaceis. — in Brasiliâ prope RioJaneiro. \dagger .

36. *M. NICÆNSIS* (All. ped. n. 1416.) caule prostrato, foliis angulatis acutis, pedicellis 3-4 erectis petiolo quadruplò brevioribus, calycibus pilosis corollâ brevioribus. $\textcircled{1}$ secus vias Nicæ et in Bætiâ. — Cav. diss. 2. t. 25. f. 1. (v. s.)

β . *montana* (Forsk. descr. 124.) foliis mollibus villosis. — in Ægypto.

37. *M. MICROCARPA* (Desf. cat. h. par.) caule erecto, foliis cordato-subrotundis subquinque lobis crenatis glabris, pedicellis subgeminis petiolo brevioribus, corollis paulò calycem superantibus. $\frac{1}{2}$ in Ægypto. Pers. ench. 2. p. 251. (v. v.)

38. *M. PARVIFLORA* (Lin. am. 3. p. 416.) caule patulo, foliis subrotundis obtusè angulatis crenatis glabriusculis, floribus axillaribus sessilibus glomeratis, corollis vix calyce longioribus. $\frac{1}{2}$ in Galliâ australi, Mauritanîâ. Cav. diss. 2. t. 26. f. 1. Carpella supernè rugosa margine dentata. (v. v.)

39. *M. VERTICILLATA* (Lin. sp. 970.) caule erecto, foliis cordatis suborbiculatis obtusè angulatis, floribus axillaribus glomeratis sessilibus, calycibus scabris subinflatis, carpellis subglabris. $\frac{1}{2}$ in Chinâ. Cav. diss. 2. t. 25. f. 3. M. glomerata hortul. (v. v.)

40. *M. MAROTICA* (Delil. ined.) caule erecto, foliis semi-rotundis 5-nerviis crenatis, summis sublobatis, floribus in capitulum terminalem subaggregatis subsessilibus, calycibus inflatis 5-fidis, lobis latissimis mucronatis integris, carpellis reticulatis pubescentibus. $\textcircled{1}$ in Ægypto. (v. s. in h. Balb.)

41. *M. TRIONOIDES*, caule erecto, foliis semirotundis 5-nerviis crenato-dentatis, summis sublobatis, floribus in capitulum terminalem approximatis subsessilibus, calycibus inflatis 5-fidis, lobis lanceolato-acuminatis ciliato-serratis, carpellis hirsutissimis. $\textcircled{1}$ Patr. ign. Involucelli foliola setacea. — Hæc et duæ priores sectionem ferè constituunt. (v. s.)

42. *M. CRISPA* (Lin. sp. 970.) caule erecto, foliis angulatis dentatis crispis glabris, floribus axillaribus sessilibus. $\textcircled{1}$ in Syriâ. — Cav. diss. 2. t. 23. f. 1. (v. v.)

43. *M. FLEXUOSA* (Horn. h. hafn. 2. p. 655.) caulibus prostratis flexuosis, foliis cordato-orbiculatis 7-lobatis crenatis, floribus axillaribus sessilibus paucis. $\textcircled{1}$ Patr. ign. (v. s.)

44. *M. TENELLA* (Cav. ic. 5. t. 422. f. 3.) caule humifuso, foliis trilobis crenatis, floribus axillaribus sessilibus ternis, corollis vix calyce majoribus. $\textcircled{1}$ in montibus Cordillera de Chili.

45. *M. BONARIENSIS* (Cav. diss. 2. t. 22. f. 1.) caule erecto ramoso, foliis trilobatis, lobis inæqualiter crenato-dentatis, floribus glomeratis axillaribus sessilibus parvis. — in agro Bonariensi. An hujus sectionis? \dagger

§. 6. Capenses, floribus purpurascentibus aut albis, foliis angulato-lobatis, caule fruticoso, calyce exteriori seu involucello 3-phylo, pedicellis solitariis 1-floris rariùs geminis ternisve aut 2-3-floris. — N. B. Plures hujus sect. species videntur, aut varietates aut hybridæ à culturâ ortæ denuò examinandæ. Omnes ad cap. Bonæ-Spei crescunt.

46. *M. AMOENA* (Sims bot. mag. t. 1998.) pedicellis 1-floris aggregatis folio bre-

vioribus, involucellis ovatis acuminatis, foliis 5-lobatis pilosis rugosis. ♂. (v. v.)

47. *M. VIRGATA* (Cav. diss. 2. p. 70. t. 18. f. 2.) pedicellis 1-floris solitariis geminisve petiolo longioribus, involucellis linearibus, foliis incisis crenatis glabris rigidis. ♂. (v. v.)

48. *M. CAPENSIS* (Cav. diss. p. 71. t. 24. f. 3.) pedicellis 1-floris solitariis geminisve petiolo longioribus, involucellis ovato-lanceolatis, foliis 5-lobis 3-lobisque crenato-dentatis glutinosis. ♂ *M. biflora* Desrous. in dict. encl. 3. p. 747. — Ab hac forsitan diversa *M. n.* 3429. Burch. cat? (v. v.)

49. *M. BAISAMICA* (Jacq. ic. rar. 1. t. 140.) pedicellis 1-floris solitariis petiolo longioribus, involucellis oblongo-linearibus, foliis ovatis subtrilobis acutis inæqualiter dentatis. ♂. (v. s.)

50. *M. OXYACANTHOIDES* (Horn. h. hafn. 2. p. 654.) pedicellis solitariis petiolo longioribus, foliis glabris cuneiformibus profunde trilobis inciso-serratis, lobo intermedio majore. ♂.

51. *M. RUGOSA* (Desrous. in dict. encl. 3. p. 747.) pedicellis 1-floris solitariis petiolo longioribus, involucellis ovatis acutis, foliis subcordatis pinnatifido-sinuatis rugosis. ♂.

52. *M. TRIDACTYLITES* (Cav. diss. 2. p. 75. t. 21. f. 2.) pedicellis solitariis 1-floris foliorum longitudine, foliis sessilibus cuneiformibus trifidis integris. ♂ *M. reflexa* Andr. bot. rep. t. 135.

53. *M. DIVARICATA* (Andr. bot. rep. t. 182.) pedicellis solitariis petiolo longioribus, foliis lobatis plicatis dentatis scabridis, ramis ramulisque divaricatis flexuosis. ♂ *Cor. alba* ad basim rubro lineata. (v. v.)

54. *M. RETUSA* (Cav. diss. 2. t. 21. f. 1.) pedicellis solitariis petiolo longioribus, involucellis lanceolatis, foliis oblongis obtusissimis trilobis dentatis tomentosis. ♂ *M. triloba* Thunb. prod. 118.

55. *M. CALYCINA* (Cav. diss. 2. p. 81. t. 22. f. 4.) pedicellis solitariis 1-floris petiolo duplò longioribus, involucellis ovatis acutis maximis, foliis cordatis crenatis piloso-scabris. ♂. (v. s.)

56. *M. FRAGRANS* (Jacq. h. vind. 3. t. 35.) pedicellis 1-floris solitariis petioli longitudine, involucellis lanceolatis, foliis cordatis 5-lobis dentatis, ramis glutinosis. ♂ Cav. diss. 2. p. 72. t. 23. f. 3. (v. s.)

57. *M. STRICTA* (Jacq. h. Schœnbr. 3. t. 294.) pedicellis solitariis 1-floris petioli longitudine, involucellis sublinearibus, foliis ovatis subtrilobis dentatis hirtis. ♂ *Cor. alba* rubro virgata.

58. *M. BRYONIFOLIA* (Lin. sp. 968.) pedicellis solitariis 1-2-floris petiolo brevioribus, foliis cordatis sub-5-lobis obtusis pube stellatà scaberrimis. ♂ *M. reflexa* Wendl. h. herrenh. 1. t. 4. *M. stellata* Thunb. prod. 119? (v. s.)

59. *M. GLOSSULARIÆFOLIA* (Cav. diss. 2. p. 71. t. 24. f. 2.) pedicellis solitariis 1-3-floris petioli longitudine, involucellis oblongo-linearibus, foliis sinuato-lobatis serratis rugosis ramisque hirsutis. ♂. (v. s.)

60. *M. ASPERRIMA* (Jacq. h. Schœnbr. 2. t. 139.) pedicellis 1-2-floris solitariis petiolo sublongioribus, involucellis linearibus, foliis 5-lobis obtusis lobo terminali elongato rugoso-dentatis scaberrimis. ♂. (v. s.)

♂. 7. Multifloræ, floribus purpurascensibus aut albis, pedunculis axillaribus multifloris, involucello 3-phyllo, foliis angulatis.

61. *M. LACTEA* (Ait. h. Kew. 2. p. 448.) frutescens, foliis angulatis acutis cordatis villosis, petalis obcordatis calyce subbrevioribus, pedunculis paniculatis. ♂ in Mexico. *M. vitifolia* Jacq. ic. 1. t. 30. (v. s.)

62. *M. CAPITATA* (Cav. diss. 5. t. 137. f. 1.) frutescens, foliis quinque-lobis, lobis pinnatifido-sinuatis dentatis, intermedio elongato, pedunculis corymboso-capitatis, carpellis birostratis. ♂ in Peru.

63. *M. MINIATA* (Cav. ic. 3. t. 278.) fruticulosa, foliis ovatis trilobis dentatis tomentosis, pedunculis axillaribus racemosis paucifloris (interdum 1-floris). $\frac{5}{2}$ Patr. ign. Corollæ miniatæ. (v. v.)
64. *M?* *OPERCULATA* (Cav. diss. 2. t. 35. f. 1.) frutescens, foliis angulato-quinque lobis, lobo intermedio majoræ, pedunculis axillaribus racemosis, floribus secundis, carpellis operculatis. $\frac{5}{2}$ in Peruvæ arenosis. An genus?
65. *M. PERUVIANA* (Lin. sp. 968.) herbacea erecta, foliis palmatifidis, spicis axillaribus secundis, carpellis denticulatis. (1) in Peru. Jacq. h. vind. t. 156. Cor. parvæ purpureæ. *M. exasperata* Mœnch. (v. v.)
66. *M. LIMENSIS* (Lin. sp. 968.) herbacea erecta, foliis 7-lobatis rugosis, spicis secundis axillaribus, carpellis lævibus. (1) in Peru. Jacq. hort. vind. t. 141. *M. lævis* Mœnch. Cor. cæruleæ. (v. v.)
67. *M. ACAULIS* (Cav. diss. 2. p. 82. t. 35. f. 2.) herbacea acaulis, foliis angulatis dentatis, dentibus bisetis, pedunculis radicalibus multifloris. $\frac{2}{4}$ in Peruvæ Andibus.

Sect. II. MALUCHIA.

Involucellum 5-6-phyllo, foliolis linearibus. Carpella 5 monosperma distincta indehiscencia.

68. *M. HIBISCOIDES* (Desrous. in dict. enc. 3. p. 748.) foliis ovatis acuminiatis subangulatis serratis molliter villosis, pedicellis petiolo paulò brevioribus, involucello 5-phyllo. — in ins. Borboniæ. Flores rubri.
69. *M. BORYANA*, foliis quinquangularibus dentatis acutis pubescentibus, pedicellis brevissimis, involucello 6-phyllo. — in ins. Borboniæ. Bory S.-Vincent. Calycis lobi basi 3-nervi. Cor. non vidi. (v. s.)

Sect. III. SPHÆROMA.

Carpella unilocularia 2-oo-sperma in fructum globulosum coalita. Pedunculi axillares sæpius multiflori. Caulis fruticosus. An genus proprium?

70. *M. UMBELLATA* (Cav. ic. 1. t. 95.) foliis subpeltatis 5-lobis obtusis, pedunculis axillaribus; umbelliferis, involucelli foliolis obovatis substipitatis deciduis. $\frac{3}{2}$ in declivitate orient. montium Novæ-Hispaniæ. Flores magni igneo-violacei. (v. s.)
71. *M. ROSEA* (fl. mex. ic. ined.) foliis subpeltatis 5-lobis acutis, pedunculis axillaribus 1-3-floris, involucelli foliolis ovatis sessilibus. $\frac{3}{2}$ in Mexico. Flores rosei; involucelli lobi 3-nervi. Priori nimis affinis.
72. *M. ARBUTILOIDES* (Lin. sp. 971.) foliis quinquangularibus tomentosis, pedunculis axillaribus bifidis paucifloris, involucelli foliolis oblongis parvis, fructibus globosis. $\frac{3}{2}$ in insulis Bahamensibus, Jacq. h. Schœnb. 3. t. 293. (v. s.)
73. *M. ELAGANS* (Cav. diss. 2. t. 16. f. 1.) foliis tripartitis canescentibus, laciniiis apice dentatis, intermediâ trifidâ, pedicellis axillaribus 1-floris petiolo ferè brevioribus, fructibus globosis. $\frac{3}{2}$ ad cap. Bonæ-Spei. Jacq. Coll. 4. t. 6. f. 1. — An verè polysperma?
74. *M. ANGUSTIFOLIA* (Cav. diss. 2. p. 64. t. 20. f. 1.) foliis lanceolatis dentatis tomentosis, pedunculis axillaribus binis paucifloris, involucelli foliolis setaceis deciduis. $\frac{5}{2}$ in Mexico.

Sect. IV. MODIOLA. (*Mench. meth.* 620.)

Carpella bivalvia disperma, valvis biaristatis introrsum inflexis et indè carpella semi-bilocularia. Pedicelli axillares 1-flori. Caulis herbaceus prostratus aut diffusus. — An genus?

75. *M. CAROLINIANA* (Lin. sp. 969.) foliis palmato-5-lobatis inciso-dentatis, pedicellis solitariis petiolo longioribus, fructu villosio. (1) in Carolinâ. Modiola multifida Mœnch. l. c. (v. s.)
76. *M. URTICÆFOLIA* (H. B. et K. nov. gen. am. 5. p. 276.) foliis subrotundo-ovatis

acutis obsolete cordatis serrato-incisis subnovemlobis pilosiusculis, pedicellis solitariis petiolo brevioribus, floribus 10-andris, capsulis 15-coccis hispidis. \mathcal{L} Prope Santa-Fé de Bogôta. Flores violacei magni. prioris cui affinis.

77. *M. PROSTAT* (Cav. diss. 2. t. 16. f. 3.) foliis palmato-5-lobatis inciso-dentatis, pedicellis solitariis petiolo longioribus, fructu glabro, petalis integris. — Ad vias circa Monte-Video.
78. *M. ERIOCARPA*, foliis palmato-5-lobis inciso-dentatis, pedicellis geminis petioli longitudine, fructu villosa. — Circa Buenos-Ayres in campis dictis Pampas. (v. s.)
79. *M. DECUMBENS* (Willd. enum. 731.) foliis ovatis inciso-dentatis sublobatis, pedicellis petiolo longioribus, fructu villosa, petalis integris. \mathcal{L} in Americâ merid. (v. v.)

†. *Species non satis notæ.*

80. *M. CHINENSIS* (Mill. dict. n. 5.) caule erecto herbaceo, foliis suborbiculatis obsolete 5-lobatis, floribus confertis alaribus sessilibus. ① in Chinâ. An *M. verticillata*?
81. *M. ORIENTALIS* (Mill. dict. n. 3.) caule erecto herbaceo, foliis lobatis obtusis crenatis. ① in Oriente. Flores magni suavè rubentes.
82. *M. ITALICA* (Poll. pl. veron. 1816.) caule hirsuto, foliis inferioribus subreniformibus 5-lobis, superioribus 5-partitis, omnibus dentatis scabriusculis. — in agro Mediolanensi.

III. *KITAIBELIA*. Willd. n. act. soc. n. scrut. ber. 2. p. 107. t. 4. f. 4.

Calyx cinctus involucello, 7-9-fido. Carpella capsularia 1-sperma in capitulum 5-lobum adgregata.

1. *K. VITIFOLIA* (Willd. l. c.) \mathcal{L} in Slavoniæ com. Sirmiensi. Herba ramosissima, 4-8-pedalis. Folia 5-loba acuta dentata. Flores albi. — Waldst. et Kit! pl. rar. hung. 1. p. 29. t. 31. (v. v.)

IV. *ALTHÆA*. Cav. diss. 2. p. 91.

Calyx cinctus involucello 6-9 fido. Carpella capsularia 1-sperma in orbem disposita.

Sect. I. *ALTHÆASTRUM*. — *Althæa*. Lin. gen. n. 839. Lam. ill. t. 581.

Carpella emarginata, margine membranaceo destituta. Involucellum sæpius 8-9-fidum.

1. *A. OFFICINALIS* (Lin. sp. 966.) foliis utrinquè molliter tomentosis cordatis ovatisve dentatis indivisis subtrilobisve, pedunculis axillaribus multifloris folio multò brevioribus. \mathcal{L} in palustribus Europæ præsertim maritimis. — Cav. diss. 2. p. 93. t. 30. f. 2. (v. v.)
2. *A. TAURINENSIS*, foliis utrinquè molliter tomentosis trifidis subtripartitis dentatis, pedunculis axillaribus multifloris folio sublongioribus strictis. \mathcal{L} in collibus circa Taurinum. *A. officinalis*, β . Willd. sp. 3. p. 771. Media inter priorem et posteriorem, ab utraq; adhuc diversa calycibus magis acuminatis. An unius alteriusve varietas? (v. v.)
3. *A. NARBONENSIS* (Poirr. in Cav. diss. 2. p. 94. t. 29. f. 2.) foliis pubescenti-subtomentosis inferioribus 5-7-partitis, superioribus trifidis, pedunculis multifloris folio longioribus laxiusculis. \mathcal{L} in agro Narbonensi et Hispaniâ. Jacq. ic. rar. 1. t. 138. (v. s.)
4. *A. CANNABINA* (Lin. sp. 966.) foliis pubescentibus subtus subincanis, inferioribus palmato-partitis, superioribus 3-partitis, lobis angustis grosse dentatis, pedunculis axillaribus multifloris laxi folio longioribus. \mathcal{L} in Galliâ australi, Italiâ, Hungariâ. Cav. diss. 2. t. 30. f. 1. (v. v.)

5. *A. HIRSUTA* (Lin. sp. 965.) foliis cordatis piloso-scabris suprà glabris, infimis obtusis, superioribus 5-lobis, caule hispido, pedicellis 1-floris folio longioribus. ① in sepibus Europæ. Cav. diss. 2. t. 29. f. 1. (v. v.)
6. *A. LUDWIGII* (Lin. mant. 98.) foliis glabris cordato-subrotundis lobatis dentatis, pedicellis axillaribus congestis 1-floris, calycibus villosis. ① in Sicilia, Ægypto. — Cav. diss. 2. t. 30. f. 3. icon. 5. t. 423. (v. s.)

Sect. II. *ALCEA*. Lin. gen. n. 840. Lam. ill. t. 581.

Carpella margine membranaceo sulcata circumdata. Involucellum 6-7-fidum.

7. *A. ACAULIS* (Cav. diss. 2. t. 27. f. 3.) subcaulis, foliis subrotundo-cordatis subquingularibus crenatis, pedicellis 1-floris petiolo multò brevioribus, petalis emarginatis basi barbatis. ① in Syria. (v. s.)
8. *A. CARIBÆA* (Sims bot. mag. t. 1916.) caule stricto glabriusculo, foliis rotundato-lobatis crenulato-serratis, floribus solitariis subsessilibus, petalis subobcordatis unguibus barbatis, stigmatibus diffusis. ② in insulis Caribæis.
9. *A. STRIATA*, caule puberulo subscabro, foliis cordatis obtusè trilobis crenatis, floribus solitariis breviter pedicellatis, involucello duplò calyce breviorè, lobis calycis 7-striatis, petalis obcordato-bilebis. — patr. ign. olim. in hort. Cels culta. Flos pallidus 2 $\frac{1}{2}$ poll. diam. (v. s.)
10. *A. PALLIDA* (Wald. et Kit. in Willd. sp. 3. p. 773.) caule erecto hispido, foliis subrotundis cordatis, involucello calycem æquante, petalis bilobis. ② in Hungaria. Flores albedo-purpurei. (v. s.)
11. *A. ROSEA* (Cav. diss. 2. t. 29. f. 3.) caule stricto hirsuto, foliis cordatis 5-7-angulatis crenatis rugosis, floribus axillaribus sessilibus ad apicem subspicatis, petalis subcrenatis, unguibus villosis. ② in Oriente. *Alcea rosea*. Lin. sp. 966. (v. v.)
- Æ? *biloba*, petalis obtusè bilobis. — hæc interdum in hortis occurrit sub nomine *A. grandifloræ*, sed flores potius minores. (v. s.)
12. *A. SINENSIS* (Cav. diss. 2. t. 29. f. 3.) caule stricto glabro infernè ramoso, foliis cordatis scabris crenatis angulatis, floribus axillaribus sessilibus ad apicem spicatis, petalis subcrenatis, unguibus villosis. ① in Chinâ. An prioris varietas? Differt radice annuâ, caule humiliore 3-4-pedalî, floribus ferè majoribus.
13. *A. AFRICANA* (Lour.-fl. coch. 2. p. 513. sub *Alceâ*.) caule fruticoso hispido, foliis trilobis crenatis, floribus solitariis axillaribus pedicellatis, involucello et calyce 6-partitis. 3 ad oras Africæ orient. ↑.
14. *A. COROMANDELIANA* (Cav. diss. 2. p. 93.) caule erecto tomentoso, foliis subtriangularibus crenatis obtusis 5-nerviis subtrilobis, floribus solitariis subsessilibus axillaribus, petalis lato-oblongis. — in Pondicheri.
15. *A. FLEXUOSA* (Sims bot. mag. t. 892.) caule subflexuoso hispido, foliis cordatis subseptem-lobis obtusis longè petiolatis, floribus axillaribus solitariis pedicellatis, petalis obcordatis. — in Indiâ orient.
16. *A. FICIFOLIA* (Cav. diss. 2. p. 92. t. 28. f. 2.) caule erecto piloso, foliis ultrà medium palmato-7-lobatis, lobis oblongis obtusis irregulariter dentatis involucellis calyce dimidiò ferè brevioribus. ② in Sibirîâ, ad Tauaim. *Alcea ficifolia* Lin. sp. 967. — Blackw. herb. t. 54. Variat in hortis floribus pedicellatis. (v. s.)
17. *A. LAVATERÆFLORA*, caule erecto piloso, foliis utrinquè tomentosis ultrà medium palmato-lobatis, lobis oblongis obtusis irregulariter dentatis, involucellis striato-nervosis calyces ferè æquantibus. — prope Seyde ad radices Libani. Meryon. (v. s.)

Sect. III. *ALPHÆA*.

Carpella nervulis rugosa, non marginata. Involucellum 5-fidum.

18. *A. BURCHELLII*, caule erecto piloso, foliis cordatis sub-5-lobis grossè dentatis velutinis, pedicellis axillaribus 1-floris pilosis petiolo vix brevioribus, involucello calyceque 5-fidis. — ad cap. Bonæ-Spei. *Urena pilosa* Burch! cat. n. 2557. (v. 6.)

19. *A. BORBONICA*, caule erecto infernè villosa supernè folisque velutinis, foliis cordatis, nonnullis sub-5-lobis grossè dentatis, pedicellis axillaribus 1-floris petiolo duplò longioribus, involucellis basi ciliatis. $\frac{5}{2}$ in ins. Borboniæ ad rivum Sancti-Dyonysii. Bory Saint-Vincent. Species priori valdè affinis. (v. 8.)

V. *LAVATERA*. *Lin. gen. n.* 842. *Lam. ill. t.* 582.

Calyx cinctus involucello 3-6-fido, foliolis nempè ad medium coalitis. Carpella capsularia, 1-sperma, in orbem circa axim variè dilatatum disposita. — An sectiones in genera mutandæ?

Sect. I. *STEGIA*. *fl. fr.* 4. *p.* 835.

Receptaculum seu axis fructus apice expansus in discum ovaria tegentem.

1. *L. TRIMESTRIS* (Lin. sp. 974.) caule herbaceo, foliis glabriusculis subrotundo-cordatis, superioribus angulatis, pedicellis solitariis, orbiculo carpella omninò tegente. ① in Syriâ, Hispaniâ. — Cav. diss. 2. t. 3. f. 1. *Stegia Lavatera fl. fr. n.* 4525. Variat flore roseo et albo. (v. v.)

2. *L. PSEUDO-OLBIA* (Poir. suppl. 3. p. 309.) caule fruticoso, foliis tomentosis, inferioribus 5-lobis, pedicellis aggregatis, orbiculo carpella semi-tegente. $\frac{3}{2}$ patr. ign. *L. undulata* Desf. arb. 1. p. 471. non Mill. Habitus ferè sectionis sequentis. (v. v.)

Sect. II. *OLBIA*. *Medik. malv. p.* 41.

Receptaculum fructus centrale conicum exsertum.

3. *L. PHENICEA* (Vent. malm. t. 120.) caule arboreo, foliis acutè 5-lobis dentatis glabriusculis, pedunculis solitariis 3-5-floris, involucello caduco. $\frac{3}{2}$ hab. Maderæ aut in Africâ. Culta in hortum Canar. undè migravit in Europæos. Flores ampli phœnicei. *L. coccinea* Dietr. (v. s.)

4. *L. ACERIFOLIA* (Cav. el. h. madr. p. 20.) caule fruticoso, foliis acutè 5-lobis subdentatis glabriusculis, pedicellis solitariis 1-floris. $\frac{3}{2}$ in Teneriffâ. Broussonet. DC. cat. h. monsp. p. 121. Lois herb. amat. t. 322. Fl. pallidè lilacini. (v. v.)

5. *L. JULII* (Burch! cat. n. 2664.) caule fruticoso, foliis 3-5-lobatis subtus cano-tomentosis, lobis obtusis, terminali longiore, pedunculis subracemosis, involucello 3-partito. $\frac{3}{2}$ ad cap. Bonæ-Spei. An forsàn inter *Malvas* potiùs adnumeranda? (v. s.)

6. *L. HISPIDA* (Desf. atl. 2. p. 118. t. 171.) caule fruticoso pilis fasciculatis hispido, foliis subcanescentibus 5-lobis, summis 3-lobis aut indivisis, floribus subsessilibus, involucello 3-partito maximo birsutissimo. $\frac{3}{2}$ in sepibus Algeriæ. (v. s.)

7. *L. AFRICANA* (Cav. diss. 5. p. 282. t. 139. f. 1.) caule fruticoso pube floccosâ subtomentoso, foliis canescentibus omnibus obtusè 5-lobis, florum pedicellis petiolo æqualibus, involucello 3-partito velutino. $\frac{3}{2}$ in Hispaniâ et Africâ septentr. Varietas prioris ex Willd. (v. s.)

8. *L. OLBIA* (Lin. sp. 972.) caule fruticoso pilis subfasciculatis distantibus subscabro foliis molliter cano-tomentosis 5-lobis, et superioribus 3-lobis, lobo medio elongato, summis oblongis subindivisis, floribus solitariis sessilibus. $\frac{3}{2}$ in sepibus circa Olbiam ubi ipse legi. — Lob. ic. t. 653. f. 2. Vereor ne icones recentiorum ad sequentem sint referendæ. (v. v.)

9. *L. UNGUICULATA* (Desf. arb. 1. p. 471.) caule fruticoso pube stellatâ tomentoso, foliis utrinquè tomentosis acutè 5-lobis, superioribus 3-lobis, floribus solitariis breviter pedicellatis. $\frac{3}{2}$ in insulâ Samo. Priori valdè affinis et pro eâ in hortis sæpè admissa. (v. v.)

10. *L. MICANS* (Lin. sp. 972.) caule fruticoso, foliis 7-angularibus acutis crenatis plicatis tomentosis, racemis terminalibus. 3 in Hispaniâ, Lusitaniâ? — Moris. ox. s. 5. t. 17. f. g. *L. bryonifolia* Mill. dict. n. 11.
11. *L. LUSITANICA* (Lin. sp. 973.) caule fruticoso, foliis 7-angularibus tomentosis plicatis, racemis terminalibus. 3 in Lusitaniâ? *L. undulata* Mill. dict. n. 10. Omnes quæ sub nomine *L. Lusitanicæ* occurrunt in hortis à descriptione Linnæanâ valdè recedunt. †.
12. *L. FLAVA* (Desf. atl. 2. p. 119. t. 172.) caule herbaceo tomentoso, foliis subtomentosis subrotundis obsolete trilobis, pedicellis axillaribus adgregatis. 2 in cultis prope Mascar et in Sicilia. *L. Agrigentina*. Tineo. pl. sic. 1. p. 13. *L. Empedoclis* Rœusch. (v. s.)
13. *L. PLEBEIA* (Sims bot. mag. t. 2269.) caule herbaceo scabro, foliis 5-lobatis subtus pubescentibus, pedunculis axillaribus adgregatis (solitariisve), petalis cuneiformibus emarginatis acutis. 2 5? in Novâ-Hollandiâ. Fruct. indescr. et ideò locus in genere dubius.
14. *L. THURINGIACA* (Lin. sp. 973.) caule herbaceo tomentoso, foliis subtomentosis, inferioribus angulatis, superioribus 3-lobis, lobo intermedio longiore, pedicellis solitariis 1-floris petiolo longioribus, petalis bilobis. 2 in Thuringiâ, Germaniâ, Tatariâ, in collibus circa Odessam. D'Urv. enum. 616. Jacq. fl. austr. t. 311. *L. Olbia* Steph. mosc. 32. non Lin. (v. v.)
15. *L. BIENNIS* (Bieb. fl. taur. 2. p. 183.) caule herbaceo folisque tomentosis, inferioribus subrotundis lobatis, floralibus 3-lobis, lobo medio lanceolato, pedicellis solitariis 1-floris petiolo longioribus, petalis subemarginatis. 2 in Caucasi orient. campestribus. (v. s.)
16. *L. PUNCTATA* (All. auct. p. 26.) caule herbaceo pube stellatâ subscabro, foliis subtomentosis, inferioribus orbiculato-cordatis, summis trilobis, pedicellis solitariis 1-floris folio longioribus. 1 in agris et cultis Gallo-provinciæ, Niceæ, Italiæ. *Olbia deflexa* Moench. suppl. p. 200. (v. v.)

Sect. III. AXOLOPHA.

Receptaculum apice truncatum in cristas tot quot carpella membranaceas laterales verticales expansum.

17. *L. MARITIMA* (Gou! ill. p. 46. t. 11. f. 2.) caule fruticoso foliisque tomentosis subrotundis obtusè angulatis crenatis, pedicellis axillaribus solitariis. 3 in Galliæ austr. et Hispaniæ rupestribus. Cav. diss. 2. t. 32. f. 3. *L. Hispanica* Mill. dict. n. 9. *L. rotundifolia* Lam. (v. v.)
18. *L. TRILOBA* (Lin. sp. 972.) caule fruticoso foliisque tomentosis subcordatis subtrilobis rotundis crenatis, pedicellis adgregatis, calycibus acuminatis. 3 in Hispaniâ. Cav. diss. 2. t. 31. f. 1. *L. calycina* Poir. suppl. 3. p. 310. (v. s.)
19. *L. SUBOVATA*, caule suffruticoso, foliis subtomentosis ovatis dentatis subtrilobis, lobo medio productiore, pedicellis 1-2 axillaribus petioli longitudine, calycis lobis acuminatis. 3 in agro Mogadoriensi, ut autumo ex specim. incompl. ad Broussonet lecto. (v. s. c.)

Sect. IV. ANTHEMA. *Med. malv. p. 42.*

Receptaculum parvum foveolatum, nec exsertum, nec in cristas expansum.

20. *L. ARBOREA* (Lin. sp. 972.) caule arboreo, foliis 7-angularibus subtomentosis plicatis, pedicellis axillaribus 1-floris confertis petiolo multo brevioribus. 2 in Italiâ, Hispaniâ, Lusitaniâ, Angliâ, Africâ bor., Canariis. — Cav. diss. 2. t. 139. f. 2. (v. v.)
21. *NEAPOLITANA* (Ten. cat. 1819. n. 125.) caule herbaceo scabro erecto, foliis subrotundis 7-nerviis obtusissimè 7-lobatis crenatis, pedunculis axillaribus adgregatis, involucllo calycè brevior, calycis lobis acuminatis. 2 in maritimis circa Neapolim. Petala obcordata cærulea. (v. s.)
22. *L. CRETICA* (Lin. spec. 973.) caule herbaceo hispido-scabro, foliis 5-lobis

acutis, pedicellis axillaribus 4-floris aggregatis petiolo multò brevioribus. ① in Cretâ. Cav. diss. 2. t. 32. f. 1. (v. s.)

23. *L. SYLVESTRIS* (Brot. fl. lus. 2. p. 277.) caule herbaceo pube stellatâ scabro, foliis obtusè angulatis subrotundis. summis subtrifidis acutis, pedicellis 1-floris geminis ternisve petiolo brevioribus, involucellis 3-partitis. ① ad sepes Lusitanicæ. An L. sicula. Tin. sic. pug. 1. p. 14? An L. Weinmanniana Trev? (v. s.)
24. *L. AMBIGUA*, caule herbaceo hispido-scabro, foliis inferioribus 5-lobis angulatis, superioribus 3-lobis, lobis lanceolatis acutis dentatis, medio duplò longiore, pedicellis solitariis folio brevioribus, carpellis denudatis. — in agro Neapolitano. Lav. sylvestris Ten. prod. 40. non Brot.

†. *Species non satis notæ.*

25. *L. LANCEOLATA* (Willd. enum. 733.) caule herbaceo punctato-scabro, foliis stellato-pubescentibus oblongo-lanceolatis serratis, superioribus integerrimis, pedunculis solitariis folio longioribus. ① patr. ign. An L. punctatæ varietas?
26. *L. TRIPARTITA*, caule suffruticoso, ramis foliisque pube stellatâ tomentoso-caescentibus, foliis tripartitis, lobis cuneatis apice grossè inciso-dentatis, pedicellis solitariis 1-floris, involucelli tubo calycis adnatis et hujus lobos paulò superantibus. ⑤ Patria et fruct. ignot. — Ab omnibus ferè Malvaceis calyce differt. (v. s.)

VI. MALACHRA. *Lin. gen. ed. Schreb. n. 1131. Lam. ill. t. 580.*

Involucrum generale 3-5-phyllum florum capitula cingens. Calyx cinctus involuculo proprio foliolis 8-12 linearibus aut setiformibus. Carpella capsularia 5, in orbem disposita, 1-sperma.

1. *M. URENS* (Poi! in Schrad. neu journ. 2. p. 293.) foliis ovatis 5-nerviis capitulis subsessilibus, involucri foliolis basi dentatis. — in Sancto-Domingo. (v. s.)
2. *M. ROTUNDIFOLIA* (Schränk. h. mon. t. 56.) foliis orbiculatis crenatis, capitulis pedunculatis 3-phyllis 5-floris, involucri foliolis reniformi-orbiculatisve. ① in Brasiliâ.
3. *M. PLUMOSA* (Desrous. in dict. enc. 2. p. 686.) foliis ellipticis dentatis, capitulis pedunculatis polyphyllis multifloris, involucri foliolis interioribus ciliato-plumosis. — in Brasiliâ. *Sida plumosa*. Cav. diss. 1. t. 12. f. 4.
4. *M. CILIATA* (Poir. suppl. 3. p. 578.) foliis lato-ovatis inæqualiter crenatis glabris, capitulis subsessilibus, involucris bracteisque ciliatis, caule pubescente. — in Porto-Ricco.
5. *M. CORDATA* (Poir. suppl. 3. p. 578.) foliis cordatis biserratis subglabris, capitulis laxiusculis vix involveratis, bracteis elongatis filiformibus pilosis-simis. ① in Porto-Ricco. (v. s.)
6. *M. CAPITATA* (Lin. syst. 518.) foliis cordatis subrotundis obtusè angulatis denticulatis, involucris pedunculatis 3-phyllis 7-floris, caule scabro. ① in ins. Caribæarum paludosis et ad Congo ex Brown. *Sida capitata* Lin. act. ups. 1743. p. 137. t. 2. (v. s.)
7. *M. FASCIATA* (Jacq. ic. rar. 3. t. 548.) foliis subrotundis obsolete lobatis, capitulis brevè pedunculatis 3-phyllis sub-5-floris, caule villosa. ① ad Caracas. Flores rosei.
8. *M. TRILOBA* (Desf. hort. par.) foliis subrotundis obtusè trilobis crenulatis, capitulis longè pedunculatis multifloris triphyllis, caule scabro. ① Patr. ign. Flores parvi albi. Poir. suppl. 3. p. 578. (v. s.)
9. *M. RADIATA* (Lin. syst. 518.) foliis palmati-lobatis, capitulis pedunculatis 5-phyllis multifloris, involucris acuminatis, calycibus caulibusque pilosis-

simis, floribus ebracteatis. — in Sanctâ-Dominicâ et Cayennâ. Cav. diss. 2. t. 33. f. 3. *Sida radiata* Lin. sp. 965. (v. s.)

10. *M. BRÆTEATA* (Cav. diss. 2. t. 34. f. 2.) foliis palmati-lobatis, capitulis longè pedunculatis 5-phyllis multifloris, involucri foliolis ovatis acutis, floribus bracteatis, caule hirsutissimo. — in Americâ. Corolla albida.

11. *M. ALCÆFOLIA* (Jacq. ic. rar. 3. t. 549.) foliis cordatis palmato-5-lobis, capitulis pedunculatis 5-phyllis sub-10-floris, caule rariter hispido. (I) ad Caracas. Flores flavi.

12. *M. HEPTAPHYLLA* (Fisch. in Horn. suppl. 78.) foliis cordatis palmato-5-lobis crispis, lobo medio elongato, capitulis subpedunculatis sub-5-phyllis multifloris (I) in Brasiliâ. Valde priori affinis.

13. *M. PALMATA* (Mœnch. meth. 615.) foliis palmato-lobatis, lobis 3-5, intermedio productiore apice latiore, caule erecto scabro lineis 2 villosis decurrentibus. (I) Patr. ign. Flores flavi. An *M. alceæfoliæ*, heptaphyllæe varietas?

14. *M?* *URENA*, foliis angulatis inæqualiter crenatis utrinquè pilosis, capitulis axillaribus multifloris, caule suffruticoso, carpellis muricatis. 5 ad Cantonem Sinarum. *Urena polyflora* Lour. coch. ed. Willd. 2. p. 508. Ob involucri multiflora hûc, suadente cl. Willdenow, retuli. Flores crocei. †.

VII. *URENA*. Lin. gen. n. 844. Lam. ill. t. 583.

Calyx cinctus involucello 5-fido, nempè foliolis 5 ad medium coalitis. Anthæræ in apice tubi staminei. Carpella capsularia 5 conniventia 1-sperma, extus aculeis apice radiatis sæpissimè echinata. — Folia subtus in nervis sæpius glandulosa.

*. *Foliis indivisis aut subtrilobis.*

1. *U. LOBATA* (Lin. sp. 974.) foliis subrotundis obtusissimè subtrilobis utrinquè molliter velutinis 7-nerviis uniglandulosis, calycibus oblongo-lanceolatis. (I) in Chinâ. — Dill. elrh. t. 319. f. 412. Plures hic videntur confusæ species et omnia alia synonyma auctorum dubia aut falsa. (v. s.)

2. *U. SCABRIUSCULA*, foliis subrotundis acutiusculè 3-5-lobatis utrinquè scabriusculis 7-nerviis 1-3-glandulosis. calycibus linearibus. (I) in Indiâ et Napauliâ. *U. lobata* Cav. diss. 6. t. 185. f. 1. — Breyn. cent. t. 35? (v. s.)

3. *U. REPANDA* (Smith in Rees. cycl. vol. 37. n. 6.) foliis lato ovatis serrato-dentatis vix sublobatis subtus reticulatis uniglandulosis, involucellis subulatis, carpellis lævibus! — in Indiâ orientali.

4. *U. LAPPAGO* (Smith in Rees. cycl. 37. n. 11?) foliis obtusè repando-sinuatis subcordatis subtus canescenti-pubescentibus uni-glandulosis, involucelli lobis 5 lanceolatis calycem æquantibus. — in Amboinâ. Icon Rumph. amb. 6. t. 25. differt foliorum lobis nimis acutis. (v. s.)

5. *U. MULTIFIDA* (Cav. diss. 6. p. 336. t. 184. f. 2.) foliis latè ovatis inciso-lobatis, sinubus angustis, lobis acutis grossè inæqualiter dentatis, supernè pubescentibus, subtus villosis pallidis 1-glandulosis 7-9-nerviis. — in ins. Bourboniâ et Mauritiî. (v. s.)

6. *U. PROCUMBENS* (Lin. sp. 975.) foliis oblongis sinuatis serratis, caule fruticoso procumbente. 5 in Chinæ montibus.

7. *U. TRICUSPIS* (Cav. diss. 6. p. 334. t. 183. f. 1.) foliis ovato-angulatis villosotomentosis trilobis, lobis acuminatis uni glandulosis, involucello striato, caule piloso. 2 in insulis Franciæ et Borboniæ. (v. s.)

8. *U. RETICULATA* (Cav. diss. 6. p. 335. t. 183. f. 2.) foliis inferioribus 3-lobis, superioribus lanceolatis subpanduriformibus subtus canis reticulatis uniglandulosis, involucello calycem superante. 5 in Cayennâ et Santo-Domingo. *U. Americana* Lin. f. suppl. 308. non Smith.

9. *U. SUBTRILOBA* (Schränk. h. mon. t. 79.) foliis uniglandulosis mollibus sub-

- orbiculatis 7-nerviis inæqualiter crenatis, inferioribus acuminatis, mediis trilobis, superioribus profundius fissis, lobis omnium acuminatis. ① in Brasiliâ.
10. *U. GRANDIFLORA* (fl. mex. ic. ined.) foliis ovatis subcordatis irregulariter subtrilobis grossè et inæqualiter dentatis, petalis basi glandulosis. — in Mexico. Flos diametro ultra-pollicari. †
11. *U. STELLATA* (Spreng. nen. entd. 2. p. 163.) foliis omnibus oblongis eglandulosis inæqualiter dentatis stellato-pubescentibus suprâ punctatis subtus incanis, floribus racemosis, fructibus glabris. 5 in Brasiliâ.
12. *U. VIMINEA* (Cav. diss. 6. p. 335. t. 184. f. 1.) foliis subrombeis dentatis subtus reticulatis canescentibus 5-7-nerviis 1-glandulosis, involucello calycem superante. 5 in Brasiliâ et Porto-Ricco.
13. *U. MICROCARPA*, foliis ovatis rhombeo-subtrilobis supernè glabriusculis subtus canescentibus eglandulosis trinerviis, fructibus echinatis, setis apice uncinulatis. — in ins. Caribæis? (v. s.)
14. *U. MONOPETALA* (Lour. coch. 2. p. 508.) foliis ovato-lanceolatis serratis tomentosis, involucello calyce breviori, corollâ monopetalâ. 5 in Cochinchinâ.
15. *U. RIBESIA* (Smith in. Rees. cycl. 37. n. 5.) foliis acutè 3-lobis basi rotundatis cordatis subtus 1-glandulosis villosis, involucelli foliolis spathulatis obtusis. — in Surinamo et ins. Barbadosibus. Sloan. jarn. 1. t. 11. f. 2.

* *Foliis profundè 3-5-lobatis, sinubus dilatatis, lobis sinuato-lobatis.*

16. *U. SWARTZII*, foliis trifidis puberulis subtus pallidis 1-glandulosis, fructibus glabriusculis setis echinatis. 5 in ins. Caribæis, Surinamo. *U. sinuata* Swartz! obs. 264. *U. Americana* Smith in Rees. cycl. 37. n. 4. non Lin. Sequentibus valdè affinis. (v. s.)
17. *U. SINUATA* (Lin. sp. 974.) foliis trifidis puberulis subtus pallidis 3-glandulosis, lobis angulatis denticulatis obtusis. 5 in Indiâ orientali. — Rheed. mal. 10. t. 2. — Cav. diss. 6. t. 185. f. 2. *U. aculeata* Mill. dict. n. 2.
18. *U. PARADOXA* (H. B. et K. nov. gen. am. 5. p. 278.) foliis profundè trifidis basi triglandulosis suprâ puberulis subtus tenuiter incano-tomentosis, laciniis argutè serrulatis, intermediâ rhombeâ profundè sinuato-trilobâ, lateralibus minoribus 2-3-lobis, alterâ sæpius oblitteratâ. 3 prope Atures ad ripam Orinoci. Flores axillares solitarii rosei; caps. echinatæ. *U. sinuatæ* affinis.
19. *U. MORIFOLIA*, foliis 5-lobis subtus pallidis velutinis 1-glandulosis supernè subpuberulis, sinubus serratis valdè dilatatis, fructibus densè tomentosis setisque echinatis. 5 in insulis Amicorum. Folia juniora densè tomentosa. (v. s.)
20. *U. MURICATA*, foliis 5-lobis hirsutis subtus pallidis 1-glandulosis, sinubus vix dilatatis denticulatis, fructibus tuberculis minimis submuricatis. — in Bengalo. (v. s. comm. à cl. Lechenault. ex hort. Calc.)
21. *U. HETEROPHYLLA* (Smith in Rees. cycl. 37. n. 8.) foliis profundè 5-lobis subtus 1-glandulosis canescenti-pubescentibus, sinubus latis, lobis mediis 3-lobis, superioribus elongatis basi contractis. 5 in Indiâ orientali. — Burm. zeyl. 150. t. 69. f. 2. Pluck. alm. t. 74. f. 1.

VIII. PAVONIA. Cav. diss. 3. p. 132. Lam. ill. t. 585.

Calyx cinctus involucello 5-15-phylo. Stigmata 10. Carpella capsularia 5 bivalvia 1-sperma.

Sect. I. TYPHALEA.

Carpella echinata spinis rigidis retrorsum scabris.

1. *P. SPINIFEX* (Willd. sp. 3. p. 854.) foliis ovatis acuminatis subcordatis duplicato-dentatis, pedicellis axillaribus 1-floris. 3 in Americâ calidiore. *Hibiscus spinifex* Lin. sp. 978. Flores lutei. (v. s.)

- α. *ovalifolia*, foliis ovatis, corollis calyce duplò triplòve majoribus. P. spinifex. Cav. diss. 3. p. 133. t. 45. f. 2.
- β. *aristata* (Cav. diss. 3. p. 133. t. 45. f. 3.) foliis subcordatis, corollis calyce ferè minoribus.
- γ. *oblongifolia* (fl. mex. ic. ined.), foliis oblongis, corollis calycis longitudine.
- δ. *grandiflora* (fl. mex. ic. ined.) foliis cordatis, corollis calyce duplò triplòve majoribus. An hic latent quædam species confusæ?
2. P. *TYPHALEA* (Cav. diss. 2. p. 134. et 6. t. 197.) foliis oblongo-lanceolatis dentatis, pedunculis axillaribus subterminalibus multifloris, floribus congestis. ♂ in pascuis Jamaicæ et Guianæ. Urena typhalea Lin. mant. 258. Swartz! obs. 294. Flores albi parvi. (v. s.)
3. P. *TYPHALÆOIDES* (H. B. et K. nov. gen. am. 5. p. 279.) foliis lanceolato-oblongis acutis basi obtusis serrato-crenatis ramisque pilosis, floribus terminalibus capitato-congestis, involucello 5-phylo, carpellis longè 3-aristatis. ♀ inter S.-Ana et Mariquita Novo-Granatensium. Priori affinis, sed folia multò minora nempe subquadripollicaria.
4. P. *URENS* (Cav. diss. 3. t. 49. f. 1. et 5. p. 283.) foliis 7-angularibus acuminatis dentatis hirtis, floribus axillaribus subsessilibus glomeratis. ♂ in insularum Mauritiæ et Borboniæ herbosis frigidis. Jacq. ic. rar. 3. t. 522. (v. s.)
5. P. *LEPTOCARPA* (Cav. diss. 6. p. 351.) foliis lanceolatis serratis, floribus sessilibus glomeratis, involucello 5-phylo, stylis 5. ♂ in Surinamo. Urena leptocarpa Lin. f. suppl. 508. An Hibisci species ob stylos quinque?

Sect. II. MALACHE. Trew. ehr. 50.

Carpella inermia. Involucellum calyce brevius, 5-15-phyllum.

6. P. *HASTATA* (Cav. diss. 3. p. 138. t. 47. f. 2.) foliis lanceolatis hastatis dentatis, pedicellis axillaribus 1-floris, involucello 5-phylo. ♂ in Brasiliâ. Flores rubri venis saturatioribus. (v. s.)
7. P. *COCCINEA* (Cav. diss. 3. p. 140. t. 47. f. 1.) foliis cordatis trilobis serratis, pedicellis 1-floris axillaribus adscendentibus, involucello 5-phylo. ♂ in S.-Dominicâ. — Plum. ed. Burm. t. 169. f. 2. Flores coccinei 2 poll. diam.
8. P. *COLUMELLA* (Cav. diss. 3. p. 138. t. 48. f. 3.) foliis quinquangularibus, lobis dentatis acuminatis, pedicellis axillaribus 1-floris petiolo multò brevioribus, involucello 5-phylo. ♂ in ins. Borboniæ. Columella Malvinda Comm. mss. Flores ex albo pupurascens. (v. s.)
9. P. *PARVIFLORA* (Desf. h. par. Poir. suppl. 4. p. 335.) foliis quinquangularibus subpilosis, lobis lanceolatis ovatisve, pedicellis axillaribus 1-floris longitudine petiolorum petiolisque pilosissimis, involucello 5-phylo. ♂ Patr. ign. Flores albo-flavescentes.
10. P. *SIDÆFOLIA* (H. B. et K. nov. gen. am. 5. p. 283.) foliis ovatis subacutis cordatis crenato-serratis suprâ hirtellis subtus incano-tomentosis, floribus axillaribus terminalibusque pedunculatis, involucelli foliolis 5-7-lanceolatis. ♀ inter Angosturam et Trapiche de Ferreras. Rami hirsuti. Petala flava.
11. P. *RACEMOSA* (Swartz. fl. ind. occ. 2. p. 1215.) foliis ovatis subcordatis acuminatis subserratis, racemo terminali nudo, involucello 8-phylo. ♀ in paludosis Jamaicæ et Guadalupæ. Radices longissimæ suprâ terram sæpè arcuatæ ut Rhizophoræ. P. spicata Cav. diss. 3. t. 46. f. 1. (v. s.)
12. P. *SPECIOSA* (H. B. et K. nov. gen. am. 5. p. 281. t. 477.) foliis ovato-ellipticis acutis cordatis denticulatis densissimè hirtis subtus canescentibus, floribus axillaribus subterminalibusque breviter pedunculatis, involucelli foliolis 7-9 lanceolato-spathulatis. ♀ prope Angostura. Petala violacea ad unguem purpurea subsesquipollicaria.
13. P. *PAPILIONACEA* (Cav. diss. 3. p. 140. t. 49. f. 2.) foliis subrotundo-cordatis

acutis dentatis, pedicellis axillaribus 1-floris petioli longitudine, style adscendente, involucllo 10-phylo. 5 in ins. Taheiti.

14. *P. PELLITA* (H. B. et K. nov. gen. am. 5. p. 282.) foliis ovatis acutis obsolete cordatis crenulatis molliter hirtis subtus fusciscenti-canescensibus, floribus axillaribus terminalibusque breviter pedunculatis, involuclli foliolis 11 lineariformibus calyce dimidiò brevioribus. 5? in humidis prope Maypure ad fl. Orinoci. Petala rosea basi violacea. Affinis *P. speciosa*.
15. *P. SESSILIFLORA* (H. B. et K. nov. gen. am. 5. p. 281.) foliis ovatis acutis subcordatis serratis suprâ hirtis subtus molliter hirtis pilosis canescensibus, floribus axillaribus subsessilibus, involuclli foliolis 11 apice uncinato-reflexis, vix calyce brevioribus. 5? in temperatis Novæ-Granatæ prope S.-Anna. Petala crocea? Carpella 5? hispida.
16. *P. PRÆMORSA* (Willd. spec. 3. p. 833.) foliis latè obovatis truncatis apice crenatis, pedicellis axillaribus 1-floris folio longioribus, involucllo 14-phylo. 3 ad cap. Bonæ-Spei. *P. cuneifolia* Cav. diss. 3. t. 45. f. 1. *Hibiscus præmorsus* Lin. f. suppl. 309. Jacq. ic. rar. 1. t. 141. (v. v.).

Sect. III. CANCELLARIA.

Carpella inermia, Involucllum 8-15-phyllum, calyce longius.

17. *P. MUTISI* (H. B. et K. nov. gen. am. 5. p. 283.) foliis ovatis subacuminatis cordatis serratis mollissimis, suprâ pubescentibus, subtus canescensibus tomentosis, floribus axillaribus terminalibusque pedunculatis, involuclli foliolis 8 calyce parum longioribus. 5 in montosis Novæ-Granatæ. *Hibiscus cordifolius* Lin. f. suppl. 309. Rami tomentosi. Fruct. ign.
18. *P. MOLLIS* (H. B. et K. l. c.) foliis subrotundo-ovatis cordatis acuminatis obsolete trilobis serratis hirtis subtus canescensibus mollibus, floribus axillaribus terminalibusque pedunculatis, involuclli foliolis 8 linearibus calyce duplò ferè longioribus. 3? in temperatis Novæ-Granatæ. Rami viscosi pilosi. Flores violacei.
19. *P. PANICULATA* (Cav. diss. 3. p. 135. t. 46. f. 2.) foliis cordatis subrotundis acuminatis crenato-dentatis interdum tricuspidatis, floribus paniculatis, genitalibus declinatis, involucllis 8-9 phyllis. 3 in Peru ad origin. flum. Maragnon et prope Caracas ad fl. Anauco. H. B. et K. nov. gen. am. 5. p. 280. *Urena fetida* L'her! mss. Flores flavi. (v. s.)
20. *P. CORYMBOSA* (Willd. sp. 3. p. 836.) foliis cordatis angulatisve serratis glabris, floribus corymbosis, pedunculis involucllisque polyphyllis pilosis. 3 in Jamaica et Hispaniolæ ripis fluviorum, ad Sanctam-Martham. Bertero. *Althæa corymbosa* Swartz. fl. ind. occ. 2. p. 1213. Flores lutei. (v. s.)
21. *P. ODORATA* (Willd. sp. 3. p. 837.) foliis ovatis subcordatis tricuspidatis subdentatis ramisque piloso-viscosis, pedicellis axillaribus 1-floris apice subracemosis, involucllis polyphyllis. — in Indiâ orientali. Flores rubri.
22. *P. SIDAÏDES* (Horn. h. hafn. 2. p. 658.) foliis subrotundo-cordatis, inferioribus apice subserratis obtusis, superioribus angulatis acutis, pedicellis 1-floris subsolitariis petiolo subbrevioribus, involucris 12-phyllis. ② in Bengalo. Tota piloso-viscosa. (v. s. ex hort. Cal.)
23. *P. ZEYLANICA* (Willd. sp. 3. p. 838. excl. syn. Seb.) foliis inferioribus subrotundo-cordatis crenatis, ceteris 3-5-lobatis, pedicellis axillaribus 1-floris, involucllis 10-phyllis setaceis ciliatis. ① in Zeylonâ. *Hibiscus Zeylanicus* Lin. sp. 981. *Hibiscus arenarius* Scop. del. ins. 3. t. 2.
- a. *Burmanni*, foliis 3-5-fidis, pedicellis folio brevioribus. — Burm. ind. t. 48. f. 2. (v. s.)
- β. *Cavanillesii*, foliis 3-5-partitis, pedicellis folio longioribus. — Cav. l. c. t. 48. f. 2. (v. s.)
24. *P. CANCELLATA* (Cav. diss. 3. p. 135.) foliis sagittato-cordatis dentatis, pedicellis axillaribus 1-floris petiolo longioribus, involucllis 20-phyllis pilosis

setaceis. ① in Surinamo, Guianâ Gallicâ et ad Caracas. Hibiscus cancellatus Lin. f. suppl. t. 311. (v. s. comm. à cl. Perrottet.)

IX. MALVAVISCUS. Dill. elth. 210. Cav. diss. 3 p. 131. — Achania. Swartz. prod. 2.

Calyx cinctus involucello polyphylo. Petala erecta convoluta. Stigmata 10. Carpella 5 baccata 1-sperma, nunc subdistincta, sæpius in fructum 5-locularem connata.

Sect. I. ACHANIA.

Petala basi hinc auriculata.

1. *M. CANDIDUS* (fl. mex. ic. ined.) foliis cordatis 5-fidis subacutis, lobo medio productiore, involucellis 10-12-phyllis subpatulis. 5 in Mexico.
2. *M. ARBOREUS* (Cav. diss. 3. t. 48. f. 1.) foliis cordatis 3-5-lobis acuminatis scabriusculis, foliolis involucelli erectis. 5 in petrosis calcareis Jamaicæ, Novæ-Granadæ et Mexici. Achania Malvaviscus. Swartz. fl. ind. occ. 2. p. 1222. Hibiscus Malvaviscus Lin. sp. 978. — Dill. elth. t. 170. (v. v.)
3. *M. MOLLIS*, foliis cordatis subtrilobis acuminatis molliter tomentosis, foliolis involucri subpatulis. 5 in Mexico. Achania mollis Ait! h. Kew. ed. 1. v. 2. p. 459. (v. v.)
4. *M. PILOSUS*, foliis cordatis crenatis obtusis aut acuminatis, ramulis petiolisque pilosis. 5 in fruticetis Jamaicæ australis. Achania pilosa Swartz! fl. ind. 2. p. 1224. (v. s.)
5. *M. BALBISII*, foliis cordiformibus acuminatis utrinquè scabris, nervis subtus petiolis ramulisque tomentoso-hirsutis, pedicellis petiolo duplò longioribus. 5 Patr. ign. M. cordatus Balb! herb. Planta speciosissima; involucelli foliola linearia acuta integra. (v. s. in h. Balb.)
6. *M. PENDULIFLORUS* (fl. mex. ic. ined.) foliis ovatis subcordatis acutis serratis, pedicellis gracilibus incurvo-nutantibus, involucellis erectiusculis. 5 in Mexico.
7. *M. PENTACARPUS* (fl. mex. ic. ined.) foliis cordato-ovatis acutis serratis vix subtrilobis, pedicellis erectis, involucellis linearibus, carpellis 5 baccatis subdistinctis. 5 in Mexico. Atlas-zopillin. Hern. mex. 117. ic.
8. *M. CILIATUS*, foliis ovatis acutis serratis, foliolis involucelli linearibus ciliatis, petalis hinc acutè auriculatis. 5 in ins. Tabago Amer. merid. Pavonia spiralis Cav. ic. 5. t. 435.
9. *M. CONCINNUS* (H. B. et Kunth nov. gen. am. 5. p. 286.) foliis ovato-oblongis acuminatis cordatis grossè crenato-serratis utrinquè puberulis, floribus 2-3 subcorymbosis, pedicellis petiolo brevioribus, involucello 7-phylo calycem æquante. 5 prope Loxam Peruvix. Petala coccinea ciliata. Ramuli pubescentes.
10. *M. GRANDIFLORUS* (H. B. et Kunth. l. c.) foliis ovato-oblongis acutis basi rotundatis vel obsolete cordatis subtrilobis serratis glabrisculis, pedunculis solitariis petiolo longioribus, involucello 8-phylo calyce brevior. 5 prope Guanaxuato Mexicanorum. Ramuli petiolique pilosi.
11. *M. ACAPULCENSIS* (H. B. et Kunth. l. c.) foliis ovatis subacuminatis profundè cordatis grossè dentatis utrinquè pilosis subtus molliter canescentibus, pedunculis solitariis petiolo brevioribus, involucello sub-7-phylo calycem subæquante. 5 prope Acapulco Mexic. an idem ac *M. pilosus*?

Sect? II. ANOTEA.

Petala basi hinc non auriculata. — Species non satis notæ, ab Hibisco stigmatibus 10, à Pavoniâ floribus convolutis, ab Achania petalis basi hinc non auriculatis distinctæ.

12? *MP. CORDATUS* (Nees et Mart. nov. act. boun. 21. p. 99. sub Achania) foliis

cordatis crenatis subtus tomentosis, involucello 8-9-phyllo calyceque setoso, genitalibus non exsertis. ♂ in Brasiliâ ad viam Felisberti. Cor. rubra. Locus in serie dubius.

13. M? FLAVIDUS (fl. mex. ic. ined.) foliis acutè quinquangularibus dentatis basi cordatis, corollis cylindraceis, genitalium columnâ erectâ petalis duplò longiore. ♀ in Mexico.
14. M? PLEUROGONUS (fl. mex. ic. ined.) foliis palmatifidis acutè 5-7-lobis dentatis basi cordatis, corollâ subcampanulatâ, genitalium columnâ longissimâ subdeclinatâ, antheris plerisque secundis. ♀ in Mexico.
15. M? PLEURANTHERUS (fl. mex. ic. ined.) foliis basi cordatis acutè 3-5-lobis palmatifidis, lobis dentatis, medio productiore, floribus subcylindraceis, genitalium columnâ petalis æquali erectâ, antheris secundis. ♀ in Mexico.

X. LEBRETONIA. Schrank. pl. rar. h. mon. t. 90.

Calyx 5-partitus involucello 5-partito subbreuiore cinctus. Petala 5 in parte exsertâ æstivatione contorta, limbo patente. Styli 10. Carpella 5, aut abortu 4, monosperma, indehiscentia. — An satis à Pavoniæ sect. II. diversa?

1. L. COCCINEA (Schrank l. c.) foliis ovatis acuminatis serratis, pediculis axillaribus 1-floris petiolo longioribus, corollâ involucello duplò longiore. ♂ in Brasiliâ. Truncus hispidus. Flores ampli coccinei. Folia suprâ pubescentia, subtus tomentosa.
2. L. LATIFOLIA (Nees et Mart. nov. act. bonn. XI. p. 98.) foliis ovatis subcordatis acutiusculis crenato-serratis, pedicellis 1-floris petiolo longioribus, corollâ involucellum subæquante. ♂ in Brasiliâ. Cor. coccinea; calyx ut in priore virescens; folia pubescentia.
3. L? SEMISERRATA, foliis oblongis apice serratis coriaceis longè petiolatis, floribus terminalibus, calycibus coloratis persistentibus. ♀ in Brasiliâ. Schouwia semiserrata. Schrad. goett. ann. 1821. p. 717. An prioris, an Gerthæ congener? Characteres indicati non sufficiunt.

XI. HIBISCUS. Lin. gen. n. 846. Lam. ill. t. 584.

Calyx cinctus involucello sæpius polyphyllo, rariùs foliolis paucis aut inter se coalitis. Petala hinc non auriculata. Stigmata 5. Carpella in capsulam 5-locularem coalita, valvis intus medio septiferis, loculis polyspermis aut rariùs 1-spermis. — An genus dividendum in genera tot quot sectiones infrâ memoratæ.

Sect. I. CREMONTIA. Comm. ined.

Corollæ (ut in Malvavisco) convolato-cylindraceæ, sed petala non auriculata, stigmata 5 et fructus loculi oo-spermi.

1. H. LILIFLORUS (Cav. diss. 3. p. 154. t. 57. f. 1.) foliis lanceolato-oblongis integris rariùs trifidis, involucello 5-phyllo, calyce 5-dentato, petalis extus subvelutinis. ♀ in ins. Borboniæ montibus sylvaticis. Malvaviscus puniceus Bory! ined. (v. s.)
2. H. BORYANUS, foliis ovatis subdentatis indivisis 5-nerviis glabris, involucello 5-7-phyllo calyce 5-dentato paulò longiore, petalis oblongis extus sub lente subvelutinis. ♀ in ins. Borboniâ. Bory Saint-Vincent. Flores albidii brevis-simè pedicellati. (v. s.)
3. H. FRAGILIS, foliis ovatis vix subtrilobis dentatis glabris, pedicellis 1-floris folii longitudine, involucello 5-phyllo. ♂ in insulâ Borboniâ. Malvaviscus fragilis Bory! ined. (v. s.)
4. H. PEDUNCULATUS (Cav. diss. 3. p. 163. t. 66. f. 2.) foliis 3-5-lobis obtusis crenatis hirsutis, pedicellis folio duplò longioribus, involucello polyphyllo,

- corollâ subcampanulâtâ. ζ ad cap. Bonæ-Spei. Cor. rosea ex Weinm. flora 1821. p. 30.
5. *H. LAMPAS* (Cav. diss. 3. p. 154. t. 56. f. 2.) foliis cordatis tricuspidatis glabris subtus punctatis, pedicellis solitariis 1-floris petiolo longioribus, involucello minimo 5-7-setoso. ζ in ins. Philippicis. (v. s.)
6. *H. MEMBRANACEUS* (Cav. diss. 3. p. 159. t. 57. f. 2.) foliis cordatis ovato-lanceolatis acuminatis dentatis, pedicellis petiolo duplò longioribus 1-floris, involucello polyphylo, interdum 5-fido, laciniis 3-nerviis. ζ Patr. ign.
7. *H. SPIRALIS* (Cav. ic. 2. p. 47. t. 162.) foliis ovatis acutis dentatis glabris basi integerrimis, pedicellis 1-floris petiolo duplò longioribus, involucello polyphylo. ζ in Mexico. Petioli subvillosi. Cor. ex luteo incarnata. (v. s.)
8. *H. TUBIFLORUS* (fl. mex. ic. ined.) foliis cordatis acutis subsinuatis dentato-serratis villosis, pedicellis 1-floris petiolo duplò longioribus, involucello polyphylo. ζ in Mexici montibus Sancti-Hieronimi. Cor. ex flavo-rubra.
9. *H. SENEGALENSIS* (Cav. diss. 3. p. 160. t. 68. f. 1.) foliis cordatis dentatis tomentosis, inferioribus angulatis, pedicellis 1-floris petiolo brevioribus, involucro decaphyllo setaceo corollam subæquante. — in Senegaliâ. Corolla parva flavescens. (v. s.)
10. *H. TUBULOSUS* (Cav. diss. 3. p. 161. t. 68. f. 2.) foliis cordatis inæqualiter dentatis subtus cano-villosis, inferioribus sub-5-lobis, superioribus acuminatis, pedicellis 1-floris brevissimis, involucello 8-phylo, foliolis linearibus subspatulatis. (1) in Indiâ orientali. Corolla subcampanulata, flavescens, in fundo purpurea. Specimen huic simillimum ex Senegaliâ habeo. (v. s.)
11. *H. HISPIDUS* (Spreng. pug. 2. p. 73.) foliis cordatis trilobis crenatis cauleque hispidis, pedicellis 1-floris petiolo brevioribus, involucello 7-phylo subulato. \mathcal{Z} Patr. ign. Corolla alba, connivens, in fundo flavescens.
12. *H. URENS* (Lin. suppl. 309.) foliis reniformibus crenatis cauleque tomentosis, pedicellis 2-3 axillaribus, involucello 12-partito, corollâ calyce brevior. \mathcal{Z} ad cap. Bonæ-Spei. Cav. diss. 3. p. 161. t. 67. f. 1.

SECT. II. PENTASPERMUM.

Corollæ expansæ. Capsulæ valvulæ medio septiferae, loculi 1-spermi e duabus valvis compositis.

13. *H. OVATUS* (Cav. diss. 3. p. 143. t. 50. f. 3.) foliis ovatis integerrimis ciliatis 3-nerviis subtus villosis, pedicellis 1-floris petiolo duplò longioribus. ζ ad cap. Bonæ-Spei.
14. *H. HASTATUS* (Cav. diss. 3. p. 144. t. 50. f. 1.) foliis hastatis oblongis serratis angustis, pedicellis 1-floris folii ferè longitudine, fructu globoso tomentoso. — in Indiâ. — Pluk. t. 127. f. 2. (v. s. in p. Balb.)
15. *H. ACUMINATUS* (Cav. diss. 3. p. 144. t. 50 f. 2. excl. Pluk. syn.) foliis cordatis acuminatis, inferioribus tricuspidatis, pedicellis 1-floris folio brevioribus, involucello 9-phylo, fructu tomentoso depresso. ζ Patr. ign.
16. *H. PENTACARPOS* (Lin. sp. 981.) foliis cordatis oblongis dentatis obtusiusculis angulatis leviter subtrilobis glabris, pedicellis petiolo longioribus, floribus cernuis, pistillis ntantibus. \mathcal{Z} in paludibus Etruriæ, Venetiæ. Jacq. ic. rar. 1. t. 143. Corolla dilutè rubra. (v. s.)
17. *H. VIRGINICUS* (Lin. sp. 981.) foliis acuminatis inæqualiter dentatis subvillosis, inferioribus indivisis cordatis, superioribus ovato-cordatis trilobis, pedicellis petiolo longioribus, floribus cernuis, pistillis ntantibus. \mathcal{Z} in Virginix et Carolinæ paludosis. Corolla duplò major quàm præcedens. Jacq. ic. rar. 1. t. 142. — Pluk. t. 6. f. 4. (v. s.)
18. *H. PENTASPERMUS* (Bert! ined.) totus pilis patentibus hispidus, foliis cordiformibus acuminatis grossè dentatis, pedicellis axillaribus 1-floris petiolo paulò longioribus, fructu pentagono in angulis hispido. — in Jamaicâ. Bertero. Flores parvi flavescens. (v. s.)

Sect. III. MANIHOT.

Carpella seu capsulæ loculamenta polysperma. Semina glabra. Involucella 4-6-phylla. Calyx spathaceus 5-dentatus longitudinaliter ruptus.

19. *H. MANIHOT* (Lin. sp. 980.) caule inermi, foliis subglabris palmato-partitis, lobis 5-7 acuminatis grossè dentatis, involucello hispido 4-6-phylo, pedicellis floridis declinatis. 5 in Indiâ orientali. — Dill. h. elth. p. 189. t. 156. — Cav. diss. 3. p. 172. t. 63. f. 2. (v. s.)

β. palmatus (Cav. diss. 3. p. 168. t. 63. f. 1.) foliis palmatifidis, radice crassâ fungosâ. 5 in Americâ calid. — Fructus ut var. *α* pyramidatus pilosus. (v. v.)

20. *H. TIMORIENSIS*, caule inermi, foliis glabris palmatifidis 7-lobatis, lobis acuminatis serratis, involucello 5-phylo glabro. 5 in ins. Timor. *H. Manihot* var. *β.* valdè affinis, sed differt notis indicatis et fructu (ex immaturo) subgloboso nec pyramidato. (v. s.)

21. *H. PSEUDO-MANIHOT*, caule inermi, foliis 3-fidis subpuberulis, lobis ovatis acuminatis serratis, involucello hispido 4-phylo. 5 in herbosis insulæ Borboniæ ad rivum Sancti-Denisii. Bory-Saint-Vincent. Fructus et flos *H. Trioni*, sed folia 3-fida nec 5-7-partita. (v. s.)

22. *H. FICULNEUS* (Lin. sp. 978.) caule tuberculis aculeato, foliis palmato-5-lobis, summis 3-lobis, lobis obtusis inæqualiter dentatis basi angustatis, involucello 5-phylo caduco. 5 in Zeylonâ. — Dill. h. elth. t. 157. f. 190. *H. sinuatus* Cav. diss. 3. t. 52. f. 2. (v. s.)

23. *H. TETRAPHYLLUS* (Roxb. ex Horn. h. hafn. 561.) caule herbaceo, foliis 5-lobis serratis, involucello 4-5-phylo. ① in Indiâ orientali.

Sect. IV. KETMIA.

Carpella seu capsulæ loculamenta polysperma. Semina glabra. Corollæ expansæ. Involucella 5-7-phylla. Calyx 5-lobus non longitudinaliter ruptus.

24. *H. MICANS* (Cav. diss. 3. p. 167. t. 60. f. 2.) caule inermi, foliis cordatis quinquangularibus dentatis acuminatis, involucello 6-partito. 5 in Javâ.

25. *H. CALYGINUS* (Willd. sp. 3. p. 817.) caule inermi, foliis cordatis angulatis subtrilobis repandis, involucello 5-phylo petalis longiore. 5 in ins. Borboniâ. *H. calyphyllus* Cav. diss. 5. p. 283. t. 140.

26. *H. COLUMNARIS* (Cav. diss. 3. p. 166. t. 59. f. 2.) caule inermi, foliis cordatis quinquangularibus obtusiusculis repandis, involucello sub-6-fido, columnâ stamineâ corollam campanulatam superante. 5 in ins. Borboniâ.

27. *H. ACERIFOLIUS*, caule inermi, foliis cordatis 5-lobatis hirtis, lobis acuminatis subrepandis, involucello 6-7-phylo setaceo. 5 in Indiâ orientali. *Pavonia platanifolia* Willd. berl. mag. 1810. p. 220. *Pavonia acerifolia* Link et Otto. abb. 1. p. 5. t. 1. Styli 5, ergo non est *Pavonia*. Fructus ignotus.

28. *H. ROSA-SINENSIS* (Lin. sp. 977.) caule inermi arboreo, foliis ovatis acuminatis glabris basi integerrimis apice grossè dentatis subincisis, pedicellis folii longitudine, involucello 7-phylo. 5 in Indiâ orientali. Variat flore purpureo albo et flavo, sæpius in hortis pleno. Cav. diss. 3. t. 69. f. 2. — Rheed. mal. 2. t. 16. (v. v.)

29. *H. SYRIACUS* (Lin. sp. 978.) caule inermi arboreo, foliis cuneiformi-ovatis trilobis dentatis, pedicellis petiolo vix longioribus, involucello 6-7-phylo. 5 in Syriâ, Carniolâ. Variat flor. purpureis, albis, rubris, variegatis, sæpe plenis. Cav. diss. 3. p. 69. f. 1. (v. v.)

30. *H. PRURIENS* (Roxb. cat. calc. 51.) caule herbaceo piloso, foliis longè petiolatis ovatis subtrilobis serratis membranaceis glabriusculis, pedicellis brevissimis, involucello et calyce basi pilosissimis corollam æquantibus utrinquè 5-partito acuminato. ① ② in Indiâ orientali. Horn. h. hafn. suppl. p. 78. (v. s.)

Sect. v. FURCARIA.

Carpella polysperma. Semina glabra. Involucelli foliola apice bifurcata aut hinc dente grosso appendiculata.

31. *H. SURRATTENSIS* (Lin. sp. 979.) caule herbaceo aculeis recurvatis scabro, stipulis semi cordatis, foliis palmato-5-lobis, pedicellis petioli longitudine, involucelli foliolis hinc obtusè appendiculatis. $\bar{\imath}$ in Indiâ orientali. Cav. diss. 3. t. 33. f. 1. — Rumph. amb. 4. t. 16. Flos luteus fundo atro purpureo. (v. s.)
32. *H. RADIATUS* (Cav. diss. 3. p. 150. t. 54 f. 2.) caule suffruticoso aculeis rigidis scabro, stipulis lanceolatis, foliis 5-7-partitis, lobis lanceolatis acuminatis serratis, pedicellis brevissimis inermibus, involucelli foliolis rigidè ciliatis vix appendiculatis. $\bar{\imath}$ in Indiâ orientali. Sims. bot. mag. t. 1911.
33. *H. FURCATUS* (Roxb. cat. calc. p. 31.) caule herbaceo petiolis calycibusque tuberculato-muricatis, foliis basi subovatis trifidis, inferioribus 5-fidis, lobis acuminatis serratis, involucelli foliolis 9 bifidis. $\bar{\imath}$ in Bengalo. Willd. enum. 736? (v. s.)
34. *H. SCABER* (Michx. fl. bor. am. 2. p. 45.) caule herbaceo scabro, foliis asperis subrotundis basi truncatis, superioribus palmati lobatis, lobis supernè dilatatis crenatis, floribus sessilibus, calycibus hispidissimis, involucelli foliolis furcellatis. $\bar{\imath}$ in paludibus Carolinæ et Georgiæ. *H. aculeatus* Walt. cat. 177. Flores ampli sulfurei. (v. s.)
35. *H. FURCELLATUS* (Desrous. dict. enc. 3. p. 358.) caule fruticoso, ramis foliisque tomentosis, foliis cordatis subangulatis, involucelli foliolis 10 cylindraceis apice furcellatis, calycibus hispidis. $\bar{\imath}$ in Guianâ. (v. s. comm. à cl. Perrottet.)
36. *H. DIONON*, caule fruticoso, ramis foliisque tomentosis, foliis cordatis 3-5-fidis lobo medio acuminato, involucelli foliolis 10 cylindraceis apice furcellatis, calycibus hispidis. $\bar{\imath}$ in Cayennâ. (v. s.)
37. *H. UNGINELLUS* (fl. mex. ic. ined.) caule fruticoso, ramulis retrorsum pilosis, foliis palmatifidis 3-5-lobis basi cordatis dentatis. involucelli foliolis 10 appendice dorsali uncinato instructis. $\bar{\imath}$ in Mexico. Flores ampli vix aperti violaceo-purpurei.
38. *H. BICORNIS* (Meyer esseq. 231.) caule fruticoso petiolisque aculeatis, foliis 3-5-lobis auriculato-lanceolatis serratis, involucelli foliolis calyce longioribus apice bilobis. $\bar{\imath}$ in pascuis et sylvis Guianæ. Corolla rosea fundo purpureo calyce sexies minor.
39. *H. BIFURCATUS* (Cav. diss. 3. p. 146. t. 51. f. 1.) caule fruticoso versùs apices ramorum scabro, pedunculis nervisque foliorum subtùs aculeis reflexis scabris, foliis 3-5-lobis acuminatis serratis, involucelli foliolis 10-12 apice bifidis. $\bar{\imath}$ in Brasiliâ et Porto-Ricco. Cor. magna purpurascens. (v. s.)

Sect. VI. ABELMOSCHUS *Med. malv. p. 45.*

Carpella polysperma. Semina glabra aut in dorso lineâ subvillosâ. Corollæ expansæ. Involucella foliolis 8-15 integris constantia.

§. 1. *Caule tuberculis aculeato.*

40. *H. LAMBERTIANUS* H. B. et K. nov. gen. am. 5. p. 291. t. 478.) caule aculeato simplici herbaceo, foliis ovato-lanceolatis acuminatis basi rotundatis serratis suprâ hirtis subtùs canescenti-tomentosis, involucello 11-phylo. $\bar{\imath}$? in humidis prov. Caracasanæ Flores ampli purpurei. Ovarii loculi oo-spermi. Stigma peltato-convexum.
41. *H. TRILOBUS* (Cav. diss. 3. p. 147. t. 53. f. 2. excl. syn. Rheed.) caule arboreo aculeato, pedicellis inermibus, foliis cordatis 3-lobis serratis, lobo intermedio longiore, involucello 12-phylo. $\bar{\imath}$ in paludosis ius. Sanctæ-Dominicæ. Plum. ed. Barm. t. 159.
42. *H. DIVERSIFOLIUS* (Jacq. ic. var 3 t. 551.) caule petiolisque aculeatis, pedicellis brevibus inermibus pilosissimis, foliis 3-5-lobis obtusis dentatis,

- superioribus oblongo-lanceolatis indivisis, involucello 9-phyll. 5 in Indiâ orientali. *H. ficulneus* Cav. diss. 3. t. 51. f. 2. Cor. flava fundo atrovioleaceo. (v. s.)
43. *H. MACULATUS* (Desrous. in dict. enc. 3. p. 349.) caule petiolisque aculeatis, foliis inferioribus palmati-partitis dentatis, superioribus ovatis sub-3-lobis, involucello polyphyll. calyce hispido maculato. 5 in Sancto-Domingo.
44. *H. HETEROPHYLLUS* (Vent. malm. t. 103.) caule fruticoso tuberculis aculeato, foliis lineari-lanceolatis acuminatis plerumque lobatis aculeato-serratis, involucello 10-phyll. 5 in Novâ-Hollandiâ. Flores magni albi fundo purpureo. *H. grandiflorus* Salisb. par. t. 22. (v. v.)
45. *H. CANNABINUS* (Lin. sp. 979.) caule aculeato, foliis palmato-5-partitis subtus 1-glandulosus, floribus subsessilibus, calyce glanduloso-piloso. ① in Indiâ orientali. Flores magni lutei fundo fusco. Cav. diss. 3. p. 148. t. 52. f. 1. *H. vitifolius* Mill. (v. s.)
46. *H. VITIFOLIUS* (Lin. mant. 569.) caule subaculeato, foliis glabriusculis dentatis 5-angularibus acuminatis, floribus cernuis, capsulis pentapteris pilosis. ① in Indiâ orient. Rheed. mal. 6. t. 46. Cav. diss. 3. p. 145. t. 58. f. 2. Cor. flava fundo atropurpureo. (v. s.)
47. *H. OBTUSIFOLIUS* (Willd. sp. 3. p. 829.) caule subaculeato, foliis subtus tomentosis crenatis cordatis, inferioribus subrotundis, superioribus acuminatis 3-lobis obtusis, capsulis pentapteris pilosis. ① in Indiâ orient. Affinis *H. vitifolio* et cum eo forsân ob capsulas pentapteras sectionem propriam constituens. (v. s.)
48. *H. HETEROTRICHUS*, caule hirsuto setisque sparsis hispido, foliis cordatis acutè 5-lobis dentatis, summis trilobis calycibusque hirsutis, pedicellis petiolo brevioribus, floribus cernuis. — Patr. et fruct. ign. flores omninò duorum priorum. (v. s.)

§. 2. *Caule inermi.** *Species annuæ.*

49. *H. ESCULENTUS* (Lin. sp. 980.) foliis cordatis 5-lobis obtusiusculis dentatis, petiolis flore longioribus, involucellis 10-phyllis deciduis, calycibus longitudinaliter rumpentibus. ① in Indiis culta. Cav. diss. 3. t. 61. f. 2. Fructus pyramidatus sulcatus. An *H. Bamia* Targ. satis differt? (v. s.)
50. *H. LONGIFOLIUS* (Willd. sp. 3. p. 827.) foliis 3-5-partito-palmatis, lobis lanceolatis inciso-dentatis, petiolis flore longioribus, involucellis sub-10-phyllis deciduis, calycibus longitudinaliter rumpentibus. ① in Indiâ orientali.

** *Species perennes.*

51. *H. MOSCHEUTOS* (Lin. sp. 975.) foliis ovatis acuminatis serratis subtus tomentosis, petiolis pedunculisque inter se coalitis, involucellis calycibusque tomentosis, capsulis glabris. 2 in Amer. boreali. Flos albus maximus in fundo purpureus. Cav. diss. 3. t. 65. f. 1. Variat flore pallidè purpureo. Bot. mag. t. 882. (v. v.)
52. *H. PALUSTRIS* (Lin. sp. 976.) foliis ovatis dentatis subtrilobis subtus cano-tomentosis, pedicellis axillaribus à petiolo liberis suprâ medium articulatis. 2 in Amer. borealis paludibus. Cor. amplæ roseæ albæ aut flavidæ. Cav. diss. 3. t. 65. f. 2. (v. s.)
53. *H. ROSEUS* (Thorl. in Lois. fl. gall. 2. p. 434.) foliis cordatis dentatis subtrilobis subtus cano-tomentosis, pedicellis axillaribus à petiolo liberis 1-floris suprâ medium articulatis. 2 in Vasconiâ ad ripas flum. Adour. DC. fl. fr. suppl. p. 626. Cor. roseæ. (v. v.)
54. *H. AQUATICUS* (DC. fl. fr. suppl. 627.) foliis ovatis dentatis subtrilobis subtus cano-tomentosis, pedicellis axillaribus à petiolo liberis 1-floris prope basim articulatis. 2 in Etruriæ paludibus. Flores albi. *H. palustris*. Savi cent. 1. p. 126. (v. s.)

55. *H. INCANUS* (Willd. sp. 3. p. 807.) foliis ovatis acuminatis obtusè serratis utrinquè incano-tomentosis, pedicellis axillaribus 1-floris à petiolo liberis medio circiter articulatis. ♀ in Carolinâ. Flor. ampli sulfurèi. Wendl. h. herr. 4. t. 24. (v. s.)
56. *H. MILITARIS* (Cav. diss. 6. p. 352. t. 198. f. 2.) foliis 3 lobo-hastatis acuminatis serratis utrinquè cauleque glabris, pedicellis medio articulatis, corollâ subcampanulatâ, capsulis ovatis acuminatis glabris, seminibus holo-sericeis. ♀ ad ripas fluminum Americæ bor. Sims. bot. mag. t. 2385. *H. lævis* Scop. del. 3. t. 17. *H. hastatus* Michx. *H. riparius* Pers. Cor. purpureæ. An potiùs *Bombicellæ* seu *Pentaspermi* sp. (v. s.)
57. *H. SPECIOSUS* (Ait. h. Kew. 2. p. 456.) foliis palmato-5-partitis cauleque glabris, lobis lanceolatis acuminatis apice subserratis, pedicellis sub apice articulatis, corollâ patente, capsulâ glabrâ pentagono-ovatâ. ♀ ad ripas fluminum in Carolinâ austr. et Floridâ. *H. coccineus* Walt. fl. car. 177. — Bot. mag. t. 360. Flores ampli coccinei. (v. v.)
58. *H. GRANDIFLORUS* (Michx. fl. bor. am. 2. p. 46.) foliis triangulato-cordatis trilobis coriaceis utrinquè eximie tomentosis subtus incanis, capsulis tomentoso-hirsutissimis subtruncatis. ♀ in paludosis salsis Georgiæ et Floridæ. Flores ampli carnei.

*** *Species frutescentes.*

59. *H. LASIOCARPUS* (Cav. diss. 3. p. 159. t. 70 f. 1.) foliis ovato-lanceolatis acutis serratis tomentosis, pedicellis axillaribus, calycibus tomentosis, involucello ciliato, capsulâ setosâ 5 Patr. ign.
60. *H. FERRUGINEUS* (Cav. diss. 3. p. 162. t. 60. f. 1.) foliis cordatis parabolicis obtusis dentatis scabris, pedicellis axillaribus petiolo longioribus, caule tomentoso-ferrugineo, capsulâ setosâ. 5 in Madagascar. (v. s.)
61. *H. SULFUREUS* (H. B. et Kunth nov. gen. am. 5. p. 289. *) foliis oblongis utrinquè obtusis integerrimis suprâ glabriusculis subtus tenuissimè canescenti-tomentosis, involucello 10-phylo calyce breviorè, ovarii loculis sub-4-spermis, stigmate multifido capitato. 3 ad Caracas. Fruct. ign.
- ℓ. acutifolius*, foliis majoribus acutis, capsulâ subglobosâ acutâ piloso-sericâ. 3 5 in monte Cocollar Novæ Andalusie.
62. *H. AFFINIS* (H. B. et Kunth l. c.) foliis oblongis acutis basi obtusis integerrimis suprâ puberulis subtus incano-tomentosis, involucello 9-11-phylo calycem æquante, ovarii loculis 4-5-spermis, stylo 3-5-fido. 3 in Americâ merid. inter Angosturam et Trapiche de Farreras. Petala ut in priore sulfurea basi purpurea.
63. *H. CAVANILLESIIANUS* (H. B. et Kunth l. c.) foliis subrhomboidèis ovato-oblongis angustato-acuminatis basi acutis grossè serratis utrinquè pube stellatâ pilosiusculis, ovarii loculis sub-4-spermis, stylo 5 fido. 3 in ripâ fl. Amazonum prope Tomependa. Flores violacei. Capsula ex Bonpl. depresso-ovata.
64. *H. LUNARIFOLIUS* (Willd. sp. 3. p. 811.) foliis subrotundo cordatis acuminatis argutè dentatis subtus hirtis, pedicellis incrassatis villosò-hirtis, involucello calycis longitudine. 5 in Indiâ orient.
65. *H. BRASILIENSIS* (Lin. sp. 977.) foliis cordatis denticulatis, ramis petiolis pediculis calycibusque hirtis, involucello calyce duplò longiore. 5 in Brasiliâ. †.
66. *H. ÆTHIOPICUS* (Lin. mant. 258.) foliis cuneiformibus sub-5-dentatis stellato-pilosis, pedicellis folio longioribus, involucello 8-10-phylo hispido. 3 ad cap. Bonæ Spei. Cav. diss. 3. t. 61. f. 1.
67. *H. MICROPHYLLUS* (Vahl symb. 1. p. 50.) foliis ovalibus hirtis anticè serratis sub-5-nerviis, pedicellis longitudine foliorum, involucello 10-phylo setaceo calyce longiore. 5 in Arabiâ felic. *H. flavus* Forsk. descr. 126. An hujus sect.

68. *H. VELUTINUS*, foliis cordatis obtusè dentatis utrinquè molliter cano-velutinis, caule suffruticoso, pedicellis petiolo paulò longioribus, involucelli foliolis 8-10 linearibus apice sublatioribus, capsulà subrotundà hispidissimà. 5 in Timor. (v. s.)
69. *H. SORORIUS* (Lin. f. suppl. 311.) foliis cordatis crenatis scabris, caule suffruticoso piloso, involucelli 10-phylli foliolis teretibus apice dilatatis. 5 in Surinamo. An hujus sect?
70. *H. MUTABILIS* (Lin. sp. 977.) foliis cordatis angulatis 5 lobis acuminatis dentatis ramisque subtomentosis, pedicellis folii ferè longitudine, involucello 8-10 phyllo, calycis lobis elongatis 5-nerviis. 5 in Indià orientali. Cav. diss. 3. t. 62. f. 1. *H. Sinensis* Mill. dict. n. 2. Corolla matutino tempore alba, meridie pallido-incarnata, vespere rosea. (v. s.)
71. *H. CLYPEATUS* (Lin. sp. 980.) foliis cordatis angulatis parè dentatis subglabris, ramis velutinis, pedicellis petiolo brevioribus, involucello 8-9-phylo, calycis lobis ovali-oblongis 3-nerviis, capsulà turbinatà truncatà hispidà. 5 in fruticetis Jamaicæ et Sancti-Domingi. Cav. diss. 3. t. 58. f. 1. Flores ex rubro flavescentes. (v. s.)
72. *H. ABELMOSCHUS* (Lin. sp. 980.) foliis subpeltato-cordatis 7-angularibus acuminatis serratis, caule hispido, pedicellis petiolo longioribus, involucello 8-9-phylo, capsulà setosà. 5 in Indià orient. et Amer. merid. Cav. diss. 3. t. 62. f. 2. Semen moschatum. An plantæ utriusque orbis omninò similes? (v. s. f. r.)
73. *H. ARIOCARPUS*, foliis glabris basi ovatis 5-nerviis trilobis, lobis acuminatis integris, pedicellis petiolo longioribus, involucello 8-9-phylo, foliolis oblongis undulatis subdentatis, capsulà subglobosà intùs extùsque hispidissimà. 5 in Bengalo. cl. Lecheuault specimina misit in hort. Calc. lecta. (v. s. sine fl.)
74. *H. DOMINGENSIS* (Jacq. ic. rar. 3. t. 550.) foliis 3-5-lobatis dentatis, caule arboreo aculeato, pedicellis inermibus hirtis, involucello sub-12 phyllo setaceo, calyce basi complanato. 5 in Santo-Domingo. Flores magni violacei.
75. *H. STRIATUS* (Cav. diss. 3. p. 146. t. 54. f. 1.) foliis 3-lobis ~~dentatis~~ serratis glabris, caule pedicellisque aculeatis, involucello 8-9-phylo setaceo, calyce basi ovato. — Patr. ign. Flores flavi. *H. Domingensis* var. ex Wild.

Sect. VII. BOMBICELLA. — Bombyx. *Medik. malv. p. 44.*

Carpella polysperma. Semina lanata seu gossypina. Corollæ sæpissimè expansæ. Involucella 5-10-phylla.

76. *H. PHÆNICEUS* (Willd. sp. 3. p. 813.) foliis ovatis acuminatis serratis, infimis subcordatis tricuspidatis, pedicellis sub apice articulatis, involucello 10-phylo. 5 in Indià orientali. Cor. purpurascens. (v. s.)
- α. involucri calyce brevior. *H. phæniceus* Lin. f. suppl. 310. Jacq. l. vind. t. 4.
- β. involucri calycem æquante. *H. hirtus* Lin. sp. 977. Cav. diss. 3. t. 67. f. 3.
77. *H. BETULINUS* (H. B. et K. nov. gen. am. 5. p. 292.) herbaceus ramosus, foliis deltoideo-ovatis acutis basi rotundatis crenato-serratis utrinquè pilosiusculis, involucello 9-11-phylo, capsulis subglobosis, loculis oligo-spermis. 4 prope Camanum. Flores albi.
78. *H. UNILATERALIS* (Cav. diss. 3. p. 158. t. 67. f. F. e.) foliis ovatis acuminatis dentatis, pedicellis folio longioribus suprà medium articulatis, involucello 9-phylo, staminibus unilateralibus. 5 in Sancto Domingo. Plum. ed. Burn. t. 160. f. 1. — *H. columbinus*. fl. mex. ic. ined. videtur idem sed stamina non unilateralia. (v. s.)
79. *H. RHOMBIFOLIUS* (Cav. diss. 3. p. 156. t. 69. f. 3.) foliis rhombeo-ovatis in-

divisis apice crenatis acuminatis, pedicellis brevissimis, involucello 7-phylo. 5 in Indiâ orientali.

80. *H. GOSSYPINUS* (Thunb. prod. 118.) foliis ovatis serratis, petiolis cauleque hispidis, pedicellis suprâ medium articulatis pilosis folio ferè longioribus, involucello parvo 7-phylo. 5 ad cap. Bonæ Spei. — Burchl. cat. n. 2364? (v. s.)
81. *H. MICRANTHUS* (Cav. diss. 3. p. 155. t. 66. f. 1.) foliis ovatis subrotundisve indivisis serratis scabris, pedicellis folio longioribus, corollâ reflexâ, involucello 7-phylo. 5 in Indiâ orientali. *H. micranthus* et *rigidus* Lin. f. suppl. 308 et 310 ex Ait. h. kew. ed. 2. v. 4. p. 226.
82. *H. CLANDESTINUS* (Cav. ic. 1. p. 1. t. 2.) foliis oblongis subcordatis dentatis scabrinis, inferioribus obsolete trilobis, pedicellis sub apice articulatis folii longitudine, involucello 6-phylo, petalis calyce inclusis. 5 in Senegal.
83. *H. OVALIFOLIUS* (Vahl. symb. 1. p. 50.) foliis ovalibus subangulatisque hispidis, caule pilis stellatis scabro, involucello 5-phylo. 5 in montosis Arabiæ felicis. *Urena ovalifolia* Forsk. descr. 124.

Sect. VIII. TRIONUM. *Medik. malv. p. 46.*

Carpella polysperma. Semina glabra. Corollæ expansæ. Involucellum polyphyllum. Calyx demum vesiculosus inflatus.

84. *H. TRIONUM* (Lin. sp. 981.) foliis dentatis, inferioribus subindivisis, superioribus tripartitis, lobis lanceolatis intermedio longissimo, calycibus inflatis membranaceis nervosis. ① in Italiâ et Carnioliâ. Flores flavi cum fundo atro-purpureo. Cav. diss. 3. t. 64. f. 1. Curt. bot. mag. t. 209. (v. v.)
- β ? *ternatus* (Cav. diss. 3. p. 172. t. 64. f. 3.) foliis ferè omnibus 3-partitis, lobis grossius dentatis, pedicellis petiolo vix longioribus, caule humili. ① ad cap. Bonæ-Spei.
- γ . *hispidus* (Mill. dict.) foliis inferioribus 3-lobis, summis 5-partitis obtusis, caule hispido. ① ad cap. Bonæ-Spei.
- δ . *cordifolius* (Moench. suppl. 202. sub Triono) foliis radicalibus cordato-subrotundis, supremis 3-partitis. ① *H. Africanus* Rojb. Beitr. 1. p. 43.
85. *H. VESICARIUS* (Cav. diss. 3. p. 171. t. 62. f. 2.) foliis dentatis, inferioribus indivisis, superioribus 5-fidis, lobis oblongis subæqualibus obtusis, calycibus inflatis membranaceis nervosis. ① in Africâ. *H. Africanus*. Mill. dict. n. 20. Flos prioris. (v. v.)

Sect. IX. SABDARIFFA.

Capsulæ loculi polyspermi. Semina glabra. Involucellum monophyllum multidentatum, foliolis nempe basi aut ad medium inter se coalitis. Plantæ herbacæ annuæ.

86. *H. SABDARIFFA* (Lin. sp. 978.) foliis dentatis, inferioribus ovatis indivisis, superioribus trilobis basi cuneatis, floribus subsessilibus, involucello 12-dentato. ① in Indiis. Caulis inermis glaber rubescens. Folia Acetosæ sapore. Cav. diss. 3. t. 198. f. 1. Bonpl. nav. t. 29. Lois. herb. amat. t. 296. (v. s.)
87. *H. DIGITATUS* (Cav. diss. 3. p. 151. t. 70. f. 2.) foliis palmatilobis, lobis lanceolatis serratis, petiolis muricato-scabris, floribus subsessilibus solitariis, involucello 7-fido. ① in Brasiliâ.
- β . *Kerrianus*, foliis aliis 3-5-fidis, aliis ovalibus indivisis dentatis. ① ad Rio-Janeiro. *H. digitatus* Ker. bot. reg. t. 608. Cor. alba fundo et extus rubra. An à specie Cav. distincta?

Sect. X. AZANZA. *fl. mex. ic. ined.*

Capsulæ semina et involucella Sabdariiffæ, sed caules fruticosi arborescentes.

88. *H. TRICUSPIS* (Cav. diss. 3. p. 152. t. 55. f. 2.) foliis incanis trilobis, lobis lanceolatis subdentatis, pedunculis axillaribus subbifloris, terminalibus subracemosis, involucello 9-dentato. 5 in insul. Societatis *H. hastatus* Lin. f. suppl. 310.

89. *H. AZANZÆ*, foliis glabris, inferioribus cordatis palmati-lobatis, mediis ovato-cordatis summisque ovatis integris, involucello 10-12-dentato. 5 in Mexico. *Azanza insignis* fl. mex. ic. ined. Flores magni coccinei.
90. *H. CIRCINNATUS* (Willd. enum. 735.) foliis orbiculato-cordatis acuminatis integerrimis subtus cano-pubescentibus, involucello 10-dentato. 5 ad Caracas.
91. *H. TILIACEUS* (Lin. sp. 976.) foliis subrotundo-cordatis acuminatis crenatis subtus cano-pubescentibus, involucello 10 dentato. 5 in Indiâ orient. ad ripas. Cav. diss. 3. t. 55. f. 1. Pariti Rheed. mal. 1. t. 30. Folia subtus in nervis 3 poro lineari instructa. (v. s.)
92. *H. ELATUS* (Swartz fl. ind. occ. 2. p. 1218.) foliis subrotundo-cordatis integerrimis subtus cano-pubescentibus, pedunculis brevissimis 1-floris, involucello 10-fido. 5 in campis montium Jamaicæ, Porto-Ricci, Guianæ gallicæ. Folia subtus in nervo medio poro lineari instructa. An à priore satis differt? (v. s.)
93. *H. GUINEENSIS*, foliis subrotundo-cordatis integerrimis subtus cano-pubescentibus, pedunculis 1-floris petiolo longioribus, involucello brevissimo 10-dentato 5 in Guineâ ad ripam Quoya. Bacle. Priori valde affinis, flos duplò major. (v. s.)
94. *H. ABUTILOIDES* (Willd. enum. 736.) foliis subrotundo-cordatis acuminatis crenatis utrinque viridibus glabris, stipulis cordato-ovatis. 5 in America calid. — Sloan. jam. 1. t. 234. f. 4. ex Bertol. exc. p. 13.
95. *H. PERNAMBUCENSIS* (Bertol. exc. p. 13.) foliis profundè cordatis subrotundo-acuminatis crenatis, senioribus utrinque glaberrimis, stipulis lanceolato-falcatis. 5 in Brasiliâ et Guadalupâ. Flores ignoti.

Sect. XI. LAGUNARIA.

Involucellum subnullum nempè reductum ad marginem prominentem subintegrum dentatumve, aut 1-phyllum deciduum.

96. *H. PATERSONII* (Ait. h. kew. 4. p. 224.) foliis lanceolato-oblongis integerrimis subtus squamis albicantibus lepidotis. 5 in insulâ Norfolk. *H. Patersonius* Andr. bot. rep. t. 286. *Lagunæa Patersonia* Sims bot. mag. t. 769. *Lagunæa squamea* Vent. malin. t. 42. (v. v.)
97. *H. CUNEIFORMIS*, foliis oblongis cuneatis obtusis glabris integerrimis. 5 in Novæ-Hollandiæ orâ orientali. Pedicelli longitudine calycis. Involucellum minutissimum dentatum. Calyx ultrâ medium 5-fidus lobis acuminatis. Caulis et folia ferè *Candolleæ cuneiformis*. (v. s.)

† *Species non satis notæ.*

98. *H. DIGITIFORMIS*, foliis digitatis subnovemlobatis, lobis inæqualibus angustissimis lanceolatis subpubescentibus, petiolis cinereo-tomentosis. 5 Patr. ign. *H. digitatus* Poir. suppl. 3. p. 220. non Cav.
99. *H. FLABELLATUS* (Poir. suppl. 3. p. 220.) spinulosus, foliis flabellatis 5-lobatis glaberrimis, lobis lanceolatis longissimis serratis. 5 in Novâ-Hollandiâ.
100. *H. SPICATUS* (Cav. diss. 3. p. 163. t. 59. f. 1.) foliis 5-lobis obtusis tomentosis, floribus sessilibus in spicam terminalem digestis. 5 Patr. ign. An potius *Althææ species*?
101. *H. FLAVESCENS* (Cav. diss. 3. p. 164. t. 70. f. 3.) foliis 5-lobo-palmatis inæqualiter dentatis hirsutis, caule petiolisque inermibus. — in Pondicheri.
102. *H. FASCICULATUS* (fl. mex. ic. ined.) foliis villosis palmato-5-fidis, lobis dentatis, ultimo-longiore, florum pedicellis in axillis confertis, unico ceteris longiore. — in Mexico.
103. *H. TRIPARTITUS* (Forsk. descr. p. 126.) foliis 3-partitis, lobis lanceolatis serratis, pedunculis axillaribus fructu quadruplò brevioribus, caule et calyce aculeato-hispidis, capsulis pilosis. — in Aegypto aut Arabiâ.

104. *H. ACETOSÆFOLIUS* (fl. mex. ic. ined.) hispidus, foliis profundè tripartitis, lobis oblongis acutis dentatis, 2 brevibus erectis, medio longissimo, pedicellis axillaribus 1-floris. ① in Mexico.
105. *H. FRATERNUS* (Lin. f. suppl. 311.) lævis, foliis trilobis, involucelli radiis teretibus apice mucronato-concavo, capsulâ pubescente. — in Surinamo.
106. *H. BORBONICUS* (Link. enum. 2. p. 216.) foliis cordato-subrotundis sublobatis inæqualiter et acutè dentatis subtus tomentosis, involucri foliolis 5 oblongis longè acuminatis basi attenuatis. 5 in ins. Borboniâ. Cor. ampla flava.
107. *H. TOMENTOSUS* (Mill. dict. n. 5.) foliis cordatis angulatis serratis tomentosis, caule arboreo, capsulis pilosis 5-cornutis. 5 in Indiâ occid. Flores flavi, demum subpurpurascentes.
108. *H. BRACTEOSUS* (fl. mex. ic. ined.) foliis cordatis suborbiculatis mucronatis integris, stipulis ovatis magnis, summis bracteiformibus, involucello 5-phylo. 5 in Mexico. Flores flavi. Habitus *H. tiliacei*.
109. *H. MACROPHYLLUS* (Roxb. ex. Horn. suppl. 149.) foliis subrotundo-cordatis cuspidatis utrinque tomentosis, floribus paniculatis, pedunculis petiolisque sericeo-lanatis. 2 in Indiâ orientali ad Chittagong.
110. *H. OXYPHYLLUS* (fl. mex. ic. ined.) glaber, foliis cordatis longè acuminatis dentatis, involucello 5-phylo, foliolis patentibus linearibus apice subdilatatis calyce longioribus. — in Mexici montibus Xochipici. Flores albi in medio incarnati.
111. *H. CYANOCYNUS* (fl. mex. ic. ined.) hispidus, foliis cordatis acutis dentatis, pedunculis axillaribus multifloris, summis subpaniculatis, involucello polyphylo patente. — in Mexico.
112. *H. BAHAMENSIS* (Mill. dict. 14.) foliis glabris subtus incanis oblongo-cordatis dentatis longè petiolatis, floribus subterminalibus amplissimis lætè purpureis. 2 in ins. Bahamensibus.
113. *H. PURPUREUS* (Forsk. descr. p. 126.) foliis cordato-oblongis acutis serratis, stigmatibus 5 longis apice capitatis, capsulâ globosâ multi supernè 7-alatâ. — in Arabiâ. Flor. purpurei.
114. *H. PANDURÆFORMIS* (Burm. ind. p. 151. t. 47. f. 2.) foliis cordato-lanceolatis denticulatis tomentosis, caule herbaceo hirtio, involucello 8-phylo. — in Indiâ orientali. Flores sulphurei. Sat diversus videtur ab *H. tubuloso*.
115. *H. PARVIFLORUS* (Weinm. in flora 1820. p. 610.) caule frutescente piloso-hispido, foliis cordatis angulatis crenatis subtus tomentosis, involucello 9-phylo, foliolis apice dilatatis. — in Americâ. Petala parva extus hispida sordidè flavescentia maculis 5 rubicundis. Affinis videtur *H. Senegalensi* et *tubuloso*.
116. *H. CHINENSIS*, glaber, foliis ovatis acuminatis dentatis, pedicellis axillaribus unifloris petiolo longioribus, involucello patente setaceo 6-8-phylo. — in Chinâ. — Braam. ic. chin. 1821. t. 24. — Flores albi. Habitus ferè *H. phænicei*.
117. *H. CUCURBITINUS* (Burch. cat. geogr. n. 1481. voy. 1. p. 278.) procumbens valdè tomentosus, foliis rotundatis repandis suprâ minus tomentosis, floribus confertis racemosis axillaribus. — ad cap. B.-Spei prope Dwaal river. Corolla rufa campanulata vix calyce major. Ab Hottentotis dicitur Wilde Kalebas.

XII. *THESPESIA*. *Corr. ann. mus. 9. p. 290.* — *Malvaviscus. Gaertn. fruct. 2. t. 135.*

Calyx truncatus, cinctus involucello 3-phylo deciduo. Capsula 5-locularis, loculis septo incompleto semi-partitis basi 4-spermis. Albumen parcum inter cotyledonum rugas. — Caulis arboreus. Folia integerrima.

1. *T. POPULNEA* (Corr. l. c.) foliis subrotundo-cordatis acuminatis. 5 in Indiâ orientali et ins. Societatis. *Hibiscus populneus* Lin. sp. 976. Cav. diss. 3. p. 152. t. 56. f. 1. — Rumph. amb. 2. t. 74. Pediculi 1 $\frac{1}{2}$ -2 pollicares. (v. s.)

β. *Guadalupensis*, pedicellis pollicaribus, petalis angustioribus basi evidentius ciliatis. 5 An in Guadalupâ culta? An species propria? Fruct. ign. (v. s.)

2. *T. GRANDIFLORA*, foliis ovatis subcordatis subacuminatis. 5 in Porto-Ricco. *Hibiscus grandiflorus* Juss! ined. Flores punicei 4-5 poll. diam.. Pedicelli 4-pollicares. Fructus ignotus. Habitus omnino *T. populneæ*. (v. s.)

XIII. *GOSSYPIUM*. Lin. gen. n. 845. Lam. ill. t. 586. — Xylon. Tourn. inst. t. 27.

Calyx cyathiformis, obtusè 5-dentatus, cinctus involucello 3-partito, foliolis basi coalitis cordatis dentato-incisis. Stigmata 3-5. Capsula 3-5-locularis polysperma. Semina lanam tomentosam gerentia!

N. B. Species omnes incertæ, ex characteribus mancis stabilitæ! conf. Rhor. cot. 1790. ed. gall. 1807. — Paris. mem. cot. 1810. — Lasteyrie cot. 1808. etc. Hic species à Botanicis admissas recenseam monens tamen hoc genus monographiæ accuratæ et ex vivo elaboratæ maximè egere.

1. *G. HERBACEUM* (Lin. sp. 975.) foliis 5-lobis subtus r-glandulosis, lobis rotundatis mucronatis, involucello serrato, caule lævi. ① Lin. ② Par. 4. Roxb. in Oriente. Petala flava unguibus purpureo-maculatis. Cav. diss. t. 164. f. 2. (v. v.)
2. *G. INDICUM* (Lam. dict. 2. p. 134.) foliis 3-5-lobis obtusis subtus eglandulosis, involucello apice subinciso, caule hirsuto. ① ② in Indiâ orient. — Rumph. amb. 4. t. 12. Cav. diss. 3. t. 169. Flores flavidis unguibus purpureis.
3. *G. MICRANTHUM* (Cav. diss. 6. p. 311. t. 193.) foliis 5-lobis obtusis glaberrimis subtus r-glandulosis, involucello multifido petalis longiore, caule glabro punctato. — in Persiâ ad Hispahan.
4. *G. ARBOREUM* (Lin. sp. 975.) foliis 5-lobis palmatis, lobis lanceolatis obtusis setulâ brevi mucronatis subtus r-glandulosis, involucellis subintegris. 5 in Indiâ arenosis. — Rheed. mal. 1. t. 31. Cav. diss. 6. t. 195. *G. rubrum*. Forsk. descr. n. 88? (v. s.)
5. *G. VITIFOLIUM* (Lam. dict. 2. p. 135.) foliis inferioribus 5-lobis palmatis, super. 3-lobis subtus r-glandulosis, involucellis laciniatis, calycibus basi 3-glandulosis, caule punctato glabro. ① Comm. 5 Roxb. in Indiâ orient. — Rumph. amb. 4. t. 13. Cav. diss. 6. t. 166. — *G. glabrum* Lam. ab hoc non differt ex Cav. (v. s.)
6. *G. HIRSUTUM* (Lin. sp. 975.) foliis superioribus indivisis cordatis, inferioribus 3-5-lobis subtus r-glandulosis, ramulis petiolisque hirsutis, involucellis apice 3-dentatis. ① ② Willd. 5 Roxb. in Americâ calid. Flores flavi. Pluk. t. 299 f. 1. Cav. diss. 6. t. 167. (v. s.)
7. *G. EGLANDULOSUM* (Cav. diss. 6. p. 354.) foliis 5-lobis eglandulosis, lobis tribus oblongis acuminatis, caule villosa, involucellis apice 3-4-dentatis. ① Patr. ign. Ad eadem ac *G. herbaceum*. †
8. *G. RELIGIOSUM* (Lin. sp. 975.) foliis superioribus 3-lobis, infer. 5-lobis, subtus r-glandulosis, ramulis petiolisque nigro-punctatis, involucello sub-3-phylo laciniato villosa, seminum lanâ pallidè croceâ. ② Cav. 4. Roth. 5 Willd. in Indiâ orient. Cav. diss. 6. t. 164. f. 1. An *G. tricuspidatum*. Lam. dict. 2. p. 136?
9. *G. LATIFOLIUM* (Murr. comm. goett. 1776. p. 32. t. 1.) foliis acutis, infimis indivisis, reliquis 3-lobis, subtus r-glandulosis. ② 5 Patr. ign.
10. *G. BARBADENSE* (Lin. sp. 975.) foliis super. 3-lobis, infer. 5-lobis subtus

triglandulosis, caule læviori, seminibus liberis. ② 5 Lin. 4 Roxb. in Barbados. Pluk. alm. t. 188. f. 1.

11. *G. PERUVIANUM* (Cav. diss. 6. p. 313. t. 168.) foliis 5-lobis 3-glandulosis, inferioribus indivisis, involucellis laciniatis basi 3 glandulosis. ② in Peru. Flores flavi ad unguis purpurei.
12. *G. PURPURASCENS* (Poir. suppl. 2. p. 369.) foliis 3-lobis subtus pubescentibus, lobis ovato-lanceolatis acutis, involucello laciniato, ramis apice puberulis, capsula 3-valvi. 5 in Americâ merid. †.
13. *G. RACEMOSUM* (Poir. suppl. 2. p. 370.) glaberrimum, foliis subcordatis 3-lobis acuminatis, floribus apice ramorum subracemosis, involucellis laciniatis, capsulâ 3-valvi. 5 in Porto-Ricco. †.

†. *Species inquirendæ.*

G. OBTUSIFOLIUM (Roxb. cat. calc. 51.)

G. ACUMINATUM (Roxb. cat. calc. 51.)

G. GLANDULOSUM (Rœusch. nom.)

G. Species 29 à cl. von Rohr. descriptæ et cum prioribus conferendæ.

G. Species 7 à cl. Paris descriptæ et cum prioribus conferendæ.

G. Species 18 à cl. Badier descriptæ et cum prioribus conferendæ.

XIV. REDOUTEA. Vent. cels. t. 11.

Calyx 5-partitus, cinctus involucello 10-12-phylo calyce breviorē *Stigmata* 3. *Capsula* 3-locularis, 3-valvis, oo-sperma, placentis 3 cum valvis alternantibus, utroque latere semina lanigera gerentibus.

1. *R. HETEROPHYLLA* (Vent. l. c.) foliis ciliatis ellipticis utrinquē rotundatis integris rariùs trifidis. ① in ins. St. Thomas et in ripâ fl. Orinoci. H. B. et Kunth nov. gen. 5. p. 293. Caulis erectus glaber ramosus. Folia 3 nervia, squamulis sparsis lepidota. Flores sulphurei unguibus purpureis pedicellis 3-gonis. (v. s.)
2. *R. TRIPARTITA* (H. B. et Kunth l. c.) foliis glabresculis profundē tripartitis, laciniis oblongis acutis basi cuneatis, intermediâ integrâ aut obsoletè trilobâ, lateralibus bifidis. 5 in ripâ flum. Amazonum. Procumbens. Flores flavi.

XV. *FUGOSIA*. Juss. gen. 274. — *Cienfugosia*. Cav. diss. 3. p. 174. t. 72. f. 2. Lam. ill. t. 577.

Calyx 5-fidus, cinctus involucello 12-phylo brevissimo setaceo. *Antheræ* paucæ in mediâ superficie tubi quasi verticillatæ. *Stigma* 1 clavatum. *Capsula* 3-locularis, globosa, 3 sperma.

1. *F. DIGITATA* (Pers. ench. 2. p. 240.) 5 in Senegal. Folia 3-5-partita, lobis linearibus obtusis; pedicelli 1-flori. *Cienfugosia digitata* Cav. l. c. *Cienfuegia* Willd. sp. 3. p. 723.

XVI. *SENRA*. Cav. diss. 2. p. 83. t. 35. f. 3. — *Senræa*. Willd. sp. 3. p. 695.

Calyx 5-dentatus, parvus, cinctus involucello 3-phylo, foliolis cordatis integris. *Antheræ* circiter 10 stipitatæ in apicē et superficie tubi. *Membrana* 4-5-crenata sub ovario. *Stigmata* 5. *Capsula* 2?-locularis, 10-sperma. — vix hujus loci ex cl. Kunth.

1. *S. INCANA* (Cav. l. c.) — in Arabiâ ad insul. Soccotara. Planta tomentosa 3-pollicaris. Folia cordata truncata 3-dentata. Flores axillares subsessiles.

XVII. *LOPIMIA*. Mart. in nov. act. bonn. xi. p. 96.

Involucellum calyce longiùs, 20-phyllum, foliolis setaceis conniven-

tibus. Cor. plana. Stam. columna subdeflexa. Stigmata 10. Antheræ 30-40. Capsula 5-coeca, coccis clausis mucilagine viscidulo illinitis. — Genus habitu Sidæ, affine Pavoniæ et Urenæ, fructu antè exsiccationem mucoso distinctissimum.

1. *L. MALACOPHYLLA* (Mart. l. c.) ♂ in Brasilæ prov. Bahiensis palustribus. Sida malacophylla. Link et Otto abb. gew. berl. 1. p. 67. t. 30. Frutex pubescentia molli pubescens. Folia orbicularia subcordata duplicato-dentata. Flores axillares solitarii coccinei.

Div. II. *Calyx basi nudus.*

XVIII. PALAVIA. Cav. diss. 1. p. 40. Lam. ill. t. 577.

Calyx nudus, 5-fidus. Carpella capsularia plurima monosperma in capitulum absque ordine adgregata. — Diff. à Sida ut Malope à Malvâ, et à Malope ut Sida à Malvâ.

1. *P. MALVÆFOLIA* (Cav. diss. 1. p. 40. t. 11. f. 4.) glabriuscula prostrata, pedunculis petioli circiter longitudine. ① in Limæ arenosis. Malope parviflora. Lher. stirp. 1. p. 105. t. 50. Palava declinata Mœnch. Flores purpurei parvi. (v. s.)
2. *P. MOSCHATA* (Cav. diss. 1. p. 41. t. 11. f. 5.) tomentosa erecta, pedunculis folio longioribus. ① in Limæ arenosis. Flores purpurascens. (v. s. in h. Balb. sub nomine *P. prostrata* All.)

XIX. CRISTARIA. Cav. ic. 5. p. 10. non Sonn.

Calyx nudus 5-fidus. Fructus orbiculato-depressus, pelliculâ tectus, constans carpellis plurimis ad centrum bialatis 1-spermis. — Vix à sequente differt ex cl. Kunth.

1. *C. BEYONICÆFOLIA* (Pers. ench. 2. p. 248.) erecta, foliis subcordatis inciso-crenatis incanis. — In Chili. Fenill. hist. 3. p. 40. t. 27.
2. *C. GLAUCOPHYLLA* (Cav. ic. 5. p. 11. t. 418.) prostrata, foliis lobatis incisim tomentosis glaucis. — in Chili in arenosis maritimis urbis Coquimbo, Petala carnea unguibus villosis.
3. *C. MULTIFIDA* (Cav. l. c.) prostrata, foliis multifidis glabris. ① in Peruvæ arenosis. Sida multifida. Cav. diss. 1. p. 25. t. 4. f. 2. Sida pterosperma Lher. stirp. 1. p. 119. t. 57. Flores albi axillares versùs terram secundi.

N. B. An huc revocandæ Sida crispifolia et S. heterophylla quarum fructus ignotus et quibus est affinitas summa ex habitu cum *Cr. glaucophylla*?

XX. ANODA. Cav. diss. 1. p. 38.

Calyx nudus 5-fidus, lobis acuminatis in fructu patentissimis. Capsula subtus subhemispherica, supernè depresso-plana stellariformis, multi-locularis, carpellis nempè connatis 1-locularibus 1-spermis. — An potius sectio inter Sidas.

* *Carpellis seu capsulæ loculis in mucrones subspinosos stellatim digestos desinentibus.*

1. *A. HASTATA* (Cav. diss. 1. p. 38. t. 11. f. 2.) foliis inferioribus cordatis acuminato 5-angulis subdentatis obtusis, superioribus hastatis acuminatis basi subdentatis, pedicellis solitariis axillaribus longitudine foliorum. ① in Mexico et Lima locis humidis. Sida hastata Willd. sp. 3. p. 763. Mirè foliorum forma variat, Petala alba cærulea aut subpurpurea. (v. s.)
2. *A. TRILOBA* (Cav. diss. 1. p. 39. t. 10. f. 3.) foliis omnibus crenatis, inferioribus subrotundo-cordatis obtusis subquingularibus, superioribus rotundo-hastatis acuminatis, pedicellis solitariis axillaribus folio longioribus. ① in Mexico. Calyx villosissimus. Petala purpurea, submarginata. Sida cristata Willd. sp. 3. p. 763. (v. s.)
3. *A. DILLENIANA* (Cav. diss. 1. p. 40. t. 11. f. 1.) foliis inferioribus triangulo-hastatis crenatis, superioribus ovato-lanceolatis subintegerrimis, pedicellis

- solitariis axillaribus longitudine foliorum petiolo duplò longioribus. ① in Mexico. — Dill. h. elth. 1. t. 2. *Sida Dilleniana* Willd. sp. 3. p. 764. S. cristata bot. mag. t. 330. excl. syn. Petala rosea apice emarginata. (v. s.)
4. *A. TRIANGULARIS*, foliis triangulari-subrhombeis acuminatis dentatis basi et apice integerrimis, pedicellis solitariis axillaribus longitudine foliorum petiolo quintuplò longioribus ① in Mexico Humb. et Bonpl. *Sida deltoidea* Horn. cat. hafa. 36. Brevitate petioli à priori facile differt. (v. s.)
5. *A. INCARNATA* (H. B. et K. nov. gen. am. 5. p. 255.*) ramis folisque utrinque hirtis, inferioribus ovato-oblongis cordato-hastatis serratis, superioribus angustis trifido-hastatis, pedicellis solitariis axillaribus folio sublongioribus. ②? in Horto Mexicano culta. Petala incarnata. Carpella 12. An eadem ac prior?
- ** *Fructu mutico.*
6. *A. ACERIFOLIA*, foliis inferioribus cordato-angulatis, summis hastato-clongatis, reliquis 5-lobo-hastatis, pedicellis folio longioribus. petalis calyce majoribus obovatis. ① in Mexico. Petala cærulea dimidiò quam in prioribus minora. *Sida acerifolia* Zucc. obs. n. 80. *Sida hastata* bot. mag. t. 154? S. quinqueloba fl. mex. ic. ined. (v. s.)
7. *A. PARVIFLORA* (Cav. ic. 5. p. 19. t. 431.) foliis inferioribus cordatis angulatis, superioribus hastatis, pedicellis folio brevioribus, petalis crepatis vix calyce majoribus. ① in valle Queretaro Novæ-Hispaniæ. Petala dilutè lutea. *A. crenatiflora* Ort. dec. p. 96. *Sida crenatiflora* Pers. (v. s.)

XXI. PERIPTERA.

Calyx nudus 5-fidus. Petala erecta, in tubum spiraliter contorta, demum libera. Capsula stellatim multilocularis, carpellis 1 spermis. — Genus proprium à *Sida* ut *Malvaviscus* ab *Hibisco* ferè distinctum?

1. *P. PUNICEA*, foliis inferioribus cordatis sub-5-lobo-hastatis, superioribus hastatis, petalis erectis spathulatis apice subdentatis calyce duplò longioribus. ② in Novâ-Hispaniâ, corolla punicea *Sida periptera* Sims! bot. mag. t. 1644. *Sida Malvaviscus* fl. mex. ic. ined. *Sida rubra* Ten. h. nap. *Anoda punicea* Lag. nov. gen. t. 21. (v. s.)

XXII. *SIDA*. Cav. diss. p. 5. — *Sida* et *Napæa*. Lin. Lam. ill. t. 578. et 579. — *Sida*, *Bastardia*, *Gaya* et *Abutilon*. Kunth malv. p. 4.

Calyx nudus, 5-fidus, sæpè angulatus. Stylus apice multifidus. Carpella capsularia 5-30 circa axim verticillata, plus minùsve inter se coalita, 1-locularia, mono aut oligosperma, apice mutica aut aristata. — Genus polymorphum, in posterum jam docente cl. Kunth, dividendum ex fructû seminisque structurâ valdè diversâ, sed adhuc intactum hùc retinui ob innumeras species non satis cognitâs ut in ordine carpologico locum obtineant.

Sect. I. MALVINDA. Medik. malv. p. 23.

Carpella 5-12 monosperma non vesicaria.

* *BREVI PEDICELLATÆ*, nempè *pedicellis petiolum non excedentibus, foliis oblongis ovatisve.*

1. *S. LINIFOLIA* (Cav. diss. 1. p. 14. t. 2: f. 1.) foliis linearibus integerrimis floris diametro multò longioribus, racemis terminalibus, carpellis 5-8 submuticis. ② in Peruvîâ, Cayennâ, S. Domingo. H. B. et K. nov. gen. 5. p. 260.* *Malva hirsuta* gramineo folio Aubl. guian. 2. p. 704. Flores parvi vix 4. lin. diam.
2. *S. BRACHYSTEMON* (fl. mex. ic. ined.) foliis linearibus integerrimis vix diametro floris longioribus, pedicellis axillaribus 1-floris stipularum petiolique longitudine. — in Mexico: Fructus ignoti. Flores albidî, in centro rosei, diam. ferè sesqui-pollicari. Stamina brevissima.
3. *S. ANGUSTIFOLIA* (Lam. dict. 1. p. 4.) foliis lineari-lanceolatis dentatis, taber-

- culo spinoso sub basi folii, pedicellis axillaribus subsolitariis, carpellis 5 bicuspidatis. 5 in ins. Bourboniæ. Flores lutei, parvi. *S. angustifolia* Cav. diss. 1. p. 14. t. 2. f. 2. non Mill. *S. ulmifolia* Retz obs. 3. p. 37. (v. v.)
4. *S. LINEARIS* (Cav. ic. 4. p. 6. t. 312. f. 1.) foliis linearibus serratis, tuberculo sub basi folii spinoso, pedicellis axillaribus solitariis, carpellis 10 muticis. 5 in Novâ-Hispaniâ. Flores lutei parvi, vix aperti. Priori valdè affinis.
5. *S. SPINOSA* (Lin. sp. 960.) foliis ovato-lanceolatis dentatis, tuberculo sub basi folii spinoso, pedicellis axillaribus solitariis stipulis petioloque brevioribus, capsulis 5 hirsutis. ① in Indiâ, AËgypto, Senegal, Jamaicâ. Cav. diss. 1. p. 11. t. 1. f. 9. *Stewartia corchoroides* Forsk. Flor. flavi. Duabus prioribus ex pulvino spinoso affinis. Variat foliis subcordatis. (v. s.)
6. *S. ACUTA* (Burm. ind. 147.) foliis lineari-lanceolatis dentatis glabris, pedicellis axillaribus solitariis stipularum petiolique longitudine, carpellis 5 birostratis. 5 in Coromandeliâ, Javâ, Cochinchinâ. — Pluk. t. 334. f. 2. opt. — Cav. diss. 1. p. 15. t. 2. f. 3. *S. scoparia*. Lour. coch. 2. p. 504. Flores pallidè lutei (v. s.)
7. *S. ALBA* (Lin. sp. 960 non Cav.) foliis oblougo ovatis subcordatis obtusis dentatis, pediculis petiolo æqualibus, carpellis 5 birostratis. ① in Indiâ orientali. Flores albi. — Dill h. elth. 2. t. 171. f. 210.
8. *S. STANTONIANA*, foliis oblougo-lanceolatis dentatis glabris subtus pallidis, pedicellis axillaribus solitariis stipularum petiolique longitudine, carpellis 7 birostratis. 5 in Chinâ. Stantton. *S. lanceolata* Willd. sp. 3. p. 736? *S. stipulata*, var. e Chinâ Cav. diss. 1. p. 23? (v. s.)
9. *S. STIPULATA* (Cav. diss. 1. p. 22. t. 3. f. 10.) foliis lanceolatis dentatis acutis pedicellis solitariis axillaribus, stipulis ciliatis dimidiò ferè brevioribus, carpellis 7-10 birostratis. — in ins. Mauritiî. Variat foliis pilosis et glabrioribus. (v. s.)
10. *S. MURICATA* (Cav. ic. 6. p. 78. t. 597. f. 2.) foliis lanceolatis serratis vix petiolo longioribus, stipulis ciliatis elongatis, floribus subcapitatis, carpellis 7 muricatis biaristatis. — in Novâ-Hispaniâ prope Chalma. Caulis pedalis villosus.
11. *S. BERTERIANA* (Balb! in litt.) foliis lanceolatis dentatis acutis subpilosis, pedicellis 1-2 axillaribus 1-floris brevissimis, stipulis ciliatis petiolo florequè duplò longioribus, carpellis 5 submuticis. — in Sancto-Domingo et Porto-Ricco in pratis. Valdè accedit ad *S. stipulatam*. (v. s.)
12. *S. GLOMERATA* (Cav. diss. 1. p. 18. t. 2. f. 6.) foliis ovato-lanceolatis serratis tomentosis, pedicellis axillaribus 1-floris brevissimis, carpellis 5 bicornibus. — Patr. ign. Calyces ciliati. Aff. *S. Jamaicensi*.
13. *S. JAMAICENSIS* (Cav. diss. 1. p. 17. t. 2. f. 5.) foliis ovatis serratis tomentosis, serraturis obtusis subaristatis, pedicellis axillaribus 1-floris brevissimis, carpellis 5 bicornibus. 5 in aridis Jamaicæ et Sancti-Domingi. (v. s.)
14. *S. BALBISIANA*, foliis ovato-lanceolatis subrhombicis acuminatis basi integris dentato-serratis ramulisque pube substellatâ minimâ scabridis, pedicellis solitariis 1-floris stipularum ciliatarum longitudine, carpellis 8-10 birostratis. 5 in Porto-Ricco. Bertero. (v. s. in h. Balbis.)
15. *S. BRACHYPETALA*, foliis ovato-lanceolatis acuminatis inæqualiter serratis pube simplici adpressâ subpubescentibus, stipulis calycibusque ciliatis, pedicellis solitariis 1-floris stipularum longitudine, corollâ calyce breviorè, carpellis 8-10 birostratis. 5 in Martinicæ arvis dormientibus et in S.-Domingo. Differt à *S. frutescente* pedicellis basi nec apice articulatis. (v. s.)
16. *S. REPANDA* (Roth. nov. sp. 328.) foliis ovatis acuminatis duplicato-serratis suprâ pilis stellatis minimis adspersis, pedicellis solitariis stipulis linearilanceolatis ciliatis brevioribus, petalis obliquè repandis, carpellis hicuspidatis. 5 Patr. ign. Corollæ sulphuræ. Affinis ex cl. Roth *S. carpinifolia*.
17. *S. BRACTEOLATA*, foliis ovato-lanceolatis acuminatis dentatis glabris, ramis teretibus puberulis, stipulis cujusque folii unâ inferiore subulatâ, alterâ linearari, racemulis brevissimis 1-3-floris bracteolatis, carpellis 7-8 birostratis. 5 in Americâ merid. in itinere à Chili ad Brasiliam semina collegit P. Schmidt-meyer. Flores flavi parvi. (v. v. in hort. Genevensi.)

18. *S. CARPINIFOLIA* (Lin. f. suppl. 307.) foliis ovato-oblongis duplicato-serratis, pedunculis axillaribus brevissimis sub-4 floris, ramulis complanatis, carpellis 8 birostratis. ♀ in Maderâ et ins. Canariis. *S. carpinifolia*. Cav. diss. 1. p. 21, 5 p. 274 (excl. descr. fruct.) t. 134. f. 1. et *S. planicaulis* Cav. diss. 1 p. 24. t. 3. p. 11. (pl. jun.) (v. s.)
 ♂? *betulina* (Lag. h. madr. 1815.) foliis latioribus basi subcordatis. — in insulis Mauritanis. — *S. carpinifolia* Jacq. ic. rar. 1. t. 135. (v. s.)
19. *S. CARPINOIDES*, foliis ovato-oblongis subduplicato-serratis, pedicellis axillaribus 1-floris solitariis, carpellis 10-11 dorso breviter bicornatis et basi internâ 1-aristatis. ♀. Patr. ign. In hortis sæpè sub nomine *S. carpinifoliz* occurrit. (v. s.)
20. *S. MUCRONULATA*, foliis infer. ovato-oblongis, super. oblongis subduplicato-dentatis, pedicellis axillaribus brevissimis 1-floris, carpellis 10 11 apice bimucronulatis basi breviter 1-aristatis. ♀ in Javâ. Priori valdè affinis, differt foliis magis oblongis glabris, fructibus magis hispidis minoribus brevius mucronatis. (v. s.)
21. *S. ORIENTALIS* (Cav. diss. 1. p. 21. t. 12. f. 1.) foliis ovatis acuminatis dentatis glabris, pedicellis axillaribus 1-floris petiolo subbrevioribus, carpellis 9-10 muticis. — in Indiâ orientali.
22. *S. CAPENSIS* (Cav. diss. 1. p. 23. t. 12. f. 3. et 2 p. 49.) foliis ovato lanceolatis ovatisve dentatis, pedicellis axillaribus solitariis petioli longitudine, stipulis ciliatis petiolo longioribus, carpellis 10 muticis. — Ad cap. Bonæ-Spei? — Pluk. alm. t. 240. f. 5.
23. *S. MICROPHYLLA* (Cav. diss. 1. p. 22. t. 12. f. 2.) foliis ellipticis dentatis pedicellis axillaribus solitariis petiolo sublongioribus, carpellis 7 birostratis ♀ in Indiâ orientali.
24. *S. TRIDENTATA* (Cav. ic. 4. t. 312.) foliis ovatis apice tridentatis subtus tomentosis, pedicellis axillaribus 1-floris solitariis petiolo stipulisque æqualibus, caule filiformi brevissimo. — in Sancto-Domingo.
25. *S. EROSA* (Link. enum. h. ber. 5. p. 203.) foliis rhombeis basi attenuatis anticè serrato-dentatis subtus stellulato-subtomentosis, pedicellis petiolo brevioribus, carpellis birostratis. ♀ in Brasiliâ. Flores flavi.
26. *S. CILIARIS* (Lin. sp. 961.) foliis ellipticis subovatis retusis apice dentatis, pedicellis axillaribus solitariis brevissimis, stipulis ciliatis flore sublongioribus, carpellis 7 muricatis breviter birostratis. — in Jamaicâ, Sancto-Domingo, S. Thomas, locis conculcatis. Cav. diss. 1. p. 21. t. 3. f. 9. et 5. t. 127. f. 2. — Sloan. hist. t. 137. f. 2. (v. s.)
27. *S. FUSILLA* (Cav. diss. 1. p. 6. t. 1. f. 4. et 5. t. 127. f. 1.) foliis subrotundo-ellipticis dentatis glabris, pedicellis axillaribus 1-floris petiolo sublongioribus, carpellis 5 muticis, caule prostrato. — in insulâ Mahe. Habitu accedit ad *S. ciliarem*.
28. *S. ALNIFOLIA* (Lin. sp. 961.) foliis infer. subrotundo-ovatis, super. oblongis dentatis basi integerrimis subcuneatis, pedicellis axillaribus pluribus petiolo brevioribus, carpellis 5-7 bicornibus. (†) in Indiâ orientali. — Dill. h. elth. t. 172. f. 211. Cav. diss. 1. p. 12. t. 1. f. 13P
29. *S. PARVIFOLIA*, foliis ovatis dentatis subtus canescentibus, pedicellis axillaribus 1-floris petiolo multò longioribus, carpellis 5-rugosis muticis. ♀ in insulâ Bourboniâ. Bory. (v. s.)
30. *S. HERMANNIOIDES* (H. B. et K. nov. gen. am. 5. p. 258.*) foliis ellipticis utrinquè rotundatis breviter petiolatis serrato-crenatis suprâ adpresso-pilosis subtus cano-tomentosis, pedicellis axillaribus solitariis 1-floris petiolo brevioribus, carpellis 5 bicuspidatis. ♀ in siccis inter Hondam et flum. Guali. Humifusa, ramis pilosis. Flores magni. Geranii molliis. Affinis *S. rotundifoliz*.

** *OBLONGIFOLIZ*, nempe *pedicellis elongatis distinctius articulatis, foliis oblongis ovatisve*.

31. *S. FRUTESCENS* (Cav. diss. 1. p. 12. t. 10. f. 1.) foliis ovato-oblongis subcordatis serratis subtus albicantibus, pediculis axillaribus 1-floris petiolo

- duplò longioribus sub apice articulatis, carpellis 5 birostratis. 5 in insulâ Mauritiî. An S. spinosæ var? (v. s.)
32. S. COMPRESSA, foliis ovato-lanceolatis acuminatis subtus canis, ramulis compressis pube stellatâ punctatis, pedicellis axillaribus 1-floris petiolo triplò longioribus, carpellis 5 pubescentibus submucronatis. — in Napauliâ? (v. s.)
33. S. BICOLOR (Cav. ic. 4. p. 6. t. 311. foliis lanceolatis valde acuminatis serratis, pedicellis axillaribus 1-floris petiolo multò longioribus, carpellis 5 muticis. 5 in Novâ Hispaniâ. Caulis 2-pedalis viridi-violaceus fruticosus erectus, ramis patulis; petala flava extus rubentia apice obliquè truncata. — S. obliqua fl. mex. ic. ined. differt tantum foliis minùs acuminatis.
34. S. CANARIENSIS (Willd. sp. 3 p. 755. foliis lanceolatis dentatis glabris, pedicellis axillaribus 1-floris folii longitudine, carpellis 7-10 birostratis. 5 in Canarias ubi ut Thea usurpatur (Cav.), in Indiâ orientali undè forsàn in Canariis migravit. S. alba Cav. diss. 1. p. 22. t. 3. f. 8. non Lin. — Specimina mea habent folia subtus caudicantia non omninò glabra. Ad sequentem nimis accedit. (v. s.)
35. S. RHOMBIFOLIA (Lin. sp. 961.) foliis oblongo-lanceolatis dentatis basi cuneatis subtus candicantibus, pedicellis axillaribus 1-floris folio brevioribus, carpellis 8-10 birostratis. 5 in Carolinâ, Brasiliâ, Cumanâ, ins. Caribæis et forsàn in Indiâ orientali? Cav. diss. 1. t. 3. f. 12. H. B. et Kunth nov. gen. am. 5. p. 261. * (v. s.)
- ρ. canescens* (Cav. diss. 1. t. 8. f. 3.) foliis subtus magis tomentosus, pedicellis folio longioribus. 5 in Senegal. Vereor ne in S. rhomboidali plures adhuc sint confusæ species. (v. s.)
36. S. HONDENSIS (H. B. et Kunth nov. gen. am. 5. p. 260. *) foliis lanceolato-oblongis subrhombeis acutis cordatis serratis subtus tenuiter cano-tomentosis, pedicellis axillaribus 1-floris folio brevioribus, carpellis 8 rugulosis post dehiscentiâ bidentatis. 5 prope Honda Novo-Granatensium. S. rhombifolia simillima.
37. S. RHOMBOÏDEA (Roxb. ex journ. bot. 1814. v. 4. p. 207) foliis lanceolatis dentatis subtus canescentibus, pedicellis axillaribus 1-floris folio brevioribus, carpellis 8-10 submuticis. 5 in Timor et Bengalo. Specimina huic valde similia sed forsàn diversa habeo ex cap. Bonæ-Spei. (v. s.)
38. S. RETUSA (Lin. sp. 961.) foliis obovatis apice dentatis retusis subtus subcanescentibus, pedicellis 1-floris, carpellis 7-9 rostratis. 5 in Indiâ orientali, ins. Mauritiî. Variat pedicellis folio brevioribus longioribusve. Carpella rostro distinctissimo nec mutica ut in S. Philippicâ. Cav. diss. 1. p. 18. t. 3. f. 4. et 5. t. 131. f. 2. (v. s.)
39. S. PHILIPPICA, foliis obovatis apice dentatis obtusis basi cuneatis integris glabrisculis, pedicellis axillaribus 1-floris folio subbrevioribus, carpellis 8-10 submuticis. 5 in ins. Philippicis. S. retusa ex ins. Philippicis Cav? (v. s.)
40. S. RECISA (Link. enum. h. ber. 2. p. 203.) foliis subrhombeis retusis antice crenatis subtus ramulisque stellulato-subtomentosis, pedicellis petiolo longioribus folio triplò fere brevioribus, carpellis muticis. 4 in Brasiliâ.
41. S. CALYXYMENIA (Gay! ined.) foliis lanceolatis obtusis dentatis utrinque cano-tomentosis, pedicellis axillaribus 1-floris petiolo vix duplò longioribus, calyce tomentoso demum membranaceo aceto. 5 in Novâ-Hollandiâ. Fructus mihi ignotus An hujus generis? (v. s.)
42. S. MICANS (Cav. diss. 1. p. 19. t. 3. f. 1.) foliis ovatis obtusis serratis tomentosus nitidis, pedicellis axillaribus solitariis petiolo multò longioribus, carpellis 9-10 breviter bicornibus. — in Indiâ orientali. A S. althæifoliâ vix differt ex cl. Kunth. nov. gen. am. 5. p. 262.
43. S. MACULATA (Cav. diss. 1. p. 20. t. 3. f. 7.) foliis ovatis obtusis dentatis ramulis calycibus stipulisque tomentosus, pedicellis axillaribus 1-floris petiolo duplò longioribus apice subracemosus, carpellis 9-10 bicornibus. 5 in Sancto-Domingo. Petala flava basi rubro maculata. S. suberosa L'her! stirp. p. 113. t. 54. ex ejus specim. et descr. ined. (v. s.)
44. S. ACUMINATA, foliis basi ovatis acuminatis dentatis ramis stipulis calycibusque tomentosus, ramulis axillaribus floriferis, pedicellis 1-floris petiolo

duplò longioribus. — in Sancto Domingo. *S. maculata* Bertero! ined. Priori affinis sed distincta, carpella mihi ignota. (v. s.)

*** *CORDIFOLIÆ, pedicellis elongatis, foliis basi cordatis dentatis non lobatis.*

45. *S. PILOSA* (Cav. diss. 1. p. 9. t. 8.) foliis ovato-cordatis obtusis dentatis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 5 birostratis. (1) in insul. Caribæis. Tota pilis hirta. Flores parvi.

♂. *glabra*, tota glabra (Cav. l. c.) in ins. Porto-Ricco. (v. s.)

46. *S. BETONICÆFOLIA* (Balb! in litt.) tota hispidissima, foliis ovatis cordatis subacutis crenatis, pedicellis solitariis 1-floris, infimis petiolo longioribus, summis brevioribus, carpellis 5 reticulato-nervosis pubescentibus birostratis. — in *S. Marthâ*. Flores albi. *S. hispida* Bertero! non Pursh. (v. s.)

47. *S. MULTICAULIS* (Cav. diss. 1. p. 10. t. 1. f. 6.) foliis subrotundo-cordatis acutis dentatis, pedicellis solitariis 1-floris petiolo duplò longioribus, carpellis 5 muticis. (1) in Malabariâ. Tota pilis longis hispida. (v. s.)

48. *S. RADICANS* (Cav. diss. 1. p. 8.) foliis subrotundo-cordatis acutis ciliato-dentatis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 5 muticis. — in Indiâ orientali. — Rheed. mal. 10. t. 69. Caulis radicans pilis albicans.

49. *S. HEDERÆFOLIA* (Cav. diss. 1. p. 8. t. 9. f. 3.) foliis subrotundo-cordatis obtusis repandis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 5 biaristatis, caule repente. — in Sancto-Domingo. — Plum. ed Burm. t. 69. f. 3. Habitus *Glechomæ* nec *Hederæ*.

50. *S. JUSSIEANA*, foliis subrotundo-cordatis dentatis utrinquè hispidis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 5 muticis, caule filiformi prostrato. (1) in Peru. Jos. de Jussieu. *S. repens* var. Cav. diss. 1. p. 7. t. 1. f. 2.

51. *S. DOMBRYANA*, foliis ovato-cordatis dentatis utrinquè pilosis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 5 birostratis, caule filiformi prostrato. (1) circa Lima. (v. s.)

52. *S. HUMILIS* (Willd. sp. 3. p. 744.) foliis subrotundo-cordatis suprâ pilosis serratis, pedicellis subsolitariis petiolo longioribus, carpellis 5. (1) in Indiâ orientali. *S. unilocularis* L'her! stirp. 1. p. 127. t. 53. bis. *S. pilosa* Retz non Cav.

α. Carpellis muticis. *S. humilis* Cav. diss. 5. t. 134. f. 2. (v. s.)

β. Carpellis submucronatis. *S. morifolia* Cav. diss. 1. p. 9. t. 1. f. 1.

γ. Carpellis biaristatis. *S. veronicæfolia* Lam. dict. n. 11. Cav. diss. 1. p. 7. t. 1. f. 3. et 5. t. 127. f. 3. An species propria?

53. *S. SUPINA* (L'her! stirp. 1. p. 109. t. 52.) foliis subrotundo-cordatis obtusiusculis crenatis molliter subvelutinis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 5 breviter bicornibus, caule procumbente. (1) in campis siccis Hispaniolæ. *S. ovata*. Cav. diss. 6. t. 196. f. 2. *S. procumbens*. Sw. fl. 2. p. 1211. (v. s.)

54. *S. DIFFUSA* (H. B. et K. nov. gen. am. 5. p. 257.*) foliis subellipticis apice rotundatis leviter cordatis longè petiolatis crenato-serratis pilosis subtus canescentibus, pedicellis axillaribus solitariis folia subæquantibus, carpellis 5 muticis. (1) prope Zelaya Mexicanorum. Caules procumbentes ramosi filiformes molliter pilosi. Folia 4 lin. longa. Flores parvi violacei.

55. *S. FLAVESCENS* (Willd. sp. 3. p. 755.) foliis ovato-cordatis obtusis inæqualiter dentatis tomentosis, pedicellis geminis 1-floris petiolo brevioribus, carpellis 5 muticis. (1) in rupibus Monte-Video. *S. flavescens* Cav. diss. 1. t. 13. f. 2. et *S. prostrata*. Cav. diss. 1. t. 13. f. 3.

56. *S. CALYCINA* (Cav. diss. 1. p. 9. t. 8. f. 2.) foliis subrotundo-cordatis acuminatis repando-dentatis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 5 muticis pyriformibus. — in ins. Borboniâ. Calyx magnus 5 partitus. (v. s.)

57. *S. HERBACEA* (Cav. diss. 1. p. 19. t. 13. f. 1.) foliis oblongis basi cordatis

- acutis dentatis pilosis, pedicellis solitariis 1-floris petiolo brevioribus, carpellis 9-10 biaristatis. — in Indiâ orientali.
58. *S. ROTUNDIFOLIA* (Cav. diss. 1. p. 20. t. 3. f. 6. et 6. t. 194. f. 2.) foliis subrotundo-ovatis cordatis dentatis tomentosis, pedicellis solitariis 1-floris petiolo multò longioribus carpellis 9-10 longè biaristatis. — in ins. Borboniâ. Aristæ pilosæ in fasciculum exsertæ.
59. *S. MACRODON*, hirsuta, foliis subrotundis cordatis grossè crenatis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 10 muticis. — in Brasiliâ. (v. s.)
60. *S. CORDIFOLIA* (Lin. sp. 961.) foliis ovatis cordatis dentatis subangulatis obtusiusculis tomentosis, pedicellis solitariis 1-floris petiolo paulò brevioribus, carpellis 9-10 birostratis. (I) in Indiâ orientali et Africâ. Cav. diss. 1. p. 19. t. 3. f. 2. — Dill. h. elth. t. 171. f. 209. (v. s.)
61. *S. BIVALVIS* (Cav. diss. 1. p. 13. t. 11. f. 2.) foliis ovato-cordatis acuminatis crenatis tomentosis, pedicellis solitariis 1-floris petiolo brevioribus, calycibus aristato-acuminatis, carpellis 5 apice bidentatis inter se cohærentibus et ideo bivalvibus. — in Sancto-Domingo.
62. *S. PELLITA* (H. B. et K. nov. gen. am. 5. p. 263.) foliis cordatis ovato-ellipticis obtusiusculis serrato-crenatis, suprâ adpresso-pubescentibus, subtus ramisque molliter villosis tomentosis, floribus brevissimè pedunculatis in ramulis spicato-conglomeratis, carpellis 10-11 birostratis. ♀ ad Orinocum prope Augustura.
63. *S. MULTIFLORA* (Cav. diss. 1. p. 18. t. 3. f. 3.) foliis ovato-cordatis obtusiusculis dentatis tomentosis, pedicellis solitariis 1-floris petiolo longioribus, carpellis 9 longè birostratis. — in Brasiliâ.
64. *S. EMARGINATA* (Willd. sp. 3. p. 757.) foliis ovato-lanceolatis cordatis dentatis apice truncatis retusis, pedicellis solitariis 1-floris petiolo duplò brevioribus, carpellis 5 birostratis. ♀ in Hispaniolâ. *S. truncata* L'her! stirp. 1. t. 51. non Cav. (v. s.)
65. *S. ARGUTA* (Swartz. fl. ind. occ. 2. p. 1205.) foliis cordatis serratis apice attenuatis, in petiolo margine et nervis subtus puberulis, ceterum glabris, pedicellis solitariis 1-floris petiolo longioribus, carpellis 5 biaristatis. ♀ in sepius Jamaicæ et Guadalupæ. *S. glabra* Mill. dict. n. 14. (v. s.)
66. *S. ULMIFOLIA* (Cav. diss. 1. p. 15. t. 2. f. 4. non Retz.) foliis ovato-cordatis acuminatis crenatis pubescentibus, pedicellis solitariis 1-floris petiolo subæqualibus, carpellis 5 longè birostratis. ♀ in Sancto-Domingo. Rostra retrorsum scabra. (v. s.)
67. *S. BO-BONICA* (Cav. diss. 1. p. 9. t. 10. f. 2.) foliis subrotundo-cordatis acutis dentatis, pedicellis solitariis 1-floris petiolo paulò longioribus, carpellis 5 longè biaristatis. ♀ in ins. Borboniâ. Caulis hirtus.
68. *S. FRAGRANS* (L'her! stirp. 2. p. 111. t. 53.) foliis subrotundo-cordatis acuminatis crenatis hirsuto-viscosis, pedicellis solitariis 1-floris petiolo brevioribus, carpellis 5 subaristatis. ♀ in Hispaniolâ. (v. s.)
69. *S. GLUTINOSA* (Cav. diss. 1. p. 16. t. 2. f. 8.) foliis subrotundo-cordatis acuminatis dentatis, pediculis geminatis subbifloris petiolo longioribus, carpellis 5 longè biaristatis. ♀ in ins. Mauritiî. *S. racemosa* Burm.
70. *S. INTERRUPTA* (Balb! in litt.) foliis cordiformibus acuminatis dentatis, adultis glabris, junioribus calycibusque cano-velutinis, stipulis setaceis deciduis, floribus in spicam terminalem interruptam superne aphyllam dispositis fasciculatis, carpellis 8-10 unirostratis. ♀ Sanctæ-Marthæ. Bertero. (v. s.)
71. *S. ALTHÆA-FOLIA* (Swartz prod. 101. fl. 2. p. 1207.) foliis cordatis subangulatis obtusis serrato-crenatis utrinquè tomentosis, pediculis petiolo brevioribus 1-5-floris, carpellis 10-12 breviter birostratis. ♀ in campis siccis Jamaicæ anstralis et Novæ-Granadæ. L'her! stirp. 1. p. 112. — Sloan. hist. 1. t. 136. f. 2. Flores fulvi. (v. s.)

ß. *aristosa*, aristis carpello longioribus retrorsum scabris. ♀ in Guianâ Gallicâ Perrotet. Specimina Guadalupeña inter duas variet. media!

72. *S. URENS* (Lin. sp. 963.) foliis ovato-cordatis acuminatis dentatis, pedunculis 3 4-floris brevissimis, carpellis 5 muticis, caule petiolisque hispidis. — in Jamaicæ et Guadalupe fruticetis aridis. Cav. diss. 1. p. 15. t. 2. f. 7. An planta Senegalensis à Cavan. citata diversa? (v. s.)
73. *S. VERTICILLATA* (Cav. diss. 1. p. 13. t. 1. f. 12.) foliis ovato-cordatis acuminatis dentatis ramisque subpilosis, floribus axillaribus plurimis subsessilibus, carpellis 5 submuticis. — in Brasiliâ ad Rio-Janeiro. (v. s.)
74. *S. PYRAMIDATA* (Cav. diss. 2. p. 11. t. 1. f. 10 et 6. t. 194. f. 1.) foliis subrotundo-cordatis acuminatis serratis glabris, pedunculis multifloris, carpellis 5 velutinis breviter bimucronatis. ♀ in fruticetis maritimis Sancti-Domingi et Porto-Ricci. (v. s.)
75. *S. DUMOSA* (Swartz fl. ind. occ. 2. p. 1209.) foliis cordatis ovatis acuminatis serratis utrinquè glabris, pedunculis multifloris, carpellis 5-6 subrotundis pube stellatâ minutissimâ scabridis. ♀ in fruticetis Jamaicæ et Sancti-Domingi. (v. s.)
76. *S. FLORIBUNDA* (H. B. et K. nov. gen. am. 5. p. 258.) foliis cordatis ovato-oblongis acuminatis serratis utrinquè molliter villosopilosis, subius fuscescenti-canescens, racemis axillaribus, carpellis 5 muticis. ♀ in aridis regni Novo-Granatensis. Ramosissima. Ramuli hirsuto-tomentosi ferruginei. Pedicelli capillares. Flores violacei. Sequenti valde affinis.
77. *S. PANICULATA* (Lin. sp. 962.) foliis ovato-cordatis dentatis acuminatis pube stellatâ puberulis, pedunculis laxè paniculatis capillaribus, calycibus enerviis, carpellis 5 breviter birostratis. — in rupibus calcareis Jamaicæ, Peruvia, Brasiliæ. *S. paniculata* et *S. capillaris* Cav. diss. 1. p. 16 et 10. t. 12. f. 5. et 1. f. 7. *S. atro-sanguinea* Jacq. ic. rar. 1. t. 136. Corolla atro-purpurea. (v. s.)
78. *S. NERVOSA*, foliis cordatis acuminatis dentatis pube stellatâ puberulis, pedunculis paniculatis, calycibus nervosis, carpellis 5 birostratis. — in Sancto-Domingo. Bertero. Habui etiam ex horto Calcutensi specimen cultum (v. s.)
- **** *PALMATILOBATÆ*, foliis nempè palmatim in lobos 3 5-7-9 fissis.
79. *S. JAVENSIS* (Cav. diss. 1. p. 10. t. 1. f. 5.) foliis subrotundo-cordatis dentatis tricuspidatis, pedicellis solitariis 1-floris petiolo longioribus, capsulis 5 bicuspidatis, caule reclinato. — in Javâ.
80. *S. OXYPHYLLA* (fl. mex. ic. ined.) foliis cordatis dentatis villosis 3-lobis, lobis lateralibus brevibus bifidis, medio longiore acuminato, pedicellis solitariis 1-floris, petioli longitudine, carpellis 5 birostratis. — in Mexico.
81. *S. PUNGENS* (H. B. et K. nov. gen. am. 5. p. 263.) foliis cordato-subovatis obsolete trilobis grossè serrato-crenatis mollibus suprâ pubescentibus subtus ramisque villosotomentosis, floribus axillaribus solitariis in apice ramulorum fasciculato-glomeratis, carpellis 12 biaristatis. ♀ ad Ormocum prope Angostura. Florès flavi.
82. *S. HETEROPHYLLA* (Cav. ic. 5. p. 12. t. 421.) foliis radicalibus ovatis sinuatis, caulinis 3-partitis, lobis incisim dentatis, intermedio longiore, pedicellis solitariis 1-floris folio longioribus. 4 (I) in montibus Chilensibus. Fructus ignotus. An *Cristaria* species?
83. *S. CRISPIFOLIA* (Cav. ic. 5. p. 11. t. 419.) foliis cordatis inciso-lobatis infernè tomentosis supernè punctato-stellatis margine crispia, pedicellis solitariis petiolo longioribus, calycibus sinuato-undulatis. ♀ in Americâ merid. ad portum Deseado. An *Cristaria* species?
84. *S. COCCINEA*, foliis trifidis canescenti-tomentosis, floribus densè racemosis, carpellis 10 muticis. — in campis, Missouriensibus vulgata. Flores coccinei. *Cristaria coccinea*. Pursh. fl. 2. p. 453, sed carpella diversa ex Nutt. *Malva coccinea*. Nutt. gen. am. 2. p. 81, sed involucellum nullum.

85. *S. TRILOBA* (Cav. diss. 1. p. 11. t. 1. f. 11 et 5. t. 131. f. 1.) foliis cordatis dentatis trilobis, lobo intermedio acuto longiore, pedicellis solitariis folio subæqualibus, carpellis 7 muticis. 5 ad cap. Bonæ-Spei. Jacq. Schœnbr. 2. t. 142. Folia inf. indivisa. Cor. alba. (v. v.)
86. *S. RIGINOIDES* (L'Her! stirp. 1. p. 115. t. 55) foliis subpeltatis 5 lobatis, lobis ovatis acutis dentatis indivisis, pedunculis sub 1-floris, carpellis 8 biseriatis. ① in Peru. *S. palmata* Cav. diss. 1. p. 20. t. 3. f. 3. *Malviada palmata* Mœnch. (v. s.)
87. *S. JATROPHOIDES* (L'Her. stirp. 1. p. 117. t. 56.) foliis subpeltatis 7-lobatis, lobis lanceolatis acuminatis pinnatifidis dentatis, pedunculis multifloris, carpellis 7-8 muticis. ① in Peruviae arenosis. *S. palmata* Jacq. ic. rar. 3. t. 547. Cav. diss. 5. t. 131. f. 3. Cor. violacea.
88. *S. NAPA* (Cav. diss. 5. p. 277 t. 132. f. 1.) foliis palmatis 5-lobatis glabris, lobis oblongis acuminatis dentatis, pedunculis multifloris carpellis 10 muticis acuminatis. 2 in Virginia. *Napæa lævis*. Lin. syst. 750. Lam. ill. t. 579. f. 1. Flores albi. (v. v.)
89. *S. DIOICA* (Cav. diss. 5. p. 278. t. 132. f. 2.) foliis palmatis 7 lobatis scabris, lobis lanceolatis inciso-dentatis, pedunculis multifloris bracteatis subcorymbosis, floribus dioicis, carpellis 10 muticis 2 in Virginia. *Napæa scabra*. Lin. syst. 750. *Napæa dioica*. Lin. sp. 965 (v. v.)

***** PINNATILORATÆ, foliis nempè pinnatifidis aut pinnati-partitis.

90. *S. PINNATA* (Cav. ic. 5. p. 13. t. 422. f. 1.) foliis pinnatipartitis imbricatis infernè tomentosis, lobis integerrimis, floribus axillaribus sessilibus solitariis. 2 ad radices montis Chimborazo. Subacaulis, radice crassâ.
91. *S. ACAULIS* (Cav. ic. 5. t. 422. f. 2.) foliis pinnatipartitis, lobis trifidis ovatis tomentosis, floribus axillaribus sessilibus solitariis, carpellis 8-10 muticis. 2 hab. cum priore cui valde affinis.
92. *S. PICHINCHENSIS* (Humb. et Bonpl. pl. equin. 2. t. 116.) cespitosa, foliis pinnatifidis suprâ incano tomentosis subtus glabris, laciniis lineari-lanceolatis tripartitis, infimâ indivisâ, caulibus unifloris medio monophyllis, carpellis 12 hirsutis 2 in summo monte iguivomo Pichincha Quitensium alt. 2356 hexap. H. B. et K. nov. gen. am. 5. p. 255.

Sect. II. ABUTILOIDES. — *Gaya* et *Bastardia Kunth. malv. p. 4.*

Carpella 15-40 monosperma vesicaria.

93. *S. OCCIDENTALIS* (Lin. sp. 964.) foliis oblongo-cordatis dentatis sublobatis, pedicellis solitariis petiolo brevioribus, carpellis 27-30 obtusis in fructum globosum digestis. ① in Americâ. — Dill. h. elth. 1. t. 6. Cav. diss. 1. p. 24. t. 4. f. 3. *Abutilon deflexum* Mœnch. Pedicellâ fructiferi deflexi. (v. s.)
94. *S. SYLVATICA* (Cav. diss. 2. p. 56. et 5. p. 276. t. 133. f. 2.) foliis ovato-cordatis acuminatis crenatis, pedicellis geminatis petiolo multò longioribus, carpellis 30-36 muticis in globum umbilicatum collectis. 5 in sylvis Peruviae prope flumen Maragnon.
95. *S. SPICATA* (Cav. diss. 1. p. 24. t. 8. f. 1.) foliis ovatis cordatis acutis dentatis, racemo terminali nudo, carpellis 30 muticis in globum umbilicatum digestis. — in Sancto-Domingo. — Plum. ed. Burm. 1. t. 2.
96. *S. GAYA*, foliis subellipticis acutis oblique cordatis argutè serratis ramisque albidto-tomentosis, pedicellis solitariis petiolo triplò longioribus, carpellis 15. 5 in decl. occid. montium Mexicanorum prope Sopilote. *Gaya hermannioides* H. B. et K. nov. gen. 5. p. 268. t. 475. Flores flavi magn. *Malvæ rotundifoliz.*
97. *S. CANDICANS*, foliis ovato-ellipticis obtusis cordatis crenato-serratis ramisque canescenti-tomentosis, pedicellis solitariis petiolo duplò triplòve longioribus, carpellis 14-15. 5 in regno Qnitensi ad ripam fl. Guallabamba. *Gaya canescens* H. B. et K. l. c. p. 269. Flores flavi.
98. *S. SUBTRILOBA* (H. B. et K. nov. gen. 5. p. 270. t. 476. sub *Gaya*.) foliis

longè petiolatis ovatis acuminatis obsolete trilobis profundè cordatis tenuissimè tomentosis subtile canescentibus, pedicellis solitariis, fructiferis petiolo ferè longitudine. 54ⁱⁿ regno Novo-Granatensi. S. occidentali affinis.

99. S. DISTICHA (Cav. ic. 5. p. 12. t. 43a.) foliis cordatis ovatisve acuminatis-simis dentatis utrinquè cano-velutinis, pedicellis solitariis 1-floris petiolo quadruplò longioribus, carpellis 10-15 muticis. 5 in Novâ-Hispaniâ. — Appendix in quoque carpello è basi orta valvæ more semen tegit. Petala flava oblongo-obovata calyce acuminato duplò longiora. (v. s.)
100. S. NUTANS (L'Her! stirp. 1. p. 119. t. 57.) foliis oblongo-cordatis acutis dentatis molliter velutinis, pedicellis subgeminatis 1-floris petiolo vix duplò longioribus, carpellis 10 obtusis depressis. 5 in Peruvix arenosis. S. calyptrata Cav. diss. 2. p. 57. et 5. t. 133. f. 1. Structura carpellorum ut in S. distichâ. Petala paulò calyce majora. (v. s.)
101. S. VISCOsa (Lin. sp. 963.) foliis ovato-cordatis acuminatis tenuissimè serratis tomentoso-viscosis, petiolis pedicellisque pilosis, stipulis subpaulis setaceis, pedicellis petiolo longioribus solitariis, carpellis 7 muticis. 5 in Jamaicâ. S. Thomas, Porto-Ricco. Sloan. hist. 1. t. 139. f. 4. Bastardia viscosa H. B. et Kunth nov. gen. am. 5. p. 256. (v. s.)
102. S. BASTARDIA. foliis cordatis subrotundo-ovatis obtusè acuminatis subcrenatis utrinquè pube tenui stellulatâ tomentosis, petiolis pedicellisque subviscido-tomentosis longitudine subæqualibus, carpellis 5 muticis. 5 in ripâ flum. Amazonum Bastardia parvifolia. H. B. et K. nov. gen. am. 5. p. 255. t. 472. Flores flavi. In hac et in priore capsula loculicido-dehiscens.
103. S. RETROFRACTA, foliis cordiformibus acuminatis dentatis petiolisque velutino-tomentosis, stipulis setaceis deflexis, pedicellis petioli longitudine ad articulationem sæpè infractis, carpellis 7 pubescentibus muticis. 5 in Martinicâ Calyx 5-fidus lobis acuminatis patentibus aut reflexis ut in ic. H. B. et K. t. 476. (v. s.)
104. S. FÆTIDA (Cav. diss. 6. p. 349.) foliis cordatis ovatis acutis dentatis utrinquè tomentosis, petiolis pedicellisque pilosis, stipulis setaceis subpaulis, pedicellis petiolo dimidiò brevioribus, carpellis 7 muticis in globum coarctatis. ① in Peruvia et Martinicâ. S. viscosa L'Her! stirp. 1. p. 52. t. 53. bis. Abutilon fætidum Moench. Pedicelli secus caulem orti. (v. s.)
105. S. BREVIPES, foliis cordatis subrotundis acuminatis dentatis velutino-pubescentibus, petiolis ramisque patentim pilosis, stipulis setaceis subrectis, pedicellis 1-2 axillaribus petiolo quintuplò brevioribus, carpellis 5-6 muticis ①? ad Sanctam-Martham. Bertero. Priori valdè affinis. (v. s.)
106. S. MAGDALENÆ, foliis cordiformibus acuminatis denticulatis petiolis ramisque velutino canescentibus, stipulis setaceis subrectis, pedicellis petiolo paulò brevioribus, carpellis 5 muticis vix vesicariis, seminibus pubescentibus. 5 ad flumen Magdalene. Bertero. Petioli fol. ramealium 6-9 lin. longi. Pedicelli 4-5 lin. longi. Flores flavi, 4-5 lin. diam. (v. s.)
107. S. DENUData (Nees et Mart. nov. act. bonn. 11. p. 500.) foliis cordato-subrotundis acuminatis duplicato-crenatis hirtis, paniculâ terminali nudâ, calyce hispido, carpellis 6-7 mucronatis. 3 in sylvestribus Brasiliæ ad Serra di mundo novo et in prov. Pi Auhienai. Habitus S. nudifloræ, sed carpella 1-sperma.

Sect. III. ABUTILON. Kunth. malv. p. 4.

Carpella 5-30 polysperma sæpè vesicaria.

* *Oligocarpæ*, nempe carpellis 5-8.

108. S. PERIFLOCIFOLIA (Lin. sp. 962.) foliis cordato-lanceolatis acuminatis integerrimis subtile tomentosis, pediculis subdivisis gracillimis petiolo (imò sæpè folio) longioribus sub flore articulatis, carpellis 5 ovatis acuminatis. ① 5? Cav. diss. 1. p. 26. t. 5. f. 2
- a. *Zeylanica*, foliis angustioribus supernè subseabris — in Ceylonâ, Indiâ orient. — Pluk. t. 74. f. 7. — Dill. h. elth. 1. t. 3. f. 2. (v. s.)

- β. *Caribæa*, foliis evidentius cordatis supernè glabris subrugosis. — in Jamaicâ. — Sloan. hist. 1. t. 139. f. 3. Swartz obs. 260. An spec. dist?
- γ? *Peruviana*, foliis evidentius cordatis supernè subvelutinis candicantibus, pedicellis simplicibus folio brevioribus. — in Peruviâ. Dombey. Media inter hanc et sequentem, sed ob pedicellos apice nec mediò articulatos, ad hanc potius referenda. An species distincta! (v. s.)
109. *S. FERRUGINEA*, foliis ovatis subcordatis acuminatis utrinquè tomentosis integerrimis, pedicellis simplicibus 1-floris folio brevioribus medio articulatis, capsulis 5 rostratis. *h* propè Loxam Peruvianorum. Aff. *S. periploci*-foliæ. *Abutilon ferrugineum* H. B. et Kunth nov. gen. am. 5. p. 271. (v. s.)
110. *S. EXCELSIOR* (Cav. diss. 1. p. 27. t. 5. f. 3.) foliis cordatis ovatis acuminatis integerrimis subtus tomentosis, racemis paniculatis, carpellis 5 bidentatis. *h* in Peruviâ calidis.
111. *S. HERNANDIÖIDES* (L'her! stirp. 1. p. 121. t. 58.) foliis subpeltatis cordatis ovatis acuminatis subintegerrimis pubescentibus, pedicellis 1-floris petiolo brevioribus, carpellis 5 muticis. *h* in Hispaniolâ. (v. s.)
112. *S. LUCIANA*, foliis cordatis ovato-rotundis acuminatis subintegerrimis supernè subvelutinis subtus cano-tomentosis, pedunculis ramosis brevissimis in paniculam interruptam dispositis, carpellis 5 subinflatis subbidentatis. — in ins. Sancta-Luciâ. Anderson. Flores albi ex And. (v. s.)
113. *S. SPICIFLORA*, foliis cordatis suborbiculatis valdè acuminatis subcrenulatis pubescentibus subtus canescentibus, spicis terminalibus gracilibus, floribus fasciculatis sub-5-gynis *h* in umbrosis prope S. Carlos del Rio-Negro prov. Guyanæ. Rami canescentes. Flores 10-andri magn. *Geranii parviflori*. Fruct. ign. *Abutilon spicatum* H. B. et Kunth. nov. gen. am. 5. p. 271.*
114. *S. LECHENAULTIANA*, foliis cordatis subrotundis acuminatis integerrimis supernè velutinis subtus canescentibus, pedicellis multifloris axillaribus petioli longitudine, ultimis racemoso-paniculatis, petalis obcordatis, carpellis 5 subbidentatis. — in Indiâ orient. (v. s. ex hort. Calcut. ab amic. Lechenault.)
115. *S. NUDIFLORA* (L'her! stirp. 1. p. 123. t. 59. bis.) foliis subrotundo-cordatis acuminatis subintegerrimis subtus tomentosis, paniculâ terminali racemosâ, carpellis 5-7 subbidentatis. *h* in Sancto-Domingo. *S. stellata*. Cav. diss. 1. t. 5. f. 4. — Plum. sp. t. 3. (v. s.)
116. *S. POLYANTHA* (Schlecht. in Link enum. h. ber. 2. p. 204.) foliis cordatis brevè acuminatis subcrenulatis utrinquè tenuissimè tomentosis virentibus, paniculâ aphyllâ. *h* Patr. ign. Priori simillima. Cor. minor flavissima.
117. *S. AURITA* (Wall. in Link enum. h. ber. 2. p. 206.) foliis sinu angusto profundè cordatis acuminatis serrulatis supernè pilosiss. nbtus canis, stipulis, latis cordato-auritis acuminatis, floribus paniculato-corymbosis; staminum tubo villosissimo, carpellis 5 hirsutis. *h* in Bengalo. (v. s. ex hort. Calc.)
118. *S. TIMORIENSIS*, foliis cordatis subrotundis acuminatis subcrenatis utrinquè velutino-albidis, paniculâ terminali laxâ, carpellis 5-6 oblongis molliter hispidis. — in ins. Timor. (v. s. ex Mus. Par.)
119. *S. TRIQUETRA* (Lin. sp. 962.) foliis cordatis acuminatis serrulatis utrinquè velutino-albidis, pedicellis solitariis 1-floris, carpellis 5 muticis, ramis triquetris. *h* in Amer. calidiori. *S. trisulcata* Jacq. am. 195. — Cav. diss. 1. t. 5. f. 1. Gaertn. fruct. 2. t. 134. — Generis proprii typus ex Kunth. Cor. lutez basi purpureæ. (v. s.)
120. *S. INCANA* (Link. enum. h. ber. 2. p. 204.) foliis incano-tomentosis cordatis subacuminatis acutè crenulatis, pedicellis 1-floris petiolo longioribus, calycibus capsulisque muticis tomentosis. *h* in ins. Sandwich. Caulis subtri-queter ut in priore cui affinis. Cor. calyce parùm major, flava.
121. *S. PULCHELLA* (Bonpl. nav. t. 2.) foliis cordatis ovato-lanceolatis grossè inæqualiter crenatis subtus pube stellatâ subtomentosis supernè scabris, racemis axillaribus petiolo longioribus, carpellis 5 biaristatis. *h* in Novâ-Hol-

landiâ. Flores albi. Carpella 2-sperma. Willd. enum. p. 724. (v. s. in h. Balb.)

122. *S. LIGNOSA* (Cav. diss. 1. p. 28. t. 6. f. 2.) foliis subrotundo-cordatis acuminatis crenatis tomentosis, pedunculis solitariis petiolo longioribus, carpellis 7-8 muticis durissimis. 5 in Sancto-Domingo.
123. *S. OBTUSA* (Cav. diss. 1. p. 34. t. 9. f. 2.) foliis cordatis ovatis obtusis dentatis tricuspidatis, pedunculis sub-4-floris umbellatis petiolo longioribus, carpellis 8 acutis. — Patr. ign.
124. *S. RAMOSA* (Cav. diss. 1. p. 28. t. 6. f. 1.) foliis ovato-cordatis inæqualiter dentatis, racemis axillaribus, carpellis 5 biaristatis. — in Senegal. — Tota pilis mollibus vestita.
125. *S. UMBELLATA* (Lin. sp. 962.) foliis subrotundo-cordatis dentatis subangulatis acuminatis, pedunculis sub-4-floris umbellatis axillaribus, carpellis 6-11 biaristatis. (1) in Jamaicâ. Jacq. hort. vind. t. 56. Cav. diss. 1. t. 6. f. 3. et 5. t. 129. f. 2. (v. s.)
126. *S. GIGANTEA* (Jacq. Schœnb. 2. p. 8. t. 141.) foliis subrotundo-cordatis crenatis acuminato-tricuspidatis utrinquè velutino albidis, floribus paniculatis, corollis reflexis, carpellis 5-7 villosis. 5 ad Caracas. Carpella 8-11 ex Jacq. (v. v.)

** *POLICARPÆ*, nempe carpellis 9 aut ultrâ.

127. *S. REFLEXA* (Cav. diss. 1. p. 36. t. 7. et 6. t. 195. f. 1.) foliis subrotundo-cordatis acuminatis crenatis tomentosis, pedicellis solitariis petiolo longioribus, petalis cuneiformibus apice dentatis reflexis, carpellis 12. 5 in Peruvîæ arenosis. *S. retrorsa* L'her! stirp. 1. t. 64. flor. rubri. (v. s.)
128. *S. PEDUNCULARIS*, foliis subrotundo-ovatis acuminatis profundè-cordatis dentato-repandis suprâ puberulis subtùs molliter canescenti-tomentosis, pedicellis solitariis longissimis, petalis oblongo-spathulatis reflexis, carpellis circ. 20. 5 in umbrosis ad ripam fl. Amazonum. Abutilon pedunculare. H. B. et Kunth nov. gen. am. 5. p. 273. Petala purpurea.
129. *S. MICROSPERMA* (Cav. diss. 1. p. 29. t. 13. f. 4.) foliis subrotundo-cordatis acutis subcrenatis, pedicellis solitariis petiolo brevioribus, carpellis 12-13 birostratis, rostris ciliatis. — Patr. ign.
130. *S. IMBERBIS*, foliis cordatis acutis crenulatis utrinquè albido-velutinis, summis sessilibus, pedicellis solitariis petiolo triplò longioribus, carpellis 12-13 subinflatis subscabris eciliatis muticis. — in Guadalupâ et Sancto-Domingo. Bertero. Sequenti affinis. (v. s.)
131. *S. CRISPA* (Lin. sp. 964.) foliis cordatis acuminatis crenatis subvelutinis, summis sessilibus, pedicellis solitariis petiolo longioribus, fructiferis deflexis, carpellis 12-13 inflatis muticis undulato-crispis carinâ hirtis. 2 in Carolinâ, Novâ-Hispaniâ, Providentiâ, Bahamâ, Peruvîâ. — Dill. h. elth. t. 5. f. 5. — Cav. diss. 1. t. 7. f. 1 et 5. t. 135. f. 2. *S. amplexicaulis* Lam. dict. 1. p. 7. (v. s.)
132. *S. AMPLEXIFOLIA* (fl. mex. ic. ined.) foliis cordatis acuminatis crenatis, summis sessilibus amplexicaulis, paniculâ terminali multiflorâ. 2 in Mexico. An satis à priore diversa? Flores flavi, pedicelli fructiferi erecti. †.
133. *S. VIRGATA* (Cav. ic. 1. p. 53. t. 73.) foliis cordatis ovatis acuminatis serratis subpuberulis, pedicellis solitariis petiolo longioribus, petalis calyce brevioribus, carpellis 9 biaristatis. 5 in Peru. (v. s.)
134. *S. ARBOREA* (Lin. f. suppl. 397. L'her! stirp. p. 131. t. 63.) foliis orbiculato-cordatis acuminatis crenatis tomentosis, pediculis petiolo longioribus, carpellis 13-15 truncatis villosis. 5 in Peru. *S. Peruviana* Cav. diss. 1. p. 36. t. 7. f. 8. et 5. t. 130. *S. grandiflora* Poir. suppl. 1. p. 31. Flores maximi. (v. s.)
135. *S. MAURITIANA* (Jacq. ic. rar. 1. t. 137.) foliis subrotundo-cordatis acuminatis dentatis subtùs tomentosis, pedicellis petiolo longioribus, carpellis 30 truncatis rostratis calyce longioribus. 5 in ins. Mauritiî. *S. planiflora*. Cav. diss. 1. p. 32. t. 7. f. 4 et 5. t. 135. f. 1.

136. *S. TILIÆFOLIA* (Fisch. cat. h. gor. 1808) foliis sinu lato subrotundo-cordatis acuminatis dentatis molliter pubescentibus, pedicellis petiolo brevioribus, carpellis 15 hirsutis biaristatis. (I) in Chinâ aut Thibeto. Willd. enum. 722. Jacq. f. ecl. 1. 1. 35. opt. Flores lutei. Folia ampla. (v. v.)
137. *S. MUTICA* (Delil. ill. eg. n. 633.) foliis orbiculato-cordatis subacuminatis grossè dentatis utrinquè molliter tomentosis, pedicellis petiolo brevioribus, carpellis 25-30 hirsutis muticis vix calyce longioribus. — in Ægypto superiore. (v. s.)
138. *S. AMERICANA* (Lin. sp. 963.) foliis cordatis oblongis indivisis tomentosis, pediculis folio brevioribus, carpellis 12 tomentosis acuminatis calycis longitudine. — in Jamaicâ.
139. *S. ABUTILON* (Lin. sp. 963.) foliis subrotundo-cordatis acuminatis dentatis tomentosis, pedunculis petiolo brevioribus, carpellis 15 truncatis birostratis pilosis. (I) in Galliâ australi, Helvetiâ, Sibirîâ, Indiâ. An species Asiatica eadem ac Europæa? Houtt. syst. 8. t. 61. Abutilon Avicennæ Gaertn. (v. s.)
140. *S. ABUTILOIDES* (Jacq. obs. 1. p. 17. t. 7.) foliis cordatis indivisis acuminatis crenatis tomentosis, pedunculis longitudine petiolorum, carpellis 10 trispermis. (I) in Jamaicâ. *Lavatera Americana* Lin. sp. 973.
141. *S. ASIATICA* (Lin. sp. 964.) folio cordatis ovato oblongis dentatis utrinquè velutinis, pedicellis petiolo longioribus, carpellis 20 truncatis acutis calyci subæqualibus. (I) in Indiâ orient. Cav. diss. 1. p. 31. t. 7. f. 2. et 5. t. 128. f. 1. L'her! stirp. p. 130. (v. s.)
142. *S. SONNE-ATIANA* (Cav. diss. 1. p. 29. t. 6. f. 4.) foliis subrotundo-cordatis acuminatis dentatis tomentosis, pedunculis foliis longioribus, carpellis 9 truncatis obtusis calyce majoribus. — ad cap. B-Spei. *S. Asiaticæ* simillima.
143. *S. IBARENSENSIS*, foliis subrotundis profundè cordatis acuminatis longè petiolatis dentato-crenatis mollissimis tomentosis suprâ incanis subtus albidis, pedunculis axillaribus bifloris, carpellis 16-18 obtusis. ♂ in regno Quitensi. Flores pallidè flavi. *Abutilon Ibarrense* H. B. et Kunth nov. gen. 5. p. 272.*
144. *S. PETIOLARIS*, foliis subrotundis profundè cordatis acuminatis longè petiolatis duplicato-dentatis mollibus suprâ pubescentibus subtus albo-tomentosis, pedicellis axillaribus 1-floris, carpellis 11 apice truncato-angulatis. ♀ in regno Novo-Grauatensi. *Abutilon petiolare* H. B. et K. l. c. Flores albi magni. *S. Abutili*.
145. *S. GEMINIFLORA*, foliis ovatis cordatis acuminatis crenato-serratis utrinquè hirtellis, pedicellis axillaribus geminis 1-floris, floribus 13-gynis, calycibus ferrugineo-tomentosis. ♀? prope Caracas. Affinis *S. sylvaticæ*, sed ovula 4 in quoque carpello. Fr. ign. *Abutilon geminiflorum* H. B. et Kunth nov. gen. 5. p. 274. t. 474. Flores ex albo flavi.
146. *S. POPULIFOLIA* (Lam. dict. 1. p. 7.) foliis subrotundo-cordatis acuminatis inæqualiter repando-dentatis tomentosis, pedunculis petiolo longioribus, carpellis 11-19 truncatis acutis calyce longioribus. (I) in Indiâ orient. Beloere Rheed. mal. 6. t. 65. *S. Beloere*. L'her! stirp. 1. p. 130. *S. populifolia* Cav. diss. 1. p. 32. t. 7. f. 9. et *S. Eteromischos*. Cav. diss. 2. p. 55. ex 5. p. 275. t. 128. f. 2. (v. s.)
147. *S. NIATA* (Lam. dict. 1. p. 7.) foliis subrotundo-cordatis acuminatis dentatis subtus albedo-tomentosis, pedunculis petiolo longioribus, carpellis 15-17 truncatis acutis calyce brevioribus. (I) in Indiâ orient. — Rumph. amb. 4. t. 10? *S. pilosa* L'her! stirp. 1. p. 130. Rami pilis hirsuti, sed multò minùs quàm in sequenti. (v. s.)
148. *S. MOLLIS* (Ort. dec. p. 65.) foliis cordato-orbiculatis duplicato-crenatis pubescenti-mollibus, pedunculis 1-3-floris petiolo brevioribus, carpellis 8-10 acuminatis calyce parùm majoribus, ramis hispidissimis. ♀ in Peru. *S. grandifolia* Willd. enum. p. 724. (v. v.)
149. *S. MOLLISSIMA* (Cav. diss. 2. p. 49. t. 14. f. 1.) foliis subrotundo-cordatis

- acuminatis dentatis velutinis. pedunculis subbifloris petiolò brevioribus, carpellis 11 truncatis bidentatis calyci æqualibus. *h* in Peruvix sylvis ad flum. Maragnou. S. cistiflora L'her! stirp. 1. p. 127. t. 61. (v. s.)
150. S. EXSTIPULÂRIS (Cav. diss. 2. p. 56. t. 14. f. 2.) foliis ovato-cordatis acuminatis utrinquè tomentosis exstipulatis, pedicellis petiolo brevioribus, carpellis 30 truncatis muticis calyce brevioribus. *h* in ins. Borboniâ.
- 151? S. TRUNCATA (Cav. diss. 1 p. 35. t. 6. f. 7.) foliis subrotundis cordatis apice truncato-obtusis dentatis tomentosis, pedicellis petiolo longioribus, carpellis 9 muticis. (1) in Sancto-Domingo. An carpella 1-sperma aut oo-sperma?
152. S. TERMINALIS (Cav. diss. 1. p. 29. t. 6. f. 6. et 6. t. 195. f. 2.) foliis ovato-lanceolatis cordatis dentatis, racemo terminali elongato bracteato. carpellis 9 muticis villosis capitatis calyce majoribus. *h* in Brasiliæ montibus prope Monte-Video.
153. S. PERMOLLIS (Willd. enum. p. 728.) foliis subrotundo-cordatis acuminatis inæqualiter crenatis tomentosis, pedunculis axillaribus solitariis, super. subracemosis, carpellis 9-10 calycem subæquantibus rostratis. *h* Patr. ign.
154. S. CORNUTA (Willd. enum. p. 724.) foliis ovato-cordatis dentatis utrinquè pubescenti mollibus, floribus paniculatis, carpellis inflatis cornutis. *h* in Americâ merid. — An hujus sectionis?
155. S. PUBESCENS (Cav. diss. 1. p. 33. t. 7. f. 6.) foliis subrotundo-cordatis acuminatis angulato-dentatis, pedicellis petiolo longioribus, carpellis 15 bidentatis truncatis calyce majoribus. — in Sancto-Domingo.
156. S. ORBICULATA, foliis ovato-orbiculatis subreniformibus dentatis subtus canescentibus, pedicellis petiolo longioribus, carpellis 13 junioribus tomentosis obtusis, adultis glabris subbidentatis. *h* in Chinâ. (v. s.)
157. S. INDICA (Lin. sp. 964.) foliis cordatis sublobatis mollibus, stipulis reflexis, pedicellis erectis petiolo triplò longioribus, carpellis 13-15 scabris calyce longioribus. (1) in Indiâ orient. Cav. diss. 1. p. 33. t. 7. f. 10. Abutilon elongatum Moench. (v. s.)
158. S. VESICARIA (Cav. diss. 2. p. 55. t. 14. f. 3.) foliis ovato-cordatis dentatis subtricuspidatis, pedicellis petiolo duplò longioribus, carpellis 10 truncatis muticis acutiusculis. *h* in Mexico. An satis à S. Indicâ differt? (v. s.)
159. S. GLAUCA (Cav. ic. 1. p. 8. t. 11.) foliis subrotundo-cordatis acuminatis grossè dentatis tomentoso-glauciscentibus mollibus, pedicellis inf. petiolo brevioribus, sup. longioribus, carpellis 12-15 obtusis calyce brevioribus, caule glanduloso piloso-hirto. *h* in Senegal. Carpella 1-3-sperma. (v. s. in h. Balb.)
160. S. ALBIDA (Willd. enum. p. 722.) foliis subrotundo-cordatis acuminatis dentatis utrinquè tomentoso-mollibus canis, pedicellis petioli longitudine, carpellis 12-15 truncatis acutis calycem subæquantibus hirsutis. *h* in insul. Canariis. Sida n. 340 Boryl! ess. isl. fort. Pedicelli petiolis nunc longiores nunc breviores. (v. s.)
161. S. CRASSIFOLIA (L'her! stirp. 1. p. 125. t. 60.) foliis ovato-cordatis dentatis subtricuspidatis utrinquè cano-tomentosis, pedicellis petioli longitudine, carpellis 9 10 bicuspidatis. *h* in Hispaniolâ. S. tricuspidata. Cav. ic. p. 30. t. 6. f. 5. (v. s.)
ℓ. tomentosa (Cav. descr. 164.) ramis tri-tetraquetris. — in Americâ. (v. s.)
162. S. MOLLICOMA (Willd. enum. p. 725.) foliis ovato-oblongis cordatis inæqualiter dentatis truncatis mollissimè tomentosis, pedicellis petiolo brevioribus, carpellis 9 acuminatis inflatis. *h* Patr. ign. — S. sericea Cav. descr. p. 164? — Sub hoc nom. vidi in h. Balb. specimen descriptioni sat congruens, sed rami hispidissimi, folia subrotundiora, pedicelli vix petiolo breviores. An eadem?
163. S. VITIFOLIA (Cav. ic. 5. t. 428.) foliis cordatis 5-7-lobatis lobis acumi-

natis serratis, pedunculis petiolo longioribus ramoso-umbellatis, carpellis 9 apice longè biaristatis. 5 in regno Chilensi. Flores magni rosei

164. *S. ACERIFOLIA* (Lag. nov. gen. p. 21.) foliis cordatis subpeltatis 3-5-lobis inæqualiter dentatis villosis, pedicellis 1-floris petiolo longioribus, carpellis 10-12 inflatis basi externâ biaristatis. 5 in Novâ-Hispaniâ. Tota hispida. Flores cærulei. *S. spinifex* et forsân *S. palmata* fl. mex. ic. ined. (v. s.)

† *Species non satis notæ, præsertim quoad fructûs structuram.* (1)

* *Foliis linearibus oblongis ovatis lanceolatisve.*

165. *S. PAUCIFOLIA*, glabra ramosissima, foliis linearibus integris, pedicellis axillaribus 1-floris folio brevioribus sub flore articulatis post anthesim persistentibus. 5 in Timor. Species distinctissima, forsân *S. linifolia* affinis, sed fructus ignotus. (v. s. ex Mus. Par.)

166. *S. MILLERI*, foliis lineari-lanceolatis dentatis subtus villosis, pedicellis axillaribus 1-floris, caule erecto ramoso. — *S. angustifolia* Mill. dict. n. 3. Flores parvi, flavi. Semen bidentis.

167. *S. FRUTICOSA* (Mill. dict. n. 18.) foliis lanceolatis inæqualiter serratis acuminatis, floribus capitatis terminalibus, carpellis 5 bidentatis. 5 in Jamaicâ. Flores parvi albi.

168. S? *HISPIDA* (Pursh. fl. bor. am. 2. p. 452.) hispido-pilosa, foliis lanceolatis serratis, pedicellis solitariis axillaribus longitudine foliorum, involucello filiformi. 2 in Georgiæ campis arenosis. Flores lutei. An *Malvæ* species Nat?

169. *S. VIMINEA* (Fisch. in Link. enum. h. ber. 2. p. 202.) foliis lanceolatis longissimis integerrimis pilosis, racemis terminalibus brevissimis. 2 in Brasiliâ. Caulis erectus calyxque strigosi. Cor. calyce major basi aurantiaca.

170. *S. SEMICRENATA* (Link. l. c.) foliis latè lanceolatis obtusis crenatis basi integerrimis trinerviis, junioribus subtus tomentosis, pedunculis axillaribus 1-floris, calycis laciniis longè acuminatis. 2 in Manilâ. Caulis erectus glaber. Calyx subtomentosus Corolla flava parùm calyce major.

171. *S. ACRANTHA* (Link. l. c.) foliis mediis oblongis obtusis anticè acutè crenatis, summis lanceolatis acutis anticè serratis, floribus terminalibus subsessilibus. 5 in Brasiliâ. Caulis erectus. Corolla flava calyce hirtò acuminato parùm major.

172. *S. SPIRÆIFOLIA* (Willd. enum. suppl. 49. Link. en. 2. p. 203.) foliis oblongo-lanceolatis serratis basi integerrimis glabris, pedunculis axillaribus solitariis petiolo longioribus, calyce glabro, laciniis acuminatis. 5 Patr. ign. Corolla flava calyce major. Capsula birostris ex Link.

173. *S. SCHRANKII*, foliis oblongo-lanceolatis serratis basi integerrimis subtus tomentosis, pedunculis in ramulis confertis, calyce subtomentoso, laciniis acutis. 2 in Brasiliâ. Priori affinis. Corolla flava parùm major. *S. brasila* Schrank. in Link. enum. h. ber. 2. p. 203. non Cav.

174. *S. BETULINA* (Horn. h. hæfn. 2. p. 646.) foliis ovato-oblongis subduplicato-serratis basi integerrimis, pedunculis axillaribus multifloris. 5 (1) in Americâ merid. ?

175. *S. PATULA* (Pers. ench. 2. p. 243.) herbacea patula flexuosa, foliis brevipetiolatis ovatis subcrenatis mollissimè tomentosis, floribus axillaribus sessilibus. — in Cayennâ. *S. mollis* Richard act. soc. h. nat. par. Flores albi.

176. *S. BRASILIENSIS* (Cav. diss. 1. p. 37. t. 34. f. 1.) foliis ovatis acuminatis 5-nerviis vix denticulatis subtus ramisque tomentosis, stipulis filiformibus, pedicellis 1-floris petiolos æquantibus, capsulis hirtis. — in Brasiliâ. Schrad. ill. pl. bras. in Gœtt. an z.

(1) Pleræque ad *Malvindas* referendæ sed ob descr. incompl. non specificècum *Malvindis* suprâ descriptis facillè conferendæ.

** *Foliis cordatis indivisis.*

177. *S. GRACILIS* (Rich. act. soc. h. nat. par. p. III.) foliis oblongo-cordatis integerrimis sursum angustatis reflexis, ramulis fasciculifloris, caule fruticoso fragili. 5 in Antillis.
178. *S. CARNEA* (fl. mex. ic. ined.) foliis ovato-oblongis subcordatis dentatis acutis, pedicellis solitariis axillaribus petiolo longioribus, carpellis 5 distinctis. 5 in Mexico. Caulis ramosissimus. Flores carnei parvi. Stigmata purpurea.
179. *S. VILLOSA* (Mill. dict. n. 6.) foliis subcordatis sessilibus serratis subvillosis, floribus axillaribus confertis, carpellis bidentatis. 5 in America calid. Fruticulus 2-3-pedalis erectus hirsutus. Flores pallide flavi.
180. *S. VERRUCULATA*, caule verruculato-pubescente, foliis cordato-lanceolatis acuminatis acutè crenatis pubescentibus, pedicellis axillaribus 1-floris filiformibus petiolo parùm longioribus. 5 in Brasilià. *S. arguta* Fisch. in Link. enum. h. ber. 2. p. 206. non Sw. Petala flava cal. parùm majora. Fruct. ign.
181. *S. GRAVEOLENS* (Roxb. ex Hortn. h. hafn. suppl. 77.) foliis subrotundocordatis subacuminatis, pedunculis petiolo brevioribus, carpellis truncatis calyce longioribus. (1) in Indià orientali.
182. *S. PURPURASCENS* (Link enum. h. ber. 2. p. 206.) foliis cordatis acuminatis crenato-denticulatis canibusque tomentoso-virentibus, pedicellis axillaribus 1-floris petiolo parùm longioribus. 5 in Brasilià. Petioli longi limbo tamen breviores. Nervi subtus flavicantes. Petala ungue angusto. Fr. ign.
183. *S. PATENS* (Andr. bot. rep. 571.) foliis cordatis acuminatis inciso-serratis, pedunculis solitariis petiolo longioribus, carpellis 5 aristatis. (2) in Abyssinià.
184. *S. CONTRACTA* (Link enum. h. ber. 2. p. 204.) foliis cordatis acuminatis repandis rariùs crenatis cano-tomentosis, paniculà contractà bracteolatà, calycibus tomentosis. 5 in Madagascar. Affinis videtur *S. nudifloræ* sed flor. et fruct. ign.
185. *S. CONFERTA* (Link. l. c. p. 207.) foliis cordatis acutis crenatis rugosis canibusque flavicanti-tomentosis, floribus sessilibus aggregatis. 5 in Brasilià. Cor. obscure flava calyce parùm longior. Calyx dense tomentosus.
186. *S. HIRUTA* (Mill. dict. n. 9.) foliis orbiculatis, cordatis crenatis, caule petiolisque hirsutis, pedunculis longis axillaribus 1-floris. — Patr. ign. Sessipedalis. Flores albi parvi.
187. *S. PERSICA* (Burm. ind. t. 47. f. 1.) foliis inferioribus petiolatis cordatis acuminatis, superioribus sessilibus lanceolatis dentatis, pedicellis solitariis 1-floris. — in Persià. Cav. diss. 1. p. 35. t. 4. f. 1.
188. *S. LASIOSTEGA* (Link enum. h. ber. 2. p. 205.) foliis cordatis acuminatis subtus caescenti-tomentosis, pedicellis axillaribus 1-floris petiolo longioribus calycibus capsulisque muticis tomentoso-villosis. 5 in Brasilià. Caulis subteres incano-tomentosus. Cor. calyce parùm major flava.

*** *Foliis cordatis tricuspidatis aut trilobis.*

189. *S. BIFLORA* (Cav. diss. 1. p. 37. t. 9. f. 1.) foliis ovato-cordatis acuminatis dentatis tricuspidatis, pedunculis geminatis axillaribus petiolo æqualibus. 5 Patr. ign.
190. *S. PENTACARPUS* (Roxb. ex Horn. suppl. 78.) foliis cordatis subtricuspidatis, pedunculis longitudine petioli, carpellis muticis hispidissimis. (1) in Indià.
191. *S. SESSEI* (Lag. nov. gen. 21.) foliis cordato-ovatis subtrilobis acuminatis dentatis subtus præcipuè tomentosis, pedunculis axillaribus solitariis geminisque petiolo longioribus subbifloris, carpellis 5. 5 in Novà-Hispanià.

**** *Foliis palmatilobatis,*

192. *S. GERANIODES*, foliis subtus cano-tomentosis superne subvelutis palmato-5-lobis, lobis inciso-dentatis medio elongato, pedicellis axill. solit. 1-floris petioli longitudine, calyce tomentoso campanulato truncato obtusè 5-6 dentato. 5. in Novâ-Hollandiâ. Fructus ignotus. Petala cal. duplò majora. Calyx in genere anomalus. An potius *Lagunæa*? (v. s. ex Mns. Par.)
193. *S. ALCEOIDES* (Michx. fl. bor. am. 2. p. 44.) foliis inferioribus triangulari-cordatis incis, superioribus palmato-multifidis, corymbo terminali, calycibus hispidis. — in glareosis Kentucky et Tenassee. Facies *Malvæ Alceæ*.
194. *S. MALVÆFLORA* (fl. mex. ic. ined.) herbacea glabrescens, foliis longè petiolatis semi-orbicularibus basi truncatis palmatim 5-9 fidis, lobis apice 3-dentatis, racemo terminali. — in Mexico. Facies *Malvæ Alceæ* ut in præced. spec.
195. *S. PHYLLANTHOS* (Cav. diss. 5. p. 276. t. 127. f. 4.) acaulis, foliis tripartitis, segmentis sessilibus 3-partitis cuneiformibus integerrimis obtusis, flore solitario petiolo alato inserto, carpellis 12. 2. in Peru. Species maximè singularis nulli affinis! Folia tomentosa inferne rufescentia. Flos magnus cæruleus ex Cav., aut pallidè violaceus ex H. B. et Kunth nov. gen. am. 5. p. 264.*

XXII. *LAGUNEA*. Cav. diss. 3. p. 171. — *Solandra Murr. comm. goett.* 1784. Lam. ill. t. 580. non Swartz.

Calyx nudus 5-fidus. Antheræ in apice et superficie tubi. Stigmata 5. Capsula 5 loc. 5-valvis, valvis medio septiferis solubilibus, axi centrali filiformi superstiti. — Est *Hibisco* ut *Sida Malvæ*.

1. *L. LOBATA* (Willd sp. 3. p. 733.) foliis cordatis trilobis, lobis ovali-oblongis acuminatis dentatis, sinu angustissimo (1) in ins. Borboniâ. *Solandra lobata*. Murr. comm. Goett. 1784. p. 21. t. 1. *Tignera acerifolia* Cav. diss. 1. p. 41. t. 11. f. ult. *Lagunea lobata* Cav. diss. 5. t. 136. f. 1. *Hibiscus solandra* L'her! stirp. 1. t. 49. (v. s.)
2. *L. SINUATA* (Horn. h. hafn. 645.) foliis cordatis trilobis, lobis oblongis acuminatis dentatis, sinubus dilatatis. (1) Patr. ign. An prioris varietas? *L. angulata* hortul. (v. s.)
3. *L. TERNATA* (Cav. diss. 5. p. 279. t. 136. f. 2.) foliis inferioribus tripartitis lobis oblongis integris medio longissimo, superioris subhastatis elongatis. — Ad Senegal. Caulis herbaceus villosus.
4. *L. ACULEATA* (Cav. diss. 3. p. 173. t. 71. f. 1.) foliis tripartitis, lobis oblongo-linearibus inciso-dentatis, caule aculeato tomentoso. — prope Pondicheri. Calyx longitudinaliter rumpens.

XXIII. *INGENHOUIA*. fl. mex. ic. ined. †.

Cal. nudus 3-partitus, lobis ovato-lanceolatis acuminatis Pet. 5. Urceolus campanulatus intra petala. Stam. ∞ monadelphæ. Stylus 1. Fruct.

1. *I. TRILOBA* (fl. mex. ic. ined.) — in Mexico. Herba habitu *Gossypii*. Folia petiolata triloba, lobis ovato-lanceolatis acutis integris. Pedicelli oppositi folii 1-flori. Flores ex flavo rubescentes.

Ordo XXV. BOMBACEÆ. *Kunth diss. malv. p. 5. nov. gen. am. 5. p. 294.*

Calyx basi nudus bracteolisve paucis involuclatus, constans sepalis 5 in tubum urceolato-campanulatum cylindraceumve concretis, apice nunc truncatis nunc irregulariter imbricatis nunc subvalvatim connatis et lateraliter rumpentibus, æstivatione ideò dubiâ nec ut in Malvaceis Buttneriaceisve regulariter valvatâ. Petala 5 regularia aut nulla et tunc calyx intus coloratus. Stamina 5?-10-15 aut plura, filamentis infernè in tubum petalorum basi adnatum concretis, apice subpentadelphis, adelphiis mono aut polyantheris, filis nonnullis interdum sterilibus. Antheræ uniloculares. Ovarium carpellis 5 (rariùs 10) nunc subdistinctis nunc arcte coalitis et variè dehiscentibus constans. Styli totidem distincti aut plus minùs coaliti. Fructus varius. Semina sæpè lanâ aut pulpâ involutâ, alia exalbuminosa cotyledonibus corrugatis convolutisve, alia albuminosa cotyledonibus planis.

Arbores aut Frutices regionum tropicarum incolæ. Folia alterna, bistipulata. Pubes partium herbacearum stellulata.

Ordo dubius et recognoscendus, Malvaceis valdè affinis ob antheras 1-loculares, stamina monadelphe, petala sæpè convoluta et habitum, sed ab iis distinctus calyce non verè valvato, staminum tubo apice pentadelpho. Bombaceæ accedunt etiam ad Buttneriaceas et Chlenaceas, sed antheræ uni nec biloculares ab iis faciliùs segregant.

I. HELICTERES. *Lin. gen. n. 1024.*

Calyx tubulosus sub-5-fidus. Petala 5 ligulato-unguiculata supernè subdentata. Stamina 5-10-15 longè monadelphe, urceolo apice multifido pilis nonnullis sterilibus. Ovarium longè stipitatum. Styli 5 basi concreti. Carpella 5 1-loc. ∞-sperma intus dehiscentia, sæpè in spiram regulariter contorta, interdum recta. Semina exalbuminosa, cotyledonibus spiraliter convolutis.

Sect. I. SPIROCARPÆA.

Carpella in spiram contorta, fructum oblongum ovatumve spiraliter 5-sulcatum constituentia.

1. *H. ISORA* (Lin. sp. 1366.) decandra, foliis cordatis ovatis dentato-serratis acuminatis scabris subtus tomentosis, floribus axillaribus, fructu cylindraceo subvelutino apice subulato. ♂ in Malabarâ et Moluccis. — Pluk. t. 245. f. 2. Rumph. amb. 7. t. 17. f. 1. *H. Jamaicensis* Lam. dict. 3. p. 87? folia Coryli. Petala obovata. (v. s.)
2. *H. BARUENSIS* (Lin. mant. 122.) decandra, foliis cordatis argutè serratis subtus tomentosis, pedunculis bifloris, calyce subbilabiato, fructu cylindraceo subvelutino apice subulato. ♂ in sylvat. maritim. ins. Baru et isthmi Panama. Jacq. am. 236. t. 149. Calycis lobis 5, in labia dua irregulariter concreti. Petala linearia.

♂? ovata, foliis ovatis vix cordatis obtusis dentatis subtus tomentosis. — Patr. ign. Fructus iconi Jacq. similis, sed flores ignoti et folia diversa. An spec. propria?

3. *H. GUAEUMÆFOLIA* (H. B. et K. nov. gen. am. 5. p. 304.) decandra, foliis obsoletè cordatis subovatis inæquilateralis acuminatis serratis suprâ hirtellis subtus tenuiter albido-tomentosis, pedunculis bifloris, calyce 5-lobato, petalis spatulatis. 5 ad ripam Oriuoci et prope Caripe Cumanensium. Similis *H. Baruensi*.
4. *H. GREWIEFOLIA*, decandra, foliis ovatis vix basi subcordatis acutis inæqualiter duplicato-serratis utrinquè tomentosis, fructu cylindraceo subvelutino apice subulato. 5 in ins. Timor. Petala longè unguiculata oblonga. Fructus *H. Baruensis* sed gracilior. (v. s. ex h. Mus. Par.)
5. *H. MEXICANA* (H. B. et K. nov. gen. am. 5. p. 305.) decandra, foliis ovatis subcordatis inæqualiter serratis mollibus suprâ pubescentibus, subtus incanis villosis-tomentosis, calyce acutè 5-dentato, fructu ovato subvelutino. 5 inter Mazatlan et Chilpancingo Mexicanorum. *H. rubra* fl. mex. ic. ined. Petala æqualia apice spatulata rubra. Calyces velutini. Genitalia vix corollâ duplè longiora.
6. *H. JAMAÏCENSIS* (Jacq. amer. 235. t. 170. f. 99.) decandra, foliis cordatis crenatis utrinquè velutino-tomentosis, floribus subterminalibus paucis corymbosis, fructu ovato densè tomentoso. 5 in Jamaïcæ et Sancti-Thomasii fruticetis. — Pluk. t. 245. f. 3. Swartz. fl. 2. p. 1156. An ab hac differt *H. althæefolia* Lam. dict. 3. p. 88. (v. s.)
7. *H. VERBASCIFOLIA* (Link. enum. 2. p. 200.) foliis cordatis acuminatis serratis tomentosis virentibus, pedunculis axillaribus paucifloris, carpophoro longissimo. 5 in Brasiliâ. Cor. rubra. Carpoph. 4 poll. longum. — Lodd. bot. cab. t. 504.
8. *H. FERRUGINATA* (Link. enum. 2. p. 199.) foliis cordato-lanceolatis crenulatis tomentoso subtus ferrugineo-candicantibus, floribus terminalibus subracemosis, carpophoro longissimo: 5 in Brasiliâ. Cor. flavescentes. Carpophorum 2-3 poll. longum.
9. *H. PENTANDRA* (Lin. mant. 294.) pentandra, foliis ovatis, floralibus coloratis. 5 ad Surinamum. Calyces hispidi setis ramosis.

Sect. II. ORTHOCARPÆA.

Carpella recta approximata nempè nec in spiram contorta nec radiantia.

10. *H. ANGUSTIFOLIA* (Lin. sp. 1366.) decandra, foliis lanceolatis integerrimis subtus tomentosis, pedicellis bifloris, fructu ovato subvelutino. 5 in Chinâ. Obs. itin. 232. t. 5. Lour. coch. 2. p. 647. (v. s.)
 11. *H. HIRSUTA* (Lour. coch. 2. p. 648.) decandra, foliis ovatis (cordatisve ex Lour.) acuminatis serratis tomentosis, pedunculis multifloris, fructu oblongo hirsutissimo. 5 in sylvis Cochinchinæ Willd. sp. 3. p. 721. Flores obscurè purpurei.
 12. *H. PRONIFLORA* (Rich. act. soc. h. n. par. p. 111.) hexandra, foliis cordatis ovalibus inæqualiter serratis acuminatis in nervis subtus petiolis ramulisque hirsutis, spicis axillaribus 3-4-floris brevibus. 5 in Cayennâ. Folia in planum unicum patentia. (v. s.)
 13. *H. CARTHAGENENSIS* (Lin. sp. 1366.) polyandra, foliis cordatis serratis utrinquè tomentosis, floribus subsessilibus subcorymbosis, fructu oblongo. 5 in Carthagenæ sylvis. Flores foetidi, purpurei.
- † *Species non satis notæ.*
14. *H. LANCEOLATA*, foliis lanceolatis utrinquè acuminatis integerrimis superne glabris subtus canis, floribus axillaribus breviter racemoso-fasciculatis, petalis linearibus genitalia superantibus, 1 ad basim limbi bidentato. 5 in Indiâ orient. Species distinctissima sed fruct. ign. (v. s. ex hort. Calcut. comm. ab amic. Lechenault.)
 15. *H. SEMITRILoba* (Bert! in litt.) foliis subcordatis lato-ovatis subtrilobis crenato-dentatis superne glabris subtus velutino-canis 5-nerviis reticulatis, floribus terminalibus corymbosis. 5 in Sancto-Domingo. Fructus ign. (v. s.)
 16. *H? UNDULATA* (Lour. coch. 2. p. 649.) foliis lanceolatis undulatis, floribus subterminalibus congestis, carpellis rectis stellatim radiatis. 5 in sylvis Cochinchinæ. Nonne potiùs *Sterculiæ* species?
 17. *H? PANICULATA* (Lour. coch. 2. p. 649.) foliis ovatis acutis, floribus sub-

terminalibus laxè paniculatis, carpellis rectis stellatim radiatis. ♂ in sylvis Cochinchinæ. Nonne ut præced. spec. Sterculiæ species?

II. MYRODIA. Schreb. gen. n. 1147.

Calyx nudus tubulosus 4-5-dentatus lateraliter rumpens. Petala oblongo-linearia. Stamina columnæ longa. Antheræ 10-15. Capsula drupacea 2-3-locularis, loculis 1-spermis. — Bracteolæ 2-3 in pedicello floris.

Sect. I. EUMYRODIA.

Antheræ omnes ad apicem columnæ stamineæ.

1. *M. TURBINATA* (Swartz fl. ind. occ. 2. p. 1227.) foliis ovato-oblongis, calycibus turbinatis, columnâ staminum petalis brevioribus, ramis sparsis. ♂ in ins. Caribæis ad torrentum ripas et in Mexico. *M. ovata*. fl. mex. ic. ined. (v. s.)
2. *M. VERTICILLARIS* (fl. mex. ic. ined.) foliis oblongis utrinquè acuminatis, calycibus turbinatis irregulariter sinuatis, staminum columnâ petalis subbrevioribus, ramis floriferis verticillatis. ♂ in Mexico.

Sect. II. QUARARIBEA.

Antheræ secus columnam stamineam sparsæ.

3. *M. LONGIFLORA* (Swartz fl. ind. occ. 2. p. 1229.) foliis lanceolato-oblongis, calycibus cylindraceis, columnâ staminum petalis longioribus. ♂ in Guianâ ad fluviorum ripas. *Quararibea Guianensis* Aubl. l. c. t. 278. Cav. diss. 3. t. 71. f. 2. (v. s.)

III. PLAGIANTHUS. Forst. gen. t. 43.

Calyx nudus 5-fidus. Petala 5 ovata, 2 approximata à reliquis remota. Stigma clavatum. Bacca †.

1. *P. DIVARICATUS* (Forst. loc. cit.) — in Novâ-Zelandiâ. Rami divaricati; folia parva, fasciculata, linearia; flores solitarii.

IV. MATISIA. Humb. et Bonpl. pl. equin. 1. t. 2. nov. gen. am. 5. p. 306.

Calyx nudus irregulariter 2-5-dentatus rumpens, persistens. Petala ovata. Staminum columnæ apice 5-fida, antheris in quoque lobo 12, ad latus externum secundis. Stigma 5-sulcatum. Drupa ovata 5-locularis, loculis 1-spermis. Albumen farinaceum? Cotyledones corrugatæ.

1. *M. CORDATA* (H. B. loc. cit.) ♂ in regni Novæ-Granatensis et Peruviae calidis. Arbor excelsa, foliis petiolatis cordatis 7-nerviis glabris, floribus cernuis fasciculatis albo-röseis.

V. POURRETIA. Willd. sp. 3. p. 844. — Cavanillesia. fl. per.

Calyx nudus 5-partitus campanulatus persistens. Pet. 5. Stam. concreta in cylindrum apice 5-adelphum. Antheræ 1-loculares. Stigma capitatum. Capsula coriaceo-membranacea 5-ptera, alis foliaceis maximis, 1-locul. indehiscens, loculis 1-spermis, plurimis sæpè abortivis. Cotyledones chrysaloïdeo-contortuplicatæ. Corr. ann. mus. 9. p. 293. t. 26.

1. *P. ARBOREA* (Willd. l. c.) foliis cordatis. ♂ in Peruviae Andibus. *Cavanillesia umbellata*. Ruiz. et Pav. prod. p. 97. t. 20. Arbor trunco crasso ad medium quasi tumefacto, ligno fungoso. Flores umbellati, rubri, fugacissimi. (v. s. fr.)
2. *P. PLATANIFOLIA* (H. et Bonpl. pl. equin. 2. p. 162. t. 133.) foliis subpeltatis 5-7-lobatis. ♂ in prov. Carthagenensi. Petala carnea externè ferrugineo-tomentosa. *Cavanillesia platanifolia*. H. B. et Kunth. nov. gen. am. 5. p. 306.

VI. MONTEZUMA. fl. mex. ic. ined.

Calyx nudus hemisphæricus truncatus sinuato-dentatus. Petala 5 subsinuata maxima. Stam. ∞ spiraliter circa stylum longè monadelphæ sulcis 5 subdistincta. Stylus in stigma clavato-ligulatum desinens. Bacca globosa 4-5 loc., loculis polyspermis. Sem.

1. *M. SPECIOSISSIMA* (fl. mex. ic. ined.) ♂ in Mexico. Folia glabra, cordiformia,

acuta, integra, petiolata. Pediculi 1-flori infrà folia e ramis orti. Flores purpureo-phænicei ampli.

VII. OPHELUS. *Lour. coch. 2. p. 501.*

Cal nudus 5-fidus, lobis acutis patenti-reflexis. Pet. 5 crassa. Stam. ∞ in tubulum basi coalita supernè subreflexa. Stigma multifidum. Bacca lignosa oblongo-ovata 12-locul. ∞ -sperma. Sem. . . . Adansonie affinis videtur.

1. *O. SITALIARIUS* (Lour. l. c.) ζ ad littora Africæ orientalis. Folia sparsa oblonga integerrima glabra petiolata. Flores albi solitarii 3 poll. diam.

VIII. ADANSONIA. *Lin. gen. n. 836.*

Calyx nudus, deciduus, 5-partitus. Petala 5 ad medium ferè coalita. Stamina urceolus supernè dilatato-expansus. Stylus longissimus. Stigmata plura. Capsula indehiscens lignosa 10 locul., loculis polyspermis pulpâ farinaceâ circa semina repletis. Folia palmatim composita.

1. *A. DIGITATA* (Lin. sp. 960.) ζ in Senegal. Cav. diss. 5. p. 298. t. 15. Lam. ill. t. 588. A baobab. Gaertn. fr. 2. p. 253. t. 135. — Baobab. Alp. æg. 66. t. 67. Arbor trunco valdè crasso diametro 25-pedali. (v. s.)

IX. CAROLINEA. *Lin. f. suppl. 314.*—*Pachira Aubl. guian. 2. p. 725.*

Calyx nudus, subtruncatus, persistens. Petala 5 oblonga longissima. Stamina basi monadelpha, supernè in adelphas plures dodecandras fasciculata. Stylus longissimus. Stigm. 5. Capsula lignosa multivalvis, 1-locularis, polysperma. Semina nec gossypio nec farinâ induta, arillo carnoso forsân cincta. Cotyl., ex Lin. f., plicatæ. — Folia palmatim composita.

1. *C. PRINCEPS* (Lin. f. suppl. 314.) foliolis 5-8 ovato-lanceolatis acuminatis. ζ in Guianæ locis aquâ marinâ submersis *Pachira aquatica* Aubl. guian. 2. p. 725. t. 291. 292. Cav. diss. 3. p. 176. t. 72. f. 1. Lam. ill. t. 589. Flores maximî speciosissimi, petalis supernè flavis, subtùs virentibus. Filamenta rubra. Anthere purpureæ. An ab hac satis differt *Pachira nitida* H. B. et Kunth nov. gen. am. 5. p. 302? (v. s.)
2. *C. FASTUOSA* (fl. mex. ic. ined.) foliolis 5 obovatis obtusis, calyce truncato integerrimo, petalis à calyce revolutis. ζ in Novæ Hispaniæ locis moderatè calidis. Xiloxochitl flore capillaceo. Hern. mex. 68. ic. opt. Flores sanguinei. Foliola apice subemarginata, interdum subpurpurascens. Stamina monadelpha nec, ut videtur ex icon., polyadelpha!
3. *C. INSIGNIS* (Swartz fl. ind. occ. 2. p. 1202.) foliolis 5-7 obovato-oblongis, calyce sinuato, petalis erectis apice summo patulis. ζ in Martinicâ et Tabago, solo arido arenoso. Flores ruberrimi. *Bombax grandiflorum*. Cav. diss. 5. p. 295. t. 154. (v. s.)
4. *C. MINOR* (Sims bot. mag. t. 1412.) foliolis 7 elliptico-oblongis utrinquè acutis, calyce truncato, petalis erectis. ζ in Mexico. *C. pompalis* fl. mex. ic. ined. *Bombax carolinoides*. Donn. cant. 156. Petala virescentia; filamenta rubra; antheræ luteæ. Pedicelli calyce longiores, sed in icone fl. mex. multò breviores quàm in Simsianâ.

†. *Species non satis nota.*

5. *C. ALBA* (Lodd. bot. cab. t. 752.) ζ in Brasiliâ. Arbor magnifica, foliis digitatis, floribus albis graveolentibus ad apices ramorum. Calyx mellifluus. Filam. innumera bifurca basi in tubum coalita.

X. BOMBAX. *Lin. gen. n. 835. excl. spec. plurim.*

Calyx nudus subquinquefidus truncatusve. Petala 5 inter se et cum staminum columnâ basi subconnexa. Stamina plurima, nunc purè monadelpha, nunc apice pentadelpha. Capsula magna, 5-locularis, 5-valvis, lignosa, loculis polyspermis. Semina lanugine densâ obvoluta, albuminosa.

1. *B. CRIBA* (Lin. sp. 959.) candice aculeato, foliis palmatis 5-foliolatis, fructu

- turbinato apice concavo. § in Americâ calidiore, Jamaicâ, etc. *B. quinatum* Jacq. am. 192. t. 176. f. 1. Arbor vastissima vulgò Ceiba.
2. *B. MALABARICUM*, caudice aculeato, foliis palmatis 7-foliolatis, foliolis integerrimis acuminatis, fructu oblongo obtuso. § in Malabariâ. Bengalo. — Rheed. mal. 3. t. 52. *B. heptaphyllum* Cav. diss. 5. p. 296. Roxb. cor. 3. t. 247. Petala intus coccinea, extus pallida; calyx irregulariter 2-3-lobatus.
3. *B. SEPTENATUM* (Jacq. am. 191. excl. syn.) caudice inermi suberoso, foliis palmatis 7-foliolatis, foliolis integerrimis. § Carthagenæ. *B. heptaphyllum* Lin. sp. 960. Fructus Eriodendria anfractuosi. †.
4. *B. BUONOPONENSE* (Beauv. fl. ow. 2. p. 42. t. 83.) caudice inermi. foliis ..., calyce integerrimo. § in regno Owariensi prope Buonopozo. Calyx cupulæformis, margine zonatus parvus. Petala intus purpurea, extus lanuginosa. (v. s. fl.)
5. *B. GLOBOSUM* (Aubl. guian. 2. p. 701. t. 281.) caudice inermi, foliis palmatis, foliolis 5 ovalibus obtusis integris, calyce obtusè 5-lobato, fructu globoso. § in Guianâ. Cav. diss. 5. p. 297. t. 55. Flores Anbletio ignoti, fasciculatim ex ramis orti, pedicellis 1-floris. Calyces glabri. Petala oblonga extus lanuginosa, intus pallida. Stam. petalis breviora. (v. s.)
6. *B. ELLIPTICUM* (H. B. et K. nov. gen. am. 5. p. 299.) caudice inermi, foliis palmatis, foliolis 5 ellipticis utrinquè rotundatis apice retusis integerrimis glabris, calyce integro. § Petala rosea extus fusciscentia. Stam. fil. rubra, longitudine petalorum.
7. *B. DISCOLOR* (H. B. et K. l. c.) caudice inermi, foliis palmatis, foliolis 5 oblongis acuminatis crenulatis supra hirsutis et viridibus, subtus albidotomentosis. § in calidis prope S. Felipe prov. Jaen de Bracamoros. Flores albi magnitudine Citri.

†. *Species non satis notæ.*

8. *B. VILLOSUM* (Mill. dict. n. 3.) foliis quinquangulatis villosis, caule geniculato. § in Americâ Hispanicâ. Flores ignoti.
9. *B. CUMANENSE* (H. B. et K. nov. gen. am. 5. p. 300.) inermis? foliis palmatis, foliolis 7 petiolatis lanceolatis acuminatis integerrimis glabris subtus pallidioribus. § prope Cumana. An à *B. septenatum* differt?
10. *B. MOMPOXENSE* (H. B. et K. l. c.) aculeatum, foliis palmatis, foliolis 7-9 subsessilibus obovato-lanceolatis acuminatis versus apicem remote serrulatis membranaceis glabris. § ad ripam fl. Magdalenæ prope Mompox. Fl. et fr. ign.
11. *B. ORINOCENSE* (H. B. et K. l. c.) inermis, foliis palmatis, foliolis 5-6 oblongis acuminatis integerrimis membranaceis glabris § in ripâ Orinoci. Fl. ign.
- XI. *ERIODENDRON*. — *Ceiba Plum. gen. 42. t. 32. Gaertn. fr. 2. p. 244. t. 133.*

Calyx nudus obtusè 5-lobatus. Petala 5 inter se et cum staminum columnâ connexa. Stam. filamenta basi in tubum brevem coalita, apice pentadelphæ, adelphiis ad summum connexis filiformibus apice antheras 2-3 gerentes lineares aut anfractuosas et antheram unicam simulant. Cetera Bombacis.

1. *E. ELIANTHERUM*, antheris rectilineis. § in Brasiliâ prope S. Sebastianum. *Bombax elianthos* Cav. diss. 5. p. 294. t. 152. f. 1. Truncus aculeatus. Folia 7-foliolata. Corollæ magnæ extus lanatæ.
2. *E. ANFRACUOSUM*, antheris anfractuosis foliolis integris, caudice sæpius aculeato. § *Bombax pentandrum* Lin. sp. 959 Cav. diss. 5. p. 293. t. 151. Truncus aculeatus aut variis inermis. Folia 7-foliolata. Corollæ minores, extus sericeo-lanatæ. — An hic 3 confusæ species, nempe:
- a. *Indicum*, floribus intus flavescentibus. — in Indiâ orientali. — Rheed. mal. 3. t. 49 51. Rumph. amb. 1. t. 80. (v. s.)
- β. *Caribæum*, floribus intus roseis fasciculatis. — in ins. Caribæis. Jacq. am. 191. t. 176. f. 70. (v. s. fl.)
- γ. *Africanum* (Brown cong. p. 10.) ad ripas flum. Congo.
3. *E. AESCULIFOLIUM*, antheris anfractuosis, foliolis argutè serratis, caudice inermi. § in Novæ-Hispaniæ littore prope Campechê. *Bombax axillare* fl.

mex. ic. ined? Foliola 7-8 lanceolato-oblonga acuminata et mucronata glabra. Petala rosea? obtusa extus hirsuta staminibus æqualia. Flores ex ic. fl. mex. axillares subsolitarii, pedicellis brevissimis crassis.

XII. CHORISIA. *Kunth diss. malv. p. 6. nov. gen. am. 5. p. 295.*

Calyx campanulatus basi 3-bracteatus persistens 3-5-lobus. Petala 5 elongata. Tubus stamineus duplex, interior tenuis teres apice antheras 10 geminatas extus gerens, exterior brevis prioris basi supra adnatus 10-lobulatus sterilis. Ovarium sessile ovatum 5-loc? Stylus 1 filiformis stam. superans. Stigma capitellatum. Capsula 5-valv. 1-loc.? polysperma. Semina lanata. — Arbores aculeatæ. Folia palmata. Flores magni, petalis subvillosis.

1. *C. insignis* (H. B. et K. nov. gen. am. 5. p. 297. t. 485. f. 1.) petalis oblongo-spathulatis margine planis apice emarginatis. § ad ripam flum. Amazonum. Foliola 5 obovato-oblonga, acuminata, apicem versus obsolete undulato-crenata, glabra, subtus glaucescentia.
2. *C. crispiflora* (H. B. et K. l. c. t. 485. f. 2) petalis linearibus obtusis margine undulato-crispis. § in Brasiliæ sylv. *C. ventricosa* Nees et Mart. nov. act. bonn. xi. p. 101. t. g. *Bombax ventricosa* Arrud. in Kost. trav. p. 489. Neuw. bras. reis. 2. p. 247. ex N. et M.

XIII. DURIO. *Lin. syst. 698. non Adans.*

Calyx nudus obtusè 5-lobus. Petala 5 calyce minora. Stam. co, pentadelphæ, antheris anfractuosus. Ovarium lepidotum. Stylus filiformis. Stigma subrotundum. Fructus subrotundus, muricatus, 5-fariam dehiscentis, 5-locul., loculis intus pulposis 4-5-spermis. — *Eriodeudro* affinis ex antheris videtur. Sem. ign.

1. *D. zibethinus* (Lin. l. cit. Lam. ill. t. 641.) § in Indiâ orientali. Rumph. amb. 1. p. 99. t. 29. Folia Cerasi supernè virentia glabra, subtus lepidota fusco-cinerea. Fructus magnitudine capitis à Zibethis voraciter appetitus. (v. s. in) h. Mus. Par.)

XIV. OCHROMA. *Swartz. act. holm. 1792. p. 148. t. 6.*

Calyx basi tubulosus subinfundibuliformis vix 5-fidus, lobis planis 5 rotundatis, 2 acutiusculis. Pet. 5 calyce majora. Antheræ anfractuosæ. Stigmata 5. Capsulæ valvulæ intus lanâ sericeâ vestitæ. Semina oblonga.

1. *O. LAGOPUS* (Swartz. fl. 2. p. 1144. t. 23.) foliis cordatis 5-angulato-sublobatis denticulatis subtus pubescentibus. § in montibus Jamaicæ, Hispaniolæ Porto-Ricci, Mexici cal. — *Bombax pyramidale*. Cav. diss. 5. p. 294. t. 155. (v. s.)
2. *O. TOMENTOSA* (Willd. enum. 695.) foliis cordatis subtrilobis repandis subtomentosis. § in Americâ merid. Humb. et Bonpl.

XV. CHEIROSTEMON. *Humb. et Bonpl. pl. eq. 1. t. 44. nov. gen. am. 5. p. 302.*
— *Chiranthodendron*. *Larr. diss. cum ic.*

Calyx subcampanulatus 5-partitus extus ad basin 3-bracteatus, sepalis deciduis crassis intus coloratis ad basin foveolatis, æstivatione quincunciali. Petala 0. Stam. filamenta in tubum apice 5-fidum connata, lobis secundis apice exserto mucronatis dorso antheras 2 lineares longitudinaliter dehiscentes gerentibus. Stylus 1. Stigma acutum. Capsula oblonga, 5-ang. 5-loc. 5-valvis, valvis medio septiferis, septis villosis. Semina in quoque loculo 15-18 ovoidea carunculata. Albumen carnosum. Embryo axilis. Cotyledones planæ. — Arbor foliis 5-7-lobatis palminerviis.

1. *C. PLATANOIDES* (H. et B. l. c.) § in Novâ-Hispaniâ prope Toluca. Tiles. act. petrop. 5. p. 321. t. 9. Fisch. p. 531. — Hern. mex. 383. f. 1. et 459. f. 2. (v. v. et s.)

Ordo XXVI. BYTTNERIACEÆ. *Brown cong. Kunth diss. malv. p. 6. nov. gen. am. 5. p. 309.* — Sterculiaceæ seu Hermanniaceæ *Vent. Juss.*

Calyx nunc nudus, nunc involucrio cinctus, sepalis 5 plus minus basi coalitis per æstivationem valvatis constans. Petala 5 hypogyna, sepalis alterna, per æstivationem convoluta, formâ in variis varia, rarius inæqualia aut nulla. Stamina numero petalorum aut sepalorum æqualia dupla tripla multipliciave, filamentis plus minus et variis modis monadelphis, nonnullis interdum sterilibus, antheris posticis bilocularibus. Carpella 5 rarissimè 3 distincta aut in ovarium unicum coalita. Styli tot quot carpella distincti aut coaliti. Albumen oleosum aut carnosum, rarius nullum. Embryo rectus, radicula inferâ, cotyledonibus nunc foliaceis planis aut plicatis aut circa plumulam convolutis, nunc (in seminibus exalbuminosis) crassissimis. — An tribus sequentes ut ordines habendæ? An potius Byttneriaceæ in unicum ordinem phalangemve conjungendæ cum Malvaceis Tiliaceis et Elæocarpeis quibuscum conveniunt habitu, calyce valvato, petalis convolutis? à Malvaceis distinguuntur tantum antheris bilocularibus et à Tiliaceis et Elæocarpeis filamentis monadelphis.

Trib. I. STERCULIÆ. — Sterculiaceæ *Kunth l. c.*

Flores abortu sæpè 1-sexuales. Calyx nudus 5-lobus deciduus. Petala 0. Stam. 5–10–15–20 basi breviter monadelpha. Ovarium sæpius pedicellatum, carpellis 5 distinctis constans. Albumen oleosum aut nullum. Embryo erectus, cotyledonibus in seminibus albuminosis planis foliaceis, in seminibus exalbuminosis crassissimis inæqualibus, radicula ovata brevi. — Tribus habitu ad Bombaceas accedens, characteribus verò Byttneriaceis potius referenda. An ordo proprius? An characterem reformato huc forsân associandæ Niota, Samadera?

I. STERCULIA. *Lin. gen. n. 1086.*

Calyx 5-lobus subcoriaceus. Stam. monadelpha in urceolum brevem sessilem stipitatumve disposita; antheræ 10, 15, 20, uni aut biseriales solitariæ aut ternatim adregatæ. Ovarium stipitatum aut sessile. Carpella follicularia 5 distincta 1-loc., 1-aut-∞-sperma, rimâ supernè latè dehiscentia. Semina albumine oleoso, cotyledonibus planis foliaceis æqualibus. — Genus aut dividendum aut in sectiones naturales ex fructificatione ordinandum, sed ob species plurimas non satis notas hic adhuc artificialiter digestum.

§. 1. *Foliis ovatis oblongisve integris aut rarissimè trilobis.*

1. S. LANCEOLATA (Cav. diss. 5: p. 287. t. 144. f. 1.) foliis integerrimis lanceo-

latis, carpellis oblongis oligospermis. 5 in Chinâ. Carp. chermesina. Semina nigra. †.

2. *S. BALANGHAS* (Lin. sp. 1438. excl. syn. Rumph.) foliis ovato-lanceolatis integris glabris, calycis laciniis linearibus apice cohærentibus, carpellis obovatis polyspermis. 5 in Malabariâ. Cavalam Rheed. mal. 1. t. 49. *S. Balanghas* Cav. diss. 5. p. 286. t. 143. Lois. herb. amat. t. 843.
3. *S. NOBILIS* (Smith in Rees cycl. n. 4.) foliis ovato-oblongis integris glabris, calycis laciniis linearibus apice cohærentibus, carpellis ovatis mucronatis 1-2-spermis. 5 in Indiâ orientali. *S. monosperma* Vent. malm. t. 91. *S. Balanghas* Ait. h. Kew. ed. 2. v. 5. p. 338. *Southwellia nobilis*. Sal. par. lond. t. 69. excl. syn. Flores viridi-flavescentes odore Vanillæ.
4. *S. ACUMINATA* (Beauv. fl. ow. 1. t. 24.) foliis oblongis acuminatis integerrimis glabris longè petiolatis, floribus axillaribus paniculatis, antheris biserialibus sessilibus, carpellis 1 spermis. 5 in Africâ æquinoctiali. Cola. Bauh. pin. 507. Semina ab Africanis valdè æstimata, quia iis manducatis aquæ etiam foetidæ gustu tolerabiles evadunt. (v. s.)
5. *S. LONGIFOLIA* (Vent. malm. n. 91. adn.) foliis ovato-oblongis glabris integerrimis, floribus paniculatis terminalibus, calycinis laciniis erectis intus hirsutis. 5 in Indiâ orientali Fruct. ign.
6. *S. RUBIGINOSA* (Vent. malm. n. 91. adn.) foliis lanceolato-oblongis supra glabris subtus tomentosis rubiginosis, carpellis acuminatis polyspermis intus rugosis et nudis. 5 in Javâ. Eadem ex cl. Smith ac *S. Balanghas*.
7. *S. GRANDIFLORA* (Vent. malm. n. 91. adn.) foliis ovatis acuminatis integris glabris, laciniis calycinis patentibus, staminum urceolo subsessili, stylis? 5 reflexis. 5 in Indiâ orient. et ins. Mauritiî.
8. *S. MITIDA* (Vent. l. cit.) foliis lanceolato-oblongis acuminatis integris glabris, calycinis laciniis patentibus, urceolo subsessili. 5 in Africâ. Forsan dioica?
9. *S. FRONDOSA* (Rich. act. soc. h. nat. par. p. 111.) foliis oblongo-ovatis obtusissimis subrepandis glabris lucidis in ramorum summitate congestis, paniculis axillaribus longè pedunculatis. 5 in Guianâ Gallicâ.
10. *S. URCEOLATA* (Smith in Rees. cycl. n. 3.) foliis ellipticis oblongis acutis subtus pallidis et tenuiter velutinis, paniculis contractis paucifloris vix petiolo longioribus, calyce lageæformi margine villosa. 5 in ins. Homineâ prope Amboinam. Clompanus minor Rumph. amb. 3. p. 169. t. 107. Carpella polysperma.
11. *S. COCCINEA* (Roxb. h. calc. p. 50.) foliis oblongis lanceolatis obtusè acuminatis glabris, racemis axillaribus et lateralibus nutantibus, calycinis laciniis linearibus patentibus. 5 in Indiâ orientali ad Pulo-Pinang. Mal. misc. 1. n. 1. p. 20. Folliculi coccinei.
12. *S. ANGUSTIFOLIA* (Roxb. h. calc. p. 50.) foliis lanceolatis supernè latioribus acuminatis subtus villosis, racemis extrâ-axillaribus nutantibus, laciniis calycinis linearibus apice connexis. 5 in Napauliâ. Roxb.; in Pulo-Pinang. Mal. misc. 1. n. 1. p. 21.
13. *S. GUTTATA* (Roxb. h. calc. p. 50.) foliis ovatis obtusis crassis glabris integris, racemis subfasciculatis patentibus folio longioribus, calycibus intus lanuginosis patentibus. 5 in Malabariæ montanis. Ramena-Pou-Maram Rheed. mal. 4. t. 61.
14. *S. OBLONGIFOLIA* (fl. mex. ic. ined.) foliis oblongis acutis basi attenuatis integerrimis glabris, floribus laxè racemosis, calycibus patentibus, genitalibus subsessilibus, carpellis subdrupaceis dispermis. 5 in Mexico.
15. *S. HETEROPHYLLA* (Beauv. fl. ow. 1. t. 40.) foliis oblongis basi attenuatis longè petiolatis integris trilobisque irregulariter dentatis, floribus paniculatis terminalibus. 5 in regno Owariensi.

16. *S. IVIRA* (Swartz. fl. ind. occ. 2. p. 1160.) foliis ovatis glabris apice acuminatis integris rariùs, trilobis, floribus paniculatis hermaphroditis decandris, carpellis basi crinitis. ζ in sylvis Amer. calid. *Ivira pruriens* Aubl. guian. t. 279. I. crinita Cav. diss. 5. t. 162. *Carpella* basi pilis setosis rigidis tecta, intùs setis urentibus minutis rufis vestita.
17. *S. IMBERBIS*, foliis ovatis glabris basi obtusissimis subcordatis acuminatis integris, carpellis basi nudis pedicellatis. ζ in Cayennâ. *S. Ivira* Richard in Pers. ench. n. 10. (v. s.)

§. 2. *Foliis cordatis integris aut subtrilobis.*

18. *S. MACROPHYLLA* (Vent. malm. n. 91. adn.) foliis cordato-subrotundis subtùs tomentosis integris crassis, carpellis ovatis intùs glaberrimis dispersis. ζ in Indiâ orientali.
19. *S. POPULIFOLIA*, foliis cordato-subrotundis obtusis integris utrinquè glabris submembranaceis, carpellis ovatis intùs glaberrimis tetraspermis. ζ in ins. Timor. Flor. ign. (v. s. ex Mus. Par.)
- β . *acutiuscula*, foliis cordatis acutiusculis utrinquè glabris submembranaceis, racemis axillaribus lateralibus petiolo brevioribus, calycibus 5-partitis elongatis velutinis. ζ in Indiâ or. *S. populifolia* Roxb. cat. calc. p. 50? (v. s. sine fruct. ex Roxb.)
20. *S. CORDIFOLIA* (Cav. diss. 5. t. 144. f. 2.) foliis lato-cordatis subacuminatis utrinquè glabris interdum obtusè trilobis, carpellis acuminatis tomentosis intùs setosis polyspermis. ζ ad Senegal.

§. 3. *Foliis cordatis quinquelobis.*

21. *S. PLATANIFOLIA* (Lin. suppl. 423.) foliis palmato-5-lobis, calycibus rotatis reflexis. ζ n Japoniâ, Chinâ. Cav. diss. 5. t. 145. *Hibiscus simplex* Lin. sp. 977. *Firmiana* Mars. act. pat. 1. t. 1. 2. *Culhamia* Forsk. descr. 96. syn. dub. ex Salisb. (v. v.)
22. *S. COLORATA* (Roxb. cor. 1. p. 26. t. 25.) foliis palmato-5-lobis, lobis acuminatis, calycibus cylindraceo-clavatis, carpellis oblongis glabris. ζ in montibus Indiæ orientalis. Calyces et *Carpella* rubro-coccinei.
23. *S. URENS* (Roxb. cor. 1. p. 25. t. 24.) foliis palmato-5-lobis, lobis acuminatis, calycibus campanulatis, carpellis ovatis hispidis. ζ in montibus Indiæ orient. Cal. et carp. virescunt.
24. *S. VILLOSA* (Roxb. cat. calc. p. 50.) foliis 5-lobis acutis dentatis tomentoso-velutinis, calycibus 5-partitis patentibus, carpellis coriaceis pube stellatâ scabris. ζ in Coromandeliâ. Smith in Rees. cycl. n. 16.*
25. *S. HELICTERES* (Pers. ench. 2. p. 240.) foliis semi-5-lobis subtùs subvillosis, lobis ovato-subrotundis acutis, calycibus campanulatis, laciniis patentibus extùs velutinis. ζ in sylvis Carthagenæ. *Helicteres apetala* Jacq. am. 238. t. 181. f. 97. Stam. 14-15, *Carpella* divaricata. (Pers.)
26. *S. PUNCTATA* (fl. mex. ic. ined.) foliis palmato-5-fidis, lobis ovatis acuminatis, ramis petiolisque punctatis, calycibus subrotatis, carpellis 4-spermis intùs setosis. ζ in Novâ-Hispaniâ.

§. 4. *Foliis palmatim compositis.*

27. *S. FORTIDA* (Lin. sp. 1431.) foliis compositis, foliolis 7-9 peltatim dispositis oblongo-acuminatis. ζ in Indiâ orient. Cav. diss. 5. t. 141. Sonn. voy. t. 132. *Clompanus major* Rumph. amb. 3. t. 107. *Antheræ* 15. *Carpella* polysperma. (v. s.)

II. *TRIPHACA*. Lour. coch. 2. p. 708.

Fl. monoïci. Calyx 5-fidus. Pet. o. σ Stam. 15 monadelphia? φ Stylus 1. *Carpella* 3, leguminiformia. †:

1. *T. AFRICANA* (Lour. l. c.) ζ in Africæ littore Mozambico. Folia longè pe-

tiolata cordata integerrima. Cymæ laterales et terminales. Flores latei. An Sterculiæ species?

III. HERITIERA. *Ait. h. Kew.* 3. p. 546. — *Balanopteris. Gaertn. fr.* 2. p. 94. — *Samandura. Lin. fl. zeyl. n.* 433.

Calyx 5-dentatus. ♂ Stam. 5-10, filamentis in tubum coalitis, antheris apice tubuli sessilibus. ♀ Antheræ 10 sessiles, 2 inter singula carpella. Carpella 5 monostyla pauci-ovulata demum drupaceo-coriacea carinato-alata, indehiscentia, abortu 1-sperma. Albumen o. Embryo crassissimus, cotyledonibus carnosus inæqualibus, radicula ovato-acuminatâ suturæ carpelli obversâ, plumulâ diphyllâ. — Arbores, foliis simplicibus alternis lepidotis integris, paniculis axillaribus.

1. *H. LITTORALIS* (Ait. l. c.) foliis ovalibus simpliciter venosis, carpellis supernè nervo longitudinaliter notatis. § in ins. Philippinis, Moluccanis, Indiciis. — Rheed. mal. 6. t. 21. Rumph. amb. 3. t. 63. — *B. Tothila Gaertn. fr.* 2. t. 99. (v. s.)

2. *H. FOMES* (Willd. sp. 4. p. 971.) foliis subtus reticulato-venosis. § ad ripas in regno Avæ Indiæ. — Symes it. ed. gall. 3. p. 100. t. 28.

3. *H. MINOR* (Lam. dict. 3. p. 229.) carpellis supernè sulco longitudinali exaratis. § in ins. Mauritii ubi verosim. culta. — *B. minor Gaertn. fr.* 2. t. 98. f. 2. An forsâ eadem ac *H. Fomes*?

Trib. II. *BYTTNERIEÆ*. — *Büttneriaceæ* veræ *Kunth l. c.*

Petala basi sæpissimè concavo-fornicata apice in ligulam expansa, Stam. 10-30 et ultrâ. Filamenta 5 petalis opposita sterilia ligulata, alia fertilia alterna solitaria ternatimve pentadelfha aut rariùs monanthera. Ovarium 5-locul. loculis sæpiùs 2-spermis. Semina nunc exalbuminosa cotyledonibus crassis, nunc albuminosa cotyledonibus foliaceis planis convolutisve.

IV. *THEOBROMA. Juss. gen.* 276. *Kunth nov. gen. am.* 5. p. 316. — *Cacao. Tourn. inst. t.* 444.

Calyx 5-sepalus. Petala 5 basi fornicata, in ligulam spathulatam producta. Stamina uterolus cornicula 5 exserens et inter ea filamenta 5 biantherifera petalis opposita. Stylus filiformis. Stigma 5-partitum. Caps. 5-locul. evalvis. Semina in pulpâ butyraceâ nidulantia. Albumen o. Cotyledones crassæ, oleosæ, corrugatæ.

1. *T. CACAO* (Lin. spec. 1100.) foliis integerrimis ovato-oblongis acuminatis utrinquè glaberrimis concoloribus. § in Americâ merid. alt. 0-200 hexap. *Cacao sativa Lam. dict.* 1. p. 553. ill. t. 653. *C. Theobroma Tuss. ant.* t. 13. *C. minus Gaertn. fruct.* 2. p. 190. t. 122. *Cat. car.* 3. t. 6. (v. s. sine fl.)

2. *T. GUIANENSIS* (Willd. spec. 3. p. 1422.) foliis acuminatis repando-dentatis subtus tomentosis. § in sylvis paludosis Guianæ. *Cacao Guianensis Aubl. guian.* 2. p. 683. t. 275. — *Fractus ovato-quinquangularis tomentosus rufescens.*

3. *T. BICOLOR* (Humb. et Bonpl. pl. eq. 1. p. 104. t. 30.) foliis oblongis obliquè cordatis subtus albicantibus 7-nerviis. § in Novæ-Granatæ vallibus calidis. *Fructus drupaceus ovalis indehiscens variè excavatus sericeus. Semina cum T. Cacao ab incolis miscentur.* (v. s.)

4. *T. ANGUSTIFOLIA* (fl. mex. ic. ined.) foliis oblongis utrinquè attenuatis, apice acuminatis basi trinerviis subtus pallidis. § in Mexico. Flores sordidè armeniaci. Petalorum limbus oblongo-stipitatus. Fila sterilia obovato-oblonga petalis ferè majora. Fructus ovatus.

5. *T. OVATIFOLIA* (fl. mex. ic. ined.) foliis ovatis integerrimis basi trinerviis peltato-subcordatis apice obtusis subtus cano-tomentosis. 5 in Mexico. Flores parvi. Sepala acuminata. Fructus ovoideus costis elevatis rugosus.

V. *A. ROMA*. Lin. fl. suppl. 341. Sal. par. 102. Kunth nov. gen. 5. p. 318.

Cal. 5-partitus. Pet. 5 unguibus basi dilatatis saccatis. Stamina urceolus 10-fidus, laciniis 5 triantheriferis, 5 alternis petaloideis. Capsula 5-loc. 5-alata polysperma. Semina arillata. Albumen carnosum. Cotyl. foliaceæ transversæ flexuosæ.

1. *A. AUGUSTA* (Lin. suppl. 341.) ramis molliter velutinis lævibus, foliis adultis subtus glabris aut pube simplicissima, capsulæ alis apice truncatis, angulo exteriori acutiusculo. 5 in Indiâ orient. *A. angusta* et *A. Wheleri* Willd. sp. 3. p. 1424 et 1425. *A. fastuosum* Jacq. vind. 3. p. 3. t. 1. Gaertn. fr. 1. p. 306. t. 64. Folia. inf. palmatifida 5-7-nervia, super. ovato-lanceolata indivisa. (v. s.)
2. *A. FASTUOSA* (Brown in h. Kew. ed. 2. v. 4. p. 409.) ramis muricatis, foliis adultis pube stellari et simplici sparsâ scabris, capsulæ alis apice subtruncatis, angulo exteriori elongato acuminato. 5 in Timor et Novâ-Hollandiâ. *A. fastuosum* Gaertn. fr. 1. p. 307. t. 64. Sal. par. lond. t. 102. Folia infer. acutè 5-loba, sup. ovata subcordata indivisa. (v. s.)
3. *A. MOLLIS*, ramis muricatis, foliis adultis pube brevissima molli confertâ et setis stellatis velutinis, capsulæ alis... 5 in Moluccis. Fol. infer. cordata subrotunda vix sublobata, super. etiam cordata brevissimè petiolata acuminata serrata, omnia tactu molliter velutina. An prioris varietas? (v. s.)

VI. *GUAZUMA*. Plum. gen. 36. t. 18. Juss. Cav. Pers. — *Bubroma*. Schreb. gen. n. 1216.

Cal. 5-sepalus aut sepalis diversè connatis 2-3-partitus. Pet. 5 apice bicornia. Stam. filamenta vix basi monadelphæ, tubo brevissimo exserto in lobos 5 steriles et in fila 5 lobis alternâ apice trifida triantherifera. Styli 5 conniventes. Capsula lignosa tuberculata evalvis 5-locularis, decuplici serie foraminulorum pertusa, polysperma. Sem. ovata. Cotyledones plicatæ ex Kunth 5. p. 320.

1. *G. UMBROSA* (Lam. dict. 3. p. 52.) foliis adultis utrinquè glabris. 5 in ins. Caribæis. *Theobroma Guazuma* Lin. sp. 1100. Plum. ed. Burm. t. 144 sed folia nimis cordata et racemi nimis elongati. Pluk. alm. t. 77. f. 5. *Bubroma Guazuma* et forsân *B. Invira* Willd. enum. 806. Folia in meis specim. basi ovata aut oblonga apice acuminata inæqualiter dentata, juniora vix in nervis pube stellatâ tomentosa. (v. s.)
2. *G. TOMENTOSA* (H. B. et Kunth nov. gen. am. 5. p. 320.) foliis suprâ pilosisculis, subtus tenuiter tomentosis albidis. 5 An à priore aut à seq. satis differt?

a. *Monpoxensis* (H. B. et Kunth l. c.) corymbis folio duplò longioribus, calyce bipartito. — Circa Monpox. ad fl. Magdalenæ et in Novâ-Andalsiâ.

æ? *Cumanensis* (H. B. et Kunth l. c.) paniculis folio duplò brevioribus, calyce tripartito. — prope Cumanam. An species propriâ?

3. *G. POLYBOTRYA* (Cav. ic. 3. p. 51. t. 299.) foliis subtus pube stellatâ velutino-tomentosis, supernè junioribus pubescentibus, adultis glabris. 5 in Novâ-Hispaniâ et Sancto-Domingo, forsân in Brasiliâ si *Bubroma polybotryum* Willd. enum. 806 huc ritè relatum. Guacimo Hern. mex. 40. f. 1. (v. s.)

VII. *GLOSSOSTEMON*. Desf. mem. mus. 3. p. 238. t. 2. H. B. et Kunth nov. gen. am. 5. p. 311. in adnot.

Cal. 5-partitus. Pet. 5 acumine filiformi terminata. Stam. 25-35 basi

monadelphæ, supernè in phalanges 5 distincta, fil. medio cujusque phalangis sterili ligulæformi. Ovarium 1, 5 locul. loculis co-spermis. Stylus 1. Stigm. 5. Capsula? setigera. — Genus affine hinc Sparmanniæ, illinc Byttneriaceis.

1. *G. BRUGUIERI*. ? in Persiâ prope Bagdad. Folia petiolata ovato-rotunda sublobata dentata hispida, pilis stellatis. Flores corymbosi, rosei. (v. s.)

VIII. *COMMERSIONIA*. Forst. gen. 43. t. 22. H. B. et Kunth nov. gen. am. 5. p. 311. in adn.

Cal. 5-partitus petaloideus persistens. Pet. 5 basi dilatato saccata apice longè mucronata. Stam. 10 filamentis basi connatis, alternis sterilibus petaloideis apice stellatim patentibus, alt. fertilibus subulatis tricornibus. Antheræ utrinquè rimâ later. dehisc. Styli 5. Ovarium 5-loc. loculis 3-4-ovulatis. Capsula 3 5 valvis echinata, setis villosis. — Inflorescentia cymosa. An à Byttneriâ satis differt?

1. *C. ECHINATA* (Forst. l. c.) caule arboreo, foliis ovato-lanceolatis supernè glabrinusculis subtùs ænescentibus. 5 in Molluccis, Novâ-Caledoniâ et insul. Societatis et Amicorum. — Rumph. amb. 3. t. 119. Variat foliis basi cordatis ovatis aut inæqualibus. (v. s.)

2. *C. PLATYPHYLLA* (Andr. bot. rep. n. 603. et t. 519. sub *C. echinata*) caule frutescente, foliis ovato-acuminatis supernè hispidulis subtùs hirsutis. 5 in Molluccis. Sims bot. mag. t. 519.

β. *Lechenaultii*. — Specimen habeo ex hort. Calc. *C. platyphyllæ* simile sed folia æqualiter serrata supernè glabra, paniculæ axillares foliosæ, ramuli floriferi foliis oppositi! An species? An status anomalus? (v. s.)

3. *C. GAUDICHAUDI* (Gay. in litt.) caule fruticoso, foliis profundè cordatis inæquilateralis, supernè hispidis, subtùs tomentosis. 5 in Novâ-Hollandiâ orâ orientali à cl. Gaudichaud lecta.

IX. *BYTTNERIA*. Læfl. itin. 313. Lin. gen. n. 268. non Duham.

Omnia Commersoniæ sed filamenta fertilia simplicia nec trifurca et capsulæ aculei pilis nullis paucissimisve ornati.

§. 1. *Species inermes* (Commersoniis affines.) *RULINGIA*. Brown.

1. *B. DASYPHYLLA* (Gay. in litt.) foliis ovato-lanceolatis inæqualiter serratis utrinquè hirsutis, petalorum mucrone calycem superante. 5 in terrâ Van-Diemen. Commersonia diphylla Andr. bot. rep. t. 603.

2. *B. PANNOSA*, foliis ovato-lanceolatis inæqualiter dentato-serratis supernè pubescentibus subtùs hirsutis, petalorum mucrone segmentis calycinis brevior. 5 in Novâ-Hollandiâ? Rulingia pannosa. Brown in bot. mag. t. 2191. Buttneria inodora. Gay. ined. Lasiopetalum tomentosum. Cels. (v. s.)

3. *B. HERMANNIÆFOLIA* (Gay! in litt.) foliis ovatis inæqualiter dentato-crenatis rugosulis subtùs tomentosis, petalorum mucrone segmentis calycinis brevior. 5 in Novâ-Hollandiâ ad Port-Jackson. (v. s.)

4. *B. MACROPHYLLA* (H. B. K. nov. gen. am. 5. p. 315.) foliis cordatis ovato-subrotundis obtusis serrato-crenatis basi maculâ rasâ notatis ramisque pubescentibus, pedunculis multifloris axillaribus subternis. 4 prope Honda Novo-Granatensium.

5. *B. HERBACEA* (Roxb. cor. 1. t. 29.) foliis cordatis acuminatis dentatis glandulosis, petalorum mucronibus brevissimis reflexis. 4 in Coromandelæ montibus Roem. et Schult. syst. 5. p. 469. excl. Willd. syn.

6. *B. GRANDIFOLIA*, foliis cordatis 7-nerviis obtusis submucronatis ovatis integris utrinquè glabris, petalorum mucronibus brevissimis. 4 in Indiâ orient.

Recepi à cl. Wallich sub nomine *B. asperæ* Roxb. quod in libris evulgatis non video. Fructus ut in aliis echinatus. Folia lævia. (v. s.)

7. *B. CATALPÆFOLIA* (Jacq. h. schoenb. 1. t. 46.) foliis cordatis integerrimis longè acuminatis glabris, caule scandente. ζ ad Caracas. Flores albi.

§. 2. *Species in caulibus ramis petiolis nervisque aculeatæ.*

8. *B. HIRSUTA* (fl. per. 3. p. 10.) foliis cordatis acutis crenatis subtus hirsutis, costâ petiolisque aculeatis, pedunculis congestis compositis. ζ in Andium montibus. Flores umbellati involucrati.

9. *B. MOLLIS* (H. B. & K. nov. gen. am. 5. p. 314. t. 481. a et b.) foliis cordatis acuminatis crenatis inermibus utrinquè molliter tomentosis basi maculâ rasâ notatis, ramis aculeatis villosis-tomentosis, umbellis 7-11-floris axillaribus et oppositifoliis solitariis ternis. ζ ? prope Santa Fe de Bogota.

10. *B. CORDATA* (Lam. dict. 1. p. 523.) foliis cordatis acuminatis serratis subtus glandulosis pubescentibus, petiolis inermibus, caulibus aculeatis, pedunculis subternis inæqualibus 7-floris. ζ in Peruviam prope Lima ad Chancay sepibus. Cav. diss. 5. p. 291. t. 150. Rœm. et Schult. syst. 1. p. 469. excl. Willd. syn.

11. *B. SULCATA* (fl. per. 3. p. 10.) foliis cordato-ovatis serratis pubescentibus costâ petiolisque aculeatis, pedunculis geminis 3-5-floris. ζ in Peruviam præruptis calidis. Caulis 5-sulcatus.

12. *B. LANCEOLATA* (fl. mex. ic. ined.) foliis ovato-lanceolatis subcordatis rari-ter serratis acuminatis petiolisque inermibus, caule-aculeato scandente anguloso, pedunculis ternis 5-7-floris. ζ in Mexici montibus. Petala alba setis purpureis.

13. *B. OVATA* (Lam. dict. 1. p. 522.) foliis ovatis dentato-serratis glabris, petiolis inermibus, ramis 5-angularibus aculeatis, caule erecto. ζ in Peru. Cav. diss. 5. p. 291. t. 149. f. 1. (v. s.)

14. *B. MICROPHYLLA* (Lin. mant. 209.) foliis ellipticis integerrimis apiceve subdentatis emarginatis, aculeis stipularibus. ζ in Americâ Lin., in Sancto-Domingo Jacq. Cav. diss. 5. p. 292. t. 148. f. 2. Rami flexuosi.

15. *B. TERETICAULIS* (Lam. dict. 1. p. 523.) foliis lanceolatis acuminatis integerrimis, rachi petiolis cauleque tereti aculeatis. ζ in Peru. Cav. diss. 5. p. 292. t. 149. f. 2.

16. *B. SCABRA* (Lin. syst. 197.) foliis lanceolatis dentatis basi subhastatis, rachi petiolis cauleque angulato subsarmentoso aculeatis. ζ in Americâ merid. Cav. diss. 5. p. 291. t. 148. f. 1. Aubl. guian. t. 96. (v. s.)

17. *B. CARTHAGENENSIS* (Jacq. amer. ed. pict. p. 41.) foliis caulinis ovatis, ramis ovato-lanceolatis obtusè acuminatis integris, costâ petiolis ramisque aculeatis. ζ ad Carthagensem amer. sylvarum margines. Bertero. *B. aculeata* Jacq. amer. 76. *Chætæa aculeata* Jacq. enum. 17. (v. s. in h. Balb.)

† *Species non satis notæ.*

18. *B. ACUMINATA* (Bred. ex Willd. rel. in Rœm. et Schult. syst. 5. p. 470.) foliis ovatis acuminatis inermibus, petiolis subaculeatis, caule tereti aculeato. ζ ad Caracas.

19. *B. SALICIFOLIA* (Humb. et Bonpl. ex Willd. rel. in Rœm. et Schult. syst. 5. p. 470.) foliis lanceolatis glabris, caule angulato petiolis rachique foliorum aculeatis. ζ in Cumana.

20. *B. CORYLIIFOLIA* (Humb. et Bonpl. l. c.) foliis oblongis subcordatis serratis acutis, caule aculeato. ζ in Americâ merid.

X. *AYENIA*. Lin. gen. n. 1020.

Calyx 5-partitus. Petala longè unguiculata arcuata supernè latiora glandulis pedicellatis superata. Staminum urceolus 10-15-dentatus,

dentibus 5-10 obtusis sterilibus, 5 alternis monantheris. Stylus 1. Stigma 5-gonum. Carpella 5 bivalvia 1-sperma conniventia in fructum subglobosum echinatum. Albumen o. Cotyledones foliaceæ convolutæ.

1. *A. PUSILLA* (Lin. sp. 1554.) foliis ovatis glabrinusculis argutè serratis, caulibus prostratis. ① 2 in ins. Caribæis et Peruvîâ. Cav. diss. 5. p. 289. t. 147. Sloan. hist. t. 132. f. 2. Dayenia. Mill. ic. t. 118. (v. s.)
2. *A. LÆVIGATA* (Swartz fl. ind. occ. 2 p. 1131.) foliis ovatis glaberrimis integris, urceolo in dentes 10 præter stamina exserto. 5 in Jamaicæ frutetis rariùs. (v. s.)
3. *A. TOMENTOSA* (Lin. sp. 1354.) foliis ovato-subrotundis tomentosis, urceolo in dentes 5 præter stamina exserto. — in Cumanâ Lœfl. it. 200. n. 3.
4. *A. SINEFOLIA* (Lœfl. it. 257.) foliis ovato-oblongis duplicato-serratis subtus tomentosis. 5 in Americâ calid. Cum priorè à Linnæo conjuncta. 7.
5. *A. MAGNA* (Lin. sp. 1354.) foliis cordatis ovato-acuminatis serratis pubescentibus, pedunculis multifloris petiolo longioribus. 5 ad Cumanam. (Lœfl. it. 199. n. 1.) et in Mexico (*A. cardiopetala* fl. mex. ic. ined.)
6. *A. CORDIFOLIA* (fl. mex. ic. ined.) foliis cordatis serratis pubescentibus, pedicellis plurimis aggregatis petiolo brevioribus. ① in Mexici montosis.

XI? *KLEINHOVIA*. Lin. gen. n. 1024. Gaertn. fruct. 2. p. 261. t. 137. H. B. et Kunth nov. gen. am. 5. p. 313.

Calyx 5-partitus. Pet. 5 quorum 1 ceteris longius apice abscissum. Stam. filamenta basi in urceolum monadelpha, supernè pentadelpha, adelphiis dorso 3 antheriferis petalis oppositis. Ovarium stipitatum, carpellis 5 quadri-ovulatis concretis constans. Stylus 1. Stigma subcrenatum. Capsula inflata turbinata 5-angul. 5-locul. 5-sperma. Semina subrotunda. Cotyledones circa plumulam spiralerè convolutæ. — Ex hoc charact. ad *Ayeniam* accedens.

1. *K. HOSPITA* (Lin. sp. 1365.) 5 in Molluccis. — Rumph. amb. 3. t. 113. Cav. diss. 5. p. 188. t. 146. Arbor glabra, foliis cordatis latis acuminatis integris, floribus racemosis purpureis. (v. s.)

Trib. III. *LASIOPETALÆ*. Gay. diss. p. 8. et Mém. du Mus. 7. p. 431. H. B. Kunth. l. c.

Calyx 5-partitus petaloïdeus persistens aut marcescens. Pet. minuta, squamuliformia, rariùs o. Stam. filam. subulata basi connata, nunc 5 petalis opposita, nunc 10 alternè sterilia et fertilia. Antheræ filam. incumbentes lobis contiguïs. Ovarium 3-5-loc., loculis 2-8-ovulatis. Carpella 5 bivalvia sæpiùs arcuè connata aut sublibera. Semina basi strophiolata. Albumen carnosum. Embryo erectus. Cotyledones planæ foliaceæ.

XII. SERINGIA. Gay. diss. p. 12.

Cal. marcescens. Pet. o. Filam. 10 alterna sterilia. Antheræ rimis dorsalibus dehiscentes. Carpella 5 bivalvia 2-3-sperma. Seminum strophiola crenata.

1. *S. PLATYPHYLLA* (Gay! l. c. p. 13. t. 1. 2.) 3 in Novæ-Hollandiæ orâ orientali. Lasioptalum arborescens Ait. h. Kew. ed. 2. v. 2. p. 56. Pedunculi oppositifolii cymam multifloram confertam gerentes. Stipulæ parvulæ deciduæ. Folia ovato-lanc. grossè dentata. (v. s.)

XIII. LASIOPETALUM. *Smith trans. lin. soc. 4. p. 216.*

Cal. persistens. Pet. 5 glanduliformia. Filam. 5 libera. Antheræ intus poris 2 dehiscentes. Ovarium 1, 3-locul., loculis biovulatis. Capsula 3-valvis, septis valvaribus, dehisc.-loculicida. Sem. strophiliola laciniata. — Stipulæ o. Folia lineari-lanceolata integerrima. Inflor. cymosa oppositifolia. (Gay. l. c.)

1. *L. FERRUGINEUM* (Smith in Andr. bot. rep. t. 208.) segmentis calycinis utrinquè tomentosus. ♂ in Novâ-Hollandiâ. Vent! malm. t. 59. Sims bot. mag. t. 1766. Gay. l. c. p. 16. t. 3. (v. s.)
2. *L. PARVIFLORUM* (Rudg. trans. soc. lin. 10. p. 297. t. 12. f. 2.) segmentis calycinis intus glabris. ♂ in Novâ-Hollandiâ. Gay. l. c. p. 17. t. 4.

XIV. GUICHENOTIA. *Gay. diss. p. 18.*

Calyx persistens. Petala 5 glanduliformia. Fil. 5 libera. Antheræ rimis lateralibus dehiscentes. Ovarium 5-locul., loculis 5-ovulatis intus densè tomentosus. — Stipulæ o. Folia ternato-verticillata lineari-lanceolata integra. Inflor. intrâ-foliacea.

1. *G. LEDIFOLIA* (Gay. diss. p. 19. t. 5.) in Novæ-Holl. orâ occidentali. Frutex totus canescens, à Lasiopetalo ledifolio Vent. nunc Boroniâ ledifoliâ, omnino diversus.

XV. THOMASIA. *Gay. diss. p. 20.*

Cal. persistens venosus. Pet. 5 squammuliformia aut o. Fil. plerumque basi connata 5 aut 10. Antheræ rimis lateralibus dehiscentes. Ovarium 3-loc., loculis 2-8-ovulatis. Capsula 3-valvis, septis valvaribus, dehisc.-loculicida. Seminum strophiliola crenata. — Stipulæ foliaceæ persistentes. Folia ovata lobata utrinquè tomentoso-hispida. Inflor. racemosa oppositifolia, bracteis 3-partitis.

§. 1. *Species pentandræ, stylo longo, seminibus in singulo loculo 2.*

1. *T. PURPUREA* (Gay! l. c. p. 22. t. 6.) foliis lineari-ellipticis integris, stipulis foliaceis, petalis 5, capsulis stipitatis glabris profundè trisulcatis. ♂ in Novæ-Hollandiæ orâ australi-occid. Lasiopetalum purpureum. bot. mag. t. 1755. L. purpurascens herb. amat. t. 294. (v. s.)
2. *T. FOLIOSA* (Gay! l. c. p. 24. t. 7.) foliis ovatis cordatis obtusè 5-7-lobis, stipulis minutis, petalis nullis, capsulis sessilibus tomentosus obiter 3-sulcatis. ♂ in Novæ-Hollandiæ orâ occidentali-australi. (v. s.)

§. 2. *Species decandræ, stylo brevi, seminibus in singulo loculo 3-8.*

3. *T. SOLANACEA* (Gay! l. c. p. 26. t. 6.) petalis quinis. ♂ in Novæ-Hollandiæ orâ australi-occid. Lasiop. triphyllum Smith in Rees. cycl. L. solanaceum Sims. bot. mag. t. 1486. Folia generis maxima (v. v.)
4. *T. TRIPHYLLA* (Gay! diss. p. 28.) foliis sinuato-angulatis dorso glabriusculis, petalis nullis, capsulâ mucronatâ. ♂ in Novæ-Hollandiæ terrâ Van-Leuwin. Lasiopetalum triphyllum Labill! Nov.-Holl. 1. p. 63. t. 88. Folia videntur ternata ob stipulas magnas petiolatas. (v. s.)
5. *T. QUERCIFOLIA* (Gay. diss. p. 29.) foliis trilobis subtus hispidotomentosis, petalis nullis, capsulâ muticâ. ♂ in Novâ-Hollandiâ. Lasiop. quercifolium. Andr. bot. rep. t. 459. Sims. bot. mag. t. 1485.

XVI. KERAUDRENIA. *Gay. diss. p. 31.*

Calyx persistens. Pet. o. Filam. 5 fertilia basi imbricato-conniventia. Antheræ rimis dorsalibus dehiscentes. Ovarium 3-locul. loc. pluri-ovulatis. Styli 3 partim connexi! Caps. echinato-tomentosa abortu 1-locul.

3-valv. Semina bina, strophiolâ integrâ. — Stipulæ minutæ persistentes. Folia sinuato-undulata. Inflor. corymbosa, pedicellis articulatis.

1. *K. HERMANNIAEFOLIA* (Gay. l. c. p. 32. t. 8.) ♂ in Novæ-Hollandiæ orâ occidentali.

Trib. IV. *HERMANNIÆ*. — *Hermanniaceæ*. *Kunth malv.*
p. 11. nov. gen. am. 5. p. 312.

Flores hermaphroditi, Calyx 5-lobus persistens, nunc nudus. nunc subinvolucratus. Pet. 5 antè florescentiam spiraliter contorta. Stam. 5 basi monadelpha, omnia fertilia, petalis opposita, antheris ovatis bilocularibus. Carpella in fructum unicum coalita. Albumen carnosum-farinaceum. Embryo inclusus curvatus, radicula inferâ ovatâ, cotyledonibus planis foliaceis integris.

XVII. *MELOCHIA*. *H. B. et Kunth nov. gen. am. 5. p. 322.* — *Melochiæ* sp.
Linn. gen. n. 829.

Calyx 5-fidus, nudus aut bracteis 1-3 calyculatus. Petala 5 patentia. Stam. 5 basi monadelpha. Styli 5. Capsula 5-locul. loculicido-5-valvis, valvis medio septiferis. Sem. 1-2 in quoque loculo. — Pedunculi multiflori oppositifolii.

1. *M. PYRAMIDATA* (Lin. syst. 510.) foliis ovato-lanceolatis dentatis glabris, pedunculis 5-6-floris petiolo longioribus, petiolis ramisque puberulis, pubescentibus, subtus tenuiter canescenti-tomentosis, umbellis 7-10-floris petiolo longioribus. ♂ in ins. Caribæis, Brasiliâ. Cav. diss. 6. t. 172. f. 1. *M. Domingensis* Jacq. vind. 1. t. 30 Flores violacei. (v. v.)

2. *M. TURPINIANA* (H. B. et Kunth l. c. p. 323. t. 432.) foliis subinæquilateris ovatis acutis basi truncato-subcordatis duplicato-crenato-serratis suprâ pubescentibus, subtus tenuiter canescenti-tomentosis, umbellis 7-10-floris petiolo longioribus. ♂ in Novâ-Granatâ. Flores violacei.

3. *M. TOMENTOSA* (Lin. sp. 932.) foliis inæquilateris ovato-oblongis acutis crenatis plicato-lineatis utrinquè ramulisque cano-tomentosis, umbellis 3-8 floris petiolo longioribus axillaribus et in ramulis oppositifoliis. ♂ in campestribus siccis ins. Caribæarum et arenosis maritimis prope Cumanam. H. B. et Kunth nov. gen. am. 5. p. 323. Flores violacei. (v. s.)

β. *frutescens*, (Jacq. obs. 2. p. 44.) à priore habitu multò duriore foliisque minoribus differt. An spec. alt?

4. *M. MACROPHYLLA* (H. B. et Kunth l. c. p. 324.*) foliis æquilateris ovatis acutis cordatis crenato-serratis, suprâ hirtellis, subtus tenuiter molliterque canescenti-tomentosis, umbellis multifloris petiolum æquantibus. ♂ in Novâ-Andalusâ prope Bordones. Flores albi.

5. *M. PARVIFOLIA* (H. B. et Kunth l. c. p. 325.*) foliis parvis oblongis aut obovato-oblongis utrinquè obtusis crenato-serratis, suprâ scabriusculis, subtus adpresso-pilosiusculis, floribus in apice ramulorum umbellato-congestis. ♂ in aridis prope Caracas. Flores albi.

XVIII. *RIEDLEIA*. — *Riedlea Vent. choix. n. 37.* — *Mongeotia Kunth diss. p. 12.*
nov. gen. am. 5. p. 326. — *sena Houtt. pfl. syst. 6. p. 287.*

Calyx 5-fidus nudus aut bracteis 1-3 calyculatus. Pet. 5 patentia. Stam. 5 basi monadelpha. Styli 5. Carpella 5 in capsulam ideò 5-loc. 5-valv. subglobosam subconcreta, demum distincta rimâ longitudinali dehiscencia cocciformia, 1-2-sperma, axi centrali libero. — Inflorescentia varia. Flores albi aut flavi.

1. *R. POLYSTACHYA*, foliis oblongis acutis basi obtusis serratis, suprâ pubescentibus.

- tibus, subtus ramisque sericeo-pilosis, paniculâ terminali aphyllâ racemis plurimis pedunculatis cymosis seu secundifloris constante. 2? prope Honda ad ripam fl. Magdalenæ. Flores flavi. Mougeotia polystachya H. B. et Kunth nov. gen. am. 5. p. 328. t. 483. f. a et b.
2. *R. MOLLIS*, foliis ovatis acutis basi rotundatis subcordatis serratis, suprâ sericeo-pubescentibus, subtus molliter canescenti-tomentosis, umbellis axillaribus multifloris petiolum multò superantibus. 5 prope Honda et Santanna Novo-Granatensium. Petala alba versùs medium flava. Mougeotia mollis H. B. et Kunth l. c.
 3. *R. TILIÆFOLIA*, foliis cordatis dentatis junioribus velutino-tomentosis, adultis glabris, corymbis terminalibus multifloris, fructibus velutinis obtusè pentagonis. 5 in ins. Timor. Habitus ferè *R. odoratæ* et velutinæ. (v. s. ex h. Mus. Par.)
 4. *R. CRENATA*, foliis subrotundis crenatis tomentosis lineatis, umbellis pedunculatis axillaribus terminalibusque. 5 in Americâ merid. *Melochia crenata* Vahl. symb. 3. p. 86. t. 68.
 5. *R. ODORATA*, foliis ovatis subcordatis duplicato serratis acuminatis glabris, corymbis axillaribus pedunculatis. — in ins. Tannâ et Fa-uwa *Melochia odorata* Forst. prod. 253. — *Wisenia indica* Gmel. syst. 5. 5. *Visena indica*. Houtt. pfl. 6. t. 46. f. 3. ex Steud. nom. 521. An potiùs ad sequentem referenda?
 6. *R. VELUTINA*, foliis inferioribus cordatis, superioribus ovatis acuminatis simpliciter serratis utrinquè molliter velutinis, corymbis axillaribus pedunculatis. 5 in Bengalo, priori valdè affinis. (v. s. ex h. Calc.)
 7. *R. CORYMBOSA* (fl. mex. ic. ined.) foliis cordatis orbiculatis apice acuminatis villosis, corymbis axillaribus laxis pedunculatis. 5 in Mexici montibus calidis. Flores purpurei duplò majores quàm in *R. odoratâ*.
 8. *R. TRUNCATA*, foliis cuneiformibus apice dentatis supernè glabris subtus pube stellatâ canescentibus, floribus solitariis, capsulis depressis. 5 in Indiâ orientali. *Melochia truncata* Willd. sp. 3. p. 601.
 9. *R. DEPRESSA*, foliis ovatis serratis velutino tomentosis, pedicellis 1-3-floris oppositifoliis, capsulis obtusè pentagonis angulis ciliatis. 5 Hab. Havannæ. Lin., ad Sanctam-Martham. Bertero. *Melochia depressa* Lin. sp. 913. syn. Mill. excl. — Icon Cav. t. 173. f. 1. à nostrâ differt stipulis patentibus lanceolatis nec erectis subovatis, pedicellis brevioribus axillaribus 1-floris. (v. s.)
 10. *R. PERUVIANA*, foliis ovatis serratis tomentosis, floribus solitariis axillaribus, capsulâ globosâ villosâ. 5 in Peru. *Melochia Peruviana* Desrous. in dict. enc. 4. p. 83. Caulis suffrutescosus, humilis. Flores citrini.
 11. *R. SUPINA*, foliis ovatis serratis, floribus capitatis, caulibus procumbentibus. — in Indiâ. Pluk. alm. t. 232. f. 4. *Melochia supina* Lin. sp. 944.
 12. *M. CORCHORIFOLIA*, foliis ovatis sublobatis serratis glabris, floribus subterminalibus capitatis sessilibus. ① in Ceylonâ, Javâ. Dill. h. elth. t. 176. f. 217. *Melochia corchorifolia* Lin. sp. 944.
 13. *R. INFLATA*, foliis ovatis subacuminatis obsoletè cordatis duplicato-serratis utrinquè adpresso-pilosis, pedunculis axillaribus trifidis multifloris petiolo dimidiò brevioribus. 2? in umbrosis humidis regni Novo Granatensis. Mougeotia inflata. H. B. et Kunth nov. gen. 5. p. 330. t. 484. Flores albi.
 14. *R. NODIFLORA*, foliis ovatis acuminatis serratis glabris, junioribus adpressè pilosis, floribus axillaribus conglomeratis sessilibus. 5 in sepibus Jamaicæ. Cubæ, Cumanæ, Mexici, Porto-Ricci. Sloan. hist. t. 135 f. 2. *Melochia nodiflora* Swartz! fl. ind. occ. 2. p. 1139. *Mougeotia nodiflora* H. B. et Kunth nov. gen. am. 5. p. 330.* *M. carpinifolia* Wendl. obs. 52 (v. s.)
 15. *R. BORBONICA*, foliis ovatis acuminatis serratis subvillosis, floribus axillaribus conglomeratis sessilibus. 5 in ins. Borboniâ. Priori valdè affinis sed distincta ex cl. Kunth. *Melochia Borbonica* Cav. diss. 6. p. 321. t. 174. f. 1.

16. R. *BERTERIANA*, foliis ovatis subcordatis serratis glabris, capitulis globosis densis pedunculo petioli longitudine donatis. — in Guadalupâ. *Melochia Berteriana* Balb! in litt. (v. s.)
17. R. *BRIANCHA*, foliis ovatis dentatis puberulis brevissimè petiolatis, floribus axillaribus congestis sessilibus, calycibus villosis, carpellis 5-distinctis: 5 in Cayennâ. (v. s. sine fl.)
18. R. *HIRSUTA*, foliis ovatis acutis villosis, capitulis terminalibus subspiciformibus coarctatis hirsutis, calycibus tribracteatis. 5 in Novâ-Granatâ et ins. Caribæis ut autumno ex specim. incompl. *Mougeotia hirsuta* H. B. et Kunth nov. gen. am. 5. p. 331. (v. s.)²
19. R. *VENOSA*, foliis ovatis serratis venosis subtus tomentosis, pedunculis distinctis terminalibus multifloris, caule piloso. 5 in aridissimis Jamaicâ. *Melochia venosa* Swartz! ind. occ. 2. p. 1137. Flores lutei majusculi. (v. s.)
20. R. *CONCATENATA*, foliis ovato-lanceolatis dentatis glabris, racemis terminalibus confertis. 2 in Indiâ utrâque? — Pluk. t. 9. f. 5. *Melochia concatenata* Lin. sp. 944. Cav. diss. 6. t. 175. f. 2.
21. R. *JAMAÏCENSIS*, foliis ovato-lanceolatis dentatis adpressè villosis, superne glabriusculis, racemo terminali aphylo interrupto. 5 in Jamaicâ. *Melochia Jamaicensis* Balb! ined. Stam. in cylindrum ferè coalita. Fructus velutinus. (v. s.)
22. R. *CARACASANA*, foliis cordatis crenatis subtus tomentosis, floribus capitatis subsessilibus axillaribus oppositifoliisque. 5 ad Caracas. *Mougeotia Caracasana* H. B. K. et nov. gen. 5. p. 329. **Melochia Caracasana* Jacq. ic. rar. 3. t. 507.
23. R. *SERRATA* (Vent! choix. t. 37.) foliis ovatis cordatis acuminatis utrinque villosis serratis, petioli ramisque hispidis, stipulis lanceolato-linearibus petioli ferè longitudine, floribus axillaribus glomeratis subsessilibus. 2 in Porto-Ricco et Sancto-Domingo. Flores purpurei, 3-5-glomerati in spicam interruptam. (v. s.)

XIX. WALTHERIA. Lin. gen. n. 827.

Calyx 5-fidus, instructus involucello laterali 3-phylo deciduo. Pet. 5. Stylus 1. Stigma penicillatum. Capsula 1-loc., 2-valvis, 1-sperma aut rectius Carpella 5, 4 abortiva, 5^{ma} superstes. — Habitus *Melochiæ*.

1. W. *AMERICANA* (Lin. sp. 941. excl. ex Smith syn. Breyn.) foliis ovatis plicatis acutè inæqualiter dentatis utrinque tomentosis, capitulis pedunculatis. 5 in Bahama, Surinamo, ins. Caribæis. W. arborescens Cav. diss. 6. t. 170. f. 1. W. *indica* Jacq. ic. rar. 1. t. 130. Variat capitulis sessilibus et tunc forsan à W. *indica* non differt. Variat etiam pedunculis elongatis florum fasciculos plurimos quasi concatenatos gerentibus. (v. s.)
2. W. *ANGUSTIFOLIA* (H. B. et Kunth. nov. gen. am. 5. p. 332 *) foliis oblongis superne angustatis acutis basi rotundatis serratis molliter tomentosis, spicis glomeratis pedunculatis petiolum multò superantibus. 5 in aridis inter Bordonos et Cumana. Priori valde affinis. An W. *angustifolia* Lin. sp. 941. excl. syn?
3. W. *CORCHORIFOIA* (Pers. ench. 2. p. 216.) foliis ovato-lanceolatis duplicato-serratis, capitulis longè pedunculatis. — ad Rio-Janeiro Brasilæ. †.
4. W. *GLABRA* (Poir. dict. 7. p. 325.) foliis ovatis mucronato-serratis canleque glaberrimis, capitulis pedunculatis, calycibus ciliatis. 5 in Guadalupâ W. lævis Schrank! pl. mon. t. 55. Link. enum. 2. p. 179. Petala flava, linearia. (v. s.)
5. W. *CORDATA* (Smith in Rees cycl. vol. 37.) foliis cordatis latè ovatis lævibus acutè et inæqualiter dentatis, capitulis solitariis rigidè pedunculatis axillaribus. — in Indiâ occidentali?
6. W. *LOPHANTHUS* (Forst. prod. n. 252.) foliis subrotundo-cordatis serratis

- sericeo-tomentosis petiolatis, capitulis pedunculatis imbricato-bracteatis. *h* in Marchionis insulis. *Lophanthus tomentosus*. Forst. car. gen. 14.
7. *W. RHIOCLADA*, foliis ovalibus plicatis dentatis tomentosis, stipulis setaceis ciliatis persistentibus, capitulis breviter pedunculatis, ramis petiolis pedunculis hirsuto-tomentosis. *h* in Brasiliâ.
8. *W. INDICA* (Lin. sp. 941.) foliis ovalibus plicatis dentatis tomentosis, stipulis deciduis, capitulis sessilibus. *h* in Indiâ or. — Burm. zeyl. 149. t. 68. Varietas *W. Americanæ* ex cl. Brown.
9. *W. OVATA* (Cav. diss. 6. p. 317. t. 171. f. 1.) foliis subrotundo-ovatis inæqualiter dentatis tomentosis, capitulis sessilibus. *h* in Peruvîâ. (v. s.)
10. *W. ELLIPTICA* (Cav. diss. 6. p. 316. t. 171. f. 2.) foliis lanceolato-oblongis obtusis dentatis tomentosis, capitulis sessilibus. *h* in Indiâ orientali. (v. s.)
11. *W. MICROPHYLLA* (Cav. diss. 6. p. 317. t. 170. f. 2.) foliis oblongis obtusis plicatis dentatis canis, capitulis sessilibus. *h* in Indiâ orient. — Pluk. alm. t. 140. f. 5. Tota planta tomento tenui induta. Folia parva.
12. *W. LONGIFOLIA*, foliis oblongis acutiusculis dentatis subtus pubescentibus, capitulis sessilibus et in ramulis quasi interrupto-spicatis. *h* in insul. Caribæis. (v. s.)

XX. ALTHERIA. *Pet. Th. nov. gen. mad. n. 64.*

Calyx 5-fidus, cinctus involucello 3-phylo. Pet. 5. Stam. 5 in tubum monadelphum, antheris extus dehiscentibus. Ovarium 5-gonum. Styli 5 coaliti. Capsulæ 5, coalitæ, 1-spermæ. Semina centro affixa.

1. *A. MADAGASCARIENSIS*. — Herba erecta, hirsuta. Folia cordato-serrata. Flores fasciculati, axillares, parvi. †

XXI. HERMANNIA. *Lin. gen. n. 628.*

Calyx subnudus campanulatus 5-fidus. Pet. 5. Stam. 5, filam. imâ basi monadelphis lanceolatis sæpè alatis. Styli 5 in 1 coaliti. Caps. 5-loc., 5-valv., loculis polyspermis. — Frutices floribus sæpius flavis, pube sæpissimè stellatâ, omnes ad cap. Bonæ-Spei habitantes.

Sect. I. TRIONELLA.

Calyx (more Trioni) per et præsertim post anthesin inflatus. Stam. filamenta amplè dilatata.

1. *H. MULTIFIDA*, foliis canis palmato-partitis, lobis pinnati-partitis, lobulis linearibus integris subcomplicatis, pedicellis 1-floris folio brevioribus, calycibus inflatis puberulis. *h* *H. bipinnata* var. Burch! cat. n. 1627. trav. 1. p. 310. (v. s.)
2. *H. HALICACABA*, foliis subtus subalbidis superne punctatis, infimis palmato-5-partitis, super. 3-partitis, lobis pinnatifidis, medio longiore, pedunculis 2-4-floris folio brevioribus, calycibus inflatis puberulis. — *H. bipinnata* var. Burch! cat. n. 2020. An præcedentis var.? (v. s.)
3. *H. COMOSA* (Burch! cat. n. 1683.) foliis tomentosis ovatis sinuato-dentatis, stipulis lanceolato-linearibus pedunculis 2-floris, calycibus inflatis appendicibus ciliatis crinito-comosis! (v. s.)
4. *H. ALTHÆIFOLIA* (Lin. sp. 941.) foliis tomentosis ovatis plicatis crenatis, stipulis ovato-lanceolatis 3-5-nerviis, pedunculis 3-floris folio longioribus, calycibus inflato-angulatis puberulis. *h* Curt. bot. mag. t. 307. Cav. diss. 6. t. 179. f. 2. *H. aurea* Jacq. schœnbr. t. 214. (v. v.)
5. *H. PLICATA* (Willd. sp. 3. p. 589.) foliis tomentoso-hirtis ovatis subcordatis rugosis denticulatis, stipulis ovatis acutis, calycibus florentibus subeylin-draceis demùm inflatis. *h* *H. althæifolia* Jacq. schœnbr. t. 213. excl. syn.
6. *H. CANDICANS* (Ait. h. Kew. 2. p. 412.) foliis albidis tomentosis subro-

- tundo-ovatis crenatis, stipulis lanceolato-subulatis, calycibus campanulatis patulis. § Jacq. h. schœnbr. t. 117. H. præmorsa Wendl. obs. 51. (v. s. sine fl.)
7. H. MOLLIS (Willd. enum. 692.) foliis molliter tomentosis albidis oblongis obtusis dentatis basi cuneatis integerrimis, pedunculis bifloris, calycibus campanulatis velutinis. § vix à priore differt ex Link. (v. s.)
8. H. DECUMBENS (Willd. enum. suppl.) foliis pubescenti-tomentosis oblongis inæqualiter dentatis utrinquè subrotundatis, stipulis ovatis subdentatis, pedunculis 3-4-floris cernuis, calycibus inflato-globosis villosissimis. — Spreng. neu entd. 1. p. 299.
9. H. HYSSOPIFOLIA (Lin. sp. 942.) foliis pubescentibus cuneiformibus lanceolatis obtusis apice dentatis, floribus adgregatis, calycibus inflatis puberalis. § Cav. diss. 6. t. 181. f. 3. (v. v.)
10. H. TRIPOLIATA (Lin. sp. 942.) foliis albo-tomentosis sessilibus cuneato-obcordatis apice subcrenatis, stipulis oblongis obtusis foliola lateraliter mutuantibus, floribus adgregatis, calycibus inflatis puberulis. § Cav. diss. 6. t. 182. f. 2. Caulis humillimus quasi repens. (v. s.)
11. H. TRIPHYLLA (Cav. diss. 6. t. 178. f. 3. non Lin.) foliis subsabris tripartitis, partitionibus cuneiformibus apice truncatis dentatis, intermediâ petiolatâ, stipulis lanceolato-subulatis. § H. triphylla Lin. est Connari species. Hæc habet folia verè tripartita, prior folia simplicia. An cal. vesicarii?
12. H. GLANDULOSA (Link. enum. 2. p. 179.) foliis ovalibus inæqualiter crenatis subpubescentibus, stipulis ovatis acutatis sæpè fere incis, caule glanduloso-pubescente. § Flor. indescr. An hujus sect.? †
13. H. FRAGRANS (Link l. c.) foliis petiolatis ovalibus obtusis undulatis crenatis cauleque hirsuto-tomentosis, stipulis lanceolatis. § Flor. ign. et idè locus incertus. †

SECT. II. HERMANNELLA.

Calyx non aut vix inflatus. Stam. filamenta non sensim dilatata.

14. H. DISERMÆPOLIA (Jacq. Schœnbr. t. 121.) foliis albidito-tomentosis lanceolatis serratis obtusiusculis margine undulatis, stipulis subulatis, pedunculis 1-floris brevissimis. § Confusa ap. Pers. errore typogr. cum trifurcatâ. (v. s.)
15. H. DISTICHA (Schr. et Wendl. sert. hann. t. 10.) foliis hispido-villosis subrotundo-ovatis obtusis dentatis, stipulis subovatis acuminatis, pedicellis 1-floris folio brevioribus, calycibus angulatis. § H. rotundifolia Jacq. schœnbr. t. 118. (v. s.)
16. H. MELOCHIOIDES (Burch! cat. n. 2957.) foliis glabris ovatis inæqualiter dentatis, stipulis ovato-acuminatis, pedicellis 1-floris folio brevioribus, filamentis filiformi-linearibus. § (v. s.)
17. H. BRYONIFOLIA (Burch! cat. n. 2141.) foliis pube stellatâ scabris cordatis ovatis inæqualiter dentatis, stipulis lineari-lanceolatis, pedicellis 1-floris patenti-cernuis. § Folia ferè Malvæ bryonifoliæ. (v. s.)
18. H. SALVIFOLIA (Lin. suppl. 302.) foliis tomentoso-hispidis rugosis oblongis obtusiusculis integerrimis subsessilibus, stipulis longis lanceolato-subulatis, pedicellis 2-3-floris, floribus nudis. § Cav. diss. 6. t. 180. f. 2.
19. H. MICANS (Schr. et Wendl. sert. hann. t. 5.) foliis tomentoso-hispidis subrugosis oblongis obtusissimis apice subdentatis breviter petiolatis, stipulis lanceolato-subulatis, floribus adgregatis involucreatis. § H. latifolia Jacq. schœnbr. t. 119. Folia tomento-stellata subflavescentia. (v. v.)
20. H. INVOLUCRATA (Cav. diss. 6. t. 177. f. 1.) foliis tomentoso-hispidis oblongis acutiusculis integerrimis subsessilibus, stipulis lanceolato-subulatis, floribus subadgregatis bracteis lanceolatis involucreatis. § duab. prior. affinis. (v. s.)
21. H. SCORDIFOLIA (Jacq. schœnbr. 1. t. 120.) foliis subulis tomentosis oblongis obtusis crenatis petiolatis, stipulis subulatis, pedicellis 1-2-floris folio sub-

- longioribus, calycibus patulis. \S Folia supernè viridia scabriuscula, subtùs subalbida tomento brevissimo. (v. s.)
22. *H. DENUDATA* (Lin. f. suppl. 301.) foliis glabris lanceolatis apice serratis acutis, stipulis ovatis acuminatis, pedicellis 2-4-floris. \S Cav. diss. 6. t. 181. f. 1. Jacq. schœnbr. t. 122. (v. s.)
23. *H. ALNIFOLIA* (Lin. sp. 942.) foliis glabris latè obovato-cuneiformibus obtusissimis crenato-emarginatis plicatis, stipulis lanceolato-subulatis, pedunculis subtrifloris in apice ramorum subracemosis. \S Curt. bot. mag. t. 299. Cav. diss. 6. t. 179. f. 1. Jacq. schœnbr. t. 291. (v. s.)
24. *H. CUNEIFOLIA* (Jacq. schœnbr. 1. t. 124.) foliis pubescentibus obovatis cuneiformibus truncatis emarginatis dentatis, stipulis ovatis acutis, racemis terminalibus, pedicellis 1-floris. \S Folia multò quàm in præcedente minora. (v. s.)
25. *H. MOLOSERICEA* (Jacq. schœnbr. t. 292.) foliis molliter albido-tomentosis oblongis cuneiformibus apice rotundatis dentatis, stipulis lanceolatis, racemis terminalibus paniculatis. \S Willd. sp. 3. p. 594.
26. *H. HIRSUTA* (Schr. et Wendl. sert. han. t. 4.) foliis subtùs albo-tomentosis oblongo-obovatis cuneiformibus apice inæqualiter dentatis, stipulis semi-cordatis acuminatis, racemis lateralibus, pedicellis elongatis bifloris. \S Rami divaricati hirsuti. Folia supernè viridia Willd. sp. 3. p. 594. (v. s. in h. Balb.)
27. *H. SCABRA* (Cav. diss. 6. t. 182. f. 2.) foliis supernè scabris subtùs tomentosis cuneiformibus oblongis inæqualiter dentatis basi integerrimis, stipulis semi-cordatis acuminatis, pedunculis bifloris. \S Jacq. schœnbr. t. 127. excl. syn. — *H. aspera* Wendl. obs. 52.
28. *H. MULTIFLORA* (Jacq. schœnbr. t. 128.) foliis glabriusculis cuneiformibus oblongis truncatis apice dentatis, stipulis oblongis acutis, racemis paucifloris, calycibus campanulatis. \S Pedicelli 1-flori. (v. s.)
29. *H. FLAMMEA* (Jacq. schœnbr. t. 129.) foliis glabris cuneiformibus lanceolatis truncatis apice dentatis, racemis terminalibus, pedunculis 1-floris, calycibus subreflexis. \S Cor. aurantii coloris. (v. s.)
30. *H. ANGULARIS* (Jacq. schœnbr. t. 126.) foliis supernè glabris subtùs hirtis cuneiformibus lanceolatis truncatis apice dentatis, pedunculis bifloris calycibus alato-5-angulatis. \S Caulis pube fasciculatâ scaber. An ad sect. primam? (v. s. in h. Balb.)
31. *H. TRIFURCATA* (Lin. sp. 942.) foliis subvelutinis cuneiformibus linearibus obtusis apice integris tridentatisve, racemis multifloris secundis, calyce campanulato. \S Cav. diss. 6. t. 178. f. 2. Jacq. schœnbr. t. 125. Flores purpureo-violacei. Folia variant tomentosa aut glabra. (v. s.)
32. *H. ODORATA* (Ait. h. kew. 2. p. 412.) foliis velutinis cuneiformibus lanceolatis obtusis inf. apice 3-5-dentatis, sup. integris, stipulis lineari-subulatis, calycibus campanulatis patulis. \S Aff. *H. lavandulæfolia*. (v. s. sine fl.)
33. *H. LAVANDULÆFOLIA* (Lin. sp. 942.) foliis velutino-tomentosis lanceolatis obtusis, integerrimis, stipulis lineari-subulatis, racemis paucifloris, calycibus angustis. \S Dill. h. elth. t. 147. f. 176. Cav. diss. 6. t. 180. f. 1. Jacq. schœnbr. t. 215. (v. s.)
34. *H. VELUTINA*, foliis velutino-tomentosis oblongo-lanceolatis integerrimis apice mucronatis, stipulis linearibus acutis petiolo duplò longioribus, racemis paucifloris. \S Burch. cat. n. 3393. Priori affinis. (v. s.)
35. *H. LINIFOLIA* (Lin. mant. 256.) foliis linearibus, pedunculis 1-floris. \S †.
36. *H. FILIFOLIA* (Lin. f. suppl. 302.) foliis glabris margine scabris lineari-triquetris integerrimis, stipulis lanceolatis magnis. \S Cav. diss. 6. t. 180. f. 3. Jacq. schœnbr. t. 123.
37. *H. PROCUMBENS* (Cav. diss. 6. t. 177. f. 2.) foliis glabriusculis oblongis

- dentato-pinnatifidis, infimis ovatis, sup. elongatis, caule procumbente, racemis paucifloris. ζ (v. s.)
38. *H. TENUIFOLIA* (Sims bot. mag. t. 1348.) foliis pinnatifidis, lobis linearibus integris acutis. ζ Flores flavi brevissimè pedicellati nutantes. Folia in icone scabra, in specim. nostro glabra. (v. s.)
39. *H. INCISA* (Willd. sp. 3. p. 599.) foliis pinnatifidis, laciniis lineari-lanceolatis integerrimis, petalis unguiculatis inciso-dentatis. ζ Filamenta cuneiformia. Habitus Maherniæ.
40. *H. PULVERULENTA* (Andr. bot. rep. t. 164.) foliis scabriusculis albicantibus bipinnatifidis, pedunculis bifloris longissimis. ζ Corollæ sordidè luteæ. Maherniæ affinis sed filamenta lanceolata.
41. *H. ARGENTEA* (Smith in Rees cycl. vol. 17.) foliis squamis stellatis lepidotis duplò pinnatifidis, lobis decurrentibus, pedunculis racemosis 1-floris. ζ Folia ferè Lavandulæ multifidæ, sed minora et lepidota. Flores aurantio et flavo mixti. Calyx brevis campanulatus.
42. *H. CORONOPOLIA* (Link. enum. 2. p. 180.) foliis linearibus pinnatifidis canosis glabriusculis, caule pubescente. ζ Flor. ign.†.

XXII. MAHERNIA. *Lin. mant. 59.*

Calyx nudus campanulatus 5 fidus. Pet. 5. limbis obcordatis spiraliter convolutis, unguibus rectiusculis. Filam. 5. basi monadelphæ, medio in tuberculum cordatum aut in cupulam dilatata. Styli 5 interdum coaliti Capsula 5-loc., 5-valv., polysperma. — Species omnes hab. ad cap. B. Spei.

1. *M. VERTICILLATA* (Lin. mant. 59.) foliis verticillatis integris trifidiave linearibus, caule decumbente. ζ Cav. diss. 6 t. 176. f. 1. Hermannia ciliaris Lin. f. suppl. 302. Pluk. t. 344. f. 3. Variat foliis omninò glabris et ciliolatis. (v. s.)
2. *M. RESERAEFOLIA* (Burch! cat. n. 2280.) foliis glabris pinnatipartitis, lobis linearibus integris, stipulis incisiss trifidis. ζ Affinis sequenti, sed stipulis lobatis ab omnibus differt. (v. s.)
3. *M. BIPINNATA* (Lin. syst. 253.) foliis glabris bipinnatipartitis, lobis linearibus, pedunculis axillaribus elongatis bifloris. ζ Cav. diss. 6. t. 176. f. 2. Hermannia pinnata Lin. sp. 943? excl. syn.
4. *M. INCISA* (Jacq. h. schœnbr. 1. t. 54.) foliis pinnati-lobatis incisis pube aliâ stellatâ aliâ simplici subglandulosâ scabris, pedunculis subbifloris. ζ Flores albi marcescentes flavidi. (v. s.)
5. *M. DIFFUSA* (Jacq. schœnbr. 2. t. 201.) foliis pinnatifidis glabris, pedunculis bifloris ramulisque erectis, caule scabro procumbente diffuso. ζ Hermannia diffusa. Lin. f. suppl. 302. Flores flavi.
6. *M. HETEROPHYLLA* (Cav. diss. 6. p. 324. t. 178. f. 1.) foliis pube stellatâ sparsâ puberulo-scabris lineari-cuneiformibus grossè dentatis, stipulis linearibus integris, pedunculis subterminalibus 2-3-floris velutinis. ζ Hermannia grossulariæfolia. Lin. sp. 943. ex herb. Burm. (v. s.)
7. *M. ERODIODES* (Burch! cat. n. 1491.) foliis glabris ovatis obtusis inciso-dentatis, lobulis inæqualibus obtusis, stipulis ovatis, pedunculis bifloris folio duplò longioribus. ζ ζ .
8. *M. SESSELI-FOLIA*, foliis pinnati-partitis, lobis filiformibus integris acutis, infimis caulibusque pilis setosis hispidulis, ramis supernè glabris virgatis, pedunculis elongatis 1-2-floris. ζ (v. s. comm. a cl. Lambert.)
9. *M. PULCHELLA* (Cav. diss. 6. t. 177 f. 3.) foliis glabris pinnatifidis, lobis integris incisive obtusis, stipulis ovatis subincisis, pedunculis bifloris folio brevioribus, caule erecto subscabro. ζ Hermannia pulchella Lin. f. suppl. 302. (v. s.)
10. *M. VERNICATA* (Burch! cat. n. 1461. trav. 1. p. 278.) tota glaberrima, foliis

- pinnatifidis, lobis integris obtusis, stipulis ovatis, pedunculis subbifloris folio brevioribus, caule erecto. ♀ Flores nutantes. (v. s.)
11. *M. GLABRATA* (Cav. diss. 6. p. 326. t. 200. f. 1.) foliis pube stellatâ sparsâ punctato-scabris lanceolatis pinnatifido-dentatis, dentibus indivisis, pedunculis bifloris longissimis. ♀ Jacq. schœnbr. 1. t. 53. *Hermannia glabrata* Lin. f. suppl. 301. *Mahernia odorata* Andr. rep. t. 85. ex Ait. h. kew. ed. 2. v. 2. p. 199. Flores cernui odore Jonquillæ.
12. *M. EXALIDIFLORA* (Burch! cat. n. 1536. trav. 1. p. 295.) foliis glabris pinnatifidis, lobis integris subacutis, stipulis ovatis acutis, ramis erectis subscabris, pedunculis 1-2-floris folio longioribus. ♀ Aff. priori, sed folia profundius incisa et flores duplò majores. (v. s.)
13. *M. GRANDIFLORA* (Burch! trav. 1. p. 295. et in bot. reg. 3. t. 224.) foliis glabriusculis lanceolato-cuneatis obtusis integris aut apice dentatis, pedunculis bifloris divaricatis calycibusque viscosis. ♀ *Hermannia grandiflora* Ait. h. kew. ed. 2. v. 4. p. 141. Paters. itin. 60. ic. Flores cernui, in genere maximi. (v. s.)
14. *M. BISERRATA* (Cav. diss. 6. p. 326. t. 200. f. 2.) foliis glabris oblongo-lanceolatis duplicato-dentatis, pedunculis bifloris foliorum longitudine. ♀ *Hermannia biserrata* Lin. f. suppl. 302. *Habitus Hermannia*.
15. *M. VESICARIA*, foliis pilis furcato-ramosis subtus hirtis lanceolatis pinnatifidis, lobis denticulatis acutis, caule piloso-scabro, fructibus inflatis. ♀ *Hermannia vesicaria* Cav. diss. 6. p. 331. t. 181. f. 2. *M. odorata* var. ? incisa Pers. ench. 2. p. 218. Filamenta libera obcordata pilosa. Pers. †.
16. *M. LINNÆOIDES* (Burch! cat. n. 1878.) foliis utrinquè pube stellatâ punctatis ovato-rotundis crenatis, stipulis acutis, pedicellis 1-2-floris folio multò longioribus. ♀ Species distinctissima habitu Linnææ borealis! (v. s.)
17. *M?* *VIOLACEA* (Burch! cat. n. 3098.) foliis ovatis obtusis dentatis glabris, pedunculis oppositifoliis 1-3-floris folio longioribus. ♀? habitu accedit ferò ad *Melochiam pyramidalam*. (v. s. sine fructu.)
18. *M. SPINOSA* (Burch. trav. 1. p. 279.) foliis minutis cuneatis apice subtridentatis, cauleque erecto ramoso glabris, pedunculis demùm spinescentibus. ♀ ad cap. B. Spei prope Dwaal River Poort.

Trib. V. *DOMBEYACEÆ*. Kunth. *malv.* p. 12.

Cal. 5-lobus. Pet. 5 plana majuscula subinæquilatera, per aestivationem convoluta. Stam. numero petalorum multiplicia, 1-serialia, monadelphæ, rarò omnia fertilia, sæpissimè alia sterilia fili aut liguliformia, alia (2-3 inter quodque stamen sterile) fertilia, plus minùs inter se coalita. Styli 3-5 coaliti aut liberi. Ovula in loculis 2 aut plura biseriata. Embryo rectus in axi albuminis carnosus. Cotylédones foliaceæ, sæpè bifidæ, contortuplicatæ aut planæ.

XXIII. RUIZIA. Cav. diss. 3. p. 117.

Cal. persistens 5-partitus, involucello 3-phylo deciduo cinctus. Pet. 5. Stam. 30-40 basi in urceolum connata, omnia antherifera. Styli 10. Carpella 10 lignoso-membranacea 2-sperma arcuè cohærentia. Semina subtriquetra non alata. — Frutices Mauritanici, foliis subtus farinaceo-tomentosis, pedunculis axillaribus bifidis corymbosis.

1. *R. CORDATA* (Cav. diss. 3. p. 117. t. 36. f. 2.) foliis cordato-lanceolatis repandis. ♀ in ins. Borboniâ.
2. *R. LOBATA* (Cav. l. c. p. 118. t. 36. f. 1.) foliis cordatis crenatis 3-5-lobatis

oblongis, lobo medio productiore acuminato. 5 in ins. Borboniâ. (v. s.)

3. *R. VARIABILIS* (Jacq. schœnbr. 3. t. 295.) foliis ramorum florescentium palmatifidis, steriliū palmatipartitis. 5 in ins. Borboniâ. *R. palmata*. Cav. diss. 3. p. 119. t. 37. f. 1.

β. *laciniata* (Cav. l. c. f. 2.) foliis palmatipartitis, lobis linearibus angustissimis pinnatifidis. 5 in ins. Borboniâ.

XXIV. PENTAPETES. *Lin. gen. n. 834.* — Brotera. *Cav. ic. 5. p. 19.*

Calyx caducus involucello 3-phyllo unilaterali cinctus. Pet. 5. Stam. filamenta antherifera 3 inter quodque sterile. Stylus unicus apice 5-dentatus aut styli 5. Capsula 5-loc. 5-valv. ∞-sperma. Semina nuda nec alata. — Herbæ annuæ.

1. *P. PÆNICEA* (Lin. sp. 958.) foliis hastato-lanceolatis serratis, floribus 1-2 axillaribus cernuis, antheris 15, stylo 1 apice 5-dentato. ① in Indiâ orientali. Mill. ic. t. 200. Ker. bot. reg. t. 575. (v. v.)

2. *P. OVATA*, foliis ovatis serratis, pedunculis axillaribus 2-3-floris, antheris 5, styli 5. ① in Novâ-Hispaniâ. Brotera ovata Cav. ic. 5. p. 19. t. 433. Sprengelia modesta Schult. obs. 134. An genericè à priore differt? (v. s.)

XXV. ASSONIA. *Cav. diss. 3. p. 120.*

Calyx persistens 5-partitus involucello laterali 3-crenato 1-phyllo cinctus. Pet. 5. Stam. antherifera 15, nempe 3 inter quodque sterile. Styli 5 brevissimi. Carpella 5 disperma in capsulam arcuè coalita. Semina subtriquetra nec alata.

1. *A. POPULNEA* (Cav. diss. 3. p. 120. t. 42. f. 3.) foliis cordatis acuminatis utrinquè glabris subserratis, pedunculis vix petiolo longioribus. 5 in ins. Borboniâ. (v. s.)

2. *A. VIRBURNOIDES*, foliis cordatis subacuminatis crenatis subtus junioribus adpressè tomentosis, pedunculis petiolo triplò longioribus. 5 in insulâ Borboniâ. (v. s. comm. à Mus. Par.)

XXVI. DOMBEYA. *Cav. diss. 3. p. 121. Gaertn. fr 2. p. 259. t. 137. non Lam.*

Calyx 5-partitus persistens, involucello 3-phyllo unilaterali cinctus. Pet. 5. Stam. 15-20, filamentis vix basi coalitis, 5 sterilia, 2-3 fertilia inter quodque sterile. Stylus 1 apice in stigmata 5 subreflexa fissus. Carpella 5 bivalvia, 1-∞-sperma in capsulam arcuè connexa. Cotyledones contortuplicatæ bifidæ.

§. 1. *Involucelli foliolis latis ovatis cordatisve.*

1. *D. PALMATA* (Cav. diss. 3. p. 122. t. 38. f. 1.) foliis glabris palmato-7-nerviis 7-fidis, lobis lanceolatis serratis, corymbo bifido. 5 in ins. Borboniâ Var. foliis 5-nerv. 5-fidis. (v. s.)

2. *D. ACUTANGULA* (Cav. diss. 3. p. 123. t. 38. f. 2.) foliis junioribus tomentosis demum glabris cordatis 5-7-nerviis subrotundo-3-5-cuspidatis crenatis, corymbo bifido. 5 in ins. Borboniâ. (v. s.)

3. *D. ANGULATA* (Cav. diss. 5. p. 123. t. 39. f. 1.) foliis tomentosis cordatis subrotundis dentatis obsolete tricuspidatis, umbellis simplicibus, pedunculis petiolo brevioribus. 5 in insul. Borboniâ. Calyx tomentosus.

4. *D. TILIÆFOLIA* (Cav. diss. 3. p. 124. t. 39. f. 2.) foliis junioribus tomentosis, adultis subglabris, cordatis 7-nerviis crenatis, corymbo bifido, pedunculis petiolis ramulisque patentim pilosis. 5 in ins. Borboniâ.

5. *D. TOMENTOSA* (Cav. diss. 3. p. 125. t. 39. f. 3.) foliis tomentosis subrotundis cordatis acuminatis 5-nerviis transversè venosis crenatis, umbellis bifidis. 5 in Madagascar.

§. 2. *Involucelli foliolis angustis lanceolatis linearibusve.*

6. *D. CORDIFOLIA*, foliis glabris cordatis acuminatis crenatis 5-7-nerviis interdum subtrilobis, pedunculis petiolo subbrevioribus. 5 in Indiâ orient. Involucrum caducissimum. Antheræ 10. (v. s. ex hort. Calo.)
7. *D. UMBELLATA* (Cav. diss. 3. p. 127. t. 41. f. 1.) foliis glabris cordatis ovato-oblongis longè acuminatis subrepandis, umbellis globosis simplicibus, pedunculis petiolo duplò longioribus. 5 in ins. Borboniâ. Lam. ill. t. 576. f. 2. (v. s.)
8. *D. FERRUGINEA* (Cav. diss. 3. p. 128. t. 42. f. 2.) foliis subtùs tomentosis, sup. glabris, ovato-oblongis 7-nerviis subcordato-peltatis dentatis, corymbis bipartitis, pedunculis petiolo duplò longioribus. 5 in ins. Mauritii. Folia in sicco subtùs subferruginea. (v. s.)
9. *D. PUNCTATA* (Cav. diss. 3. p. 125. t. 40. f. 1.) foliis subtùs subtomentosis, sup. perne punctato-scabris, subcordatis ovato-oblongis acuminatis, 5-nerviis, corymbo conferto, pedunculis petiolo vix longioribus. 5 in ins. Borboniâ ad Cafrorum planitiem. (v. s.)
10. *D. OVATA* (Cav. diss. 3. p. 127. t. 41. f. 2.) foliis subtùs ferrugineo-tomentosis, sup. perne subglabris, 3-5-nerviis ovatis acutis integris, corymbo bifido, stylo minimo. 5 in ins. Borboniâ. (v. s.)

XXVII. MELHANIA. *Forsk. descr. 64. Ait. h. kew. ed. 2. v. 4. p. 146.*

Omnia Dombeyæ sed stamina 10, 5 alterna sterilia, 5 alterna fertilia 1-2-antheras gerentia. — An satis à Dombeyâ differt?

§. 1. *Filamentis fertilibus biantheriferis.*

1. *M. DECANThERA*, foliis glabris ellipticis utriusque acuminatis repando-dentatis, pedunculis multifloris umbelliferis. 5 in Madagascar. Dombeya decanthera. Cav. diss. 3. p. 126. t. 40. f. 2.

§. 2. *Filamentis fertilibus monantheris.*

2. *M. ERYTHROXYLON* (Ait. h. Kew. ed. 2. v. 4. p. 146.) foliis ovato-cordatis subpeltatis acuminatis crenulatis subtùs tomentosis reticulatis, pedunculis subtrifloris. 5 in memoribus Sanctæ-Helenæ. Corolla alba magna. Pentapetes Erythroxylon. Ait. h. kew. ed. 1. v. 2. p. 138. Dombeya Erythroxylon bot. mag. t. 1000.
3. *M. MELANOXYLON* (Ait. h. kew. ed. 2. v. 4. p. 146.) foliis cordatis integerimis utriusque tomentosis lævibus, pedunculis subtrifloris. 5 in Sanctæ-Helenæ. Pluk. mant. 6. t. 333. f. 5. In hac et priore involucella caducissima.
4. *M. VELUTINA* (Forsk. descr. 64.) foliis ovato-lanceolatis serratis tomentosis, umbellis axillaribus simplicibus 4-floris. 3 in Arabiæ-Felicis monte Melhân (undè generis nomen.) Pentapetes velutina Vahl. symb. 1. p. 49. Dombeya velutina Willd.
5. *M. BURCHELLII*, foliis oblongo-linearibus obtusiusculis subserratis utrinque cano-velutinis, pedicellis axillaribus 1-floris petioli longitudine. 5 ad cap. Bonæ-Spei. Melhania Burch! cat. n. 2417. (v. s.)
6. *M? PROSTRATA*, foliis linearibus integris, sup. perne glabris, subtùs adpressè canis, pedicellis axillaribus 1-floris petiolo duplò triplòve longioribus. 3 ad cap. Bonæ-Spei. Dombeya prostrata Burch! n. 2153. Priori valdè affinis, sed flores non vidi. (v. s.)

XXVIII. TROCHETIA. *DC. mem. mus. 10. p. 106.*

Calyx 5-partitus, patens, extùs nudus. Pet. 5. Stam. 20, filam. basi in urceolum monadelphis, 5 sterilibus. Ovarium 1 subrotundum lepidotum. Stylus filiformis. Capsula 5-locularis, 5-valvis. Semina parva subrotunda, aptera. — Frutices foliis integris subtùs ferrugineis lepidotisve, pedunculis axillaribus 1-3-floris reflexo-pendulis.

1. *T. UNIFLORA* (DC. l. c. cum ic.) foliis ovatis subacutis latitudine duplò longioribus, pedunculis 1-floris. 5 in ins. Borboniâ. (v. s.)
2. *T. TRIFLORA* (DC. l. c. cum ic.) foliis ovato-lanceolatis acuminatis latitudine quadruplò longioribus, pedunculis 3-floris, floribus ex apice pedunculi ortis, medio nudo, lateralibus 1-bracteatis. 5 in ins. Borboniâ. (v. s.)

XXIX. PTEROSPERMUM. *Schreb. gen. n. 1124. DC. mem. mus. 10. p. 111.*

Calyx 5-partitus, basi subtubulosus, nudus involucratusve. Pet. 5. Stam. 20, quorum 5 sterilia. Stylus cylindraceus. Stigma crassiusculum. Capsula lignosa, 5-loc. 5-valvis. Semina in alam producta, albumine parco aut nullo.

Sect. I. VELAGA. *Adans. fam. 2. p. 389.*

Involucellum nullum.

1. *P. ACERIFOLIUM* (Willd. sp. 3. p. 729.) foliis cordatis obtusis dentatis. 5 in Indiâ orientali. *Pentapetes acerifolia* Lin. sp. 939. Cav. diss. 3. p. 131. t. 44. *Velaga xylocarpa* Gaertn. fr. 2. p. 245. t. 133. — *Amm. act. petr. 8. t. 16-17.*
2. *P. SUBERIFOLIUM* (Willd. sp. 3. p. 728.) foliis oblongis acuminatis apice grossè subdentatis, pedicellis vix petiolo longioribus. 5 in Indiâ or. *Pentapetes suberifolia* Lin. sp. 959. Cav. diss. 3. t. 43. f. 2. *Sims bot. mag. t. 1526.* — *Amm. act. petr. 8. t. 14.*
3. *P. LANCEFOLIUM* (Roxb. cat. calc. 50.) foliis ovali-oblongis acuminatis integerrimis, pedicellis petiolo multò longioribus. 5 in Indiâ orient. (v. s. ex hort. Calc.)

Sect. II. PTEROLENA. *DC. l. c.*

Involucellum 3-phyllum, foliolis maximis cordato-rotundis fimbriato-laceris.

4. *P. SEMISAGITTATUM* (Roxb. cat. calc. 50.) foliis oblongis acuminatis integris basi cordatis, lobulis inæqualibus, alio brevi obtuso, altero longè producto acuminato. 5 in Indiâ orientali. — *Link. enum. 2. p. 200. DC. l. c. p. 113. cum icon.* — *Stirps speciosissima, involucello et formâ foliorum distinctissima. Flos involucello ablato valdè P. acerifolio similis. Fructus mihi ignotus.* (v. s. ex hort. Calc.)

XXX. ASTRAPAEA. *Lindl. coll. bot. t. 14.*

Flores umbellati involucro communi polyphylo cincti, foliolis subrotundo-ovatis, 2 ext. oppositis. Cal. 5 sepalus, extus 1-bracteatus. Petala 5 more *Malvavisci* convoluta-clausa. Stam. in tubum longè coacta, 5 sterilia, 20 antherifera. Ovarium 5-loc. Styl. 1. Stigm. 5. Ovula in loculis pauca inappendiculata.

1. *A. WALLICHII* (Lindl. l. c.) 5 in Indiâ orient. *Ker. bot. reg. t. 691. Arbor speciosissima, foliis magnis cordatis latè dentatis, stipulis foliaceis ovatis acuminatis, pedunculis longis hirsutis, corollis coccineis.* (v. s. in h. Mus. Par.)

†. *Dombeyaceæ incertæ.*

XXXI. KYDIA. *Roxb. cor. 3. p. 11.*

Calyx campanulatus 5-dentatus, involucello 4-6-phylo cinctus, foliolis involucelli calyci adnatis. Pet. 5 obliquè obcordata, calyce longiora. Stam. monadelphæ tubo longo, antheris in adelphas 5 tetrantheras ad apicem tubi dentium dispositis. Ovarium 1. Stylus 3-fidus. Stigmata dilatata. Capsula 3-loc., 3-valvis, 3-sperma. Semina in fundo loculorum adfixa. — *Arbores Indicæ foliis alternis 5-nerviis sub-5-lobis, floribus albis paniculatis.* — *Locus in ordine incertus.*

1. *K. CALYCINA* (Roxb. cor. 3. t. 215.) involucello 4-phylo calyce multò lon-

- giore. 5 in orâ Coromandelianâ et in Indiâ. — Antheræ subsessiles ad apicem tubi. Stylus extrâ tubum exsertus.
2. *K. FRATERNA* (Roxb. cor. 3. t. 216.) involucello 6-phylo calyce brevior. 5 in Indiâ montibus Circariensibus. — Antheræ ad apicem filamenti ultrâ tubum longè exserti adgregatæ. Stylus longitudine tubi.

XXXII? GLUTA. *Lin. mant.* 293.

Calyx campanulatus membranaceus deciduus nudus. Pet. 5, lanceolata, calyce longiora, toro longo stipitiformi adglutinata. Stamina circa torum monadelpha adglutinata, ex apice libera exserta. Antheræ subrotundæ, versatiles. Ovarium obovatum. Styl. 1. Fruct. — Arbor foliis lato-lanceolatis nudis, floribus paniculatis ferè ut in Clematide Flammulâ.

1. *G. BENGHAS* (Lin! mant. 293.) 5 in Javâ. Rami gemmæque pubescentes. Folia obtusa integra utrinquè lævia. (v. s. in h. Lin. apud cl. Smith.)

Trib. VI. *WALLICHIEÆ*. *DC. mem. mus.* 10. p. 102.

Calyx 5-lobus, involucello à floresubdistante 3-5-phylo cinctus. Pet. 5 plana. Stam. ∞ , filam. longè monadelphis, exterioribus minoribus, in columnam Malvacearum more dispositis, antheris erectis bilocularibus. Fruct. ign.

XXXIII. *ERIOLENA*. *DC. mem. mus.* 10. p. 102.

Calyx valdè tomentosus 5-partitus acuminatus, cinctus involucello 5-phylo, foliolis tomentosis laciniatis, 3 interioribus majoribus, 2 accessoriis minoribus, omnibus calyce brevioribus. Pet. 5 unguiculata calyce breviora. Stam. multiserialia monadelpha, exteriora breviora, omnia fertilia. Stylus villosus. Stigmata plurima parva in capitulum adgregata. — Genus novum sed fructu ignoto haud ritè determinandum (Wall! in litt.). Involucelli structurâ ad *Pterolænâ* accedens.

1. *E. WALLICHII* (DC. l. c. cum ic.) 5 in Indiâ orientali. Rami teretes. Folia petiolata cordata acuminata dentata, supernè pubescentia, subtùs villosa. Pedicelli villosi, petiolo duplò longiores 1-flori. (v. s. comm. ex h. Calc. à cl. Wallich.)

XXXIV. *WALLICHIA*. *DC. mem. mus.* 10. p. 104.

Involucellum 3-4-phyllum à flore intervallo distans, parvum. Calyx 4-partitus, lobis oblongo-linearibus, alabastro oblongo. Petala 4 patenti-reflexa, unguibus crassis velutinis. Stam. circiter 20 in tubum conicum monadelpha, exteriora breviora. Ovarium ovatum 8-loculare. Stylus 1. Stigmata 8. Capsula? loculis monospermis.

1. *W. SPECTABILIS*. (DC. l. c. cum icon.) 5 in Napanliâ. Ex sicco videtur arbor Tiliæ facie. Ramuli velutini. Folia petiolata latè ovata subcordata serrata subtùs velutina. Pedunculi axillares foliis longiores multiflori, paniculam foliaceam quasi constituentes. (v. s.)

XXXV. *GOETHEA*. *Nees et Mart. nov. act. bonn.* XI. p. 91.

Cal. campanulatus abbreviatus 5-fidus, involucello amplo vesicario 4-6-partito cinctus et obvellatus. Pet. 5, basi subconcreta, æstivatione spiraliter convoluta. Filam. ∞ in columnam longè coalita. Antheræ ovatæ hiloculares. Stylus elongatus, in stigmata 8-10 apice fissus. Cocco 5 coriacea 1-sperma. — Arbores aut Fruticeæ, foliis glabris-

culis coriaceis, petiolis barbatis, stipulis angustis, floribus speciosis axillaribus in pedunculis 1-floris nutantibus, involucellis coloratis reticulatis vesicariis. — Genus à cl. auct. ad Malvaceas relatum, sed ob antheras ab iis biloculares dictas potius huc pertiuens.

1. *G. SEMPERFLORENS* (Nees et Mart. l. c. p. 92. t. 7.) foliis ellipticis apice serratis, floribus subterminalibus, involucello 6-fido. 5 in Brasiliæ sylvis primævis juxta viam Felisbertiam ad ædes Ferreiræ Campos. Invol. fusco et corol. purpurecente flores Calycanthorum referunt.
2. *G. CAULIFLORA* (N. et M. l. c. p. 93 t. 8.) foliis oblongis integerrimis, floribus è trunco lateralibus, involucello 4-partito. 5 in Brasiliâ ad flumen Ilhéos. Folia conspicuè in spiram disposita. Flores involucello coccineo.



Ordo XXVII. TILIACEÆ. *Juss. gen. p. 290. §. 2. excl. gen. Kunth malv. p. 14. nov. gen. am. 5. p. 354.*

Calyx extus nudus, 4-5-sepalus, sepalis per æstivationem valvatis. Petala tot quot sepala, iis alterna, sæpè ad unguem foveolata, integra, rarissimè nulla. Stam. hypogyna, libera, numero sæpiùs indefinita, rarò definita. Antheræ ovales aut subrotundæ, biloculares, rimâ duplici longitudinali dehiscen-tes. Glandulæ tot quot petala, iis oppositæ, stipiti ovariifero adnatæ. Ovarium 1 carpellis 4-10 coalitis constans. Styli toti-
dem in 1 coaliti, stigmatibus sæpiùs liberis. Capsula plurilo-
cularis. Semina in loculis plurima. Albumen carnosum. Em-
bryo erectus, cotyledonibus planis foliaceis. — Arbores fruti-
cesve, rariùs herbæ. Folia simplicia bistipulata, sæpiùs den-
tata. Flores axillares.

I. SPARMANNIA. *Thunb. nov. gen. 88. Lin. f. suppl. 41.*

Calyx 4-sepalus. Pet. 4, subrotunda. Fila sterilia plurima torulosa basi subadhærentia. Stam. ∞. Capsula echinata, 5-angularis, 5-locul., loculis 2-spermis.

1. *S. AFRICANA* (Lin. f. suppl. 265.) ♂ ad cap. Bonæ-Spei in lateralibus montium. Retz obs. 5. n. 65. t. 3. excl. syn. et patr. Vent. malm. t. 78. Sims. bot. mag. t. 726. Flores albi. Fila sterilia flavo et fusco mixta. (v. v.)

II? ABATIA. *Ruiz et Pav. syst. p. 135. prod. t. 14. H. B. et Kunth nov. gen. am. 5. p. 358.*

Calyx 4-partitus. Pet. 0. Fila sterilia plurima staminiformia. Stam. ∞. Stigma 1. Capsula 2-valvis, 1-locularis, polysperma. Receptaculum lineare per singulas valvas excurrans. Semina striata. — Calyx valvatus ut in Tiliaceis. Fila sterilia ut in Sparmanniâ. Fructus ferè Bixinarum. locus ideò incertus.

1. *A. RUGOSA* (Ruiz et Pav. l. c.) foliis oblongo-lanceolatis rugosis, racemis terminalibus. ♂ in Peruvîæ collibus frigidis.
2. *A. PARVIFLORA* (Ruiz et Pav. l. c.) foliis lanceolato-oblongis acutis planis, racemis ex dichotomiâ. ♂ in Peruvîæ runcationibus vulgò Carpales dictis. A. verbascofolia H. B. et Kunth l. c. p. 358. t. 486. Folia utrius speciei colorem atrum præbent.

III. HELIOCARPUS. *Lin. gen. n. 606.*

Calyx 4-sepalus. Petala 4. Stam. 12-20. Stylus 1, bifidus, laciniis recurvatis. Capsula stipitata subcompressa utrinquè longitudinaliter ra-
diata, 2-loc. 2-valv. 2-sperma.

1. *H. AMERICANA* (Lin. sp. 643.) foliis etiam secùs nervos glabris, ramis cymæ glabris. ♂ in Verâ-Cruce. Montia Houst. Heliocarpus Lin. hort. cliff. 211. t. 16. Trew. ehr. t. 45. Folia cordata subtriloba serrata. (v. s.)
2. *H. POPAYANENSIS* (H. B. et Kunth nov. gen. am. 5. p. 341.) foliorum nervis venisque subtus hirsutis, ramis cymisque floccoso-tomentosis. ♂ in mon-
tosis prope Popayan. An prioris varietas?

IV. ANTICHORUS. *Lin. f. fasc. p. 3. t. 2.*

Cal. 4-sepalus. Pet. 4. Stam. 8. Stylus 1. Capsula subulata, 4-valv., 4-locul., loculis polyspermis. — Corchoro valdè affinis.

1. *A. DEPRESSUS* (Lin. mant. 64.) ① in Arabiâ, Jussiea edulis Forsk! descr. 210. Carictera Scop. Corchorus antichorus Rœusch. — Planta pusilla, foliis ovalibus grossè serratis, floribus axillaribus binis, fructibus reflexis. (v. s.)

V. CORCHORUS. *Lin. gen. n. 675.*

Cal. 5-sepalus deciduus. Pet. 5. Stam. ∞. Stylus subnullus. Stigmata 2-5. Capsula subulæformis aut rotunda 2-5-valvis, 2-5-locul., valvis medio septiferis. Semina hiserialia. — Corchorus Japonicus format genus inter Rosaceas à me jam anno 1816 Kerriadictum.

Sect. I. CORETA. *P. Browne. Kunth.*

Capsulæ siliquæformes bivalves biloc., apice in cornua non desinentes.

1. *C. siliquosus* (Lin. sp. 746.) capsulis linearibus compressis bivalvibus glabris, foliis lanceolatis æqualiter serratis, caule ramosissimo glabriusculo. ② in Americâ merid. — Plum. ed. Burm. t. 103 f. 1. Flores ex Lin. vernaes apetalæ 4-sepalæ 4-andræ, autumnales 5-sepalæ 5-petalæ polyandri. *C. linearis* Mill. dict. n. 5. et *C. secundiflorus* fl. mex. ic. ined. huc pertinere videntur. (v. s.)
2. *C. hirtus* (Lin. sp. 747.) capsulis linearibus compressis bivalvibus cauleque pilosis, foliis oblongis æqualiter serratis. — in Amer. calidiore. — Plum. ed. Burm. t. 103 f. 2. (v. s.)
3. *C. pilosus* (Link. enum. h. berol. 2. p. 72.) capsulis linearibus compressis subfalcatis hirtis stylo junioribus acuminatis, foliis oblongis subcordatis acutè crenatis glabriusculis, caule supernè hirtio. ② Patr. ign. Cum priore sæpè in hortis confusa.
4. *C. ORINOCENSIS* (H. B. et Kunth nov. gen. am. 5. p. 337.) capsulis siliquosis teretiusculis bilocularibus puberulis, foliis lanceolato-oblongis, superioribus lineari-lanceolatis acutis crenatis glabris subciliatis, caule subsimplici erecto. ① ad ripam fl. Orinoci, prope Angostura.
5. *C. ARGUTUS* (H. B. et Kunth l. c.) capsulis siliquosis compressis bilocularibus puberulis, foliis lanceolatis acutis argutè serratis glabris, caulibus ramosis, ③ prope Ibaguè Novo-Granatensium. Calyx extus pubescens.
6. *C. PILIOSUS* (H. B. et Kunth l. c. p. 338. t. 487.) capsulis siliquosis bilocularibus pilosis, foliis lanceolatis acutis serrato-crenatis utrinquè ramisque molliter pilosis. ③ prope Ibaguè Novo-Granatensium.
7. *C. MOMPOXENSIS* (H. B. et Kunth l. c.) capsulis siliquosis rectis ancipitibus bilocularibus pilosiusculis, foliis oblongo-lanceolatis angustato-acuminatis serratis glabriusculis, ramis glabris hinc lineà pubescente. ③ prope Mompox ad ripam fl. Magdalenzæ.

Sect. II. CORETOIDES.

Capsulæ siliquæformes 3-6-valves, 3-6-loculares, apice in cornua non desinentes.

8. *C. TRILOCULARIS* (Lin. mant. 77.) capsulis subulato-trigonis 3-valvibus scabris, foliis ovato-oblongis serratis, serraturis infimis sæpè setaceis. ① in Arabiâ. Jacq. h. Vind. t. 173. *C. æstuans* Forsk. descr. 101. (v. v.)
9. *C. OLITORIUS* (Lin. sp. 746.) capsulis oblongis subteretibus glabris 5-locularibus, foliis ovato-oblongis serratis, serraturis infimis longè setaceis. ① inter tropicos Asiæ, Africæ, Americæ, in hortis et ruderalis. Lam. ill. t. 478. f. 1. (v. s.)
10. *C. ÆSTUANS* (Lin. sp. 746.) capsulis oblongis 3-valvibus 6-sulcatis, foliis subcordatis ovato-acuminatis serratis, serraturis infimis longè setaceis. ① in Amer. calidiore. Jacq. h. Vind. t. 85. Pluk. t. 127. f. 3. (v. s.)
11. *C. SERRÆPOLIUS* (Burch! cat. geogr. n. 1962. voy. 1. p. 537.) capsulis lineari-teretibus scabris sæpius tortuosis sex-valvibus, pedunculis lateralibus trifidis trifloris, foliis linearibus grossè serratis. ② ad cap. B.-Spei, in planitie prope

Gattskamma trahs fl. Gariep. Caules procumbentes elongati villosi; alabastra acuta. (v. s.)

12. *C. ASPLENIFOLIUS* (Burch! cat. geogr. n. 1737. voy. i. p. 400) capsulis. . . . , pedunculis oppositifoliis trifidis trifloris, foliis elongato-ovatis simpliciter crenato-serratis, alabastris subglobosis. ♀ ad cap. B.-Spei prope vadum Salinarum fl. Gariep. An prioris var? (v. s. sine fr.)

Sect. III. CERATOCORETA.

Capsulæ elongatæ, angulatæ apice in cornua 3-5 divergentia desinentes.

13. *C. TRIDENS* (Lin. mant. 566. excl. syn. Burm.) capsulis gracilibus 2 3-valvibus 2-3-angulatis 2-3-cornutis, foliis oblongis serratis, serraturis infimis setaceis. — in Indiâ Lin., Senegaliâ. Variat foliis angustioribus et latioribus. *C. Senegalensis* Juss. — Pluk. t. 127. f. 4. (v. s.)
14. *C. ACUTANGULUS* (Lam. dict. 2. p. 104.) capsulis oblongis acutè angulatis in cornua 3-5 integra bifidave desinentibus, foliis ovatis serratis, hinc basi 1-setosis. ♀ in Sancto-Domingo, et Indiâ or. — Pluk. t. 44. f. 1. *C. tetragonus* Mill. dict. n. 4? An forsan 2 species hic confusæ? (v. s.)

Sect. IV. GANJA. *Rumph. amb. 5. t. 78. f. 1.*

Capsulæ subglobosæ depressæ rugoso-muricatæ.

15. *C. CAPSULARIS* (Lin. sp. 746.) capsulis subglobosis depressis rugoso-muricatis glabris, foliis oblongis acuminatis serratis, serraturis infimis setaceis. (1) in Indiâ orient. Gaertn. fr. t. 129. Jacq. f. eclog. 2. t. 120. — Pluk. t. 275. f. 4. (v. s.)

Sect. V. GUAZUMOIDES.

Capsulæ ovatæ, aculeis mollibus setoso-echinatæ subvelutinæ.

16. *C. HIRSUTUS* (Lin. sp. 747.) capsulis ovatis lanatis, foliis ovatis obtusis tomentosis æqualiter serratis crenatis integrisve. ♂ in Americâ calid. — Plum. ed. Burm. t. 104. *C. frutescens* Lam. dict. 2. p. 105. (v. s.)
17. *C. ARENARIUS* (H. B. et Kunth nov. gen. am. 5. p. 339.) capsulis oblongis trigonis trilocularibus, foliis parvis subfasciculatis oblongis utrinquè obtusis undulato-crenatis subcomplicatis coriaceis tomentosis, ramis floccoso-tomentosis. ♂ in arenosis Novæ Andalusie circa castellum Arayæ.
18. *C. TOMENTOSUS* (Thunb. fl. jap. 228.) capsulis oblongis lanatis, foliis ovatis obtusis tomentosis æqualiter serratis. ♂ in Japoniæ montibus. Flores aurantiaci.
- 19? *C. FASCICULARIS* (Lam. dict. 2. p. 104.) capsulis ovato-oblongis lanatis fasciculatis subsessilibus, foliis lanceolato-oblongis glabris æqualiter serratis. ♀ in Indiâ or. — Pluk. t. 439. f. 6.

†. *Species non satis notæ.*

20. *C? SCANDENS* (Thunb. act. soc. lin. lond. 2. p. 335.) foliis ovatis setaceo-serratis oppositis, caule ramisque flexuoso-scandentibus, floribus terminalibus solitariis. — in Japoniâ.
21. *C. SERRATUS* (Thunb. l. c.) foliis oblongis serratis cuspidatis, ramis glabris. — in Japoniâ. Rami purpurei.
22. *C. FLEXUOSUS* (Thunb. l. c.) foliis duplicato-serratis cuspidatis obliquè cordatis villosis, caule flexuoso. — in Japoniâ.
23. *C? JAVANICUS* (Burm. ind. 123. t. 36. f. 3.) capsulis subrotundis hispidis, foliis ovato-lanceolatis æqualiter serratis, calycibus aristatis. — in Javâ. An *Meiochis* affiniore?
24. *C. BURMANNI*, capsulis subulato-trigonis 3-lob. 3-valv., foliis lineari-oblongis serratis, serraturis infimis setaceis. ♀ in Indiâ or. *C. trilocularis* Burm. ind. p. 123. t. 37. f. 2. Stirps ad *C. trilocularem* aut ad *tridentem* forsan referenda, sed icon. ab utroque differt.
25. *C. BIFURCATUS* (Mill. dict. n. 6.) capsulis linearibus compressis, apicibus furcatis, foliis cordatis serratis. — in Jamaicâ.

VI. HONCKENYA. Willd. in *Ust. del. op.* 2. p. 201. t. 4.

Cal. 5-sepalus coriaceus, extus hirsutus, intus coloratus. Pet. 5 oblonga. Stam. 8, antheris oblongis. Fila. ∞ capillacea. Ovar. oblongum. Stylus 1. Stigma 6-dentatum? Capsula aculeis echinata, 5-loc., 5-valvis, valvulis medio septiferis. Semina ∞ arillata. — Affinis hinc Sparmanniæ, illinc Apeibæ.

1. *H. FICIFOLIA* (Willd. l. c.) 5 in Guineâ. Folia subtus tomento fulvo tecta, summa spathulato-oblonga dentata, inferiora obtusè 3-5-loba. Flores terminales terni cæruleo-violacei.

VII. TRIUMFETTA. *Lin. gen. n.* 600.

Cal. 5-sepalus, obtusus aut sæpè sub apice apiculatus. Pet. 5 aut rarius 0. Stam. 10-30 libera aut imâ basi vix subcoalita. Ovar. subrotundum. Stylus 1. Carpella 4, plus minus arcuè coalita in capsulam setis apice uncinatis echinatam. Semina in loculis bina aut solitaria. Embryo inversus.

Sect. I. LAPPULA. — *Triumfetta Gærtn. fruct.* 111.

Flores apetal. Carpella in capsulam non partibilem arcuè coalita. Semina in loculis solitaria.

1. *T. LAPPULA* (Lin. sp. 637.) foliis basi cordatis subrotundis inæqualiter dentatis subvillosis 5-nerviis 3-lobis, lobis acuminatis. 5 in insul. Caribæis et Bermudianis. *T. Plumieri* Gærtn. 2. p. 137. — Plum. ed Burm. t. 255. (v. s.)
2. *T. HETEROPHYLLA* (Lam. dict. 3. p. 420.) foliis basi indivisis, inferioribus inciso-trilobis, superioribus oblongis repandis. 5 in Sancto-Domingo. — Pluk. t. 425. f. 3.
3. *T. PILOSA* (Roth. nov. sp. p. 223.) foliis ovato-acuminatis integris inæqualiter serratis pilosis subtus sericeo-tomentosis basi vix emarginatis, capsularum echinis ciliatis. — in Indiâ orient. — Valdè accedit ad *T. Lappulam*.

Sect. II. BARTRAMEA. — *Bartramia Gærtn. fr. t.* 111. *non Hedw.*

Flores 5-petal. Carpella non arcuè coalita et capsula ideò matura 4-partibilis. Semina sæpè in loculis bina et tunc embryonis Cotyledones margine nec lateribus axi obversæ.

4. *T. RUBRICAULIS* (H. B. et Kunth nov. gen. am. 5. p. 342.) foliis oblongis acuminatis pubescentibus serratis, serraturis infimis glandulosis, ramis puberulis, pedunculis axillaribus 2-3 apice trifloris. 5 in umbrosis siccis ad Caracas. Rami nigro-purpurei. Stam. 24. Fructus sphaerici hirtelli 6-locul. indehiscentes. Pet. 5. An sect. propria?
5. *T. BOGOTENSIS*, foliis ovato-oblongis acuminatis argutè duplicato-dentatis ramisque pilosis, pedunculis axillaribus 2-3 apice trifloris. 5 prope Santa-Fé de Bogota. Stam. 10. Fructus globosi triloculares echinati. *T. pilosa* H. B. et K. nov. gen. am. 5. p. 342. non Roth.
6. *T. ROTUNDFOLIA* (Lam. dict. 3. p. 421.) foliis rotundatis inæqualiter crenatis subtus albedo-tomentosis, floribus subsapicatis, fructibus hirsutis echinatisque. 4 in Indiâ or. Folia 4-5 lin. diam.
7. *T. SUBORBICULATA*, foliis suborbiculatis integris eroso-crenatis supra nudiusculis subtus cinereo-tomentosis, racemo terminali laxo, fructibus uncinato-hispidis. — in Indiâ or. — Caules erecti sesquipedales. — *T. rotundifolia* Roth. nov. sp. 222. non Lam.
8. *T. GLANDULOSA* (Lam. dict. 3. p. 421.) foliis subrotundis lato-ovatisve indivisis subtus velutino-tomentosis dentatis, dentibus infimis subtus glandulosis, floribus axillaribus summis subsapicatis, calycibus pubescentibus apicu-

latis, fructibus tomentosis echinatisque. 5 in insulâ Mauritiî. *Bartramia* Lam. ill. t. 400. f. 1. (v. s.)

9. *T. Vahlî* (Poir. supp. 3. p. 300.) foliis ovato-lanceolatis serratis utrinquè pube stellatâ villosis subtus incanis, superiorum dentibus infimis glandulosis, stipulis subulatis. 5 in Arabiâ-Felici. *T. glandulosa* Forsk. cat. ar. 297. Vahl symb. 3. p. 62. An à priore differt?

10. *T. OVATA*, foliis ovatis indivisis inæqualiter dentatis supernè glabriusculis subtus cano-velutinis eglandulosis 7-nerviis, floribus interruptè spicatis.—in insul. Caribæis. (v. s.)

11. *T. TRICHOCLADA* (Link. enum. h. ber. 2. p. 5.) foliis ovatis acuminatis 7-nerviis serratis hirtis, ramis lineâ pilosâ, floribus confertis, calycibus apice hirtis 4-sepalis, fructibus echinatis. (1) in Napauliâ. *T. oblonga* Wall in litt. caulis ramosus 2-3-pedalis. Bractæ stipulæque hirtæ. (v. s.)

12. *T. OBLONGATA* (Link. l. c.) foliis oblongis serratis 5-nerviis molliter hirtis, caule ramoso supernè villoso, floribus terminalibus confertis. (1) in Napauliâ. Caulis 2-3-ped. Pili bulbo fusco. Stipulæ et bractæ lineares villosæ. Calyx hirtus.

13. *T. ANNUA* (Lin. mant. 73.) foliis ovatis acuminatis dentatis indivisis, supernè glabris, subtus in nervis subpilis, stipulis pedunculisque glabris, calycibus apiculatis. (1) in Javâ. Mill. ic. t. 298. Sims. bot. mag. t. 2296. — *T. indica* Lam. dict. 3. p. 420? Folia interdum sublobata pedunculi axill. 3-flori. Fructus aculeis nudis hamato-aculeati ex Sims. (v. s.)

14. *T. ANGULATA* (Lam. dict. 3. p. 421. var. α .) foliis basi ovatis 5-nerviis apice angulatis 3-lobis acuminatis inæqualiter dentatis subtus subvillosis, dente utrinquè infimo glanduloso, calycibus hispidis apiculatis, fructibus glabriusculis echinatis. — in Indiâ orient. Pluk. t. 41. f. 5. *Bartramia*. Lam. ill. t. 400. f. 2. (v. s.)

15. *T. SEMITRILoba* (Lin. mant. 73.) foliis basi ovatis trilobis calloso-serratis utrinquè velutinis, inferioribus sub-5-lobis, superioribus oblongis indivisis, calycibus pubescentibus subapiculatis, fructibus globosis glabris echinatis, setis retrorsum pilosulis. 5 in insulis Caribæis. (v. v.)

16. *T. HAVANENSIS* (H. B. et Kunth nov. gen. am. 5. p. 345.) foliis obovatis trilobis tenuiter membranaceis serratis ramisque pubescentibus, lobis lateralibus rotundatis, intermedio acuminato. 5 in ins. Cuba prope Havanam. Stam. 11. Fructus globosi echinati 3-loc. Valdè affinis *T. semi-trilobæ*.

17. *T. MOLLISSIMA* (H. B. et Kunth l. c. p. 345. t. 488.) foliis ovatis cordatis trilobis argutè duplicato-serratis mollibus suprâ tomentoso-pubescentibus subtus canescenti-tomentosis, lobo intermedio longissimè acuminato. 5 prope Santa-Fé de Bogota. Stam. 20. Fructus globosus cano-tomentosus echinatus.

18. *T. VELUTINA* (Vahl symb. 3. p. 62.) foliis ovatis acuminatis subangulatis inæqualiter serratis subtus velutinis, dentibus infimis subtus glandulosis, floribus axillaribus, calycibus pubescentibus subapiculatis, fructibus echinatis subpilosiusculis. — in ins. Mauritiî et regno Owariensi. Valdè accedit ad *T. glandulosam*. (v. s.)

19. *T. RHOMBOÏDEA* (Jacq. am. 147. t. 90.) foliis rhomboïdeis, infer. sub-5-lobis, mediis sub-3-lobis, summis oblongis, omnibus calloso-serratis utrinquè velutinis basi subcuneatis, floribus axillaribus, calycibus apiculatis, fructibus echinatis subpilis. 5 in insulis Caribæis et in Peruviâ. *T. rhombæifolia* Swartz prod. 76. *T. rhomboïdea* Lindl. coll. t. 29. Stylus apice tricuspis. An planta Peruana, à cl. Lindley depicta, omninò eadem ac Caribæa? (v. s.)

20. *T. HIRTA* (Vahl symb. 3. p. 63.) foliis ovatis trilobis supernè glabris, subtus velutinis, paniculæ terminalis ramis dichotomis hirtis. 5 ad Sanctam-Martham Amer. mer.

21. *T. ALTHÆOIDES* (Lam. dict. 3. p. 420.) foliis lato-ovatis indivisis acuminatis utrinquè molliter tomentosis inæqualiter serratis, serraturis infimis

exsertis calloso-glandulosis, floribus axillaribus, calycibus tomentosis apiculatis, fructibus echinatis setis retrorsum pilosulis. 5 in Cayennâ. T. macrophylla Vahl. ecl. 2. p. 34. (v. s.)

β. *subtriloba*, foliis obsolete trilobis, supra pubescentibus. 5 in Novâ Andalusâ prope Caripe. T. althæoides H. B. et Kunth nov. gen. am. 5. p. 343.*

22. T. *ACUMINATA* (H. B. et Kunth l. c.) foliis ovato-oblongis leviter cordatis angustato-acuminatis duplicato-serratis supra pubescentibus subtus molliter cano-tomentosis, ramis pilosis, floribus axillaribus. 5 in regno Novæ-Granatæ prope Mariquita. Stam. circiter 20. Fructus echinatus cano-tomentosus.

23. T. *GANDIFLORA* (Vahl! ecl. 2. p. 34.) foliis ovatis subcordatis acuminatis pilosiusculis inæqualiter serratis, dentibus infimis subtus glandulosis, calycibus glabris apiculatis, fructibus longè echinatis glabris. 5 in ins. Caribæis. (v. s.)

24. T. *POLYANDRA* (fl. mex. ic. ined.) foliis ovatis subcordatis acuminatis pilosiusculis serratis, calycibus glabris apiculatis, fructibus longè echinatis, setis pilosis. 5 in Mexico. Flores magni flavi. Genitalia stipite crasso insidentia. Fructus 5-loculares. Priori affinis.

† *Species non satis notæ.*

25. T. *PROCUMBENS* (Forst. prod. n. 204.) foliis subrotundo-cordatis subtrilobis tomentosis, caule procumbente. — in Societatis insulis.

26. T. *TRILOCULARIS* (Roxb. ex Horn. suppl. 140.) foliis subrotundo-ovatis trilobatis serratis, superioribus ovato-lanceolatis, capsulis trilocularibus. 5 in Indiâ orientali.

27. T. *INDICA* (Lam. dict. 3. p. 420.) foliis ovato-rhomboides indivisis subtus tomentosis, fructibus axillaribus, aculeis nudis. 5 in Indiâ or. An eadem ac T. annua?

28. T. *OXYPHYLLA* (fl. mex. ic. ined.) foliis cordatis acuminatis serratis villosis, paniculæ ramis subdichotomis, sepalis lineari-acuminatis petala oblonga duplò superantibus. 2 in montibus Novæ Hispaniæ.

29. T. *OBLIQUA* (Roth. nov. sp. p. 224.) foliis obliquè cordato-oblongis acuminatis indivisis inæqualiter serratis, supra atomis stellatis scabriusculis, subtus mollissimè tomentosis, capsularum echinis ciliatis. 5 Patr. ign. Flores completi.

N. B. *Triumfetta Bartramia* (Lin. sp. 638.) indeterminata remanet, cum synonyma omnia ad diversas species pertineant.

VIII. GREWIA. Juss. ann. 2. p. 82. — *Grewia* et *Microcos*. *Lin.*

Calyx 5-sepalus coriaceus intus coloratus. Pet. 5, basi intus glandulosa aut squamulosa, imo toro stipitifirmi inserta. Stam. ∞ ex apice tori orta, libera, antheris subrotundis. Stylus 1. Stigma 4-lobum. Drupa 4-loba, 4-pyrena aut abortu 2-3-pyr. Nuces biloculares, 2-spermæ, ant abortu 1-spermæ. Embryo erectus.

§. 1. *Petalis brevissimis aut nullis; foliis tri-rariùs 5-nerviis, sepalis 1-nerviis.*

1. G. *APETALA* (Juss. l. c. p. 93. t. 49. f. 3.) foliis 3-nerviis obovatis basi cuneatis crenulatis subasperis, pedunculis subsolitariis ramosis, floribus apetalis 4-fidis. 5 in Javâ.

2. G. *MULTIFLORA* (Juss. l. c. p. 89. t. 47. f. 1.) foliis 3-nerviis ovato-lanceolatis serratis glaberrimis, pedunculis adgregatis 3-4 floris. 5 in ins. Philipinis.

3. G. *GUAZUMÆFOLIA* (Juss. l. c. p. 89. t. 48. f. 3.) foliis 3-nerviis ovato-oblon-

gis acuminatis suprà glabris subtùs tomentosis crenato-dentatis, crenis inf. glandulosis, pedunculis solitariis 2-3-floris. 5 in Javà.

4. *G. TOMENTOSA* (Juss. l. c. p. 89. t. 49. f. 1.) foliis 3-nerviis ovato-lanceolatis hinc basi productionibus utrinquè tomentosis inæqualiter dentatis, pedunculis brevibus multifloris. 5 in Javà. *G. hirsuta*. Smith in Rees. cycl. n. 9.
5. *G. CELTIDIFOLIA* (Juss. l. c. p. 93.) foliis 5-nerviis ovato-oblongis acuminatis basi hinc productionibus supernè ad nervos pilosis subtùs tomentosis, pedunculis brevibus 2-3-floris. 5 in Javà. Flores ferè prioris. (v. s.)
6. *G. HIRSUTA* (Vahl. symb. 1. p. 34.) foliis 3-nerviis ovato-lanceolatis acuminatis basi rotundatis hinc subproductionibus utrinquè molliter villosis, pedunculis brevissimis subternis, petalis ciliatis. 5 in Javà. (v. s.)
7. *G. BRACTEATA* (Roth. nov. sp. 243.) foliis elliptico-lanceolatis obtusè serratis glabris, nervis venis ramisque ferrugineo-pilosis, pedunculis solitariis axillaribus 3-floris, floribus subsessilibus basi involucreatis, petalis nullis. 5 in Indià orient.

§. 2. *Petalis oblongis, sepalis 3-nerviis, foliis basi 3-nerviis aut penninerviis.*

8. *G. MALLOCOCCA* (Lip. f. suppl. 409.) foliis cordatis ovato-oblongis crenatis scabris, pedicellis axillaribus trifloris. 5 in ins. Tongatabu. *Mallococca crenata*. Forst. gen. t. 39 (v. s.)
9. *G. OPPOSITIFOLIA* (Buchan! in h. Lamb.) foliis ovatis acuminatis dentatis scabris, pedunculis folio oppositis multifloris. 5 in Napaulià. Roxb. cat. calc. p. 42. Genus proprium ex Wallich. (v. s. sine fruct.)
10. *G. UMBELLATA* (Roxb. cat. calc. p. 42.) foliis ellipticis subdenticulatis utrinquè glaberrimis, pedunculis 3-floris ad apices ramorum corymbosis. 5 in Indià or. Stipes genitalia gerens multò longior quàm in aliis speciebus. (v. s. ex hort. Calc.)

§. 3. *Petalis oblongis, sepalis enerviis aut 1-nerviis, foliis basi 3-nerviis aut penninerviis.*

11. *G. EXCELSA* (Vahl. symb. 3. p. 35.) foliis oblongis suprà glabris subtùs tomentoso-incanis obtusè serratis, pedunculis solitariis 1-3-floris, petalis calyce brevioribus. 5 in Ægypto et Arabià. *Chadara arborea* Forsk. descr. p. 105.

f. verrucosa (Juss! l. c. p. 90.) foliis hinc indè supernè verrucosis subrepandis. 5 in Javà. Verrucæ formatæ sùnt ab Erineo quodam inedito. (v. s.)

12. *G. BICOLOR* (Juss! l. c. p. 90 t. 50. f. 2.) foliis ovato-oblongis argutè serrulatis suprà glabris subtùs incanis, pediculis solitariis 1-3-floris. 5 in Senegal. Petala sepalis paulò breviora, staminibus æqualia. Fructus globosi glabri pìso dimidiò minores. (v. s.)
13. *G. ROTHII*, foliis ovato-oblongis acuminatis obtusis argutè serrulatis, suprà glabris, subtùs incanis, stipulis lanceolatis petiolo longioribus, pedunculis subternis subtrifloris, petalis staminibus duplò brevioribus. 5 in Indià orient. — *G. bicolor* Roth. nov. sp. p. 240. non Juss.
14. *G. SALVIFOLIA* (Roth. nov. sp. p. 239.) foliis oblongo-ovatis obtusis tennissimè crenulatis basi subintegris, suprà punctatis pubescentibus, subtùs niveo-tomentosis, pedunculis solitariis 3-floris, ovariis tomentosis. 5 in Indià orient. *G. bicolor* affinis. *G. salvifolia* Lin. est Alangii sp.
15. *G. FLAVA* (DC. cat. h. monsp. p. 113.) foliis ovali-cuneiformibus obtusissimis crenulatis glabriusculis subtùs pubescenti-canescensibus, pedicellis 1-floris, petalis bifidis calyce brevioribus. 5 in plagis orientalibus cap. Bonæ-Spei ultrà Orange flumen. Burchell. (v. v. et. s. sp.)
16. *G. CUNEIFOLIA* (Juss. l. c. p. 90. t. 49. f. 2.) foliis cuneiformibus crassius-

- culis glabris apice rotundato-crenatis basi integris, pedunculis solitariis bifloris, sepalis angustis petala paulò superantibus. 5 in Madagascar.
17. *G. NITIDA* (Juss. l. c. t. 47. f. 2.) foliis ovato-oblongis crenulatis glabris nitidis, pedunculis solitariis brevibus 1-2 floris, sepalis petalisque ovatis. 5 in Chinâ.
18. *G. MEXICANA*, foliis ovato-oblongis subserrulatis supernè glabris subtùs velutinis, pedunculis multifloris racemosis, petalis calycem æquantibus. 5 in Novâ-Hispaniâ. Alabastra velutina obovato-globosa. Unica species americana! Fructus ignotus! An vera Grewia? (v. s. comm. à cl. Lagasca.)
19. *G. OVALIFOLIA* (Juss. l. c. p. 90.) foliis ovalibus crenulatis glabris apice attenuatis, pedunculis 2-3-floris solitariis aut ad apices subracemosis, sepalis angustis petala duplò superantibus. 5 in Coromandelâ.
- 20? *G. Microcos* (Lin. syst. ed. 12. p. 602.) foliis ovato-oblongis acuminatis glabris subintegris, floribus paniculatis terminalibus. 5 in Indiâ or. *Microcos paniculata* Lin. sp. ed. 1. p. 514. — Burm. zeyl. t. 74. Rheed. mal. 1. t. 56. Petala calyce triplò breviora. (v. s.)
21. *G. PANICULATA* (Roxb. cat. p. 93.) foliis elliptico-oblongis obtusis cum acumine obtusè serratis subtùs villosis, floribus paniculatis terminalibus, stipulis sæpiùs bifidis. 5 in Indiâ or. *Microcos tomentosa* Smith in Rees. cycl. n. 2. Mal. misc. 1. n. 1. p. 13.
22. *G. ORIENTALIS* (Lin. sp. ed. 1. p. 964.) foliis ovatis oblongis crenatis subtùs subasperis obtusè acuminatis, pedunculis axillaribus 3-floris, sepalis angustis petala triplò superantibus stamina æquantibus. 5 in Indiâ or. — Rheed. mal. 5. t. 46. Pluk. t. 50. f. 4. (v. s.)
23. *G. COLUMNARIS* (Smith in Rees. cycl. v. 17. u. 5.) foliis ovato-oblongis crenatis utrinquè scabris, pedunculis 3-floris, genitalium stipite villosissimo elongato. 5 in Indiâ or. Stipulæ subulatæ petiolo æquales.
24. *G. PILOSA* (Lam. dict. 3. p. 43. excl. syn.) foliis ovatis crassiusculis subasperis crenatis, pedunculis 2-6-floris axillaribus et terminalibus, fructibus pilosis. 5 in Indiâ or. *G. orientalis* Gaertn. Vahl. ex Smith in Rees. cycl. n. 4.
25. *G. GLANDULOSA* (Vahl symb. 1. p. 34.) foliis ovato-lanceolatis acuminatis utrinquè lævibus crenulatis, crenis inferioribus glandulosis, pedunculis axillaribus petiolo brevioribus sub-3-floris. 5 in ins. Mauritiî. Juss. l. c. t. 48. f. 1.
26. *G. SERRULATA*, foliis ovato-oblongis acuminatis utrinquè lævibus serrulatis, serraturis infimis glandulosis, pedunculis axillaribus strictis petiolo longioribus subtrifloris. 5 in Bengalo. (v. s. ex h. Calc.)
27. *G. LÆVIGATA* (Vahl symb. 1. p. 34.) foliis ellipticis acuminatis utrinquè lævibus apice serratis basi integerrimis, pedunculis longis 3-floris. 5 in Indiâ or. Eadem ex Smith cycl. n. 7. ac *G. damina* Gaertn. et forsân *G. ovalifolia*. Juss.
28. *G. MOLLIS* (Juss. l. c. p. 91.) foliis ovato-lanceolatis dentato-serratis mollihus subtùs tomentosis, pedunculis subsolitariis subtrifloris petiolo longioribus. 5 in Senegaliâ et Owariâ.
29. *G. FLAVESCENS* (Juss. l. c. p. 91.) foliis ovato-oblongis acutis inæqualiter serratis utrinquè pilis stellatis scabris, stipulis linearibus petiolo longioribus, pedunculis subsolitariis 3-floris petioli longitudine. 5 in Indiâ or. Petala flavescentia.
30. *G. ACUMINATA* (Juss. l. c. p. 91. t. 48. f. 2.) foliis ovato-oblongis abruptè acuminatis crenulatis basi rotundatis glabris, pedunculis subgeminatis longis 2-3-floris. 5 in Javâ. Sepala pollicaria petalis triplò longiora. (v. s.)
31. *G. CARPINIFOLIA* (Juss! l. c. p. 91. t. 51. f. 1.) foliis ovatis basi cordatis apice acutis serratis subtùs subasperis, pedunculis 1-2 brevibus subtrifloris.

5 in Guineâ et Owariâ. Sepala angusta, staminibus æqualia, petalis longiora. Fructus lævis pisiformis. (v. s.)

♀. *rharnifolia* (Roth. nov. sp. p. 244.) foliis ovatis vix subcordatis. 5 in Indiâ orient.

32. *G. COMMUTATA*, foliis oblongo-ovatis acuminatis acutis basi subcordatis æqualibus inæqualiter serratis, suprâ stellato-pubescentibus, subtùs tomentosis, pedunculis 2-3 subcorymbosis petiolum subæquantibus. 5 in Indiâ orient. *G. carpinifolia* Roth. nov. sp. 245. non Juss.

33. *G. MEGALOCARPA* (Beauv! fl. ow. 2. p. 69. t. 102.) foliis oblongis obtusè acuminatis serratis lævibus, pedunculis 1-3-floris, fructu glabro 4-lobo. 5 in Owariâ. (v. s.)

34. *G. OBLIQUA* (Juss. l. c. p. 92.) foliis lanceolatis obliquis dentatis utrinquè pilis stellatis scabris, stipulis linearibus petiolo æqualibus, pedunculis solitariis 3-floris. 5 in Indiâ or.

35. *G. OCCIDENTALIS* (Lin. sp. ed. 1. p. 964.) foliis subrotundo-ovatis obtusis dentatis serratis, pedunculis solitariis 1-floris. 5 in Æthiopiâ et ad cap. Bonæ-Spei. Seba. thes. 1. t. 26. f. 3. Delaun. herb. amat. t. 95. Sepala petalis æqualia, staminibus paulò longiora. (v. v.)

§. 4. *Petalis oblongis, foliis 5-7-nerviis.*

36. *G. POPULIFOLIA* (Vahl symb. 1. p. 33.) foliis orbiculatis inæqualiter obtusè dentato-crenatis suprâ glabris subtùs pubescentibus, pedunculis solitariis longis 1-floris. 5 in Ægypto et Arabiâ. Chadara tenax Forsk. descr. 105. Gr. Chadara Lam. dict. 3. p. 44.

37. *G. BETULÆFOLIA* (Juss. l. c. p. 92. t. 50. f. 1.) foliis cordatis serratis subvillosis, pedunculis solitariis longis abortu 1-floris. 5 in Senegaliâ. An prioris var?

38. *G. ROTUNDIFOLIA* (Juss. l. c. p. 92. t. 50. f. 3.) foliis rotundis crenulatis albido-subtomentosis, pedunculis 2-3 petiolo longioribus 2-5-floris. 5 in Coromandeliâ. Folia ferè Betulæ pumilæ, subsessilia. *G. orbiculata* Roth. nov. sp. 247.

39. *G. ASIATICA* (Lin. mant. 122.) foliis cordatis subrotundis inæqualiter serratis suprâ glabris subtùs canescentibus, petiolo brevi incrassato, pedunculis 2-4 petiolo duplò triplòve longioribus 3-floris. 5 in Coromandeliâ, ins. Mauritiî. Sonn. voy. 2. t. 138. Folia omninò Tiliæ argenteæ.

40. *G. SUBINÆQUALIS*, foliis orbiculatis basi obliquis palmato-5-7-nerviis apice acuminatis serratis utrinquè pubescentibus, pedunculis 2-4 petiolo longioribus 2-3-floris. 5 in Indiâ or. Habui sub nom. *G. Asiaticæ* Roxb. sed differt foliis nec cordatis nec subtùs pube adpressissimâ canis, junioribus subtùs velutinis subalbidis, adultis concoloribus. An *G. arborea* Roth. nov. sp. 247. non Forsk? (v. s.)

41. *VILLOSA* (Roth. nov. sp. 248.) foliis subrotundo-ovatis breviter acuminatis basi subcordatis duplicato-serratis, suprâ stellato-villosis, subtùs stellato-tomentosis cinereis, utrinquè rufo-tomentosis, pedunculis confertis 1-floris petiolo brevioribus. 5 in Indiâ orient. Priori valdè affinis.

42. *G. ABUTILIFOLIA* (Juss. l. c. p. 92.) foliis lato-cordatis apice sinuato-angulatis inæqualiter dentatis, suprâ scabris, subtùs subtomentosis, pedunculis 2-3 petiolo duplò brevioribus 3-floris. 5 in Javâ. *G. arbutifolia* Pers. n. 31. Affinis sequenti.

43. *G. ASPERA* (Roth. nov. sp. 245.) foliis subrotundo-ovatis acuminatis subtrifidis basi obtusis inæqualiter serratis utrinquè stellato-pilosis asperis, pedunculis ternis 1-floris petiolo brevioribus. 5 in Indiâ orient. *G. abutilifoliæ* affinis.

44. *G. TILIÆFOLIA* (Vahl symb. 1. p. 35.) foliis cordato-subrotundis utrinquè glabris obtusè serratis hinc basi productioribus, pedunculis 2-3 petiolo bre-

vioribus 4-floris. 5 in Indiâ or. et Zeylonâ. *Microcos lateriflora* Lin. sp. 734. G. arborea Roxb. ex Willd. nov. act. nat. cur. 1813. p. 205.

45. G? *SAPIDA* (Roxb! cat. calc. 42.) foliis ovalibus 5-nerviis utrinquè obtusis grossè dentatis glabriusculis, pedunculis 3 petiolo quadruplò longioribus 3-floris. 5 in Bengalo. (v. s.)

46. G. *SCABRA* (Smith in Rees. cycl. 23. n. 3. sub *Microco.*) foliis oblongis basi cordatis 5-nerviis transversim reticulatis subtus scabris, paniculâ amplâ diffusâ, floribus involucreatis. 5 in Amboinâ. Petala lanceolata calycem æquantia.

47. G. *VILLOSA* (Willd. nov. act. nat. cur. 1813. p. 205.) foliis subrotundocordatis scâbris subtus tomentosis ciliato-serratis, pedunculis umbellatis petiolo brevioribus. 5 in Indiâ or. G. orbiculata Rottl. ex Willd. Eadem videtur ac G. villosa Smith. in Rees cycl. 17. n. 13. qui suam etiâ à Rottlero habuit, sed pedunculos petiolo æquales dicit.

48. G? *THEREBINTHINACEA* (DC. cat. h. monsp. p. 114.) foliis 5-nerviis cordatis acuminatis dentatis rariùs 3-lobis utrinquè molliter villosis, petiolis compressis, floribus racemosis, pedicellis adgregatis. 5 Patr. igu. Ramuli in ramis floriferis foliis oppositi. Folia trita odorem Pelargonii therebinthinacei spirant. Fructus ignotus. Interdum in hortis sub nomine *Heliocarpi* occurrit. An potius *Heliocarpi* sp? sed ovarium 4-loc.

49. G? *ERIOCARPA* (Juss. l. c. p. 93.) foliis ovatis 5-nerviis obtusè serratis subtus tomentosis, pedunculis 1-2 sæpè 3-floris, petalis angustissimis non glandulosis, toro non elongato. 5 in Javâ.

†. *Species non satis notæ.*

50. G. *VELUTINA* (Vahl symb. 1. p. 35.) foliis ovalibus utrinquè mollissimis subtus incanis obtusè serratis basi obliquis, pedunculis 2-3 brevibus 3-floris. 5 in Arabiâ. *Chadara velutina* Forsk. descr. 106.

51. G. *OBTUSIFOLIA* (Willd. enum. 566.) foliis oblongo-ellipticis utrinquè obtusis hirtis acutè inæqualiter dentatis. 5 in cap. Bonæ-Spei regionibus interioribus.

52. G. *GLABRA* (Mal. misc. 1. n. 1. p. 14.) foliis trinerviis glabris, ramulis tomentosis, floribus paniculatis. 5 in Indiæ insulâ Carnicobar. Aff. G. paniculatæ.

53. G. *AFRICANA* (Mill. dict. n. 2.) foliis ovato-lanceolatis serratis. 5 in Senegal. An eadem ac G. mollis?

N. B. Plures adhuc species indicatæ sunt à cl. Roxburghio in cat. hort. calc. p. 42 et 93. sed characteribus destitutæ et nonnullæ ad priores verosimiliter referendæ.

IX. COLUMBIA. ³ *Pers. ench. 2. p. 66. Colona. Cav. ic. 4. p. 47. t. 370.*

Cal. 5-sepalus deciduus intus coloratus. Pet. 5, intus ad basim squamulâ aucta. Torus pentagonus stipitiformis. Stam. ∞ libera. Stylus staminibus duplò longior. Fructus globosus 4-locul. 4-alatus, alis duplicatis, carpellis nempè coalitis 2-alatis 2-spermis. — Affinis Greviæ.

1. C. *SERRATIFOLIA*. 5 in insulis Philippinis prope Bannos. Columbia Americana. *Pers. l. c. Colona serratifolia. Cav. l. c. H. B. et Kunth nov. gen. am. 5. p. 335. in adn. Folia ovato-lanceolata serrata. Flores racemosi. (v. s.)*

X. TILIA. *Lin. gen. n. 660.*

Calyx 5-partitus, deciduus. Pet. 5 intus aut nuda aut squamulâ aucta. Stam. ∞, fil. liberis aut subpolyadelphis. Ovar. globosum villosum 1-stylum 5-loc., loculis 2-spermis. Nux coriacea, abortu 1-loc. 1-2-sperma. Cotyledones sinuato-dentatæ. — Arborea cortice flexili, ligno lævi. — Plures species variant foliis peltatis cucullatis.

§. 1. *Petalis nudis*. — Species omnes Europææ. — T. Europæa. Lin.
sp. 733.

1. T. MICROPHYLLA (Vent! diss. p. 4. t. 1. f. 1.) foliis cordato-subrotundis acuminatis argutè serratis glabris ad axillas venarum subtùs barbatis petiolo vix longioribus, fructibus subglobosis vix costulatis tenuissimis fragilibus. § in Europæ sylvaticis submontanis. T. Europæa γ Lin. sp. 733. T. ulmifolia Scop. carn. n. 642. T. sylvestris Desf. cat. hort. par. p. 152. T. parvifolia Ehrh. ex Sut. fl. helv. 1. p. 317. Variat ut plures aliæ species foliis peltato-cucullatis. (v. v.)
2. T. INTERMEDIA (Hayn. et Sv. bot. t. 40.) foliis cordatis acuminatis serratis glabris ad axillas venarum subtùs barbatis petiolo duplò longioribus, fructibus membranaceis oblongis difformibus dispermis. § in lucis Sueciæ T. Europæa borealis Wahl ups. 181. OEd. fl. Dan. t. 553 ex quâ fructus videtur 5-costatus.
3. T. RUBRA (DC. cat. h. monsp. p. 150.) foliis cordatis basi inæqualibus subtùs petiolis turionibusque pilosiusculis, axillis venarum subtùs barbatis, fructibus globosis lævibus. § in Tauriâ (Steven), forsan in Etruriâ, Græciâ. T. triflora Puer. in Horn. cat. 2. p. 493. T. Corinthiaca. Bosc. dict. agr. 13. p. 139. Ramuli rubri. — An T. corallina hort. kew. huc referenda? (v. v.)
4. T. PLATYPHYLLA (Scop. carn. n. 641.) foliis cordato-subrotundis acuminatis argutè serratis subtùs puberulis, fructibus turbinatis prominulo-costatis lignosis etasis. § in Europæ sylvaticis submontanis. Vent. l. c. p. 6. t. 1. f. 2. T. cordata Mill. dict. n. 1. T. cordifolia Bess. gal. 1. p. 343. T. Europæa Desf. cat. 152. T. grandifolia. Ehrh. ex sut. fl. helv. 1. p. 317. (v. v.)

§. 2. *Petalis intùs basi squamulâ auctis*. — Species pleræque Americæ.

5. T. GLABRA (Vent. diss. p. 9. t. 2.) foliis profundè cordatis abruptè acuminatis argutè serratis subcoriaceis glabris, petalis apice truncatis crenatis, stylo petalis æquali, fructu ovato subcostato. § in Americâ bor. T. Americana. Lin. sp. 733 Michx. f. arb. am. 3. p. 311. t. 1. T. Canadensis Michx. fl. bor. 1. p. 306. (v. v.)
 6. T. LAXIFLORA (Michx. fl. bor. am. 1. p. 306.) foliis cordatis sensim acuminatis serratis membranaceis glabris, paniculis laxifloris, petalis emarginatis, stylo petalis longiore, fructu globoso. § in Americâ à Marylandiâ ad Georgiam. Species distinctissima ex Pursh. fl. bor. am. 2. p. 363. (v. s.)
 7. T. PUBESCENS (Ait. h. kew. ed. 1. v. 2. p. 219.) foliis basi truncatis subcordatis obliquè denticulato-serratis subtùs pubescentibus, petalis emarginatis, stylo petalis longiore, fructu globoso lævi. § in Americâ à Virginiâ ad Georgiam. Vent. diss. p. 10. t. 3. Duham. ed. nov. t. 51. Michx. f. arb. am. 3. p. 318. t. 3. T. Caroliniana. Mill. dict. n. 4. T. Americana. Walth. car. 153. (v. s.)
- β. *leptophylla* (Vent. diss. p. 11.) foliis tenuissimis profundè et rariter serratis. T. Mississipensis hort. Par. An species propria?
8. T. HETEROPHYLLA (Vent. l. c. p. 16. t. 5.) foliis ovatis subtùs tomentosis basi nunc cordatis nunc obliquè aut æqualiter truncatis, nuce globosâ 5-costatâ. § in Americâ ad ripas Ohio et Mississipi. Pursh fl. 2. p. 363. T. alba. Michx. f. arb. am. 3. p. 315. t. 2. An Ait. ed. 1. non ed. 2? — Habeo specimen huic simillimum ex agro Mexicano. (v. s.)

9. T. ARGENTEA (Hort. Par. et DC. cat. h. monsp. p. 150.) foliis cordatis subacuminatis basi subinæqualibus serratis; subtùs niveo-tomentosis, supernè glabris, petiolo quadruplò serè longioribus, fructibus ovatis costis non prominulis. § in Hungariâ. T. alba. Wadst. et Kit. pl. rar. 1. p. 3. t. 3. T. rotun-

difolia Vent. diss. p. 13. t. 4. Duham. ed. nov. t. 52. T. *tomentosa* Moench. Weiss. 136. (v. v.)

10. T. PETIOLARIS, foliis cordatis acuminatis serratis supernè glabris subtùs pube adpressissimâ albis petiolo duplò longioribus. 5 in horto Odessano culta. Priori valdè affinis, sed petiolo duplò longiore differt. Species recognoscenda! (v. s. sine fl. et fr.) †.

XI. DIPLOPHRACTUM. Desf. mem. mus. 5. p. 34. t. 1. Kunth malv. p. 15. nov. gen. am. 5. p. 334. in adn.

Calyx 5-sepalus. Pet. 5 basi intùs squamulâ ad basim aucta. Stam. ∞, libera. Stylus 1. Stigmata 5 aggregata. Capsula globosa indehiscens 5 alata 10-locul., loculis septis transversis in loculamenta monosperma partitis. Semina arillata parietibus adfixa. Embryo ad basim albuminis carnosì.

1. D. AURICULATUM (Desf. l. c.) 5 in Javâ. Folia sessilia oblonga rugosa subtùs tomentosa versus apicem serrata basi obliqua. Stipulæ aristatæ.

XII. MUNTINGIA. Lin. gen. n. 651.

Cal. 5-7 partitus, deciduus. Pet. 5-7. Stam. ∞, libera. Ovarium sessile globosum basi pilis creberrimis cinctum. Stylus 0. Stigma capitato-radiatum persistens. Bacca 5-loc. ∞-sperma. Semina in pulpâ nidulantia minima. Embryo parvus. (Gaertn. fr. 1. t. 59.) Genus affine Apeiba.

1. M. CALABURA (Lin. sp. 728. Jacq. am. t. 107.) 5 in ins. Caribæarum collibus calcareis. Folia subsessilia basi obliqua oblonga acuminata. Flores axillares albi. (v. s.)

XIII. APEIBA. Aubl. guian. 1. p. 538. — Aubletia Schreb. gen. n. 889. — Oxytaendrum Neck. elem. n. 1005.

Cal. 5 rariùs 4-partitus. Pet. 5 rariùs 4 unguiculata, calyce subminora. Stam. ∞, filam. brevibus antheris longis apice lobulis 2 foliaceis apiculatis. Stylus 1 dilatatus in stigma infundibuliforme dentatum. Capsula sphæroïdea depressa coriacea setis mollibus tuberculisve rigidis scabra 8-24-locul. Semina in loculis ∞ minuta recept. carnosò centrali adfixâ. — Arbores pube stellulatâ. Folia integra Pedunculi oppositifolii. — Genus ad Elæocarpeas paululùm antheris accedens.

1. A. TIMOURBOU (Aubl. guian. 1. p. 538. t. 213.) foliis cordato-lanceolatis serratis subtùs hirsutis, fructibus densè setosis 10-locularibus. 5 in collibus Guianæ Cayennæ, Tabago et prov. Caracasanae. Rich! act. soc. hist. nat. par. p. 110 H. B. et Kunth nov. gen. am. 5. p. 347.* Mareg. bras. t. 124. Slossneæ Lœff. it. 311. A. hirsuta. Lam. dict 1. p. 208. (v. s.)
2. A. PETOUMO (Aubl. l. c. p. 543. t. 215) foliis oblongis subcordatis serratis subtùs incanis, fructibus densè setosis 18-20 locularibus. 5 in sylvis Guianæ. Rich! act. soc. hist. nat. par. p. 110. A. hispida Gaertn. fr. 2. t. 121, ambigit inter hanc et priorem, cum Tibourbou longitudine setarum, cum Petoumo numero loculorum conveniens; sed in specim. Richardi apud am. Delessert servato, folia subtùs ferè omninò glabra. (v. s. in h. Deless.) — An species propria?
3. A. ULMIFOLIA (H. B. et Kunth nov. gen. am. 5. p. 347.) foliis elliptico-oblongis subacuminatis basi rotundatis argutè dentatis suprâ puberulis subtùs tenuissimè ferrugineo-tomentosis et arachnoïdeis. 5 in ripâ flum. Orinoci. Affinis videtur A. Petoumo. Fruct. iga.

4. *A. ASPERA* (Aubl. l. c. p. 545. t. 216.) foliis oblongis subcordatis integerrimis subtus pubescentibus, fructibus muricatis 8-10-locularibus, tuberculis conicis brevibus pubescentibus. 5 in Cayennâ, ubi cum specie secundâ dicitur Petonomo. Lam. dict. 1. p. 209. ill. t. 470. f. 3. Gaertn. fr. 2. p. 188. t. 121. (v. s. s. fr.)
5. *A. ECHINATA* (Gaertn. fr. 2. p. 189. t. 121.) foliis . . . , fructibus muricatis 8-10-locularibus tuberculis longis pyramidatis glaberrimis. 5 Patr. et arbor ipsa ignot. †.
6. *A. GLABRA* (Aubl. l. c. p. 541. t. 214.) foliis oblongo-obovatis acuminatis integerrimis glabris, fructibus scabris 8-10-locularibus. 5 in Cayennâ. Lam. ill. t. 470. f. 1. Petala obtusa virescentia.
7. *A? DISCOLOR* (Spreng. neu. entd. 2. p. 166. sub Aubletia.) foliis ovato-oblongis integerrimis subtus hirsutis discoloribus, antheris apice receptaculoque setosis. 5 in Brasiliâ. Flores subterni axillares. Calyx extus pilis ferrugineis sericeus. Fruct. ignot.

XIV. SLOANEA. *Lin. gen. n. 655. non Læfl.*

Cal. 4-7-sepalus, sepalis lanceolato-linearibus extus subvelutinis intus coloratis, nunc imâ basi, nunc ad medium ferè connexis. Pet. o. Stam. ∞, filam. subnullis, antheris longissimis acumine parvo superatis. Ovar. 1. Styl. 1 filiformis. Capsula coriaceo-lignea subrotunda setis sublignosis confertissimis echinata, 4-5-locul., 4-5-valv., loculis 1-3-spermis. Semina arillo carnoso tecta. — Arbores ex Americâ æquinoctiali. Folia alterna, ampla. Flores 1-bracteolati. — An Genus dividendum?

Sect. I. SLOANEA. *Plum. gen. t. 15.*

Calyx 6-7-fidus. Stylus longus simplex. Capsula 4-valvis, setis variè inflexis. Semina arillo carnoso involuta.

1. *S. DENTATA* (Lin. sp. 730.) foliis ovatis acutis dentatis, stipulis cordato-triangularibus serratis. 5 in Americâ merid. — *Plum. ic. ed. Burm. t. 244?* *Castanea Sloanea* Mill. dict. *Sloanea grandiflora* Smith in Rees cycl. n. 2. — *Icon. Plumerii* exhibet folia cordata acutè dentata. Meum specimen è Cayennâ, ad quod pertinet *S. Plumerii* Aubl. guian. 1. p. 536, habet folia ovata obtusè sinuato-dentata. — An duæ species hic adhuc confusæ? (v. s.)

Sect. II. GYNOSTOMA.

Calyx 5-partitus, lobis æqualibus. Torus incrassatus. Antheræ extus hirsutæ. Stylus subulatus. Stigma perforatum, vix denticulatum. Capsula 4-valvis setis variè inflexis, valvis à basi ad apicem dehiscentibus.

2. *S. MASSONI* (Swartz fl. ind. occ. 2. p. 938.) foliis cordato-ellipticis obtusis integris, stipulis linearibus. 5 in ins. Caribæis. (v. s.)

Sect. III. MYRIOCHÆTA.

Cal. 5-partitus, lobo 1 ceteris minore. Torus villosus. Stylus brevis. Stigmata 4-5 simplicia. Caps. 4-5-loc., 4-5-valv., setis rigidis confertissimis tecta.

3. *S. SINEMARIENSIS* (Aubl. guian. 534. t. 212.) foliis subrotundo-ovatis integris, stipulis longis acuminatis deciduis. 5 in Guianâ et ins. Sancti-Christophori. Lam. ill. t. 469. dict. 6. p. 21. *S. Aubletii* Swartz fl. ind. occ. 940.* An *Ablania* species? (v. s.)

Sect. IV. OXYANDRA.

Cal. 5-partitus, lobis lanceolato-linearibus acuminatis. Torus parvus. Stylus

filiformis longus simplex. Capsula (ex ovario) verosim. setis destituta. An genus proprium?

4. S? CORYMBIFLORA, foliis ovatis acutis integris, stipulis deciduis, pedunculis axillaribus apice ramoso-corymbosis multifloris. 5 in Guianâ Gallicâ. (v. s.)

Sect? v. FOVEOLARIA. DC. — non R. et P. fl. per.

Cal. 4-partitus, lobis ovato-lanceolatis obtusiusculis. Torus distinctè foveolatus. Stam. filam. longiuscula hirsuta, antheræ elongatæ puberulæ vix acutæ. Ovarium ovatum villosum. Stylus basi villosus. Stigmata 3. — Verosimiliter genus proprium.

5. S? BERTERIANA (Chois. mss.) 5 in Sancto-Domingo. Rheedea lateriflora Bertl. in litt. Folia alterna ovalia utrinquè attenuata, petiolo basi et apice tumidulo. Racemi axillares pauciflori petiolo subbreviares. Flores parvi. Fruct. ign. (v. s.)

†. Genera Tiliaceis affinia, plurima non satis nota.

- XV. ABLANIA. Aubl. guian. 1. p. 585. t. 234. — Trichocarpus Schreb. gen. n. 923.

Cal. 4-5-partitus (an per æstiv. valvatus?) Pet. o. Stam. ∞ hypogyna, filam. liberis, antheris subrotundis. Ovarium ovatum. Styli 2 bifidi. Capsula 4-valvis, 1-loc., valvis extus setis rigidis hispidis. Semina ∞ placentæ liberæ affixæ arillata. — Genus non satis notum affine Sloaneæ et Bixæ.

1. A. GUIANENSIS (Aubl. l. c. t. 234.) 5 in sylvis Guianæ. — Lam. ill. t. 479. A. laurifolia Pers. ench. 2. p. 81. Trichocarpus laurifolia Willd. sp. 2. p. 1224.

XVI. GYROSTEMON. Desf. mem. mus. 6. p. 16 et 8. p. 115.

Flores dioici. Cal. patens 6-7-lobus. Pet. o. ♂ Stam. ∞ in gyros concentricos disposita, filam. o, antheris recept. nudo impositis bilocularibus sub 4-loc. ♀ ovaria 18 20 monostyla verticillata. Carpella capsularia totidem membranacea bivalvia 1 sperma. Semen incurvum, transv. striatum, axi fructus centrali libero supernè adnatum. Embryo incurvus in albumine corneo, radic. inferâ, cotyl. linearibus parallelis. — Frutices è Novâ-Holl. glabri ramosissimi — An Malvaceis, an Euphorbiaceis affinius?

1. G. RAMULOSUM (Desf. mem. 6. p. 17. t. 6.) foliis linearibus subsessilibus. 5 in Novæ-Hollandiæ ins. Sterilibus dictis. Frutex siccus Ephedræ facie. (v. s.)
2. G. COTINIFOLIUM (Desf. mem. 8. p. 116. t. 10.) foliis ovato-rotundatis petiolatis. 5 in Novâ-Holl. loco dicto Baye des Chiens Marins. (v. s. in h. Mus. Par.)

XVII. CHRISTIANA. — Gen. nov. Brown. cong. p. 9.

Cal. 3-lobus. Petala 5. Stamina indefinita. Carpella capsularia 5, 1-sperma, basi tantum connexa. — Genus vix notum, certè Tiliaceum, ex cl. Brown, affine Ventenatiæ.

1. C. AFRICANA. — in Africâ ad Congo flumen. Descr. nondum edita. — nomen dedi ex cl. et infortunato detectore Christiano Smith.

XVIII. ALEGRIA. fl. mex. ic. ined.

Involucellum patens 10-12-phyllum, foliolis lanceolatis acutis. Cal.

5-sepalus, sepalis per æstiv. valvatis, demum apice revolutis. Pet. 5 plana obovalia. Stam. ∞ , filam. liberis aut basi subcoalitis, antheris subrotundis. Fila sterilia ∞ inter stam. et petala. Ovar. ovatum. Stylus 1. Stigmata 5, subrotunda, approximata. Caps. 5-angul., 5-valvis, 5-locul., valvis medio intus septiferis. Semina in alam expansa. — Genus involucello et fructu ad Malvaceas, staminum filamentis liberis et filis sterilibus ad Tiliaceas accedit.

1. *A. CANDIDA* (fl. mex. ic. ined.) 5 Arbor speciosissima, foliis brevè petiolatis ovatis serratis, floribus albis Sparmanniæ æmulis, magnitudine Rosæ.

XIX. LUHEA. Willd. act. soc. nat. scrut. berol. 3. p. 409. t. 5.

Involucellum breve 9-phyllum. Calyx 5-partitus. Pet. 5. Stam. ∞ , filamentis subulatis basi pilosis et infra connatis in phalanges 5 quibus basi adnati sunt processus penicilliformes, antheris subrotundis. Stylus crassus. Stigma truncatum. Fruct. ign. — Car. ex Willd. et Juss. mss. Affinis videtur hinc Alegriæ, illinc Grewiæ.

1. *L. SPECIOSA* (Willd. l. c. et sp. 3. p. 1434.) 5 Folia alterna 3-nervia venosa subcordata obtusa inæqualiter dentata. Petioli breves pubescentes. Racemi terminales pauciflori. Flores albi.

XX. VATICA. Lin. mant. 2. p. 152.

Cal. 5-fidus, lobis acutis per æstiv. forsan valvatis? Pet. 5 obovato-oblonga æstiv. contorta. Antheræ 15 sessiles ovatæ 4 locales, 3 antè petalum quodque. Ovarium 5-angulare. Stylus 1. Stigma sub-3-lobum? Caps? 3-locul? loculis 1-spermis? (Smith.) — Genus, ob caracteres non satis notos et forsan inter ordines plures ambiguos, in serie dubium.

1. *V. CHINENSIS* (Lin. l. c.) 5 in Chinâ. — Smith ic. ined. t. 36. Lam. dict. 8. p. 418. ill. t. 397.

XXI. ESPERA. Willd. act. soc. nat. cur. berol. 3. p. 449.

Cal. 4-partitus patens. Pet. 6 triplò longiora persistentia. Stam. ∞ , filamentis capillaribus, antheris subrotundis. Stylus 1. Stigma 1. Capsula oblonga 4-6-alata 4-6-locularis, loculis 1-spermis. Semina subrotunda hirsuta. — Affinis Humiriæ et Sloanæ ex Juss. mss.

1. *E. CORDIFOLIA* (Willd. l. c.) ♂ Frutex foliis alternis petiolatis cordatis integerrimis, floribus paniculatis terminalibus.

XXII. WIKSTROEMIA. Schrad. gætt. anz. 1821. p. 710.

Cal. 5-sepalus inæqualis deciduus. Cor. 5-petala. Stam. ∞ toro inserta. Ovarium 1. Stigmata plura. Capsula 10-locularis 10-valvis, valvulis introflexis et margine adfixis columellæ cylindraceæ. Semina valvularum marginibus adfixa. — Genus Tiliaceis affine ex cl. Schrader, sed ob calycem inæqualem et ideò forsan non valvatum meo sensu locus dubius.

1. *W. FRUTICOSA* (Schrader. l. c.) ♂ in Brasiliâ. — 10-petalis, flores albi terminales. Fructus pollicares Rhododendri facie.

XXIII. BERRYA. Roxb. cat. calc. p. 42.

Cal. sepala 5 antè floresc. coalita, dein irregulariter valvatum de-

hiscentia, extus tomentosa, intus colorata, decidua. Pet. 5 oblonga. Stem. ∞ , filam. liberis (aut imâ basi subconnexis?), antheris parvis bilocularibus. Ovarium 1 sessile 3-alatum. Stylus 1. Stigma trigonocapitatum. Capsula subrotunda, 3-locularis, 3-valvis, 6-alata, valvis nempe alas 2 horizontales dorso gerentibus, intus medio septiferis. Semina in quoque loculo 2 ovato-globosa grossa pilis rigidis undique tecta (in meis intus vacua).

1. B. AMOMILLA (Roxb. l. c.) 5 in Zeylonâ. Rami teretes glabri. Folia alterna exstipulata petiolata ovata acuminata integra glabra basi 7-nervia. Panicula terminales. (v. s. ex hort. Calc. comm. à cl. Wallich.)
-

Ordo XXVIII. ELÆOCARPEÆ. *Juss. ann. mus.* 11. p. 223.

Calyx extus nudus, 4-5-sepalus, sepalis per æstivationem valvatis. Petala 4-5, sepalis alterna, hypogyna, apice lobata fimbriatave. Torus glandulosus subexsertus. Stam. 15-20, filamentis brevibus liberis, antheris elongatis, filiformi-tetragonis, bilocularibus, loculis poro oblongo apice dehiscentibus. Ovarium pluri-loculare. Stylus 1. Semina in loculis 2 aut plura. Albumen carnosum. Embryo erectus, cotyledonibus planis foliaceis. — Frutices arboresce. Folia alterna simplicia.

Ordo Tiliaceis valdè affinis, et tantum distinctus petalis lobatis et antheris apice biporosis.

I. ELÆOCARPUS. *Lin. gen.* 663.

Cal. 5 sepalus. Pet. 5 apice lacera. Antheræ hinc setigeræ. Drupa nucleo sæpius rugoso-sulcato, intus 5-loc. aut abortu 1-locularis.

1. *E. serratus* (Lin. sp. 734.) foliis lanceolato-ellipticis serratis, racemis axillaribus, fructibus globosis, nucleo rugoso-sulcato. 5 in Indiâ or. — Rumph. amb. 3. t. 101. Burm! zeyl. 39. t. 40. Ganitrus spherica Gaertn. fr. 2. p. 271. t. 139. f. 6. (v. s.)
2. *E. PERIM-KARA*, foliis ovato-lanceolatis serratis, floribus racemosis, fructibus ovatis, nucleo læviusculo. 5 in Malabariâ Rheed. h. mal. 4. t. 24. racemi in icone terminales videntur, sed verosim. pictoris incuriâ. †.
3. *E. CYANÆUS* (Sims bot. mag. t. 1737.) foliis oblongo-lanceolatis serratis reticulato-venosis, racemis axillaribus confertifloris, fructibus subglobosis, nucleo læviusculo. 5 in Novâ-Hollandiâ. Flores albi. Drupæ ceruleæ. *E. reticulata* Smith in Rees cycl. ex Ker. bot. reg. t. 657. *E. cyanæus* herb. amat. t. 237. (v. s.)
4. *E. MONOCERAS* (Cav. ic. 6. p. 1. t. 501.) foliis lanceolatis basi cuneatis apice serratis, racemis axillaribus. 5 in ins. Luzon et ad oppidum Bannoâ. Ovarium bilocul. ? Fructus ignotus. Antheræ apice 1-setæ. Flores ferrugineo-rubri.
5. *E. INTEGRIFOLIUS* (Lam. dict. 2. p. 604.) foliis ovato-oblongis obtusis integerrimis, racemis axillaribus folio longioribus, floribus 4-fidis. 5 in ina. Mauritiî.
6. *E. OBLONGUS* (Smith in Rees cycl. n. 2.) foliis ovato-oblongis acutis integerrimis, drupis ovato-oblongis 1-spermis. 5 in insul. Moluccanis. Ganitrum oblongum Rumph. amb. 3. p. 161. t. 102. Gaertn. fr. 1. p. 202. t. 43. Flores ex Rumphio *E. serrato* similes. An satis à priori differt. ?
7. *E. INTEGRISSIMUS* (Lour. coch. 1. p. 412.) foliis lanceolatis integerrimis, floribus congestis axillaribus. 5 in Cochinchinâ. Flores aurei suaveolentes. Drupæ nigre subovatæ.
8. *E. SYLVESTRIS* (Poir. suppl. 2. p. 704.) foliis ovato-lanceolatis serratis, spiciis subterminalibus, glandulis tori bilobis, drupis 1-spermis. 5 in Cochinchinæ sylvis. Adenodus sylvestris Lour. fl. coch. 1. p. 361. Rami patentæ. Flores albi rubro varii.
9. *E. NITIDUS* (Mal. misc. 1. n. 2. p. 41.) foliis ovato-lanceolatis serratis, racemis axillaribus folia brevioribus, staminibus 15, nuce 5-loculari, loculis 4 plerumque abortivis. 5 in Pulo-Pinang. Flores albi. Semina albuminosa, cotyl. planis, radiculâ superâ? An diversus ab *E. sylvestri*?
10. *E. ELLIPTICUS* (Smith in Rees cycl. n. 3.) foliis lævibus eglandulosis, calycibus acutis, ovario globoso ad basin 5-squamuloso. — Lin. mant. 2. p. 401. in adnot.

II. ACERATIUM. †.

Cal. 5-sepalus. Pet. 5. apice simbriato-lacera, ungue lato villis ciliato. Antheræ puberulæ setis terminalibus destitutæ. Stylus 1. Fructus ignotus.

1. *A. OPPOSITIFOLIUM*. 5 in Amboinâ. Folia opposita elliptico-oblonga in

nervis subpubescentia, serraturis mucronatis raris instructa. Pedunculi terminales triflori. (v. s.)

III. DICERA. Forst. gen. t. 40.

Cal. 4-5 sepalus. Pet. 4-5 apice triloba. Antheræ 12-20 lineares setis 2 superatæ. Capsula (in *D. dentatâ* generis typo) 2-locul., loculis oo-spermis.

1. *D. DENTATA* (Forst. l. c. p. 80.) foliis alternis oblongis supernè dentato-serratis, racemis axillaribus, floribus 1-gynis, capsulis 2-locul., loculis oo-spermis. 5 in Novâ-Zelandiâ *Elæocarpus dentatus* Vahl symb. 3. p. 67.
2. *D. SERRATA* (Forst. l. c.) foliis oppositis ovatis duplicato-serratis, racemis compositis, floribus 4-gynis, bacca 4-locul., loculis 2-spermis. 5 in Novâ-Zelandiâ. *Elæocarpus Dicera* Vahl. symb. 3. p. 67. An genus proprium? An *Friesiæ* species?
3. *D. CRASPEDUM* (Gmel. syst.) foliis oblongis crenatis apice acuminato reflexo, spicis subterminalibus congestis, floribus 1-gynis, bacca 1-locul. polysperma. 5 in sylvis Cochinchinæ. *Craspedum tectorium* Lour. coch. 2. p. 411. *Elæocarpus tectorium*. Poir. suppl. 2. p. 104. An genus proprium?

IV. FRIESIA. (1)

Cal. 4-partitus. Pet. 4 apice triloba. Antheræ 12 cordato-oblongæ acuminatæ apice dehiscentes. Bacca sicca substipitata indehiscens 2-4-sulca 2-4-loc., loculis 2-spermis.

1. *F. PEDUNCULARIS*. 5 ad cap. Van-Diemen. Folia opposita lanceolata serrata; pedicelli axillares 1-flori patentes subnantes. *Elæocarpus peduncularis* Labill. nov. holl. 2. p. 15. t. 155.

V. VALLEA. Mut. in Linn. suppl. 266.

Cal. 5-sepalus. Pet. 5 trifida. Glandula tori ovarium annulatim ambiens. Squamæ 5 sub unguibus petalorum adnatæ. Stam. 30-40, antheris muticis apice dehiscentibus. Stylus sulcatus. Caps. 4-5-valvis 4-5-locul., valvulis muricatis patentibus, loculis 2-spermis. — Arbusculæ, stipulis reniformibus, foliis alternis cordatis petiolatis integris.

1. *V. STIPULARIS* (Mutis in Linn. f. suppl. 266.) ramis pedicellis petiolisque glabris, foliis cordatis obtusis subtus in axillis venarum et ad basin villosobarbatis, stipulis breviter petiolatis. 5 ad Santa-Fé de Bogota. H. B. et Kunth nov. gen. am. 5. p. 350. t. 489. *. Stipulæ reniformes. (v. s.)
2. *V. CORDIFOLIA* (Ruiz. et Pav. syst. fl. per. 132.) ramis pedicellis petiolisque puberulo-hirsutis, foliis cordatis acutis subtus præsertim ad basin subhirsutis, stipulis breviter petiolatis. 5 in Peruvix memoribus. (v. s.)
3. *V. PUBESCENS* (H. B. et K. nov. gen. am. 5. p. 350.) ramis pedicellis petiolisque ferrugineo-tomentosis, foliis cordatis obtusis subtus fuscescenti-pubescentibus, stipulis sessilibus. 5 prope Santa-Fé de Bogota.

VI. TRICUSPIDARIA. Ruiz. et Pav. syst. p. 112. prod. t. 36. — *Tricuspis Pers. ench. 2. p. 9.*

Cal. 5 dentatus. Pet. 5. tricuspidata. Glandula tori annularis decagona. Antheræ 15 muticæ apice dehiscentes. Caps. 3-locul., 3-valvis, valvulis septiferis. Semina pauca.

1. *T. DEPENDENS* (Ruiz. et Pav. l. c.) 5 in R. Chilensis memoribus et inundatis. Folia opposita, ovato-oblonga, serrata. Pedicelli axillares, solitarii. (v. s.)

VII. DECADIA. Lour. coch. 1. p. 385.

Cal. 3-partitus, persistens. Pet. 10 subovata subserrata. Stam. 30 petalorum basi insistentia. Antheræ subrotundæ. Stylus filiformis. Drupa nuce 5-locul. — An stamina calyci inserta et tunc ad Rosaceas forsân referenda?

1. *D. ALUMINOSA* (Lour. l. c.) 5 in sylvis Cochinchinæ. — Rumph. amb. 2. t. 100. Folia alterna lanceolata serrata glabra. Racemi subsimplices. Flores albi.

(1) *Friesia Spreng.* est *Crotanopsis Michx.*

Ordo XXIX. CHLENACEÆ. *Pet. Th. hist. veg. afr. austr.*
p. 46.

Involucrum 1-2-florum, persistens, formâ et consistentiâ varium. Calyx 3-sepalus parvus. Pet. 5-6, hypogyna, basi latiora, interdum imâ basi subcoalita. Stam. nunc plurima, nunc rariùs 10, filamentis basi in tubulum subcoalitis aut petalorum tubo adnatis, antheris subrotundis adnatis aut liberis bilocularibus. Ovarium unicum intùs 3-loculare. Stylus 1 filiformis. Stigma triplex. Capsula 3-locul. aut abortu 1-locul. Semina in loculis solitaria aut plurima, centro adfixa, inversa. Albumen carnosum (ex Juss.), aut corneum (ex P. T.) Embryo viridis centralis. Cotyledones foliaceæ undulatæ.

Arbusculæ aut Frutices Madagascarienses. Folia alterna, stipulacea, penninervia, integra. Stipulæ deciduæ. Flores paniculati aut racemosi.

Ordo affinis, suadente cl. Petit-Thouars, Malvaceis ex florum involucro et staminum monadelphiâ, et, sentiente cl. Jussieu, Ebenaceis aut Symplocëis ex petalis basi coalescentibus et seminibus albuminosis.

I. SARCOLÆNA. *Pet. Th. l. c. p. 37.*

Involucrum carnosum urceolatum 5-dentatum ferrugineo-tomentosum. Cal. intrâ involucrum. Pet. 5 basi in tubum conniventia. Stam. ∞ basi tubi inserta. Antheræ terminales. Ovarium 3-loc., loculis 2-spermis. Capsula intrâ involucrum ampliatus baccatum intùs villis prurientibus instructum. Albumen tenue. — Rami decumbentes. Folia juniqra plicata undè adulta quasi 5-nervia videntur.

1. *S. GRANDIFLORA* (P. Th! l. c. p. 40. t. 9.) paniculâ pauciflorâ, foliis acutis subtùs ferrugineo-tomentosis, involucro depresso pilis scabro. 5 in Madagascar. (v. s.)
2. *S. MULTIFLORA* (P. Th! l. c. p. 40. t. 10.) paniculâ confertâ, foliis acutis subtùs in nervis tantùm subpilosis, involucro pilis scabro trilobo. 5 in Madagascar. (v. s.)
3. *S. ERIOPHORA* (P. Th. l. c. p. 40.) paniculâ pauciflorâ axillari, foliis obtusis plicis longitudinalibus destitutis, involucro pilosissimo. 5 in Madagascar.

II. LEPTOLÆNA. *Pet. Th. hist. veg. afr. austr. p. 41.*

Involucrum subcarnosum cylindricum parvulum. Sepala 3 involucro longiora. Pet. 5 in tubum conniventia. Stam. 10. Antheræ dorso adfixæ. Ovarium 3-locul., loculis 2-spermis. Capsula involucro scabro cincta, abortu 1-loc. 1-sperma. — Arbor elegans. Folia undulata glabra. Flores corymbosi.

1. *L. MULTIFLORA* (P. Th! l. c. p. 41. t. 11.) 5 in Madagascar ad Foulpointe. (v. s.)

III. SCHIZOLÆNA. *Pet. Th. hist. veg. afr. austr. p. 43. gen. nov. n. 55.*


Involucrum biflorum, non carnosum, fimbriatum, demùm ampliatus

laciniatum viscosum connâvens. Sep. 3. Pet. 5. Stam. ∞ , filam. tenuibus, antheris adnatis. Capsula intrâ involucrium inclusa, 3-loc., loculis ∞ -spermis. — Arbusculæ elegantes. Folia ovata glabra. Flores paniculati vel racemosi.

1. *S. ROSA* (P. Th. l. c. p. 43. t. 12.) paniculis terminalibus, involucrio amplissimo irregulariter fissio. 5 in Madagascar. (v. s.)
2. *S. ELONGATA* (P. Th. l. c. p. 44.) paniculis terminalibus, involucrio 5-lobo capsulâ paulò longiore. 5 in Madagascar.
3. *S. CAULIFLORA* (P. Th. l. c. p. 44.) floribus racemosis, racemis è trunco aut ramis majoribus ortis. 5 in Madagascar.

IV. RHODOLÆNA. Pet. Th. hist. veg. afr. austr. p. 47. gen. nov. n. 56.

Involucrium bracteolis 2 calyci adpressis constans. Sepala 3 concava carnosâ glutinosa. Pet. 6 urceolata magna, per æstivationem spiraliter contorta. Stam. ∞ petalis breviora, basi urceolo brevi juncta, antheræ 4-angulæ dorso insertæ. Ovarium 3-locul. ∞ -spermum. — Frutex scandens. Folia ovalia acuta nervo medio mucronata. Pedunculi axillares nudi apice biflori. Flores purpurei, maximi.

1. *R. ALTIVOLA* (P. Th. l. c. p. 48. t. 13.) 5  in Madagascar. — Fructus ignotus.

†. Genus *Chlenaceis* affine.

V? HUGONIA. Lin. gen. n. 831. Gaertn. fruct. 1. p. 281. t. 58.

Calyx extus nudus, persistens, 5-partitus, sepalis nempe 5 imâ basi coalitis inæqualibus æstivatione quincunciali imbricatis. Petala 5 sepalis alterna unguiculata æstivatione contortâ. Stam. 10, filamentis basi in urceolum monadelphis, dein liberis filiformibus; antheræ ovatæ aut didymæ (Lin.) Ovarium subrotundum. Styli 5, distincti. Drupa carnosâ fovens carpella 5 ex (Cav. 10 ex Gaertn.) 1-sperma, inter se adhærentia. Semen in loculo quoque pendulum. Embryo inversus in axi albuminis carnosi, radiculâ superâ brevi, cotyledonibus planis foliaceis. — Frutices, foliis alternis, prope flores sæpè congestis et suboppositis. stipulis 2 subulatis, pedunculis axillaribus 1-floris nonnullis abortientibus in spinas uncinatas mutatis!

Genus *Malvaceis* *Buttneriaceis*ve affinis, at Calyx non valvatus sed imbricatus. An inter *Bombaceas* potius recensendum, sed antheræ à Linnæo didymæ dictæ verosimiliter hi nec uniloculares sunt? Ergo ad calcem *Chlenacearum* etiamsi styli plures dicti et involucrium nullum, donec lux fiat, adjungo.

1. *H. MYSTAX* (Lin. spec. 944.) foliis ovalibus glabris integerrimis, spinis suboppositis uncinatis. 3 in Zeylonâ, Malabariâ. — Rheed. mal. 2. p. 29. t. 29. (v. s.)
2. *H. SEARATA* (Lam. dict. 3. p. 149.) foliis ovali-oblongis serratis, adultis utrinquè glabris, spinis suboppositis uncinatis. 5 in ins. Mauritiî. *H. Mystax* Cav. diss. 3. p. 177. t. 73 f. 1. excl. syn. In speciminibus meis spinæ nullæ. (v. s.)
3. *H. TOMENTOSA* (Cav. diss. 3. p. 178. t. 23 f. 2.) foliis oblongis serratis utrinquè tomentosâ, spinis nullis? 5 in ins. Mauritiî. — Lam. dict. 3. p. 150.

Ordo XXX. TERNSTROEMIACEÆ. DC. *mem. soc. h. nat. gen. vol. 1.* — Ternstroemiæ. *Mirb. bull. philom. 1813. p. 381.*

Calycis Sepala 5 concava inæqualia coriacea imbricata obtusa persistentia, sæpè bracteolis 2 stipata. Pet. 5 disco hypogynò inserta, nunc libera, nunc basi subcoalita. Stam. ∞ hypogyna, petalis inter se coalitis subadnata, filam. subulatis brevibus, antheris erectis bilocularibus. Ovarium ovatum. Styli 2-5 liberi aut plus minùs coaliti. Fructus ovato-globosus intùs radiatim divisus in loculos tot quot sunt styli, nunc sicco-baccatus indehiscens, nunc capsularis dehiscens. Semina ∞ ad placentam centram adfixa, nunc arcuata, nunc ovata aut angulata, in tribu primâ donata albumine carnoso parco aut nullo, embryone tereti axili arcuato replicatove, cotyledonibus oblongis, plumulâ inconspicuâ, radiculâ longâ hilo adversâ, in ceteris non satis nota. — Arbores fruticesve. Folia alterna exstipulata penninervia coriacea indivisa. Pedunculi axillares aut terminales.

Ordo ex structurâ calycis ad Camellieas accedit, ex floribus à Symploceis Ebenaceisve parùm recedit, sed ob fructus non satis notos adhuc in serie subincertus et forsàn in posterum dividendus.

Trib. I. TERNSTROEMIEÆ.

Calyx basi bibracteolatus. Petala inter se basi coalita sepalis opposita. Antheræ adnatæ. Stylus 1. Stigma simplex. Ad hanc tribum pertinent characteres spermici suprâ indicati.

I. TERNSTROEMIA. *Lin. f. suppl. 39.* — Toanabo. *Aubl. guian. 1. p. 569.*

Calyx 5-6-sepalus basi bibracteolatus. Petala 5-6. Stam. ∞ duplici serie. Antheræ oblongæ glabræ. Bacca subexsucca 2-locul., loculis 3-4-spermis. — Pedicelli 1-flori, basi squamulâ instructi, subflore bibracteolati. — Species ferè omnes Americanæ.

1. T. BREVIFEX (DC. l. c.) foliis integerrimis obovatis subemarginatis, pedicellis axillaribus flore vix longioribus. ζ in Amer. calidiore. T. meridionalis Mutis? (v. s.)
2. T. PEDUNCULARIS (DC. l. c.) foliis integerrimis ovato-oblongis obtusis, pedicellis lateralibus flore triplo longioribus. ζ in ins. Caribæis. T. meridionalis Swartz obs. 207? (v. s.)
3. T. LINEATA (DC. l. c.) foliis integerrimis oblongis subacutis, pedicellis lateralibus cernuis flore vix longioribus. ζ in Mexico. T. meridionalis fl. mex. ic. ined.
4. T. ELLIPTICA (Swartz prod. 81.) foliis integerrimis ellipticis acutis, pedicellis lateralibus petiolo duplo longioribus. ζ in ins. Caribæis. (v. s. sine fl.)
5. T. PUNCTATA (Swartz prod. 81.) foliis integris oblongis subemarginatis margine punctato-denticulatis, pediculis axillaribus petiolo multo longioribus. ζ in Guianæ sylvis. Toanabo punctata Aubl. guian. 1. p. 571. t. 228.

6. *T. CLUSIEFOLIA* (H. B. et Kunth nov. gen. am. 5. p. 207. t. 463. f. 1.) foliis oblongis obtusis integerrimis subtus nigro-punctulatis, pedunculis axillaribus petiolo paulò longioribus. 5 in Andibus Popayanensium.
7. *T. DENTATA* (Swartz. prod. 81. non Mirb.) foliis dentato-serratis ovali-oblongis acuminatis, pedunculis axillaribus lateralibusque petiolo paulò longioribus. 5 in Guianæ sylv. *T. dentata* Aubl. Guian. 1. p. 569. t. 227. (v. s.)
8. *T. SALICIFOLIA* (DC. l. c.) foliis serrulatis oblongis acuminatis subaveniis, pedicellis 2-3 axillaribus petiolo ferè brevioribus. 5 in sylv. Guadalupe. Bertero (v. s.)
9. *T. VENOSA* (Spreng. neu. entd. 1. p. 162.) foliis serrulatis oblongis venosis, pedicellis adgregatis axillaribus petiolo ferè brevioribus. 5 in Brasiliâ.
10. *T. P. QUINQUEPARTITA* (Ruiz et Pav. syst. 180.) foliis obsolete serrulatis obovatis, corollis 5-partitis. 5 in Andium sylvaticis frigidis. Stam. pentadelphæ. An genus diversum?
11. *T. P. GLOBIFLORA* (Ruiz et Pav. syst. 180.) foliis integerrimis oblongis, corollis globosis 5-dentatis. 5 in Andium sylvaticis frigidis. Stam. pentadelphæ. An genus proprium?
12. *T. RUBIGINOSA* (mal. misc. 1. n. 2. p. 39.) foliis ovatis spinuloso-serratis subtus incanis, floribus lateralibus et axillaribus fasciculatis monadelphis, pedunculis calycibusque glanduloso-pilosis, fructu 3-locul. 5 in Sumatrâ.
13. *T. PENTAPETALA* (mal. misc. 1. n. 2. p. 40.) foliis obovato-lanceolatis spinuloso-denticulatis glabris, floribus lateralibus fasciculatis, pedunculis glabris, fructu 3-loculari. 5 in Pulo-Pinang. An hæc et prior sint veræ Ternstroemiæ dubito?
14. *T. P. CORYMBOSA* (Smith in Rees. cycl. 35.) foliis oppositis ellipticis acuminatis integris, paniculis furcato-corymbosis multifloris terminalibus, calycibus ebracteolatis. 5 in Guianâ.

Trib. II. *FREZIEREÆ*.

Cal. basi bibracteolatus. Pet. inter se libera, sepalis alterna Antheræ adnatæ. Styl. 1. Stigm. 2-5 distincta. Semina aptera, albumine carnoso, embryone subcurvato.

II. *CLEYERA*. Thunb. fl. jap. p. 12.

Cal. 5-sepalus 2-bracteolatus. Pet. 5 libera. Antheræ retrorsum setoso-hispidæ stylus filiformis. Stigmata 2-3. Bacca 2-3-locularis. — Habitus Ternstroemiæ. Species omnes Asiaticæ.

1. *C. JAPONICA* (Thunb. fl. jap. p. 224.) foliis oblongo-lanceolatis aveniis apice serrulatis. 5 in Japoniâ prope Nagasaki. — Kœmpf. am. 5. p. 774. ic. Ternstroemia Japonica Thunb. act. soc. Lin. 2. p. 335. (v. s.)
2. *C. OCHNACEA* (DC. l. c.) foliis ovali-oblongis utrinquè acutis supernè venosis integris.
 - a. *Kœmpferiana*. — Sakaki kœmpf. am. 777. Banks ic. Kœmpf. t. 33. 5 in Japoniâ.
 - β. *Wallichiana*. 5 in Napauliâ. (v. s. comm. à cl. Wallich.)

III. *FREZIERA*. Swartz. fl. ind. occ. 2. p. 971. — Eroteum. Swartz. prod. 85.

Cal. 5-sepalus 2-bracteolatus. Petala 5 basi latiora. Antheræ glabræ subcordatæ. Stylus apice 3-5-fidus. Bacca 3-5-locularis. — Arborea Americanæ Lauri habitu, pedicellis axillaribus.

1. *F. THÆOIDES* (Swartz. fl. 972.) foliis ovato-lanceolatis serrulato-dentatis utrinquè glabris, pedicellis solitariis unifloris. 5 in montibus Jamaicæ. Eroteum thæoides Swartz. prod. 85. (v. s.)
2. *F. UNDULATA* (Swartz. fl. 974.) foliis elliptico-lanceolatis acuminatis serratis utrinquè glabris, floribus axillaribus confertis. 5 in sylv. ins. Caribæarum. *E. undulatum* Swartz. prod. 85. (v. s.)

3. *F. NERVOSA* (Humb. et Bonpl. pl. eq. 1. p. 31. t. 9.) foliis lanceolatis dentatis supernè glabris subtùs pubescentibus, pedicellis plurimis fasciculatis. 5 in frigidis prov. Pasto.
4. *F. SERICEA* (Humb. et Bonpl. l. c. p. 29. t. 8.) foliis elliptico-lanceolatis acuminatis serrulatis subtùs argenteis, floribus 2-3 axillaribus sessilibus. 5 in frigidis inter Quito et Popayan.
5. *F. CHRYSOPHYLLA* (Humb. et Bonpl. l. c. p. 27. t. 7.) foliis lanceolato-oblongis acuminatis supernè glabris subtùs tomento sericeo aureo villosissimis, pedicellis axillaribus paucis brevibus. 5 in frigidis Andium Peruviae ad Popayan.
6. *F. CANESCENS* (Humb. et Bonpl. l. c. p. 25. t. 6. nov. gen. am. 5. p. 211. t. 463. f. 2.) foliis elliptico-oblongis serrulatis subtùs tomentosis incanis, pedicellis 1-2 axillaribus. 5 in Peruviae Andibus.
7. *F. RETICULATA* (Humb. et Bonpl. l. c. p. 23. t. 5.) foliis ovato-lanceolatis serratis subtùs tomentosis, pedunculis 3-5 axillaribus fasciculatis. 5 in Andibus Peruviae prope Almager. (v. s.)

IV. *EURYA*. Thunb. fl. jap. p. 11. Brown. pl. chin. diss. p. 7.

Flores polygami. Cal. 5-sepalus 2-bracteolatus. Pet. 5 basi subconcreta subrotunda. Antheræ glabræ tetragonæ. Stylus apice 3-5-fidus. Bacca (Br.) Capsula (Th.) 3-5-locul. — Frutices Asiaticæ.

1. *E. JAPONICA* (Thunb. jap. p. 191. t. 25.) ramulis etiam junioribus glabris, foliis ellipticis acutis, floribus axillaribus. 5 in Japoniâ prope Nagsaki. (v. s.)
2. *E. CHINENSIS* (Brown. l. c. cum icon.) ramulis ultimis pubescentibus, foliis cuneato-ovalibus, floribus axillaribus. 5 in Chinæ prov. Kiang-si et Quang-tong, in campis et collibus.
3. *E. MULTIFLORA* (DC. l. c.) ramulis pilosiusculis foliis elliptico-oblongis, floribus secùs ramulos sæpius infrâ folia fasciculatis. 5 in Napauliâ. Wallich. (v. s.)
4. *E. ACUMINATA* (DC. l. c.) ramulis villosis, foliis elliptico-oblongis acuminatis, floribus axillaribus paucis. 5 in Napauliâ. Wallich. (v. s.)

V. *LETTSOMIA*. Ruiz et Pav. prod. fl. per. p. 77. t. 14.

Cal. 5-sepalus 2-bracteolatus. Petala 5 infernè equitantia, interiora angustiora. Stylus brevissimus. Stigm. 3-5. Bacca 3-5-locularis.

1. *L. TOMENTOSA* (Ruiz et Pav. syst. 134.) foliis lanceolatis integerrimis subtùs tomentoso-sericeis, baccis 5-locularibus. 5 in Peruviae nemoribus.
2. *L. LANATA* (Ruiz et Pav. syst. 135.) foliis lanceolatis obsolete serrulatis lanatis, baccis 3-locularibus. 5 in Peruviae nemoribus.

Trib. III. SAURAUJÆ.

Calyx extùs ebracteolatus. Petala sepalis alterna, basi plus minùs inter se in corollam monopetalam coalita. Stam. ∞ imæ corollæ adhærentia. Antheræ dorso insertæ incumbentes nec adnatæ. Styli 3-5 ex ovario distincti. — Semina non satis nota.

VI. *SAURAUJA*. Willd. nov. act. soc. nat. cur. berol. 3. p. 406. t. 4.

Calyx ebracteolatus. Petala 5 ad medium inter se coalita. Styli 3-5. Capsula 5-locularis. — Arbores fruticesve albifloræ Americanæ aut Asiaticæ.

1. *S. EXCELSA* (Willd. l. c.) foliis oblongo-obovatis acutiusculis integerrimis suprâ scabridis subtùs ad venas piloso-hirtis, pedunculis fusco-villosis longis apice trichotomo-paniculatis. 5 in sylvis montanis prov. Caracasana.
2. *S. VILLOSA* (fl. mex. ic. ined. sub Davya.) foliis ellipticis utrinquè acuminatis à medio ad apicem hinc indè serratis, subtùs ramulis pedunculisque villosis. 5 in Mexico.

3. *S. SERRATA* (fl. mex. ic. ined. sub Davya.) foliis ellipticis basi attenuatis acutis serratis glabris, ramis petiolis pedunculisque tomento ferrugineo velutinis. 5 in Mexico.
4. *S. NAPAULENSIS* (DC. mem. soc. gen. 1. p. . . .) foliis oblongis utrinque acuminatis serratis, supernè glabris, subtùs in nervis ramulisque strigosopilosis, pedunculis racemosis petiolo multò longioribus. 5 in Napauli. Wallich. (v. s.)
5. *S. LANCEOLATA* (DC. l. c.) foliis oblongo-lanceolatis acuminatis minutissimè serrulatis, adultis glabris, junioribus squamulas rufas in nervis gerentibus, pedunculis axillaribus petiolo subæqualibus apice umbellatis. 5 in Javâ. *Vanalphimia lanceolata*. Lech! mss. (v. s. in herb. Mus. Par.)
6. *S. NUDIFLORA* (DC. l. c.) foliis obovatis subacutis subserratis, adultis utrinque glabris, pedunculis lateralibus 1-floris solitariis. 5 in Javâ. (v. s. in h. Deless.)
7. *S. BRACTEOLATA* (DC. l. c.) foliis ovalibus utrinque attenuatis subserratis glabris, pedunculis lateralibus simplicibus ramosive bracteolas minimas sparsas gerentibus. 5 in Javâ. (v. s. in h. Juss.)
8. *S. BRACTEOSA* (DC. l. c.) foliis ovalibus basi-cordatis acuminatis subserratis, pedunculis axillaribus trichotomis, bracteis oblongis pedicelli longitudine. 5 in Javâ. (v. s. in h. Deless.)
ℓ? punctata, foliis magis serratis, supernè pube sparsâ punctatis.
9. *S. TRISTYLA* (DC. l. c.) foliis ellipticis utrinque acutiusculis tenuissimè serratis glabris, petiolis pediculisque strigoso-squamulosis, pedicellis axillaribus fasciculatis simplicibus trifidisve. 5 in Moluccis. (v. s.)
10. *S. GIGANTEA* (Nor. ic. ined. sub Scaphâ.) foliis ovalibus basi cordatis acuminatis serratis subtùs velutino-rufis, pedunculis axillaribus trichotomis folio dimidiò brevioribus subbracteatis. 5 in Javâ.
11. *S. CRENULATA* (DC. l. c.) foliis obovatis basi cuneatis apice subacutis crenulato-serratis, pedunculis 1-floris in ramis veteribus aggregatis. 5 in Javâ. *Vanalphimia Djinite* Lech. mss. (v. s. in h. Mus. Par.)
12. *SP. CAULIFLORA* (Nor. ic. ined. sub Scaphâ.) foliis oblongis acuminatis aristato-serratis supernè glabris subtùs villosis ferrugineis, pedicellis 1-floris in trunco adgregatis. 5 in Javâ ad ripas Sudani. Fructus baccatus et ideò videtur genus proprium.

VII. *APATELIA*. DC. mem. soc. gen. vol. 1. — Palava. Ruiz. et Pav. prod. fl. per. p. 88. t. 22. non Cav.

Cal. ebracteolatus. Pet. 5 basi imâ inter se subcoalita. Stam. ∞ in phalanges 5 disposita. Antheræ apice biperforatæ. Styli 5. Capsula 5-locul. 5-valvis.

1. *A. LANCEOLATA* (Ruiz et Pav! syst. 181. sub Palavâ.) foliis ovali-lanceolatis utrinque acuminatis serratis subtùs ferrugineis, ramis petiolis pedunculisque strigoso-pilosisimis. 5 in Peruviz montibus circa Chincao. (v. s. in h. Deless.)

ℓ. peduncularis, foliorum limbo basi ovato, pedunculis petiolo quadruplò longioribus. 5 in Peruvâ. (v. s. in h. Deless.)

2. *A. GLABRATA* (Ruiz et Pav! syst. 181. sub Palavâ.) foliis oblongis acutis serrulatis utrinque subglabris, ramis petiolis pedunculisque strigosis minimis subpilosis. 5 in Munzè runcationibus. (v. s. in h. Deless.)
3. *A. BISERRATA* (Ruiz et Pav. syst. 181. sub Palavâ.) foliis obovato-oblongis biserratis, racemis brachiatis, pedicellis trifloris. 5 in Peruviz nemoribus.

Trib. IV. *LAPLACEÆ*.

Calyx extùs ebracteolatus, 3-4-sepalus. Petala sepalorum numerum pluriès superantia. Stam. ∞, filam. liberis, antheris basi affixis. Styli in unicum coaliti. Fructus 5-locul. Albumen carnosocorneum.

VIII. COCHLOSPERMUM. *Kunth diss. malv. p. 6.*

Cal. 5-sepalus persistens demum reflexus; sepalis ovali-oblongis obtusis inæqualibus. Pet. 5 subovata apice emarginata. Stam. ∞ , filam. imâ basi submonadelphis?, antheris acuminatis. Stylus simplex. Capsula ovato-globosa 5 locul. Semina ∞ , subcochleata, lanigera. — Arbores foliis palmatifidis, floribus flavis.

1. *C. Gossypium*, foliorum lobis integerrimis. § in Indiâ orient. *Bombax grandiflorum*. Sonn. voy. ind. 2. p. 235. t. 133. *Bombax Gossypium*. Lin. syst. 517. Cav. diss. 5. p. 297. t. 157. *Bombax Congo* Burm. ind. 145. Folia fere *Gossypii religiosi*, lobis ovato-lanceolatis integerrimis. Flores flavi.
2. *C. serratifolium*, foliorum lobis serratis. § in Mexico et Brasiliâ? *Bombax serratifolium* fl. mex. ic. ined. *Bombax vitifolium* Willd. enum. 720?

IX. LAPLACEA. *H. B. et Kunth nov. gen. am. 5. p. 207. t. 461.*

Cal. 4-sepalus persistens, sepalis suborbiculatis imbricatis. Pet. 9 subæqualia. Stam. ∞ subtriseriata basi petalorum adnata, filamentis liberis, antheris reniformibus bilocularibus. Ovarium sessile 5-loc., loculis 3-spermis. Styli 5 in unum connati. Capsula 5-loc. loculicido-5-valvis lignosa. Semina angulo interno adfixa, pendula, in alam elongatam desinentia. Embryo linearis rectus in axi albuminis.

1. *L. speciosa* (H. B. et Kunth l. c.) 5 rara in sylvis Americæ mer. inter Gonzananam et Lozam Folia oblonga integerrima coriacea glabra. Flores axillares solitarii pedunculati albi.

X. VENTENATIA. *Beauv. fl. ovar. 1. t. 17.*

Calyx 3-lobus concavus deciduus. Pet. 11-12 oblonga basi attenuata obtusa patentia. Stam. ∞ , filam. liberis filiformibus basi petalorum subadnatis, antheris oblongis bilocul. Ovarium ovatum. Stylus stam. longior. Bacca ovato-globosa, longitudinaliter sulcata, 5-locularis, loculis ∞ -spermis. Semina ignota et ideo locus in ord. nat. incertus. Calyx imbricatus nec valvatus à Tiliaceis hoc genus removet. An potius Chlenaceis adnumerandum?

1. *V. GLAUCA* (Beauv. l. c.) § in regno afric. Beniniensi prope Agathou. Arbuscula foliis exstipulatis petiolatis ovatis acuminatis glaucis penninerviis.

Trib. V? GORDONIÆ.

Cal. sepala 5 libera aut inter se coalita. Petala sæpè basi connexa. Stam. ∞ , filamentis imâ basi monadelphis filiformibus, antheris ovalibus oscillatoriis. Carpella 5 plus minus coalita. Styli 5 aut distincti aut basi tantum connexi, ad apicemve fere coaliti. Carpella ideo capsularia nunc subdistincta, nunc in capsulam arcuè coalita, 1-2-sperma, valvis capsulæ mediò septiferis. Semina pauca; albumen nullum. Embryo erectus, radicula oblonga, cotyledonibus foliaceis longitudinaliter plicato-rugosis, plumula inconspicua.

Arbores fruticesve Americanæ rarissimè Asiaticæ. Folia alterna sæpius decidua, ovalia-oblongave integra penninervia exstipulata. Flores affines ex habitu hinc Camelliæ illinc Cydoniæ.

Tribus olim cum Malvaceis Tiliaceisve obcotyledones plicato-rugosas confusus, sed calyce imbricato nec valvato stipulisque nullis distinctus; affinis Camelliæ et Ternstroemiæ, à prioribus cotyledonibus foliaceis corrugatis, à posterioribus albumine nullo

discrepans; sed forsân cum iis classem naturalem constituens. Car. ex Gordoniâ, alia genera ex analogiâ autopsiæ relata.

XI. MALACHODENDRON. Cav. diss. 5. p. 302. f. 2. Juss. gen. 275.

Cal. 1-bractæatus. Pet. 5-6 limbo crenulato. Ovarium 5 sulcatum. Styli 5 à basi liberi. Stigmata capitata. Carpella capsularia 5 connexa 1-sperma. Semina ignota.

1. *M. OVATUM* (Cav. l. c.) 5 in Virginiâ. *Stuartia pentagyna* L'her. stirp. nov. 1. p. 155. t. 74. Arborescens foliis ovatis acutis, floribus solitariis sphe-ssilibus. (v. s.)

XII. STEWARTIA. Cav. diss. 5. p. 303. t. 159. f. 2.

Cal. sepala 5 ad medium ferè coalita, persistentia. Pet. 5. Ovarium subrotundum. Stylus 1 filiformis. Stigma capitatum sub-5-crenatum. Capsula lignosa, 5-loc. 5-valv. loculis 1-2-spermis. Semina aptera ovata lævia, in meis spec. crassa.

1. *S. VIRGINICA* (Cav. l. c.) 5 in Carolinæ et Virginiæ maritimis. *S. Malachodendron* Lin. sp. 982. L'her! stirp. rar. 1. t. 73. — Lam. ill. t. 593. *S. Marylandica* Andr. bot. rep. t. 73. Flores ampli albi, filamentis staminum purpureis, antheris cæruleis (v. s.)

XIII. GORDONIA. Ellis. act. angl. 1770. Cav. diss. 307.

Cal. sepala 5 coriacea rotundata. Petala 5 staminum urceolo subadnata. Stylus 1. Stigm. 5. Capsula 5-loc. 5-valvis, loculis 2-spermis. Semina in alam foliaceam desinentia.

Sect. I. LASIANTHUS.

Petala basi subconnexa. Stamina ferè pentadelphea. Stylus 1.

1. *G. LASIANTHUS* (Lin. mant. 570.) pedicellis axillaribus folio dimidiò brevioribus, foliis oblongis coriaceis glabris serratis, calyce sericeo. 5 à Virginiâ ad Floridam in uliginosis maritim. Cav. diss. 6. t. 161. Sims bot. mag. t. 666. Cat. car. t. 44. *Hypericum Lasianthus* Lin. sp. 1101. (v. s.)
2. *G. WALLICHII*, pedicellis axillaribus folio quintuplò brevioribus, foliis ovalibus acuminatis integris, calyce subciliato. 5 in Napauliâ. *Alabastrum corollinum globosum* calyce brevi cinctum. (v. s.)

Sect. II. HÆMOCHARIS. Salisb.

Petala distincta. Styli 5.

3. *G. HÆMATOXYLON* (Swartz. fl. ind. occ. 2. p. 1109.) pedicellis crassis brevissimis, foliis ovatis acuminatis serratis, stylis 5 distinctis, fructu ovato subpentagono. 5 in Jamaicæ occid. montibus. An genus à Gordoniâ ut *Malachodendron* à *Stewartiâ* distinctum? (v. s.)

Sect. III. LACATHEA. Salisb.

Petala basi connexa. Filamenta libera. Stylus 1.

4. *G. PUBESCENS* (Pursh fl. bor. am. 451.) floribus subsessilibus, foliis obovato-lanceolatis subtus pubescentibus subserratis membranaceis, petalis sepalisque extus subsericeis. 5 in Carolinâ merid. Lois. herb. amat. t. 236. *Lacathea florida* Sal. par. lond. t. 56.

a. *velutina*, foliis oblongis subtus velutinis. *G. pubescens*. L'her! stirp. p. 156. Vent. malm. t. 1. Cav. diss. 6. t. 162. *Franklinia Americana* Marsh. (v. s.)

β. *subglabra*, foliis obovato-oblongis subtus glabresculis. *G. Franklini* L'her! stirp. 1. p. 156. *Franklinia Alatomaha*. Marsh. arb. 48. (v. s.)

Ordo XXXI. CAMELLIÆ. DC. *theor. elem. ed.* 1. 1813.
février. — Theaceæ. Mirb. *bull. phil. decemb.* 1813.

Cal. Sepala 5-7 per æstivationem imbricata, interiora sæpius majora, subconcava, coriacea, decidua. Pet. 5-6-9 sepalis numero æqualia, iis alterna, sæpè basi subcohærentia. Stam. ∞ , filamentis filiformibus imâ basi polyadelphis monadelphisve, antheris ellipsoideis rotundatisve versatilibus. Ovarium 1 ovato-rotundum. Styli 3-6 filiformes plus minus inter se coaliti. Capsula 3-locul. 3-valvis dehiscens, ovulorum plurium abortu 3-sperma, valvis (dehiscentiæ variæ lusu?) nunc medio septiferis, nunc marginibus introflexis. Semina septorum margini centrali adfixa, magna, crassa, pauca. Albumen nullum. Embryo cotyledonibus crassis magnis oleo pingui repletis plano-convexis basi quasi articulatis, radiculâ brevissimâ obtusâ hilo adversâ retractâ, plumulâ vix conspicuâ adscendente. — Frutices arboresve è Chinâ, Japoniâ, Cochinchinâ aut Indiâ ortæ, glabræ, sempervirentes. Flores axillares speciosi.

Ordo ex structurâ floris accedens ad Ternstrœmiaceas, sed semine exalbuminoso discrepans. Forsan tamen hi duo ordines in posterum, si genera nova intermedia accedant, in unicum conjungendi erunt?

I. CAMELLIA. Linn. *gen. n.* 848.

Calyx imbricatus, nempe bracteis sepalisve nonnullis accessoriis cinctus. Stam. basi polyadelphe aut monadelphe. Antheræ ellipsoideæ. Capsula valvis medio septiferis, axim triquetrum liberum post dehiscentiam relinquentibus.

1. C. JAPONICA (Linn. sp. 982.) foliis ovatis acuminatis acutè serratis, floribus terminalibus subsolitariis. $\frac{1}{2}$ in Japoniâ. Cav. diss. 6. t. 160. J. cq. ic. rar. 3. t. 553. Duham. ed. nov. t. 71. Andr. bot. rep. t. 25. Lodd. bot. cab. t. 329 et 455. Herb. amat. t. 43. 44. 45 et 46. Variat fl. simp. et plenis, rubris albis, flavidis variegatisve. (v. v.)
2. C. SESANQUA (Thunb. fl. jap. p. 273. t. 30.) foliis ovato-oblongis obtusè serratis, floribus terminalibus subsolitariis, petalis obcordatis. $\frac{1}{2}$ in Japoniâ. Kœmpf. amœn. 853. Sims bot. mag. t. 2080. Ker. bot. reg. t. 567. Staunt! emb. chin. 2. p. 466. ic. Variat fl. albo, carneo, roseo aut rubro, simplici aut pleno. (v. v.)
3. C. AXILLARIS. (Roxb. ex Ker. bot. reg. t. 349. et Sims bot. mag. t. 2047.) foliis obovato-oblongis serrulatis, superioribus integerrimis, floribus solitariis subsessilibus subaxillaribus. $\frac{1}{2}$ in ins. Pulo-Pinang. Flos albus subflavescens ferè magnit. C. Japonicæ. Styli 4 vix apice summo liberi.
4. C. KISSI (Wall! asiat. res. 13. p.) foliis ovato-oblongis acuminatis acutè serratis, floribus tristylis subsolitariis sessilibus axillaribus subterminalibusve. $\frac{1}{2}$ in Napauliâ. Wallich. Calyces sericeæ. Styli 3 vix basi juncti. (v. s.)
5. C? DRUPIFERA (Lour. coch. 2. p. 499.) foliis ovato-oblongis subcrenatis, fl.

ribus, binis ternatisve terminalibus octopetalis, drupâ 4-locul. 5 in Cochinchinâ. Oleum e seminibus vulgò elicitur. Videtur genus proprium!

II. THEA. Linn. gen. n. 668.

Cal. 5-6-sepalus. Pet. 6-9 imâ basi subcobærentia 2-3-serialia. Stam. basi sublibera, Antheræ subrotundæ. Capsula 3-coeca, septis valvaribus nempè à valvularum marginibus introflexis formatis. (Gaertn. fr. t. 95.)

1. *T. CHINENSIS*. (Sims bot. mag. t. 998.) floribus 5-6-sepalis 6-9-petalis axillaribus solitariis erectis, fructibus nutantibus dehiscentibus. 5 in Chinâ.

a. viridis (Linn. sp. 735.) foliis lanceolatis planis latitudine triplò longioribus. Letts. mon. t. 1. Blackw. herb. t. 351. (sed flos ruber.) *T. Bohea stricta* Ait. h. Kew. ed. 2. v. 3. p. 303? (v. v.)

β. Bohea (Linn. sp. 743.) foliis elliptico-oblongis subrugosis latitudine duplò longioribus. Sims bot. mag. t. 998. Blackw. herb. t. 352. Letts. mon. ed. 1. ic. Lois. herb. amat. t. 255. *T. Cantoniensis* Lour. coch. 1. p. 414. (v. v.)

An species diversæ aut aliæ adhuc hic confusæ? intereà ex auctoritate cl. Sims eas conjunxi.

2. *T. COCHINCHINENSIS* (Lour. coch. 1. p. 413.) floribus 3-5-sepalis 5-petalis solitariis terminalibus, fructibus apice dehiscentibus. 5 in Cochinchinâ boreali. Folia lanceolata. Capsula 3-loba sæpè 1-sperma.

3. *T. OLKOSA* (Lour. coch. 1. p. 414.) floribus 6-sepalis 6-petalis, pedunculis 3-floris axillaribus, fructibus indehiscentibus. 5 circa Cantonem Sinarum. Folia lanceolata.

Ordo XXXII. OLACINÆÆ. *Mirb. bull. philom.* 1813. n. 75.
p. 377.

Calyx 1-sepalus subdentatus per anthesin parvus, demùm sæpè ampliatus baccatusve, forsan pro involucro habendus. Petala (sepalave) 4-6, hypogyna subcoriacea, in æstivatione valvata, nunc libera, nunc omnia aut nonnulla binatim coalita et sic semibifida visa. Appendices piliformes aut filiformes simplices bifidæve è disco petalorum aut juxtà eorùm marginem ortæ. Stam. 3-10, nempè petalorum numero nunc dimidia, nunc dupla, nunc non ritè congruentia, hypogyna aut epipetala, filamentis compresso-subulatis antheris cordato-oblongis erectis bilocularibus. Ovarium liberum, ex cl. Mirbel 3-4 locale (in Olace ex cl. Brown 1-locul.) demùm abortu in fructum 1-locul. mutatum, loculis uni-ovulatis. Styl. 1. filiformis. Fructus subdrupaceus indehiscens, sæpè calyce ampliato aut baccato cinctus, abortu 1-loc. 1-sperinus. Semen pendulum basi umbilicatum. Albumen carnosum magnum. Embryo parvus intrà albumen inclusus ovoïdeus basilaris, radicula ad umbilicum directà, cotyledonibus continuis.

Arbores fruticesve glabri, foliis simplicibus petiolatis alternis exstipulatis integris, rarissimè nullis, floribus parvulis axillaribus.

Ordo distinctissimus, sed locus in serie naturali incertus. An calyx pro involucro et corolla pro calyce habenda et tunc, suadente cl. Brown et habitu satis congruente, Olacineæ inter Monochlamydeas non longè à Santalaceis ordinandæ sunt? An calyce et corollâ verà admissis, corollâ ut monopetala habenda tunc, suadente cl. Jussieu, ordo inter Calycifloras prope Sapoteas collocandus venit? An tandem, admittente ut videtur cl. Mirbel, Petala ut distincta considerata etiamsi variè juncta et tunc inter Thalamifloras Olacineæ admittuntur?

§. 1. *Petala staminibus antheriferis numero dupla aut circiter dupla.*

I. OLAX. *Linp. amœn.* 1. p. 387. *Gaertn. f. carp.* 3. p. 119. t. 201.

Calyx parvus integer, post anthesin ampliatus baccatus. Petala 6 per paria ad medium juncta, seu 3 semibifida. Appendices filiformes 6 bifidæ, 1 nempè disco medio cujusque petali insertæ. Stam. 3 filamenta petalis inter se coalitis adnata et ideò antheram ovatam subsessilem infrà fissuram gerentia. Drupa exsucca 1-sperma, calyce cincta. — Flores parvi racemosi. Folia ovato-oblonga 1-nervia integra:

1. *O. ZEYLANICA* (Linn. sp. 49.) foliis glabris, ramis angulatis flaccidis, caule inermi. 5 in Zeylonâ. Vahl Symb. 3. p. 7. enum. 2. p. 33. (v. s. sine fl.)
2. *O. SCANDENS* (Roxb. cor. 2. t. 102.) foliis subtus pubescentibus, ramis teretibus scandentibus, caule aculeis validis armato. 5 in Coromandelâ. (v. s.)
3. *O. IMBRICATA* (Roxb. fl. ind. 1. p. 169.) scandens, foliis ovato-lanceolatis oblongisve bifariis integris nitidis, racemis axillaribus antè anthesin bifariam imbricatis, drupis ovatis. 5 in Indiâ or. ad Chittagong.

II. *SPERMATOPHYTES*. Labill. nov.-holl. 2. p. 84. — *Oxal. Brown. prod.* 358.

Calyx parvus integer, post anthesin non ampliatus. Petala 5 nempe 4 per paria filamentis staminum mediantibus connexa et ideò semibifida, quinto libero integro. Appendices filiformes simplices. Stam. 3, 2 petalis coalitis adnata, tertio libero. Ovarium 1-loc., 3-ovulatum, ovulis ad apicem columnæ centralis filiformis pendulis. Drupa exsucca 1-sperma. — Folia disticha et secus ramos ferè ut foliola pinnatorum foliorum secus petiolum disposita, rarius nulla. Flores abortu polygami.

§. 1. *Phyllanthoidæ*, foliis distichis, pedunculis 1-floris.

1. *S. PHYLLANTHI* (Labill! l. c. t. 233.) foliis ovalibus retusis. 5 in Novâ-Holl. terrâ Van-Leuwin. *Oxal. Phyllanthi. Brown. l. c. (v. s.)*
2. *S. STRICTUM*, foliis oblongo-linearibus mucronatis. 5 in Novâ-Holland. ad Port-Jackson. *Oxal. stricta Brown! prod.* 358. (v. s.)

§. 2. *Osyroidæ*, foliis squamaceis subnullis, floribus ad squamarum axillas sessilibus et spicam terminalem sic constituentibus.

3. *S. APHYLLUM*, foliis subnullis, floribus sessilibus subspicatis. 5 in Novâ-Hollandiâ inter tropicos. *Oxal. aphylla Brown! prod.* 358. (v. s.)

III. *FISSILIA*. Comm. in Juss. gen. 260.

Calyx integer, post anthesin ampliatus. Petala 5, nempe 4 per paria mediantibus staminum filamentis juncta et ideò semibifida, quinto libero. Appendices filiformes simplices. Stam. 3, 2 petalis coalitis longè adnatis, tertio libero, filam. complanatis, antheris ellipsoideis oscillatoriis. Ovarium 3-locul. 3-spermum. Drupa sicca, calyce cupulæformi cincta, 1-sperma.

1. *F. PSITTACORUM* (Lam! dict. et ill. t. 28.) 5 in insulâ Mauritianâ. *Oxal. psittacorum. Vahl enum. 2. p. 33.* Rami rigidi. Folia ovato-oblonga aut lanceolata enervia persistentia. Racemuli axillares pauciflori. (v. s.)

§. II. *Stamina petalorum numero dupla.*

IV. *HEISTERIA*. Linn. gen. n. 535. non Berg.

Calyx minimus 5-fidus, post anthesin demùm valdè ampliatus cupulæformis, limbo expando. Petala 5 distincta. Stam. 10, filam. planis, antheris subrotundis. Stylus brevis, stigma 3-fidum. Ovarium 3-locul., 3-ovulatum. Drupa olivæformis, 1-sperma, calyce semi-cincta.

1. *H. COCCINEA* (Jacq. am. 126. t. 81.) foliis ellipticis oblongis basi rotundatis

- lucidis, floribus axillaribus, calyce fructifero lato patente obtusè sublobato. $\frac{5}{5}$ in Martinicæ sylvis ad torrentes. Vulgò *Bois Perdrix*. Lam. ill. 354. Smith in Rees. cycl. 17. n. 1. (v. s.)
2. *H. CAULIFLORA* (Smith in Rees, cycl. n. 2.) foliis oblongis interdum obovatis basi attenuatis, floribus è caule ramisve denudatis, calyce fructifero clauso profundè lobato. $\frac{5}{5}$ in Guianâ Batavâ. Flores ignoti.
3. *H. PARVIFOLIA* (Smith l. c. n. 3.) foliis ovatis acutis lucidis, floribus axillaribus, calyce fructifero patente, lobis ovatis profundis. $\frac{5}{5}$ in Sierrâ-Leonâ. Rami subcompressi. Calycis sinus reflexi.

V. *XIMENIA*. *Plum. gen.* 6. t. 21.

Calyx minimus 4-fidus persistens non ampliatus. Pet. 4, intus pilosa, basi conniventia, supernè revoluta. Stam. 8, filam. capillaribus, antheris adnatis elongatis linearibus. Ovarium 4-locul. 4-spermum. Styl. 1. Drupa ovata 1-sperma. — Arborès fruticesve, foliis ovatis aut lanceolatis.

1. *X. AMERICANA* (Linn. sp. 497.) pedunculis multifloris sæpè inferioribus in spinas mutatis. $\frac{5}{5}$ in Americâ calid. (v. s.)
a. ovata, foliis ovatis. — in ins. Caribæis. *X. multiflora* Jacq. am. 106. t. 277. f. 31. Lam. ill. t. 297. f. 1.
β. oblonga, foliis oblongis. Heymassoli spinosa Aubl. Guian. 1. p. 324. t. 125. — Lam. ill. t. 297. f. 2.
2. *X. ELLIPTICA* (Forst. prod. n. 162.) pedunculis multifloris inermibus, foliis elliptico-lanceolatis. $\frac{5}{5}$ in Novâ-Caledoniâ.
3. *X. INERMIS* (Linn. sp. 497.) pedicellis 1-floris, foliis ovatis. $\frac{5}{5}$ in Jamaicâ. Lunan hort. jam. 2. p. 156.*
4. *X? LANCEOLATA*, floribus corymboso-racemosis, foliis oppositis ovato-lanceolatis. $\frac{5}{5}$ in Cochinchinâ. Gela lanceolata Lour. coch. 1. p. 285. Flores luteo-virides, petalis linearibus.

†. *Genera non satis nota.*VI. *PSEUDALEIA*. *Pet. Th. gen. mad.* n. 51.

Calyx minimus subinteger. Pet. 3 in tubum conniventia. Filam. 6, petalis arcuè adplicata undè vid. epipetala. Capillamenta apice bifurca ex utroque latere petalorum. Ovarium conicum. Stylus longitudine corollæ. Stigma 3-lobum. Drupa spherica, 1-sperma. Semen exalbuminosum! Embryo cotyledonibus carnosus indistinctis oleosis. — Ex flore videtur inter Olacem et Heisteriam media, sed semen omninò differt.

1. *P. MADAGASCARIENSIS*. $\frac{5}{5}$ in Madagascar. Arbuscula ramosa, foliis alternis lævibus, pedunculis axillaribus paucifloris. *Olax Pseudaleia* Willd. in Steud. nom. phau.

VII. *PSEUDALEIODES*. *Pet. Th. gen. mad.* n. 52. †.

Cal. minimus integer. Petala 4 basi lata conniventia inæqualia. Stam. 6, filamenta lata, petalis dispariter applicata ut ibi appareant inserta. Antheræ apice insertæ. Ovarium 1-spermum? stylus longitudine corollæ. Stigmata 3 globosa. Fruct. ign. — Genus priori affine.

1. *P. THOUARSII*. $\frac{5}{5}$ in Madagascar. Frutex debilis, folia alterna, racemi 1-laterales pauciflori. *Olax Pseudaleioides*. Willd. in Steud. nom. phau.

GENUS OLACINEIS AFFINE.

VIII. ICACINA. *Adr. Juss. mem. soc. h. nat. par. 1823. 1. p. 174.*

Cal. brevis 5-fidus sæpè persistens. Pet. 5. per æstivationem valvata, lobis cal. alterna, iis triplo longiora, basi intus villosa. Stam. 5 petalis alterna erecta, antheris cordatis medifixis bilocularibus, loculis longitudinaliter dehiscentibus. Stylus simplex, incurvus, apice truncatus. Ovarium disco glanduloso insidens, 1-locul. Capsula apice dehiscens, abortu sæpius 1-sperma. Sem. ign. — Arbor Senegalensis. Folia simplicia, alterna, extipulacea, breviter petiolata, ovata integra, reticulato-nervosa. Flores paniculati terminales.

1. I. SENEGALENSIS (Ad. de Juss. l. c. t. g.) 5 in Senegaliâ. Habitus ferè Chrysobalani Icaco, unde nomen genericum.

Ordo XXXIII. AURANTIACEÆ. *Corr. ann. mus. 6. p. 376.*
Mirb. bull. philom. 1813. p. 379.

Calyx urceolatus campanulatusve, disco subadnatus, brevis, 3-5-dentatus, marcescens. Petala 3-5 basi lata, nunc libera, nunc inter se subcoalita, disco hypogyno extus inserta, per æstivationem marginibus leviter imbricata. Stam. numero petalorum æqualia, dupla aut multiplicia, disco hypogyno inserta; filamenta basi complanata, nunc libera, nunc inter se variè adnexa polyadelpha, nunc longè monadelpha, semper apice libera subulata; antheræ terminales, basi insertæ, erectæ. Ovarium ovatum, multiloculare. Stylus 1 teres. Stigma subdivisum crassiusculum. Fructus (Aurantium DC.) constans 1°. Indusio crasso glanduloso evalvi indehiscente, verosimiliter pro toro continuato habendo. 2°. Carpellis plurimis (rariùs abortu solitariis) circa axim rationalem verticillatis, sine laceratione sæpè separabilibus, membranaceis, nunc semina tantum foventibus, nunc intus pulposis carnosisve; pulpâ in sacculis innumeris è parietibus ortis reclusâ. Semina in carpellis sita, ad eorum angulum interiorem adfixa, nunc plurima nunc solitaria, exalbuminosa, sæpiùs pendentia, interdum plures embryones foventia. Spermodermis raphe et chalazâ sæpiùs cupulari notata. Embryo rectus, radiculâ retractâ superâ ad hilum versâ, cotyledonibus magnis crassis basi biauriculatis, plumulâ conspicuâ.

Arbores aut frutices ferè omnes glaberrimæ, in foliis, calycibus, petalis, stam. filamentis, fructus indusio et cotyledonibus glandulis vesicularibus oleo volatili foetis instructæ. Folia alterna, super caulem articulata, tardè decidua nunc composita pinnata multi aut unijuga imo foliolo unico terminali super petiolum dilatato-foliaceum articulato instructa, nunc simplicia nempè forsàn ad petiolum dilatatum reducta et foliolo ultimo abortivo orbata. Spinæ axillares in ramos non abeuntes.

I. ATALANTIA. *Corr. ann. mus. 6. p. 383.*

Flores proportionè partium quarternariâ. Stam. 8 longè monadelpha, filam. apice liberis, antheris terminalibus. Pistillum villosum. Fructus sphæricus 4-locul. 4-spermus.

1. A. MONOPHYLLA. 5 in Indiâ orient. Folia ovato-oblonga apice emarginata. Spinæ parvæ simplices. Pedunculi axillares. Limonia monophylla Roxb. cor. 2. p. 59. t. 83. (v. s.)

II. TRIPHASIA. *Lour. coch. 1. p. 189.*

Flores proportionè partium ternariâ. Stamina 6 libera, rariùs 5 et

tunc quintum oppositum petalo majori. Antherae subsagittatae. Fructus 3-locul. 3-spermus. Embryones in semine plures. — Spinæ axill. rectæ.

1. *T. MONOPHYLLA*, foliis simplicibus oblongis, 5 in ins. Timor. Folia subsessilia obtusa. Racemi parvi ad axillas spinarum folio breviores. Flores minimi. Fructus ignotus. Habitus Atalantiæ, flos Triphasiae. (v. s.)
2. *T. TRIFOLIATA*, foliis 3-foliolatis. 3 in Indiâ, Cochinchinâ, Chinâ. Limonia trifoliata Linn. mant. 237. Jacq. ic. rar. t. 463. Andr. bot. rep. t. 143. *T. Auran-tiola* Lour. l. c. Folia breviter petiolata, apice 3-foliolata, foliolis ovatis, terminali sæpius emarginato. Spinæ rectæ parvæ. (v. s.)

III. LIMONIA. Linn. gen. n. 524.

Flores proportionem partium quaternariâ aut quinariâ. Calyx 4-5-partitus. Petala 4-5. Stâm. libera, petalorum numero dupla, interdum æqualia. Fructus baccatus pulposus 4-5-loc., loculis 1-spermis. — Genus forsân species heterogeneas colligens.

* *Foliis simplicibus*. — Species dubiæ quoad charact. genericum.

1. *L. LUCIDA*. (Forst. prod. n. 191.) foliis simplicibus, pedunculis axillaribus. 5 in insulâ Mallicollo.
2. *L. LAUREOLA*, foliis simplicibus, floribus terminalibus corymboso-capitatis. 5 in Napanliâ. Folia exactè *Daphnes Laureolæ*. Cal. 5-partitus. Pet. et stamina 5! huc relata ex auctor. clar. Wallich. (v. s.)

** *Foliis trifoliolatis*. — Species habitu *Toddaliæ*.

3. *L. CITRIFOLIA*. (Willd. enum. 448.) foliis simplicibus trifoliolatisque, foliolis ovato-oblongis acuminatis, pedunculis axillaribus petiolo brevioribus. 3 in Chinâ. Flores parvi.
4. *L. DIACANTHA*, foliis trifoliolatis, foliolis ovalibus subemarginatis, spinis axillaribus duabus rectis, pedicello 1-floro inter spinas iis brevior. 5 in Pondicheri. Flores parvi 5-andri? (v. s.)
5. *L. MAURITIANA* (Lam. dict. 3. p. 517.) foliis 3-foliolatis, foliolis ovalibus submucronatis, paniculis axillaribus, spinis nullis. 5 in ins. Mauritiæ. Flores 4-fidi, 8-andri. (v. s.)

*** *Foliis pinnatis, petiolo aptero, spinis nullis*.

6. *L. MADAGASCARIENSIS* (Lam. dict. 3. p. 517.) foliis pinnatis, foliolis 4-5 alternis lanceolatis dentatis, paniculis brevibus axillaribus. 5 in Madagascar. Rhoem refert.
7. *L. PARVIFLORA* (Sims bot. mag. t. 2416.) foliis bijugis, foliolis elliptico-lanceolatis integerrimis, corollis campanulatis, fructibus ovato-sphæroideis obliquis. 3 in Chinâ.
8. *L. MINUTA*. (Forst. prod. n. 190.) foliis pinnatis, racemis corymbosis terminalibus. 5 in insulis Amicorum.

**** *Foliis pinnatis, petiolo alato, spinis axillaribus*. — Limoniæ veræ.

9. *L. ACIDISSIMA* (Linn. mant. 380.) foliis pinnatis bijugis, foliolis obovatis subemarginatis, spinis solitariis, fructibus ovoideis. 5 in Indiâ orient. Rumph. Amb. 2. t. 43. Flores 5-fidi.
10. *L. CRENULATA*. (Roxb. cor. 1. p. 86.) foliis pinnatis 2-3-jugis, foliolis elliptico-oblongis crenulatis, spinis solitariis, fructibus globosis. 3 in Coromandelâ. Rheed. mal. 4. t. 14. Flores 4-fidi 8-andri.
11. *L. AMBIGUA*, foliis pinnatis, spinis geminis, foliolis rotundato-ovalibus crenatis. 5 in Floridâ orient. culta? *L. acidissima*? Nutt. in Sillim. am. journ. 5. 1822. p. 295. An *L. acidissimæ*, an *crenulatæ* var? †.

IV. COOKIA. *Sonner. voy. 2. p. 136. t. 131.*

Flores proportione partium quinariâ. Cal. 5-fidus. Pet. navicularia villosa. Stam. 10, filam. linearibus liberis, antheris subrotundis. Ovarium villosum. Fructus baccatus subglobosus 5-locul., abortu 1-2-locul., loculis 1-spermis. — Arbusculæ foliis pinnatis, foliolis alternis basi inæqualibus obliquisve.

1. *C. PUNCTATA* (Retz. obs. 6. p. 29.) foliis ovato-lanceolatis acuminatis vix basi inæqualibus. 5 in Chinâ et ins. Moluccanis. — Jacq. schoenb. 1. t. 101. Lam. ill. t. 354. Quinaria Lansium Lour. coch. 1. p. 334. — Rumph. amb. 1. t. 55. Arbor mediocris fructu eduli. An 2 aliæ Lansii icones Rumphianæ sint hujus varietates aut species distinctæ? (v. s.)
2. *C. FALCATA*, foliis lanceolato-falcatis, petalis intus 4-sulcatis. 3 in sylvis Cochinchinæ. Aulacia falcata Lour. coch. 1. p. 335. Arbuscula 8-pedalis, ramis patentibus.

V. MURRAYA. *Kæn. in Schreb. gen. n. 717. — Marsana Sonn. voy. ind. 2. t. 139.*

Flores proportione partium quinariâ. Cal. 5-partitus. Cor. campanulata. Stam. 10, filam. lineari-subulatis, antheris subrotundis. Fructus carnosus-baccatus 2-loc., sæpè abortu 1-loc., loculis 1-spermis. Semen appendens. Spermodermis crassa lanata. Cotyledonum auriculæ minimæ. (Mirb. l. c.). — Folia impari-pinnata.

1. *M. EXOTICA* (Linn. mant. 563.) foliolis ovatis, pedunculis multifloris corymbosis, baccis subrotundis sæpius 1-spermis. 5 in Indiâ orient. — Murr. comm. goett. 9. p. 186. t. 1. Lam. ill. t. 352. Chalcas Japonensis Lour. coch. 332. Marsana buxifolia Sonner. it. ind. t. 139. (v. s.)
2. *M. PANICULATA* (mal. misc. 1. n. 2. p. 31.) foliolis ovatis acuminatis, floribus terminalibus axillaribusque subsolitariis, baccis oblongis sæpius dispersis. 5 in Indiâ orient. Chalcas paniculata Lour. coch. 331. — Rumph. amb. 5. p. 26. t. 17.

VI. AGLAIA. *Lour. coch. 216.*

Flores proportione partium quinariâ. Calyx 5-partitus. Stamina 5, filam. ut in Meliaceis in tubum ovatum coalitis, antheris inclusis. Stigma latum sessile. Ovarium 1-locul. 2-ovulatum. Bacca 1-sperma. (mal. misc. 1. n. 2. p. 33.)

1. *A. ODORATA* (Lour. l. c.). 3 in Cochinchinâ et Chinâ. Folia pinnata 5-7-foliolata nitida. Flores parvi flavi fragrant. Camunium Sinense Rumph. amb. 5. p. 28. t. 18.

VII. BERGERA. *Kæn. in Schreb. gen. n. 718.*

Flores proportione partium quinariâ. Stam. 10, filamentis subulato-compressis, antheris subrotundis. Stylus obconicus. Ovarium 2-locul. 2-ovul. Fructus baccatus, sæpius 1-locul., 1-spermus. Semen appendens; spermodermis membranacea. Cotyledonum auriculæ maximæ. (Mirb. l. c.) An à Murraya satis differt? Folia impari-pinnata, foliolis alternis acuminatis pubescentibus.

1. *B. KOENIGII* (Lin. mant. 565.) foliolis serratis. 5 in Indiâ orient. — Rumph. amb. 1. p. 149. t. 53. f. 1. Racemi plurimi ad apicem ramorum corymbosi.
2. *B. INTEGERRIMA* (Roxb! in herb. Lamb.) foliolis integerrimis, 5 in Indiâ orient. Paniculæ subcorymbosæ terminales. (v. s.)

VIII. CLAUSENA. *Burm. ind. p. 89.*

Flores proportione partium quaternariâ. Cal. 4-dentatus. Pet. subrotunda. Stam. 8, filam. subulatis basi dilatato-excavatis, antheris subrotundis. Stylus cylindraceus (Burm.) Ovar. 4-locul., loculis 1-spermis. Fructus subsiccus, abortu 1-locul., 1-spermus. Semen appendens. Spermodermis membranacea. Cotyledonum auriculæ maximæ (Mirb. l. c.) — Folia impari-pinnata, foliolis petiolatis pubescentibus. Flores minimi laxè paniculati.

1. *C. EXCAVATA* (Burm. l. c. t. 29.) foliolis 13-15. 5 in Javâ C. Javensis Roëseh. (v. s.)
2. *C² PENTAPHYLLA*, foliolis 5-7. 5 in Indiâ orient. *Limonia pentaphylla* herb. Lamb. non Roxb. (v. s. sine fructu.)

IX. GLYCOSMIS. *Corr. ann. mus. 6. p. 384.*

Flores proportione partium quinariâ. Stam. 10. filam. subulato-complanatis, antheris ellipsoideis. Stylus brevis cylindraceus. Ovarium 5-locul., loculis 1-spermis. Fructus carnosus 1-2-locul. Semina pendentia. Spermodermis membranacea. Cotyledonum auriculæ brevissimæ. (Mirb. l. c.) — Arbores foliis impari-pinnatis glabris.

1. *G. ARBOREA*, foliolis 5-7 oblongo-linearibus alternis denticulatis. 5 in sylvis Coromandelæ et in ins. Mauritianâ. *Limonia arborea* Roxb. cor. 1. t. 85. (v. s.)
2. *G. PENTAPHYLLA*, foliolis 5 ovatis integerrimis. 5 in Coromandelâ. *Limonia pentaphylla* Retz. obs. 5. p. 24? Roxb. cor. 1. t. 84.

X. FERONIA. *Corr. act. soc. lin. 5. p. 224.*

Flores proportione partium quinariâ. Calyx planus 5-partitus. Petala 5 oblonga. Stamina 10, filamentis liberis basi dilatatis villosis, antheris oblongis. Discus tori elevatus. Fructus baccatus multilocularis, loculis polyspermis, carne spongiosa obvolutis.

1. *F. ELEPHANTUM* (Corr. l. c. Roxb! cor. 2. t. 141.) foliis opacis, petiolo communè angustè marginato glabro. 5 in Coromandelæ sylvis montosis. Spinæ simplices. Folia impari-pinnata, foliis 5-7 obovatis glabris. Paniculæ parvæ axillares terminalesve. Bacca globosa pomi magnitudine. (v. s. sine fr.)
2. *F. PELLUCIDA* (Roth. nov. sp. p. 384.) foliis pellucido-punctatis, petiolo communi tereti pubescente. 5 in Indiâ orientali.

XI. ÆGLE. *Corr. act. soc. lin. 5. p. 222. — Belou Adans. fam. 2. p. 408.*

Flores proportione partium 3-5-nariâ. Calyx 3-5-dentatus. Pet. 3-5 patentia. Stam. 30-36 libera, antheris longis linearibus mucronatis. Stigma subsessile. Fructus baccatus demùm lignosus turbinato-globosus multilocularis, loculis polyspermis. Spermodermis carnosa mucobducta. Cotyledonum auriculæ brevissimæ. (Mirb. l. c.) — Frutices spinis simplicibus, foliis 3-foliolatis denticulatis.

1. *Æ. MARMELOS*. (Corr. l. c. Roxb. cor. 2. t. 143.) foliolo medio petiolato, fructu 12-loculari. 5 in Indiâ orient. *Cratæva Marmelos* Lin. sp. 637. Fructus pulpâ gratissimâ. (v. s.)
2. *Æ² SEPIARIA*, foliolo medio sessili, fructu 7-loculari. 5 in Japoniâ. *Citrus trifoliata* Lin. sp. 1101. — Ssi Kœmpf. amoen. 801. t. 802. ex habitu et staminibus liberis ut videtur in icone huc relatâ. Flores axillares solitarii. Fructus pulpâ ingrâtâ.

XII. CITRUS. *Lin. gen. n. 1218.*

Flores proportione partium sæpius quinariâ. Calyx urceolatus 3-5-fidus. Petala 5-8. Stam. 20-60, filamentis compressis plus minùsve basi coalitis polyadelphis, antheris oblongis. Stylus teres. Stigma hemisphæricum. Fructus baccatus 7-12-locul. loculis polyspermis pulposis. Spermodermis membranacea. Cotyledonum auriculæ brevissimæ. — Arbores fruticesve spinis axillaribus foliis apice petioli 1-foliolatis petiolo sæpè alato.

1. *C. MEDICA* (Risso ann. mus. 20. p. 199. t. 2. f. 2.) petiolis nudis, foliis oblongis acutis, floribus 40-andris sæpè agynis, fructuum oblongorum rugosorum cortice crasso, pulpâ acidulâ. 5 in Asiâ. Culta speciatim in Mediâ et in Europâ anstr. jam à temporibus antiquissimis. Ferr. hesp. t. 59. 60. 63. — *C. medica Cedra*. Gall. citr. p. 87. var. 1. 2. 5 et 7. Ramuli teneri violacei. Cortex fructus adhærens odore suavi. Petala extûs purpurascentia. Varietates 3 enum. cl. Risso. Vulgò *Cedrat*, *Cedrot*. (v. v.)
2. *C. LIMETTA* (Risso ann. mus. 20. p. 195. t. 2. f. 1.) petiolis nudis, foliis ovatis rotundatis serratis, floribus 30-andris, fructuum globosorum umbone obtuso coronatorum cortice firmo, pulpâ dulci. 5 in Asiâ. Culta in Italiâ. — *C. medica Limon*. Gall. citr. n. 10. 25-38. Ferr. hesp. t. 230. 395. 233. 321. Sequenti valdè affinis. Corolla utrinquè alba. Vulgò *Limo dulcis*. *Limetta*, *Bergamotta*, *Peretta*. Var. 7 enum. Risso. (v. s.)
3. *C. LIMONUM* (Risso ann. mus. 20. p. 201.) petiolis subulatis, foliis oblongis acutis dentatis, floribus 35-andris sæpè agynis, fructuum oblongorum cortice tenuissimo, pulpâ acidissimâ. 5 in Asiâ, nunc in Europâ austr. culta. *C. medica Limon*. Gall. citr. 105. excl. variet. sub n. 2 citatis. — Ferr. hesp. t. 247. 211. 253. 223. 229. 293. 255. 265. 105. 225. 207. 243. 219. 301. 307. 215. Petala extûs purpurascentia. Ramuli violacei. Fructus ovatus, cortice adhærente flavo. Vulgò Gall. *Citron*. Ital. *Limone*. Var. 25 enum. à Risso, aliæ fructibus ovoideis umbone obtusè terminatis, aliæ fructibus oblongis grossè umbonatis. (v. v.)
4. *C. AURANTIUM* (Risso ann. mus. 20. p. 181. t. 1. f. 1. et 2.) petiolis subnudis, foliis ovato-oblongis acutisque, floribus 20-andris, fructuum globosorum cortice tenni, pulpâ dulci. 5 in Asiâ orient. nunc in Europâ austr. culta. *C. Aurantium sinense* Gall. citr. 119. — Ferr. hesp. t. 427. 399. 401. 385. Rami in comam sphæricam. Petala alba. Fructus aureus. Vulgò *Aurantium dulce*. Var. 19 enum. à cl. Risso. (v. v.)
5. *C. VULGARIS* (Risso ann. mus. 20. p. 190.) petiolis alatis, foliis ellipticis acutis crenulatis, floribus 20-andris, fructuum globosorum cortice tenui scabroso, pulpâ acri amarâ. 5 in Asiâ, nunc in Europâ anstr. culta. *C. Aurantium Indicum*. Gall. citr. 122. *C. Bigaradia* Duham. ed. nov. 7. p. 99. Ferr. hesp. t. 409. 589. 391. 430. 433. *C. Sinensis* Pers. ench. 2. p. 74? *C. Aurantium* bot. reg. t. 346. *C. Calot* Lag. gen. et sp. 17. Petala alba. Variet. 11 enum. cl. Risso. Vulgò *Bigarade*, *Aurantium amarum*. (v. v.)

†. *Species non satis notæ.*

6. *C. DECUMANA* (Lin. sp. 1100.) petiolis alatis, foliis obtusis emarginatis, fructibus maximis crassè corticatis. 5 in Indiâ orient. et nunc in Americâ calid. Pampel-mœs Rumph. 2. t. 24. f. 2. (v. s.)
7. *C. HYSTRIX* (DC. cat. h. monsp. p. 97. petiolis latè alatis, foliis ovatis vix petiolo majoribus, ramis spinosissimis. 5 in Indiâ. Lemo ferus Rumph. 2. t. 28? (v. v. sine fl. et fr.)
8. *C. SPINOSISSIMA* (Meyer esseq. 247.) petiolis alatis, foliis ovalibus utrinquè obtusiusculis crenatis. 5 in sylvis arenosis Guianæ. Flores gemini. Fructus acidus, magnitudine nucis Juglandis.

9. *C. JAPONICA* (Thunb. fl. jap. 292.) petiolis alatis, foliis acutis, caule angulato, floribus axillaribus solitariis geminisve. ♀ in Japoniâ. Thunb. ic. jap. t. 15. Fructus colore et formâ Aurantii, sed minutissimi, dulces et edules.
10. *C. FUSCA* (Lour. coch. 571.) petiolis cordato-alatis, foliis lanceolato-ovatis, ramis spinosis, fructibus globosis asperis. ♀ in Cochinchinâ, Chinâ, Moluccis. — Rumph. amb. 2. t. 33. Bacca fusco-viridis, pulpâ ingratâ acidâ.
11. *C. NOBILIS* (Lour. coch. 569.) petiolis sublinearibus strictis, ramis adscendentibus inermibus, fructibus depressis crassè corticatis. ♀ in Cochinchinâ et Chinâ. Ker. bot. reg. t. 211. Andr. bot. rep. t. 608. Fructus intus et extus ruber, succo dulci.
12. *C. MARGARITA* (Lour. coch. 570.) petiolis linearibus, foliis lanceolatis, ramis adscendentibus spinosis, fructibus oblongis 5-locularibus tenuiter corticatis. ♀ ad Cantonem Sinarum. Fructus rubro-luteus, 8 lin. longus, pulpâ dulci.
13. *C. MADURENSIS* (Lour. coch. 570.) petiolis linearibus, foliis lato-lanceolatis, ramis diffusis inermibus angulatis, fructibus globosis lævibus. ♀ in Cochinchinâ, Chinâ, Madurâ. — Rumph. amb. 2. t. 31. Fructus semipoll. diam. pulpâ amaricante.
14. *C. ANGULATA* (Willd. sp. 3. p. 1426.) petiolis nudis, foliis ovatis acutis, fructibus angulatis. ♀ in Amboinâ. — Rumph. amb. 2. t. 32.
15. *C. BUXIFOLIA* (Poir. dict. 5. p. 681.) petiolis linearibus brevissimis, foliis ovatis retusis, floribus racemosis. ♀ in Chinâ.
-

Ordo XXXIV. HYPERICINEÆ (1). *DC. theor. elem. ed. 1. p. 214. fl. fr. 4. p. 860. Chois. prod. hyp. 32. — Hyperica Juss. gen. p. 254.*

Calyx monosepalus 4-5-partitus, seu 4-5-sepalus, persistens, sæpè inæqualis, nempè 2 sepalis minoribus externis, 2 aut 3 majoribus, sæpè punctatus, aut dentato-glandulosus. Petala 4-5 hypogyna, calycis lobis alterna, contortè æstivata, vulgò lutea venata, aliquandò nigro-punctata. Stamina numerosa, sæpiùs indefinita, basi polyadelphe, rarò libera aut monadelphæ; filamenta longa; antheræ oscillantes, luteæ minutæ. Ovarium unicum, liberum. Styli plurimi longi aliquandò in unum coaliti. Stigmata simplicia, rarò capitata. Capsula baccæ multivalvis multilocularis; loculi numero stylos æquant. Placenta integra centralis seu multipartita margini introflexo valvularum adfixa. Semina numerosissima, vulgò teretia, rarò complanata. Embryo rectus; radícula infera; albumen nullum. — Herbæ, frutices, suffrutices et arbores, sæpè succu resinoso donatæ aut variis glandulis obsitæ. Folia opposita, integra, rarissimè, alterna, crenata, sessilia aut breviter petiolata, pellucido-punctata, aut nigro-punctata, aut impunctata, penninervia. Flores aut terminales, aut axillares, aut pedunculati, aut sessiles, aut foliosi, aut nudè paniculati, sæpè bracteati.

Subordo I. HYPERICINEÆ VERÆ. *Chois. prod. hyp. 32.*

Semina teretia Styli sæpiùs 3-5.

Trib. I. *VISMIEÆ*. *Chois. prod. hyp. 33.*

Bacca. Flores in paniculis distinctis aphyllis racemosis vel corymbosis terminalibus. Frutices et arbores. Folia petiolata.

I. HARONGA. *Pet. Th. gen. nov. madag. n. 49. Chois. prod. hyp. 33. — Harongana Lam. ill. t. 645. — Arongana Pers. ench. 2. p. 91.*

Bacca drupacea; loculi quinque 2-3-spermi. Styli et stygmata 5. Stam. 15 altè 5-adelphe. Squamulæ 5 adelphiis alternæ. Sepala et petala 5. — Frutices. Caulis ramosus. Flores in paniculis nunc brachiatis ramosissimis nunc paucifloris dispositi.

§. 1. *Foliis integris.*

I. H. MADAGASCARIENSIS (*Chois. prod. hyp. 34.*) caule tereti, foliis elliptico-

(1) Ordinis hujus enumerationem proprio Marte conscripsit et mecum communicavit cl. et am. J. D. CHOISY.

lanceolatis amplis longè petiolatis, corymbo florum terminali amplissimo ramosissimo. ζ in ins. Madagascar et Borboniâ. (v. s.)

β . *pubescens*, foliis minoribus magis pubescentibus. H. *pubescens*? Poir. encycl. suppl. 4. p. 721.

2. H. *LANCEOLATA* (Chois. mss.) caule anguloso, foliis ovato-lanceolatis 1-2 pollices longis utrinquè glaberrimis, brevissimè petiolatis, corymbo terminali paucifloro. ζ in ins. Madagascar. (v. s.)

3. H. *REVOLUTA* (Chois. mss.) foliis ovato-oblongis obtusis margine revolutis amplis in nervis rufescentibus, corymbo florum paucifloro stricto. ζ in ins. Madagascar. (v. s.)

§. 2. *Foliis crenatis.*

4. H. *MOLLUSCA* (Pers! ench. 2. p. 91.) foliis lanceolato-acuminatis obsolete crenatis, pedunculis axillaribus 3-4-floris. ζ in Madagascar. (v. s.)

5. H. *CRENATA* (Pers! l. c.) foliis ovatis obtusis amplis latè crenatis, pedunculis axillaribus multifloris. ζ in Madagascar. (v. s.)

II. *VISMIA*. Vand. in Ræm. script. hisp. p. 138. t. vii. f. 4. Chois. prod. hyp. 34.

Bacca membranacea. Stylî 5. Stigmata peltata. Stamina 10, 5-adelpha; glandulæ 5 adelphiis alteruæ. Sepala 5. Petala 5, intus plerumque villosa. — Frutices et arbores, ramis quadrangulis, foliis sæpius tomentoso-rufescentibus, rarò pellucido-punctatis: flores in paniculis ramosis dispositi; gemmæ globosæ aut oblongæ.

1. V. *GLABRA* (Ruiz et Pav. syst. fl. per. p. 183.) caule 3-orgyali, ramulis compressis, foliis elliptico-lanceolatis glabris, petiolo brevi compresso, gemmâ globosâ, calyce ovato-oblongo obtuso glabro, paniculis laxis, baccâ globosâ. ζ in Peruvîâ. (v. s.)

2. V. *SESSILIFOLIA* (Pers. ench. 2. p. 86.) caule anguloso, foliis elliptico-lanceolatis acutis basi cordatis glabris, petiolo brevissimo crasso, gemmâ globosâ, calyce ovato-oblongo, paniculis multifidis. ζ in Guianâ. Hypericum sessilifolium. Aubl. guian. 2. p. 787. t. 312. f. 2. (v. s.)

3. V. *RETICULATA* (Chois. prod. hyp. p. 34.) ramulis rufescentibus, foliis elliptico-oblongis longissimis subobtusis basi subcordatis reticulatis rufescentibus, petiolo brevissimo, gemmâ ovato-globosâ, calyce obtuso extus villosus. ζ in Guianâ. Hypericum reticulatum Poir. suppl. dict. 3. p. 694. (v. s.)

4. V. *MACROPHYLLA* (H. B. Kunth! nov. gen. am. 4. p. 184.) ramulis glabris, foliis elliptico-oblongis acuminatis cordatis subtus pilosisculis, petiolis 8 lineas longis, calycibus ferrugineo-tomentosis. ζ ad ripam fluminis Cassiquiare. (v. s.)

5. V. *GUIANENSIS* (Pers. ench. 2. p. 86.) caule quadrangulo, foliis ovato-lanceolatis acuminatis basi dilatatis subtus rufescentibus suprâ glabris, petiolo brevi, calyce ovato obtuso villosus, floribus corymbosis, baccâ ovatâ. ζ in Guianâ. Hypericum Guianense Aubl. guian. 2. p. 784. 4. t. 311. (v. s.)

β . *glabrata*, totâ plantâ minùs rufescente candidiore, foliis magis acuminatis, gemmis magis globosis. An species distincta? Hypericum bacciferum. Margr. bras. 96. f. 1. Moc. et Sess. ic. fl. mex. ined. — in Mexico et Surinam.

6. V. *RUFESCENS* (Pers. ench. 2. p. 86.) ramulis tetragonis, foliis oblongo-lanceolatis longissimè acuminatis basi attenuatis infrâ subpilosis suprâ glabris, petiolo brevi canaliculato, calyce acuto glabro. ζ in Guianâ. Chois. prod. hyp. p. 35. t. 1. (v. s.)

7. V. *DEALBATA* (H. B. et Kunth! nov. gen. am. 4. p. 184. t. 454.) ramulis tetragonis albedo-tomentosis, foliis ovatis acuminatis basi rotundatis subtus albo-tomentosis, petiolo 7-8 lineas longo sulcato, calyce ferrugineo-tomentoso, ovario subgloboso-ovato. ζ ad ripam fl. Nigri et Cassiquiare. (v. s.)

8. V. *FERRUGINEA* (H. B. et Kunth! nov. gen. am. 4. p. 183.) ramulis tetragonis

calycibusque ferrugineo-tomentosis, foliis elliptico-oblongis acuminatis supra subnitidis subtus tenuissimè aureo-tomentosis, petiolis 6-9-lineas longis canaliculatis, ovario oblongo. ζ ad Orinocum. (v. s.)

9. V. LAURIFORMIS (Chois. prod. hyp. p. 35.) foliis ovatis acutis glabris, calyce ovato-lanceolato concavo, corollâ longitudine calycis, adelphis penicelliformibus, stylis filiformibus. ζ in Novâ-Granadâ. *Hypericum lauriforme*. Lam. dict. 4. p. 152.

10. V. BRASILIENSIS (Chois. prod. hyp. p. 35. t. 2.) caule tereti, ramulis apice subcompressis, foliis ovato-lanceolatis acutis breviter acuminatis subtus rufescentibus supra glabris, petiolo canaliculato, floribus corymbosis, bacca globosâ. ζ in Brasiliâ. (v. s.)

11. V. CAYENNENSIS (Pers. ench. 2. p. 86.) caule tereti, foliis ovato-lanceolatis acuminatis subtus punctatis supra nigris, petiolo canaliculato, calyce obtuso, gemmis globosis, paniculis paucifloris. ζ in Guianâ et Cajennâ. *Hypericum Cayennense*. Linn. amœn. 8. p. 321.

12. V. ACUMINATA (Pers. l. c.) foliis subtus hispidulis vix punctatis apice maximè acuminatis, gemmis parvis ovatis, calyce subuloso, bacca globosâ. ζ in Guianâ. *Hypericum acuminatum* Lam. dict. 4. p. 150. (v. s.)

β . *caparosa*, undiquè multò hispidiore. In monte Hyguerote. V. *caparosa*. H. B. et Kunth nov. gen. am. 4. p. 182. (v. s.)

13. V. GUINEENSIS (Chois. prod. hyp. p. 36.) caule tereti, ramis divaricatis, foliis ovato-lanceolatis acutis subtus punctatis mollibus, petiolo tenui, paniculis patulis, calyce ovato-lanceolato, corollâ glabrâ. ζ in Guineâ, Sierra-Leonâ. *Hypericum Guineense* Lin. amœn. 8. p. 32. t. 8. f. 1. (v. s.)

†. *Species minus notæ aut dubii generis.*

14. V. LATIFOLIA (Chois. prod. hyp. p. 36.) arborescens, foliis ovato-acuminatis subcordatis infra breviter tomentoso-rufescentibus supra viridibus, petiolo brevi crasso. ζ in Guianâ. *Hypericum latifolium* Aubl. guy. 2. p. 787. t. 312. f. 1.

15. V. TOMENTOSA (Ruiz et Pav. syst. fl. per. p. 183.) foliis ovatis acutis subtus tomentosis, racemis terminalibus. ζ in Peruvîâ.

16. V? PETIOLATA (Chois. prod. hyp. p. 36.) floribus 3-gynis, foliis ovatis subtus tomentosis, caule tetragono compresso. ζ in Brasiliâ. *Hypericum petiolatum* Linn. sp. 1102.

17. V? ARBORESCENS (Chois. prod. hyp. p. 36.) floribus 3-gynis, foliis ellipticis parùm acuminatis, racemis brachiatis, calycibus corollisque glabris, fructu capsulari? ζ in Indiâ orient. *Hypericum arborescens* Vahl symb. 2. p. 86. t. XLIII.

Trib. II. *HYPERICINÆ*. Chois. prod. hyp. 37.

Capsula. Flores terminales aut axillares. Herbæ aut suffrutices: folia plerùmque sessilia.

III. *ANDROSÆMUM*. All. ped. n. 1440. Chois. prod. hyp. 37.

Capsula baccata sub-1-locularis. Calyx 5-partitus, lobis inæqualibus. Petala 5. Styli 3. Stamina ∞ basi coalita. — Suffrutex, folia sessilia; flores terminales pedunculati.

1. A. OFFICINALE (All. l. c.) char. id. ac generis. — ζ in humidis Europæ borealis, Caucasi et Græciæ. Blackw. t. 94.

IV. *HYPERICUM*. Linn. gen. n. 902. Juss. gen. p. 255. Chois. prod. hyp. 37.

Capsula membranacea. Styli 3-5 in quibusdam numero variabiles

Stamina ∞ basi polyadelphia, raro numero subde finita. Pet. 5. Sepala 5 basi plus minusve coalita et inæqualia. — Herbæ aut suffrutices. Folia opposita, sæpè pellucido-punctata, aut margine nigro-punctata. Flores variè dispositi.

Sect. I. ASCYREIA. *Chois. prod. hyp. 38.*

Sepala basi coalita et inæqualia. Stamina ∞ . Styli 3-5. — Suffrutices. Folia sæpè maxima. Flores vulgò terminales, magni, pauci.

* *Stylis vulgò 3.*

1. *H. ELATUM* (Ait. h. kew. ed. 2. v. 3. p. 104.) caule juniore alato-rubricante, foliis ovato-oblongis acutis basi dilatatis submarginatis margine subrevolutis, floribus corymbosis, pedunculis 2-bracteatis, sepalis ovato-oblongis. ζ in Americâ boreali. Juss! ann. mus. 3. p. 162. t. 17. (v. s.)
2. *H. FRONDOSUM* (Michx! fl. bor. am. 2. p. 81.) ramis ancipitibus, foliis ovato-oblongis apice obtusis basi angustatis, floribus magnis subsolitariis, calyce petala æquante etiam superante, stylis coadunatis. \mathcal{L} . ad saxa prope fluvium Tennassée. (v. s.)
3. *H. AMOENUM* (Pursh fl. bor. am. 2. p. 375.) ramis ancipitibus, foliis oblongo-ellipticis apice obtusiusculis basi attenuatis margine crispato-revolutis, floribus terminalibus subsolitariis calyce ovato-acuminato corollam nunquàm æquante, stylis coadunatis. \mathcal{L} in Carolinâ et Georgiâ. (v. s.)
4. *H. GRANDIFOLIUM* (Chois. prod. hyp. p. 38. t. 3.) caule tereti rubescente, foliis ovato-oblongis cordato-amplexicaulibus apice acutis reticulato-pellucidis, floribus corymbosis, pedunculis 2-bracteatis, calyce subacuto post inflorescentiam in pedunculum reflexo corollâ multò breviora. ζ in ins. Teneriffâ. (v. s.)
5. *H. HIRCINUM* (Linn. sp. 1103.) ramis alatis, foliis basi submarginatis dilatatis sessilibus apice acutis ovato-lanceolatis margine glandulosis, pedunculis 2-bracteatis, staminibus corollam superantibus, seminibus 2 appendiculatis. ζ in regionibus Mediterraneis. Schk. bot. handb. 3. t. cxxiii. 3.
 β . *obtusifolium*, foliis multò obtusioribus. ζ in petrosis humidis montium Corsicorum. (v. v.)
6. *H. INODORUM* (Willd. sp. 3. p. 1449.) caule suffruticoso tereti, ramis subulatis, foliis ovatis obtusissimis, calyce lanceolato acuto, pedunculis aliquandò 2-bracteatis, staminibus corollam æquantibus. ζ in Græciâ, Oriente. Herba inodora. (v. s.)
7. *H. FOLIOSUM* (Ait. h. kew. ed. 1. v. 3. p. 104.) ramis alatis, foliis sessilibus apertis ovato-oblongis subacutis tenuiter perforatis, calyce lanceolato caduco. ζ in insulis Açoris.
8. *H. FLORIBUNDUM* (Ait. h. kew. ed. 1. v. 3. p. 104.) caule tereti, foliis sessilibus lanceolatis impunctatis numerosis, pedunculis apice dilatatis subcompressis, calyce obtuso, corollâ staminibusque marcescentibus. ζ in ins. Canariis et Maderâ. H. frutescens, etc. Comm. h. amst. p. 137. t. 68. (v. v.)
9. *H. OLYMPICUM* (Linn. sp. 1102.) caule tereti, foliis elliptico-ovatis subobtusis pellucido-punctatis, calyce ovato acuto, pedunculis 2-bracteatis, corollâ et staminibus marcescentibus. ζ ad montem Olympum, et in Chinâ. Smith exot. bot. 2. p. 71. t. 96. Sims bot. mag. t. 1867. (v. s.)
10. *H. CANARIENSE* (Linn. syst. p. 575.) caule obsolete quadrangulo, ramis compressis, foliis ovato-lanceolatis acutis, calyce ovato obtuso, stylis 3-4 divergentibus. ζ in insulis Canariis. (v. v.)
 β . *triphyllum*, foliis ternis.
 γ . *salicifolium*, foliis lineari-lanceolatis longioribus.

** *Stylis vulgò 5.*

11. *H. CHINENSE* (Linn. amœn. 8. p. 323.) caule tereti, foliis ellipticis obtusis paulò nigro punctatis, pedunculis 2-bracteatis, calyce oblongo obtuso nigropunctato, stylis coadunatis. ζ in Indiâ, Japoniâ, Chinâ, et ad cap. Bonæ-Spei. *H. monogynum* Linn. sp. 1101. Curt. bot. mag. 334. (v. v.)
12. *H. CORDIFOLIUM* (Chois. mss.) caule tereti fruticoso, ramis teretibus, foliis ovatis amplexicaulibus cordatis impunctatis confertis, floribus paucis, calyce lanceolato acuto impunctato, staminibus brevibus, stylis liberis rectis. ζ in Napauliâ. Wallich. (v. s.)
13. *H. ALTERNIFOLIUM* (Vahl symb. 2. p. 85. t. 42.) ramis teretibus, foliis alternis lanceolatis, pedunculis 3-bracteatis, stylis reflexis. — in Indiâ orient.
14. *H. PYRAMIDATUM* (Ait. h. kew. ed. 1. v. 3. p. 103.) caule alato, foliis amplexicaulibus oblongo-lanceolatis acutis margine revolutis, pedunculis brevibus crassis. calyce subobtusato, stylis 5-7 brevibus crassis basi coadunatis. \mathcal{L} in Americâ boreali. Vent. rcalm. t. 118. (v. s.)
15. *H. ROSTRATUM* (Rafin. fl. lud. p. 87.) caule frutescente ancipiti, foliis sessilibus ovatis utrinquè acutis margine revolutis punctatis, corymbis axillaribus terminalibusque, calyce 2-bracteato, ovario rostrato 5-stylo. \mathcal{L} in Louisianâ.
16. *H. ASCYRON* (Linn. sp. 1102) caule tetragono herbaceo simplici erecto, foliis amplexicaulibus lanceolatis acutis pellucido-punctatis, floribus paucis, calyce obtuso, stylis basi coadunatis. \mathcal{L} in Sibirîâ. Gmel. fl. sib. 4. p. 178. t. 69. (v. s.)
17. *H. ASCYROIDES* (Willd. sp. 3. p. 1443.) caule basi alato apice tetragono herbaceo simplici, foliis oblongis lanceolatis acutis, calyce ovato-lanceolato, stylis liberis longitudine staminum. \mathcal{L} in Pensylvaniâ. (v. s.)
18. *H. LANCEOLATUM* (Lam. dict. 4. p. 145.) caule tereti fruticoso, foliis lanceolatis margine nigro punctatis apice subobtusis, floribus solitariis, calyce leviter obtuso, stylis coadunatis. ζ ad insulas Borboniæ et Caffrorum. Chois. prod. hyp. p. 41. excl. syn. Jacq. et Cat. (v. s.)
19. *H. ANGUSTIFOLIUM* (Lam. dict. 4. p. 145.) caule tereti fruticoso, foliis lineari-lanceolatis approximatis margine revolutis impunctatis, floribus solitariis maximis, calyce subacuto, stylis coadunatis. ζ in ins. Borboniâ. (v. s.)
20. *H. PATULUM* (Thunb. jap. p. 295. ic. 17.) caule tereti suffruticoso purpurascente, foliis ovato-lanceolatis acutis basi attenuatis margine revolutis obscure pellucido-punctatis, floribus corymbosis, pedunculis 2-bracteatis, calyce obtusissimo suborbiculati. ζ in Japoniâ et Napauliâ. Hyp. Uralum Don. mss. in Curt. bot. mag. t. 2375 (v. s.)
 \mathcal{L} . *attenuatum*, omnibus partibus brevioribus tenuioribus confertioribus, ramulis compressis.
21. *H. KALMIANUM* (Lam. dict. 4. p. 148.) ramis tetragonis, foliis lineari-lanceolatis, floribus 3-7 in corymbo terminali, calyce lanceolato subobtusato. ζ in Virginiâ.
22. *H. VENOSUM* (Lam! dict. 4. p. 146) caule tetragono erecto? rubicante simplici, foliis ovato-oblongis leviter amplexicaulibus subobtusis. floribus terminalibus maximis, pedunculis bracteatis, calyce ovato rotundato, stylis staminibus superantibus. ζ hab. An diff. ab *H. calycino*? (v. s.)
23. *H. LENSCHENAUThii* (Chois. mss.) caule tereti suffruticoso, foliis oblongo-ellipticis obtusis pellucido-punctatis, calyce ovato-lanceolato acuto, staminibus brevissimis, stylis staminibus superantibus corollâque brevioribus. ζ in Javâ. Sequenti maximè affine. (v. s.)
24. *H. OBLONGIFOLIUM* (Chois. prod. hyp. p. 42. t. 4.) caule tereti suffruticoso,

ramis compressis, foliis elliptico-lanceolatis confertis margine paulò revolutis tenuissimè pellucido punctatis, calyce oblongo subobtusò, stylis staminibus superantibus. ♀ in Indià, Napaulià. (v. s.)

25. *H. CALYCINUM* (Linn. mant. 106.) caule tetragono humifuso, foliis ovatis coriaceis pellucido-punctatis, floribus solitariis maximis, calyce maximo post inflorescentiam patulo ovato, capsulâ nutante. ♀ in Oriente, monte Olympo etc. Curt. bot. mag. 146. (v. v.)

β. *acutifolium*, foliis magis acutis ad basin ramulorum usquè decrescen-
tibus.

26. *H. BALEARICUM* (Linn. sp. 1101.) caule quadrangulo verrucoso, foliis ovatis obtusis subamplexicaulibus verrucosis. ♀ ad ins. Majorcam. Curt. bot. mag. t. 137. (v. v.)

Sect. II. TRIDESMOS. *Chois. mss.*

Calyx 5-sepalus, sepala æqualia integra. Stamina altissimè 3-adelpha; adelphiæ apice subpenicilliformes. Styli 3. — Suffrutices. Flores axillares longè pedunculati.

27. *H. BIFLORUM* (Lam. dict. 4. p. 171.) caule glabro subtereti griseo, foliis ovato-ellipticis glabris acutiusculis petiolatis venosis, calyce ovato obtuso; stylis stamina æquantibus. ♀ in Chinâ, prope fretum Bouton. *H. Chinense* Retz obs. bot. 5. p. 27. (v. s.)

28. *H. ARTICULATUM* (Lam. dict. 4. p. 159.) caule tereti aut subtetragono griseo, foliis ovato-oblongis aut ellipticis obtusis petiolatis venosis subtus punctatis, calyce ovato obtuso, adelphiis tomentosis, valvulis fructûs carinatis. ♀ in ins. Madagascar. (v. s.)

Sect. III. ELODEA. *Chois. prod. hyp. 43.*

Calyx 5-sepalus, sepala æqualia integra. Styli 3. Filamenta numero subdefinita 9-15-18 altissimè coalita. — Herbæ purpurascentes. Flores aliquandò rubri axillares aut confertim terminales.

29. *H. PALUDOSUM* (*Chois. prod. hyp. p. 43.*) caule herbaceo tereti, foliis oblongis obtusis in petiolum attenuatis pellucido-punctatis, floribus in pedunculis axillaribus elongatis, calyce obtuso, staminibus 9-10. ♀ in paludibus Carolinæ. *H. petiolatum* Walth. car. 191. *Elodea petiolata* Pursh am. bor. 2. p. 379. (v. s.)

30. *H. VIRGINICUM* (Linn. sp. 1104.) caule tereti suffruticoso, foliis oblongis obtusis subamplexicaulibus pellucido-punctatis, floribus in pedunculis axillaribus, calyce lanceolato, staminibus 9-12. ♀ in Americâ boreali. Andr. bot. rep. t. 552. (v. s.)

31. *H. TUBULOSUM* (Walth. car. 191.) foliis sessilibus, corollis tubulosis?, staminibus plus quàm ad medium connatis. ♀ in Carolinâ. *Elodea tubulosa* Pursh fl. am. bor. 2. p. 379.

Sect. IV. PERFORARIA. *Chois. prod. hyp. p. 44.*

Calyx 5-sepalus; sepala æqualia, nunc integra, nunc dentata, aut dentato-glandulosa. Stamina ∞. Styli vulgò 3. — Herbæ aut suffrutices. Flores axillares, aut paniculati. Folia rarò linearia.

* *Sepalis integris.*

32. *H. MICRANTHUM* (*Chois. prod. hyp. p. 44. t. 5.*) caule tereti punctato purpurascente, ramplis rectis, foliis oblongis obtusis nigro-punctatis, floribus confertis terminalibus, calyce minimo obtuso, corollâ nigro-punctatâ, antheris nigro-punctatis. ♀ in Carolinâ. (v. s.)

33. *H. ANGULOSUM* (Michx. bor. am. 2. p. 78.) caule herbaceo quadrangulo erecto, foliis distantibus elongatis ovatis subamplexicaulibus margine sinuatis acutis impunctatis, floribus axillaribus solitariis, calyce lanceolato acuto, stylis coadunatis. ♀ in aquaticis Carolinæ. (v. s.)

34. *H. ULIGINOSUM* (H. B. et Kunth! nov. gen. am. 5. p. 194.) caule herbaceo stricto tetragono, foliis oblongis acutiusculis superioribus lanceolatis punctulatis, calyce lineari-lanceolato acuminato, stylis 3-4, stigmatibus capitatis. \mathcal{L} prope la Venta grande Caracasanorum. (v. s.)
ß. multiflorum, floribus minoribus copiosioribus. — in monte Saraguru prope Loxam.
35. *H. PUNCTATUM* (Lam! dict. 4. p. 164.) caule tereti nigro punctato, foliis ovato-lanceolatis subacutis amplexicaulibus nigro-punctatis floribus corymbosis, calyce lanceolato et corollâ nigro-punctatis. \mathcal{L} in Carolinâ. H. maculatum Walth. car. 189. H. corymbosum Willd. sp. 3. p. 145. (v. s.)
36. *H. FORMOSUM* (H. B. et K! nov. gen. am. 5. p. 196. t. 460.) caule herbaceo tereti impunctato glabro, foliis ovato-oblongis obtusis subamplexicaulibus subtus punctulatis, floribus corymbosis, calyce ovato-lanceolato petalisque obovato-cuneatis glanduloso-punctatis. \mathcal{L} prope Pazcuaro. Vix diff. à præcedente. (v. s.)
37. *H. DOLABRIFORME* (Vent! h. cels. p. 45.) caule erecto purpureo, foliis lineari-lanceolatis reflexis pellucido-punctatis, floribus corymbosis, pedunculis dichotomis, calyce inæquali acuto reflexo punctato margine revolutato. \mathcal{L} in Kentucky. (v. s.)
38. *H. CISTIFOLIUM* (Lam! dict. 4. p. 158.) caule anguloso, foliis ovato-oblongis, subtus nigro-punctatis acutiusculis subamplexicaulibus margine revolutis, floribus in corymbis dichotomis, calyce ovato, stylis coadunatis. \mathcal{L} in Amer. boreali. (v. s.)
39. *H. DENSIFLORUM* (Pursh bor. am. 2. p. 374.) ramosissimum, ramis subteretibus, foliis lineari-lanceolatis obtusiusculis basi attenuatis, calyce brevissimo ovato deciduo, stylis coadunatis. \mathcal{L} in Virginie montibus.
40. *H. PROCUMBENS* (Mich! bor. am. 2. p. 81.) caule procumbente subtetragono herbaceo, foliis lineari-lanceolatis subobtusis margine revolutis pellucido-punctatis, paniculâ terminali dichotomâ, calyce cuneato-lanceolato acuto, stylis coadunatis. \mathcal{L} in Amer. boreali. (v. s.)
41. *H. ROSMARINIFOLIUM* (Lam! dict. 4. p. 159.) caule tereti recto, foliis obtusis amplexicaulibus ovatis margine revolutis, calyce lineari-obtusato punctato, stylis coadunatis. \mathcal{L} in Carolinâ. (v. s.)
42. *H. VIRGATUM* (Lam! dict. 4. p. 158.) caule recto quadrangulo, foliis ovato-lanceolatis leviter amplexicaulibus nigro-punctatis margine revolutis, paniculâ pauciflorâ dichotomâ, calyce lanceolato, stylis 2-3, stigmatibus capitatis. \mathcal{L} in Amer. boreali. (v. s.)
43. *H. NEDYOTIFOLIUM* (Poir. suppl. dict. 3. p. 700.) caule recto quadrangulo, foliis sessilibus decussatis lanceolatis subacutis adpressis non pellucido-punctatis subtus nigro-punctatis, calyce lineari-lanceolato nigro-punctato, stylis 3-4-stigmatibus capitatis. \mathcal{L} in Amer. boreali. (v. s.)
44. *H. MYRTIFOLIUM* (Lam! dict. 4. p. 180.) caule tereti, foliis ovato-cordatis amplexicaulibus aut cuneato-lanceolatis margine revolutis, floribus cymoso-dichotomis, calyce subinæquali ovato corollam superante, stylis coadunatis. \mathcal{L} in Amer. boreali. (v. s.)
45. *H. BRASILIENSE* (Chois. mss.) caule herbaceo glabro, ramulis tetragonis, foliis lineari-lanceolatis acutiusculis patulis leviter nigro-punctatis, paniculis dichotomis paucifloris, calyce lineari-subulato, corollâ parvâ, stylis 5 liberis rectis. \mathcal{L} ad Rio-Janeiro. (v. s.)
46. *H. PROLIFICUM* (Linn. mant. 106.) caule tereti, ramulis angulosis, foliis lineari-lanceolatis margine revolutis pellucido-punctatis, floribus in corymbo paucifloro, calyce ovato-lanceolato, staminibus numerosissimis, stylis sæpè coadunatis. \mathcal{L} in Amer. boreali. H. foliosum Jacq. h. schœnb. 3. p. 27. t. 299. (v. s.)
47. *H. CLAUCEUM* (Mich! bor. am. 2. p. 78.) caule tereti, foliis cordatis am-

plexicaulibus obtusis margine revolutis glaucis pellucido-punctatis, floribus in paniculis folioissimis, calyce ovato obtuso, staminibus numerosissimis. \mathcal{L} in Americâ boreali. (v. s.)

48. *H. nudiflorum* (Michx. bor. am. 2. p. 78.) caule tetragono alatoque, foliis ovato-oblongis obtusis acupunctatis non pellucido-punctatis, paniculâ florum nudâ, calyce acuto lineari-lanceolato, stylis coadunatis. \mathcal{L} in Americâ bor. (v. s.)

β . *ovatum*, calyce ovato, foliis amplexicaulibus. *H. lævigatum*? Ait. h. kew. ed. 2. v. 4. p. 425.

γ . *ramosum*, caule magis ramoso et folioso.

49. *H. sphærocarpon* (Michx. bor. am. 2. p. 78.) caule anguloso erecto, foliis lineari-lanceolatis obtusis sessilibus impunctatis, paniculâ nudâ dichotomâ, calyce lineari-lanceolato acuto, stylis coadunatis, capsulâ globosâ. \mathcal{L} in Americâ boreali, Kentucky. (v. s.)

50. *H. gramineum* (Forst! prod. 53.) caule tetragono herbaceo dichotomo, foliis ovato-lanceolatis amplexicaulibus sessilibus non pellucido-punctatis, pedunculis terminalibus solitariis unifloris, calyce lineari-lanceolato, stylis brevibus liberis, stigmatibus capitatis. \mathcal{L} ad Novam-Caledoniam. (v. s.)

51. *H. fastigiatum* (H. B. et Kl. nov. gen. am. 5. p. 195.) caule herbaceo recto superne ramoso, foliis oblongo-lanceolatis margine revolutis, calyce lineari-lanceolato, stigmatibus subinfundibuliformibus, ramulis fastigiatis. \mathcal{L} in Mexico prope Pazcuaro. (v. s.)

52. *H. quadrangulum* (Linn. sp. 1104.) caule herbaceo quadrangulo recto ramoso, foliis ovatis obtusis pellucido-punctatis margine nigro-punctatis, florum paniculis multifloris, calyce lanceolato, antheris nigro-punctatis. \mathcal{L} in humidis Europæ et in montibus Delphinatûs, Jurassi. etc. — Fl. dan. t. 640. (v. s.)

β . *dubium*, caule minus ramoso, calyce vix punctato, corollâ nigro-punctatâ, foliis vix pellucido-punctatis minus obtusis. *H. Delphinense* Vill. delph. 3. p. 497. t. 44.

γ . *maculatum*, caule magis ramoso obsolete quadrangulo contorto, foliis pluribus vix perforatis, floribus numerosioribus, corollâ variè nigro-punctatâ. *H. maculatum* All. ped. n. 1433. t. 83. f. 1.

δ . *confertum*, caule subsimplici alato non nigro-punctato, foliis brevibus confertioribus variè nigro-punctatis, calyce magis acuto pellucido-punctato, corollâ impunctatâ.

ϵ . *undulatum*, foliis margine undulatis. *H. undulatum* Schousb. in Willd. enum. p. 81 f.

53. *H. attenuatum* (Chois. prod. hyp. p. 47. t. 6.) caule tereti nigro-punctato recto, foliis ovato-oblongis obtusis subamplexicaulibus nigro-punctatis, calyce lanceolato nigro-punctato, corollâ apice nigro-punctatâ. \mathcal{L} ad Sibiriam, Dahuriam. (v. s.)

54. *H. repens* (Linn. sp. 1103.) caule suffruticoso tereti purpurascente ascendente et prostrato, foliis ovato-linearibus obtusis approximatis vix perforatis, corymbo paucifloro, calyce ovato-lanceolato acuto nigro-punctato. \mathcal{L} in Oriente et in Europæ regione mediterraneâ. (v. s. sp. in h. Balbis è regione inter Cannes et Antibes.)

55. *H. connatum* (Lam. dict. 4. p. 168.) caule suffruticoso tereti, foliis deltoideis perfoliatis obtusis coriaceis subtus punctatis et venoso-reticulatis margine inflatis, calyce lanceolato acuto, stylis 5, stigmatibus capitatis. \mathcal{L} in Monte-Video. (v. s.)

56. *H. erectum* (Thunb. fl. jap. p. 296.) caule herbaceo tereti purpureo recto, foliis lanceolatis acutis amplexicaulibus margine revolutis, calyce lanceolato. \mathcal{L} ad montes Nagasaki.

57. *H. japonicum* (Thunb. fl. jap. p. 295. t. 31.) caule debili tetragono glabro de-

cumbente, foliis ovatis subcordatis obtusis margine revolutis subtus vix punctatis, floribus laxè pedunculatis, calyce oblongo acuto corollam et stamina adæquante, stylis 3 brevibus divergentibus. \mathcal{L} ad insulam Nipon Japoniæ et in Napanliâ. (v. s.)

β . *ramosum*, caule ramoso, foliis floribusque confertis.

58. *H. dichotomum* (Lam. dict. 4. p. 167.) caule basi suffruticoso tereti dichotomè ramosissimo, foliis minimis lineari-elongatis obtusis basi attenuatis non nigro-punctatis obscure pellucido-punctatis, calyce ovato-lanceolato acuto staminibus 9 liberis? \mathcal{L} in Sancto-Domingo. An ad *Elodeas* referendum. (v. s.)

59. *H. crispum* (Linn. mant. 106.) caule tereti ramosissimo, foliis sessilibus lanceolatis basi undulato-sinuatis pellucido-punctatis, calyce minimo obtuso. \mathcal{L} in regionibus mediterraneis. — Bocc. mus. 2. p. 34. t. 12. (v. s.)

60. *H. pilosum* (Walth. car. 190.) caule virgato simplici, foliis dilatatis ovatis aliquandò basi attenuatis, paniculis terminalibus panicifloris. \mathcal{J} in Americâ boreali. *H. Virginianum*. etc. Pluk. alm. t. 245. f. 6. (v. s.)

61. *H. simplex* (Michx. bor. am. 2. p. 80.) caule simplici herbaceo tereti lacunosus, foliis oblongis cauli adpressis canaliculatis pubescentibus, floribus paucis subsessilibus, calyce ovato ciliolato, stylis 3 liberis, stigmatibus subcapitatis. \odot in Amer. boreali. Pluk. amalg. p. 120. t. 421. f. 3. (mala.) (v. s.)

62. *H. heterophyllum* (Vent! h. cels. t. 68.) caule suffruticoso tereti, foliis lineari-lanceolatis pellucido-punctatis, inferioribus confertis imbricatis brevissimis obtusis, calyce acuto subinæquali. \mathcal{J} in Persiâ. (v. s.)

63. *H. ægyptiacum* (Linn. sp. 1103.) caule fruticoso tereti, foliis minimis ovatis confertis impunctatis, floribus paucis subsessilibus, calyce acuto lanceolato, stylis minimis divergentibus. \mathcal{J} in Ægypto. Linn. amœn. 8. p. 323. t. 8. f. 3. *H. maritimum* Sieb. exs. (v. v.)

64. *H. nanum* (Poir. suppl. dict. 3. p. 699.) caule suffruticoso, foliis ovato-subrotundis obtusissimis mucronulatis pellucido-punctatis, calyce tenuissimo acuto lanceolato. \mathcal{J} ad Syriam. (v. s.)

65. *H. cuneatum* (Poir! suppl. dict. 3. p. 699.) caule herbaceo, ramulis teretiusculis glabris, foliis obovatis basi cuneatis obtusis subpetiolatis margine parçè nigro-glandulosis, floribus paucis, calyce lineari-lanceolato corollam superante, staminibus 15-20. \mathcal{L} in Oriente (v. s.)

66. *H. humifusum* (Linn. sp. 1103.) caule herbaceo humifuso, foliis oblongis obtusis tenuissimè pellucido-punctatis et nigro-punctatis, floribus cymosis terminalibus, calyce lineari-lanceolato corollam superante, staminibus 15-20. \mathcal{L} in arenosis Europæ totius et Caucasi. — OEd. fl. dan. t. 141. (v. s.)

β . *Liottardi*, caule magis minuto erectiore bisannuo, floribus aliquandò 4-partitis, calyce longiore. *H. Liottardi* Vill. dauph. t. 44.

67. *H. pusillum* (Chois. prod. hyp. p. 50.) caule debili prostrato tetragono, foliis ovatis obtusis pellucido-punctatis, calyce lanceolato, petalis rectis, stigmatibus capitatis. \mathcal{L} in Australasiâ. *Ascyrum humifusum*. Lab. ill. nov.-holl. 2. p. 33. t. 175. (v. s.)

68. *H. involutum* (Chois. prod. hyp. p. 50.) caule ascendente 4-alato tenui, foliis ovato-oblongis obtusis pellucido-punctatis, paniculâ laxâ dichotomâ, calyce lanceolato acuto, corollâ involutâ, stigmatibus capitatis. \mathcal{L} in Australasiâ. *Ascyrum involutum* Lab. ill. nov.-holl. 2. p. 32. t. 174. (v. s.)

69. *H. asperum* (Lam. dict. 4. p. 166.) caule suffruticoso tereti alato, foliis ovatis obtusis, tenuissimè obtusis pellucido-punctatis margine et subtus nigro-punctatis, calyce acuto subinæquali quandòque serrato. \mathcal{J} ad Barbariam, Hispaniam, Syriam. (v. s.)

70. *H. perforatum* (Linn. sp. 1105.) caule ancipiti, foliis obtusis ovato-ellipticis pellucido-punctatis, floribus paniculatis, calyce lanceolato pellucido-punctato, antheris nigro-punctatis, stylis divergentibus. \mathcal{L} Ubiquè vulgatissimum. Turp. in dict. sc. nat. ic. opt. (v. v.)

β. *latiglandulosum*, glandulis latioribus paucioribus, foliis apice confertioribus, paniculis strictioribus paucifloris.

γ. *elatum*, caule elatiore, internodiis distantioribus, foliis minoribus margine revolutis, ramis rectioribus, paniculis laxis paucifloris.

δ. *punctatum*, caule humiliore, calyce obtusiore aliquandò margine glanduloso, corollâ nigro-punctatâ.

ε. *microphyllum*, omnibus partibus minoribus confertioribus, paniculis strictis multifloris.

ζ. *albiflorum*, floribus albis.

71. H. QUINQUEMNERVIUM (Walth. car. 190.) caule herbaceo tetragono, foliis subamplexicaulibus ovatis obtusis obscure 5-nerviis pellucido-punctatis subtus acropunctatis, corymbo dichotomo, calyce lineari-lanceolato. 2℥ in Americâ bor. et monte Quindiu. H. B. K! nov. gen. am. 5. p. 196. H. parviflorum Willd. sp. 3. p. 1456. (v. s.)

72. H. TRYMPOLIUM (H. B. K! nov. gen. am. 5. p. 186. t. 455.) caule fruticoso, ramis dichotomis, foliis oblongis parvis obtusiusculis pellucido-punctatis, calyce oblongo, stylis liberis, stigmatibus peltatis. 5 prope Santa-Fé de Bogota? (v. s.)

73. H. BREVISSYLUM (Chois. prod. hyp. p. 51. t. 7.) caule herbaceo prostrato tenni, foliis oblongo-lanceolatis subobtusis pellucido-punctatis, paniculis paucifloris, calyce lanceolato, stylis coadnatis brevissimis, stigmatibus capitatis. 2℥ in Amer. meridionali. (v. s.)

74. H. DENTICULATUM (H. B. K! nov. gen. am. 5. p. 191. t. 458.) caule herbaceo ascendente, foliis lineari-lanceolatis margine retrorsum denticulatis, floribus decandris, calyce oblongo-lanceolato, petalis obovato-cuneatis, stylis liberis brevissimis, stigmatibus subcapitatis. 2℥ in Mexico prope Guanajuato. (v. s.)

75. H. LALANDII (Chois. mss.) caule herbaceo tetragono nigro-punctato, foliis lanceolatis acutis margine revolutis cauli adpressis subamplexicaulibus, paniculâ pauciflorâ dichotomâ, calyce lineari-lanceolato corollam superante, stylis 3 liberis. 2℥ ad cap. Bonæ-Spei. (v. s.)

76. H. SILENOIDES (Juss! ann. mus. 3. p. 162. t. 16. f. 3.) caule herbaceo, foliis lanceolatis margine revolutis pollicaribus, floribus secundis, calyce angusto acuto glanduloso. 5 in Peruvîâ et Andibus Novo-granatensibus. (v. s.)

77. H. IMECORUM (H. B. K! nov. gen. am. 5. p. 193.) caule herbaceo basi ramoso, ramis elongatis tetragonis, floribus sub-5-andris, foliis oblongo-lanceolatis acutis punctulatis, stigmatibus subcapitatis. 2℥ prope Loxam. (v. s.)

β. *paniculatum*, caule stricto multifloro, floribus minoribus 5-7-andris, petalis linearibus. — prope Ario. H. paniculatum H. B. K! l. c. p. 195. t. 459.

78. H. CANADENSE (Linn. sp. 1104.) caule herbaceo recto 4-alato, foliis linearibus subobtusis tenuissimè pellucido-punctatis subtus nigro-punctatis, paniculâ elongatâ dichotomâ, calyce lanceolato, stylis brevissimis, capsulâ conicâ. 2℥ ad Canadam et in regno Mexicano. (v. s.)

β. *minimum*, totâ staturâ minore, floribus paucioribus. Varietates hujus mihi videntur H. thesiifolium, pauciflorum et Moranense. H. B. et K! nov. gen. 5. p. 192. et 193.

79. H. TARQUENSE (H. B. K! nov. gen. am. 5. p. 193.) caule herbaceo recto ramoso dichotomo corymboso-multifloro, foliis lanceolato-oblongis punctulatis margine revolutis, calyce lanceolato acuminato, stylis 3-5, stigmatibus subcapitatis. 2℥ in locis montosis Quito. (v. s.)

80. H. GALIOIDES (Lam. dict. 4. p. 161.) caule suffruticoso tereti recto, foliis lineari-lanceolatis basi attenuatis apice latioribus acutis margine revolutis punctatis, calyce acuto post inflorescentiam reflexo, stylis primum coadnatis, capsulâ conicâ acutissimâ. 5 in Amer. boreali. (v. s.)

81. H. AXILLARE (Lam! dict. 4. p. 160.) caule fruticoso tereti diffuso, foliis lanceolato-linearibus basi angustatis margine subtus revolutis, calyce subinæ-

- quali, stylis coadunatis postea liberis. ζ in Amer. boreali. H. fasciculatum Willd. Pursh Nutt. (v. s.)
82. H. ORIGANIFOLIUM (Willd. sp. 3. p. 1467.) caule ascendente tomentoso, foliis ovatis obtusis pellucido-punctatis pubescentibus, floribus magnis paucis, calyce ovato obtuso glabro apice paulo nigro-punctato, corollâ nigro-punctatâ, staminibus numerosissimis. \mathcal{L} in Oriente, ad Constantinopolim. (v. s.)
 ** *Sepalis dentatis sæpè dentato-glandulosis.*
83. H. ELODES (Linn. sp. 1106.) caule herbaceo villosa tereti, foliis subrotundis ovatis sessilibus supra emarginatis, paniculis laxis paucifloris, calyce serrato-glanduloso lineari-acuto, staminibus paucis, stigmatibus capitatis. \mathcal{L} in paludosis Europæ borealis. Schk. bot. hand. 3. ccxiii. 5. (v. s.)
84. H. TOMENTOSUM (Linn. sp. 1106.) caule herbaceo tomentoso tereti ascendente, foliis ovatis obtusis subamplexicaulibus margine nigro-punctatis, paniculis laxis dichotomis, calyce acuminato, stigmatibus simplicibus. \mathcal{L} in pratis humidiusculis regionis Mediterraneæ. H. supinum, etc. Moris. hist. 2. p. 470. s. 5. t. 6. n. 5. (v. s.)
85. H. LANUGINOSUM (Lam. dict. 4. p. 171.) caule suffruticoso tereti lanuginoso recto, foliis amplexicaulibus ovatis obtusis margine nigro-punctatis, corymbo florum vasto dichotomo multifloro, calyce obtuso, antheris nigro-punctatis. \mathcal{L} in Oriente, Græciâ. (v. s.)
86. H. HIRSUTUM (Linn. sp. 1105.) caule tereti suffruticoso hirsuto recto, foliis ovato-oblongis nervatis pellucido-punctatis, paniculâ longâ racemosâ, calyce lineari subacuto, stylis divergentibus. \mathcal{L} in umbrosis Europæ, Caucasii, etc. — *Æd. fl. dan. t. 802.* (v. v.)
87. H. NUMMULARIUM (Linn. sp. 1106.) caule tereti ascendente herbaceo, foliis orbiculatis petiolatis, calyce ovato obtuso. \mathcal{L} in saxosis montanis Galliæ meridionalis, Pedemontii, etc. Lam. ill. t. 643. (v. s.)
88. H. ELEGANS (Steph. in Willd. sp. 3. p. 1469.) caule herbaceo recto leviter alato nigro-punctato, foliis ovato-lanceolatis subamplexicaulibus obtusiusculis pellucido punctatis, calyce ovato lanceolato acuto pellucido-punctato, antheris nigro-punctatis. \mathcal{L} in Sibiria et prope Hales. H. Kohliannu Spreng. fl. hal. n. 864. t. 9. (v. v. in h. Genev. missum à clar. Steven et apprimè congruens cum specimine H. Kohliani in herb. DC.)
89. H. GLANDULOSUM (Ait. h. Kew. ed. 1. v. 3. p. 107.) caule fruticoso tereti recto ramoso, foliis elliptico-lanceolatis acutis margine glandulosis pellucido-punctatis, calyce lanceolato acuto. ζ in Maderâ, Teneriffâ. (v. s.)
90. H. REFLEXUM (Linn. f. suppl. p. 346.) caule fruticoso tereti apice villosulo, foliis amplexicaulibus lanceolatis acutis pellucido-punctatis margine nigro-punctatis sæpè reflexis, paniculis laxis paucifloris, calyce lineari-lanceolato. ζ in Teneriffâ. (v. s.)
91. H. PULCHRUM (Linn. sp. 1106.) caule herbaceo recto tereti, foliis amplexicaulibus cordatis obtusis pellucido-punctatis margine subrevolutis, calyce ovato obtuso, corollâ glandulosâ. \mathcal{L} in umbrosis Europæ. — *Æd. fl. dan. t. 73.* (v. s.)
92. H. ELODEOIDES (Chois. mss.) caule herbaceo tereti glabro simplici impunctato, foliis amplexicaulibus basi dilatatis lanceolatis punctatis, auriculis amplexicaulibus sæpè dentato-ciliatis, paniculis confertis, calyce lineari-lanceolato acuto pellucido-punctato, corollâ apice nigro-punctatâ. \mathcal{L} in Napanliâ. Wallich. (v. s.)
93. H. BARBATUM (Linn. amœn. 8. p. 323.) caule herbaceo tereti, foliis amplexicaulibus oblongo-lanceolatis nigro-punctatissimis, calyce acuto barbato fimbriato, corollâ et antheris nigro-punctatis. \mathcal{L} ad regionem mediterraneam. Jacq. austr. 3. p. 33. t. 259. (v. s.)
ß. Calabricum, foliis strictioribus canli appressis. H. Calabricum Spreng. neue entd. 3. p. 300.

94. *H. DENTATUM* (Lois. fl. gall. p. 499. t. 17.) caule suffruticoso tereti adscendente, foliis amplexicaulibus oblongis subobtusis pellucido-punctatis, superioribus aliquandò dentatis, calyce lanceolato acuto, corollâ et antheris nigropunctatis. \mathcal{L} in insulis Stæchadum. (v. s.)
95. *H. MONTANUM* (Lin. sp. 1105.) caule herbaceo tereti recto, foliis amplexicaulibus ovatis obtusis tenuissimè pellucido punctatis margine nigro-punctatis, internodiis crescentibus, calyce lineari-lanceolato acuto, antheris nigro-punctatis. \mathcal{L} in montanis Europæ. — OEd. fl. dan. n. 173. (v. v.)
 β . *triphyllum*, foliis ternis.
96. *H. FIMBRIATUM* (Lam. dict. 4. p. 148.) caule herbaceo tereti purpurâscente simplici, foliis amplexicaulibus ovatis margine nigro-punctatis, non pellucido punctatis, calyce ovato acuto ciliato, stylis 3-4-5. \mathcal{L} in Alpibus, Jarrasso, Pyrenæis. H. Richeri Vill. delph. 3. p. 501. t. 44. (v. v.)
 β . *androsæmifolium*, caule magis repente, foliis magis ovatis. Vill. delph. 3. p. 502. t. 44. H. alpinum. Waldst. et Kit. pl. rar. hung. 3 p. 294. t. 265.
 γ . *Burseri*, caule altiore, foliis obtusioribus, bracteis longioribus minùs ciliatis, calyce breviorè minùs ciliato. H. Burseri DCI fl. fr. supp. p. 630.
 δ . *pentagynum*, caulibus crassioribus, foliis obtusioribus, calyce longiore, floribus omnibus 5-gynis.
97. *H. ÆTHIOPICUM* (Thunb. prod. 138.) caule herbaceo tereti glabro rubescente, foliis ovatis subamplexicaulibus margine revolutis pellucido-punctatis, paniculâ terminali dichotomâ, calyce acutissimo, corollâ nigro-punctatâ, antheris nigro-punctatis. \mathcal{L} ad Cap. Bonæ-Spei. (v. s.)
98. *H. CILIATUM* (Lam. dict. 4. p. 170.) caule herbaceo tereti leviter alato, foliis amplexicaulibus subcordatis ovato-oblongis obtusis pellucido punctatis margine nigro-punctatis, calyce subacuto ciliato, antheris nigro-punctatis. \mathcal{L} in Europæ regionibus mediterraneis. Androsæmum Sambac perfoliato folio Bocc. mus. 2. t. 127. (v. s.)
 β . *acutifolium*, foliis minoribus acutioribus, floribus longiùs pedicellatis. H. myrti acuto folio Bocc. mus. 2. p. 117 t. 91. §2.
99. *H. SERPYLLIFOLIUM* (Lam. dict. 4. p. 176.) caule suffruticoso tereti, foliis ovatis obtusis minimò petiolo donatis margine revolutis, calyce ovato obtuso. \mathcal{L} in Persiâ. H. folio breviorè Moris. hist. 2. p. 469. s. 5. t. 6. f. 2. (v. s.)
100. *H. CONFERTUM* (Chois. prod. hyp. p. 55. t. 8.) caule herbaceo tereti basi villosò, foliis lanceolatis acutis villosis pellucido-punctatis margine revolutis, calyce lanceolato. \mathcal{L} in Oriente. (v. s.)
101. *H. TRIPLINERVE* (Vent! h. cels. p. et t. 58.) caule herbaceo biangulari basi decumbente, foliis linearibus patentissimis obtusis margine revolutis, calycibus ovatis acutis, petalis inæqualibus. \mathcal{L} in Americâ boreali. Vix diff. à sequente. (v. s.)
102. *H. HYSOPIFOLIUM* (Vill. delph. 3. p. 505. t. 44.) caule suffruticoso tereti adscendente, foliis oblongo-lanceolatis subobtusis utrinquè attenuatis pellucido-punctatis in axillâ fasciculatis, calyce subobtusò, stylis 3-4. \mathcal{L} in Galliâ austr. Tanriâ. (v. s.)
 β . *pauciglandulosum*, foliis latioribus, caule minore, calyce acutiorè longiore minùs glanduloso.
103. *H. LINEARIFOLIUM* (Vahl symb. 1. p. 65.) caule herbaceo recto tereti, foliis linearibus obtusis margine nigro-punctatis non pellucido-punctatis, calyce lanceolato nigro-punctato. \mathcal{L} in Galliâ occidentali et australi. (v. s.)
104. *H. ORIENTALE* (Lin. sp. 1103.) caule fruticoso biangulari gracili recto aut geniculato, foliis amplexicaulibus linearibus obtusis rectis margine capitato-glandulosis, calyce ovato oblongo. \mathcal{L} in Oriente. Tourn! voy. lev. 2. p. t. 220. (v. s.)
105. *H. NAPAULENSE* (Chois. mss.) caule herbaceo tereti tenuissimo simplici

- minimo, foliis ovato-lanceolatis amplexicaulibus confertis margine sæpè ciliato-glandulosos pellucido-punctatis, calyce lineari-lanceolato acutissimo. 2ℓ in Nepauliâ. Wallich. (v. s.)
106. *H. SCABRUM* (Lin. amœn. 4. p. 287.) caule tereti glanduloso-scabro, foliis linearibus glabris acutiusculis, umbellâ terminali densiflorâ, calyce minimo ovato obtuso non glanduloso, stylis 3 liberis. 2ℓ in Arabiâ. (v. s.)
107. *H. CAPITATUM* (Chois. prod. hyp. p. 57. t. 9.) caule herbaceo tereti recto apice subalato, foliis linearibus obtusis margine plicatis pellucido-punctatis in axillâ fasciculatis, calyce minimo obtuso, corollâ nigro-glandulosâ. 2ℓ prope Bagdad. (v. s.)
108. *H. EMPETRIFOLIUM* (Willd. sp. 3. p. 1452.) caule suffruticoso tereti, ramulis subulatis, foliis linearibus ternis margine revolutis, calyce minimo obtuso, corollâ non glandulosâ. 3 in regione Mediterraneâ. (v. s.)
109. *H. CORIS* (Linn. sp. 1107.) caule suffruticoso adscendente tereti, foliis verticillatis linearibus margine revolutis, calyce lineari subobtus. 2ℓ in Oriente et siccis Europæ australis. Curt. bot. mag. 178. *H. multicaule* Lam! dict. 4. p. 178. (v. s.)
110. *H. LUSITANICUM* (Poir! suppl. dict. 3. p. 762.) caule villosa tereti gracili, foliis minimis ellipticis obtusis hispidis, calyce lineari-lanceolato mucronato, antheris nigro-punctatis, stylis 3 filiformibus. 3 in Lusitaniâ.
111. *H. ERICOIDES* (Linn. sp. 1104.) caule suffruticoso tereti tortuoso minuto, foliis teretibus acutis confertissimis punctatis glaucis minimis, calyce acuto vix glanduloso. 3 in Hispaniâ, Lusitaniâ, Oriente. Cav. ic. 2. p. 20. t. 122. (v. s.)

Sect. v. *BRATHYS*. *Chois. prod. hyp.* 58.

Calyx 5-sepalus; sepala integra æqualia foliis sæpius simillima. Stamina ∞. Styli 3-5. — Suffrutices. Flores solitarii axillares; folia imbricata sæpius lineari-subulata, verticillata aut conferta.

* *Stylis 3; stigmatibus simplicibus.*

112. *H. STRUTHIOLEFOLIUM* (Juss! ann. mus. 3. p. 160. t. 16. f. 2.) caule suffruticoso tenui cicatrisato, ramis adpressis, foliis oppositis margine revolutis impunctatis apice imbricatis, calyce lanceolato impunctato, petalis lanceolatis. 3 in Peruviâ. (v. s.)

β. *minutum*, staturâ minore.

γ. *strictum*, ramis longioribus strictis. *H. strictum* H. B. et K! nov. gen. am. 5. p. 190.

113. *H. ACEROSUM* (H. B. et K! nov. gen. am. 5. p. 187. t. 457.) caule fruticoso, ramulis scabris tetragonis, foliis omnibus imbricatis acerosis crassiusculis, calyce lanceolato-oblongo, petalis inæquilateri-obovatis. 3 in Andibus Quintensium. (v. s.)
114. *H. ACICULARE* (H. B. K! nov. gen. am. 5. p. 190.) caule fruticoso, ramulis scabris strictis, foliis acerosis patulis, floribus monogynis? mediocribus, calyce lanceolato, petalis cuspidatis. 3 prope Loxam. An sp. à præc. distincta? (v. s.)
115. *H. LANCEIFOLIUM* (Juss! ann. mus. 3. p. 160. t. 16. f. 1.) caule suffruticoso tereti scabro, ramulis patulis, foliis lanceolatis utrinquè attenuatis confertis, calyce lanceolato impunctato, petalis lanceolatis, stylis longissimis, stigmatibus crassis. 3 in Peruviâ. (v. s.)
116. *H. THUYOIDES* (H. B. K! nov. gen. am. 5. p. 187. t. 456.) caule fruticoso, ramulis scabris tetragonis patulis, foliis decussato-oppositis ovatis acutius-

culis, floribus solitariis, calyce elliptico-oblongo, petalis obovatis, stigmatibus obtusis. ♂ in Andibus Quindiuensibus. (v. s.)

117. *H. CARACASANUM* (Willd. sp. 3. p. 1458.) caule suffruticoso, ramis tetragonis cicatratis, foliis ovato-oblongis acutis dense approximatis subimbricatis, floribus terminalibus postea lateralibus, stigmatibus truncato-incrassatis. ♂ prope Caracas. (v. s.)

118. *H. MEXICANUM* (Linn. amœn. 8. p. 322. t. 8. f. 2.) caule fruticoso, ramis subtetragonis cicatratis, foliis imbricatis ovato-oblongis obtusis flabellonervosis glabris viscosis subtilis punctulatis, floribus solitariis terminalibus, stigmatibus depresso-capitatis. ♂ ad Santa-Fé de Bogota. *H. Mutisianum* H. B. et Kunth! (v. s.)

119. *H. REVOLUTUM* (Vahl symb. 1. p. 66.) caule fruticoso, ramis cicatratis infra subtetragonis apice biangularibus, foliis lineari-lanceolatis approximatis basi revolutis, stylis coadunatis. ♂ ad montes Arabiæ-Felicias. *H. Scabrum?* Linn. amœn. 4. p. 287.

120. *H. NITIDUM* (Lam. dict. 4. p. 160.) caule tereti fruticoso, ramis subangulosis, foliis impunctatis angustissimis, calyce subinæquali, stylis coadunatis. ♂ hab.

121. *H. FASCICULATUM* (Lam. dict. 4. p. 160. non Lapeyr.) caule suffruticoso tereti apice compresso, ramis erectis, foliis densis impunctatis canaliculatis margine subrevolutis, calyce æquali recto, stylis coadunatis. ♂ in America boreali. *H. aspalathoides* Willd. Pursh, Nutt. (v. s.)

♂. *laxifolium*, ramulis angulosis, foliis obtusioribus minus fasciculatis. Mich. bor. am. 2. p. 80. *H. tenuifolium* Pursh fl. bor. am. 2. p. 377. *H. Michauxii* Poir. dict. suppl. 3. p. 696.

** *Stylis 5; stigmatibus capitatis.*

122. *H. BRATHYS* (Lam. dict. 4. p. 152.) caule tereti, ramis adpressis, foliis densis apice canaliculatis non pellucido-punctatis, calyce margine submembranaceo, stylis 5 raro 4. ♂ in Novâ-Granadâ, prope Santa-Fé de Bogota. Smith ic. ined. t. XL. (v. s.)

♂. *juniperinum*, foliolis robustioribus, floribus majoribus. *H. juniperinum* H. B. K! nov. gen. am. 5. p. 189.

† *Species dubiæ.*

123. *H. COCHINCHINENSE* (Lour. fl. coch. 2. p. 577.) floribus 3-gynis, foliis subpetiolatis densissimis, caule arboreo, pedunculis sub-5-floris axillaribus. ♂ in sylvis Cochinchinæ. Flores coccinei. Calyces acutissimi. Stamina 3-adelphæ. Capsula 3-locularis. (Lour.)

124. *H. EMARGINATUM* (Lam! enc. 4. p. 154.) caule fruticoso, floribus 3-gynis, foliis ovato-oblongis obtusis amplexicaulibus emarginatis, calyce lanceolato substriato. — Hab. Inter Elodeas recensendum?

125. *H. VERTICILLATUM* (Thunb. prod. 137.) caule herbaceo, foliis quatuor verticillatis. — Ad cap. Bonæ-Spei. (Thunb.)

126. *H. PULGIDUM* (Rafin. fl. lud. p. 88.) caule frutescente tetragono, ramis gracilibus, foliis sessilibus linearibus patulis, floribus subsessilibus axillaribus terminalibusque. In Louisianâ. — (Rafin.)

††. *Species vix notæ.*

127. *H. Creticum supinum* folio subrotundo flore magno. Tourn. cor. 18.

128. *H. Orientale* foliis subrotundis auritis cauli adhærentibus. Tourn. cor. 18.
 129. *H. Androsæmo* hirsuto simile sed glabrum. Tourn. cor. 18.
 130. *H. linariæfolio*. Tourn. cor. 18.
 131. *H. latifolium* subhirsutum caule purpureo villosa. Tourn. cor. 18.
 132. *H. undulatum* Schousb. Willd. Zucc.
 133. *H. nigricans* Zucc. h. flor.

V. LANCRETIA. *Del. fl. eg. p. 69. t. 25.*

Calyx 4-5-sepalus; sepala æqualia. Petala 4-5. Stamina 10 libera, quorum 5 petalis opposita breviora. Styli 4-5. — Frutex. Folia sessilia dentata seu crenata. Flores terminales.

1. *L. suffruticosa* (Del. l. c.) ♂ in Ægypto, Nubiâ. Ascyroides africanum frutescens Chamædryos folio. Lippi mss. in bibl. Juss., ex cl. Delile. (v. s.)

VI. ASCYRUM. *Linn. gen. 903. Juss. gen. p. 254. Chois. prod. hyp. 60. — Hypericoïdes Adans. fam. 2. p. 443.*

Calyx 4-sepalus; sepala 2 exteriora minora, 2 interiora majora. Petala 4. Stamina ∞ vix basi coalita. Styli 1-3. — Suffrutices. Folia sessilia integra non pellucido-punctata subtus sæpius nigro-punctata. Flores pauci terminales aut axillares.

1. *A. pumilum* (Michx. bor. amer. 2. p. 77.) caule pusillo simplici quadrangulo, foliis ovalibus obtusis fasciculatis, pedicellis 6 lineas longis reflexis, floribus 1-2-gynis. Ad Georgiam. (v. s.)

2. *A. pauciflorum* (Nutt. gen. am. 2. p. 15.) decumbens diffusum tenue pluricaule, foliis approximatis lineari-oblongis obtusis, floribus raris, pedicellis reflexis basi 2-bracteatis, stylo 1. — in sylvis Georgiæ.

3. *A. CRUX-ANDRÆ* (Linn. sp. 1107.) caule suffruticoso tereti, ramis erectis, foliis ovato-linearibus obtusis in axillâ sæpè fasciculatis, calycinis sepalis interioribus suborbiculatis, pedicellis 2-bracteatis, stylis 1-2. ♂ in Virginiâ. *H. frutescens* humifusa Plum. ed. Burm. am. p. 146. t. 152. Chois. prod. hyp. p. 61. (v. s.)

β. *angustifolium*, ramulis magis acutiusculis, foliis angustioribus oblongo-linearibus. *A. multicaule* Michx. *A. hypericoïdes* Ait. h. Kew. An potius sequentis varietas?

4. *A. HYPERICOÏDES* (Linn. sp. 1108.) caule suffruticoso tereti, ramis obliquis, foliis oblongo-linearibus obtusis basi 2-glandulosis, calycis sepalis interioribus suborbiculatis, stylis 3. ♂ in Americâ boreali. *Hypericoïdes* frutescens erecta, etc.? Plum. ed. Burm. am. p. 146. t. 152. f. 1. Chois. prod. hyp. p. 61. (v. s.)

5. *A. STANS* (Michx. bor. am. 2. p. 77.) caule fruticoso alato recto, ramis rectis, foliis ovato-ellipticis obtusis glaucis, calycis sepalis interioribus cordatis orbicularibus, staminibus basi coalitis, stylis 2. — hab. . . Ventx. h. malm. p. 90. *H. floridanum* Pluk. t. 412. f. 5? et *H. tetrapetalum* Lam. dict. 4. p. 153. ex Chois. prod. hyp. p. 61.

6. *A. AMPLEXICAULE* (Michx. l. c.) caule dichotomo paniculato, foliis ovato-cordatis brevioribus crispato-sinuatis, corymbo nudo, stylis 3. ♂ in Carolinâ Pursh. fl. bor. am. 2. p. 374. *H. stans* var. β. Chois. prod. hyp. p. 61. (v. s.)

Subordo II. **HYPERICINÆ ANOMALÆ.** *Chois. prod. hyp.* 61.

Styli ultra 5. Semina plana alata. — **Arbores.**

VII. **CARPODONTOS.** *Lab. nov. holl.* 2. p. 122. *Chois. prod. hyp.* 61.

Styli 5-8. Sepala et petala 4. Capsula lignosa; placentæ filiformes; loculi cymbiformes; ovarium villosum. Stamina ∞ . — Arbor. Folia petiolata opposita. Flores axillares solitarii. Pedunculus basi 2-squamatus.

1. *C. LUCIDA* (Labill! voy. rech. Lapeyr. 2. p. 16. t. 18.) \S ad terram Van Diemen. (v. s.)

VIII. **EUCRYPHIA.** *Cav. ic.* 4. p. 49. t. 372. *Chois. prod. hyp.* 62.

Styli ferè 12. Corolla 5-petala. Calyx 5-sepalus. Stamina ∞ basi subcoalita. Capsulæ cymbæformes è funiculis pendulæ. — Arbor. Folia opposita. Flores axillares solitarii.

1. *E. CORDIFOLIA* (Cav! l. c.) \S prope San Carlos de Chiloe. (v. s.)

CHOISY.

Ordo XXXV. GUTTIFERÆ. *Juss. gen. p. 243. (1)*

Calyx 2-6-sepalus sæpius persistens, sepalis rotundatis membranaceis imbricatim oppositis, quandòque inæqualibus et coloratis. Petala hypogyna 4-10 sæpius lutea. Flores hermaphroditi, monoïci, dioïci aut polygami. Stamina numerosa hypogyna, rariùs definita; filamenta variæ longitudinis; antheræ elongatæ adnatæ longitudinaliter intùs dehiscentes, rarò minimæ et poris duobus similes, in Symphonieis filamentorum dorso adnatæ extùs dehiscentes. Ovarium 1 liberum. Stylus nullus aut unicus sæpius brevis. Stigma sessile peltato-radiatum aut apice styli multilobum. Capsula baccata aut bacca aut drupa pericarpio corticoso crasso multivalvi: valvulæ nunc margine introflexæ placentæ uni centrali aut placentis peculiaribus crassis adfixæ et dissepimenta efformantes, nunc simplices et tunc dissepimenta nulla. Fructus aut 1-locularis oligospermus siccus vel pulpâ repletus in quâ nidulant semina, aut multilocularis loculis mono-aut polyspermis. Spermodermis tenua membranaceis. Albumen nullum. Embryo rectus. Cotyledones crassæ nunc separabiles, nunc in unum coadunatæ.

Frutices et arbores aliquandò parasiticæ, succos resinosos emittentes; folia opposita rarissimè alterna coriacea breviter petiolata integra nervum 1 intermedium aliosque laterales pinnatos referentia. Flores sæpius racemosi axillares, aut paniculati terminales, aut conferti laterales.

Ordo variis suis sectionibus affinis Hypericineis, Meliaceis et Aurantiaceis. In Indiis orientalibus occidentalibusque suavissimi saporis fructus præbent istæ plantæ.

Trib. I. *CLUSIÆ*.

Fructus multilocularis, loculis polyspermis (aff. Hypericineis.)

IP MAHUREA. *Aubl. guy. 1. p. 558. Bonnetia. Schreb. gen. p. 363.*

Calyx 5-sepalus, sepalis imbricatis. Corolla 5-petala æstivatione contortâ. Stamina ∞ libera; antheræ oblongæ. Stylus 1. Stigma simplex. Capsula conica 3-valvis, valvulis margine introflexis placentas peculiare attingentibus. Semina minutissima numerosissima compressa. — Arbores alternifoliæ; flores racemosi; pedunculi in primâ specie è 2 squamis orientes.

Genus adhuc incertioris sedis, accedit ad Bixineas habitu, ad Hypericineas fructu et seminibus, ad Guttiferas stylo staminibus et

(1) Ordinis enumerationem hanc methodicam mecum communicavit cl. et am. CHOIST. Confer ejus circa Guttiferas dissertationem mox edendam in *Mém. soc. b. nat. Par. vol. 1. part. 2.*

antheris; ad Carpodonton speciatim accedit pedunculis basi 2-squamatis, ab utroque Guttiferarum Hypericorumque ordine foliis alternis distat.

1. *M. PALUSTRIS* (Aubl. guy. l. c. t. 222.) flore in diametro vix pollicari, antheris adnatis minutis. ♂ in paludibus Guyanæ. (v. s.)
2. *M. SPECIOSA* (Chois. mss.) flore luteo in diametro 2 pollicari, antheris elongatis tetragono-sulcatis basi infixis. ♂ (Desc. ex flore unico à clar. Balbis sub nomine *Marilæ racemosæ* misso reperiit in Sanctâ-Marthâ cl. Bertero.)

II. MARILA. Swartz. prod. p. 84.

Calyx 4-sepalus; sepala cruciata, 2 exterioribus florem involventibus. Corolla 4-petala. Stamina numerosissima basi subcoacta; antheræ adnatæ. Stylus 1, brevis. Stigma capitato-adpressum. Fructus columnaris stylo persistente coronatus, 3-4-locularis, 3-4-valvis, valvulis margine introflexis placentæ centrali affixis, post dehisceniam contortuplicatis margine membranaceis. Semina numerosissima membrulâ flavâ fimbriatâ cincta.

Genus certè intermedium inter Hypericineas et Guttiferas ut optimè censuit Ill. Jussieu; affinius Hypericineis seminibus et fructu, sed ad Guttiferas inflorescentiâ Calophyllorum simillimâ, stylo et stigmate unico, antheris adnatis, multò magis accedit.

1. *M. RACEMOSA* (Swartz l. c.) ♂ in insulis Caribæis. (v. s.)

III. GODOYA. Ruiz et Pav. prod. fl. per. p. 101.

Calyx 5-phyllus coloratus. Stamina definita aut indefinita. Antheræ poris duobus pollen effundentes. Stigma 5-angulare. Capsula 5-locularis. Semina imbricata alata.

1. *G. SPATHULATA* (Ruiz et Pav. l. c. p. 102.) foliis spathulatis crenatis, staminibus circiter 40. ♂ in Chinchao et Cuchero nemoribus.
2. *G. OBOVATA* (Ruiz et Pav. l. c. p. 102.) foliis ovato-ellipticis crenatis, staminibus 10. ♂ ad Cuchero in Peruviam. (v. s.)

IV. CLUSIA. Linn. gen. n. 1154. Juss. gen. p. 256. — Quapoya Aubl. et Xantho Willd. Havetia H. B. et Kunth nov. gen. am. 5. p. 203.

Calyx 4-8-sepalus imbricatus coloratus. Corolla 4-8-petala. Stamina numerosa, rarò definita. Stylus 0. Stigma radiato-peltatum. Flores vulgò polygami et in femineis ovarium nectario stamineo brevi crasso circumdatum. Capsula carnea coriacea 5-12-valvis, costis ab apice dehiscenibus. Placentæ 3-angulares valvulis introflexis affixæ; anguli interni placentarum unam efficiunt centralem placentam angulosam. Semina teretia aut placentæ centrali aut lateralibus angulis affixa. Cotyledones separabiles. — Arbores parasiticæ, foliis oppositis, caulibus sæpè tetragonis.

Genera *Clusia*, *Quapoya*, *Havetia* habitu, existentie modo, formisque nimis affinia, internâ florum fructuumque fabricâ nimis ignota, ut unico tantum variationi obnoxio caractere, nempe staminum numero aut formâ, ea seungere audeam.

§. 1. *Staminibus numerosis*. *Clusia*.

1. *C. ROSEA* (Linn. sp. 1495.) floribus polygamis, calyce 6-phylo, corollâ 6-petalâ roseâ, nectarii densi apicibus subulatis, stigmatibus 8-12. foliis obovatis obtusis aliquando emarginatis breviter petiolatis striatis. ♂ in Carolinâ, Sancto-Domingo. Jacq. amer. 270. Catesb. car. p. et t. 99. (v. s.)

2. *C. ALBA* (Linn. sp. 1495.) floribus sæpius hermaphroditis, calyce polyphyllo, corollâ 5-8-petalâ albâ, nectarîi densi apicibus retusis amputatis aut staminibus brevibus 5-10, stigmatibus 5-6, foliis præcedentis non emarginatis. ζ in Americâ calid. Jacq. amer. 271. t. 166. Plum. am. t. 87. f. 1.
3. *C. FLAVA* (Linn. sp. 1495.) floribus polygamis, calyce polyphyllo, corollâ 4-petalâ flavâ, staminibus numerosis brevibus, stigmatibus circiter 12, foliis ut in præcedentibus. ζ in Jamaicâ. Jacq. am. 272. t. 167. *C. pedicellata*? Forst. prod. n. 390.
4. *C. VENOSA* (Linn. sp. 1495.) calyce 4-phylo, corollâ 4-petalâ albâ, staminibus numerosis, stigmatibus 5, foliis obovatis obtusis venosis. ζ in sylvis montosis humidis Martinicæ. — Plum. amer. t. 87. f. 2.
5. *C. RETUSA* (Poir. dict. 5. p. 183.) calyce 8-sepalo inæquali, corollâ 6-petalâ, stigmatibus numerosissimis, capsulâ globulosâ, foliis obovatis retusis venosis. ζ in Americâ. Lam. ill. t. 852.
6. *C. NEMOROSA* (Mey. prim. esseq. p. 203.) floribus polygamis, calyce 6-8-phylo, corollâ 4-petalâ albâ intus purpureâ, staminibus sterilibus subulatis cum aliis intermixtis, stigmatibus 5, foliis oblongo-obovatis acutis approximatis venosis. ζ in ins. Wacanama.
7. *C. PARVIFLORA* (Humb. et Bonpl. in Willd. sp. 4. p. 976.) corollis albis 5-petalis calyci subæqualibus, foliis obovatis obtusis aveniis, omnibus partibus minoribus quàm *C. albæ*. ζ in ins. Margaritâ.
8. *C. ELLIPTICA* (H. B. et K. nov. gen. am. 5. p. 199.) floribus albis, calyce 4-sepalo, stigmatibus 6, foliis sessilibus ellipticis utrinquè rotundatis, pedunculis 2-3-floris. ζ in Andibus Peruvianis prope Loxam.
9. *C. MULTIFLORA* (H. B. et Kunth nov. gen. am. 5. p. 200.) calyce 4-sepalo, corollâ 5-petalâ calycem paulò superante, stigmatibus 5, foliis sessilibus obovatis apice rotundatis, pedunculis multifloris. ζ in Andibus Quindiuensibus.
10. *C. VOLUBILIS* (H. B. et K. l. c.) stigmatibus 6, caule volubili, foliis petiolatis ellipticis utrinquè obtusis crasso-coriaceis. ζ cum præcedente.
11. *C. SESSILIS* (Forst. prod. n. 391.) floribus axillaribus solitariis subsessilibus 4-petalis, foliis obovatis ellipticisque. ζ in Tongatabu.

§. 2. *Staminibus definitis* 4. 5. 6. Quapoya.

12. *C. QUAPOYA* (Chois. mss.) floribus dioicis pedunculatis, calyce 4-6-sepalo, corollâ 4-6-petalâ luteâ, nectarîo brevi 4-6-lobo, stigmatibus 5, fructu globoso, foliis obovatis acutis. ζ in sylvis Synamari. Quapoya scandens Aubl. guian. 2. p. 898. t. 343.
13. *C. PANAPANARI* (Chois. mss.) floribus dioicis subsessilibus, fructu oblongo, foliis ovato-oblongis acutis minoribus. ζ in sylvis Guianæ. Q. Panapanari Aubl. guian. 2. p. 901. t. 344. Flores prioris.
14. *C. TETRANDRA* (Humb. et Bonpl. in Willd. sp. 4. p. 976.) paniculis axillaribus, foliis obovatis subacuminatis subtus venosis; calyce 4-sepalo, corollâ 4-petalâ, antheris? 4 disco carnoso immersis 1-locularibus papillæformibus. ζ in Andibus Popayanensium. Havetia laurifolia H. B. et Kunth nov. gen. am. 5. p. 204. t. 462.

†. *Species non satis notæ, floribus sessilibus axillaribus.*

15. *C. SESSILIFLORA* (Poir. dict. 5. p. 183.) foliis obovatis subvenosis, floribus sessilibus. ζ in Madagascari.
16. *C. ACUMINATA* (Spreng! in herb. Balb.) foliis orbiculatis basi subattenuatis mucrone spinoso acuminatis subtus venosis, floribus sessilibus. ζ in ins. Porto-Rici montibus editioribus. Bertero. (v. s. in h. Balb.)

Trib. II. *GARCINIEÆ*.

Fructus multilocularis, loculis monospermis (Aff. Aurantiaceis.)

V. *OCHROCARPOS*. *Pet. Th. gen. nov. mad. p.* 15.

Calyx 2-sepalus. Stamina numerosa in triplici ordine disposita basi coalita. Stylus nullus. Stigma planum sessile peltatum 4-6 lobum. Bacca oliuæformis 4-6-locularis. Semina arillata pseudo-monocotyledonea. — Arbor. Folia opposita sæpè approximata et quasi verticillato-ternata; flores axillares; pedunculi pauciflori.

1. *O. MADAGASCARIENSIS* (Pet. Th. 1. c.) ♀ in insulâ Madagascar. (v. s.)

VI. *MARIALVA*. *Vand. in Rœm. script. bras. p.* 118. *Tovomita. Aubl. guy. 2. p.* 956. *Beauharnoisia. Ruiz et Pav. ann. mus. 11. p.* 71.

Calyx 2-sepalus. Corolla 4-petala. Stamina 1 serialia basi coalita; antheræ minimæ ovatæ poros simulant. Stylus brevis in stigmata 4 simplicia divisus. — Arbores.

1. *M. GUYANENSIS* (Chois. mss.) foliis ovatis acutis subamplexicaulibus coriaceis 4-6-pollices longis, pedunculis corymboso-dichotomis crassis terminalibus, pedicellis articulatis. ♀ in Guyanâ et Brasiliâ. *Vand. t. VIII. f. 6. Tovomita Guyanensis Aubl. l. c. t.* 364. (v. s.)

2. *M. FRUCTIPENDULA* (Chois. mss.) foliis lanceolatis obtusiusculis 3-pollices longis, pedunculis filiformibus axillaribus terminalibusque 2-3-floris elongatis, pedicellis articulatis. ♀ in Peruvîâ et Cayennâ. *Beauharnoisia fructipendula Ruiz et Pav. l. c. t.* 9.

3. *M. UNIFLORA* (Chois. mss.) foliis præcedentis, pedunculis axillaribus unifloris vix semipollicaribus. ♀ in Guyanâ. (v. s. s.)

VII. *MICRANTHERA*. *Chois. mem. hist. nat. Par. I. t. 11 et 12.*

Calyx 2-sepalus persistens coriaceus. Petala 8-10 calyce paulò longiora. Stamina numerosissima in masc. apice minimas 2 locales antheras gerentia, in fœm. sterilia ovarium circumdantia. Bacca stigmatè 6-dentato peltato-connata 6-locularis, loculis 1-spermis. Semina centro affixa exalbuminosa. Embryo pseudo-monocotyledoneus. — Arbor dioica; folia opposita; flores racemosi.

1. *M. CLUSIÆFOLIA* (Chois. l. c.) ♀ in Cayennâ. *Clusia longifolia Richl act. soc. h. nat. par. 1. p.* 113.

VIII. *GARCINIA*. *Lin. gen. 594. Juss. gen. 256* *Garcinia et Cambogia Lin. Juss. Mangostana Gaertn. Oxycarpus Lour. Brindouia Aub. Pet. Th.*

Calyx 4-sepalus persistens. Corolla 4-petala. Stamina numerosa libera aut basi polyadelphe; antheræ ovatæ. Stylus 0. Stigma 4-8-lobum. Flores polygami aut dioici. Ovarium fœmineorum sine nectario. Bacca succulentissima 4-8-locularis. Semina arillata. Cotyledones crassæ coadunatæ. Arbores foliis ovatis aut ellipticis.

Sect. I. *MANGOSTANA*.

Flores monoici aut hermaphroditi; stamina libera.

1. *G. MANGOSTANA* (Lin. sp. 635.) foliis ovatis acutis venosis, floribus terminalibus solitariis, corollis rubris, stigmatè 6-8-lobis, baccis speciosissimis edulibus, pericarpio fungoso. ♀ in insulis Moluccis. — *Rumph. amb. 1. t.* 43. *p.* 132. *Gaertn. fr. 2. t.* 105.

2. *G. CORREA* (Lin. syst. 368.) foliis ovatis acutis venosis, floribus terminalibus solitariis cernuis, stigmate integro, baccâ pruniformi. \S in Indiâ. — Rumph. amb. 3. p. 55. t. 30.
3. *G. CAMBOGIA* (Desrous in Lam. dict. 3. p. 701.) foliis ovatis acutis, floribus terminalibus paucis, corollis luteolis, stigmate 8-lobis, baccâ 8-sulcatâ. \S in Indiâ. — Rheed. mal. 1. p. 41. t. 24. Gaertn. fr. 2. p. 106. t. 105.
4. *G. MORELLA* (Desrous. l. c.) baccâ parvâ 4-loculari striatâ, stigmate scabro 4-lobis. — Hab. Gaertn. fr. 2. p. 106. t. 105.

Sect. II. BRINDONIA. *Pet. Th. dict. sc. nat.* 5. p. 339.

Flores dioici aut hermaphroditi; stamina florum masc. basi 1-adelpha, florum herm. polyadelpha.

5. *G. COCHINCHINENSIS* (Chois. mss.) foliis ovato-oblongis subacutis, ramulis 4-gonis, floribus lateralibus congestis brevissimè pedunculatis albidis, baccâ luteo-rubrà pyriformi. \S in Chinâ et Indiâ. — Rumph. amb. 3. p. 58. t. 32. *Oxycarpus Cochinchinensis* Lour.
6. *G. ELLIPTICA* (Chois. mss.) ramulis tetragonis, foliis ellipticis ovatis venosis acutis maximis viridibus lucidis nigro-punctatis, floribus lateralibus fasciculatis minimis breviter pedunculatis, staminibus 5-adelphis, form. ignotis. \S in ins. Timor et Javâ. An satis differt à præcedente? (v. s.)
7. *G. CELEBICA* (Lin. sp. 635.) foliis ovato-lanceolatis acutis, ramulis subtetragonis, floribus terminalibus solitariis brevissimè pedunculatis, stigmate 8-lobis, baccâ globosâ. \S in ins. Celebum. — Rumph. amb. 1. p. 134. t. 44. *Brindonia Celebica* Pet. Th. dict. sc. nat. 5. p. 339.
8. *G. INDICA* (Chois. mss.) foliis ovatis acuminatis, floribus terminalibus, masculis 4-5 congestis, herm. solitariis brevius pedunculatis, baccâ globosâ 5-6-loculari. \S in Indiâ. *Brindonia Indica* Pet. Th. l. c.
9. *G. COWA* (Roxb. cat. p. 42.) foliis ovatis acuminatis, ramulis teretibus, floribus masculis lateralibus 4-5 congestis, floribus herm. solitariis terminalibus brevius pedunculatis, stigmate integro rugosâ 6-sulcato, baccâ ovato-globosâ. \S in Indiâ. *G. dioica*? Smith in Rees cycl. vol. 15. (v. s.)

Trib. III. *CALOPHYLLÆ*.

Fructus unilocularis drupaceus aut baccatus; semina pauca nunc intrâ siccum pericarpium, nunc in pulpâ nidulantia. (Aff. Meliaceis.)

IX. MAMMEA. *Lin. gen.* 1156. *Juss. gen.* p. 257.

Calyx 2-sepalus coloratus. *Pet.* 4. ovata coriacea. Stamina numerosa brevissima, antheræ minutæ oblongæ. Stylus 1 teres persistens. Stigma capitatum. Bacca carnosa, styli rudimento sæpè coronata. Semina 4 aut abortu 2-3. — Arbores; flores hermaphr. aut masculi; folia venis transversis pinnatis confertis sæpius distincta.

1. *M. AMERICANA* (Lin. sp. 731.) foliis obovatis obtusissimis integerrimis, fructibus maximis 4-spermis. \S in ins. Caribæis. Sloan. jam. 2. p. 123. t. 217. f. 3. (v. s.)
2. *M. EMARGINATA* (Moc. et Sess. fl. mex. ic. ined.) foliis obovatis obtusissimis apice emarginatis, fructibus globosis minoribus 2-spermis. — in Mexico. An sp. distincta? (v. ic.)
3. *M. HUMILIS* (Vahl eol. 2. p. 40.) foliis acutis, pedunculis petiolo longioribus, baccis 3-spermis. \S in ins. Monserrat.

X. XANTHOCHYMUS. Roxb. *corom.* 2. p. 51.

Calyx 5-partitus inæqualis imbricatus. Corolla 5-petala ovato-rotunda. Nectaria 5 petalis opposita Stamina 15-20 altissimè 5 adelpha petalis alterna; antheræ oblongæ. Stylus brevissimus. Stigma 5 lobum longum patulum. Bacca maxima pericarpio luteo coriaceo. Semina 3-4. — Arbor foliis lineari-lanceolatis acutis oppositis integris petiolatis; flores fasciculati laterales pedunculati.

1. X. TINGTORIUS (Roxb. l. c. t. 196.) § in Indiâ.

XI. STALAGMITIS. Murr. *comm. gætt.* 9. p. 175.

Calyx 4-6-sepalus caducus inæqualis. Pet. 4-6 Stamina ferè 30 receptaculo carnosio quadrangulâri inserta 5-adelpha? Stylus 1 brevissimus. Stigmata 3-4 obcordata patentia. Bacca globosa stylo et stigmatibus coronata. Semina 3-4, abortu 1. — Arbor; flores hermaphr. aut masculi cum rudimento styli.

1. S. CAMBOGIENSIS (Murr. l. c.) § in Zeylonâ et Cambogiâ.

XII. MESUA. Lin. *gen. n.* 665. Juss. *gen. p.* 253.

Calyx 4-sepalus persistens. Petala 4. Stamina numerosa basi connata. Stylus 1. Stigma crassum concavum. Nux 4 gona aut conica coriaceo-fungosa scissurâ 4-valvis, 1-4-sperma. — Frutices.

1. M. FERREA (Lin. sp. 734.) foliis elliptico-lanceolatis acutis, floribus pedunculatis, petalis subunguiculatis regularibus, nuce maturâ abortu 1-sperma. § in Javâ et Indiâ. Calophyllum Nagassarium Burm. ind. 121. Rumph. amb. 7. p. 3. t. 2. (v. s.)

2. M. SPECIOSA (Chois. mss.) foliis lineari-lanceolatis subacutis longioribus, floribus vix pedunculatis, petalis rotundatis regularibus, nuce maturâ 4-sperma. § in Indiâ. Char. ex Rheed. mal. 3. p. 63. t. 53.

XIII. CALOPHYLLUM. Lin. *gen. n.* 658. Juss. *gen. p.* 258.

Calyx 0-2-4-sepalus coloratus. Corolla 4-petala lutea. Stamina numerosa basi polyadelpha aut libera; antheræ oblongæ. Stylus crassus. Stigma simplex aut capitatum. Drupa globosa aut ovata foeta nuce abortu 1 sperma, rarè 2-sperma. — Arbores; folia nervis confertis transversis parallelis distincta; flores racemosi.

Genus pessimè notum; species vix distinctæ.

§. 1. Calyce 4-sepalo.

1. C. INOPHYLLUM (Lin. sp. 732.) foliis obovatis sæpius emarginatis, ramulis teretibus, floribus laxè racemosis, racemis axillaribus, pedunculis 1 floribus sæpius oppositis. § in Indiâ. Rumph. amb. 2. p. 211. t. 71. Rheed. mal. 4. p. 79. t. 38. (v. s.)

2. C. TAGAMANIA (Willd. mag. ber. 1811 p. 79.) foliis ovato-ellipticis acutiusculis, rarius emarginatis, cæteris præcedentis. § in insulâ Bourbonâ et Madagascar. C. inophyllum Lam. dict. 1. p. 552. — Pluk. alm. 41. t. 147. f. 3. (v. s.)

3. C. SPECTABILE (Willd. l. c. p. 80.) foliis elliptico-lanceolatis aut rarius ovato-ellipticis utrinquè sæpius acutis, cæteris præcedentium. § in Indiâ et ins. Mauritiî. C. acuminatum Lam. dict. 1. p. 553? — Rumph. amb. 2. p. 218. t. 72? C. Soulati Burm. ind. p. 121. (v. s.)

§. 2. Calyce 2-sepalo.

4. C. CALABA (Jacq. am. p. 269. t. 165.) foliis ovatis obtusis, floribus hermaphr. aut masculis, racemis lateralibus brevioribus. § in ins. Caribæis. (v. s.)

5. *C. MADBUNNO* (H. B. et Kunth nov. gen. am. 5. p. 202.) foliis oblongis utrinque acutis, pedunculis paucifloris, capsulâ maximâ sæpius 2-spermâ. \S in montosis regni Novo-Granatensis.

\S 3. *Calyce nullo?*

6. *C. SPURIUM* (Chois. mem.) foliis ovatis obtusis, pedunculis elongatis. \S in Indiâ. *C. Calaba* Linn. sp. 732. *C. apetalum* Willd. l. c. p. 79. — Rheed. mal. 4. p. 81. t. 39.

†. *Species minus notæ.*

7. *C. LONGIFOLIUM* (Willd. l. c. p. 80.) foliis oblongo-ellipticis rotundato-obtusis. \S in Amer. merid. (Humb. et Bonpl.)
 8. *C. ACUMINATUM* (Willd. l. c.) foliis oblongis acuminatis. \S in Americâ merid. (Humb. et Bonpl.)
 9. *C? CURI* (H. B. et Kunth nov. gen. am. 5. p. 203) foliis oblongis utrinque obtusis. \S in inundatis prope Sancta-Balthasar Amer. merid.

Trib. IV. *SYMPHONIEÆ.*

Fructus multilocularis, loculis mono aut polyspermis. Antheræ extrorsæ filamentis staminum in tubum aut in adelphas connatis adnatæ (aff. Meliaceis).

XIV. *CANELLA*. *Browne jam.* 215. — *Winterana* Linn. gen. 598.

Calyx 5-sepalus. Petala 5 subcoriacea glauco-cærulea contortè æstivata. Stamina in tubum coalita; antheræ 15 sulcum simulant. Stigma 3. Bacca 3-locularis abortu quandoque 1-locularis, loculis 1-2-spermis. Embryo (ex Gaertnerò forsan errante) perispermo carnoso involutus, curvulus, cotyledonibus linearibus.

1. *C. ALBA* (Murr. syst. p. 443.) \S in sylvis ins. Caribæarum et Amer. calid. *Browne jam* t. 37. f. 3. *Cat. rar.* 2. t. 50. *Winterana Canella* Linn. sp. 636. *C. alba* Sw. act. see Linn. lond. 1. p. 96. t. 8. Folia alterna obovata basi concava, nunc (verosimiliter seniora) coriacea impunctata, nunc (juniora?) pellucido-punctata. (v. s.)

XV. *MORONOBÆA*. *Aubl. guian.* 2. p. 79. — *Symphonia* Linn. f. suppl. 302.

Calyx 5-sepalus. Pet. 5 coriacea æstivatione contortâ. Stam. 3-5-adelpha, ovarium circumdantia et spiraliter sulcantia; antheræ extrorsæ lineares. Stylus 1. Stigmata 5. Bacca 5-locularis. — Arbores; folia opposita.

1. *M. COCCINEA* (Aubl. l. c? t. 313.) gemmâ globosâ, stylo brevissimo, foliis oblongis utrinque acutis coriaceis impunctatis. \S in sylvis udis et montibus Guianæ *Symphonia globulifera* Linn. f. l. c. An *S. esculenta* Arrud. ab hac diversâ? (v. s.)
 2. *M. GRANDIFLORA* (Chois. mem. cit.) gemmâ ovato-conicâ, stylo longissimo, floribus omni parte majoribus, foliis elliptico-lanceolatis. \S in Guianâ. (v. s.)

XVI. *CHRYSOPIA*. *Pet. Th. gen. mad. n.* 48.

Calyx 5-6-sepalus crassus. Petala 5 coriacea contorta. Stamina in urceolum apice 5 fidum coalita, basi nectario circumdata. Styli 5 minimi. Eructus pyriformis carnosus 5-locularis; loculis 6-spermis. Semina centro affixa. — Arbor succum flavum edens; folia coriacea opposita.

1. *C. FASCICULATA* (Pet. Th. l. c.) \S in Madagascar. Rami apice umbellatim conferti; flores apice fasciculati purpurei. (v. s.)

†. *Genera incertiora aut minùs nota.*

XVII? MACANEA. *Juss. gen. p. 257. Macahanea Aubl. guian. suppl. p. 6.*

Bacca magna pyriformis extùs gibba inæqualis cortice coriaceo 1-locularis intùs pulposa 4-6-sperma, seminibus ad receptacula lateralia affixis ovatis coriaceis in pulpâ nidulantibus. — Arbor; folia opposita dentata.

1. M. GUIANENSIS (Aubl. guian. l. c. t. 371.) ♂ in Guianâ. (v. s.)

XVIII? SINGANA. *Aubl. guian. l. c. p. 574. Juss. gen. p. 257.*

Calyx 3-5-partitus. Petala 3-5 unguiculata limbo serrulato. Stamina numerosa. Antheræ subrotundæ. Stylus 1 apice incurvus. Stigma capitatum concavum. Capsula longa cylindrica 1-locularis polysperma. Semina magna sibi mutuò incumbentia pulpâ involuta ad receptacula 3 lateralia affixa. — Arbor scandens.

1. S. GUIANENSIS (Aubl. l. c. t. 230.) ♂ in sylvis Guianæ. (v. fr.)

XIX? RHEEDIA. *Linn. gen. 641. Juss. gen. p. 258. — Van. Rheedia Plum. gen. 45.*

Calyx 0? Pet. 4. Stamina numerosa; antheræ oblongæ. Stylus 1. Stigma infundibuliforme. Bacca ovata 1-locularis. Semina 2-3 ovato-oblonga carnosa crassa in pulpâ nidulantia. — Folia opposita obtusa petiolata.

1. R. LATRIFLORA (Lind. sp. 719) ♂ in Martinicâ. *Burm. am. t. 257.*

XX? MACOUBEA. *Aubl. guian. suppl. 2. p. 17. Juss. gen. p. 257.*

Fructus aurantiiformis subcompressus, cortice tenui punctato subscabro, 1-locularis polyspermus. Semina oblonga subcurva hinc sulcata, arillata parietibus fructûs affixa. — Arbor lactescens; rami oppositi; folia opposita nervis secundariis transversis; fructus racemosi.

1. M. GUIANENSIS (Aubl. l. c. t. 378.) ♂ in Guianâ.

XXI? CHLOROMYRON. *Pers. ench. 1. p. 73. Verticillaria Ruiz et Pav. syst. fl. per. p. 140.*

Calyx 6-sepalus coloratus. Corolla 0. Stylus 0. Stigma concavum 9-lobum. Capsula 3-locularis 3-sperma. — Arbor. Folia oblonga acuminata integra. Rami tenues. Genus verosimiliter ad Garcinieas pertinet.

1. C. VERTICILLATUM (Pers. l. c.) ♂ in Peruviâ. — Ruiz et Pav. fl. per. f. 4. p. 81. t. xv.

(D. CROIST.)

Ordo XXXVI. MARCGRAVIACEÆ. *Juss. ann. mus.* 14.
p. 397. (1).

Calyx 2-7-sepalus, sepalis ovatis sæpiùs coriaceis imbricatis. Corolla hypogyna, nunc monopetala calyptræformis integra aut apice lacera, nunc 5-petala, basi post inflorescentiam circumscissa et caduca. Stamina definita aut sæpiùs numerosa, nunc receptaculo, nunc membranæ hypogynæ inserta; filamenta basi dilatata; antheræ elongatæ basi infixæ intùs dehiscentes. Ovarium unicum liberum sæpè sulcatum. Stylus 1 variæ longitudinis. Stigma simplex aut capitatum. Capsula coriacea vulgò subglobosa multivalvis vix dehiscens; valvulæ medio septiferæ; dissepimenta (in *Marcgraviâ*) tenuia arcuata basi apiceque coalita, centro distantia et fructus idèò 1-locularis. Semina minutissima numerosissima in pulpâ nidulantia. Embryo ignotus.

Frutices aliquandò adscendentes; folia alterna; flores umbellati aut spicati; pedunculi nudi aut sæpiùs bracteis, nunc simplicibus, nunc extùs concavis, nunc cuculliformibus muniti.

Ordinis hujus locus adhuc incertus remanet; ad Ebenaceas accedit corollâ monopetalâ basi circumscissâ, antheris basi infixis, foliis alternis, ad Ericæas antheris discoque generis *Antholomæ*; ad Hypericineas et Guttiferas staminibus hypogynis, corollâ quorundam generum polypetalâ, placentatione seminibusque numerosis, et idèò cel. de Jussieu prope *Clusiam* hunc collocavit ordinem.

Subordo I. MARCGRAVIÆ.

Corolla calyptræformis. Stamina receptaculo inserta.

I. ANTHOLOMA. *Labill. nov.-holl.* 2. p. 121.

Calyx 4-sepalus caducus ovatus, rariùs 2-sepalus. Corolla ovato-cylindræa margine crenata subdentata. Stamina circiter 100, disco fungoso alveolato inserta. Antheræ oblongæ apice dehiscentes. Ovarium obtusè 4-gonum. Stylus longus. Stigma acutum. Fructus immaturus 4-locularis verosimiliter capsularis. — Arbor; folia ad apicem ramulorum sparsa elliptico-oblonga coriacea petiolata; racemi florum axillares subumbellati reflexi; pedunculi nudi.

1. A. MONTANA (*Labill. voy. t.* 41.) ♂ in *Novâ-Caledoniâ*. (v. s. sp.)

II. MARCGRAVIA. *Plum. amer.* 29. *Linn. gen.* 640. *Juss. gen.* p. 244.

Calyx 6 partitus persistens ovato-rotundatus coriaceus imbricatus inæqualis. Corolla coriacea conica integra. Stamina 1-serialia mem-

(1) Auctor hujus monographiæ est suprà laudatus CHORSY.

brunulæ ovarium circumdanti inserta; antheræ oblongæ longitudinalliter dehiscentes. Stylus subnullus. Stigma crassum persistens. Capsula coriacea baccata subglobosa. — Frutices vulgò adscendentes ut Hedera, et tunc rami floriferi penduli. Folia alterna. Pedunculi bracteati.

1. *M. UMBELLATA* (Linn. sp. 503. foliis sessilibus ovato-ellipticis acutis vix venosis, sterilium ramorum ovatis obtusis, pedunculis umbellatis sæpè tuberculosi, bracteis cuculliformibus. ♂ in ins. Caribæis et Amer. cal. Jacq. am. p. 156. t. 96. Plum. gen. 7 t. 173. f. 1. Lam. ill. t. 447. (v. s.)
2. *M. CORIACEA* (Vahl ecl. 2. p. 39.) foliis ellipticis coriaceis aveniis, pedicellis verticillatis 17-18. patentissimis tuberculatis, ovario depresso globoso, stigmate 6-fido convexo-radiato. ♂ in Guianæ sylvis.
3. *M. SPICIFLORA* (Juss! ann. mus. 14. p. 402. t. 25.) foliis ovatis obtusis rariùs acuminatis, pedunculis spicatis, bracteis simplicibus, stigmate 4-lobo. ♂ in Guianâ (Rich.) (v. s. sp.)
4. *M. PICTA* (Willd. mag. berl. 1808. p. 172.) foliis sterilium ramorum ellipticis obtusis cordatis pollicaribus rotundato obtusissimis, suprâ saturatè viridibus reticulato venosis, venis albo-coloratis, subtus pallidè viridibus concoloribus (Willd.) ♂ Hab.

Subordo II. NORANTEÆ.

Corolla 5-petala. Stamina corollæ adpressa et huic quasi inserta.

III. NORANTEA. Aubl. guian. 1. p. 554. Juss. gen. p. 245. — *Ascyum* Vahl Willd.

Calyx 5-7-partitus rotundatus coriaceus. Corollæ petala reflexa. Stamina minima numerosa 1-serialia, filamentis inter se et cum corollâ subconnatis. Ovarium conicum sulcatum. Stylus nullus. Stigma capitatum aut acutum. — Frutices; folia alterna coriacea; pedunculi densi spicati bracteis cuculliformibus muniti.

1. *N. GUIANENSIS* (Aubl. l. c. t. 210.) pedunculis brevissimis, cucullis pollicaribus, antheris minutissimis vix exsertis. ♂ in Guianâ. *Ascyum violaceum* Vahl eclog. p. 41. (v. s.)
2. *N. BRASILIENSIS* (Chois. mss.) pedunculis 15 lineas longis, cucullis vix semipollicaribus, antheris exsertis longis. ♂ in Brasiliâ. Langsdorff. (v. s. in h. cl. Kunth.)

IV. RUYSCHIA. Jacq. amer. p. 75. *Souroubea* Aubl. guian. 1. p. 244.

Stamina 5 petalis alterna. Cætera præcedentis.

1. *R. SOUROUBEA* (Willd. sp. 1116.) foliis obovatis obtusis brevissimè petiolatis, floribus laxè spicatis, pedunculis elongatis, cucullis sessilibus calyci approximatis et longè 2-auriculatis. ♂ in Guianâ. *Souroubea Guianensis* Aubl. l. c. t. 97. §. Aubletii Mey. prim. esseq. p. 119. (v. s.)
2. *R. CLUSIEFOLIA* (Jacq. am. p. 75. t. 51. f. 2.) foliis ovato-ellipticis, floribus densè spicatis, pedunculis brevissimis, bracteis non cucullatis extus concavis. ♂ in Guianâ et ins. Caribæis. (v. s.)

(D. CHOISY.)




Ordo XXXVII. HIPPOCRATEACEÆ. *H. B. et Kunth nov. gen. am. 5. p. 135. — Hippocrateæ. Juss. ann. mus. 18. p. 486.*

Cal. Sepala 5 (rariùs 4-6), minima, ad medium coalita persistentia. Pet. 5 (rariùs 4-6) æqualia, hypogyna? æstivatione subimbricata. Stam. 3, rariùs 4-5, filamentis apice distinctis, basi dilatatis et usquè ad apicem ovarii in tubum crassum urceolum discumve hypogynum simulantem connexis. Antheræ (ex cl. Kunth) 1-loculæres apice transversè dehiscentes aut bi-imò quadriloculares. Ovarium intrà tubum immersum trigonum liberum. Stylus 1. Stigmata 1-3. Fructus nunc carpellis 3 samaroïdeis constans, nunc baccatus 1-3-locularis. Semina in quoque loculo 4 axi geminatim adfixa, nonnulla interdum abortientia, erecta exalbuminosa (1). Embryo rectus. Radicula infera seu ad basin tendens. Cotyledones planæ elliptico-oblongæ subcarnosæ, inter se in sicco subadherentes. — Frutices arborescentes aut scandentes sæpiùs glabri. Folia opposita, simplicia, integra aut dentata, penninervia, subcoriacea. Racemi axillares corymbosi aut fasciculati. Flores parvi indecori.

Ordo non satis notus, olim à cl. Jussæo Acerineis adjunctus, sed staminum numero ternario, filamentis in urceolum connexis et fructûs ipsius structurâ diversus. An imò Acerineis Malpighiaceisve, suadente Jussæo, verè affinis? aut habitu affinior, ex cl. Brownio, Celastrineis, mediantibus et Elæodendro in quo albumen vix visibile et Ptelidio in quo albumen ad membranam tenuissimam reductum est. — Intereà genera præsertim baccata ulteriork egent examine.

I. HIPPOCRATEA. *Linn. gen. n. 54. — Coa. Plum. gen. p. 8. t. 35.*

Petala sæpiùs apice foveata. Stam. 3; antheræ 1-loculares transversè apice dehiscentes. Carpella 3 (aut abortu 1-2 samaroïdea, bivalvia, valvis carinato-compressissimis. Semina deorsum alata, funiculo nempè umbilicali latissimè expanso! .

1. *H. OBCORDATA* (Lam. ill. 1. p. 100. t. 28. f. 1.) carpellis obcordatis, racemis corymbosis foliis brevioribus, foliis ovato-lanceolatis serratis.  in ins. Caribæis et Novâ-Hispaniâ. *H. scandens* Jacq. am. 9. t. 9. Petala ovata obtusa.
2. *H. UNIFLORA* (fl. mex. ic. ined.) carpellis obcordatis, pedunculis unifloris, foliis oblongis basi cordatis integris.  in Mexico. Petala ovata obtusa.
3. *H. OVATA* (Lam. ill. 1. p. 100. t. 28. f. 2.) carpellis ovatis, paniculis axillaribus subdichotomis, foliis oblongo-ovatis ellipticisve serratis.  in Sancto-

(1) Hippocrateæ ovatæ spermodermium et cotyledones intus singulari modo filis innumeris tracheiformibus stuposi! quod etiam in Calypsois suæ pericarpio vidit cl. Petit-Thouars.

- Domingo. Vahl enum. 2. p. 27. *H. volubilis* Linn. syst. 84. Swartz! obs. 28.
H. paniculata Bert! in Schult. mant. 1. p. 396. — Plam. ed. Burm. 76. t. 88?
 sed carpella magis rotundiora et folia angustiora. (v. s.)
 ♀ *oblongifolia*, foliis magis oblongis. 5 in Porto-Ricco. (v. s. sine fr.)
4. *H. LAEVIGATA* (Rich. in Vahl enum. 2. p. 27.) carpellis . . . , paniculis axillaribus subdichotomis, pedicellis 2 unifloris in divisiis, foliis ovato-oblongis obsolete crenatis. 5 in Cayennâ. (v. s. sine fr.)
5. *H. DISCOLOR* (Meyer pr. esseq. 19.) carpellis ovatis, paniculis axillaribus subdichotomis rufo-velutinis, ramis minoribus multifloris, foliis ovalibus breviter acuminatis obsolete crenato-serratis. 5 in sylvis littoreis Guianæ. An à priori satis diversa? Folia utriusque glabra subtus ferruginea. (v. s.)
6. *H. VIRIDIS* (Ruiz et Pav. fl. per. 1. p. 44. t. 74. f. a.) carpellis ovalibus emarginatis, corymbis dichotomis folio brevioribus, foliis ovatis obtusè acuminatis integris serratisque. 5 in Andium nemoribus. Petala obtusa virescentia.
7. *H. ACUTIFLORA* (fl. mex. ic. ined.) carpellis obovatis. corymbis dichotomis folio brevioribus, petalis acutis, foliis oblongo-ovalibus dentatis. 5 in Mexico. Petala virescentia acuta, apice quasi bifoveolata. Antheræ 4-lobæ. Stigma simplex.
8. *H. CELASTROIDES* (H. B. Kunth nov. gen. am. 5. p. 136) carpellis subrotundo-obovatis, corymbis dichotomis glabris folio dimidiò brevioribus, petalis rotundato-obtusis, foliis lanceolato oblongis acuto-cuspidatis. 5 in Mexico prope Venta del Estola. Flores virides.
9. *H. ACAPULCENSIS* (H. B. Kunth l. c.) carpellis obovatis, pedunculis dichotomis multifloris glabris folia subæquantibus, petalis ovatis, foliis obovato-oblongis brevissimè cuspidatis, ramis verrucosis. 5 prope Acapulco. Flores virides.
10. *H. VERRUCOSA* (H. B. Kunth l. c.) carpellis obovatis, pedunculis ramosissimodichotomis multifloris glabris folio subæqualibus, petalis ovatis, foliis elliptico-oblongis acutiusculis remotè serrulatis, ramis verrucosis. 5 in regno Novo-Granatensi. Flores virides.
11. *H. MALPIGHIEFOLIA* (Rudg. pl. guian. 10. t. 8.) carpellis . . . , paniculis axillaribus terminalibusque, foliis angustè ovalibus integerrimis acuminatis subtus ramisque asperis. 5 in Guianâ. Antheræ 4-lobæ.
12. *H. COMOSA* (Swartz! fl. ind. occ. 1. p. 77.) carpellis oblongis obovatisve, paniculæ pedunculis multifidis capillaribus, foliis ovatis acuminatis integerrimis basi obtusis aut subcordatis. 5 in sylvis interioribus Hispaniolæ. Carpella 2-3-poll. longa. Flores albi.
13. *H. EMARGINATA* (Rudg. pl. guian. p. 11. t. 9.) carpellis . . . , paniculis axillaribus brevibus dichotomis, foliis obovatis integerrimis emarginatis, caule pustulis minutis aspero. 5 in Guianâ. Antheræ subrotundæ.
14. *H. PANICULATA* (Vahl enum. 2. p. 28.) carpellis . . . , paniculis axillaribus dichotomis folio brevioribus, foliis oblongis utrinquè acutis obtusè serratis. 5 in Sierrâ-Leonâ. Rami patentissimi apice compressi. (v. s.)
15. *H. MACROPHYLLA* (Vahl enum. 2. p. 28.) carpellis . . . , paniculis axillaribus dichotomis folio brevioribus, foliis ovatis integerrimis nitidis basi obtusis apice acuminatis. 5 in Sierrâ-Leonâ. Folia ut in priore pallidè viridia. (v. s.)
16. *H. VELUTINA* (Afz. ex Spreng. neu. entd. 3. p. 234) carpellis . . . , racemis axillaribus, foliis oblongis acuminatis integerrimis subtus venosis rufo-villosiusculis. 5 in Guinæâ. Fructus ex Afzelio tergeminus.
17. *H. INDICA* (Willd. sp. 1. p. 193.) carpellis oblongis 2-spermis, paniculis corymbosis dichotomis folii fere longitudine, foliis ovalibus acutis serratis nitidis. 5 in sylvis montosis Indiæ. — Roxb. cor. 2. t. 130.
 ♀ *disperma* (Vahl enum. 2. p. 28.) carpellis lanceolatis utrinquè obtusis,

- paniculis folio brevioribus, foliis ellipticis acuminatis serrulatis. 5. *evonymoides* (Vahl l. c.) foliis oblongis obovatisve apice integris emarginatisve.
18. *H. OBTUSIFOLIA* (Roxb. fl. ind. 1. p. 170.) carpellis obovatis 4-spermis, corymbis terminalibus axillaribusque, foliis elliptico-oblongis integris obtusis, ramulis cirrhosis. 5 in Coromandeliâ. Antheræ 2-lobæ.
19. *H. ARBorea* (Roxb. cor. 3. t. 205.) carpellis lineari-oblongis subnatis 2-spermis, corymbis axillaribus dichotomis, foliis ellipticis serrulatis mucrone cuspidatis, ramis subadscendentibus. 5 in Indiæ prov. interioribus. Flores 4-6-fidi. Antheræ 4-lobæ.
20. *H. CASSINOIDES*, carpellis. . . . , corymbis axillaribus dichotomis folio triplò brevioribus, foliis ellipticis basi et apice acuminatis subserrulatis submembranaceis, ramulis apice compressis. 5 in insulâ Timor. (v. s.)
21. *H. PAUCIFLOA*, carpellis. . . . , corymbis axillaribus dichotomis 5-7-floris vix petiolo longioribus, foliis elliptico-oblongis obtusissimè subcrenulatis. 5 in insulâ Timor. Flores 4 lin. diam. Antheræ sub-4-lobæ. (v. s.)
22. *H. ELLIPTICA* (H. B. et Kunth nov. gen. am. 5. p. 138.) carpellis. . . . , pedunculis tenuissimè tomentosis dichotomo-quadrifidis folio brevioribus, foliis ellipticis acutis remotè crenato-serrulatis, ramulis quadrangularibus. 5 in regno Mexicano. Flores albi.
23. *H. EXCELSA* (H. B. et Kunth l. c.) carpellis. . . . , pedunculis bifidis multifloris pulverulento hirtellis folio triplò quadruplòve brevioribus, foliis elliptico-oblongis acuminatis undulato-crenatis, ramulis junioribus 4-angulis, caule arboreo. 5 in Mexico prope Masatlan. Flores albo-virides.

II. *ANTHODON*. Ruiz et Pav. fl. per. 1. p. 45. — *Anthodus* Mart. in Schult. mant. 1. p. 253.

Cal. 5-lobus, lobis rotundatis. Pet. 5 inæquilatera dentata. Stam 3 inter discum et ovarium inserta, filam. infernè dilatatis, antheris 1-locul. apice transversè dehiscentibus. Ovarium 3-loculare; ovula circiter 8 in quoque loculo axi affixa. Stylus brevissimus. Bacca subglobosa 2-3-locul., loculis abortu 1-spermis. Semina ovata mucilagine involuta. — Char. floris ex Kunth nov. gen. am. 5. p. 140, fructus ex Mart. l. c. — An genus satis à Salaciâ differt?

1. *A. DECUSSATUM* (Ruiz et Pav. fl. per. 1. t. 74. f. 2.) foliis oblongo-ovatis subacuminatis obtusè serratis nitidis, paniculis axillaribus dichotomis, petalis ovato oblongis argutè serratis. 5 in Andium nemoribus ex R. et P., prope Angustura ex H. B. et Kunth nov. gen. am. 5. p. 140. t. 443. *Hippocratea*? *Anthodon* Pers. ench. 1. p. 40. *Tonsella decussata* Vahl enum. 1. p. 30.
2. *A. PANICULATUM* (Mart. l. c.) foliis ovatis obtusiusculis obsolete et subundulato-crenatis, floribus paniculatis, petalis ovalibus remotè et inæqualiter dentatis. 5 ad Rio-Janeiro. Cæt. ut in omnibus sequentibus ignota.
3. *A. UNDULATUM* (Mart. l. c.) foliis ovatis acutis utrinquè attenuatis obtusè subundulato serratis, floribus 2-3 axillaribus breviter pedicellatis, petalis obovato-subrotundis fimbriato-dentatis. 5 ad Rio-Janeiro.
4. *A. ELLIPTICUM* (Mart. l. c.) foliis crassis ellipticis utrinquè rotundatis margine incrassato integerrimis, floribus axillaribus lateralibusque breviter pedunculatis fasciculatis, petalis obovato-orbiculatis serrulatis. 5 ad Rio-Janeiro.
5. *A. GRANDIFOLIUM* (Mart. l. c.) foliis crassis oblongis obtusis integerrimis, floribus axillaribus lateralibusque aggregatis brevissimè pedunculatis, petalis ovato-suborbiculatis denticulatis. 5 ad Rio-Janeiro.
6. *A. OBLONGIFOLIUM* (Mart. l. c.) foliis crassis oblongis obtusiusculis, antiè remotè et subundulato-crenatis, floribus axillaribus lateralibusque aggregatis,

petalis obovato-oblongis denticulatis. ζ in sylvis deserti ad fluv. Sancti-Francisci prov. Bahiæ.

7. *A. GLOMERATUM* (Mart. l. c.) foliis crassis oblongo-lanceolatis obtusiusculis, floribus glomeratis axillaribus lateralibusque sessilibus, petalis orbiculatis vix denticulatis. ζ in sylvis deserti ad fluv. Sancti-Francisci prov. Bahiæ.
8. *A. MICRANTHUM* (Mart. l. c.) foliis crassis lineari-oblongis obtusis integerrimis, paniculis axillaribus terminalibusque paucifloris, petalis obovatis vix denticulatis. ζ in sylvis Catingas dictis prov. Bahiensis.
9. *A. CRASSIFOLIUM* (Mart. l. c.) foliis crassis submarginatis lineari-ellipticis, floribus glomeratis sessilibus axillaribus lateralibusque, petalis obovato-orbicularibus integerrimis. ζ in altis campis desertis versus Vaô la Paraxa prov. Goyazanæ.

III. RADDISIA. *Leand. Sacram. in denkschr. münch. akad. 7. p. 244. t. 15. ex Schult. mant. 1. p. 252.*

Cal. 5-sepalus. Corolla rotata 5 fida. Annulus extrâ stamina ovarium cingens. Stam. 3, filam. linearibus, antheris bilocularibus? Pistillum staminibus longius. Stylus brevis. Capsula 3-locul., loculis polyspermis. Semina subglobosa axi centrali adfixa. — An ab Anthodo satis differt? an potius Raddia dicenda?

1. *R. ARBOREA* (Leand. l. c. Schrank l. c. p. 244. ex Schult. mant. 1. p. 347.) ζ ad ripas insularum Rio-Janeiro. Folia elliptica acuta glabra serrata breviter petiolata. Flores axillares solitarii aggregative.

IV. SALACIA. *Lin. mant. 293. — Tontelea Aubl. guian. 1. p. 31. — Tonsella Schreb. gen. n. 74. — Calypso Pet. Th. veg. afr. 1. p. 29. t. 6. — Sicelium Brown ex Poir. suppl. 5. p. 146?*

Cal. 5-partitus. Pet. 5 ungue lato calyci? inserta. Urceolus carnosus inter petala et pistillum. Stam. 3 disco intus inserta, filam. basilatis conniventibus; antheris apice adnatis bilocularibus, loculis divergentibus. Ovarium 3-locul., loculis polyspermis. Stylus crassus brevissimus. Bacca rotundata, polysperma. Semina ovata coriacea. — Char. ex Pet. Th. l. c. An omnibus speciebus ritè conveniens? An genera hic plura confusa? descriptiones non sufficiunt!

1. *S. SCABRA*, paniculis axillaribus et terminalibus dichotomis, foliis ovato-oblongis subintegerrimis subtus venosis et asperis. ζ in Cayennâ et insulâ Trinitatis. Tontelea scandens Aubl. t. 10. Hippocratea aspera Lam. ill. 1. p. 101. Tonsella scabra Vahl enum. 29.
2. *S. LÆVIGATA*, paniculis axillaribus multifidis, petalis ciliatis, foliis petiolatis oblongis acutis subundulatis integerrimis glabris subtus subasperis, ramis glabris. ζ Patr. ign. Tonsella lævigata Hoffmans. in Link jahrb. 3. p. 68. ex Schult. mant. 1. p. 347. Priori affinis.
3. *S. MULTIFLORA*, pedunculis congestis subtrifloris, foliis obovatis integerrimis nitidis ζ in Cayennâ. Hippocratea obovata Rich. in act. soc. h. nat. par. 1. p. 106. H. multiflora Lam. ill. 1. p. 101.
4. *S. AFRICANA*, pedunculis congestis 1-floris, foliis ovalibus nitidis extrorsum denticulatis, dentibus subglandulosis. ζ in Guineâ. Tonsella Africana Willd. sp. 1. p. 194.
5. *S. SENEGALENSIS*, pedunculis congestis è tuberculo axillari ortis 1-floris, foliis oblongis acuminatis lævibus nitidis extrorsum serratis, ramis scabris. ζ in Senegaliâ. Hipp. Senegalensis Lam. ill. p. 101. H. verticillata a. Pers. ench. 1. p. 40. Tonsella Senegalensis Vahl enum. 2. p. 31. (v. s.)
6. *S. MADAGASCARIENSIS*, pedunculis congestis è tuberculo axillari ortis 1-floris, foliis lanceolato-oblongis subintegerrimis acuminatis lævibus nitidis, petalis

linearibus. ζ in Madagascar. Hipp. Madagascariensis Lam. ill. t. p. 101. H. verticillata β . Pers. l. c. Tonsella Madagascariensis Vahl l. c.

7. δ ? CALYPSO, pedunculis subternis ϵ tuberculo axillari ortis 1-floris, foliis ovali-oblongis obtusè subdentatis venosis. petalis ovatis. ζ in Madagascar. Calypso Pet. Th. veg. afr. 1. p. 29. t. 6. Bacca globosa mucronata. Semina aluminosa ex Pet. Th.
8. S. PRINOIDES, pedunculis axillaribus confertis elongatis 1-floris?, foliis ellipticis obtusis apice obsolete serratis. ζ in Indiâ orient. Tonsella prinoïdes Villoid. act. nat. cur. ber. 4. p. 184. Bacca 1-sperma
9. S. COCHINCHINENSIS (Lour. coch. 2. p. 642.) pedunculis axillaribus congestis 1-floris, foliis ovatis subacuminatis subserratis, petalis subrotundis. ζ in dumetis Cochinchinæ.
- 10? S. CHINENSIS (Lin. mant. 293.) pedunculis axillaribus congestis 1-floris, foliis alternis? ovalibus integerrimis lævibus, ramis angulatis. ζ in Chinâ. idem ac præcedens ex adnotat in herb. Banksiano.

V. JOHNIA. Roxb. flor. ind. 1. p. 172.

Antheræ 3 ad apicem urceoli sessiles. Fructus haccatus, junior 3-locul. ovulis 1-2 in loco quoque peltatis, adultus oligospermus. — Arbusculæ pedunculis axillaribus 1 floris.

1. J. SALACIOIDES (Roxb. l. c.) foliis integris. calyce 5-partito. petalis sessilibus. ζ in Bengala orientali. Flores aurantii parvi. Bacca edulis 2-3-sperma.
2. J. COROMANDELIANA (Roxb! l. c.) foliis serrulatis, calyce 5-dentato, petalis unguiculatis. ζ in sylvis montosis Coromandelie. Bacca 1-sperma Cerasi parvi facie. (v. s.)

†. Hippocrateaceæ spurie; staminibus 5 et ultrâ.

VI. TRIGONIA. Aubl. guian. 1. p. 390.

Cal. 5-partitus inæqualis persistens. Pet. 5 basi lata, inæqualia, superiore fornicato, 2 lateralibus alæformibus, 2 inf. in carinæ modum connexis. Stam. 5-10, nonnulla sterilia, basi irregulariter connexa. Ovarium 3-gonum 3-locul., loculis ∞ -ovulatis. Stylus 1. Capsula 3-gona, 3-valvis, 3-locul., valvis intus lanuginosis. Semina ∞ lanuginosa axi adfixa. — Frutices sarmentosi, foliis oppositis integris bistipulatis, floribus paniculato-racemosis.

1. T. SERICEA (H. B. et Kunth nov. gen. am. 5. p. 141.) foliis obovatis subtus tomentoso-sericeis argenteis. ζ in Andibus Quindiuensium.
2. T. VILLOSA (Aubl. l. c. t. 149 Lam. ill. t. 347.) foliis subtus albis velutino-tomentosis, nervis nervulisque reticulatis. ζ in Cayennâ. Tres stirpes huc verosimiliter referendas reperit indefessus Perrotet.
 - a. obtusata, foliis elliptico-obovatis utrinquè obtusis. (v. s.)
 - β . cuneata, foliis obovatis basi cuneatis. (v. s.)
 - γ . oblonga, foliis oblongis utrinquè acuminatis. (v. s.)
3. T. MOLLIS, foliis ellipticis basi attenuatis apice acuminatis supernè pubescenti-villosis subtus vix albidis molliter villosis, nervulis non reticulatis. ζ in Brasiliâ. Martius. (v. s.)
4. T. LÆVIS (Aubl. l. c. t. 150. Vahl ecl. 2.) foliis utrinquè glabris nitidis. ζ in Guianâ. Folia oblonga acuminata. (v. s.)

VII. LACEPEDEA. H. B. et Kunth nov. gen. am. 5. p. 142. t. 444. — Tricera β Willd. in Ræm. et Schult. syst. 4. p. 863.

Cal. 5-partitus inæqualis. Pet. 5 brevissimè unguiculata. Stam.

5 filamentis liberis, antheris bilocularibus longitudinaliter dehiscen-
tibus. Ovarium 3-loculare, loculis 8-ovulatis. Stylus 3-sulcatus, demum
tripartibilis. Bacca stylis tricuspidata 6-9-sperma. — Arbor foliis
serrulatis, paniculis terminalibus, floribus albis odoris. — Trigonæ
affinis.

1. *L. insignis* (H. B. et Kunth l. c.) 5 prope Xalapa Mexicanorum. *Triceraja*
tinifolia Willd. in Schult. l. c.

Ordo XXXVIII. ERYTHROXYLEÆ. *H. B. et Kunth nov. gen.*
am. 5. p. 175.

Sepala 5 basi concreta persistentia. Petala 5 hypogyna, basi lata, intus aucta per squamam è plicis formatam, sepalis alterna, æqualia, antè florescentiam marginibus incumbentia. Stam. 10, filam. basi in urceolum concretis, antheris basi adfixis erectis bilocularibus longitudinaliter latere dehiscentibus. Ovarium 1-loculare ovulo solitario pendulo, aut 3-loculare loculis lateralibus vacuis. Styli 3 distincti, stigmatibus subcapitatis, aut ferè ad apicem in 1 concreti. Drupa 1-sperma. Semen angulatum. Albumen corneum. Embryo linearis rectus centralis. Cotylédones lineares planæ foliaceæ. Radicula supera teres recta. Plumula inconspicua.

Frutices aut arbores. Ramuli juniores stipulis acutis sæpè imbricatis obtekti et compressi. Stipulæ axillares. Folia alterna rariùs opposita, in plerisque, forsàn in omnibus speciebus, glabra. Flores parvi albidi aut flavo-virescentes. Pedunculi basi bracteati.

Ordo à Malpighiaceis segregatus à cl. Kunthio ob petala appendiculata, semen albuminosum, fructum sæpiùs abortu 1-locularem, et habitum proprium.

I. ERYTHROXYLUM. *Lin. gen. n. 575. Cav. diss. 8. p. 399. Lam. ill. t. 383.*
H. B. et Kunth l. c.

Calyx 5-partitus imà basi 5-angularis. Styli 3 à basi discreti nec in unicum coaliti.

§. I. PENNINERVIA, *foliis nempe penninerviis, nervulis apice tantùm conniventibus.*

* *Pedicellis solitariis.*

1. E. HYPERICIFOLIUM (Lam. dict. 2. p. 394.) foliis obovatis sæpè emarginatis, pedicellis flore triplò longioribus, fructu triloculari. 5 in insulis Mauritanis ubi vulgò dicitur *Bois d'huile*, *Bois de Dames*. Venelia Comm. herb. — Cav. diss. 8. p. 400. t. 230. Folia Spireæ hypericifoliæ submembranacea, subtùs pallida. (v. s.)
2. E. BREVIPES, foliis obovatis sæpè emarginatis, pedicellis lateralibus vix flore longioribus, fructu 1-loculari. 5 in rupestribus maritimis ins. Porto-Ricci et Sancti-Domingi. Bertero. Priori valdè affine. Folia coriacea, 5-6-lin. longa. Pedicelli folio quadruplò breviores. (v. s.)
3. E. BUXIFOLIUM (Lam. dict. 2. p. 394.) foliis lanceolato-obovatis obtusis submucronatis, nervo medio elevato, pedicellis flore duplò longioribus, fructu 1-loculari. 5 in Madagascar, et forsàn ins. Mauritanis. — Cav. diss. 8. p. 403. t. 231. f. 1. (v. s.)

emarginatis coriaceis glabris reticulato-nervosis. 5 in Sancto-Domingo. Fl. ignoti. Pedicelli 4-5 adgregati. Folia penninervia, 4-5 lineas longa et lata.

N. B. Præter species suprâ indicatas, plurimæ adhuc in herbariis exstant indescriptæ et ob flores fructusque deficientes non tutò describendæ.

II. SETHIA. *H. B. et Kunth nov. gen. am. 5. p. 175. in adn.*

Calyx 5-lobus. Styli 3 in unicum concreti, stigmatibus tantum adhuc inter se distinctis.

1. *S. INDICA.* 5 in montibus Indiæ Circarsensibus. *Erythroxyton monogynum* Roxb. cor. 1. t. 88. Folia obtusa penninervia obovato-lanceolata. Pedicelli vix flore longiores. Petala flava ungue albo.
-

Ordo XXXIX. MALPIGHIACEÆ. *Juss. gen.* 252. *ann. mus.* 18. p. 479.

Cal. 5-partitus, sæpius persistens. Petala 5 lobis calycinis alterna, disco hypogyno inserta, unguiculata, interdum subinæqualia, rarissimè nulla. Stam. 10 petalis alterna, ibidem inserta, rariùs pauciora, interdum solitaria. Filamenta nunc libera, sæpius basi breviter coalita. Antheræ subrotundæ. Ovarium 1 sæpius trilobum, carpellis 3 plus minùs junctis constans. Styli 3 distincti aut in 1 coaliti. Fructus 3-carpellatus 3-locularisve, interdum abortu 1-2-locularis, siccus baccatusve, loculis 1-spermis. Semina in loculis pendula (an semper?), exalbuminosa. Embryo plus minùs curvatus, aut rectus, radiculâ brevi, lobis foliaceis crassiusculisve.

Frutices aut arbusculæ. Rami interdum scandentes. Folia opposita, aut in Banisterieis paucissimis alterna, simplicia, impunctata, sæpius utrinquè stipulata. Flores racemosi aut corymbosi, pedicellis sæpè medio articulatis et bisquamulosis.

An forsàn ad hunc ordinem referenda *Passiflora pallida* Lour. fl. coch. 2. p. 544. excl. syn.?

Trib. I. *MALPIGHIEÆ*. — *Malpighia* Lin. *gen. n.* 572.

Styli 3 distincti aut in 1 coaliti. Fructus carnosus indehiscens. — Folia opposita.

I. *MALPIGHIA*. *Rich. in Juss. ann. mus.* 18. p. 480.

Calyx 5 partitus, basi extùs 8-10 glandulis instructus. Pet. unguiculata. Stam. filamenta basi breviter monadelphæ. Styli 3 distincti. Drupa nucleos 3 monospermos fovens. — Pedunculi axillares, aut 1-flori, aut pedicellos umbellatos gerentes.

§. 1. *Urentes, setis nempe foliorum centro adfixis, horizontalibus, rigidis, utrinquè pungentibus et urentibus.*

1. *M. FUCATA* (Ker. bot. reg. t. 189.) ramis glabris, foliis ellipticis nitidis subtùs decumbenti-hispidis supernè nudiusculis, pedicellis axillaribus umbellatim corymbosis folio quadruplò brevioribus, petalo summo submajore margine lacinialato. \S in Indiâ occid? *M. macrophylla* Desf. cat. 160. non *Juss.* (v. s.)
2. *M. URENS* (Linn. sp. 609.) ramis glabris, foliis oblongo-ovatis subtùs decumbenti-hispidis supernè glabris, pedunculis 1-floris corymbiferisve folio duplò brevioribus, petalis æqualibus. \S in Americâ calid. *Mill. dict.* t. 181. f. 1. *Cav. diss.* t. 235. f. 1. *Lam. ill.* t. 381. f. 1. (v. v)
3. *M. CRUDA* (Spreng! neu. entd. 3. p. 51.) ramis hirsutis, foliis elliptico-oblongis acutis supernè hirsutis, margine et subtùs decumbenti-hispidis, pedunculis 1-floris adgregatis folio duplò brevioribus. \S in Sancto-Domingo. *Bertero.* (v. s.)
4. *M. SETOSA* (Bert! ined. Spreng! neu. entd. 3. p. 50.) ramis glabris, foliis elliptico-oblongis obtusis subtùs decumbenti-hispidis supernè glabris, pedi-

- cellis axillaribus 1-floris folii ferè longitudine suprà medium bracteolatis. ζ in Sancto-Domingo. M. Martinicensis Jacq. amer. 136. (v. s.)
5. M. CUBENSIS (H. B. et Kunth. nov. gen. am. 5. p. 145.) ramulis pilosiusculis, foliis oblongo-lanceolatis submucronatis basi rotundatis integerrimis, suprà nitidis glabris, subtùs adpresso-setosis. ζ in ins. Cuba prope Havanam. fl. et fr. ign. — Affinis M. angustifoliæ.
6. M. ANGUSTIFOLIA (Linn. sp. 610.) ramis glabris, foliis lineari-lanceolatis acutis, utrinqùe decumbenti-hispidis, pedunculis axillaribus umbellatis. ζ in Americà merid. Cav. diss. 8. p. 408. t. 236. f. 1. M. linearis Jacq. am. 135.
7. M. AQUIFOLIA (Linn. sp. 611.) ramis glabris, foliis lanceolatis dentato-spinosis subtùs decumbenti-hispidis. ζ in Americà calid. — Plum. ed. Burm. t. 168. f. 1. — Cav. diss. 8. p. 409. t. 236. f. 2. (v. s.)

§. 2. *Apyræ, nempe aut glabræ aut pilis nec urentibus nec centro adfixis donatæ.*

8. M. COCCIFERA (Linn. ed. Reich. 2. p. 371.) foliis subovatis dentato-spinosis glabris. ζ in Americà calid. Cav. diss. t. 235. f. 2. Jacq. ic. rar. 3. t. 470. Lois. herb. amat. t. 215. M. Coccigrya Linn. sp. 611. Junior habet folia elliptica integra (Lam. ill. t. 381. f. 2.) et tunc M. microphylla interdum dicta. (v. v.)
9. M. GLABRA (Linn. sp. 609.) foliis ovatis integerrimis glabris, pedunculis axillaribus umbellatis. ζ in Americà calid. — Mill. ic. t. 181. f. 2. Cav. diss. t. 234. f. 1. (v. s.)
10. M. BIFLORA (Poir. dict. 4. p. 326.) foliis ovato-lanceolatis acutis glabris, pedunculis axillaribus bifloris. ζ in sylvis Americæ calid. M. puniceifolia Cav. diss. 8. p. 406. t. 234. f. 2.
11. M. PUNICEIFOLIA (Linn. sp. 609.) foliis ovatis integerrimis glabris, pedunculis axillaribus 1-floris. ζ in Americà merid. — Plum. ed. Burm. t. 166. f. 2. (v. s.)
12. M? FAGINEA (Swartz fl. ind. occ. 2. p. 850.) foliis oblongo-ovatis integerrimis, subtùs sericeo-nitentibus, pedunculis axillaribus tripartito-umbellatis. ζ in ins. Granatâ Americæ. Flores lutei. Fruct. ign. An Banisteriæ species?

†. *Species non satis notæ.*

13. M. EMARGINATA (fl. mex. ic. ined.) foliis ovatis obtusis undulatis apice emarginato-bifidis glabris, pedunculis axillaribus in pedicellos 4-5 umbellatos à basi divis. ζ in Mexico.
14. M. NITIDA (Mill. dict. n. 5.) foliis ovatis acutis integris glabris, pedunculis umbellatis axillaribus terminalibusque. ζ ad Carthagenam Amer. An à M. glabrâ satis differt? An eadem ac M. nitida Linn. sp. 609?
15. M. INCANA (Mill. dict. n. 3.) foliis lanceolatis subtùs incanis, pedunculis axillaribus umbellatis. ζ in agro Campechiano. M. Campechiensis Poir. dict. 4. p. 333. Affinis videtur M. fagineæ sed flores rosei.
16. M? GRANDIFOLIA (Jacq. amer. p. 137.) foliis lanceolato-oblongis integris, pedunculis axillaribus racemoso-corymbosis. ζ in Martinicæ sylvis. Folia pedalia. Flores lutei. An potiùs Bunchosiæ species?
17. M? DUBIA (Cav. diss. 8. p. 413. t. 242.) foliis ovato-acutis cartilagineis integris glabris, ramis punctatis, racemis axillaribus paniculatis. ζ in ins. Sancti-Dominici. Flores lutei minimi. Fruct. ignot. An Banisteria? An Bunchosia? (v. s. sine fruct. in herb. Balb. ex Jamaicâ, at stylis brevibus nec exsertis donata et ideò forsân distincta?)
18. M? OBOVATA (H. B. et Kunth nov. sp. gen. 5. p. 146.) foliis obovatis acutis cordatis integerrimis coriaceis suprà nitidis subtùs puberulis, umbellis axillaribus geminis 4-floris. ζ in ripâ fl. Magdalenzæ prope Nares. — Fr. ign.

Squamulæ 5 hypogyæ inter stamina et ovarium. Flores flavi. An genus proprium?

19. *M. TERNIFOLIA* (H. B. et Kunth l. c.) foliis ternatim verticillatis oblongo-lanceolatis acutis basi rotundatis integerrimis, suprà pilosiusculis nitidis, subtùs molliter hirtò-tomentosis, umbellis axillaribus compositis 3-4-radiatis multifloris. 5 propè Pandi Novo-Granatensium. Fr. ign. An Banisteriæ sp? Flores flavi.

- II. *BYRSONIMA*. *Rich. in Juss. ann. mus.* 18. p. 481. *H. B. et Kunth nov. gen. am.* 5. p. 147.

Cal. 5-partitus, basi extùs glandulis 8-10 grossis instructus. Pet. unguiculata. Stam. 10 basi breviter monodelpha. Styli 3. Drupa nucleo 3-loculari 3-spermo. — Racemi spicati terminales, simplices aut ramosi,

* *Foliis subtùs velutinis tomentosive.*

1. *B. VERBASCIFOLIA*, foliis lanceolato-obovatis integerrimis utrinquè tomentosis, racemis terminalibus, trunco crasso nodoso humili. 5 in campis littoreis Guianæ. *Malpighia verbascifolia* Lin. sp. 610. *Aubl. guian.* 1. p. 460. t. 184. *Cav. diss.* 8. p. 411. t. 240. Inter pubem paginæ fol. super. adsunt pili nonnulli centro adfixi. Folia cinerea. (v. s. sine fl.)
2. *B. LAURIFOLIA* (H. B. et Kunth l. c.) foliis ovato-oblongis apice angustatis basi cuneatis subcomplicatis suprà puberulis, subtùs tomentoso-pubescentibus et incanis, ramulis tomentoso-hirsutis, calycibus sericeo-villosis. 5 in Llanos de Cumana. Affinis *B. verbacifoliæ*.
3. *B. ENIOPODA*, foliis ovalibus obtusis attenuatis coriaceis, supernè demùm lucidis, subtùs et supernè in nervis hirsutis, pedunculis hirsutissimis. 5 in Americà calid. *Malpighia rufa* Poir. dict. 4. p. 332? (v. s.)
4. *B. NERVOSA*, foliis ovalibus utrinquè obtusis supernè lucidis rugosis subtùs rufo tomentosis penninerviis, nervis nervulisque prominulis, racemo confertifloro rufo-tomentoso. 5 in Brasiliâ. *Malpighia macrophylla* Juss. in Pers. ench. 1. p. 506? (v. s.)
5. *B. COTINIFOLIA* (H. B. et Kunth nov. gen. am. 5. p. 162. t. 447.) foliis obovatis obtusis basi attenuatis submembranaceis, junioribus tomento rufo subdeciduo utrinquè velutinis, adultis supernè glabris subtùs subvelutinis, racemo simplici confertifloro subvelutino, glandulis calycinis continuis. 5 in Mexico ad Acapulco. (v. s.)
6. *B. FERRUGINEA* (H. B. et Kunth nov. gen. am. 5. p. 151. t. 446.) foliis obovato-subrotundis brevissimè acuminatis basi cuneatis suprà molliter pubescenti-hirtellis, subtùs ramulis calycibusque ferrugineo-tomentosis, racemis simplicibus, glandulis calycinis 10 distinctis. 5 in sylvis Orinocensibus.
7. *B. ALTISSIMA*, foliis ovato-oblongis, subtùs rufo-tomentosis, supernè pilis centro adfixis bisetosis hispidis, racemo rufo-hirsuto. 5 in Guianæ sylvis. *Malpighia altissima* *Aubl. guian.* 1. p. 455. t. 181. Arbor 60-80-ped. Flores albi. Drupa rubescens subexsucca.
8. *B. CRASSIFOLIA*, foliis ovatis, supernè demùm glabris, subtùs rufo-velutinis obtusis, racemo erecto elongato rufo-velutino. 5 in Guianâ et Cayennâ et ad Orinocum? *Malpighia crassifolia* Lin. sp. 610. *Aubl. guian.* 1. p. 457. t. 182. *B. crassifolia* H. B. et K. l. c? Flores lutei. Drupæ virescentes. (v. s.)
β. Moureila (*Aubl. l. c.* t. 183.) foliis acutis. 5 in Guianâ et Cayennâ. — *Cav. diss.* 8. p. 412. t. 241. (v. s.)
9. *B. CRYSOPHYLLA* (H. B. et K. l. c.) foliis oblongis breviter acuminatis basi acutis margine subundulatis et revolutis, suprà glabris, subtùs sericeo-tomentosis et aureo-ferrugineis, racemis simplicibus, calycibus eglandulosis. 5 ad S. Carlos del Rio-Negro in Mission. Orinocens.

10. *B. MONTANA* (H. B. et Kunth l. c.) foliis elliptico-oblongis utrinque acutis suprâ glabris subtus pubescentibus, ramulis calycibusque ferrugineo-tomentosis, racemis ramosis. 5 in monte Cocollar Cumanensium. Fr. ign.
11. *B. LANCEOLATA* (Poir. dict. 4. p. 332.) foliis lanceolato-oblongis supernè demùm glabris, subtus rufo-velutinis, racemo erecto elongato supernè composito. 5 in Cayennâ. An à *B. crassifolia* satis differt? Fr. ign. (v. s?)
12. *B. PULCHRA*, foliis ovali-oblongis utrinque acutis subtus rufo-velutinis, racemis erectis simplicibus, drupis nucleisque ovatis apice umbonato-mucronatis. 5 in Mexici montibus. Malp. pulchra fl. mex. ic. ined. Drupæ coccineæ. Petala cordata, nascentia flava, dein rubescentia.
13. *B. SERICEA*, foliis elliptico-oblongis utrinque acuminatis, supernè glabris lucidis, subtus sericeo-rufo-velutinis, racemis subramosis subvelutinis. 5 in Brasiliâ. Theresmin. (v. s.)
14. *B. CINEREA*, foliis angusto-lanceolatis supernè lucidis, junioribus utrinque subrufo-velutinis, adultis subtus ramisque cinereo-tomentosis, racemis spicatis simplicibus. 5 in Guianâ Gallicâ. Perrottet. Malp. cinerea Poir. suppl. 4. p. 7.
15. *B. RHOPALEFOLIA* (H. B. et Kunth nov. gen. am. 5. p. 148.) foliis ellipticis obtusiusculis basi rotundatis complicatis glabris, junioribus subtus ramulis calycibusque ferrugineo-tomentosis, racemis simplicibus. 5 in calidis prov. Cumanensis et Caracasanae. Fr. ign.
16. *B. NITIDISSIMA* (H. B. et K. l. c.) foliis subsessilibus ellipticis utrinque rotundatis, suprâ puberulis nitidissimis, subtus ramulisque rufo-pubescentibus, racemis simplicibus, calycibus rufo-tomentosis 10-glandulosis. 5 in Islâ Panumanâ miss. Orinoc.

** *Folius adultis utrinque glabris.*

17. *B. SPICATA*, foliis lanceolatis glabris subtus ferrugineis, racemis spicatis erectis confertis subvelutinis, petalis hastatis obtusissimis. 5 in Sancto Domingo, Guadalupâ, Porto-Ricco, Brasiliâ. Malp. spicata Cav. diss. 8. p. 409. t. 237. (v. s.)
18. *B. LUCIDA*, foliis obovato-cuneiformibus glabris aveniis nitidis, racemis spicatis erectis abbreviatis glabris, petalis hastato-reniformibus. 5 in ins. Caribæis, S. Thomas, Guadalupâ. Malp. lucida Swartz. fl. ind. occ. 2. p. 852. (v. s.)
19. *B. COCCOLORÆFOLIA* (H. H. et Kunth nov. gen. am. 5. p. 148.) foliis ramulisque glabris sessilibus subrotundo-ovatis obtusis obsolete cordatis, racemis simplicibus, calycibus glabriusculis. 5 in aridis prov. Caracasanae prope Villa de Cura. Fr. ign.
20. *B. LÆVIGATA*, foliis ovatis obtusis lævibus, racemis spicatis confertifloris elongatis. 5 in Cayennâ et forsan in Guadalupâ. Malp. lævigata Poir. dict. 4. p. 332. Petala albo-flavida.
21. *B. ELEGANS*, foliis oblongis vel oblongo-lanceolatis membranaceis glaberrimis, racemis elongatis nutantibus, pedicellis glandulosis. 5 in littore ins. Guianensis Arowabisch. Malp. elegans Meyer prim. esseq. 178. Petala carnea.
22. *B. ANGUSTIFOLIA* (H. B. et Kunth nov. gen. am. 5. p. 153. t. 449.) foliis lineari-oblongis apice retusis basi acutiusculis glabris subtus cretaceo-pruinosis, stipulis infernè connatis, racemis ramosis; calycibus sericeo-pubescentibus 10-glandulosis. 5 ad confluentem fl. Atabapi et Orinoci.
23. *B. DENSEA*, foliis ovato-lanceolatis coriaceis utrinque lucidis, racemis dense spicatis, pedicellis pilosis. 5 in Guianâ. Malp. densa Poir. suppl. 4. p. 7. Petala rubra.
24. *B. CORIACEA*, foliis ovatis acutis integerrimis utrinque glabris, racemis dense spicatis pubescentibus erectis. 5 in fruticetis montosis Jamaïcæ

- australis. Flores albi odorati. Malp. coriacea Swartz fl. ind. occ. 2. p. 854. — Sloan. hist. 2. p. 20. t. 163. f. 1. Flores albi odorati. (v. s.)
25. *B. BRACHYSTACHIA*, foliis ovalibus utrinquē acutis integerrimis glabris, racemo spicato erecto rufo-velutino foliis duplō breviorē. 5 in Brasiliā. (v. s.)
26. *B. PALLIDA*, foliis ovatis utrinquē glabris suprà lucidis valdē coriaceis pallidis, racemis spicatis, pedicellis glabris brevibus, drupis parvis globoso-pyriformibus. 5 in Cayennā. Malp. pallida Poir. suppl. 4. p. 7.
27. *B. RETICULATA*, foliis ovato-lanceolatis reticulatis glabris supernē lucidis, racemis amplis compositis apice reflexis, bracteis ovatis. 5 in Cayennā. Malpighia reticulata Poir. suppl. 4. p. 8. Petala albo-flavescentia.
28. *B. MOLLIS*, foliis ovato-oblongis glabris lævibus coriaceis, racemo paniculato brevi glabro. 5 in Cayennā. Malp. mollis Poir. suppl. 4. p. 6.
29. *B. DIPHYLLO* (Jacq. am. 136.) foliis ovalibus glabris integerrimis, racemis simplicibus in ramusculis diphyllis. 3 in sylvaticis maritimis Carthagenæ. Flores flavi. Fructus rubri.
30. *B. VOLUBILIS*, foliis ovalibus acuminatis glabris lucidis, ramis volubilibus sæpē tuberculatis, racemis corymboso-umbellatis terminalibus. 5 in Indiā occid. Malp. volubilis Sims bot. mag. t. 809. Flores lutei. Fruct. ign. An Banisteriæ species?

III. *BUNCHOSIA*. Juss. ann. mus. 18. p. 481. *H. B. et Kunth nov. gen. am.* 5. p. 183.

Cal. 5-partitus, extūs ad basin glandulis 8-10 instructus. Pet. unguiculata. Stam. 10 basi monadelphæ. Stylus 1 simplex aut apice bi-trifidusve. Drupa nucleos 2 rarissimē 3, hinc planos illinc convexos fovens. — Racemi axillares laxē spicati aut subpaniculati.

* *Foliis, petiolis, pedicellisve glandulosis.*

1. *B. GLANDULOSA*, foliis ovato-ellipticis acuminatis glabris, petiolis apice biglandulosis, racemis simplicibus folio brevioribus. 5 in Antillis. Malpighia glandulosa Cav. diss. 8. p. 411. t. 239. f. 2.
2. *B. GLANDULIFERA* (H. B. et Kunth nov. gen. am. 5. p. 154.) foliis elliptico-ovatis brevissimē petiolatis undulatis utrinquē pubescentibus basi subtūs 4-glandulosi, racemis simplicibus, pedicellis 1-glandulosi. 5 in sylvis ad Caracas et in Guadalupā. Malpighia glandulosa Jacq. ic. rar. t. 469. M. glandulifera Jacq. coll. 4. p. 207 et 5. p. 5. f. 3. Flores flavi. Fruct. rubri.
3. *B. CORNIFOLIA* (H. B. et K. l. c.) foliis ellipticis acuminatis basi acutis utrinquē ramulisque pilis adpressis argenteis conspersis basim versus 2-glandulosi, racemis simplicibus, pedicellis 1-glandulosi. 5 in regno Novo-Granat. inter Turbaco et Carthagena. Petala alba. Drupa flava.
4. *B. POLYSTACHYA*, foliis oblongis acutis glabris nitidis subtūs basi biglandulosi, racemis subpaniculatis, pedicellis 1-glandulosi, stigmatē capitato bilobo. 5 in Indiā occid. Malp. polystachya Andr. bot. rep. t. 604.
5. *B. MEDIA*, foliis oblongo-lanceolatis acutis glabris subtūs glandulis 2 à basi remotiusculis costæ approximatis, racemis simplicibus, pedicellis medio uniglandulosi, stigmatē capitato bilobo. 5 in Indiā occid. Malp. media Ait. h. Kew. ed. 2. v. 3. p. 103.
6. *B. PILOSA* (H. B. et K. l. c.) foliis subsessilibus oblongis acutis basi rotundatis suprà pilosis, subtūs ramulisque hispido-pilosis et canescentibus, marginem versus 7-9-glandulosi, racemis simplicibus, pedicellis 1-glandulosi. 5 prope Turbaco Novo-Granatensium.
7. *B. TUBERCOLATA*, ramis tuberculatis, foliis ovato-lanceolatis subpubescentibus, racemis folio triplō brevioribus, pedicellis suprà basim 1-glandulosi. 5 ad Caracas Amer. Malpighia tuberculata Jacq. schœubr. 1. p. 54. t. 104. Flores flavi. Fructus rubri.

8. *B. ARGENTEA*, ramis puberulis, foliis lanceolatis subtus sericeo-argenteis, racemis oppositis simplicibus pubescentibus, pedicellis 1-glandulosis. 5 ad Caracas. Malpighia argentea Jacq. fragm. 186. t. 83. Flor. flavi. Glandulæ calycinæ elongatæ.

9. *B. GLAUCA* (H. B. et Kunth nov. gen. am. 5. p. 155.) foliis elliptico-oblongis acuminatis basi acutis, supra glabrescentibus, subtus adpresso-pilosis et glaucescentibus, marginem versus 4-11-glandulosis, ramulis verruculosis, racemis oppositis pubescentibus, pedicellis 1-glandulosis. 5 prope Hondam Novo-Granatens. Affinis *B. nitidæ*. Petala flava.

** *Foliis, petiolis, pedicellisque eglandulosis.*

10. *B. NITIDA*, foliis oblongis acuminatis glabris eglandulosis, racemis simplicibus folio sub brevioribus. 5 in Jamaicæ, Guadalupæ, Sancti-Domingi apricis cretaceis. Malp. nitida Lin. sp. 609. Cav. diss. 8. p. 411. t. 239. f. 1. (v. s.)

11. *B. ARMENIACA*, foliis ovato-oblongis acutis coriaceis glabris, racemis elongatis folii ferè longitudine, stylis subdistinctis 5 in Peruvizæ prov. Chanca. Malp. Armeniaca. Cav. diss. 8. p. 410. t. 238. Drupa refert Prunum Armeniacam. Nucleus dicitur venenosus. Folia 4-5-poll. longa!

12. *B. ODORATA*, foliis ovatis emarginatis utrinque tomentosis, racemis oppositis, stigmatibus 2-3-fido. 5 in fruticetis Carthagenæ. Malp. odorata Jacq. am. 136. t. 177. f. 41. Flores flavi suaveolentes.

†. *Species non satis notæ.*

13. *B? PANICULATA*, foliis oblongis cordatis acuminatis glabris, racemis paniculatis alaribus terminalibusque. 5 in Jamaicâ. Flores purpurei. Malp. paniculata Mill. dict. n. 6. †.

14. *B. CANESCENS*, foliis oblongis obtusis pubescentibus, racemis axillaribus compositis. 5 in Indiâ occid. Malp. canescens Ait. h. kew. 2. p. 105. †.

15. *B? SESSILIFOLIA*, foliis oblongo-lanceolatis basi attenuatis subsessilibus apice acuminatis, racemis oppositis simplicibus, stylo crassiusculo, stigmatibus orbiculato. 5 in Mexico. Petala flava margine subfimbriata. Fruct. ign.

IV. GALPHIMIA. Cav. ic. 5. p. 61.

Cal. 5-partitus eglandulosus. Pet. unguiculata ovalia aut oblonga. Stam. 10 ferè libera. Styli 3. Fructus (ex Cav.) 3-locul? 3-coccus, coccis dorso dehiscentibus 1-spermis. — Flores flavi. Racemi terminales ut in Byrsonimâ, à quâ differt tantum defectu glandularum calycis.

1. *G. HIRSUTA* (Cav. ic. 5. p. 62.) foliis ovatis acutis utrinque hirsutis breviter petiolatis, 5 in regno Mexicano.

2. *G. GLAUCA* (Cav. ic. 5. p. 61. t. 489.) foliis ovatis obtusis glabris subtus glaucis basi utrinque 1-dentatis, petiolis eglandulosis. 5 in Mexico.

3. *G. GLANDULOSA* (Cav. ic. 6. p. 43. t. 563.) foliis ovali-lanceolatis glabris, petiolis grossè ad apicem biglandulosis, petalis oblongis. 5 in Mexici occidentalis declivitate montium. H. B. et Kunth nov. gen. am. 5. p. 172. Malp. bi-glandulosa Poir. suppl. 4. p. 7. An hic forsitan species plurimæ, nempe:

a. *ovalifolia* (fl. mex. ic. ined.) foliis ovalibus obtusis.

β. *oblongifolia* (fl. mex. ic. ined.) foliis ovali-oblongis utrinque attenuatis.

γ. *lanceolata* (Cav. l. c.) foliis lanceolatis utrinque valdè acuminatis.

†. *Species non satis notæ, calycibus glandulosis donatæ.*

4. *G? LONGIFOLIA* (H. B. et Kunth nov. gen. am. 5. p. 173.) foliis lanceolato-oblongis acutis basi cuneato-angulatis glabris, petiolis 1-2-glandulosis, racemis terminalibus ramosis. 5 in umbrosis ad flumen. Cassiquiare.

5. *G? MOLLIS* (H. B. et Kunth l. c.) foliis oblongis utrinque acutis coriaceis

suprà glabris nitidis subtùs molliter canescenti-pubescentibus, petiolis biglandulosis, umbellis axillaribus? paucifloris. § in temperatis Mexici.

V. CAUCANTHUS. *Forsk. descr.* 91. †.

Cal. campanulatus, 5-partitus, eglandulosus. Pet. unguiculata, concava. Stam. 10 filam. subulatis. Antheræ margine prominulo. Styli 3 subulati. Fruct. ign. An satis à Galphimiâ differt?

1. C. ARABICUS (Lam. dict. 1. p. 658.) §. 5 in montibus Arabiæ ad Tæes. Malp. *Caucanthus* Poir. suppl. 4. p. 6. Folia opposita orbiculata glabra sæpè emarginata. Corymbi terminales. Flores albi. Fruct. dicitur ovi columbini magnitudine; an idèò carnosus?

Trib. II. HIPTAGÆÆ.

Stylus 1, seu 3 in 1 concreti. Carpella fructûs sicca indehiscencia 1-sperma sæpè in alas variè expansa. — Folia opposita aut verticillata.

- VI. HIPTAGE. *Gaertn. intr.* p. 126. *fruct.* 2. p. 169. *t.* 116. — *Gærtnera Schreb. gen. n.* 735. non Lam. — *Molina Cav. diss.* 9. p. 435. non Juss.

Calyx 5-partitus, basi 5-glandulosus. Pet. fimbriata. Stam. 10, unico ceteris multò longiore. Carpella 3 (abortu sæpè 1-2) 4-alata, alis inæqualibus.

1. H. MADABLOTA (Gaertn. l. c.) foliis ovato-lanceolatis acuminatis, racemis terminalibus. § in Indiâ orient. Madablota Sonn. voy. ind. 2. t. 135. Molina racemosa Cav. l. c. t. 263. Gaertnera racemosa Roxb. cor. 1. p. 19. t. 18. Bannisteria Benghalensis Linn. B. unicusularis Lam. Calophyllum Akara Burm! ind. 121. monente cl. Choisy. (v. s.)
2. H. OBTUSIFOLIA, foliis obovatis obtusis submucronulatis, racemis terminalibus et axillaribus. § in Chinâ. Gaertnera obtusifolia Roxb. cat. calc. 32. (v. s.)

- VII. TRISTELLATEIA. *Pet. Th. gen. mad.* p. 14. n. 47. — *Zymum Nor. Juss. ann.* 18. p. 482.

Cal. 5-fidus. Pet. inflexa unguiculata. Stam. 10, 5 alterna minora. Ovarium poris 3 impressum. Carpella 3 appendicibus 6 coronata. Embryo convolutus.

1. T. MADAGASCARIENSIS (Poir. suppl. 5. p. 367.) § in Madagascar. Frutex volubilis, foliis inf. quaternatim verticillatis, sup. oppositis basi glandulosis integris, floribus racemosis luteis.

VIII. THRYALLIS. *Lin. gen. n.* 533.

Cal. 5-partitus persistens. Pet. subrotunda. Stam. 10 subulata. Capsula 3 quetra tripartibilis, loculis exteriori angulo dehiscentibus. †.

1. T. BRASILIENSIS (Lin. sp. 554.) § in Brasiliâ. — *Marcg. bras.* p. 79. f. 3. Rami articulati. Folia ovata petiolata. Racemus terminalis. Flores parvi, lutei. †.

- IX. ASPICARPA. *Rich. in mem. mus.* 2. p. 399. — *Acosmus Desv.*

Cal. 5-partitus connivens. Pet. o. Stam. 1 inclusum. Ovarium subrotundum semifissum biloculare. Styl. 1 in fissurâ brevissimus. Fructus siccus indehiscens, abortu loculi unius transversus irregularis 1-spermus. Semen orbiculatum, exalbuminosum. Embryo recurvus, radi-

culâ basilari ad hilum lateralem factum versâ. Cotyled. crassæ inæquales.

1. *A. URENS* (Lag. nov. gen. et sp. 1. n. 5.) γ \mathcal{U} in Novâ-Hispaniâ. Caulis suffruticulosus volubilis filiformis. Folia cordato-ovalia, pilis centro adfixis adpressissimis. Flores fasciculati subsessiles axillares.

X. GAUDICHAUDIA. *H. B. et Kunth nov. gen. am. 5. p. 156. t. 445. Saint-Hil. bull. philom. 1823. p. 132.*

Cal. 5-partitus 5-fidusve, extus 10 sæpius 8-glandulosus. Pet. 5 interdum perigyna, subrotunda, patentia. Stam. 5 cum petalis inserta, filam. basi complanata connata, inæqualia, 2 antheris minoribus aut abortivis, connectivo tunc incrassato. Stylus 1. Carpella 3 libera aut basi coalita, 1-sperma, sæpius 1 abortivo Samaræ 2 inferius in membranam calcariformem productæ, dorso alatæ. Semen erectum ad apicem funiculi penduli. Album. o. Embryo rectus. — Frutices, fol. oppositis integris, floribus flavis.

1. *G. CYNANCHOIDES* (H. B. et Kunth l. c.) caule volubili fruticoso, foliis petiolatis, racemis axillaribus terminalibusque confertifloris. γ \bigcirc in Mexico prope Valladolid.

2. *G. GUARANITICA* (Saint-Hil. l. c.) caule volubili fruticoso, foliis petiolatis ovato-oblongis obtusis breviter mucronatis, petiolis apice 2-glandulosis, umbellis axillaribus 2-4-floris. γ \bigcirc in Brasiliâ.

3. *G. LINEARIFOLIA* (Saint-Hil. l. c.) caule suffruticoso erecto, foliis linearibus subsessilibus distantibus, umbellis terminalibus. γ in Brasiliâ.

4. *G. SERICEA* (Saint-Hil. l. c.) caule suffruticoso erecto, foliis breviter petiolatis subtus sericeis, intermediis ovato-lanceolatis, pedunculis filiformis 1-floris, rarius 2-floris. γ in Brasiliâ.

XI. CAMAREA. *Saint-Hil. bull. philom. 1823. p. 133.*

Cal. 5-partitus 5-fidusve, lobis basi 2-glandulosis interdum nudis. Pet. 5 subperigyna subinæqualia. Stam. 6; 3 fertilia, filam. ad apicem fere coalitis, antheris rotundis; 3 basi vix coalita, medio fertili, lateralibus sterilibus massulas petaloideas contortuplicatas antheræ vicem gerentibus. Styl. 1. Carpella 3 aut abortu 2, subconnata, 1-sperma, indehiscencia, dorso cristata, lateribus cristato-rugosa. Semen ad extrem. funiculi erecti pendulum. Album. o. Embryo rectus inversus. — Frutices, omnes Brasilienses, à cl. auctore sic indicati :

1. *C. HIRSUTA* (Saint-Hil. l. c.) foliis lanceolatis vel oblongo ovatove lanceolatis hirsutis margine sericeis, umbellis terminalibus.

2. *C. AXILLARIS* (Saint-Hil. l. c.) foliis lanceolatis acutis basi cordatis villosis patentibus, floribus solitariis axillaribus.

3. *C. SERICEA* (Saint-Hil. l. c.) foliis longis lineari-lanceolatis acutis sericeis, umbellis terminalibus.

4. *C. ERICOIDES* (Saint-Hil. l. c.) foliis parvis linearibus angustis confertis, floribus umbellatis.

5. *C. LINEARIFOLIA* (Saint-Hil. l. c.) foliis linearibus subdistantibus, floribus umbellatis.

Trib. III. *BANISTERIEÆ.*

Styli 3 distincti. Fructus carpella sicca indehiscencia monosperma, variè in alas expansa. — Folia opposita, rarius verticillata.

XII. HIRÆA. H. B. et Kunth nov. gen. am. 5. p. 167.

Cal. 5-sepalus glandulosus aut eglandulosus. Pet. subrotunda unguiculata Stam. 10, filamentis subulatis imâ basi subcohærentibus, alternis longioribus. Styli 3. Carpella samaræformia 3, rarius duo, 1-sperma, dorso cristata, alâ membranaceâ circumdata, basi et apice sæpius emarginata.

§. 1. *Calycibus eglandulosis*. — Hiræa. Jacq. am. 137.

*. *Carpellis subrotundis*.

1. H. RECLINATA (Jacq. am. 137. t. 176. f. 42.) foliis obovatis utrinquè obtusis supernè pubescentibus infernè glabris. ζ in sylvia Carthagenæ.
2. H. ODORATA (Willd. sp. 2. p. 743.) foliis ovatis acutis supernè glabris subtùs tomentosis. ζ in Guineâ. — Fructus Flabellarie paniculatæ Cav. diss. 9. p. 436. t. 264. Forsan huc pertinet, sed folium, monente cl. Brownio (bot. congo. p. 7.), ad Pierocarpum quemdam referendum est.
3. H. GLABRA (Spreng. neue entd. 2. p. 154.) foliis ovato-oblongis utrinquè glaberrimis nitidis, paniculâ terminali trichotomâ, ramulis calycibusque adpresso-pilosis. ζ in Brasiliâ.
4. H. HYPERICOIDES (Burch! cat. n. 2486 et 2531.) foliis linearibus utrinquè glabris. ζ ad Cap. Bonæ-Spei. Fructuum alæ rotundatæ semi-orbiculatæ. (v. s.)

** *Carpellis ovali-oblongis*.

5. H. NUTANS (Roxb! cat. calc. p. 34.) foliis ovatis acuminatis supernè glabris subtùs pilos (Malpighiaceos) adpressissimos centro adfixos gerentibus. ζ in Bengalo. Samaræ 2-3 ovali-oblongæ. (v. s.)
6. H. INDICA (Roxb. cat. calc. p. 90.) foliis subrotundo-ovatis subcordatis acuminatis nitidis glabris. ζ in montibus sylvis Coromandelie. Triopteris Indica Willd. spec. 2. p. 744. Roxb. cor. 2. p. 32. t. 160. Samaræ 3 oblongæ priori simillimæ. Calyces pubescentes.

§. 2. *Calycibus extùs glandulosis*. MASCAGNIA Bert. mss.

7. H. MACRADENA, foliis ovatis acuminatis supernè glabris subtùs ramulis petiolisque eglandulosis adpressè sericeo-villosis, floribus racemosis, pedicellis basi articulatis. ζ in Americâ cal. ad S.-Martham. Mascagnia Americana Bertero! Fructus alæ orbiculatæ utrinquè emarginatæ. Cal. glandulæ magnæ. (v. s.)
8. H. OBLONGIFOLIA, foliis ovato oblongis acuminatis glabris, nervis petioliis ramulisque velutino-pubescentibus, floribus paniculatis, pedicellis basi articulatis. ζ ad Sanctam-Martham cum priore cui valdè accedit. Mascagnia oblongifolia Bert! iued. (v. s.)
9. H. DIVARICATA (H. B. et Kunth nov. gen. am. 5. p. 169.) foliis oblongis acuminatis basi acutis subquintuplinerviis glabris, petiolis 3-7-glandulosis, paniculis axillaribus folium duplò superantibus è racemis 5-7 compositis, calycibus adpresso-pilosis. ζ prope Cumanam.
10. H. OVATIFOLIA (H. B. et Kunth l. c.) foliis ovatis acuminatis basi rotundatis glabris suprâ nitidis, petiolis eglandulosis, paniculis axillaribus, fructiferis folium subæquantibus, fructibus orbiculatis. ζ in umbrosis prope Cumanam. Flores ignoti.

†. *Species fructibus ignotis dubiæ*.

11. H? PRUNIFOLIA (H. B. et Kunth l. c.) foliis ellipticis utrinquè acutis margine glandulosis glabris, junioribus adpresso-puberulis, petiolis medio biglandulosis, racemis terminalibus, calycibus pubescentibus 8-glandulosis. ζ in prov. Caracasana prope La Victoria.
12. H? COMPLICATA (H. B. et Kunth l. c.) foliis ellipticis acutis basi rotundatis

subcordatis complicatis glabris, petiolis 2-4-glandulosis, racemis terminalibus et axillaribus, calycibus fusco-sericeis 10-glandulosis, petalis concavis dorso carinato-alatis. 5 in ripâ Orinoci prope Carichana.

13. H? *NIVIDA* (H. B. et Kunth l. c.) foliis ovato-ellipticis acutiusculis basi rotundatis subcoriaceis glabris supra nitidis, racemis axillaribus solitariis et terminalibus subquinis paniculatis, calycibus ferrugineo-hirsutis 8-glandulosis, petalis concavis dorso carinato-alatis. 5 prope S. Barbara del Alto-Orinoco.

††. *Species Mexicanæ non satis notæ, ad Mascagniam referendæ?*

14. H? *MUCRONATA* (fl. mex. ic. ined.) foliis ovalibus apice acutè mucronatis basi vix subcordatis; petiolis longiusculis eglandulosis, pedicellis plurimis axillaribus 1-floris. 5 in Mexico. Habitus Banisteriæ. Fructus ferè Hirææ. Calyx Triopteridis.
15. H? *CYCLOPTERA* (fl. mex. ic. ined.) foliis ovali-oblongis utriusque acuminatis subvillosis, petiolis eglandulosis, pedunculis villosis axillaribus subcorymbosis. 5 Scandens. Fructuum alæ magnæ ferè orbiculatæ ut in Hirææ, sed cal. biglandulosus?
16. H? *OXYOTA* (fl. mex. ic. ined.) foliis ovali-oblongis acutis villosis basi cordatis, auriculis parvis acutissimis, petiolis eglandulosis pedunculisque villosis. 5 in Mexico. Carpella 3 triala more Hirææ in calyce pedicellata. Calyx biglandulosus.
17. H? *PODOCARPA* (fl. mex. ic. ined.) foliis lanceolatis glabris breviter petiolatis, pedunculis 3-fidis. 5 in Mexico. Carpella 3 alata, in calyce pedicellata. Calyx forsan eglandulosus.
18. H? *ACUMINATA* (fl. mex. ic. ined.) foliis ovalibus acuminato-mucronatis villosis, petiolis eglandulosis, pedunculis villosis folio duplò longioribus apice corymbiferis. 5 in Mexico. Scandens. Samaræ bialatæ utrinque emarginatæ. Calyx biglandulosus.
19. H? *MACROPTERA* (fl. mex. ic. ined.) foliis ovali-oblongis utriusque acuminatis glabris, petiolis apice biglandulosis, racemis axillaribus. 5 in Mexico. Fructus exactè Hirææ ut pinxit Cav. diss. g. t. 264. sed ferè triplò major. Calycis lobi basi grossè biglandulosi.

XIII. TRIOPTERIS. *Lin. gen. n. 574. Cav. diss. g. p. 431.*

Cal. 5-partitus, laciniis basi extus biglandulosis. Pet. subrotunda unguiculata. Stam. 10, filamentis subulatis imâ basi subcohærentibus alternis majoribus. Styli 3. Carpella 3 basi subcoalita, 1-sperma, expansa in alas 3 nempe 2 superiores, 1 inferiorem et interdum in cristam minimam dorsalem. — Frutices volubiles.

1. *T. RIGIDA* (Swartz fl. ind. occ. 2. p. 859.) foliis subrotundis acuminatis marginatis lineatis coriaceis rigidis glaberrimis, petiolis eglandulosis 5 in fruticetis montosis Sancti-Domingi. Rami subvolubiles. (Swartz.) Fructus juniores puberuli. (v. s.)
2. *T. OVATA* (Cav. diss. g. p. 431. t. 259.) foliis ovatis obtusiusculis subcordatis glabris, petiolis apice biglandulosis. 5 in Sancto-Domingo. (v. s.)
3. *T. LUCIDA* (H. B. et Kunth nov. gen. am. 5 p. 167. t. 451.) foliis subrotundo-ellipticis utrinque acutis coriaceis glabris nitidis, petiolis eglandulosis, paniculis axillaribus et terminalibus simplicibus. 5 in umbrosis ins. Cubæ.
4. *T. HAVANENSIS* (H. B. et Kunth l. c.) foliis oblongis submucronatis basi acutis coriaceis glabris nitidis, petiolis eglandulosis. 5 prope Havanam. Fl. et fruct. ignoti.
5. *T. JAMAÏCENSIS* (Lin. spec. 612.) foliis oblongis acuminatis venosis glabris, petiolis eglandulosis. 5 ad sæpes Jamaicæ, Hispaniolæ. Rami volubiles. (v. s.)

6. *T. BRASILIENSIS* (Poir. dict. 8. p. 105.) foliis ovato-lanceolatis acutis coriaceis glabris nitidis, petiolis brevibus eglandulosis. $\frac{5}{5}$ in Brasiiliâ. An à priori satis differt? (v. s.)
7. *T. LINGULATA* (Poir. dict. 8. p. 104.) foliis lineari-lingulatis obtusis mucronatis utrinquè glabris superne nitidis. $\frac{5}{5}$ in Sancto-Domingo. Panícula terminalis, ramis divaricatis longissimis.
8. *T. FLORIBUNDA* (Bilberg in flora 1821. p. 331.) foliis oblongis obtusis glabris. $\frac{5}{5}$ in Brasiiliâ. Panícula terminalis.

XIV. *TETRAPTERIS*. Cav. diss. 9. p. 433. — *Triopteris* sp. Willd. spec. 2. p. 743.

Omnia *Triopteridis*, sed *Carpella expansa* in alas 4 oblongas, 2 superiores, 2 inferiores sæpius minores et insuper ex dorso cristas 1-4 alæformes aut subulæformes proferentia. — Petioli eglandulosi.

1. *T. BUXIFOLIA* (Cav. l. c. t. 262. f. 1.) foliis ovatis obtusis subsessilibus parvis glaberrimis, floribus umbellatis terminalibus. $\frac{5}{5}$ in Sancto-Domingo. Folia 8-9 lin. longa, 5-6 lata. Alæ fructuum in meo spec. purpurascentes. *Triopteris buxifolia* Willd. sp. 2. p. 745. (v. s.)
2. *T. ACAPULCENSIS* (H. B. et Kunth nov. gen. am. 5. p. 168.) foliis ovato-ellipticis obtusis brevissimè petiolatis cordatis coriaceis nitidis glabris, umbellis axillaribus solitariis 4-floris. $\frac{5}{5}$ in littore Mexicano prope Acapulco. Fl. ign.
3. *T. MUCRONATA* (Cav. l. c. t. 262. f. 2.) foliis ovatis acuminato-mucronatis breviter petiolatis glaberrimis, paniculis axillaribus ad apices ramorum subumbellatis. $\frac{5}{5}$ in Cayennâ. *Triopteris acuminata* Willd. sp. 2. p. 745. Petioli 3-lin. longi. Folia 4-poll. longa, 2-poll. lata. (v. s.)
4. *T. ACUTIFOLIA* (Cav. l. c. t. 261.) foliis ovato-lanceolatis acuminatis, nervo medio subtus pubente villis adpressissimis centro adfixis, paniculis terminalibus. $\frac{5}{5}$ in Cayennâ. *Triopteris acutifolia* Willd. sp. 2. p. 744. Cl. Cavan. dixit plantam suam glahram. An diversa? An pubes in descriptione omisa? An pili decidui? sed meum specimen omninò iconem refert. (v. s.)
5. *T. CITRIFOLIA* (Pers. ench. 1. p. 508.) foliis ovato-oblongis acutis glabris, umbellis axillaribus pedunculatis, alis fructûs binis inferioribus duplò brevioribus. $\frac{5}{5}$ in sylvis montium Jamaicæ. *Tetrapteris inæqualis* Cav. l. c. t. 260. *Triopteris bifurca* Gaertn. fr. 2. p. 163. t. 116. *Triopteris citrifolia* Swartz. fl. ind. occ. 2. p. 857. (v. s. ex Cayennâ? an eadem?)
6. *T. DISCOLOR* (Meyer prim. eas. 182. sub *Triopteride*.) foliis oblongis acuminatis utrinquè glabris subtus discoloribus, umbellis paniculatis terminalibus axillaribusque. $\frac{5}{5}$ in sylvis arenosis Guianæ prope Essequibo. Alæ fructûs exter. aliis triplò majores.
7. *T?* *PAUCIFLORA* (Meyer l. c. 183. sub *Triopteride*.) foliis oblongo-lanceolatis utrinquè glabris brevissimè petiolatis, umbellis axillaribus folio brevioribus. $\frac{5}{5}$ in dumetis Guianæ prope Essequibo.

XV. *BANISTERIA*. H. B. et Kunth nov. gen. am. 5. p. 161. — *Banisteriæ* capsulis discretis. Lin. gen. n. 573. Cav. diss. 9. p. 421.

Cal. 5-partitus, extûs ad basin 8-10-glandulosus. Petala unguiculata subrotunda. Stam. 10 filamentis subulatis basi cohærentibus. Styli 3 apice sæpius in foliola explanati! *Carpella* indehiscentia 3-1-sperma subdistincta, demum separabilia, desinentia in alam simplicem membranaceam ad latus superius incrassatam. Semen 1 pendulum in quoque loculo. Cotyled. crassæ inæquales. Radicula supra ad hilum versa. (1) — Arbores fruticesve sæpè scandentes.

(1) Sic in *B. angulatâ* vidi. Gaertnerus hilo oppositam descripsit in *B. laurifoliâ*.

* *Foliis basi cordatis lobatis aut angulato-dentatis.*

1. *B. PALMATA* (Cav. diss. 9. p. 430. t. 257. f. 2.) foliis palmatifidis subtus tomentosis, lobis acuminatis, petioli apice biglandulosi. ζ in Sancto-Domingo et Cayennâ. An *B. angulosæ* varietas? (v. s.)
2. *B. ANGULOSA* (Lin. sp. 611.) foliis cordatis obtusè angulosis subpanduriformibus, subtus tomentosis aut sericeo-pubescentibus, petioli apice biglandulosi. ζ in Sancto-Domingo, Cayennâ. — Plum. ed. Burm. t. 92. Cav. diss. t. 252. (v. s.)
3. *B. SINUATA*, foliis cordatis ovato-rotundis obtusissimè sinuatis supernè glabris subtus vix pubescentibus, petioli apice biglandulosi. ζ in Gujanâ. Perrotet. Carpella ad latus internum appendice majore quàm in *B. angulosâ* donata, alæ laterales subnullæ ad cristas reductæ. (v. s.)
4. *B. VARIFOLIA*, foliis cordatis subtus villosis canescentibus, aliis ovatis indivisis aliis trilobis, apice acutis, petioli apice biglandulosi. ζ ad Sanctam-Martham. *B. diversifolia* Bertero! ined. non Kunth. *B. vitifolia* f. mex. ic. ined. forsân eadem. (v. s.)
5. *B. SAGITTATA* (Cav. l. c. t. 257. f. 3.) foliis hastato-sagittatis subtus tomentosis acutè subangulatis, petioli apice biglandulosi, auriculis truncatis. ζ in Sancto-Domingo.

** *Foliis cordatis integris.*

6. *B. AURICULATA* (Cav. diss. p. 428. t. 255.) foliis profundè cordatis subsagittatis glabris, lobis rotundatis, petioli apice glandulosi, floribus umbellatis. ζ in Brasiliâ prope Rio-Janeiro.
7. *B. CILIATA* (Lam. dict. 1. p. 369.) foliis cordatis orbiculatis glabris ciliato-denticulatis, petioli apice biglandulosi, floribus umbellatis. ζ in Brasiliâ.
8. *B. SPLENDENS*, foliis cordatis reniformi-orbiculatis subtus tomentoso-sericeis, petioli apice biglandulosi, racemis axillaribus dichotomis divaricatis. ζ in Americâ calid. — Sloan. hist. 2. t. 162. f. 2. *B. fulgens* Lam. dict. 1. p. 368. Cav. diss. 9. p. 426. t. 253. *B. heterophylla* Willd. sp. 2. p. 742. Sed non *heterophylla*; folia floralia tantum orbiculata subsessilia. (v. s.)
9. *B. HUMBOLDTIANA*, foliis subrotundo-ovatis cordatis subacuminatis mucronatis membranaceis, suprâ glabrinsculis, subtus ramulisque molliter canotomentosis, basi biglandulosi, umbellis alaribus lateralibus et terminalibus sessilibus. ζ inter Carthagenam et Cerro de la Popa. *B. tiliaefolia* H. B. et Kunth nov. gen. am. 5. p. 162. non Vent.
10. *B. DICHOTOMA* (Lin. sp. 612.) foliis cordatis subovatisve acuminatis utrinquè glabris, petioli apice biglandulosi, racemis axillaribus folio longioribus dichotomis subdivaricatis. ζ in Americâ calid. — Plum. ed. Burm. t. 13. *B. convolvulifolia* Cav. diss. 9. p. 428. t. 256. (v. s.)
11. *B. TIMORENSIS*, foliis cordatis rariùs subovatis acuminatis utrinquè glabris, petioli apice biglandulosi, racemis axillaribus folio brevioribus dichotomo-subumbellatis. ζ in ins. Timor. Priori affinis. (v. s.)
12. *B. EMARGINATA* (Cav. diss. 9. p. 425. t. 249.) foliis ellipticis basi cordatis apice mucronatis aut emarginatis subtus tomentosis, petioli brevibus apice biglandulosi, racemis terminalibus axillaribusque. ζ in Guadalupâ (v. s.)

*** *Foliis ovatis oblongisve.*

13. *B. MICROPHYLLA* (Jacq. obs. 3. p. 7. t. 56.) foliis ellipticis utrinquè rotundatis utrinquè glabris rigidis breviter petiolatis, racemis terminalibus. ζ in Carolinâ? †.
14. *B. UMBELLULATA*, foliis ellipticis basi rotundatis apice emarginatis utrinquè glabris rigidulis, petioli brevibus apice biglandulosi, umbellis subcorymbosis terminalibus. ζ in Sancto-Domingo. Bertero. An eadem ac præcedens, sed florum dispositio diversa et fructus ignoti. (v. s.)

15. B? FLORIBUNDA, foliis ovalibus obtusis emarginatis coriaceis glabris, petiolis pubescentibus eglandulosis, racemis corymbosis pedunculatis folii longitudine. 5 in Porto-Ricco. Bertero. Flores numerosi ampli, ex sicco flavi. (v. s. in h. Balbis.)
16. B. SERICEA (Cav. diss. 9. p. 429. t. 258.) foliis ovatis obtusis cum mucrone, junioribus utrinquè, adultis subtus, tomentosis, petiolis medio biglandulosis, ramis ancipitibus, floribus racemosis. 5 in Brasiliâ. Tomentum folii aureum fulgidum.
17. B? TOMENTOSA (Desf. cat. 160.) foliis ovatis obtusis cum mucrone subtus tomentosis, pube ramosâ, petiolis apice biglandulosis, corymbis subpaniculatis. 5 in Antillis. Stamina 15 basi monadelpha, 5 antè petala majora, cetera subabortiva vidi in specim. florido in hort. Mus. Par. 1806. (v. v.)
18. B. TILIÆFOLIA (Vent. choix t. 50.) foliis orbiculatis acuminatis subtus tomentosis, petiolis elongatis apice biglandulosis, umbellis axillaribus pedunculatis compositis, petalis subsessilibus. 5 in Javâ. Fl. purpurei.
19. B. QUAPARA (Aubl. guian. 1. p. 464. t. 196.) foliis ovatis acutis subtus petiolisque tomentosis supernè puberulis, pilis centro adfixis utrinquè productis adpressis, umbellis axillaribus pedunculatis, fructuum alis erectis. 3 in Guianâ ad margines pratorum. Cav. 1. c. t. 250.
f. mucronulata, foliis minoribus mucronatis. 3 in Amer. merid. Bertero. (v. s. sine fr.)
20. B. SINEMARIENSIS (Aubl. guian. 1. p. 462. t. 185.) foliis ovatis acuminatis supernè glabris subtus et in petiolo apice biglanduloso setis adpressissimis centro adfixis tectis, corymbis racemosis axillaribus longè pedunculatis. 5 in Guianâ ad pratorum margines, et in Sancto-Domingo. Cav. 1. c. t. 251. (v. s.)
21. B? LAURIFOLIA (Lin. sp. 611.) foliis ovato-oblongis, acutis subcoriaceis glabris, petiolis eglandulosis, racemis paniculatis terminalibus ferrugineo-tomentosis, stigmatibus dimidiatis. 5 in fruticetis montosis Jamaicæ et Hispaniolæ. Swartz obs. 182. * An Heteropteris? (v. s. sine fr.)
22. B. MURICATA (Cav. 1. c. p. 423. t. 246. f. 2.) foliis ovatis subtus tomentosis breviter petiolatis, racemis paniculatis terminalibus, carpellis intus sculeo auctis. 5 in Peru.
23. B. LEONA (Cav. diss. 9. p. 424. t. 247. excl. fruct. et sp. amer.) foliis ovato-oblongis acuminatis coriaceis reticulato-venosis glabris subtus subglaucescentibus, petiolis eglandulosis brevibus, paniculis rufo-velutinis terminalibus folio brevioribus. 5 in Sierrâ-Leonâ. Smeathman. (v. s.)
24. B. MULTIFLORA, foliis ovato-oblongis acuminatis coriaceis reticulato-venosis glabris supernè lucidis subtus glaucis, petiolis eglandulosis brevibus, paniculis terminalibus rufo-tomentosis folio longioribus, fructibus junioribus velutinis. 5 in Cayennâ. Patris. Valdè accedit ad B. Leonam. (v. s.)
25. B. MUCRONATA, foliis ovalibus acuminato-mucronatis utrinquè glabris submembranaceis, petiolo brevi apice biglanduloso nervisque adpressissimè pubescentibus, corymbis axillaribus paucifloris. 5 in Novâ-Hispaniâ? (v. s. sine fr.)
26. B. FERRUGINEA (Cav. diss. 9. p. 424. t. 248.) foliis ovatis acuminatis supernè glabris nitidis subtus ferrugineis petiolisque adpressè puberulis, glandulis in parte limbi inferiore sitis, racemis pauciculatis, fructuum pubescentium alis erectis. 5 in Brasiliâ. Folia 2 poll. longa. (v. s.)
f. fulgens (Meyer. pr. esseq. p. 181.) foliis 4-5 poll. longis. 5 in Guianæ sylvis.
27. B. PERILOCAEOTIA (Desf. cat. h. par. ed. 2. p. 160?) foliis ovatis oblongis mucronatis subcoriaceis utrinquè glabris, petiolo apice biglanduloso, corymbis terminalibus, carpellis basi cristatis pilosisque. 5 in Porto-Ricco. Bertero.
a. subovata, foliis ovato-oblongis. (v. s.)

* *Foliis basi cordatis lobatis aut angulato-dentatis.*

1. *B. PALMATA* (Cav. diss. 9. p. 430. t. 257. f. 2.) foliis palmatifidis subtus tomentosis, lobis acuminatis, petiolis apice biglandulosi. ζ in Sancto-Domingo et Cayennâ. An *B. angulosæ* varietas? (v. s.)
2. *B. ANGULOSA* (Lin. sp. 611.) foliis cordatis obtusè angulosis subpanduriformibus, subtus tomentosis aut sericeo-pubescentibus, petiolis apice biglandulosi. ζ in Sancto-Domingo, Cayennâ. — Plam. ed. Burm. t. 92. Cav. diss. t. 252. (v. s.)
3. *B. SINUATA*, foliis cordatis ovato-rotundis obtusissimè sinuatis supernè glabris subtus vix pubescentibus, petiolis apice biglandulosi. ζ in Gujanâ. Perrotet. Carpella ad latus internum appendice majore quàm in *B. angulosâ* donata, alæ laterales subnullæ ad cristas reductæ. (v. s.)
4. *B. VARIIFOLIA*, foliis cordatis subtus villosis canescentibus, aliis ovatis indivisis aliis trilobis, apice acutis, petiolis apice biglandulosi. ζ ad Sanctam-Martham. *B. diversifolia* Bertero! ined. non Kunth. *B. vitifolia* f. mex. ic. ined. forsân eadem. (v. s.)
5. *B. SAGITTATA* (Cav. l. c. t. 257. f. 3.) foliis hastato-sagittatis subtus tomentosis acutè subangulatis, petiolis apice biglandulosi, auriculis truncatis. ζ in Sancto-Domingo.

** *Foliis cordatis integris.*

6. *B. AURICULATA* (Cav. diss. p. 428. t. 255.) foliis profundè cordatis subsagittatis glabris, lobis rotundatis, petiolis apice glandulosi, floribus umbellatis. ζ in Brasiliâ prope Rio-Janeiro.
7. *B. CILIATA* (Lam. dict. 1. p. 369.) foliis cordatis orbiculatis glabris ciliatodenticulatis, petiolis apice biglandulosi, floribus umbellatis. ζ in Brasiliâ.
8. *B. SPLENDENS*, foliis cordatis reniformi-orbiculatis subtus tomentoso-sericeis, petiolis apice biglandulosi, racemis axillaribus dichotomis divaricatis. ζ in Americâ calid. — Sloan. hist. 2. t. 162. f. 2. *B. fulgens* Lam. dict. 1. p. 368. Cav. diss. 9. p. 426. t. 253. *B. heterophylla* Willd. sp. 2. p. 742. Sed non *heterophylla*; folia floralia tantum orbiculata subsessilia. (v. s.)
9. *B. HUMBOLDTIANA*, foliis subrotundo-ovatis cordatis subtus mucronatis membranaceis, suprâ glabriusculis, subtus ramulis tomentosis, basi biglandulosi, umbellis alaribus lateralibus sessilibus. ζ inter Carthagenam et Cerro de la Popa et Kunth nov. gen. am. 5. p. 162. non Vent.
10. *B. DICHOTOMA* (Lin. sp. 612.) foliis cordatis subovatisve glabris, petiolis apice biglandulosi, racemis axillaribus dichotomis subdivaricatis. ζ in Americâ calid. — Plam. ed. Burm. t. 92. Cav. diss. 9. p. 427. *B. convolvulifolia* Cav. diss. 9. p. 427. (v. s.)
11. *B. TIMORIENSIS*, foliis cordatis ovatisve glabris, petiolis apice biglandulosi, chotomo-subumbellatis. ζ in Timori.
12. *B. EMARGINATA* (Cav. diss. 9. p. 427.) foliis mucronatis aut emarginatis subtus biglandulosi, racemis terminalibus.

*** *Foliis*

13. *B. MICROPHYLLA* (Jacq. obs. 3. p. 17.) foliis utrinquè glabris rigidis breviter Carolinâ? f.
14. *B. UMBELLULATA*, foliis ellipticis glabris rigidulis, petiolis brevibus basi terminalibus. ζ in Sancto-Domingo, sed florum dispositio diversa

7. *H. NITIDA* (H. B. et Kunth l. c.) foliis ovato-oblongis acuminatis supra nitidis subtus argenteis, petioliis eglandulosis, paniculâ terminali foliosâ. ζ in Brasilâ, Banist. nitida Lam. dict. 1. p. 369. Cav. diss. 9. t. 244 Flor ign.
8. *H. CORNIFOLIA* (H. B. et Kunth l. c.) foliis ellipticis utrinquè acutis subcoriaceis, supra glabris nitidis, subtus adpresso-pilosiusculis et glaucescentibus, petioliis eglandulosis, paniculis axillaribus et terminalibus. floribus racemoso-corymbosis, calycibus sericeis. ζ prope Ibague Novo-Granatensium.
9. *H. PLATYPTERA*, foliis ovato-oblongis utrinquè obtusis coriaceis glabris, petioliis brevissimis eglandulosis, paniculis terminalibus, fructuum alis latis. ζ in Guadalupâ. Folia 7-8 poll. longa, 3 lata. An eadem ac *H. brachiata*? (v. s.)
10. *H. APPENDICULATA*, foliis ovalibus in acumen obtusum elongatis subcoriaceis supernè glabris subtus pallidis, in nervis adpressè puberulis, petioliis brevissimis eglandulosis, corymbis axillaribus, carpellis intus ad basin alâ appendiculatis. ζ in ins. Sancti-Vincentii ubi forsan culta. Banisteria appendiculata Lamb! herb. (v. s.)
11. *H. FLORIBUNDA* (H. B. et Kunth l. c. 5. p. 166.) foliis oblongis utrinquè acutis subcoriaceis glabris, ramulis petioliis paniculis axillaribus et terminalibus calycibusque ferrugineo-tomentosis eglandulosis. ζ in montibus Mexicanis. Affinis *H. cæruleæ* sed fructus ign. ex Kunth.
12. *H. LONGIFOLIA* (H. B. et Kunth l. c.) foliis oblongis utrinquè acutis subcoriaceis glabris, petioliis eglandulosis, paniculis axillaribus calycibusque ferrugineo-tomentosis glandulosis, floribus racemosis. ζ prope urbem Mexici.

Sect. II. *Foliis alternis*. ANOMALOPTERIS.

13. *H. SMEATHMANNI*, foliis alternis ovali-oblongis acuminatis integerrimis glabris, petioliis eglandulosis, fructibus introrsum exappendiculatis. ζ in Sierra-Leonâ. Frustula tantum habeo sine flore. Ad hanc forsan alludit cl. Browa (bot. cong. p. 7.) dicens, sed sine descr., genus quoddam flore à Banisteria diversum et alternifolium ex Africâ æquinociali vidisse. (v. s. sine fl.)

†. *Genus Malpighiaceis affine.*

XVII? NIOTA. Lam. ill. t. 299. — Samadera Gaertn. fr. 2. p. 352. t. 156. — Biporeia Pet. Th. gen. mad. p. 14.

Cal. 4-5-partitus, persistens, lobis 2 externè glandulosis. Pet. 4-5 obtusa calyce multò longiora. Stam. 8-10 libera hypogyna basi squamulâ aucta Ovarium turbinatum supernè 4-5-lobum. Stylus 1 filiformis inter lobos ortus. Capsulâ 4-5 aut abortu 1-3, compressæ 1-locul. 1-spermæ. — Folia alterna! — An Banisteriæ, ut vult cl. Petit-Thouars, an Bala-nopteridi affinis? An, suadente cl. Jussieu et confirmante fructus amaritia Ochnaceis Simarubeisve referenda?

1. *N. TETRAPETALA* (Lam. ill. t. 299.) floribus 4-petalis 8-andris. ζ in Madagascar. Maudita penduliflora Comm! ined. Vitmannia elliptica Vahl symb. 3. p. 51. t. 62. Folia ovalia subcoriacea venosa. Pedunculi laterales nantes apice pedicellos 5-7 umbellatos 1-flores gerentes. (v. s.)
2. *N. PENTAPETALA* (Poir. dict. 4. p. 490.) floribus 5-petalis 5-andris? ζ in Indiâ orientali. — Karin njotti Rheed. mal. 6. t. 18. An prioris varietas? An verè 5-andra? Fructus intensè amarus.

Ordo XL. ACERINEÆ. DC. *théor. ed.* 2. p. 244. — *Acera.* Juss. *gen.* 50. *ann. mus.* 18. p. 477. *excl. sect.* 1 et 3. — *Acer.* Tourn. *inst.* 386. Linn. *gen. n.* 1115. Gaertn. *fr.* 2. p. 166. t. 116.

Cal. 5-rariùs 4-9-partitus. Petala totidem circà discum hypogynum inserta, lobis calycinis alterna, sæpiùs concolora, rariùs nulla. Stam. disco hypogyno inserta, sæpiùs 8 rariùs 5-12. Antheræ oblongæ. Ovarium didymum. Stylus 1. Stigmata 2. Fructus constans carpellis 2 (rarissimè 3) demùm separabilibus indehiscentibus samaroidis, nempè 1-locul. 1-2-spermis compressis superiùs desinentibus in alam membranaceam divergentem ad latus inferius incrassatam. Semina basi loculamentorum adfixa oblonga. Endoplevrum subcarnosum. Albumen nullum. Embryo curvatus convolutusve, cotyledonibus foliaceis irregulariter rugosis, radiculâ teretiusculâ ad basin loculi directâ.

Arbores. Folia opposita, sæpiùs simplicia ut in Malpighiaceis, rariùs composita ut in Hippocastaneis et Sapindaceis. Flores racemosi aut corymbosi axillares, sæpiùs abortu dioici aut polygami, interdùm apetalì.

Ordo omninò medius inter duos præcedentes et duos sequentes.

I. ACER. *Mœnch. method.* 334. et *Nutt. gen. am.* 1. p. 253.

Flores polygami. Calyx 5-lobus interdùm 5-partitus. Stam. rariùs 5, sæpiùs 7-9. — Folia simplicia.

§. 1. *Floribus racemosis.*

1. A. OBLONGUM (Wall! in litt.) foliis oblongis acuminatis integerrimis glabris, racemis compositis, alis fructus parallelis glabris separatis. 5 in Napauliâ. Folia subtus subglaucescentia. Fructus juniores in disco pubentes, alis glabris.
2. A. TATARICUM (Linn. sp. 1495.) foliis cordatis indivisis serratis lobis obsoletis, racemis compositis confertis erectis, fructuum alis parallelis junioribus subpuberulis. 5 in Tataria. — Pall. fl. ross. t. 3. Tratt. arch. 1. n. 1. ic. (v. v.)
3. A. STRIATUM (Lam. dict. 2. p. 381.) foliis cordatis trilobis acuminatis tenuiter et acutè serratis, racemis simplicibus pendulis, petalis ovalibus, fructibus glabris, alis subdivergentibus. 5 in Americâ bor. à Canadâ ad Carolinam. Truncus lineis albis eleganter striatus. Michx. f. arb. 2. t. 17. Petala ovalia. A. Pensylvanicum Linn. sp. 1496 Tratt. arch. 1. n. 11. ic. A. Canadense Duham. arb. 1. t. 12. Variat foliis indivisis, sed rariùs. (v. v.)
4. A. SPICATUM (Lam. dict. 2. p. 381.) foliis cordatis 3-5-lobis acuminatis subtus pubescentibus inæqualiter grossè serratis, racemis spiciformibus erectis, petalis linearibus, fructibus glabris, alis subdivergentibus. 5 in Canadâ et Alleghany mont. Flores minimi. — A. montanum Ait. h. kew. 3. p. 435. Tratt. arch. 1. n. 13. ic. A. pensylvanicum Duroi. harbk. t. 2. (v. s.)
5. A. PSEUDO-PLATANUS (Linn. sp. 1496.), foliis cordatis 5-lobis glabris, lobis

acuminatis inæqualiter dentatis, racemis pendulis subcompositis rachi staminumque filamentis hirsutis, fructibus glabris, alis subdivergentibus § in Europæ sylvis. Var. foliis variegatis. Duham. arb. 1. t. 36. Tratt. arch. 1. n. 2. ic. (v. v.)

β. *subobtusum*, foliorum lobis obtusioribus. A. opulifolium Thuill! fl. par. 538.

6. A. HYBRIDUM (Bosc dict. agr. 5. p. 251.) foliis subcordatis semicoriaceis glabris profunde trifidis, lobis inæqualiter grossè dentatis, racemis pendulis, fructibus glabris, alis divergentibus. § Patr. ign. (v. s.)

7. A. MACROPHYLLUM (Pursh am. 267.) foliis digitato-5-lobis, sinubus rotundatis, lobis subtrilobatis repando-dentatis subtus pubescentibus, racemis erectis, filamentis 9 hirsutis, ovariis hirsutissimis. § in Americâ bor. ad flum. Columbia et in Californiâ boreali.

8. A. CAMPESTRE (Linn. sp. 1497.) foliis cordatis 5-lobis, lobis grossè subdentatis, racemis erectis, fructuum alis divaricatissimis. § in Europâ et Tauriâ. Bieb. suppl. 643. An tres distinguendæ varietates, nempe:

α. *hebecarpum*, fructibus velutino-pubescentibus. A. campestre Vallr! in litt. Tratt. arch. 1. n. 7. ic.

β. *collinum*, (Wallr! in litt.) fructibus glabris, foliorum lobis obtusis, floribus minoribus. — in Galliâ.

γ. *Austriacum* (Tratt. arch. 1. n. 6. ic.) fructibus glabris, foliorum lobis subacuminatis, floribus majoribus. — in Austriâ, Podoliâ et Tauriâ.

§. 2. *Floribus corymbosis fasciculatisve.*

9. A. OPULUS. (Ait. h. kew. 3. p. 436.) foliis cordatis subrotundis 5-lobis, lobis obtusis obtusè grossèque dentatis, corymbis pedunculatis erectis, ovariis pilosis, fructibus glabris, alis subdivergentibus. § in Italiâ. A. Italum Lauth. ac. n. 8. A. rotundifolium Lam. dict. 3. p. 382. (v. v.)

10. A. OPULIFOLIUM (Vill. dauph. 4. p. 802.) foliis cordatis subrotundis 5-lobis, lobis obtusis obtusè grossèque dentatis, corymbis subsessilibus, ovariis fructibusque glabris, alis subdivergentibus. § in lapidosis apricis Valesiæ, Delphinatûs, Pedemontii, Cataloniæ. Tratt. arch. 1. n. 13. ic. A. Hispanicum Pourr. act. toul. 3. p. 305. A. vernum Reynier. A. montanum C. B. pin. 431. (v. v.)

11. A. OBTUSATUM (Kit. in Willd. sp. 4. p. 984.) foliis cordatis subrotundis 5-lobis, lobis obtusiusculis repando-dentatis subtus velutinis, corymbis pendulis, pedicellis pilosis, fructibus subpilosis, alis subdivergentibus. § in siccis montosis Hungariæ et Croatia. Tratt. arch. 1. n. 14. ic.

β. *Neapolinatum*. Ten! att. ac. neap. 1819, 1. p. 121. ic. — ad Neapolim (v. s.)

12. A. OBTUSIFOLIUM (Sibth. et Smith fl. græc. t. 361. ex prod. 1. p. 263.) foliis rotundatis obtusè trilobis denticulato-crenatis longitudine ferè petiolorum. § in montibus Sphaciolicis Cretæ. A. Cretica Tourn. cor. 43.

13. A. CRETICUM (Linn. sp. 1497.) foliis perennantibus basi cuneatis acutè trilobis, lobis integris denticulatisve, lateralibus brevioribus, corymbis paucifloris erectis, fructibus glabris, alis vix divergentibus. § in montibus Cretæ et in Archipelago Græciæ. — Tratt. arch. 1. n. 19. ic. Ait. h. kew. ed. 2. v. 5. p. 418. An var. sequentis quocum cl. Lamarck junxit sub A. trilobati nomine? (v. v.)

14. A. MONSPRESSULANUM (Linn. sp. 1497.) foliis annuis basi cordatis trilobis, lobis subintegerrimis æqualibus, corymbis paucifloris erectis, fructibus glabris, alis vix divergentibus. Vulgò §, interdum 5, imò § in apricis lapidosis Europæ mediæ et australis. Tratt. arch. 1. n. 20. ic. A. trifolia Duham. arb. 1. t. 10. f. 8. (v. v.)

α5. A. IBERICUM (Bieb. fl. taur. 2. p. 447.) foliis nitidis subtus glaucis obtusè trilobis, lobis 1-2-dentatis, lateralibus nervo medio ad petioli insertionem nudo notatis, petiolis folio paulò brevioribus. § in Iberiâ.

16. *A. HETEROPHYLLUM* (Willd. arb. 10. t. 1. f. 1.) foliis sempervirentibus ovatis integris trilobisque obsolete serratis glabris. § in Oriente. *A. sempervirens*. Linn. mant. 128.
17. *A. BARBATUM* (Michx. fl. am. 2. p. 252.) foliis ovatis subcordatis breviter 3-lobis inæqualiter serratis subtus glaucis, corymbis sessilibus, femineorum pedicellis simplicibus, masculorum ramosis, calycibus intus barbatis, fructibus glabris, alis vix divergentibus. 5 à Novâ Cæsareâ ad Carolinam in pinetis humidis. *A. Carolinianum*, Walt. car. 251.
18. *A. PLATANOIDES* (Linn. sp. 1496.) foliis cordatis glabris 5-lobis, lobis acuminatis, dentibus paucis grossis acutis, corymbis pedunculatis erectiusculis fructibusque glabris, pedicellis divaricatis. 5 in Europæ sylvaticis. Duham. arb. 1. t. 10. f. 1. Tratt. arch. 1. t. 4. Petioli abrupti lactescunt! Gemmarum squamæ patent-reflexæ. (v. v.)
 β. *laciniatum*, (Ait. h. kew. 3. p. 435.) foliorum lobis inciso-laceris, dentibus acuminato-setaceis. *A. crispum* Lanth. acer. n. 4. — Tratt. arch. 1. n. 5. ic. (v. v.)
19. *A. SACCHARINUM* (Linn. sp. 1496.) foliis cordatis glabris subtus glaucis palmato-5-lobis, lobis acuminatis sinuato-dentatis, corymbis breviter pedunculatis nutantibus, pedicellis pilosis, fructibus glabris, alis divergentibus. 5 in vallibus à Canadâ ad Pensylvaniam. Michx. f. arb. 2. t. 15. Tratt. arch. 1. n. 3. ic. Accedit ad *A. platanoidem*. (v. s.)
20. *A. NIGRUM* (Michx. f. arb. 2. p. 238. t. 16.) foliis sinu clauso cordatis subtus pubescentibus, corymbis sessilibus nutantibus, fructibus glabris basi turgidis, alis divergentibus. § in montosis à Noveboraco ad Carolinam. *Saccharinum* succum ut præcedens fundit.
21. *A. ERIOCARPUM* (Michx. fl. am. 2. p. 253.) foliis basi truncatis subtus glabris glaucisque palmato-5 lobis, sinubus obtusis, lobis inæqualiter inciso-dentatis, floribus conglomeratis breviter pedicellatis apetalis pentandris ovarii tomentosis. 5 à Novâ-Angliâ ad Georgiam. — Desfl. ann. mus. 7. p. 412. t. 25. *A. dasycarpum* Willd. sp. 4. p. 985. Tratt. arch. 1. n. 8. ic. *A. tomentosum* Hort. Par. (v. v.)
22. *A. RUBRUM* (Linn. sp. 1496.) foliis basi cordatis subtus glaucis inæqualiter inciso-dentatis palmato-5-lobis, sinubus acutis, floribus conglomeratis pentapetalis pentandris, ovarii glabris. 5 à Canadâ ad Floridam. Michx. arb. 2. t. 14. Desfl. ann. mus. 7. p. 413. t. 25. Tratt. arch. 1. n. 9. ic. (v. v.)
- §. III. *Umbellis pedunculatis*. Species Japonicæ non satis notæ.
23. *A. DISSECTUM* (Thunb. fl. jap. p. 160.) foliis 5-7-partitis, lobis oblongis acuminatis inciso-serratis pinnatifidisve, umbellis 4-6-floris. 5 in Japoniâ. — Tratt. 1. n. 18. ic. Variat fl. rubris.
24. *A. JAPONICUM* (Thunb. l. c. p. 161.) foliis subrotundis villosis palmatim multifidis, lobis 11-13 acuminatis serratis, umbellis multifloris. 5 in Japoniâ. Tratt. arch. 1. n. 16. ic.
25. *A. PALMATUM* (Thunb. l. c. p.) foliis glabris palmatim ultra medium 5-7-fidis, lobis oblongis acuminatis serratis, umbellis 5-7-floris. 5 in Japoniâ. Tratt. arch. 1. n. 17. ic.
26. *A. SEPTEMLOBUM* (Thunb. l. c. p. 162.) foliis glabris 7-lobis, lobis acuminatis æqualiter acutè serratis. 5 in Japoniâ.
27. *A. PICTUM* (Thunb. l. c. p. 162.) foliis glabris palmatim 7-lobis, lobis acuminatis integris. 5 in Japoniâ. Rami cinerei. Folia maculata. — Tratt. arch. 1. n. 15. ic.
28. *A. TRIFIDUM* (Thunb. l. c. p. 163.) foliis indivisis trifidisque integris. 5 in Japoniâ.

†. *Species non satis notæ.* *

29. *A. CIRCINNATUM* (Pursh. fl. am. 1. p. 267.) foliis orbiculatis basi subcor-

dati 7-lobis inæqualiter acutè dentatis utrinquè glabris, nervis venisque ad axillas pilosis. 5 in Americâ bor. ad flumen Columbia et in Californiâ boreali.

II. NEGUNDO. *Mæench. meth.* 334. — *Negundium Rafn.*

Flores dioïci. Cal. minimus inæqualiter 4 5 dentatus. Pet. o. — ♂ Flores fasciculati, pedicellis filiformibus. Antheræ 4-5 lineares sessiles. — ♀ Flores racemosi. — Folia composita impari-pinnata.

1. *N. FRAXINIFOLIUM* (Nutt. gen. am. 1. p. 253.) foliis 3 5-foliolatis, foliolis oppositis grossè rariterque dentatis, impari sæpius trilobo. 5 in Americâ bor. à Pennsylvania ad Carolinam. *Acer Negundo* Lin. sp. 1497. Michx. fil. arb. 2. t. 16. Tratt. arch. 1. n. 10. ic. *N. aceroides* Mæench. l. c. (v. v.)
2. *N. MEXICANUM*, foliis omnibus trifoliolatis. 5 in Mexico. *Acer ternatum* fl. mex. ic. ined. An prioris varietas?
3. *N? COCHINCHINENSE*, foliis sub-4-jugis, foliolis alternis integerrimis. 5 in sylvis Cochinchinæ. *Acer pinnatum* Lour. coch. 2. p. 797. — Petala 5 alba. Carpellorum alæ carnosæ. An potius Sapindacearum genus?

Ordo XLI. HIPPOCASTANÆ. *DC. theor. ed. 2. p. 244. — Castaneaceæ. Link. enum. 1. p. 354. — Æsculus. Linn. gen. n. 462.*

Cal. campanulatus 5-lobus. Petala 5 aut uno deficiente 4, inæqualia, hypogyna. Stam. 7-8 disco hypogyno inserta, libera, inæqualia. Antheræ subincumbentes. Ovarium subrotundo-trigonum. Stylus 1 filiformi-conicus acutus. Capsula junior 3-locul. 3-valvis, loculis biovulatis, valvis medio septiferis, ovulis septis adfixis; adulta coriacea subglobosa abortu 2-3-valvis, 2-3-locularis, 2-4-sperma. Semina (ferè Castanæ) grossa, subglobosa, variè compressa et angulata, testâ nitidâ glaberrimâ ferrugineâ, hilo lato fusco-cinereo basilari. Albumen nullum. Embryo curvatus inversus, cotyledonibus carnis crassissimis gibbis conferruminatis per germinationem intrâ spermodermim hypogæis, plumulâ grandiusculâ diphyllâ, radiculâ conicâ curvatâ ad hilum directâ, sed ob seminum abortus quoad fructum situ variâ. — Arbores fruticesve. Folia opposita composita quinato-septenatove palmata, foliolis penninerviis. Racemi terminales subpaniculati, pedicellis articulatis.

I. ÆSCULUS. *Linn. gen. n. 462. excl. sp. — Hippocastanum. Tourn. inst. t. 612.*

Calyx campanulatus. Petala 4-5 expansa, limbo ovato. Stam. filamenta introrsum recurva. Capsulæ echinatæ. Foliola sessilia aut subsessilia.

1. *Æ. HIPPOCASTANUM* (Linn. sp. 488.) capsulis echinatis, floribus 5-petalis 7-andris, foliolis 7 obovato-cuneatis acutis dentatis. ♀ in Indiâ boreali. Petala alba rubro et luteo maculata. — Riv. pentap. irr. t. 123. — H. vulgare Gaertn. fr. 2. t. III. (v. v.)
β. flore pleno.
2. *Æ. RUBICUNDA* (Herb. amat. t. 367.) capsulis echinatis, floribus 4-petalis 8-andris, unguibus calyce brevioribus, foliolis 5-7 obovato-cuneatis acutis inæqualiter biserratis. ♀ Patr. ign. *Æ. carnea* hortul. Flores coccinei pulcherrimi. Species priori affinis sed minor. (v. v.)
3. *Æ. GLABRA* (Willd. enum. 405.) capsulis echinatis, corollis 4-petalis patulis, unguibus longitudine calycis, staminibus corollâ longioribus, foliolis 5 glaberrimis. ♀ in Americâ bor. à Pensylv. ad Virginiam. Cor. viridiflavæ. (Linn.)
4. *Æ. OHIOENSIS* (Michx. arb. 3. p. 242.) capsulis echinatis, corollis., foliolis 5 glabris ovali-acuminatis irregulariter dentatis. ♀ in Americâ bor. ad ripas fl. Ohio. Flores albi numerosi racemosi. Fructus dimidiò minores quàm in *Æ. Hippocastano*. †.
5. *Æ. PALLIDA* (Willd. enum. 406.) capsulis echinatis, corollis 4-petalis patulis, unguibus calyce brevioribus, staminibus corollâ duplò longioribus, foliolis 5. ♀ in Americæ bor. sylvis Kentuckyensibus.

II. PAVIA. *Boerh. lugd. b. t. 260.*

Calyx tubulosus. Petala 4 erecta angusta. Stam. recta. Capsulae inermes. — Foliola petiolulata.

1. *P. MACROSTACHYA* (Herb. amat. t. 212.) capsulis inermibus, staminibus corollâ 4-petalâ multò longioribus, racemo longissimo, foliis 5. $\frac{5}{3}$ in Americâ bor. ad ripas fluminum Georgiæ. *Æsculus macrostachya* Michx. fl. bor. am. 1. p. 220. Jacq. ecl. 1. t. 9. *Æ. parviflora* Walt. car. 128. *Pavia alba* Poir. dict. 5. p. 95. *Pavia edulis* Poit. arb. fr. t. 88. — Flores albi. Frutex radice stoloniferâ. (v. v.)
 2. *P. RUBRA* (Lam. ill. t. 273.) capsulis inermibus, staminibus corollâ 4-petalâ brevioribus, foliis 5-elliptico-oblongis utrinquè acutis petiolisque glabris, axillis nervorum subtùs pilosis. $\frac{5}{3}$, et in *Æsculo Hippocastano* inserta $\frac{5}{3}$. in Americâ bor. *Æsculus Pavia* Lin. sp. 488. — Duham. arb. 2. t. 19. Flores sordidè rubri. — Confer an stirps *Brasiliensis* sit eadem? (v. v.)
 3. *P. HYBRIDA*, capsulis inermibus, staminibus corollâ tetrapetalâ brevioribus, petiolis glabris, foliis 5 elliptico-oblongis utrinquè acuminatis subtùs pubescenti-velutinis. $\frac{5}{3}$ in Americâ bor. ad Georgiam occid. *Æsculus hybrida* DC. h. monsp. 1813. p. 75. *Æ. discolor* Pursh fl. bor. am. 1. p. 255. bot. reg. t. 310. Flores ex albo et rubro sordidè variegati. Media inter priorem et sequentem. (v. v.)
 4. *P. FLAVA*, capsulis inermibus, staminibus corollâ 4-petalâ brevioribus, petiolis pubescentibus supernè planiusculis, foliis 5-7 elliptico-oblongis utrinquè acutis subtùs et in nervis supernè pubescentibus. $\frac{5}{3}$ in Virginîæ, Carolinæ et Kentucky montibus. *Æsculus flava* Ait. h. kew. 1. p. 494. *Æ. lutea*. Wang. act. nat. scr. ber. 8. p. 133. t. 6. *Pavia lutea* Poir. dict. 5. p. 94. — Flores pallidè flavi. (v. v.)
-

Ordo XLII. RHIZOBOLÆ.

Calyx constans sepalis 5 plus minus coalitis et ideò vulgò dicendus 1-phyllus 5-dentatus 5-partitusvè. Petala 5 crassiuscula sepalis alterna, disco hypogyno cum staminibus inserta et eorum tubo basi adnata, inter se subinæqualia. Stam. numerosissima duplici ordine è disco prodeuntia interiora sæpè breviora et castrata, filamentis basi breviter monadelphis, ceterum filiformi-subulatis, antheris subrotundis. Ovarium liberum subglobosum subtetragonum 4-loculare 4-spermum. Styli 4. Stigmata simplicia. Fructus constans nucibus 4 adglutinis, sæpius nonnullarum abortu 1-3 tantum. Nux quaque rotundata indehiscens 1-locularis, putamine crasso membranis 2 constante, exteriore membranaceâ setas rigidas ex interiore lignosâ ortas carne carnosâ oleosâ intermixtas obvelante. Funiculus dilatatus in carunculam spongiosam bilobam. Semen reniforme dorso carinatum utroque apice attenuatum exalbuminosum. Embryo radiculâ maximâ adscendente totam nucleï substantiam amygdaloideam efficiente, cauliculo (Tigellâ Rich.) ancipiti longiusculo apice cotyledones 2 foliaceas parvas ovato-lanceolatas gerente, in sulco radiculæ superficiali aut, obs. cl. Richard, intrâ radiculæ canalem (in *C. butyroso*) locato.

Arbores foliis oppositis petiolatis palmatim compositis 3-5-foliolatis, floribus racemosis ebracteatis.

Ordo distinctissimus, hinc ad Therebinthinaceas et speciatim ad Mangiferam accedens, sed ob flores hypogynos et fructum potius Sapindaceis forsân consociandus, paululùm etiam Hippocastaneis ob folia opposita palmato-composita affinis, in Hippocastaneis autem radícula parva et cotyledones maximæ, in Rhizoboleis radícula maxima et cotyledones parvæ, in utroque ordine albuminis substantia à variis embryonis partibus quasi hausta.

I. CARYOCAR. *Lin. mant.* 247. — *Rhizobolus Gaertn. fr.* 2. p. 93. — *Pekoa et Saouari Aubl. guian.* 1. p. 594 et 599.

Cal. 5-lobus. Pet. 5. Stam. ∞ basi monadelpha. Styli 4. Drupa nucleis 4 aut abortu 1-3 reticulatis.

§. 1. *Foliis trifoliolatis Saouari Aubl.*

1. *C. NUCIFERUM* (*Lin. mant.* 247.) foliis trifoliolatis, calyce corollâque purpureis. antheris oblongis, drupis capitis humani magnitudine. § in Berbices et Essequibo Americæ meridionalis.

2. *C. GLABRUM* (*Pers. ench.* 2. p. 84.) foliis trifoliolatis glabris, foliolis ovatis

acuminatis subdentatis, drupis ovi gallini magnitudine. § in Guianæ sylvis. Saouari glabra. Aubl. guian. 2. p. 599. t. 240. Rhizobolus Saouari Cor. ann. mus. 8. p. 394. t. 5. f. 2.

3. *C. VILLOSUM* (Pera. ench. 2. p. 84.) foliis trifoliatis, foliis ovatis subrotundis subacutis subtus tomentosis. § in Guianæ sylvis. Saouari villosa. Aubl. Flores et Fructus ignoti.

4. *C. AMYGDALIFERUM* (Cav. ic. 4. p. 37. t. 361 et 362.) foliis trifoliolatis glabris, foliolis lanceolatis serratis in nervorum axillis subtus fasciculo pilorum notatis. antheris subrotundis. § in sylvis Mariquitæ Santa-Fe de Bogota. Arbor 180-240 pedes alta!

§. 2. *Foliis 5-foliolatis. Pekea Aubl.*

5. *C. BUTYROSUM* (Willd. sp. 2. p. 1243.) foliis 5-foliolatis glabris, foliis integris ovali-lanceolatis acuminatis, drupis lævibus. § in Guianæ sylvis. Pekea butyrosa. Aubl. guian. 1. p. 594. t. 238. Lam. ill. t. 486.

6. *C. TOMENTOSUM* (Willd. sp. 2. p. 1244.) foliis 5-foliolatis subtus tomentosis, foliolis integris ovalibus acuminatis, drupis obtusè tuberculosi. § in Guianæ sylvis. Pekea tuberculosa. Aubl. guian. 1. p. 397. t. 239. Rhizobolus Pekea Gaertn. fr. 2. p. 93. t. 98. f. 1. (v. fr.)
-

Ordo XLIII. SAPINDACEÆ. *Juss. ann. 18. p. 476.* — *Sapindi Juss. gen. 246.*

Cal. Sepala 4-5 nunc libera nunc basibus subcoalita æqualia, æstivatione imbricatâ? Petala sæpius tot quot sepala, interdum 1 deficiente, interdum nulla, hypogyna, nunc nuda, nunc intus medio villosa aut glandulosa, nunc laminâ petaloïdeâ intus aucta. Annulus glandulosus inter petala et stamina. (an in omnibus?) Stamina petalorum numero dupla, disco hypogyna inserta, filamentis distinctis. Ovarium subrotundum. Stylus 1 seu 3. Fructus drupaceus capsularisve, 3-locularis aut abortu 1-2-locularis. Semina in loculis solitaria, angulo interiori adfixa, exalbuminosa. Embryonis radícula ad loculi fundum versa, cotyledonibus 2 plus minùs suprâ radiculam replicatis, interdum rectis.

Arbores, Frutices erecti aut scandentes, rariùs herbæ scandentes. Folia alterna sæpius composita et Therebinthacearum folia in sicco præsertim simulantia.

Trib. I. *PAULLINIEÆ. H. B. et Kunth nov. gen. am. 5. p. 99.*

Petala intus ad unguem laminâ petaloïdeâ appendiculata. Glandulæ distinctæ inter petala et stamina. Ovarium 3-loculare loculis 1-spermis. — Herbæ aut frutices scandentes et cirrhosæ.

I. *CARDIOSPERMUM. Lin. gen. n. 498. — Corindum. Tourn. inst. t. 246.*

Cal. 4-sepalus, sep. 2 minoribus. Petala 4 intus appendiculata, appendicibus inæqualibus. Glandulæ duæ hypogynæ inter petala et stamina. Stam. 8. Styli 3. Carpella 3 inflata membranacea ad axim connata dorsò alata evalvia. Semina globosa hilo lato cordato. Cotyl. crassæ inæquales (Gaertn. fr. 1. p. 381. t. 79.) Caules herbacei subscandentes. Pedunculi apice racemosi, pedicellis infer. 2 abortivis cirrhosis.

*. *Glandulis 2 hipogynis abbreviatis rotundatis.*

1. *C. HALICACABUM* (Lin. spec. 925.) caule petiolis foliisque glabris, foliis bitermatim sectis, segmentis petiolatis inciso-dentatis. (1) in Indiâ orient. et forsân in insul. Caribæis. — *Rumph. amb. 6. t. 24. f. 2. Lam. ill. t. 317. (v. v.)*
2. *C. MICROCARPUM* (H. B. et Kunth nov. gen. am. 5. p. 104.) ramis glabris, foliis bitermatim sectis, segmentis serrato-incisis utrinquè adpresso-puberulis, capsulis subtrbinatis pubescenti-hirtellis. 2 in humidis Orinocensis provinciæ.
3. *C. MOLLE* (H. B. et Kunth l. c.) ramis villosis, foliis suprâ decompositis, segmentis argutè serrato-incisis trifidis utrinquè adpresso pilosis subtus canescentibus, capsulis subglobosis molliter hirtis. 2 prope Guanaxuato Mexicanorum.
4. *C. LOXENSE* (H. B. et Kunth l. c.) ramis albo-lanatis, foliis subternatim sectis, segmentis grossè crenato-serratis, suprâ adpresso-pilosis, subtus piloso-sericeis et canescentibus, infimis confluentibus, capsulis subglobosis pubescenti-hirtellis. 2 prope Loxam Peruvianorum. (v. s. sine fr.)

** *Glandulis 2 hypogynis elongatis linearibus.*

5. *C. GRANDIFLORUM* (Swartz fl. ind. occ. 2. p. 698.) caule basi frutescente, petiolis foliisque pubescentibus, foliis triternatim sectis, capsulis acuminatis maximis tomentosis. ζ in fruticetis sepibusque Jamaicae. (v. s.)
6. *C. ELEGANS* (H. B. et Kunth nov. gen. am. 5. p. 99. t. 439.) caule basi frutescente, ramis puberulis, foliis biternatim sectis, segmentis grossè serratis glabris, capsulis elliptico-subglobosis glabris. ζ in regno Peruviano.
7. *C. COLUTROIDES* (H. B. et Kunth nov. gen. am. 5. p. 100.) foliis biternatim sectis, segmentis grossè crenato-incisis suprâ puberulis subtus ramisque molliter pubescentibus capsulis elliptico-oblongis glabris. ζ prope Caracas in montosis. *C. vesicarium* Humb. rel. hist. 1. p. 39.
8. *C. MACROPHYLLUM* (H. B. et Kunth l. c.) foliis biternatim sectis, segmentis argutè serrato-incisis, subtus ramisque molliter pubescentibus et canescentibus. ζ prope Atures in prov. Orinocensi. Fruct. ign.
9. *C. HISPIDUM* (H. B. et Kunth l. c.) ramis petiolisque hirsuto-hispidis, foliis biternatim sectis, segmentis grossè serrato-crenatis glabris. ζ in ripâ fl. Amazonum, prope Tomependa. Fruct. ign.
10. *C. HIRSUTUM* (Willd. spec. 3. p. 467.) caule petiolisque hirsutis, foliis biternatim sectis, segmentis glabris petiolatis ovatis acuminatis dentatis, axillis venarum subtus pubescentibus. $\textcircled{1}$? in Guineâ.

†. *Species non satis notæ.*

11. *C. CORINDUM* (Lin. spec. 526.) foliis subtus tomentosis biternatim sectis, segmentis subpetiolatis incisis obtusis. $\textcircled{1}$ in Brasiliâ. *C. villosum* Mill. dict. n. 3.
12. *C. PUBESCENS* (Lag. gen. et spec. 14.) frutescens, totum pubescens, capsulis obtusis. ζ in Novâ-Hispaniâ. Corolla sanguinea duplò major quàm in *C. Halicacabo*.

II. *URVILLÆA*. *H. B. et Kunth nov. gen. am. 5. p. 105. t. 440.*

Cal. 5-sepalus persistens, sep. 2 exter. minoribus. Pet. 4 unguiculata, 2 magis distantia. Stam. 8 filam. liberis. Ovarium substipitatum, 3-locul., ovulis solitariis adscendentibus. Glandulæ 2 maculæformes ad basin stipitis. Capsula membranacea 3-alata carpellis 3 samaroides axi adnatis constans. — Frutex sarmentosus, foliis ternatis, racemis basi cirrhis axillaribus spicæformibus, floribus albidis.

1. *U. ULMACEA* (H. B. et Kunth l. c.) foliolis ovatis acuminatis regulariter serratis subtus petiolis ramisque pubescentibus. ζ in montanis prope Caracas. *Kœlreuteria triphylla* Pers. ench. 1. p. 414.
2. *U. BERTERIANA*, foliolis ovatis grossè et irregulariter dentatis, medio basi angustè cuneato, subtus petiolis ramisque glabresculis. ζ in Sanctâ-Marthâ, Bertero. *Paullinia Berteriana* Balb! in litt. (v. s.)

III. *SERJANIA*. *Plum. gen. 34. t. 35.* — *Seriana*. *Schum. act. soc. nat. hafn. 3. part. 2.*

Cal. 5-sepalus. Pet. 4. intus appendiculata, quinti deficientis sede vacuâ. Stam. 8. Carpella 3 (samaræ) longitudinaliter connexa, infernè in alam membranaceam dilatata, supernè 1-sperma. — Frutices scandentes, foliis ternato-compositis.

* *Foliis ternatis.*

1. *S. SINUATA* (Schum. l. c. p. 126. t. 12. f. 1.) foliis ternatis, foliolis ovato-lanceolatis dentato-sinuatis, carpellorum alis posticè dilatatis. ζ in insul. Caribæis. — *Plum. ic. ed. Burm. t. 113. f. 2. Paullinia seriana* Lin. sp. 524. (v. s.)

2. *S. MOLLIS* (H. B. et Kunth nov. gen. am. 5. p. 108.) foliis ternatis, foliolis ovatis crenato-serratis basi subrotundis suprâ sericeo-pilosis subtus sericeo-tomentosis, terminali 3-lobo, fructibus hirsutis. ζ in Andibus Peruv. inter Querocotillo et Montañ.
 3. *S. ACAPULCENSIS* (H. B. et Kunth l. c.) foliis ternatis, foliolis oblongis basi rotundatis subauriculatis lateralibus apice obsolete crenatis. ζ in Mexico inter Acapulco et la Venta del Exido. An var. sequentis. Fruct. ign.
 4. *S. EMARGINATA* (H. B. et Kunth l. c.) foliis subbinternatis, foliolis integerrimis glabris apice rotundatis et emarginatis, lateralibus subauriculatis, intermedio trilobo aut trisecto, fructibus glabris. ζ in montibus Mexicanis.
- ** *Foliis biternatis.*
5. *S. DIVARICATA* (Schum. l. c. p. 126. t. 12. f. 2.) foliis biternatis, foliolis ovatis integerrimis subacutis lucidis petiolatis, petiolis nudis, carpellorum alis posticè dilatato-rotundatis. ζ in Jamaicae sylvis. Paullinia divaricata Swartz! fl. ind. 2. p. 696. (v. s.)
 6. *S. CARACASANA* (Willd. spec. 3. p. 465.) foliis biternatis, foliolis oblongis utrinquè acutis remotissime dentatis glaberrimis, petiolis nudis, carpellorum alis posticè rotundatis. ζ ad Caracas et in Guadaloupâ. Paullinia Caracasana Jacq. Schœnbr. 1. p. 52. t. 99. (v. s. sine fl.)
 7. *S. GLABRATA* (H. B. et Kunth nov. gen. am. 5. p. 110.) foliis biternatis glabris, foliolis elliptico-oblongis acuminatis basi cuneatis grossè serrato-crenatis, petiolis nudis, carpellorum alis posticè rotundatis. ζ in temperatis Peruviae. Priori valdè affinis.
 8. *S. RACEMOSA* (Schum. l. c. p. 127. t. 12. f. 5.) foliis biternatis, foliolis ovatis acutis profundè serratis, petiolis subnudis, carpellorum alis posticè dilatatis anticè subsinnatis. ζ in Verâ-Cruce.
 9. *S. SPECTABILIS* (Schum. l. c. p. 127. t. 12. f. 4.) foliis biternatis, foliolis obovatis integris terminalibus retuso-emarginatis, petiolis alatis, carpellorum alis posticè dilatatis. ζ in Americâ.
 10. *S. PAUCIDENTATA*, foliis biternatis, foliolis ovalibus utrinquè attenuatis ad apicem dentè 1-2-grosso utrinquè notatis, petiolis subulatis, carpellorum alis posticè rotundato-dilatatis. ζ in Cayennâ aut Guianâ-Galicâ.
 11. *S. OSSANA*, foliis biternatis glabris supernè atro-virentibus subnitidis subtus pallidis, foliolis ovatis apice subdentatis obtusis mucronulatis, petiolis nudis, carpellorum alis posticè obtusis. ζ ex Havânâ misit cl. de la Ossa. (v. s.)
 12. *S. PUBESCENS* (H. B. et Kunth l. c.) ramis villosis pubescentibus, foliis biternatis, foliolis ovato-oblongis obtusis submucronatis integerrimis aut apice grossè crenatis suprâ glabriusculis subtus pubescentibus. ζ in prov. Caracasanae convallibus Araguensium.
 13. *S. MEXICANA* (Willd. spec. 3. p. 465.) foliis biternatis, foliolis obovatis integris dentatisve sæpè emarginatis, petiolis subulatis, caule aculeato, racemis compositis, carpellorum alis rotundato-subdilatatis. ζ in Mexico. Paullinia Mexicana Linn. spec. 525. Schum. l. c. p. 124. t. 11. f. 3. fl. mex. icon. ined. ex quâ caract. emend.
 14. *S. ANGUSTIFOLIA* (Willd. spec. 3. p. 466.) foliis biternatis, foliolis lineari-lanceolatis integerrimis acutis, petiolis alatis. ζ in Americâ calid. — Plum. ic. ed. Burm. t. 113. f. 1. Paullinia Mexicana Jacq. obs. 3. p. 11. t. 61. f. 5. folium.
 15. *S. PANICULATA* (H. B. et Kunth nov. gen. am. 5. p. 111. t. 441.) ramis pubescentibus, foliis biternatis, foliolis ovato-ellipticis acuminatis grossè crenatis glabris, subtus axillis venarum barbatis, racemis paniculatis, fructibus apice pubescentibus. ζ in calidis prov. Caracasanae. Affinis *S. Caracasanae*.

16. *S. LUPULINA* (Schum. l. c. p. 127. t. 12. f. 5.) foliis biternatis, foliolis crenatis subtus ferrugineis terminalibus subrhombicis, lateralibus ovatis, petiolis subulatis, carpellorum alis semi-ovalibus. 5 in Americâ calid.
17. *S. LUCIDA* (Schum. l. c. p. 128.) foliis biternatis, foliolis ovatis acutis serratis, petiolis subnudis, carpellorum alis semi ovalibus. 5 in ins. Sanctæ-Crucis. P (v. s.)
18. *S. OXYPHYLLA* (H. B. et Kunth l. c.) ramis glabris, foliis biternatis, foliolis ovato-oblongis acuminato-mucronatis remotè serratis glabris, summo trifido aut trisetis, fructibus glabriusculis. 5 in Andibus Peruvianorum.

*** *Foliis ternatis.*

19. *S. TRITERNATA* (Willd. sp. 3. p. 466.) foliis triternatis, foliolis ovatis obtusis repandis, petiolis alatis, racemis paniculatis, carpellorum alis rotundato-dilatatis. 5 in Americâ calid. — Plum. ed. Burm. t. 112. Paullinia triternata Lin. mant. 236. P. polyphylla Lin. sp. 525. Jacq. obs. 3. t. 61. f. 10? (v. s.)

**** *Foliis impari-pinnatis.*

20. *S. PARVIFLORA* (H. B. et Kunth nov. gen. am. 5. p. 113.) foliis impari-pinnatis, foliolis trijugis, jugo infimo utrinquè trifoliolato, ovato-oblongis crenato-serratis apice acutiusculo-mucronatis, suprâ glabris, subtus hirtello-puberulis, fructibus glabris. 5 in ripâ fluminis Amazonum.
21. *S. HETEROPHYLLA*, foliis inferioribus impari-pinnatis bijugis, superioribus biternatis, foliolis ovatis subdentatis glabris, terminalibus basi cuneatis, petiolis apice subulatis, racemis pedunculatis, carpellorum alis posticè rotundatis. 5 in Sanctâ-Marthâ. Bertero. (v. s. in h. Balbis.)

IV. *PAULLINIA*. Schum. act. soc. h. nat. hafn. 3. part. 2. Juss. ann. mus. 4. p. 340. — Cururu. Plum. gen. 34. t. 35.

Cal. 5-sepalus subinæqualis. Pet. 4 intus appendiculata, quinti deficientis sede vacuâ. Stam. 8 subinæqualia. Capsula coriacea 3-locularis, 3 valvis. Semina arillo membranaceo semitecta. — Frutices scandentes, foliis compositis.

* *Capsulâ pyriformi, valvis apteris, foliis ternatis.*

1. *P. TURBACENSIS* (H. B. et Kunth nov. gen. am. 5. p. 114.) capsulis pyriformibus densissimè hirtellis, foliis ternatis, foliolis acuminatis grossè serrato-dentatis glabris nitidis, petiolo nudo. 5 in temperatis regni Novo-Grenatensium ad Turbaco.
2. *P. CURURU* (Linn. spec. ed. 1. p. 365.) capsulis pyriformibus obtusis, foliis ternatis, foliolis oblongis dentato-serratis subacuminatis, petiolo alato. 5 in insul. Antillanis. — Plum. ed. Burm. t. 111. f. 2.
3. *P. NITIDA* (H. B. et Kunth nov. gen. am. 5. p. 115.) capsulis pyriformibus, foliis ternatis, foliolis oblongis subacuminatis grossè serrato-crenatis glabris nitidissimis, petiolo nudo. 5 in ripâ fluminis Orinoci.
4. *P. RIPARIA* (H. B. et Kunth l. c.) capsulis pyriformibus, foliis ternatis, foliolis oblongis acutis grossè serratis nitidis glabris, petiolo alato, racemis folium subæquantibus. 5 in ripâ fluminis Magdaleuz.

** *Capsulâ pyriformi apterâ, foliis pinnatis.*

5. *P. PINNATA* (Linn. spec. ed. 1. p. 366.) capsulis pyriformibus, junioribus apice 3 tuberculatis, foliis pinnatis bijugis cum impari, foliolis ovato-lanceolatis sessilibus crenatis, petiolo alato. 5 in Brasiliâ, Mexico, Guianâ, Antillis et Africâ æquinoctiali. — Plum. ed. Burm. 76. t. 91. (v. s.)
6. *P. MACROPHYLLA* (H. B. et Kunth l. c.) foliis impari-pinnatis, foliolis bijugis oblongis obtusis remotè et obsolete dentatis membranaceis glabris, nervisque puberulis, petiolo rachique alatis. 5 in ripâ fluminis Magdaleuz.

*** *Capsulâ pyriformi, valvis carinatis alatis.*

7. *P. TETRAGONA* (Aubl. guian. 1. p. 355.) capsulis tricarinatis 3-cornibus, foliis pinnatis, foliolis ovatis subserratis acutis glabris, petiolo nudo, caule tetragono. ζ in Cayennâ. *P. pinnatæ* affinis.
8. *P. VESPERTILIO* (Swartz. fl. ind. occ. 2. p. 695.) capsulis apice alatis, alis horizontalibus, foliis pinnatis, foliolis ovatis remotè crenatis glabris, petiolis nudis. ζ in ins. Sancti-Christophori. Schum. act. soc. h. n. hafn. 3. p. 122. t. 11. f. 1.
9. *P. ACUTANGULA* (Pers. ench. 1. p. 443.) capsulis obovatis triangulo-alatis, foliis pinnatis, foliolis oblongo-ovatis acuminatis dentato serratis. ζ in Peruvizæ nemoribus. Semarillariæ acutangula. fl. per. syst. p. 93. (v. s.)
10. *P. MELIÆFOLIA* (Juss. ann. 4. p. 347. t. 66. f. 2.) capsulis pyriformibus apice 3-alatis, alis parvis in stylum coeuntibus, foliis pinnatis trijugis cum impari, foliolis subsessilibus lanceolatis remotè dentatis subtus pubescentibus. ζ in Brasiliâ.
11. *P. VELUTINA*, tota tomentoso-velutina, capsulis 3-alatis basi breviter attenuatis, foliis biternatis, foliolis ovato-lanceolatis utrinquè acuminatis grossè dentatis petiolis nudis. ζ in prov. Sanctæ-Marthæ. Bertero. (v. s.)
12. *P. THALICTRIFOLIA* (Juss. an. 4. p. 347. t. 66. f. 1.) capsulis à basi 3-alatis, alis apice latioribus divaricatis, foliis subtripinnatis, summis simplicibus, foliolis ovatis parvis integris aut subdentatis. ζ in Brasiliâ et Sancto-Domingo. (v. s. sine fl.)

**** *Capsulâ ovatâ.*

13. *P. OBOVATA* (Pers. ench. 1. p. 443.) capsulis obovatis, foliis pinnatis bijugis cum impari, foliolis lanceolatis serratis. ζ in Peruvizæ nemoribus. Semarillaria obovata. fl. per. syst. 1. p. 93.
14. *P. TOMENTOSA* (Jacq. am. 37. obs. t. 61. f. 13.) capsulis obovatis foliisque tomentosis, pinnatis bijugis cum impari, foliolis ovato-rhombeis repandis obtusis, petiolo marginato. ζ in ins. Antillanis. Folia vetusta subglabrata.
15. *P. SENEGALENSIS* (Juss. ann. 4. p. 348.) capsulis ovatis obtusè trigonis, foliis pinnatis bijugis cum impari, foliolis sessilibus ovatis remotè crenatis, impari majore, petiolis marginatis, caule angulato subtomentoso. ζ in Senegal.
16. *P. CUPANA* (H. B. et Kunth prov. gen. am. 5. p. 117.) capsulis ovatis acutis, foliis imparipinnatis, bijugis, foliolis ovato-oblongis acuminatis grossè serrato-crenatis coriaceis glabris, subtus in nervo venisque puberulis. ζ in ripâ fluminis Orinoci.
17. *P. SPHÆROCARPA* (Rich. in Juss. ann. 4. p. 348.) capsulis sphæricis lævibus, foliis pinnatis bijugis cum impari, foliolis subsessilibus ovatis glabris remotè crenatis, acumine obtuso. ζ in Guineâ.
18. *P. SUBROTUNDA* (Pers. ench. 1. p. 443.) capsulis subrotundis, foliis pinnatis bijugis cum impari, foliolis oblongis supernè serratis. ζ in Peruvizæ nemoribus. — Semarillaria subrotunda fl. per. syst. p. 92. Arillus edulis.
19. *P. PTEROPODA* (fl. mex. ic. ined.) capsulis subsphæricis trisulcis, foliisque villosis pinnatis bijugis cum impari, foliolis grossè dentatis ovatis, terminali cuneato, petiolo marginato. ζ in Mexico.
20. *P. CURASSAVICA* (Lin. sp. ed. 1. p. 366.) capsularum valvis semi-obcordatis, foliis biternatis, foliolis ovalibus crenatis, imparibus basi cuneatis, petiolo intermedio marginato. ζ in Curassao et prope Caracas. — Plm. ed. Burm. t. 111. f. 2. — Jacq. obs. t. 61. f. 8. (v. s?)
21. *P. BARBADENSIS* (Jacq. enum. 36. obs. t. 61. f. 9.) capsularum valvulis semi-ovatis villosiusculis, foliis biternatis, foliolis ovalibus integerrimis serratisque coriaceis, petiolis submarginatis ζ in Barbados et ins. Antillanis. (v. s.)

22. *P. POLYPHYLLA* (Schum. l. c. p. 462.) capsularum valvis obovatis, foliis suprà decompositis, foliolis ovato-cuneatis apice crenatis suprà glabris subtùs petiolisque villosis. ζ in Amer. austràli. — Pluk. alm. t. 168. f. 5? Petioli villosi. Caveas ne confund. cum *Serjaniâ triternatâ*. An hujus congener?

†. *Species fructibus ignotis.*

* *Foliis ternatis.*

23. *P. NODOSA* (Jacq. enum. 35. obs. t. 61. f. 3.) foliis ternatis, petiolis nudis, foliolis ovatis subdentatis glabris, medio obovato, racemis hinc indè incrassato-nodosis. ζ in Antillis.
24. *P. DENSIFLORA* (Smith in Rees. cycl. n. 3.) foliis ternatis, petiolis alatis, foliolis obscurè serratis, racemis adgregatis ramosissimis petiolo quadruplè brevioribus. ζ in Novâ-Granadâ. Caulis 3-angularis sulcatus.

** *Foliis biternatis.*

25. *P. CARTHAGENENSIS* (Jacq. obs. 3. p. 11. t. 61. f. 6.) foliis biternatis, foliolis ovato-oblongis sinuatis, petiolis marginatis, caule inermi. ζ in Americâ ad Carthagenam.
26. *P. CARIBÆA* (Jacq. obs. 3. t. 62. f. 7.) foliis biternatis, foliolis ovalibus apice subdentatis, mediis obovato-cuneatis, petiolis marginatis, ramis aculeatis. ζ in Caribæis Jacq.; in prov. Caracasanâ? H. B. et Kunth nov. gen. am. 5. p. 118. An diversæ species?
27. *P. PUBESCENS* (H. B. et Kunth l. c.) foliis biternatis, foliolis grossè crenato-serratis membranaceis glabris subtùs molliter tomentoso hirtis aureo fuscescentibus, terminalibus ovato-oblongis acuminatis, petiolis partialibus submarginatis. ζ ad ripam fl. Amazonum.
28. *P. MOLLE* (H. B. et Kunth l. c.) foliis biternatis, foliolis grossè crenato-serratis membranaceis suprà in nervo venisque hirtis, subtùs molliter hirtotomentosis terminalibus subrotundo-ovatis acutis, petiolis nudis. ζ in ripâ flum. Magdalenæ.

*** *Foliis triternatis.*

29. *P. TRITERNATA* (H. B. et Kunth l. c.) foliis triternatis, foliolis 3-4-dentatis coriaceis glabris nitidis, terminalibus obovatis, petiolis partialibus rachique alatis. ζ in ripâ fluminis Orinoci.

**** *Foliis decompositis bipinnatisve.*

30. *P. DIVERSIFOLIA* (Jacq. obs. 3. t. 62. f. 14.) foliis suprà decompositis, infimè pinnatis, reliquis ternatis, petiolis marginatis, foliolis ovatis glabris subdentatis ζ in Antillis. An *Serjaniæ* species? Affinis. *S. triternatæ*. An eadem ac *S. heterophylla*?
31. *P. HISPIDA* (Jacq. h. schœnbr. 3. t. 268.) foliis infernè bipinnatis, apice pinnatis. foliolis lanceolatis acuminatis subserratis, ramis angulatis hispidis, stipulis ovatis ciliatis racemis adgregatis. ζ ad Caracas.
32. *P. BIPINNATA* (Poir. dict. 4. p. 99.) foliis bipinnatis, foliolis ovatis subsessilibus apice subcrenatis, petiolis nudis ramisque subtomentosis, capsulis alatis apice bilobatis. ζ in Brasiliâ ad Rio-Janeiro. An *Serjaniæ* species?

***** *Foliis pinnatis.*

33. *P. CAULIFLORA* (Jacq. ic. rar. 3. t. 450.) foliis pinnatis, pinnulâ infimâ ternatâ, petiolo basi nudo ad apicem marginato, floribus corymboso-capitatis axillaribus et terminalibus. ζ ad Caracas.
34. *P. JAPONICA* (Thunb. fl. jap. 170.) foliis pinnatis bijugis cum impari, foliolis petiolulatis marginatis, infimis subsessilibus trilobis, caule herbaceo inermi. ζ in Japoniâ.
35. *P. CUPANLEFOLIA* (Rich. in Juss. ann. mus. 4. p. 349.) foliis pinnatis bijugis

eum impari, foliolis sessilibus ovato-oblongis remotè crenatis glaberrimis, petiolo basi nudo inter foliola marginato. ζ in sylvis Guianæ.

36. *P. CONNARIFOLIA* (Rich. in Juss. l. c. p. 349.) foliis pinnatis bijugis cum impari, foliolis sessilibus coriaceis ovatis obtusè mucronatis subintegris glabris ramisque ferrugineis, petiolo nudo, pedunculis cirrho destituti. ζ in sylvis ripariis Guianæ.
37. *P. FIBULATA* (Rich. in Juss. l. c.) foliis pinnatis bijugis cum impari, foliolis brevè petiolatis ovatis subcoriaceis subdenticulatis subtùs subtomentosis ramisque ferrugineis, petiolo nudo, pedunculis cirrho incrassatis fibuliformis floriferis. ζ in sylvis ripariis Guianæ; ex cirrho dicta *Patte de chauve-souris*.
38. *P. RUFESCENS* (Rich. in Juss. l. c.) foliis pinnatis bijugis cum impari, foliolis subpetiolatis ovato-lanceolatis remotè crenatis subtùs tomentoso-ferrugineis, suprà nigro-virentibus, petiolis nudis, ramis subangulatis ferrugineis. ζ in sylvis ripariis Guianæ. Pedunculi cirrho nunc instructi nunc destituti.
39. *P. INGÆFOLIA* (Rich. in Juss. l. c.) foliis pinnatis trijugis cum impari, foliolis coriaceis glabris, infimis 3-foliolatis, petiolo basi nudo apice marginato, ramis striatis punctato scabris. ζ in Guianæ sylvis. flum. Amazonio conterminis. Pedunculi nunc ramoso-paniculati, nunc simplices ultrà medium cirrhosi et floriferi.

Trib. II. *SAPINDEÆ*. *H. B. et Kunth nov. gen. am. 5. p. 121.*

Petala intùs barbata glandulosa aut nuda non verè appendiculata. Discus annularis aut rariùs glandulæ basi connatæ interpetala et stamina. Ovarium 2-3-loculare, loculis 1-spermis. — Arbores aut Frutices non scandentes.

V. *SAPINDUS*. *Lin. gen. n. 499.*

Cal. 4-5-sepalus coloratus, sep. 2 exterioribus. Petala 4-5 esquamata, intùs glandulosa aut barbata. Stam. 8. filam. villosis. Stylus 1. Stigmata 3. Carpella 3 globosa carnosà connata, 2 sæpè arbotiva. Semina sphærica. — Arbores inermes (1), foliis abruptè pinnatis et abortu unius folioli sæpè impari-pinnatis.

* *Rachi alata.*

1. *S. SAPONARIA* (Lin. sp. 526. excl. syn. Pluk.) rachi foliorum decurrente latè alatà, foliolis integerrimis lanceolatis 3-4-jugis, terminalibus longè acuminatis ζ in ins. Caribæis et Americà calid. — Comm. hort. 1. t. 94. Fructus globosi. Frutex nascent, quem viv. in h. Genev. video, foliis simplicibus elongatis ferè Dodonæam refert. (v. s. sine flore.)
2. *S. MARGINATUS* (Willd. enum. 432.) inermis, rachi supernè angustè marginatà, foliolis lanceolatis 6-jugis. ζ in Georgià. *S. Saponaria* Michx. fl. am. bor. 1. p. 242. Fructus therebinthinacei.
3. *S. FORSYTHII*, rachi angustè alatà, foliolis integerrimis coriaceis subpetiolulatis ellipticis utrinquè attenuatis 3-5-jugis. ζ in arenosis maritimis insulæ Granatæ. Folia odorem Allii spirant. Fructus globosi. Flores parvi albi numerosissimi densè paniculati terminales. (v. s.)
4. *S. STENOPTERUS*, rachi angustissimè alatà, foliolis integerrimis coriaceis ovato-oblongis acuminatissimis 4-5-jugis. ζ in Sancto-Domingo. Bertero. Panicula terminalis laxè paniculata. Flores parvi. Affinis *S. rigido* sed differre videtur petiolo alato glabro. (v. s.)

(1) *S. spinosus* Linn. videtur *Fagara* affinis ex Lun. h. Jam. 1. p. 443.

** *Rachi non alata.*

5. *RIGIDUS* (Ait. h. Kew. 2. p. 36) rachi apterâ pubescente, foliolis ovato-oblongis glabris 3-jugis. 5 in Americâ calid. — Pluk. alm. t. 217. f. 7. Gaertn. fr. 1. t. 70. f. 3. An eadem stirps Borbonica à Poiretio descripta sub eodem nomine?
 6. *S. ARBORESCENS* (Aubl. guian. 1. p. 357. t. 139.) rachi apterâ, foliolis ovatis brevè acuminatis glabris 3-jugis, racemis axillaribus subsimplicibus. 5 in Guianæ sylvis prope Amnem Galibiensem. Flores ign. Fructus rubri parvi.
 7. *S. FRUTESCENS* (Aubl. guian. 1. p. 355. t. 138.) rachi apterâ foliolis oblongo-lanceolatis acuminatis glabris 7-jugis, racemis axillaribus subsimplicibus. 5 in Guianæ sylvis et campis cultis. Fructus globosi coccinei. Pruni ferè magnitudine. (v. s.)
 8. *S. SURINAMENSIS* (Poir. dict. 6. p. 665.) rachi apterâ, foliolis elliptico-lanceolatis glaberrimis membranaceis 6-8-jugis, paniculis divaricatis. 5 Surinami. Fructus globosi vix Cerasi magnitudine.
 9. *S. INÆQUALIS*, rachi apterâ, foliolis ovato-lanceolatis acuminatis glabris ad latus anticum latioribus 4-jugis, paniculis divaricatis. 5 in Guadalupâ. Bertero. Lam. ill. t. 307. f. 1. male in Dict. ad *S. Saponariam* relata. *S. laurifolius* in h. Balb. non Vahl. (v. s.)
 10. *S. ANGULATUS* (Poir. dict. 6. p. 665.) rachi apterâ, foliolis lato-ovatis obtusis coriaceis supernè glabris lucidis subtùs pubescentibus 3-jugis, fructibus 3 carpellis ad dorsa angulato-carinatis. 5 Patr. ign.
 11. *S. SENEGALENSIS* (Poir. dict. 6. p. 666.) rachi complanato-striatâ subpubescente, foliolis ovato-lanceolatis glabris utrinquè nervosis 2-4-jugis. 5 in Senegaliâ. Fructus globosi Fragariæ magnitudine.
 12. *S. TETRAPHYLLUS* (Vahl symb. 3. p. 54.) rachi apterâ, foliolis lanceolato-oblongis glabris bijugis, racemis subsimplicibus, petalis glabris. 5 in Indiâ orientali.
 13. *S. SALICIFOLIUS*, rachi apterâ, foliis lineari-lanceolatis acuminatis glabris bijugis, racemis compositis, petalis glabris, intùs barbatis. 5 in insulâ Timor. (v. s. sine fr.)
 14. *S. LAURIFOLIUS* (Vahl symb. 3. p. 54.) rachi apterâ, foliolis ovato-oblongis attenuatis glabris 3-jugis, paniculâ laxâ, petalis margine tomentosis. 5 in Malabariâ. — Rheed. mal. 4. t. 19. *S. trifoliata* Lin. sp. 625.
 15. *S. LONGIFOLIUS* (Vahl symb. 3. p. 53.) rachi apterâ, foliolis lanceolatis glabris 5-jugis, unico terminali. 5 in Indiâ orient?
 16. *S. EMARGINATUS* (Vahl symb. 3. p. 54.) rachi apterâ, foliolis oblongis emarginatis subtùs villosis 2-3-jugis, paniculis decompositis pubescentibus, petalis margine tomentosis. 5 in Indiâ orient.
 17. *S. RUBIGINOSUS* (Roxb. cor. 1. p. 44. t. 62.) rachi apterâ, foliolis oblongo-lanceolatis acutis subtùs villosis 3-5-jugis, paniculis terminalibus, petalis glabris. 5 in montibus Coromandelicæ et in Pulo-pinang. Mal. misc. 1. n. 1. p. 11. Baccæ 3-oblongæ.
 18. *S. PRAXINIFOLIUS*, rachi apterâ, foliolis oblongo-lanceolatis acutis subtùs petiolisque pubescentibus 3-5-jugis, paniculis terminalibus, petalis glabris, intùs barbatis. 5 in ins. Timor. Priori affinis. (v. s.)
- †. *Species non satis notæ.*
19. *S. RARAK*, rachi apterâ, foliolis oblongis acuminatis glaberrimis membranaceis 10-12-jugis. 5 in Javâ. — Rarak. Rumph. amb. 2. p. 134. *S. Saponaria* Lour. coch. 1. p. 293? *S. pinnatus* Mill. dict. n. 3? Folium unicum habeo in Javâ à cl. Labaye lectum. (v. s.)
 20. *S. ABRUPTUS* (Rour. coch. 1. p. 293.) rachi. . . . foliolis lanceolatis glabris, racemis subterminalibus, floribus campanulatis 4-sepalis 4-petalis, stylo 3-sulcato. 5 ad Cantonem Sinarum.

21. *S. MUKOROSHI* (Gaertn. fr. 1. p. 342. t. 70.) rachi. . . . , foliolis ovatis lanceolatis glabris alternis 6-jugis teneris, baccis ovatis utrinque acuminatis. 5 in Japoniâ circa Iedo. Thunb. fl. jap. 356. n. 37.~

22. *S?* *INDICUS* (Poir. dict. 6. p. 667) foliis subsessilibus glabris venoso-reticulatis, sæpius lineari-lanceolatis indivisis, interdum basi in lobos 2 oppositos ovato-lanceolatos pinnatis-partitis. 5 in Indiâ. Cultus in hort. Paris. ubi nondum floruit.

VI. *BLIGHIA*. *Kœnig in ann. bot.* 18. 6. 2. p. 571. — *Akeesia Tussac fl. ant.* 1. (1808) p. 66. — *Bonannia Raf. specch.* (1814) 15. p. 115.

Cal. 5-partit. Petala 5 intus vix appendiculata? Stam. 8. Stylus brevissimus. Stigm. 3. Carpella 3 connata carnosa apice dehiscentia? Semina solitaria, arillo maximo carnoso insidentia. — Arbor habitu Sapindi aut Cupaniæ.

1. *B. SAPIDA* (Kœn. loc. cit. 2. t. 16. 17.) 5 in Guineâ undè in ins. Caribæas migravit. *Akeesia Africana Tussac ant.* t. 3. *Bonannia nitida Raf. l. c.* p. 116. *Akce Lunan. h. jam.* 2. p. 335. — *Arillus edulis. Rachis aptera pubescens. Foliola ovato-lanceolata venosa 3-4-juga.* (v. s.)

VII. *TALISIA. Aubl. guian.* 1. p. 349.

Flores abortu monoico-polygami. Cal. 5-partitus. Petala 5 intus ad basin glandulosa. Stam. 8 10 breviora. Ovarium 4-5-locul. Stylus brevis indivisus. Fruct. . . . Arborea pinnatifoliæ.

1. *T. HEXAPHYLLA* (Vahl ecl. 2. p. 29.) racemis simplicibus, calycibus petala æquantibus, foliis 2-3-jugis, foliolis oblongo-lanceolatis nitidis utrinque glabris. 5 in Americâ calid.

2. *T. GUIANENSIS* (Aubl. l. c. t. 136.) racemis decompositis, calycibus petalis brevioribus, foliis multijugis, foliolis ovali-oblongis acuminatis subtus in nervis nervulisque hirsutis. 5 in Guianâ ad ostium flum. Sinemari. Lam. ill. t. 310. *T. rosea Vahl ecl.* 2. p. 30. Cum sequente forsâ ab Aubletio confusa. (v. s.)

3. *T. GLABRA*, racemis decompositis, calycibus petalis brevioribus, foliis multijugis, foliolis ovato-lanceolatis acuminatis coriaceis utrinque glaberrimis. 5 in Guianâ seu Cayennâ. Patris. (v. s. sed florib. malè conserv. non dessicatis.)

4. *T?* *ACLADODEA*, racemo terminali, trunco simplici, foliis abruptè pinnatis, foliolis oblongo-linearibus acuminatis. 5 in Peruviz nemoribus. *Acladodea pinnata fl. per. prod.* 133. t. 29. syst. p. 262. Arbor habitu Palmarum. Flores fœminei ignoti. †.

VIII? *MATAYBA. Aubl. guian.* 1. p. 331. t. 128. — *Ernstingia Neck. elem.* → *Ephielis Schrebr. gen.* n. 647.

Cal. 5-partitus. Petala 5 basi intus glandulis duabus aucta. Stam. 8 filamentis villosis. Stylus 0. Stigma 1. Capsula oblonga 1-locul. 2-valvis, valvulâ alterâ nudâ, alterâ intus medio 2-spermâ. Semina arillata reniformia.

1. *M. GUIANENSIS* (Aubl. l. c. Lam. ill. t. 298.) foliis glabris abruptè pinnatis 3 4-jugis, petiolo aptero, racemis paniculatis. 5 in Guianæ sylvis et in Sancto-Domingo, ex Bertero. *Ephielis fraxinea Willd. sp.* 2. p. 328. *E. Guianensis Pers.* Folia abruptè pinnata. Paniculæ divaricatæ. (v. s. sine fr.)

2. *M. PATRISIANA*, foliis pinnatis bijugis cum impari, foliolis subtus villosis, petiolo subalato, racemis simplicibus. 5 in Guianâ aut Cayennâ. Patris. Fructus priori videtur simillimus. (v. s. sine fl.)

3. *M?* *VOUA-RANA*, foliis pinnatis glabris, foliolis 6 alternis, ultimo terminali,

tibus, paniculis laxis, baccis globosis sublævibus. 5 in Chinâ et Cochinchinâ. *Dimocarpus Longan* Lour. coch. 1. p. 288. — *Buchoz* ic. col. t. 99. Fructus flavescentes. (v. s.)

3. *E. INFORMIS*, foliolis. . . . , paniculis paucifloris, baccis tuberculosi irregularibus. 5 in Cochinchinâ sylvis. Folia ferè *Louganæ*. Baccæ carnosæ acerbæ sæpius geminæ.
4. *E. NEPHÉLIUM*, foliolis lanceolatis, baccis subovatis crinitis. 5 in Indiâ orient. *Nephelium lappaceum* Lin. syst. 4. p. 236. Lam. ill. t. 764. Marsd. sum. ic. *Dimocarpus crinita* Lour. l. c. — Vulgò *Rambutan* seu *Ramposan*. Bont. jav. f. 109. Fructus sæpè geminati.

XII. *THOUINIA*. *Poit. ann. mus.* 3. p. 70. t. 6. (non *Thunb. nec Smith.*)

Cal. 4-5-partitus. Petala 4-5 medio intus pilorum fasciculo instructa. Stam. 8-10 libera. Styl. 1. Stigm. 3. Carpella 3 stylo connata 1-sperma samaroidæ nempè in alam membranaceam supernè desinentia. — Flores sæpè abortu masculi.

* *Foliis simplicibus.*

1. *T. SIMPLICIFOLIA* (*Poit. ann.* 3. p. 71. t. 6.) foliis simplicibus grossè dentato-serratis subtus subtomentosis, racemis simplicibus. 5 in Sancto-Domingo prope urbem Cap. Racemi axillares foliis paulò breviores.
2. *T. INTEGRIFOLIA*, (*Spreng. neue entd.* 2. p. 155.) foliis simplicibus integerrimis utrinquè glabris, racemis paniculatis. 5 in Brasiliâ. Folia subtus discolora. Panicula terminalis flaccida.

** *Foliis trifoliolatis.*

3. *T. TRIFOLIATA* (*Poit. ann.* 3. p. 72. 5. t. 27.) foliis trifoliolatis, foliolis subsessilibus ovalibus basi attenuatis subserratis glabris, axillis venarum subtus pilosis, racemis simplicibus. 5 in Sancto-Domingo circa Fort-Dauphin. (v. s.)
4. *T. TOMENTOSA*, foliis trifoliolatis, foliolis elliptico-oblongis serratis superne glabris subtus velutino-tomentosis albidis, racemis simplicibus. 5 in Sancto-Domingo. Bertero. Folia et carpella dimidiò minora quàm in priore. (v. s.)
5. *T? VILLOSA* (fl. mex. ic. ined.) foliis trifoliolatis villosis, foliolis ovatis apice serratis acutis, racemis paniculatis. 5 in Novâ-Hispaniâ circa Quauahua. Carpella 3 samaroidæ à basi distincta. An generis?

*** *Foliis pinnatis.*

6. *T. PINNATA* (*Turp. ann. mus.* 5. p. 401. t. 26.) foliis pinnatis, foliolis obovatis subemarginatis, floribus paniculatis 5 petalis 8-andris. 5 in Sancto-Domingo inter Monte-Christ et San-Yago.
7. *T. POLYGAMA* (*Mey. prim. esseq.* 156.) foliis pinnatis, foliolis oblongo-ovatis, floribus racemosis villosis 4-petalis 8-andris. 5 in sylvis arenosis Guianæ.
8. *T. DECANDRA* (*H. et Bonpl. pl. eq.* 1. p. 198. t. 56.) foliis pinnatis sexjugis foliolis lanceolatis dentatis, floribus paniculatis 5-petalis 10-andris. 5 in calidis Mexici circa Acapulco.

XIII. *TOULICIA*. *Aubl. guian.* 1. p. 359. t. 140. — *Ponæa Schreb. gen. n.* 682.

Cal. 5-partitus. Pet. 4 disco intus lævi. Stam. 8. Ovarium 3-gonum. Styli 3. Capsula 3-alata, 3-locul., alis bivalvibus, loculis 1-spermis. — An satis à *Cupaniâ* differt?

1. *T. GUIANENSIS* (*Aubl. l. c. Lam. ill. t.* 317.) 5 in Guianæ sylvis. *Ponæa saponarioides* Willd. sp. 2. p. 470. Folia abruptè pinnata 8-juga, foliol. oppositis lanceolatis muticè ad basin latioribus. (v. s.)

XIV. *CUPANIA*. *Plum. gen.* 49. t. 19. *Pers. ench.* 1. p. 413.

Cal. 5-sepalus. Pet. 5 intus glabra cucullata. Stam. 8. Stylus 3-fidus.

Capsula 5-locul. 3-valvis, valvis medio septiferis, loculis 1-2-spermis. Semina erecta arillata. — Arbores foliis abruptè pinnatis, floribus sæpè abortu masculis.

Sect. I. TRIGONIS. *Jacq. am.*

Petala apice convoluta-cucullata. — Species Americanæ.

1. *C. TOMENTOSA* (Swartz fl. ind. occ. 657.) foliis 3-4-jugis, foliolis obovatis retusis subtus tenuissimè velutino-tomentosis ad apices nervorum lateralium dentato-serratis, subtus tomentosis extimis majoribus. 5 in montibus sylvaticis Hispaniolæ, Guadalupæ, et ad ripam fl. Magdalenæ. Trigonis tomentosa Jacq. am. 102. — Plum. ed. Burm. 1. 110? et ideò *C. Americana* Lin. sp. 290. Petala triangularia intus hirsuta. (v. s. sine fr.)
2. *C. EXCELSA* (H. B. et Kunth nov. gen. am. 5. p. 125.) foliis pinnatis, sub-5-jugis, foliolis oblongis obtusiusculis remotè denticulatis coriaceis suprà glabriusculis subtus pubescentibus. 5 in montosis Mexici. Fr. ign.
3. *C. LATIFOLIA* (H. B. et Kunth l. c. p. 126.) foliis pinnatis bijugis, foliolis obovato-oblongis profundè retusis dentato-repandis coriaceis glabris subtus in venis puberulis. 5 in ins. fl. Magdalenæ *isla de Brugas* dicta. Fr. ign.
4. *C. SCROBICULATA* (H. B. et Kunth l. c. p. 127.) foliis pinnatis sub-4-jugis oblongis retusis remotè et argutè dentatis submembranaceis glabris, subtus in axillis venarum scrobiculatis. 5 prope Turbaco Novo-Granatensium. Affinis *C. glabræ*.
5. *C. GLABRA* (Swartz fl. ind. occ. 659.) foliis 3-4-jugis, foliolis ovatis obtusis crenatis glabris basi acuminatis. 5 in montibus Jamaicæ et Hispaniolæ.
6. *C. SAPONARIA* (Pers. ench. 1. p. 413.) foliis 5-6-jugis, foliolis oblongis attenuatis integris, subtus scabris pubescentibus. 5 in Indiâ occid. *C. saponarioides* Swartz fl. ind. occ. 2. p. 661.
7. *C. GEMINATA* (Poir. suppl. 2. p. 419.) foliis unijugis, foliolis ovatis integris suprà glaucis subtus puberulis, nervis lateralibus prominulis apice confluentibus. 5 in Cayennâ. Patris. An *C. diphylla* Vahl? Capsulæ juniores velutinæ obtusè trigonæ. (v. s. sine fl.)
8. *C. NITIDA*, foliis 1-2-jugis cum impari!, foliolis ovalibus acuminatis integris coriaceis utrinquè glaberrimis. 5 in Guianâ-Gallicâ seu Cayennâ. Capsulæ juniores velutinæ, adultæ calvescentes apice 3-alatæ. — An potius *Pauliniae* species? (v. s. sine fl.)

Sect. II. MOLINÆA. *Juss. gen. 245. Lam. ill. t. 305.*

Petala planiuscula calyce paulò majora. Filamenta brevissima villosa. — Species Mauritianæ aut Indicæ.

9. *C. LÆVIS* (Pers. ench. 1. p. 413.) foliis 2-3-jugis, foliolis oppositis oblongis integerrimis glaberrimis coriaceis, nervulis vix prominulis. 5 in insulâ Bourboniâ. — *Molinæa*. Lam. ill. t. 305. f. 1. *M. lævis* Willd. sp. 2. p. 329. Capsulæ obovatæ triquetro-alatæ. (v. s. sine fr.)
10. *C. ALTERNIFOLIA* (Pers. l. c.) foliis 4-5-jugis, foliolis subalternis oblongis integerrimis glaberrimis coriaceis, nervulis vix prominulis. 5 cum priore cujus mera varietas videtur. *Molinæa* Lam. ill. t. 305. f. 2. *M. alternifolia* Willd. sp. 2. p. 329. (v. s. sine fr.)
11. *C. VENULOSA*, foliis 1-2-jugis, foliolis oppositis oblongis integerrimis glaberrimis nervulis prominulis reticulatis. 5 in insulâ Bourboniâ seu Mauritiî. Foliola ferè *Vaccinii* aut *Vitis-ideæ* similia. (v. s. sine fr.)
12. *C. CANESCENS* (Pers. ench. 1. p. 413.) foliis bijugis, foliolis oppositis elliptico-oblongis integerrimis nitidis lævibus. 5 in Coromandelis montosis. Paniculæ axillares. Capsulæ ovatæ 3-sulcatæ non alatæ.

XX. KOELREUTERIA. *Laxm. nov. comm. petr.* 16. p. 561. t. 18. non Murr.

Calyx 5-sepalus. Petala 4 irregularia, basi subsquammata. Stam. 8. Stylus 1 crassus. Capsula 3 locul., inflata. Semina ovato-globosa, endoplevrâ intus penetrante et axeos loco centrum embryonis spiraliter convoluti occupante. — Folia impari-pinnata multijuga, foliolis ovatis grossè dentatis, floribus paniculatis luteis.

1. K. PANICULATA (Laxm. l. c.) § 5 in Chinâ. *Sapindus Chinensis* Linn. f. suppl. 228. K. paullinioides L'her! sert. 18. t. 19. K. paniculata Duham. ed. nov. t. 36. (v. v.)

XXI. AMIROLA. *Pers. ench.* 2. p. 565. — *Llagunoa fl. per. prod.* 126. t. 28.

Flores abortu-monoïco-polygami. Calyx 4-5-fidus lateraliter fissus, intus ad basin membranâ subcarnosâ vestitus. Pet. o. Stam. 8. Ovarium trigonum. Stylus filiformis 3-sulcus. Capsula inflata 3 valv., 3-loc., valvis membranaceis medio septiferis, axi non libero. Semina in loc. solitaria rotunda. Cotyledones spiraliter convolutæ, endoplevrâ non axeos vicem fungente (conf. Kunth nov. gen. am. 5. p. 442. ic. dext.)

1. A. NITIDA (Pers. l. c.) foliis glabris serratis ovatis acutis indivisis aut utrinquè lobo ad basin auctis et ideò subtrisectis. § in Peruvîæ sylv. Llagunoa nitida fl. per. l. c. Pedunculi trifidi, ramuli laterales sæpè abortivi. (v. v.)
2. A. PRUNIFOLIA (H. B. et Kunth nov. gen. am. 5. p. 131. sub Llagunoa) foliis ovato-ellipticis argutè dentatis, suprâ glabris, subtus hirtellis, nervo venisque hirtis-tomentosis. § prope Loxam. An à priori diversa?
3. A. MOLLIS (H. B. et Kunth l. c.) foliis ellipticis serratis, suprâ hirtellis, subtus molliter canescenti-tomentosis. § prope Loxam.

XXII. DODONÆA. *Lin. gen. ed.* 1. n. 855. non Plum.

Flores sæpè abortu polygami aut dioici. Cal. 4-partitus deciduus. Pet. o. Stam. 8, filam. brevissimis, antheris oblongis linearibusve. Stylus filiformis, ab alis capsulæ distinctus, apice subtrifidus. Capsula 2-3-valvis., 2-3-loc., 2-3-alata, angulo centrali 2-3-angulato in faciebus seminifero. Semina bina subglobosa. — Frutices foliis oblongis sæpius viscosis.

1. D. VISCOSA (Lin. mant. 238. excl. syn. plur. ex Meyer esseq. 157.) foliis obovato-oblongis basi cuneatis viscosis, floribus racemosis, fructibus bi aut triatatis pedicello longioribus. § in arenosis Americæ calid., ins. Carib., et regni Owariensis. — Plum. ed. Burm. t. 247. f. 2. Sloan. hist. 2. t. 162. f. 3. Variat foliis acutiusculis obtusis aut emarginatis. Capsulæ utrinquè profundè emarginatæ 7-9 lin. longæ, 8-12 latæ Conf. quoad var. Kunth nov. gen. am. 5. p. 134. An hic plures species confusæ? D. viscosa Forst. prod. 27. ex D. viscosa, spathulatâ, et triquetrà juxtâ cl. Smith confusis constat. (v. s.)
2. D. SPATHULATA (Smith in Rees cycl. v. 12. n. 2.) foliis lineari-obovatis viscidis, sepalis ovatis acutis nudis, floribus dioicis. § in insulis Sandwich. Minor quam D. viscosa.
3. D. JAMAÏCENSIS, foliis oblongo-lanceolatis utrinquè attenuatis margine sub-revolutis subviscosis, floribus breviter racemosis, fructibus pedicello brevioribus. § in frigidioribus Jamaïcæ. — Browne jam. t. 18. f. 1. D. angustifolia Swartz! obs. 150. D. viscosa Cav. ic. t. 327. (v. s. sine fr.)
4. D. BIALATA (H. B. et Kunth nov. gen. am. 5. p. 134. t. 442.) foliis lanceolatis utrinquè attenuatis viscosis, racemis subramosis, fructibus constanter bialatis pedicelli longitudine. § in arenosis maritimis prope Camanam.
5. D. BURMANNIANA, foliis oblongis basi cuneatis acutiusculis viscosis, floribus

- racemosis, fructibus pedicello longioribus. ♂ in Indiâ orient., Ceylonâ, Timor. — Burm! zeyl. t. 23. *Ptelea viscosa* Burm! ind. 36. *D. angustifolia* Roxb. cat. 28. Folia nunc obtusa nunc acutiuscula. Capsulæ 6 lin. longæ, 9 lin. latæ. (v. s.)
6. *D. MICROCARPA*, foliis oblongo-linearibus basi attenuatis apice obtusis emarginatisve, floribus racemosis, fructibus pedicello brevioribus. ♂ in ins. Bourboniâ. Lam. ill. t. 304. f. 2. Folia 3 lin. lata. Capsulæ 3 lin. longæ, 4 lin. latæ. (v. s.)
7. *D. SALICIFOLIA*, foliis oblongo-linearibus utrinquè acuminatis viscosis, floribus racemosis, fructibus... ♂ forsàn ex Novâ-Hollandiâ? *D. angustifolia* Lam. dict. 3. p. 292. in hortis Europ. culta sub nomine *Bois de Reinette*. Folia odorata, 4 lin. lata. (v. s. sine fr.)
8. *D. DIOICA* (Roxb. cat. 28. DC. mem. soc. genev. 1. p. 445.) foliis obovato-oblongis basi cuneatis apice acutis non viscosis, ramis junioribus compressis, adultis teretiusculis, floribus dioicis racemosis. ♂ in Indiâ orient. *D. oblongifolia* Link enum. 1. p. 381? *D. heterophylla* hortul. Sequenti valdè affinis. An ad hanc aut ad *D. triquetram* referenda icon. Rumph. amb. 4. t. 50. (v. v. et s. ♂.)
9. *D. TRIQUETRA* (Andr. rep. t. 230.) foliis lanceolatis utrinquè attenuatis, ramulis triquetris, floribus dioicis racemosis, fructibus angustè alatis pedicello brevioribus. ♂ in Novâ-Hollandiâ. (v. v.)
10. *D. CUNEATA* (Smith in Rees cycl. n. 5. Rudg. lin. tr. Lin. soc. lond. 11. p. 296. t. 19.) foliis oblongo-cuneatis acuminatis aut apice subtridentatis, ramulis vix angulatis, floribus subsessilibus. ♂ in Novâ-Hollandiâ ad Port-Jackson. Fructus ferè *D. viscosæ* (v. s.)
11. *D. ASPLENIFOLIA* (Rudg. l. c. p. 297. t. 20.) foliis oblanceolatis basi attenuatis apice tridentatis viscosis, floribus subracemosis, ramis triquetris. ♂ in Novâ-Hollandiâ ad Port-Jackson.
12. *D. ERIOCARPA* (Smith in Rees cycl. n. 6.) foliis elliptico-lanceolatis undulatis, ramulis fructibusque pilosis, calycis sepalis ellipticis reflexis. ♀ in insularum Sandwich montibus.

† *Species non satis notæ.*

13. *D. FILIFORMIS* (Link enum. 381.) foliis tenuissimis (*Ericæ*) 6-8-lin. longis. ♂.....
14. *D. ANGUSTISSIMA* (hortul.) foliis linearibus latitudine decuplò longioribus punctis albidis supernè stigmatosis. ♂ Patr. flor. et fr. ign. (v. s.)
15. *D. ELÆAGNOIDES* (Rud. in Schrad. neu. jour. n. 2. p. 292.) foliis oblongo-ovatis supernè squammoso-furfuraceis. ♂ in Sancto-Domingo.
16. *D. SERRULATA*, foliis elliptico-oblongis petiolatis serrulatis coriaceis glabris, capsulis coriaceis 3-alatis. ♀ in Monte-Video. Fl. et sem. ign. (v. s.)
17. *D. PINNATA* (Smith in Rees cycl. n. 7.) foliis impari-pinnatis pilosis 6-7-jugis, petiolo interruptè alato. ♀ in Novâ-Cambriâ australi. (v. s. fl. mac. comm. à cl. Lambert.)

XXIII. ALECTRYON. *Gaertn. fr. 1. p. 216. t. 46.*

Flos. Bacca sicca 1 locul. margine aut apice alata. Semen exalbuminosum arillo incompleto cinctum basi loculi adfixum erectum. Cotyledones spiraliter convolutæ, et radícula deorsum spectans utrin Dodonæâ.

1. *A. EXCELSUM* (Gaertn. l. c.) fructu apice alâ cristato. ♀ *Evonymoides excelsa* Sol. in coll. Banks.
2. *A. CANESCENS*, fructu undiquè alâ marginato ♀ in Novâ-Hollandiâ orâ orientali. Specimen meum floribus caret. Folia oblonga obtusa adpressissime pubescentia. Racemi axillares folii longitudine. Fructus ferè *Samerariæ*. Stylus alæ utrinquè connexus! Semen oblongum in meis fructibus crassum. (v. s.)

† *Genera non satis nota, Sapindaceis affinia.*

XXIV. EYSTATHES. *Lour. coch. 1. p. 289.*

Cal. 5-sepalus. Pet. 5 ovata calyci æqualia. Stam. 8. Ovarium subrotundum. — Stylus filiformis. Stigma obtusum. Bacca globosa carnosâ 1-loc. 4-sperma. — Dodonæe forsân affinis sed 5-petala.

1. *E. SYLVESTRIS* (*Lour. 1. c.*) 5 in montibus Cochinchinæ. *Valentinia sylvestris* *Raesch.* Flores albi racemosi. Folia ovato-oblonga acuminata integerrima glabra. Rami patentes. Lignum durum.

XXV? RACARIA. *Aubl. guian. suppl. t. 24. f. 382.*

Flos. . . . Drupa ovata 1-locul. Nuclei 3 oblongi trigoni, testâ fragili. +.

1. *R. SYLVATICA* (*Aubl. 1. c.*) 5 in Guianæ sylvis ad radices montis *Serpentis*. Truncus spinis suprâ foliorum cicatrices horridus. Lignum durum. Folia abruptè pinnata 3-juga, foliolis ovatis integris.

XXVI. VALENTINIA. *Swartz fl. ind. 687. t. 14.*

Cal. 5-partitus coloratus patens persistens. Pet. o. Stam. 8. Ovarium subrotundum. Styl. 1. Capsula baccata, intus pulposa, in valvulas 3-4 revolutas dehiscens. Semina 3 4 oblonga. — *Swartzius* ad *Rhamneas*, *Jussæus* ad *Samydeas*, referunt hoc genus; sed ad *Sapindaceas* pertinere videtur ex analogiâ foliorum cum *Thouiniâ simplicifoliâ*, petalis deficientibus ut in *Schleicherâ* et *Amirolâ*, partium numero *Sapindaceis* proprio.

1. *V. ILICIFOLIA* (*Swartz 1. c.*) 5 in sterilissimis petrosis Hispaniolæ et Cubæ. Folia ferè *Aquifolii* sed magis oblonga, ad *Thouiniam simplicifoliam* accedentia. (v. s. sine fl.)

XXVII. PEDICELLIA. *Lour. fl. coch. 2. p. 805.*

Flores polygamo-dioici. Cal. 5-partitus, lobis acutis parvis. Pet. o. Stam. 8. Ovarium subrotundum pedicellatum. Stylus subnullus. Stigm. 3 reflexa. Capsula 3-valvis. Semen 1 pedicello proprio insistent. — *Arbuscula*, foliis oppositis lanceolatis integerrimis, racemis terminalibus. — Ordo dubius ob folia opposita.

1. *P. OPPOSITIFOLIA* (*Lour. 1. c.*) 5 in sylvis Cochinchinæ.

XXVIII. RATONIA.

Cal. parvus 5-sepalus persistens. Flores ignoti. Fructus compressi subcoriacei indehiscens aut vix dehiscens biloculares obcordati substipitati glabri stylo brevissimo apice bifido terminati. Semina in loculis solitaria erecta, funiculo crasso. — Arbor aut Frutex glaber. Folia alterna exstipulata abruptè pinnata 3-juga, petiolo aptero apice in acumen molle producto, foliolis oppositis obovato-oblongis obtusis integerrimis. Paniculæ terminales, ramis paucis elongatis racemosis.

1. *R. DOMINGENSIS*. 5 in Sancto-Domingo ubi ab Hispanis *Raton* dicitur ex cl. Bertero. Fructus refert capsulam *Badieræ*. (v. s. sine fl.)

XXIX? ENOUREA. *Aubl. guian. 1. p. 588. t. 235.*

Cal. 4-5-partitus, lobis 2 majoribus. Pet. 4 fundo calycis inserta, 2 majora, omnia squammâ petaloideâ intus aucta. Glandulæ 2 ad basin petal. major. Stam. 13 inæqualia basi connata subunilateralia. Ovarium subrotundum. Styl. o. Stigm. 3. Capsula 1-loc., 3-valv., 1-sperma. Semen erectum pulpâ farinosâ vestitum. — Frutices scandentes, foliis impari-pinnatis, cirrhis axillaribus, racemis paniculatis. An potius ob succum lacteum et petala perigyna dicta ad *Terebinthaceas* repellenda?

1. *E. CAPEOLATA* (*Aubl. guian. 1. c.*) foliis bijugis cum impari, foliolis coriaceis subtus puberulis ferrugineis. 5 in insulâ Guianæ *Sinemari*. (v. c.)

Ordo XLIV. MELIACEÆ. *Juss. gen.* 263. *mem. mus.* 3. p. 436.

Cal. Sepala 4-5 plus minus inter se in calycem monophyllum coalita. Pet. totidem sepalis alterna, hypogyna, ungue lato sæpius basi conniventia aut coalita, æstivatione sæpius valvatâ. Stam. numero petalorum dupla, (rariùs æqualia, tripla quadrupla); filamenta in tubum apice dentatum longè coalita, antheris ad faucem tubi sessilibus intus adnatis. Ovarium 1. Styl. 1. Stigmata distincta aut coalita. Fructus varius, baccatus, drupaceus capsularisve multilocularis, abortu sæpè 1-locul., valvis (dum absunt) medio septiferis. Semina nunc albuminosa, nunc exalbuminosa, embryo dicotyledoneo formâ vario.

Arbores aut frutices è regionibus tropicis ortæ. Folia alterna, exstipulata, simplicia aut composita.

Ordo non satis definitus et forsân typos plurimos diversos colligens, sed ob descriptiones plurium generum mancās, in statu scientiæ præsentî extricatu difficillimus et botanicis peritis commendandus! An hujus ordinis Geruma caractere manco tantum definita et forsân Malvaceis analogâ; Sandoricum stylo nullo et stigmatibus 5 donatum forsân Zanthoxylo affine? An tres tribus infra indicatæ ut ordines proprii habendæ? An *Styrax*, *Quivisiæ* et *Turrææ* habitu similis, huc revocanda? An inter *Cedreleas* genera exalbuminosa et præsertim *Carapa* ritè aliis consociata?

Trib. I. MELIÆÆ.

Fructus loculi 1-2-spermi. Semina in alam non expansa, exalbuminosa. Embryo inversus. Cotyledones planæ, foliaceæ. — Arbores aut frutices. Folia alternâ, in generibus 5 prioribus simplicia, in 2 posterioribus composita.

I? GERUMA. *Forsk. deter.* 62.

Cal. planus 5 dentatus. Petala 5 lanceolata, patentia. Stam. 5 erecta, in anulum crassum basi coalita. Antheræ trigonæ. Styl. 1. Stigm. 3. Caps. ovalis 4-5-locul., 4-5-valvis. Sem. in loculis 2 pulpæ trigonæ inserta (*Forsk.*)

1. *G. ALBA* (*Forsk. l. c.*) 5? in Arabiâ circa Hadie. Folia alterna ovali-oblonga subserata.

II. HUMIRIA. *Homiri Aubl. guian.* 1. p. 564. t. 225. — *Myrodendron Schreb. gen. n.* 901.

Cal. obtusè 5-dentatus. Pet. 5. oblonga, æstivatione valvatâ. Stam. 20 in tubum apice dentatum monadelphum, antheris erectis. Stylus 1. Stigma capitatum sub-5-radiatum. Pericarp. (ex *Aubl.*) 5-locul., loculis 1-spermis.

1. *H. BALSAMIFERA* (Aubl. l. c.) 5 in sylvis Guyanæ et Cayennæ. Myrodermum amplexicaule Willd. spec. 2. p. 1171. Cortex crassus balsamum rubrum fundens. Folia ovato-oblonga integra semi-amplexicaulia, dorsali nervo decurrentia, juniora involutiva. Flores corymbosi. (v. s.)

III. TURRAEA. *Lin. mant. n. 1306.*

Cal. 5-fidus. Pet. 5 longissima ligulæformia. Stam. 10, tubo longissimo apice 10-fido, antheris ad loborum basim aut inter eos insertis. Styl. 1. Stigma crassiusculum. Capsula 5-locul., *locul. 2-spermis, valvis medio septiferis.

1. *T. VIRENS* (Lin. mant. 237.) foliis elliptico-lanceolatis acuminatis emarginatis glaberrimis, calycibus fructibusque sericeo-villosis. 5 in Indiâ orient. ad scorias montium quondam ardentium. Smith ic. ined. 1. t. 10.

ℒ? *Billardierii*, foliis ovato-lanceolatis acuminatis junioribus pubescentibus demum glabris, calycibus pubescentibus, stigmatate longè exserto. 5 in Indiâ orient. insulis. An species propria? (v. s.)

2. *T. PUBESCENS* (Willd. spec. 2. p. 555.) foliis ovatis emarginatis subtus pubescentibus, calycibus villosis. 5 in ins. Mainam. Hellen in act. holm. 1788. p. 296. t. 10. f. 3.

3. *T. MACULATA* (Smith ic. ined. 1. t. 11.) foliis ovatis acutis glabris, calycibus ciliatis, petalis glabris 5 in Madagascar. T. glabra Cav. diss. 7. p. 360. t. 204. Flos 2-3 poll. longus.

4. *T. SERICEA* (Smith ic. ined. 1. t. 12.) foliis ovatis obtusiusculis utrinque tomentosis, calycibus pedunculis petalisque villosis. 5 in Madagascar. T. tomentosa Cav. l. c. p. 361. t. 205. f. 2. Petala linearia ferè 3 poll. tubo stamineo longiora.

5. *T. LANCEOLATA* (Cav. diss. 7. p. 361. t. 205. f. 1.) foliis lanceolatis utrinque obtusiusculis glabris, laciniis calycinis tubo longioribus lanceolatis, petalis linearibus tubo stamineo brevioribus. 5 in Madagascar. Flos pollicem longus.

6. *T. RIGIDA* (Vent. choix t. 48.) foliis ellipticis acuminatis margine revolutis rigidis lucidis, calycibus petalisque glabriusculis, ramis strictis. 5 in ins. Mauritiî. Petala pollicaria tubo paulò longiora subpilosa.

7. *T. HETEROPHYLLA* (Smith in Rees cycl. 36. n. 6.) foliis ovatis indivisis 3-lobisve, venis villosis, calycibus hirsutis breviter dentatis, petalis subspathulatis tubo stamineo longioribus. 5 in Africâ ad Cape-Coast Anglorum.

†. *Species non satis nota.*

8. *T? HERBACEA* (Poir. dict. 8. p. 147.) foliis elliptico-lanceolatis obtusis glabris, floribus solitariis, calycibus striatis glabris. (T)? in Rio-Janeiro. Petala apice laciniata, filamenta corollæ breviora. A. Turraea, inod. à Meliaceis omninò, ex descriptione, diversa videtur!

IV. QUIVISIA. *Comm. in Juss. gen. 264. — Gilibertia Gmel. syst. 682.*

Calyx urceolatus 4-5-dentatus. Pet. 4-5 breviter extus sericea. Stam. 8-10, tubo brevi. Stigma capitatum. Capsula coriacea 4-5 locul. apice in valvulas 4-5 dehiscent, valvis medio septiferis, loculis 2-spermis.

1. *Q. DECAUNDA* (Cav. diss. 7. p. 367. t. 211.) foliis alternis oblongis integris utrinque attenuatis, pedunculis axillaribus racemosis. 5 in ins. Mauritiî. *Q. racemosa* Pers. ench. 1. p. 468. *Gilibertia decandra* Willd. spec. 2. p. 551. (v. s.)

2. *Q. OPPOSITIFOLIA* (Cav. diss. 7. p. 368. t. 214.) foliis oppositis ovatis integerrimis glabris, pedunculis axillaribus 3-5-floris. 5 in insulâ Mauritiî. *Gilibertia oppositifolia* Willd. l. c. (v. s.)

3. *Q. OVATA* (Cav. diss. 7. p. 368. t. 212.) foliis alternis obovatis obtusis integris, pedicellis ternis axillatibus 1-floris, fructibus tomentosis. ζ in ins. Borboniâ. *Gilbertia ovata* Willd. sp. 2. p. 552. — *Gilbertia rutilans* Smith in Rees cycl. 16. n. 4. ab hac vix differt nisi floribus vix racemosis. An distincta?

4. *Q. HETEROPHYLLA* (Cav. l. c. t. 213.) foliis alternis ovalibus obovatisve integris sinuato-dentatis pinnatifidisve, pedicellis geminis axillaribus 1-floris. ζ in ins. Mauritii. Bory! voy. 1. p. 198. (v. s.)

V. *STRIGILIA*. Cav. diss. 7. p. 358. — *Foveolaria Ruiz et Pav. syst. fl. per.* 99. — *Tremanthus Pers. ench.* 1. p. 467.

Cal. campanulatus 5-dentatus. Petala 5, basi subcoalita, lac. linearibus extus sericeis. Stam. 10, filam. in tubum coalitis, antheris post dehisc. punctis setaceo-stellatis asperis. Drupa obovata subtrilocularis abortu monosperma (R. et P.) verosim. 6 loc., loculis 1-spermis (Cav.) — Valdè accedit ad *Styracem* sed differt antheris apice tubi sessilibus nec filamento apice libero.

1. *S. RACEMOSA* (Cav. diss. 7. p. 358. t. 201.) foliis oblongis brevi acuminatis, subtus eglandulosis velutinis, nervis ferrugineis, racemis solitariis erectis. ζ in Peruvîâ. *Tremanthus ferrugineus* Pers. ench. 1. p. 467. *Foveolaria ferruginea* Ruiz et Pav. l. c. (v. s.)

2. *S. OBLONGA* (Ruiz et Pav. l. c. sub *Foveolaria*) foliis oblongis acuminatis glabris, racemis erectis solitariis geminisve. ζ in Peruvîâ.

3. *S. OVATA* (Ruiz et Pav. l. c. sub *Foveolaria*) foliis ovato-oblongis acuminatis glabris, foveolis minimis glanduliferis. ζ in Peruvîâ.

4. *S. CORDATA* (Ruiz et Pav. l. c. sub *Foveolaria*.) foliis cordato-ovatis acutis subtus helvolis, foveolis glanduliferis. ζ in Peruvîâ nemoribus.

VI. *SANDORICUM*. Cav. diss. 7. p. 359.

Cal. brevis obtusè 5-dentatus. Pet. 5 linearia. Stam. 10 tubo 10-dentato, intrâ faucem antherifero. Stigm. 5 bifida. Bacca pomiformis fœta nucibus 5 ovato-compressis basi bivalvibus 1-spermis. Semina hilo arcuato.

1. *S. INDICUM* (Cav. l. c. t. 202. 203.) ζ in Philippinis Moluccis et Indiâ orient. Arbor foliis trifoliolatis, non palmatis sed imparipinnatis unijugis. Racemi axillares subpaniculati. *Sandoricum* Rumph. amb. 1. t. 61. Lam. ill. t. 350. Hantol Lam. dict. (v. s. sine fr.)

VII. *MELIA*. Lin. gen. n. 576.

Calyx parvus 5-fidus. Petala 5 oblongo-linearia patentia. Stam. 10. filam. coalitis in tubum apice 20-dentatum intus ad faucem antheriferum. Ovarium toro subelevato insidens. Styl. filiformis. Stigma capitatum 5-angularè. Drupa ovata, nucleo 5-sulcato 5 locul., loculis 1-spermis. Albumen carnosum. Cotyledones planæ foliaceæ. Radicula supera. — Arbores. foliis impari-pinnatis aut bipinnatis.

1. *M. AZEDARACH* (Lin. sp. 550.) foliis bipinnatis, foliolis levibus incisis subquinatis. ζ in Zeylonâ, Syriâ. Cav. diss. 7. p. 363. t. 207. Lam. ill. 372. Comm. hort. 1. t. 70. Folia autumnò decidua hyemem Europæ australis sub dio lætè fert. (v. v.)

2. *M. SEMPERVIRENS* (Swartz fl. ind. occ. 2. p. 737.) foliis bipinnatis, foliolis incisis rugosiusculis subnovenis nitidis, petiolo basi teretiusculo. ζ in sepius Jamaicæ. — bot. reg. t. 643. *M. Azedarach* β . Lin. sp. 550. Priore minor, florens jam biennis, folia tardiùs autumnò deponens et tepidarium perhyemem in nostris hortis requirens. (v. v.)

3. *M. COMPOSITA* (Willd. spec. 3. p. 559.) foliis subbipinnatis, in apice petiolorum simpliciter pinnatis, foliolis subintegris, pedunculis calycibus petalis subvelutinis. \S in Indiâ orient., ins. Timor. Culta Teneriffæ. (v. s.) — *M. dubia* Cav. diss. 7. p. 364. forsan diversa, sed non satis nota.
4. *M. ARGUTA*, foliis bipinnatis, foliolis oblongo-lanceolatis acuminatis argutè aristato-serratis, calycibus pedicellisque subvelutinis. \S in Moluccis. (v. s. sine fr. comm. à cl. Lambert.)
5. *M. AZADIRAGHTA* (Lin. sp. 550.) foliis pinnatis, petiolo tereti, foliolis dimidiato-ovatis basi inæqualibus acuminatis dentato-serratis. \S in Indiâ or. — Cav. diss. 7. p. 364. t. 208. — Burm. zeyl. t. 15. — Rheed. mal. 4. t. 52.
6. *M. BACCIFERA* (Roth nov. sp. p. 218.) foliis pinnatis, petiolo tereti, foliolis ovatis suprâ medium grossè obtusè serratis basi æqualibus. \S in Indiâ orientali.
7. *M. EXCELSA* (mal. misc. v. 1. n. 1. p. 12.) foliis pinnatis, foliolis integerrimis, paniculis coarctatis axillaribus folio paulò longioribus. \S in Palu-Pinang. Flores albi.

Trib. II. *TRICHILIEÆ*.

Fructus loculi 1-2-spermi. Semina exalbuminosa alâ destituta. Embryo inversus. Cotyledones crassissimæ.

VIII. TRICHILIA. Lin. gen. n. 528. — *Trichilia*, *Elcaja* et *Portesia* Juss.

Cal. 4-5 dentatus. Pet. 4-5 ovata aut suboblonga. Stam. 8-10, filam. nunc subdistinctis nunc in tubum apice intus antheriferum arcuè coalitis. Capsula 3-valvis, 3-locul. (rarius 2-valv., 2-locul.) valvis medio septiferis, loculis 1 2-spermis. Semina baccato-arillata exalbuminosa. Embryo inversus. Cot. 2 crassissimæ.

* *Foliis pinnatis.*

1. *T. HIRTA* (Lin. sp. 550.) foliis impari-pinnatis, foliolis 3-4-jugis ellipticis acuminatis glabris, racemis confertis, staminum filam. omninò in tubum coalitis. \S in Jamaicâ. Swartz obs. 171. — Sloan. hist. 2. t. 220. f. 1. (v. s.)
2. *T. SPONDIODES* (Swartz! fl. ind. occ. 730.) foliis impari-pinnatis, foliolis 7-10-jugis ovato-lanceolatis margine subhirsutis, racemis axillaribus, staminum filam. subdistinctis. \S in montosis Jamaicæ et Hispaniolæ. Jacq. h. schœnbr. 1. t. 102. — Sloan. hist. 2. t. 210. f. 2. 3. (v. s.)
3. *T. TOMENTOSA* (H. B. et Kunth nov. gen. am. 5. p. 215.) foliis impari-pinnatis, foliolis 12-15-jugis oblongis subacuminatis suprâ glabriusculis subtus tenuiter canescenti-tomentosis, paniculis axillaribus petiolo longioribus. \S in Andium Peruvicæ decliv. occid.
4. *T. QUADRIFUGA* (H. B. et Kunth nov. gen. am. 5. p. 215.) foliis impari-pinnatis, foliolis sub-4-jugis oblongis acuminatis glabris, paniculis axillaribus petiolo longioribus, filamentis internè hirsutis, capsulis subpyriformibus. \S in Amer. mer. ad ripam fl. Magdalenæ.
5. *T. EMETICA* (Vahl symb. 1. p. 31.) foliolis impari-pinnatis, foliolis 4-jugis ellipticis subtus villosis, floribus confertis, staminum filam. ad medium coalitis. \S in Arabiæ montibus Yemen. Elcaja Forsk descr. 127.
6. *T. HAVANENSIS* (Jacq. am. 129. t. 175. f. 38.) foliis impari-pinnatis, foliolis 2-3-jugis obovatis glabris, extimis majoribus, racemis cymosis axillaribus confertis petiolo brevioribus. \S ad Havanam in sylvis montosis et prope Xalapam H. B. et Kunth nov. gen. am. 5. p. 216. T. glabra Lin. syst. nat. 13. p. 294. (v. s.)
7. *T. PALLIDA* (Swartz! fl. ind. occ. 733.) foliis pinnatis, foliolis 3 jugis membranaceis ovato-lanceolatis acutis glabris, racemis axillaribus subterminali-

busque, floribus 8-andris, capsulis bivalvibus 5 in fruticetis montosis Hispaniolæ, Mexici. *Portesia ovata* Cav. diss. 7. p. 369. t. 215. Lam. ill. t. 302. f. 1. (v. s.)

8. *T. MOSCHATA* (Swartz! fl. ind. occ. 735.) foliis pinnatis, foliolis alternis ovatis acuminatis glabris, racemis axillaribus, floribus subdecandris monopetalis, capsulis submonospermis 5 in Jamaica bor. sylvis. (v. s.)

β. *odorata* (Andr. bot rept. t. 637) foliolis lanceolatis undulatis, floribus 4-petalis 5 culta in horto Sancti-Vincentii.

9. *T. TERMINALIS* (Jacq. amer. 130.) foliis pinnatis, foliolis 3-jugis lanceolatis obtusis nitidissimis, racemis paniculatis, lobis calycinis profundè sectis patentissimis, staminum filam. distinctis. 5 in Jamaica prope Kingston. Fruct. ign.

10. *T. GLANDULOSA* (Smith in Rees cycl. 36. n. 10.) foliis pinnatis, foliolis 5-7 elliptico-lanceolatis obtusè acuminatis, axillis venarum subtus hirsutis, floribus 8-andris, tubo staminum integro, stigmate depresso. 5 in Novâ-Hollandiâ ad Port-Jackson. *T. octandra* Sol. mss.

11. *T. SPECTABILIS* (Forst. prod. 188.) foliis pinnatis, foliolis obovatis, racemis axillaribus compositis. 5 in Novâ-Zelandiâ.

12. *T. ALLIACEA* (Forst. prod. n. 189.) foliis pinnatis, foliolis lanceolatis acutis, racemis axillaribus suprâ decompositis. 5 in ins. Namoka.

13. *T. HETEROPHYLLA* (Willd. sp. 2. p. 554.) foliis pinnatis ternatisque, foliolis ovatis acuminatis glabris, racemis axillaribus, floribus 8-andris. 5 in Madagascar. *Portesia mucronata* Cav. diss. 7. p. 370. t. 216. Lam. ill. t. 302. f. 2.

** *Foliis trifoliolatis.*

14. *T. TRIFOLIATA* (Lin. sp. 551.) foliis trifoliolatis, foliolis obovatis nitidis. 5 in Amer. merid. et in Curaçao locis siccis graminosis. — Jacq. amer. 129. t. 82. *T. Halesia* Læff. it. 188.

15. *T. NERVOSA* (Vahl symb. 1. p. 31.) foliis 3-foliolatis, foliolis ovatis. 5 in Javâ. *Melia Koatjapa* Burm. ind. 101. excl. syn.

*** *Foliis simplicibus. — Species dubiæ et forsân ad genera diversa referendæ.*

16. *T?* *SPINOSA* (Willd. sp. 2. p. 554.) foliis simplicibus ovatis emarginatis, ramis spinosis, bacca triloculari. 5 in Indiâ orientali. *Turraea virens* Hell. act. holm. 1788. p. 294. t. 10. f. 1. An potius *Limonæ* species?

17. *T?* *INERMIS* (Spreng. neue entd. 1. p. 285.) foliis simplicibus obovatis retusis coriaceis nitidis integerrimis, ramis inermibus, capsulâ 3-loculari. 5 in Brasiliâ. An *Turraæ* species?

18. *T?* *SCANDENS* (Lunan. h. jam. 2. p. 319.) foliis simplicibus lanceolatis ovatis margine revolutis, floribus spicatis 4-gyniis. 5 in Jamaica.

IX. EKEBERGIA. *Sparm. act. holm. 1779. p. 282. t. 9.*

Cal. 4-fidus. Pet. 4. Stam. 10? tubo brevissimo integro, antheris intrâ tubum subsessilibus. Stigma capitatum. Bacca globosa 5-sperma. Sem. ignota. — Huc relata ex summâ analogiâ cum *Trichiliâ* et *Guareâ*.

1. *E. CAPENSIS* (Sparm. l. c.) 5 ad cap. Bonæ-Spei. Arbor foliis impari-pinnatis 5-jugis, foliolis ellipticis utrinque acuminatis glabris. — Lam. ill. t. 358. *Trichilia Capensis* Pers. ench. 1. p. 468. (v. s.)

X. GUAREA. *Lin. mant. n. 1305.*

Cal. minimus 4-dentatus. Pet. 4, oblonga obtusa. Stam. 8, tubo apice dentato, intus ad faucem antherifero. Stigma capitatum. Capsula subglobosa, 4-loc., 4-valv. Semina solitaria arillata exalbuminosa. — Arbores foliis abruptè pinnatis.

1. *G. GRANDIFOLIA*, foliolis multijugis ovali-oblongis penninerviis, nervis lateralibus 10-15 subtus prominentibus, racemis elongatis. ζ in Guianâ Gallicâ et ins. Caribæis. — Plum. ic. t. 147. f. 2. Melia Guara Jacq. am 126. t. 176. f. 37. Trichilia Guara Lin. sp. 551. *G. trichilioïdes* Lin. mant. 228? excl. syn. Br. et Maregr. *G. macrophylla* Vahl? — Folia ampla, foliolis 8-9 poll. longis, infimis minoribus. Petala extus sericea 4-5 lin. longa. Stam. tubus apice integer. Fructus mihi ignotus, ex Jacq. subrotundus. (v. s.)
 2. *G. SWARTZII*, foliolis 2-4-jugis lanceolato-ovatis acuminatis penninerviis, nervis lateralibus 6-7 subtus prominalis, racemis elongatis. ζ in insulis Caribæis Elutheria Browne jam. 369. n. 7. — Sloan. hist. 2. t. 170. f. 1. Guarea trichilioïdes Swartz! obs. 146. (v. s.)
- N. B. G. trichilioïdes* Cav. videtur tertia species à duabus prioribus distincta ob folia multijuga et flores parvulos, et Jito arbor Pis. et Mareg. quarta etiam distinguenda ex foliis paucijugis fructuque turbinato. An hùc etiam referenda Plumea seu Plume-tree Lunan. hort. jam. 2. p. 77?
3. *G. BRACHYSTACHYA* (fl. mex. ic. ined.) foliolis 4-5-jugis ovalibus acutis, racemis axillaribus solitariis folio quadruplò brevioribus, fructibus globosis, ramulisque tuberculatis. ζ in Novâ Hispaniâ.
 4. *G. HUMILIS* (Bertero! in litt.) foliolis 2-3-jugis ovalibus acuminatis, racemis axillaribus solitariis folio quadruplò brevioribus, ramulis levibus. ζ in Porto-Ricco. *G. glabra* Vahl? (v. s. sine fr.)
 5. *G. RAMIFLORA* (Vent. choix. t. 41.) foliolis 2-jugis ovato-lanceolatis, racemis lateralibus secus ramos lateraliter ortis brevissimis, capsulis globosis. ζ in Porto-Ricco.

XI. HEYNEA. Roxb. in Sims bot. mag. t. 1738.

Cal. 5-dentatus. Pet. 5. Filam. 10 in tubum cylindricum apice antheriferum coalita. Styl. 1. Ovarium biloculare, loculis 2-ovulatis, ovulis intus adfixis. Capsula 2-valvis abortu 1-locul. 1-sperma. Semen arillatum apterum. Embryo exalbuminosus inversus. Cotyledones crassissimæ.

1. *H. TRIJUGA* (Roxb. in Sims l. c.) ζ in Napauliâ. Folia impari-pinnata trijuga. Paniculæ axillares longè pedunculatæ. Flores parvi albi. (v. s.)

Trib. III. CEDRELEÆ. Brown. gen. rem. 64.

Fructus loculi polyspermi. Semina sæpius in alam expansa. Albumen carnosum tenue. Embryo erectus. Cotyledones foliaceæ.

XII. CEDRELA. Lin. gen. n. 277. Cedrus. Mill.

Cal. minimus 5-dentatus. — Torus elevatus stipitiformis. Petala 5 basi lata approximata toro adnata. Stam. 5, filam. toro insertis distinctis brevibus, antheris oblongis. Styl. 1. Stigma capitatum. Capsula lignosa, 5-loc., 5-valvis. Semina ∞ compressa deorsum imbricata in alam membranaceam desinentia. Albumen carnosum. Embryo inversus. Cotyledones planæ foliaceæ. (Gaertn. fr. 2. p. 84. t. 95.) — Arbores foliis abruptè pinnatis multijugis, racemis axillaribus paniculatis.

1. *C. ODORATA* (Lin. sp. 289.) foliolis ovato-lanceolatis integris subsessilibus subtus concoloribus. ζ in ins. Caribæis et Barbadosibus. — Sloan. hist. 2. t. 220. f. 2. — Brown. jam. 158. t. 10. f. 1. — Lam. ill. t. 137. (v. s.)
2. *C. ANGUSTIFOLIA* (fl. mex. ic. ined.) foliolis oblongis acuminatis integris longè petiolatis subtus concoloribus. ζ in Novâ-Hispaniâ. (v. s.)
3. *C. TOONA* (Roxb. cor. 3. t. 238.) foliolis ovato-lanceolatis acuminatis subserratis subtus pallidè glaucis. ζ in Indiâ orient. Roxb. in Napauliâ, Wallich. — Willd. act. nat. cur. ber. 4. p. 198. Rœm. et Schult. syst. 5. p. 464. (v. s.)

†. *Species non satis notæ.*

4. *C. VELUTINA*, foliolis ovato-lanceolatis integris glabris subtus concoloribus, petiolis ramulisque pube brevissimâ velutinis 5. An *C. villosa* Roxb? (vidi specim. sicc. floribus fructibusq. destitutum olim in horto kewvensi cultam et à l'Heritiero subnomine *C. odorata* desiccatum.) †.
- 5? *C. ALTERNIFOLIA* (Stend. nom. phan. 170.) foliis alternis simplicibus cordato-ovatis acutis, fructu pentagono mucronato. 5 in agro Campechiano. *Cedrus alternifolia* Mill. dict. n. 3. A genere, imo ab ordine omnino diversa videtur. †.

XIII. SWIETENIA. *Lin. gen. n. 575.*

Cal. minimus 4-5-fidus deciduus. Petala 4-5. Stam. 8-10, filam. coalitis in tubum apice dentatum intus antheriferum. Styl. 1. Stigma capitatum. Capsula oviformis lignosa 5 loc. ∞ -sperma, valvis à basi aut ab apice dehiscentibus, margine appositis ad angulos placentæ centralis 5-gonæ. Semina deorsum imbricata, in alam expansa. Albumen carnosum. Embryo rectus. Cotyl. planæ foliaceæ. (Gaertn. fr. 2. p. 89. t. 96.) — Arbores foliis alternis abruptè pinnatis paucijugis.

1. *S. MAHOGONI* (Lin. sp. 271.) foliis sub-4 jugis, foliolis ovato-lanceolatis basi inæqualibus apice acuminatis, racemis axillaribus paniculatis. 5 in Amer. calidioris petrosis. Cav. diss. 7. p. 365. t. 209. *Cedrus Mahogoni* Mill. dict. n. 2. — Cat. car. 2. t. 81. — in icone Gaertneri et fl. mex., capsulæ ab apice dehiscentes indicantur. An species 2 hic confusæ? An dehiscentia varia? (v. s.)
2. *S. FEBRIFUGA* (Roxb. cor. 1. p. 18. t. 7.) foliis sub-4-jugis ovalibus obtusis emarginatisve basi subobliquis, racemis ex axillis supremorum foliorum abortientium ortis paniculam terminalem constituentibus. 5 in montibus Indiæ orientalis. *S. soyimida* Dunc. tent. Edinb. 1794. Valvulæ fr. ab apice dehiscentes.
3. *S. SENEGALENSIS* (Desrouss. in Lam. dict. 3. p. 679.) foliis 3-jugis, foliolis ovali-oblongis coriaceis obtusiusculis, paniculis terminalibus, floribus 8-andris, fructibus globosis 4-valvibus. 5 in Senegalâ.

XIV. CHLOROXYLON.

Cal. minimus 5-fidus. Pet. 5. Stam. 10, filam. imâ basi coalito-appendiculatis, è dorso urceoli exsertis, ceterum subulatis liberis radiatim patentibus. Capsula 3-valvis, 3-locul., valvis medio septiferis. Semina in loculo quoque 4 in alam desinentia, valvularum septo adfixa.

1. *C. SWIETENIA*. 5 in Indiâ Orient. *Swietenia chloroxylon* Roxb. cor. 1. p. 46. t. 64. — Folia abruptè pinnata multijuga. foliolis glaucis basi valdè inæqualibus ovatis subrhombeis obtusis. Paniculæ terminales. (v. s.)

XV. FLINDERSIA. *Brown gen. rem. p. 63. t. 1.*

Cal. 5 fidus brevis. Petala 5. Stam. 10 è dorso urceoli ex eorum basibus dilatatis facti exserta, filam. subulatis patentibus alternis sterilibus. Capsula 5-loc., 5-valvis, 5-partibilis, segmentis singulis divisis dissepimento longitudinali demum libero utrinquè 2-spermo. Sèmina erecta apice alata. Albumen o. Cotyledones transversæ crassæ foliaceæ. — Arbores foliis impari-pinnatis, fructibus echinatis.

1. *F. AUSTRALIS* (Brown l. c.) foliis impari-pinnatis 1-3-jugis, floribus paniculatis, fructibus ovatis utrinquè obtusissimis. 5 in Novæ-Hollandiæ orâ orient. Folia pellucido-punctata ut in *Aufantiaceis*.
2. *F. AMBOINENSIS* (Poir. suppl. 4. p. 650.) foliis impari-pinnatis 3-7-jugis, floribus subsolitariis?, fructibus ovato-oblongis utrinquè attenuatis. 5 in ins. Hitoe et Cerama. Rumph. amb. 3. p. 201. t. 129.

XVI. CARAPA. *Aubl. guian. suppl. p. 33. t. 387.* — *Xylocarpus. Schreb. gen. n. 646.* — *Persoonia Willd. sp. 2. p. 331.*

Cal. coriaceus 4-5-lobus. Petala 4-5 coriacea. Stam. 8-10, filam. coalitis in tubum apice dentatum intus ad faucem antheriferum. Stylus brevis. Stigma latum truncatum margine sulcatum. Drupa sicca globosa intus lignosa 4-5-sulcata, 4-5-valvis, nucleis 1-spermis fungosis. Semina crassa exalbuminosa. — Arbores foliis abrupte pinnatis coriaceis. Staminibus et habitu ad Meliaceas, semine ad Guttiferas accedit.

1. *C. GUIANENSIS* (Aubl. l. c.) foliolis 8-10-jugis alternis oppositisve elliptico-oblongis acuminatis coriaceis nitidis. \S in sylvis Guianæ. Lam. ill. t. 301. *Persoonia guareoides* Willd. sp. 2. p. 331. Oleum è seminibus elicitur. (v. s.)
2. *C? PROCERA*, foliolis 4-5-jugis oblongis basi cuneatis obtusis coriaceis, floribus 5-petalis decandris. \S in ins. Caribæis. *Trichilia procera* Forsyth! ined. in herb. L'her., et ideò ad Carapam ex habitus analogià tantum relata. (v. s. sine fructu.)
3. *C. MOLUCCENSIS* (Lam. dict. 1. p. 621.) foliolis subtrijugis oppositis ovatis acutis. \S in insulis Moluccanis. *Granatum littoreum* Rumph. amb. 3. p. 92. t. 61. *C. indica* Juss. dict. sc. nat. 7. p. 31. *Xylocarpus Granatum* Koen. nat. turf. 20. p. 2.

Ordo XLV. AMPELIDEÆ. *H. B. et Kunth nov. gen. am. 5. p. 223.* — *Vites. Juss. gen. 267.* — *Sarmentaceæ. Vent. tabl. 3. p. 167.* — *Viniferæ. Juss. mem. mus. 3. p. 144.*

Calyx parvus margine integer aut subdentatus. Petala 4-5 dentibus calycinis alterna, disco ovarium ambiente extus inserta et idè inter hypogyna et perigyna ambigua, basi latiora, rarius in corollam 1-petalam coalita, per æstivationem intus apice inflexa subvalvata. Stamina tot quot petala, ante petala disco inserta, interdum abortu sterilia; filam. libera aut basi subcoalita; antheræ ovatæ birimosæ dorso insertæ oscillatoriæ. Ovarium globosum liberum. Styl. 1 brevissimus aut subnullus. Stigma simplex. Bacca globosa, junior bi-ocularis, loculis 2-spermis, adulta sæpè dissepimentis evanidis 1-ocularis, aquosa aut subcarnosa, epicarpio non secedente. Semina 4-5 aut abortu pauciora (imò interdum nulla) axi centrali funiculo brevi adfixa erecta, ossea. Albumen carnosum durum. Embryo erectus dimidiò albumine brevior, radicula teretiusculà inferà, cotyledonibus lanceolatis hinc subcarinatis indè planis. — Frutices sarmentosi scandentes. Folia basi stipulacea, inferiora opposita, superiora alterna pedunculis opposita petiolata, simplicia, compositave. Pedunculi racemosi, interdum florum abortu in cirrhos evadentes. Flores parvi virescentes.

Ordo accedit hinc inter Thalamifloras ad Geraniaceas et Meliaceas, illinc inter Calycifloras ad Araliaceas Caprifoliaceasve et speciatim ad Hederam.

Trib. I. *VINIFERÆ* seu *SARMENTACEÆ*.

Corolla polypetala. Stamina petalis opposita. Fructus et semina ut in char. ordinis. — Pedunculi sæpè cirrhosi.

I. *CISSUS. Lin. gen. n. 147.*

Cal. subinteger. Pet. 4 ab apice ad basim more solito abscedentia. Stam. 4. Ovarium 4-locul. Bacca 1-4-sperma.

* *Foliis simplicibus.*

1. *C. VITIGINEA* (Lin. sp. 170.) foliis cordato-subrotundis sublobatis angulato-repandis subtus ferrugineo-pubescentibus. ♀ in Indiâ. — Pluk. mant. 27. t. 337. f. 4. Vahl symb. 3. p. 18. Baccæ pyriformes, nigre, 1-2-sperme.
 ♀? *Cochinchinensis*, foliis glabris denticulatis 3-lobis, baccis subrotundis.
 ♀ *C. vitiginea* Lonr. coch. 1. p. 106.
2. *C. REPANDA* (Vahl symb. 3. p. 18.) foliis cordatis integris sublobatis repandis adultis utrinquè glabris. ♀ in Indiâ orient. Rami jun. tomentosi, adulti glabri. Baccæ pyriformes.
3. *C. ADNATA* (Roxb. fl. ind. 1. p. 423.) foliis rotundato-cordatis acuminatis setaceo-serratis supernè glabris subtus ramulisque teretibus velutinis, stipulis

- ovato-orbiculatis adnatis centro gibbis margine-scariosis, floribus nutantibus. ζ in Indiâ orient. prope Dacca. Bacca nigra Pisi magnitudine.
4. *C. LATIFOLIA* (Vahl symb. 3. p. 18. excl. syn. Lam.) foliis cordatis ovatis acuminatis setaceo-serratis supernè glabris subtùs ferrugineo-villosis, ramis tetragonis, stipulis oblongis. ζ in Indiæ orient. sylvis. — Rumph. amb. 5. t. 164. f. 1. (v. s.)
 5. *C. GLAUCA* (Roxb. fl. ind. 1. p. 425.) foliis cordatis acuminatis interdum sublobatis setaceo-serratis utrinquè pétiolis pedunculisque glabris, stipulis lato-sulcatis obtusis, cymis decompositis, baccis 1-spermis. ζ in Madagascar et Indiâ orient. Rheed. mal. 7. t. 11. *C. latifolia* Lam. dict. 1. p. 30. non Vahl.
 6. *C. INDICA* (Rottl. et Willd. nov. act. nat. cur. 4. p. 183.) foliis subrotundo-cordatis acuminatis setaceo-serratis subtùs pubescentibus, ramis teretiusculis petiolisque tomentosis. ζ in Indiâ orient.
 7. *C. CORDIFOLIA* (Lin. sp. 170.) foliis cordatis integerrimis subtùs pubescentibus, pedunculis 3-fidis dichotomis. ζ in Americâ calid. — Plum. ic. t. 259. f. 3. Baccæ cæruleæ 1-spermæ.
 8. *C. ROTUNDIFOLIA* (Vahl symb. 3. p. 19.) foliis cordato-subrotundis glabris serratis, umbellis simplicibus. ζ *Sælanthus rotundifolius* Forsk. descr. ic. t. 4. Baccæ 1-spermæ.
 9. *C. CÆSIA* (Afz. rem. guin. ex Spreng. neue entd. 3. p. 234.) foliis cordatis angulato-sinuatis setaceo-dentatis utrinquè rugosis scabris subtùs subtomentosis sarmentis, teretibus glauco-pruinosis. ζ in Guineâ. Cirrhi bibracteati.
 10. *C. REPENS* (Lam. dict. 1. p. 31.) foliis cordato-ovatis subdentatis ramisque glabris, floribus umbellatis, caule repente. — in Malabariâ. — Rheed. mal. 7. t. 48. *C. cordata* Roxb. fl. ind. 1. p. 425. Caules succulenti. Baccæ rubellæ 1-spermæ. Stipulæ ovals. Flores atro-purpurei.
 11. *C. PUNCTICULOSA* (Rich act. soc. h. nat. par. 106.) foliis lato-cordatis obtusissimis, acumine abrupto obtuso, serris remotis setaceis, ramis punctatis. ζ in Cayennâ.
 12. *C. UVIFERA* (Afz. rem. guin. 69. ex Spreng. neue entd. 3. p. 235.) foliis subpeltato-cordatis integerrimis glabris basi maculatis, sarmentis teretibus glabris. ζ in Guineâ. Drupæ? pulposæ.
 13. *C. TILIACRA* (H. B. et Kunth nov. gen. am. 5. p. 222.) glabra, foliis subrotundo-ovatis cordatis argutè dentatis subcoriaceis, ramis tetragonis, cymis dichotomis. ζ prope urbem Mexici.
 14. *C. QUADRANGULARIS* (Lin. mant. 39.) foliis subcordato-ovatis serratis glabris carnosis, caule tetragono alato. \mathcal{L} ab Arabiâ ad Cochinchinam. Radix tuberosa. — Rumph. 5. t. 44. *Sælanthus quadragonus* Forsk. descr. 33. ic. t. 2. Roxb. ind. 1. p. 426. Baccæ rubræ 1-spermæ magnitudine Pisi.
 15. *C. COMPRESSICAULIS* (fl. per. 1. p. 64. t. 100.) foliis cordatis acutè serratis pubescentibus, floribus umbellatis, pedunculis partialibus dichotomis, caule tetragono compresso. — in ruderatis prope Chancay. Bacca nigro-purpurea.
 16. *C. JAVANA*, foliis subcordatis ovato-lanceolatis acuminatis glabris subserratis, serraturis setaceis adpressis, ramulis subteretibus sulcatis, pedunculis trifidis subcorymbosis. — in Javâ. (v. s. comm. ab. an. Coulon.)
 17. *C. PENTAGONA* (Roxb. fl. ind. 1. p. 426.) foliis cordatis rariter sublobatis serrulatis acuminatis lævibus, ramulis lucidis pentagonis, angulis rotundatis, lateribus sulcatis. ζ in sylvis Chittagong Indiæ orient. Cæc. ign.
 18. *C. SICYOIDES* (Lin. sp. 170.) foliis cordatis ovatis glabris crassiusculis setaceo-serratis, serraturis adpressis, ramulis teretibus. \mathcal{L} in Jamaicâ et Guadalupe. Sloan. hist. 1. t. 144. f. 1. Lam. ill. t. 84. f. 1. (v. s.)
 19. *C. SMILACINA* (H. B. et Kunth nov. gen. am. 5. p. 224.) foliis ovato-oblongis subacuminatis basi obliquè truncatis suprâ glabris, subtùs ramisque sulcato-

- angulatis hirtell^{is}, cymis pedunculisque glabris. ζ ad ripas umbrosas fluminis Magdalenæ. C. sicyoides. Poir. suppl. 1. p. 104 ex H. B. et K.
20. C. OVATA (Lam. ill. n. 1619.) foliis ovatis acuminatis utriusque glabris serrato-serratis, serraturis paucis adpressis, ramulis teretibus. ζ in Guadalupâ. Badier! Irsiola Browne jam. t. 4. f. 1. 2. ex Lam. Valdè accedit ad sicyoidem sed folia ovata nec cordata. An C. smilacina Willd. enum. 163? sed folia mea non cordata. (v. s.)
21. C. OBSCURA, foliis ovato-lanceolatis acuminatis glabris, serraturis paucis aristatis adpressis. — in Americâ merid? Paniculæ aliæ magnæ ramosissimæ steriles, aliæ parvæ fertiles. An forsân hûc revocanda C. ovata Rich. act. soc. h. nat. par. p. 105. non Lam. in Cayennâ indigena? (v. s. in h. L'her.)
22. C. FULIGINEA (H. B. et Kunth nov. gen. am. 5. p. 224.) fuligineo-ferruginea foliis subrotundo-ovatis profundè cordatis remotè denticulatis suprâ hirtellis, subtus molliter tomentosis, infer. 3-5-lobis, super. integris, ramulis teretibus pedunculisque hirsuto-tomentosis. ζ ad ripam fl. Magdalenæ.
23. C. ANTARCTICA (Vent. choix t. 21.) foliis ovatis subcordatisve laxè serratis glabrinusculis, nervis basi glandulosis, petiolis ramisque ferrugineis suprâ pubescentibus. ζ in Novâ-Hollandiâ. C. glandulosa Poir. suppl. p. 105. Vitis Kanguruh hortul. C. Baudiniana Brouss. (v. v.)
24. C. UMBELLATA (Lour. coch. 1. p. 106.) foliis ovatis glabris integerrimis, umbellis compositis, corollis intus lanatis. ζ ad Cantonem Sinarum.
25. C. CANESCENS (Lam. ill. n. 1620.) foliis ovato-oblongis inæqualiter cordatis dentatis subtomentosis canescentibus, ramis angulatis, pedunculis cymisque villosotomentos. ζ in Peruvîâ. H. B. et Kunth nov. gen. am. 5. p. 223.
26. C. UMBROSA (H. B. et Kunth nov. gen. am. 5. p. 223) foliis inæquilatero-ovatis obliquè cordatis argutè denticulatis, suprâ pilosiusculis, subtus tomentoso-pubescentibus et canescentibus, ramis subtetragonis hirtellis, pedunculis cymisque glabrinusculis. ζ in ripis umbrosis fluminis Magdalenæ.
27. C. GLANDULOSA (Gmel. syst. 256.) foliis ovatis acutis serrato-dentatis car-nosis, pedicellis et calycis basi glandulosis. ζ in Arabiâ. Sælanthus glandulosus Forsk. descr. 34. Radix tuberosa. Caules teretes. An eadem C. glandulosa Horn. cat. h. hafn. 1. p. 143 pedicellis calycibusque hispidis.
28. C. PRODUCTA (Afz. rem. guin. 63. ex Spreng. neue entd. 3. p. 234.) foliis ovato-oblongis acuminatis subsetaceo-denticulatis, floribus paniculatis, samentis longissimis striatis compressis. — in Guineâ.
29. C. TUBEROSA (fl. mex. ic. ined.) foliis obovatis glabris grossè serratis aut trifidis, lobis incis, radice tuberosâ. — in Novâ-Hispaniâ. Accedit ad C. sicyoidem et ovatam. An varietas?
30. C. MICRANTHA (Poir! suppl. 1. p. 105.) foliis glabris ovatis subtrilobis, lobis subserratis acutis, terminali longiore, pedunculis apice 5-fidis umbellatis petioli longitudine. — in Sancto-Domingo. (v. s.)
31. C. RUGOSA, foliis cordatis trifidis 5-fidiave rugosis, floribus racemosis. ζ in Jamaicâ montibus. C. nov. sp. Lunan. hort. jam. 2. p. 216. Baccæ nigræ majores 4-spermæ. Folia ferè Vitis viniferæ. †.
32. C. TOMENTOSA (Lam. ill. n. 1613.) foliis subpentagonis obtusè dentatis suprâ glabris subtus tomentoso-ferrugineis, petiolis compressis. ζ in ins. Borboniâ. Fl. et fr. ignoti.
33. C. ANGULATA (Lam. ill. n. 1614.) foliis ovatis subpentagonis angulato-lobatis subtus tomentos, pedunculis planis folio longioribus. ζ in Indiâ orient. Baccæ nigro-purpureæ.
34. C. CAPENSIS (Willd. sp. 1. p. 655.) foliis subcordatis 5-angularibus dentatis subtus ferrugineo-tomentosis, floribus subcapitatis. ζ ad cap. Bonæ-Spei. Vitis Capensis Thunb. fl. cap. 2. p. 105.
35. C. TRILOBATA (Lam. dict. 1. p. 31.) foliis trilobatis 3-sectisve, lobis ovatis

acutis dentatis subcarnosis, ramis teretibus. ζ in Malabariâ. — Rheed. mal. 5. t. 45.†

†† *Folii 3-foliolatis.*

36. *C. ACIDA* (Lin. sp. 170.) foliis 3-foliolatis carnosis glabris, foliolis obovato-cuneatis apice dentatis basi integerrimis. ζ in Americâ calid. Jacq. schomb. 1. t. 33. — Plum. ed. Burm. t. 259. f. 3. *Sicyos angulata* Lin. sp. ed. 1. p. 1013. (v. v.)
37. *C. SETOSA* (Roxb. fl. ind. 1. p. 428.) foliis 3-rariter-5-foliolatis sessilibus carnosis glabris, foliolis ovalibus undulatis grossè et inæqualiter setaceo-serratis, stipulis cordatis, caulibus teretibus setoso-glandulosis. \mathcal{Z} in sylvis et sepibus Indiæ or. ad Rajamundree Circar. Baccæ 1-spermæ, radiceæque acidissimæ.
38. *C. CARNOSA* (Lam. dict. 1. p. 31.) foliis 3-foliolatis carnosis glabris, foliolis ovatis obtusis serratis, ramis petiolisque teretibus. ζ in Indiâ orient. Rumph. amb. 5. t. 166. f. 2. Rheed. mal. 7. t. g. — Roxb. fl. ind. 1. p. 427. Baccæ nigre 4-spermæ.
39. *C. TRIFOLIATA* (Jacq. am. 23.) foliis 3-foliolatis carnosis glabris, foliolis obovatis serrato-incisis acutis, ramis subulatis. ζ in sepibus Jamaicæ, Surinami. Swartz obs. p. 50. Sloan. jam. t. 142. f. 5. 6? An differt à *C. acida*? *C. trifoliata* Lour. altera spec. videtur!
40. *C. CAUSTICA* (Tuss. fl. aut. t. 16.) foliis 3-foliolatis, foliolis ovatis obtusis, ramis teretibus geniculatis succulentis, petiolis canaliculatis. ζ in ins. Caribæis. Flores sanguinei corymbosi.
41. *C. SALUTARIS* (H. B. et Kunth nov. gen. am. 5. p. 225.) foliis 3-foliolatis, foliolis oblongis argutè serratis pellucido-punctatis suprâ hirtellis subtus ferrugineo hirtis, ramis teretibus pedunculisque hirtellis striatis. ζ in aridis Novæ-Andalusie prope Quetepe et Cumana. Radix contra hydropysia utilis.
42. *C. OBOVATA* (Vahl symb. 3. p. 19.) foliis 3-foliolatis glabris membranaceis, foliolis obovatis integerrimis mucronatis, pedunculis 3-chotomis folio longioribus. — in ins. Sanctæ-Crucis.
43. *C. PAUCIFLORA*, foliis 3-foliolatis glabris rigidulis, foliolis obovatis subintegris basi cuneatis apice obtusis, pedunculis paucifloris folio brevioribus. ζ ad cap. Bonæ-Spei. Burchell! cat. n. 3009. (v. s.)
44. *C. TERNATA* (Gmel. syst. 1. p. 256.) foliis sessilibus 3-foliolatis rigidis glabris, foliolis ovato-cordatis acutis serratis, ramis teretibus. ζ in Arabiæ arvis. *Sælanthus ternatus*! Forsk. descr. 35. Petiolus communis nullus!
45. *C. LUCIDA* (Poir. suppl. 1. p. 106.) foliis 3-foliolatis lucidis glabris, foliolis ovatis denticulato-setaceis, ramis compresso-tetragonis subulatis glandulosis. ζ in Cayennâ. Aff. *C. alata*.
46. *C. QUADRIALATA* (H. B. et Kunth l. c.) foliis trifoliolatis glabris, foliolis obtusis crenato-denticulatis pellucido-punctatis, intermedio ovato basi acuminato, lateralibus inæquilateris, ramis tetragonis quadrialatis, pedunculo cymisque pilosiusculis. ζ in Islâ Panumanâ mission. Orinocens.
47. *C. MICROCARPA* (Vahl ecl. 1. p. 16.) foliis 3-foliolatis membranaceis glabris, foliolis oblongis serratis mucronulatis, ramis angulatis. ζ in insulis Caribæis. — Plum. ic. ed. Burm. t. 259. f. 4. Baccæ oblongæ.
48. *C. ACUTIFOLIA* (Poir. suppl. 1. p. 106.) foliis 3-foliolatis membranaceis glabris, foliolis ovatis dentato-sublobatis, lobis acutis, ramis angulosis subcompressis. ζ in Indiâ orient.
49. *C. EROSA* (Rich. act. soc. h. nat. par. 106.) foliis 3-foliolatis membranaceis glabris, foliolis oblongo-ovatis basi subacutis eroso-subdentatis, petiolo submarginato, cymis longè pedunculatis. — in Cayennâ.
50. *C. TIMORIENSIS*, foliis 3-foliolatis membranaceis glabris, foliolis petiolo-latis ovatis acutis latè dentatis, paniculis laxis 3-chotomis. ζ in Timor. (v. s.)

51. *C. ALATA* (Jacq. am. 23. t. 182. f. 10.) foliis 3-foliolatis hirsutis, foliolis ovato-cordatis acutis serratis, ramis membranaceo-angulatis. ζ in Jamaicâ et Carthagenâ. Scandit in arbores. — Sloan. hist. 1. p. 233. t. 144. *C. trifoliata* Lin. sp. 170?
52. *C. PUBESCENS* (H. B. et Kunth nov. gen. am. 5. p. 226.) foliis 3-foliolatis supra puberulis subtus canescenti-pubescentibus, foliolis acutis argutè denticulatis, intermedio majore petiolato ovato-oblongo basi cuneato-angustato, ramis quadrangularibus striatis hirtis. ζ Frequens ad ripam fl. Magdalenzæ inter Mompoz et Buenavista. Fl. et fruct. ign.
53. *C. RHOMBIFOLIA* (Vahl ecl. 2. p. 10.) foliis 3-foliolatis pubescentibus, foliolis serrulatis lateralibus semi-cordatis, medio rhombeo-ovato, ramis angulatis. ζ in insulâ Trinitatis.
54. *C. CINEREA* (Lam. ill. n. 1624.) foliis 3-foliolatis crassiusculis pubescentibus, foliolis dentatis ovatis, lateralibus subcordatis, ramis angulosis, petiolis teretibus. ζ in Indiâ orient.
55. *C. CIRRHOSA* (Pers. ench. 1. p. 142.) foliis 3-foliolatis carnosis villosis, foliolis ovatis serratis, ramis petiolisque compressis, baccis hirsutis. ζ ad cap. Bonæ-Spei. *Vitis cirrhosa* Thunb. fl. cap. 2. p. 106. Variat tota glabra ex cl. Thunb. An dist. species?
56. *C. CRENATA* (Vahl symb. 3. p. 19.) foliis 3-foliolatis junioribus villosis, foliolis subrotundis crenatis. ζ in Indiâ orient. *Vitis trifolia* Lin. sp. 253. — Rumph. amb. 5. t. 166. f. 2. *C. obtusifolia* Lam. dict. 1. p. 31?
57. *C. ANGUSTIFOLIA* (Roxb. fl. ind. 1. p. 427.) dioica, foliis 3-foliolatis utrinquè lævibus, foliolis lanceolatis serratis, stipulis ovatis acutis, cymis petiolo brevioribus, caulibus compressissimis. ζ in Sumatrâ. Baccæ albæ sphaericæ 1-2-spermæ, Pisi magnitudine.
- *** *Foliis 5-foliolatis palmatis.*
58. *C. QUINATA* (Ait. h. kew. ed. 2. v. 1. p. 260.) foliis 5-foliolatis, foliolis obovato-cuneiformibus supernè serratis. ζ ad cap. B.-Spei. †
59. *C. PENTAPHYLLA* (Willd. sp. 1. p. 659.) foliis palmato-5-foliolatis glabris, foliolis ovatis serratis acuminatis. ζ in Japoniâ. *Vitis pentaphylla* Thunb. fl. jap. 105.
60. *C. STRIATA* (fl. per. 1. p. 64. t. 100.) foliis palmato-5-foliolatis glabris, foliolis sessilibus oblongo-lanceolatis serratis, caule striato flexuoso, cymis coarctatis. ζ in Peruvîâ.
61. *C. MEXICANA* (fl. mex. ic. ined.) foliis palmato-5-foliolatis glabris, foliolis petiolulatis oblongo-lanceolatis serratis, caule subtragono. ζ in Mexico.
62. *C. PALMATA* (Poir. suppl. 1. p. 107.) foliis palmato-5-foliolatis glabris, foliolis sessilibus lanceolatis angustis dentato-setaceis, ramulis tetragonis. ζ in ins. Franciæ.
63. *C. GRANULOSA* (fl. per. 1. t. 101.) foliis palmato-5-foliolatis glabris, foliolis obovatis dentatis, caule angulato granuloso, cymis patentibus. ζ in Peruvîæ præruptis.
64. *C. DIGITATA* (Lam. ill. n. 1627.) foliis palmato-5-foliolatis glabris, summis 3-foliolatis, foliolis ovatis serratis, ramis teretibus. — in Arabiâ. *Sælæthus digitatus* Forsk. descr. 35. ic. t. 3.
65. *C. DIVERSIFOLIA*, foliis glabris, radicalibus simplicibus, mediis 3-foliolatis, summis 5-foliolatis, foliolis acuminatis attenuatis acutè crenatis, petiolis lineâ pubescente. ζ Patr. et flor. ign. *C. heterophylla* Link enum. 1. p. 143. non Poir.
66. *C. FOEMINÆA* (Roxb. fl. ind. 1. p. 428.) foliis digitatis lucidis, foliolis petiolatis lato-lanceolatis integris subcrenatisve, stylo nullo, petalis stigmatibusque 4-5. ζ in sylvis montosis Indiæ orientalis.
67. *C. ELONGATA* (Roxb. fl. ind. 1. p. 429.) foliis digitatis lævibus, foliolis

petiolulatis oblongis acutis remotè serratis, stipulis bracteisque cordatis, cymis trichotomis, ramulis carnosis lævibus subcompressis — in montibus Coromandelæ et Bengali. Baccæ magnitudine Cerasi, rotundato-turbinatæ, nigrae.

68. *C. AURICULATA* (Roxb. fl. ind. 1. p. 430.) foliis 5-foliolatis, foliolis petiolulatis oblongis acutis obtusè serratis supernè lævibus subtùs villosis, stipulis auriculæformibus, ramulis teretibus villosis. 5 in Mysore. Caulis arbores excelsissimas scandit. Baccæ rotundæ læves magnit. et colore Cerasi 1-spermæ.

*** *Foliis pedatis 5-9-foliolatis, foliolo medio solitario, lateralibus geminis ternisve.*

69. *C. JAPONICA* (Willd. sp. 1. p. 659.) foliis pedato-5-foliolatis glabris, foliolis ovalibus aristato-serratis, lateralibus obtusis, pedunculis folio longioribus. 2 in Japoniâ.

70. *C. NAPAULENSIS*, foliis pedato-5-foliolatis glabris, foliolis ovalibus acutis basi attenuatis aristato-serratis, pedunculis petiolo brevioribus. 2 in Napauliâ. Wallich. (v. s.)

71. *C. CORIACEA*, foliis pedato-5-foliolatis glabris subcoriaceis, foliolis ovalibus utrinquè subattenuatis latè et distanter crenatis, pedunculis petiolo brevioribus. 5 in insulâ Timor. Variat foliolis angustioribus et latioribus. (v. s.)

72. *C. LANCEOLARIA* (Roxb. fl. ind. 1. p. 430.) foliis pedato-5-foliolatis glabrisculis subcoriaceis, foliolis lanceolatis acuminatis irregulariter subser-ratis, cymis pedunculatis petioli ferè longitudine. 5 in Indiâ orient. *C. dioica* Roxb! in h. Lamb. — Rheed. mal. 7. p. 15. t. 8. Flores dioici, feminei ign. (v. s. ♂)

73. *C. SERRULATA* (Roxb. fl. ind. 1. p. 432.) foliis pedato-5-foliolatis, foliolis oblongis serrulatis, caule lævi. 5 in Chittagong Indiæ orient. Cæt. ign.

74. *C. HETEROPHYLLA* (Poir. suppl. 1. p. 107.) foliis pedato-5-foliolatis glabris, foliolis superioribus rhomboideis crenatis, inferioribus obliquis obtusis, caule fistuloso striato. 5 in Javâ.

75. *C. PEDATA* (Lam. dict. 1. p. 31.) foliis pedato-7-9-foliolatis subtùs pubescenti-tomentosis, foliolis lanceolatis acuminatis serratis, ramis petiolisque incanis. 5 in Indiâ orient. Rheed. mal. 7. t. 10? sed flores 5-fidi dicuntur. Roxb. fl. ind. 1. p. 431. Baccæ 4-spermæ albæ depresso-4-lobæ. Stipulæ cordatæ.

**** *Foliis pinnatis bipinnatisve, foliolis oppositis.*

76. *C. PINNATA* (Russ. besch. alep. ex Roem. et Schult. syst. 3. p. 317.) foliis pinnatis 3-foliolatisque glabris membranaceis, foliolis ovatis incis. — Aleppo. †.

77. *C. OBLIQUA* (fl. per. 1. p. 65. t. 101. f. 6.) foliis simpliciter pinnatis pubescentibus, foliolis 2-3 obliquè cordatis. — in Pernviâ.

78. *C. MAPPIA* (Lam. ill. n. 1631.) foliis subbipinnatis lævibus subcarnosis, foliolis ovatis integerrimis. — in insulâ Franciæ.

79. *C. ORIENTALIS* (Lam. ill. p. 332. t. 84. f. 2.) foliis bipinnatis glabris, foliolis ovatis serratis. 5 in Oriente. (v. s.)

80. *C. CONNIVENS* (Lam. ill. n. 1630.) foliis subbipinnatis glabris, foliolis ovatis obtusiusculis subdentatis, petalis conniventibus 5 in Madagascar. A C. Orientali præcipuè differt foliolis minoribus paucioribus.

II. AMPELOPSIS. Michx. fl. bor. am. 1. p. 159.

Cal. subinteger. Petala 5 (ut in Vite) ab apice ad basim (more Cissi) abscedentia. Stam. 5. Styl. 1. Stigma capitatum. Ovarium disco non immersum, 2-4-spermum (Kunth nov. gen. am. 5. p. 222.) — Genus medium inter Cissum et Vitem.

* *Foliis simplicibus.*

1. *A. CORDATA* (Michx. l. c.) foliis cordatis acutis dentatis subtrilobis, nervis subtus villosis, racemis duplicato bifidis. 5 in Americæ borealis dumetis. *Cissus ampelopsis* Pers. ench. 1. p. 142. *Vitis indivisa* Willd. baumz. 538. (v. v.)
2. *A. BOTRIA*, foliis cordatis crenatis 3-5-lobis tomentosis, racemis subdivisis. 5 ad littora Oræ Zanguebariæ, in Africâ. *Botria Africana* Lour. coch. 1. p. 191. Flores rubescentes. Baccæ nigre edules.

** *Foliis 3-5-foliolatis palmatis.*

3. *A. HEDERACEA* (Michx. fl. 1. p. 160.) foliis palmato 3-5-foliolatis utrinque glabris, foliolis petiolatis oblongis acuminatis mucronato-dentatis, racemis dichotomo-corymbosis. 5 à Canadâ ad Virginiam. *Hedera quinquefolia* Linn. sp. 292. *Vitis quinquefolia* Lam. ill. n. 2815. *Cissus quinquefolia* Pursh. *Vitis hederacea* Willd. Caulis radicans scandens. (v. v.)
4. *A. HIRSUTA* (Donn cat. 166.) foliis palmato 3-5-foliolatis utrinque pubescentibus, foliolis ovatis acuminatis grosse dentatis. 5 in montibus Alleghany. An prioris var? ex Pursh fl. 1. p. 170.
5. *A. ? TERNATA*, foliis trifoliolatis subtus cinereo-tomentosis, foliolis oblongo-ovalibus mucronatis duplicato-serratis, lateralibus obliquis, caule cum cirrhis tomentoso. 5 in Indiâ orient. *Vitis ternata* Roth in Rœm. et Schult. syst. 5. p. 319.

*** *Foliis pinnatis bipinnatisve.*

6. *A. PINNATA* (Rœm. et Schult. syst. 5. p. 322.) foliis pinnatis glabris, foliolis quinis dentato-serratis. Patr. ign. *Vitis pinnata* Vahl symb. 3. p. 43.
7. *A. BIPINNATA* (Michx. fl. bor. am. 1. p. 160.) foliis bipinnatis glabris, foliolis inciso-lobatis, racemis pedunculatis subduplicato-bifidis. 5 in Carolinâ et Virginîâ. *Vitis arborea* Willd. sp. 1. p. 1183. *Cissus stans* Pers. ench. 1. p. 183. Caulis arborescens. Baccæ globosæ ochroleucæ. (v. s.)

III. VITIS. Lin. gen. n. 284.

Cal. sub-5-dentatus. Petala 5 apice cohærentia et instar calyptræ basi abscedentia simulque decidua. Stam. 5. Stylus o. Bacca 2-loc., 4-sperma, loculis seminibus sæpè abortivis. — Folia simplicia. — Species pleræque ulteriori egent examine.

* *Hermaphroditæ, in orbe veteri indigenæ.*

1. *V. VINIFERA* (Lin. sp. 293.) foliis lobatis sinnato-dentatis nudis aut tomentosis. 5 in Asiâ australi indigena, nunc ubiquè ferè culta. Blackw. herb. t. 154. Variat foliis plus minus lobatis, glabris, pubescentibus aut tomentosis, planis aut crispis, pallidis aut intense viridibus; ramis prostratis scandentibus erectisve. teneris durisque; racemis laxis aut confertis, ovatis cylindricisve; acinis rubris pallidis aut albis, aquosis aut carnosis, globosis ovatis aut oblongis, saccharatis moschatis aut subacerbis; seminibus numero variis imò interdum omnibus abortivis. Conf. quoad varietates innumeras Duham. arb. fr. 2. t. 1-6. Rox. Clem. var. vit. and. 1. vol. 8. Madrid 1807. Chapt. vign. 2. vol. 8. Paris 1801. Rœm. et Schult. syst. 5. p. 300. Conf. etiam catalogum horti agrarii Luxemburgiensi Parisiensis in quo ferè 1400 varietates coluntur et catalogum arb. fructif. horti Genevensis in quo circiter 600 alatur. (v. v.)
2. *V. LACINIOSA* (Lin. sp. 293.) foliis 5-sectis, segmentis petiolulatis multifidis. 5... Potius prioris mera varietas? Vulgò gall. *Cioutat*. — Corn. can. t. 183. *V. vinifera* var. 24. Rœm. et Schult. syst. 5. p. 314. (v. v.)
3. *V. DENTATA* (Link enum. 1. p. 236.) foliis cordatis sub-5-lobis inæqualiter

micronato-serratis scabriusculis. ♀ Patr. fl. et fr. ign. Petioli glandulis induratis scabri. †.

4. *V. FLEXUOSA* (Thunb. act. soc. lin. 2. p. 332.) foliis cordatis dentatis subtus villosis, caule flexuoso, paniculis elongatis. ♀ in Japoniâ. *V. Indica* Thunb. fl. jap. 103.
5. *V. WALLICHII*, foliis basi cordato-truncatis apice acuminatis acutè dentatis utrinquè levibus, racemis folio brevioribus. ♀ in Nepauliâ. Folia ferè lucida, duplò triplovè minore quàm in *V. vinifera*. Racemi cum pedunculo suo vix pollicem longi. Fruct. ignoti. (v. s.)
6. *V. GLABRATA* (Roth nov. sp. 156. Rœm. et Schult. syst. 5. p. 318.) foliis cordatis subtrilobis serratis glabris, serraturis æqualibus cum lobis obtusis, cirrhis paniculiferis. ♀ in Indiâ orient. Proxima *V. vulpinæ* sed foliorum serraturæ breviores.
7. *V. HEYNEANA* (Rœm. et Schult. syst. 5. p. 318.) foliis cordatis acuminatis indivisis dentatis, suprâ glabris, subtus ferrugineo-tomentosis, paniculis elongatis ♀ in Indiâ orient. *V. cordifolia* Roth non Michx.
8. *V. TRIFIDA* (Roth l. c.) foliis cordato-subrotundis apice trifidis, suprâ griseo subtus ferrugineo-tomentosis repando-serrato-dentatis, corymbis bifidis glomeratis. ♀ in Indiâ orient.
9. *V. TRILoba* (Roth l. c.) foliis cordato-trilobis, suprâ pubescentibus, subtus ferrugineo-tomentosis, inciso-serrato-dentatis acuminatis, lobis subæqualibus, racemis ovatis tomentosis. ♀ in Indiâ orient. An *V. Labrusca* Lour. coch. 1. p. 193?
10. *V. TOMENTOSA* (Roth l. c.) foliis cordato-trilobis tomentosis serratis, lobo intermedio ovato, lateralibus semilunaribus, racemis ovatis densis tomentosis. ♀ in Indiâ orient. *V. trilobæ* affinis, sed folia simpliciter serrata et petioli duplò triplovè longiores.
11. *V. HETEROPHYLLA* (Thunb. fl. jap. 103.) foliis 3-5-lobis aut supremis indivisis serratis glabris paniculâ subdichotomâ glabrâ. ♀ in Japoniâ. Calycis dentes obtusi. Stylus filiformis persistens. Bacca globosa viridis magnitudine Pisi. (v. s. sine fl.)
12. *V. INDICA* (Lin. sp. 293.) foliis cordatis acuminatis subdentatis supernè Incidis subtus pubescentibus. ♀ in Indiâ orient. — Rheed. mal. 7. t. 6. Fructus ex Rheed. acinis confertis globosis nigris, ex Lour. (fl. coch. 1. p. 192.) fusco-viridibus. An omninò eadem?

** *Dioïcæ aut polygamæ, Americanæ.*

13. *V. CARIBÆA*, foliis cordatis acuminatis exsertè et acutè dentatis supernè glabrinusculis subtus pedunculisque tomentosis. ♀ in Jamaicæ dumetis asperis. Fructus magn. Uvæ Corinthiacæ. *V. Indica* Swartz obs. 95. Poir. dict. 8. p. 607. — Sloan. hist. 2. p. 104 t. 210. f. 4. Huc verosim. pertinet *V. Indica* H. B. et Kunth nov. gen. am. 5 p. 227. in Mexico lecta. An à *V. indicâ* certè distincta? An cum *Labruscâ* potius confereunda. (v. s.)
14. *V. LABRUSCA* (Lin. sp. 293.) foliis cordatis subtrilobis acutè dentatis subtus pedunculisque tomentosis subferrugineis. ♀ à Canadâ ad Floridam. Jacq. schoenb. t. 426. — Plum. ic. t. 259. f. 1. Folia vetusta subtus calvæscunt. Racemi fertiles parvi. Baccæ magnæ. (v. s.)
15. *V. ÆSTIVALIS* (Michx. fl. am. bor. 2. p. 230.) foliis lato-cordatis 3-5-lobis, junioribus subtus arachnoideo-tomentosis, adultis glabris. ♀ à Virginiâ ad Carolinam. *V. vulpinâ* Willd. sp. 1. p. 1181. Jacq. schoenb. t. 425. *V. Labrusca* Walt. car. 242. Racemi fertiles oblongi. Baccæ parvæ æstate maturæ.

℞. *sinuata*, foliis sinuato-palmatis grossè dentatis, sinubus rhomboideis. Pursh fl. bor. am. 1. p. 169.

16. *V. CORDIFOLIA* (Michx. fl. am. bor. 2. p. 231.) foliis cordatis acuminatis

inciso-dentatis utrinquē glabris. *h* à Canadâ ad Floridam ubi vulgò anglicè dicitur *Winter-grape* seu *Chicken-grape*. *V. incisa* Jacq. h. schœn. t. 427. *V. vulpina* Walt. car. 243. Racemi laxè multiflori. Baccæ parvæ virescentes serotinæ.

17. *V. RIPARIA* (Michx. l. c.) foliis cordatis breviter 3-fidis inæqualiter inciso-dentatis, petiolo margine nervisque pubescentibus. *h* secus flumina à Pennsylvania ad Carolinam. Vulgò dicitur *Vigne des Battures*. *V. odoratissima* Donn cat. 66. ex Pursh fl. bor. amer. 1. p. 169. (v. s. sine fl.)
18. *V. ROTUNDIFOLIA* (Michx. l. c.) foliis utrinquē lucidis reniformi-cordatis subæqualiter dentatis, racemorum floribus pluries capitulatis. *h* ad ripas fluminum à Virginia ad Floridam. Baccæ magnæ intensè cærulæ. Vulgò ab Anglo-Amer. dicitur *Bullet-grape*.

†. *Species Americanæ non satis notæ.*

19. *V. PALMATA* (Vahl symb. 3. p. 42.) foliis cordatis palmatis glabris, laciniis lanceolatis incisis, umbellis racemosis. *h* in Virginia.
20. *V. VIRGINIANA* (hort. par. ex Poir. dict. 8. p. 608.) foliis ovato-cordatis glabris profundè 5-lobatis, lobis inæqualiter latèque crenatis, racemis subsimplicibus. *h* in Virginia.
21. *V. TILIÆFOLIA* (Humb. et Bonpl. ex Willd. in Rœm. et Schnlt. syst. 5. p. 320.) foliis cordatis serratis subtus incanis. *h* in Americâ calid. Kunthio ignota.

Trib. II. *LEEACEÆ.*

Corolla monopetala. Stamina, petalis alterna? sæpè monadelphæ. Fructus et semina vix noti. Pedunculi in cirrhum non conversi.

IV. *LEEÆ*. *Lin. mant.* 124. — *Aquilicia* *Lin. mant.* 211.

Calyx 4-dentatus. Cor. 5-fida, laciniis revolutis. Stamina urceolus 5-lobus, filamentis extus urceolo adnatis inter lacinias, antheris ovatis glabris. Stylus simplex. Bacca 4-6-locul., loculis seminibusve non nullis abortivis. Semina (ex Gaertn.) in loculis solitaria erecta albumine cartilagineo 5-lobo!, embryo tereti acuminato arcuato subexcentrico.

1. *L. SAMBUCINA* (Willd. sp. 1. p. 1177.) caule sulcato-angulato glabro, foliis subbipinnatis. *h* in Indiâ orient. et in insulis Mauritanis ubi dicitur *Bois de Source*. *Aquilicia sambucina* *Lin. mant.* 211. Cav. diss. 7. t. 218. *Staphylea indica* Burm. ind. t. 24. f. 2? Rheed. mal. 2. p. 43. t. 26. (v. s.)
2. *L. CRISPA* (*Lin. mant.* 124.) caule angulato, angulis fimbriatis, foliis pinnatis, foliolis ovalibus grossè et æqualiter serratis. *h* ad cap. B.-Spei. Lam. dict. 3. p. 460. Ait. h. kew. ed. 2. v. 2. p. 51. *L. pinnata* Andr. bot. rep. t. 355. (v. s. sine fl. ex herb. Van-Royen.)
3. *L. ÆQUATA* (*Lin. mant.* 124.) caule tereti pubescente, foliis pinnatis, foliolis lanceolatis acuminatis serratis, junioribus pubescentibus. *h* in Indiâ orient. Corymbi 3-chotomi.
4. *L. FIRTA* (Horn. h. hafn. 1. p. 231.) caule sulcato angulato petiolis muricatis, foliis pinnatis, foliolis 3-5-ovatis serratis utrinquē scabris. *h* in Indiâ orient. et in Napauliâ. *L. scabra* Roxb? (v. s.)
5. *L. MACROPHYLLA* (Roxb. ex Horn. l. c.) caule angulato petiolis folisque glabris, foliis pinnatis, foliolis lato-ovatis grossè serratis. *h* in Indiâ orient. Corymbi laxi multiflori. (v. s.)

†. *Species non satis nota.*

6. *L. OTTILIS*, fructibus 6-spermis pallidis sapore fatuis. — in Zeylonâ: Aquilicia *Ottilis* Gaertn. fr. 1. p. 275. *Ottilis Zeylanica* Gaertn. fr. 1. t. 57.

V. *LASIANTHERA*. *Beauv. fl. ow. 1. p. 85. t. 51.*

Cal. 5-dentatus extus bracteolatus. Cor. 5-fida. Stam. 5 imæ corollæ inserta, filam. latis cum lobis corollæ alternis, antheris oblongis hirsutis. Stylus brevis.

7. *L. AFRICANA* (Beauv. l. c.) 5 prope Chama ad ripas flum. Saint-Yago. Suffrutescens scandens. Folia ovato-oblonga integra cuspidata. Pedunculi folio oppositi apice umbellatim ramosi. Flores capitati ferè ut in *Araliis*.
-

Ordo XLVI. GERANIACEÆ. DC. fl. fr. 4. p. 838. Sweet
geran. 8°. — *Gerania* Juss. *gen.* 268.

Calyx persistens 5-sepalus, sepalis plus minùs inæqualibus æstivatione imbricatis, interdùm uno, basi in calcar cavum cum pedunculo arctè connatum, producto. Petala 5 (rariùs 4 uno abortivo, rarissimè nulla) sepalis alterna, unguiculata, æqualia aut inæqualia, in prioribus hypogyna, in posterioribus sæpiùs calyci inserta aut cum eo connexa. Stamina filamenta rarissimè libera, ferè semper basi plus minùs monadelpha, simplici serie disposita, hypogyna aut perigyna, numero petalorum dupla aut in Monsoniâ tripla, interdùm nonnulla sterilia, æqualia aut inæqualia. Ovarium primâ fronte 5-loculare in stylum longum crassum et stigmatibus 5 coronatum desinens, sed revera torus elongatus in axim subpentagonum gracilem. Carpella 5 (impropriè arilli cocculivè dicta) submembranacea indehiscentia 1-locularia biovulata ad basim tori adpressa, apice gerentia stylos (aristas auct.) filiformes totidem secùs tori sulcos arctè adnatos et apice summo in stigmata 5 breviter acuta simplicia desinentes, post anthesim styli à basi ad apicem variis modis tortiles carpella è calyce extrahentes et cum iis demùm à toro secedentes. Semina in carpellis solitaria, pendula, exalbuminosa; embryo curvus, radicula deflexa ad fundum carpelli directâ, cotyledonibus foliaceis convoluto-aut flexuoso-plicatis, interdùm lobatis. — Herbæ aut suffrutices, caulibus saltem junioribus articulato-nodosus ad articulationes, ut in Ampelideis, solubilibus. Folia ad nodos inferiores opposita, aut alterna pedunculis tunc folio oppositis, ut in Vite, sed nunquàm cirrhosis. Flores ad apicem pedunculi solitarii aut umbellati.

I. RHYNCHOTHECA. fl. per. prod. p. 142. t. 15. H. B. et K. nov. gen. am. 5. p. 232.

Cal. 5-sepalus æqualis. Pet. 0. Stam. 10., filamentis liberis. Stylus brevissimus adpressus. Stigmata 5 longa crassa. Carpella 5 supernè caudata, infernè dehiscentia. Ovula 2 in quoque carpello axi adfixa pendula. Receptaculum columnare pentagonum. Semina subcarinata. Embryo rectus inversus in albumine carnosus. — Frutices ramosissimi, ramulis spinescentibus. Ramis oppositis tetragonis. Flores pedunculati in apice ramorum fasciculati. — Genus à Geraniaceis differt petalis nullis, staminibus liberis seminibusque albuminosis.

1. R. INTEGRIFOLIA (H. B. et Kunth l. c. t. 464.) foliis oblongis integris, sepalis submucronatis. 3 in temperatis Quitensium prope urbem Alausi. Vix à seq. differt.

2. R. DIVERSIFOLIA (H. B. et K. l. c. t. 465.) foliis obovato-oblongis integra vel

trifidis, sepalis mucronato-aristatis. § Cum priore ad Pillao et Pamallactam. R. spinosa Ruiz et Pav. syst. p. 142. Frutex sepiarius.

II. MONSONIA. Lin. f. suppl. p. 342.

Calyx 5-sepalus, sepalis æqualibus apice aristato-mucronatis. Pet. 5 æqualia calyce duplò maiora. Stam. 15 basi monadelpha, sæpius in fasciculos 5 tristemones semi-coalita. Fructus Erodii et Pelargonii. — Omnes species ad cap. Bonæ-Spei crescunt.

Sect. I. SARCOCAULON.

Caulis fruticoso-carnosus, spinis horridus. Folia ovata aut oblonga integra aut subdenrata. Pedunculi 1-flori ad basin bracteolis 2 minimis stipati. Petala integra. Stam. (ex ic. L'her.) imâ basi tantum monadelpha nec in fasciculos 5 coalita. — An genus?

1. *M. L'HERITIERI*, foliis ovatis mucronatis integris aliis subsessilibus aliis longè petiolatis. § *M. spinosa*. L'her. ger. t. 42. Lam. dict. 3. p. 270. Willd. sp. 3. p. 719. Flores diam. bipollicari. Petioli persistentes spinescentes.
2. *M. PATERSONII*, foliis oblongo-cuneatis obtusis integris, aliis sessilibus, nonnullis breviter petiolatis. § *Paters. itin.* t. 14. Flores diam. pollicari. Pedunculi forsan ex icone persistentes spinescentes.
3. *M. BURMANNI*, foliis oblongo-cuneatis crenatis subcomplicatis. § *Burm afr.* p. 81. t. 31. Ger. spinosum *Burm.* ger. u. 2. Cav. diss. 4. p. 195. t. 75. f. 2. Folia videntur ex icone alia sessilia, alia minora longè petiolata, petiolis demum persistentibus spinescentibus. Flores diam. semi pollicari. Stam. nota satis nota sed dum 10 adsint, adhuc à Geranio fructus aristis more Erodii tortilibus differe videtur, et habitu ad 2 priores accedit.

Sect. II. OLOPETALUM.

Caulis herbaceus. Folia subovalia dentata. Stipulæ bracteolæque subulatæ subinduratæ. Pedunculi 1-2-flori, medio bracteolas 2-4 gerentes. Petala ovata integra. Stam. pentadelpha.

4. *M. OVATA* (Cav. diss. 4. p. 193. t. 113. f. 1.) foliis ovato-oblongis subcordatis crenatis undulatis, stipulis rigidulis, pedunculis 1-floris 2-bracteolatis. (2) Ger. emarginatum Lin. f. suppl. 306. *M. emarginata* L'her! ger. t. 41. In ipso L'heritieri specimine vidi stamina pentadelpha, ut in icone Cavendishii sunt depicta, nec monadelpha ut videntur in Geraniologiâ. (v. s.)
5. *M. BIFLORA*, foliis elliptico-oblongis obtusis dentatis basi subconnatis, stipulis spinescentibus, pedunculis bifloris quadribacteolatis. — Burch! cat. afr. austr. n. 2611. Flores magn. prioris. Sepala brevissimè mucronata. (v. s.)

Sect. III. ODONTOPETALUM.

Caulis herbaceus. Folia lobata aut multifida. Pedunculi longi uniflori medio bracteis 6-8 verticillatis stipati. Petala oblonga apice grossè dentata. Stam. pentadelpha.

6. *M. LOBATA* (Mont. act. goth. 2. p. 1. t. 1. ex Willd. spec. 3. p. 718.) foliis cordatis 5-7-fidis, lobis obtusis serratis, subtus petiolis calycibusque pilosisculis. 2. bot. mag. t. 385. herb. amat. t. 5. *M. filia* Lin. f. suppl. 341. Cav. diss. 3. p. 180. t. 74. f. 2. Ger. anemonoides Thunb. prod. 112. Fructus rostrum longissimum. Folia plus minus pilosa. (v. s.)
7. *M. PILOSA* (Willd. enum. 717.) foliis palmatim 5-partitis, laciniis tripartito-pinnatifidis, subtus petiolis calycibusque pilosis. 2. *M. filia* Pers. ench. n. 3. Geranium *Monsonia* Thunb. prod. 112. *M. speciosa* Sweet ger. t. 77. Petala extus virescentia, intus albida, basi rubro maculata. Sequenti affinis. (v. s.)
8. *M. SPECIOSA* (Lin. f. suppl. 342.) foliis palmatim 5-partitis, laciniis tenuiter bipinnatifidis, petiolis calycibusque glabris. 2. Curt. bot. mag. t. 73. Cav. diss. 3. t. 74. f. 1. Ger. speciosum Thunb. Petala purpurea. (v. s.)

III. GERANIUM. *L'her. ger. DC. fl. fr. 4. p. 844.*

Sepala 5 æqualia. Pet 5 æqualia. Stam. 10 fertilia alterna majora. Glandulæ nectariferæ ad basin stam. majorum. Carpellorum aristæ intus glabræ, demum elasticæ à basi ad axeos apicem circinnatim revolutæ. — Herbæ rarissimè suffrutescentes, foliis palmato-lobatis, pedunculis 1-2-floris.

* *Perennia, pedunculis 1-floris.*

1. *G. SESSILIFLORUM* (Cav. diss. 4. p. 198. t. 77. f. 2.) acaule, pedunculis radicalibus petiolo multò brevioribus, foliis reniformibus 5-7-partitis, lobis 3-5-fidis. \mathcal{U} ad fretum Magellanicum. *G. brevipes* L'her. mss. (v. s.)
2. *G. ACAULE* (Willd. mss. in H. B. et K. nov. gen. am. 5. p. 231.) acaule, pedunculis radicalibus brevibus, foliis multipartitis, laciniis linearibus integerrimis. \mathcal{U} in Andibus Quitensium alt. 2100 hexap. \dagger .
3. *G. CUCULLATUM* (H. B. et K. l. c.) cespitosum, ramis foliosis, pedunculis folio parùm longioribus, foliis 5-partitis cucullatis ciliatis glabris, laciniis intermediis 2-3-fidis, calycibus mucronatis. \mathcal{U} in frigidis montium Popayanensium alt. 1690 hexap. Fl. ign.
4. *G. DIFFUSUM* (H. B. et K. l. c.) caulis ramosis diffusis reflexo-pilosis, pedunculis petiolo paulò brevioribus, foliis profundè 5-fidis subtus adpresso-pilosis, laciniis 3-lobis, calycibus mucronatis. \mathcal{U} in montibus Peruviae. Fl. ign.
5. *G. AGAVACENSE* (Willd. in H. B. et K. l. c.) caule divaricato glabro, foliis 5-partitis, laciniis lanceolatis bipartitis, intermedià bifidâ, floribus axillaribus? longè pedunculatis. \mathcal{U} prope Ayavaca Peruvianorum. Flor. albi. \dagger .
6. *G. CHILOENSE* (Willd. in H. B. et K. l. c.) caule ramoso sericeo glanduloso piloso, foliis 5-lobis, lobis trifidis oblongis dentatis pubescentibus, floribus axillaribus? \mathcal{U} prope Chillo Quitensium. Flor. albi. \dagger .
7. *G. PHILONOTHEUM*, caulis prostratis, pedunculis petiolo longioribus imà basi bracteolatis sæpè tortilibus, foliis reniformibus 3-5-partitis, lobis cuneatis, apice 3-dentatis. \mathcal{U} in Novæ-Hollandiæ orâ merid. Habitus ferè Ranunculi parviflori seu Philonotis, sed glabrum. (v. s.)
8. *G. POTENTILLOIDES* (L'her! mss.) caulis prostratis, pedunculis petiolo longioribus suprâ medium bibracteolatis, foliis reniformibus 5-partitis, lobis 3-fidis. \mathcal{U} in Novâ-Hollandiâ. Banks. (v. s.)
9. *G. RADICANS*, caulis prostratis ad nodos radicanibus, pedunculis petioli longitudine imà basi bracteolatis, foliis tripartitis, lobis cuneato-ovatis grosse dentatis. \mathcal{U} in Napauliâ. Wallich. (v. s.)
10. *G. SIBIRICUM* (Lin. sp. 957.) caule erecto subdiffuso ramoso, pedunculis petiolo longioribus ferè infrâ medium bibracteolatis, foliis 5-partitis, lobis oblongis incisio-dentatis. \mathcal{U} in Sibiria, Caucaso, China. — Gmel. sib. 3. t. 67. Cav. diss. 4. t. 77. f. 1. Cor. alba striis purpureis notata. (v. s.)
11. *G. SANGUINEUM* (Lin. sp. 958.) caule erecto aut diffuso ramoso, pedunculis axillaribus petiolo multò longioribus medio bibracteolatis, foliis oppositis 5-partitis, lobis 3-fidis, lobulis linearibus. \mathcal{U} in Europæ sylvaticis. Cav. diss. 4. t. 76. f. 1. flor. dan. t. 1107. Flores sanguinei ampli. (v. s.)
 β . *prostratum* (Cav. diss. 4. p. 196. t. 76. f. 3.) caule prostrato ad nodos subtumido, floribus roseis variegatis. *G. Lancastriense* With. brit. 600. (v. s.)
 γ . *villosissimum* (DC. fl. fr. n. 4541.) caule prostrato foliisque villosissimis, floribus purpureis, angue albo.
12. *G. POTENTILLÆFOLIUM*, caule subdiffuso ramoso, pedunculis axillaribus petiolo multò longioribus suprâ medium bibracteolatis, foliis oppositis subtus albo-tomentosis breviter petiolatis 5-partitis, lobis laciniatis, lobulis linearibus. \mathcal{U} in Novâ Hispaniâ. Sequenti valdè affine, sed pedunculi 1-flori. (v. s. et ic. fl. mex.)

** *Perennia, pedunculis bifloris.*

13. *G. INCANUM* (Lin. sp. 957.) caule diffuso, foliis subtus albo-tomentosis 7-partitis, lobis multifidis linearibus, pedunculis elongatis. calycibus adpresse sericeo-villosis, petalis integris. \mathcal{L} ad cap. Bonæ-Spei. — Burm. ger. 26. t. 1. Cav. diss. 4. t. 82. f. 2. Folia ferè Potentillæ argenteæ. (v. s.)
14. *G. CANESCENS* (L'her! ger. t. 38.) caule diffuso, foliis subtus canescentibus 5-partitis, lobis oblongis inciso-dentatis, pedunculis longissimis calycibusque glanduloso-pilosis, petalis emarginatis. \mathcal{L} ad cap. Bonæ-Spei. Affine hinc *G. incano*, illinc asphodeloidi. (v. s.)
15. *G. SUBCAULESCENS* (L'her! mss.) caule brevissimo, foliis subradicalibus petiolatis villosis subcineris 5-partitis, lobis 3-dentatis obtusis, petiolorum pedunculorumque pube patulâ, petalis obtusissimis calyce villosos longioribus. \mathcal{L} in summo monte Parnasso. Sibthorp. *G. asphodeloides* fl. gr. prod. 2. p. 40. ic. t. 661. Habitus *G. canescentis*. An var. *G. asphodeloidis*. (v. s.)
16. *G. ARGENTUM* (Lin. amœn. 4. p. 324.) caule brevissimo, foliis subradicalibus longè petiolatis utrinque sericeo-canis 5-7-partitis, lobis 3 fidis, lobulis linearibus, pedunculis subradicalibus, petalis emarginatis. \mathcal{L} in Alpibus Gallo-provinciæ, Pedemontii, Carinthiæ. Flores magni venosi. (v. s.)
17. *G. CINEREUM* (Cav. diss. 4. p. 204. t. 89. f. 1.) subcaule, foliis subradicalibus petiolatis glauco-pubescentibus 5-partitis, lobis cuneiformibus 3-fidis, pedunculis subradicalibus, petalis emarginatis. \mathcal{L} in Pyrenæis. *G. varium* L'her. ger. t. 37. *G. cineraceum* Lapeyr. pyr. t. 2. Radix ut in priore crassa subliguosa. (v. v.)
18. *G. ANEMONIFOLIUM* (L'her! ger. t. 36.) caule fruticoso, foliis glabris palmato-5-sectis, segmentis bipinnati-fidis, pedunculis oppositis erectis pilosis. \mathcal{Z} in Magerâ et Teneriffâ. *G. palmatum* Cav. diss. 4. t. 84. f. 2. *G. lævigatum* Burm. ex L'her. mss. Flores ampli, purpurascentes. (v. v.)
19. *G. MACRORHIZON* (Lin. mant. 343.) caule basi suffruticoso apice dichotomo, foliis glabris 5-partitis, lobis apice dentatis, calycibus globoso inflatis, petalis integris, staminibus nutantibus. \mathcal{L} in Italiâ, Carinthiâ, Græciâ. Jacq. ic. rar. 1. t. 134. Cav. diss. 4. p. 212. t. 25. Sims bot. mag. t. 2420. (v. s.)
20. *G. TUBEROSUM* (Lin. sp. 953.) radice subglobosâ, caule à basi ad bifurcationem nudo, foliis 5-partitis, lobis linearibus pinnatim inciso-serratis. \mathcal{L} in campestribus à Massiliâ ad Tauriam. Lob. ic. t. 661. f. 2. Moris. oxon. 5. t. 16. f. 21. Cav. diss. 4. t. 78. f. 1. Sweet ger. 2. t. 155. *G. radiculatum* Bieb. fl. taur. 2. p. 134. excl. syn. Moris. (v. s.)
21. *G. LINEARILOBUM* (DC. fl. fr. suppl. p. 628. in adn.) radice subglobosâ, caule subpubente à basi ad bifurcaturam nudo; foliis multipartitis, lobis linearibus subdivisis integerrimis. \mathcal{L} in nemorosis et campestribus ad Wolgam infer. et in promont. Caucasico. *G. tuberosum* Bieb. fl. taur. 2. p. 135. excl. syn. Lin. Cav. Lam. Mor. (v. s.)
22. *G. GYMNOCaulon*, caule glabro à basi ad bifurcaturam nudo, foliis super 3-partitis, lobis pinnato-incisis, calycibus ciliato-villosis, petalis integris. \mathcal{L} in Iberiâ. (Steven) Radix et folia radicalia ignota. Flores ampli cærulei. (v. s.)
23. *G. IBERICUM* (Cav. diss. 4. p. 209. t. 124. f. 1.) caule villosa dichotomo, foliis 5-7-partitis, lobis pinnatim incisis, lobulis dentatis, calycibus villosissimis, petalis obcordatis subtrifidisve. \mathcal{L} in Iberiâ. bot. mag. t. 1386. Sweet ger. t. 84. *G. grandiflorum* Guld. it. 1. p. 426. Flor. ampli cærulei. (v. s.)
24. *G. NOBISUM*. (Lin. sp. 953.) caule tetragono, foliis infer. 5-lobis, super. 3-lobis, lobis oblongis acuminatis serratis, petalis emarginatis. \mathcal{L} in Angliâ, Galliâ, Italiâ. Cav. diss. 4. t. 80. f. 1. Carpella undiquè pilosa. Petala striatovenosâ. Folia subtus lucida. (v. v.)
25. *G. HERNANDESI* (fl. mex. ic. ined.) caule teretiusculo, ramis petiolisque patentem pilosis, foliis infer. 5-lobis, super. 3-lobis, lobis oblongis acumi-

- natis serratis, petalis oblongo-cuneatis subintegris. \mathcal{U} in Mexico. Flores pallidi. Hern. mex. 293. f. 2. An à sequente satis differt?
26. *G. MEXICANUM* (H. B. et K. nov. gen. am. 5. p. 230.) caule erecto reflexo piloso, foliis pedato-5-fidis utrinque adpresso-pilosis, laciniis apice 3-7-fidis, pedunculis glanduloso-pilosis, calycibus mucronatis. \mathcal{U} inter Guanajuato et Santa-Rosa Mexicanorum.
27. *G. ANGULATUM* (Curt. bot. mag. t. 203.) caule angulato, foliis radicalibus 7-lobis, caulinis 5-lobis, lobis oblongis acuminatis dentatis, petalis emarginatis. \mathcal{U} Patr. ign. — *G. venosum* Pers. ench. 2. p. 235. Petala striato-venosa ut in *G. nodoso* ad quodvaldè accedit.
28. *G. WALLICHIANUM* (Sweet ger. t. 90.) caule erecto subangulato, foliis oppositis 5-lobis, lobis cuneato-ovatis lobato-dentatis, stipulis cujusque folii in unicum ovatam integram coalitis, petalis emarginatis, stigmatibus longissimis. \mathcal{U} in Nepalicæ montibus. Pet. magna purpurea. Herba sericeo-villosa. Sims bot. mag. t. 2377.
29. *G. VLASSOVIANUM* (hortul. et DC. in mem. soc. gen. 1. p. 441.) caule tereti, foliis 5-lobis, lobis ovali-acuminatis inciso dentatis, stipulis cujusque folii in unicum apice bifidam connatis, petalis obovatis integris. \mathcal{U} in Sibiria? Petala striato-venosa ut in *G. striato*, stipulæ connatæ ut in *G. Wallichiano*. (v. v.)
30. *G. STRIATUM* (Lin. amœn. 4. p. 282.) caule tereti, foliis infer. 5-lobis, summis 3-lobis, lobis ovatis acutis inciso-dentatis, stipulis liberis, petalis emarginato-bilobis. \mathcal{U} in Italiâ et Græciâ. Cav. diss. 4. t. 79. f. 1. Curt. bot. mag. t. 55. Delann. herb. am. t. 9. Petala reticulato-venosa. Carpella glabrinacula basi villosa. Folia ad sinus supernè fusco-maculata. (v. s.)
31. *G. REFLEXUM* (Lin. mant. 257.) caule tereti, foliis alternis 5-7-lobis inciso-dentatis, summis sessilibus, petalis reflexis apice dentato-laceris, staminibus glabris, carpellis transversè plicatis. \mathcal{U} in Italiâ, Galliâ? Cav. diss. 4. t. 81. f. 1. (v. s.)
32. *G. PHÆUM* (Lam. dict. 2. p. 658.) caule tereti, foliis 5-lobis inciso-dentatis, summis sessilibus, petalis patulis integris, staminum filamentis basi pilosis, carpellis transversè plicatis. \mathcal{U} in montosis Galliæ, Helvetiæ.
- a. *vulgatius*, petalis livide rubro-fuscis basi albo-maculatis. *G. phæum* Lin. sp. 933. Cav. diss. 4. t. 89. f. 2. (v. v.)
- β. *fuscum*, (Lin. mant. 97.) petalis atro-fuscis.
- γ. *lividum*, (L'her! ger. t. 39.) petalis roseo-violaceis subemarginatis. *G. patulum* Vill. dauph. 3. p. 371. *G. subcæruleum* Schleich. cat. 25. (v. v.)
33. *G. ERIOSTEMON* (Fisch! ined.) caule tereti simplici, foliis 5-lobis, lobis ovatis grossè dentatis, inferioribus longè petiolatis alternis, summis sessilibus oppositis, petalis patulis integris, staminum filamentis pilosis. \mathcal{U} in betuletis Dahuriæ. Calyces aristati. Flores cærulei. (v. s.)
34. *G. ERIANTHUM*, caule tereti subsimplici infernè nudo, foliis omnibus petiolatis palmato-5-7-lobis, lobis incisic acutè serrato-laceris, pedunculis confertis abbreviatis, calycibus villosissimis, petalis integris, staminum filamentis pilosis. \mathcal{U} in Kamschatkâ et Amer. bor. et occidentali. Nelson. (v. s.)
35. *G. SYLVATICUM* (Lin. sp. 954.) caule tereti erecto glabro, foliis sub-7-lobis, lobis oblongis inciso-serratis, pedunculis subcorymbosis, petalis subemarginatis, staminum filamentis subulatis ad medium ciliatis. \mathcal{U} in Europæ sylvis et pascuis montosis. Engl. bot. t. 121. Carpella pilosa. (v. v.)
- β. *batrachioïdes* (Cav. diss. 4. p. 211. t. 85. f. 1.) floribus majoribus, petalis integris.
36. *G. PRATENSE* (Lin. sp. 954.) caule tereti erecto tomentoso, foliis sub-7-lobis, lobis lineari-oblongis inciso-serratis, pedunculis subcorymbosis, petalis integris, staminum filamentis glabris basi valdè dilatatis. \mathcal{U} in Europæ

- dumetis. engl. bot. t. 424. Cav. 4. t. 87. f. 1. Delaun. herb. am. t. 118: Flores ampli cærulei. Petala 2-3 alba, alia cærulea, in var. vidi. (v. v.)
37. *G. LONGIPES* (DC. mem. soc. gen. 1. p. 442.) caule tereti erecto glabro, foliis palmato-subpeltatis 5-7 lobis, lobis oblongis grossè incisis, pedunculis è ramorum dichothomiâ longissimis, petalis integris, staminum filamentis subulatis glabris. \mathcal{L} Patr. ign. *G. Londesii* Link enum. h. ber. 2. p. 196. Flores magnitudine prioris violaceo-lilacini. Cal. aristati. (v. v.)
38. *G. MACULATUM* (Lin. sp. 955.) caule subangulato erecto dichotomo retrorsum pubescente, foliis 3-5-partitis inciso-dentatis, radicalibus longè petiolatis, summis oppositis sessilibus, petalis integris, staminum filamentis vix basi subciliatis. \mathcal{L} à Canadâ ad Carolinam. — Dill. h. elth. t. 132. f. 159. *G. maculatum* Cav. 4. t. 86. f. 2. Bigel. am. med. bot. 1. p. 84. t. 8. Flores purpurei nunc magnitudine *G. pratensis*, nunc ferè dimidiò minores. (v. s.)
39. *G. COLLINUM* (Bieb. fl. taur. 2. p. 137. suppl. 455.) caule angulato diffuso retrorsum pubescente, foliis palmato-5-partitis, lobis subtrilobis inciso-serratis, pedunculis calycibusque villosis, petalis integris subrotundis, staminibus subulatis glabris. \mathcal{L} in Tauriæ et Caucasi pratis montosis. Species varietatibus intricata, nec satis definita. Antheræ semper violaceæ. (v. s.)
40. *G. ASPHODELOIDES* (Willd. in journ. Schrad. 2. p. 26. t. 1.) caule flaccido diffuso pilis retrorsis hispido, foliis 5-lobis, lobis 3-fidis subincisis, pedunculis calycibusque villosis, petalis emarginatis, staminibus subulatis glabris. \mathcal{L} in Græciæ, Byzantii et Orientis pratis. — Tourn! voy. 1. p. 526. ic. *G. Orientale* Mill. dict. n. 10. (v. s.)
41. *G. PALUSTRE* (Lin. amœn. 4. p. 323.) caule decumbente pilis patulis subhispido, foliis 5-7-lobis, lobis inciso-dentatis, pedunculis longissimis pilosis, pedicellis declinatis, petalis integris, staminibus subulatis glabris. \mathcal{L} in pratis paludosis subsylvaticis à Galliâ ad Dahuriam. Sweet ger. t. 3. — Oeder fl. dan. t. 596. (v. s.)
42. *G. ACONITIFOLIUM* (L'her! ger. t. 40.) caule subadscendente glabriusculo, foliis subpeltato-7-partitis, lobis laciniatis, pedunculis calycibusque villosis, petalis integris, staminibus subulatis glabriusculis. \mathcal{L} in Alpibus Delphinatûs et Helvetiæ ad rivulos. *G. rivulare* Vill. dauph. 3. t. 40. Radices fasciculatæ. Flores albi lineati dimidiò minores quàm in *G. pratense*. (v. v.)
43. *G. DAHURICUM*, caule basi nudo erecto glabro, foliis caulinis oppositis 3-5-partitis, lobis incisis acutis, pedunculis folio triplò longioribus fructiferis deflexis, calycibus glabriusculis, petalis integris basi valdè barbatis, staminibus subulatis ciliatis. \mathcal{L} in turfosis Dahuriæ. Patrin. Radix tuberosis elongatis fasciculatis ut in *G. aconitifolio* cui affine. (v. s.)
44. *G. BIFOLIUM* (Patr! ined.) caule glabro erecto infra bifurcationem nudo, foliis radicalibus binis petiolatis caule triplò brevioribus, ceteris oppositis sessilibus, omnibus adpressè villosis 7-partitis, lobis linearibus pinnato-dentatis, calycibus adpressè villosis, petalis integris, staminibus ciliatis. \mathcal{L} in Dahuriæ pinetis circa Barnaoul. (v. s.)
45. *G. CÆRULEUM* (Patr! ined.) caule glabro basi nudo dichotomo, foliis caulinis oppositis 5-7-partitis, lobis pinnato-incisis acutis, calycibus villosissimis, petalis integris basi vix puberulis, staminibus ciliatis calyce dimidiò brevioribus. \mathcal{L} in Dahuriæ campestribus. Flores parvi lætè saphirini. Tribus prioribus affine; petala ut in iis parùm calyce longiora. (v. s.)
46. *G. FILOSUM* (Forst. prod. 531.) caulibus subdecumbentibus ramosis petiolis pedunculisque pilis patulis hispidis, foliis 3-5-partitis, lobis in lacinis lineares obtusos trifidis, calycibus ciliatis, staminibus glabriusculis. \mathcal{L} ? in Novâ-Zelandiâ. Sweet ger. 2. t. 119. Petala obovata subemarginata pallidè purpurascientia. (v. s.)
47. *G. PARVIFLORUM* (Willd. enum. 716.) caulibus subdecumbentibus, petiolis

pedunculis calycibusque glabriusculis, foliis caulinis oppositis 3-5-partitis, lobis 3 fidis dentatis, petalis calyce subbrevioribus, staminibus glabris. \mathcal{L} in Novæ-Hollandiæ terrâ Van-Diemen. Banks. (v. s.)

48. *G. NAPAULENSE* (Sweet ger. t. 12.) caule prostrato compresso, foliis oppositis 5-lobis, lobis oblongis inæqualiter dentatis, pedunculis elongatis calycibusque pilosis, petalis obovatis subemarginatis calycis longitudine, carpellis pilosis. \mathcal{L} ? in Napauliâ. (v. s. comm. à cl. Wallich.)

49. *G. CRISTATUM* (Stev! mem. soc. h. nat. mosc. 4. p. 50. t. 5.) caule flaccido simplici, foliis reniformibus 7-lobis, lobis 3-fidis, lobulis 3-dentatis, pedunculis elongatis calycibusque hispidis, petalis emarginatis calyce majoribus, carpellis cristatis (ut in *Onobrychide Caput-Galli*). \mathcal{L} in Albaniâ Ibericâ, ad rivum Jucharibasch. *G. Albanum* Bieb. fl. taur. 2. p. 137. (v. s.)

50. *G. PYRENAÏCUM* (Lin. mant. 97.) caule erecto ramoso, foliis reniformibus 7-lobis, lobis oblongis obtusis 3-fidis, lobulis 3-dentatis, floralibus 3-partitis, petalis emarginatis calyce paulò majoribus, carpellis ecristatis pubescentibus. \mathcal{L} in submontosis ab Hispaniâ et Angliâ ad Iberiam. — Engl. bot. t. 405. Cav. diss. 4. t. 99. f. 2. Variat flore albo et pallide purpureo. (v. v.)

β . *umbrosum* (Kit! pl. hung. 2. p. 131. t. 124.) caule flaccidiore subnudo, carpellis subglabris. \mathcal{L} in umbrosis Hungariæ.

γ ? *nemoriosum* (Ten. cat. 1819. p. 60.) foliis profundius lobatis, petalis integerrimis. \mathcal{L} in nemoribus regni Neapolitani.

*** *Annua, pedunculis bifloris.*

51. *G. MOLLE* (Lin. sp. 955.) foliis reniformibus, radicalibus 9-lobis, caulinis 7-lobis, lobis 3 fidis, petalis bifidis longitudine calycis mutici, carpellis glabris rugosis, seminibus lævibus. $\textcircled{1}$ in ruderalis Europæ. — Vaill. bot. t. 15. f. 3. fl. dan. t. 679 Flores rarissime abortu 5-antheri. (v. v.)

52. *G. FUSILLUM* (Lin. sp. 957.) foliis subreniformibus 7-lobis, lobis 3-fidis, petalis emarginatis longitudine calycis mutici, carpellis pubescentibus non rugosis, seminibus lævibus. $\textcircled{1}$ in Europæ ruderalis. *G. malvæfolium* Lam. fl. fr. 3. p. 18. — Vaill. bot. t. 15 f. 1. Stamina sæpissimè 5 sterilia. (v. v.)

β . *humile* (Cav. diss. 4. t. 83. f. 2.) foliis tenuius incis, caule humiliore. (v. v.)

53. *G. ROTUNDIFOLIUM* (Lin. sp. 957.) foliis radicalibus reniformibus 7-lobis, caulinis subrotundis basi truncatis 5-lobis, lobis 3-fidis, petalis integris longitudine calycis aristati, carpellis hirsutis, seminibus reticulatis $\textcircled{1}$ in pascuis et pratis. Cav. diss. 4. t. 93. f. 2. engl. bot. t. 157. Flores parvi rosei. (v. v.)

54. *G. PALLENS* (Bieb. fl. taur. 2. p. 138.) foliis infer. petiolatis, summis sessilibus palmato-5-partitis, lobis 3-lobis, petalis integris calyce aristato paulò longioribus, carpellis hirsutis, seminibus reticulatis. $\textcircled{1}$? in Iberiâ occid. Habitus *G. rotundifolii*. (v. s. comm. à cl. Steven.)

55. *G. COLUMBINUM* (Lin. sp. 956.) foliis 5-partitis, lobis multifidis linearibus, petalis emarginatis longitudine calycis aristati, carpellis glabris lævibus, seminibus reticulatis. $\textcircled{1}$ in Europæ arvis. — fl. dan. t. 1222. Cav. diss. 4. t. 82. f. 1. (v. v.)

56. *G. DISSECTUM* (Lin. sp. 956.) foliis 5-partitis, lobis 3-fidis linearibus, petalis emarginatis longitudine calycis aristati, carpellis pilosis, seminibus reticulatis $\textcircled{1}$ in Europæ et Iberiæ arvis. — fl. dan. t. 936. Cav. diss. 4. t. 78. f. 2. Vaill. bot. t. 15. f. 2. Petala purpurea. (v. v.)

57. *G. CAROLINIANUM* (Lin. sp. 956.) foliis ultrà medium 5-lobis, lobis inciso-3-5 fidis, pedunculis apice confertis, petalis emarginatis longitudine calycis aristati, carpellis pilosis, seminibus lævibus. $\textcircled{1}$ in arvis à Georgiâ ad Virginiam. Cav. diss. 4. t. 84. f. 1. et t. 124. f. 2. Petala alba venosa. — *G. lanuginosum* Jacq. schœnb. 2. t. 140. ab hoc differre videtur caule petiolis pedunculisque hispidis et petalis purpurascentibus. (v. s.)

58. *G. VILLOSUM* (Ten. cat. 1819. p. 60.) foliis orbiculato-cordatis 9-lobis, lobis 3-fidis obtusis, corollis calyce duplò longioribus, petalis emarginatis, caule erecto. ① in paludibus Silæ Neapolitani agri. †.
59. *G. RETRORSUM* (L'her! mss.) foliis 5-partitis, lobis 3-fidis sublinearibus, lobulis 3-dentatis obtusis, pube caulis retrorsum adpressâ, petalis obtusis calyce aristato vix longioribus, carpellis villosis, seminibus reticulatis? ① in Novâ Zelandiâ. Banks. (v. s. cum semin. immat.)
60. *G. BOHEMICUM* (Lin. amœn. 4. p. 323.) foliis 5-lobis, lobis cuneato-ovatis inciso-dentatis, pube caulis patente hispida, petalis emarginatis longitudine calycis longè aristati, carpellis hirsutis, seminibus lævibus. ① in Bohemiâ, Silesiâ, Valesiâ montosis sylvaticis. — *G. lanuginosum* Lam. dict. 2. p. 655. è Numidiâ ortum huc ex descr. pertinere videtur. (v. s.)
61. *G. DIVARICATUM* (Ehrh. beitr. 7. p. 164.) foliis 5-lobis super. 3-lobis, lobis oblongis grossè et irregulariter dentatis, pube caulis patente hispida, petalis emarginatis calyce subaristato brevioribus, carpellis scabris supernè rugosis, seminibus lævibus. ① in Hungariâ, Valesiâ, Caucaso. *G. novum* Wint. ic. t. 2. *G. Winterli* Roth cat. 2. p. 78. — pl. hung. 2. t. 123. (v. s.)
62. *G. LUCIDUM* (Lin. sp. 955.) glaberrimum, foliis rotundatis 5-lobis, calycibus pyramidato-angulatis transversè corrugatis, carpellis dorso sulcato-muricatis. ① in Europæ rupibus umbrosis. — fl. dan. t. 218. Cav. diss. 4. t. 80. f. 2. Flores rosei. (v. v.)
63. *G. ROBERTIANUM* (Lin. sp. 955.) foliis 3-5-partitis, lobis trifido-pinnatifidis, petalis integris calyce angulato aristato duplò longioribus, carpellis glabris reticulato-rugosis, seminibus lævibus. ① in rupibus Europæ frequens. (v. v.)
- β. *purpureum* (Vill! dauph. 3. t. 40.) petalis calyce paulò majoribus. — in apricis.
- γ. *Mosquense*, caule erectiore corymboso-paniculato, odore magis therebenthinaceo. — in paludosis circa Mosquam. Goldbach mem. soc. mosc. 5. p. 133. (v. s.)
- δ. *inodorum*, floribus pallidioribus, odore subnullo. — in Amer. bor. Pursh fl. bor. am. 2. p. 449.

†. *Gerania non satis nota.*

64. *G. RANUNCULOIDES* (Burm. ger. n. 4.) pedunculis longissimis 1-floris, foliis orbiculato-multifidis, radice tuberosâ, ramis dichotomis. 2/ in Africâ. An à *G. canescente* diversum?
65. *G. LUPINOIDES* (Burm. ger. n. 65.) pedunculis radicalibus binis bifidis petiolo duplò longioribus, foliis orbiculatis incisè tomentosis, lobis linearibus, radice tuberosa. 2/ ad cap. Bonæ-Spei. — Pluk. t. 186. f. 3. An à *G. argenteo* diversum?
66. *G. ARABICUM* (Forsk. descr. 124.) pedunculis binis, foliis rotundis incis. — in Yemen frequens. Variat flore albo et rubro basi fuscescente.

IV. *ERODIUM*. L'her. ger. icon. et text. ined. DC. fl. fr. 4. p. 838.

Sep. 5 æqualia in calcar seu tubum nectariferum nulla producta. Pet. 5 regularia aut irregularia. Stam. 10, filamentis basi monadelphis, 5 antheriferis, 5 alternis sterilibus. Glandulæ 5 ad basin staminum sterilium. Carpellorum aristæ intus barbatæ demùm elasticè spiraliter tortæ. — Herbæ, aut suffrutices, foliis variis, pedunculis sæpius multirarissimè 1-floris.

* *Foliis pinnati sectis aut pinnati-partitis.*

1. *E. TATARICUM* (Willd. sp. 3. p. 625.) acaule, pedunculis subbifloris, foliis pubescentibus pinnatisectis, segmentis pinnatifidis, lobis linearibus, rachi inter segmenta nudâ, petalis obovatis calyce duplò longioribus. 2/ in Tataria et Sibiria ulteriore. Sequenti affinis. Petala obscurè cærulea, Willd.

2. *E. SUPRACANUM* (L'her! ger. t. 2.) acaule, pedunculis subbifloris, foliis supra canis pinnati sectis, segmentis pinnatifidis, lobis lanceolato-linearibus, rachi inter segmenta dentatâ, petalis retusis calyce duplò longioribus. \mathcal{L} in saxosis montis Serrati prope Barcinonam. G. rupestre Cav. diss. 4. t. 90. f. 4. (v. s.)
3. *E. PETRÆUM* (Willd. sp. 3. p. 625.) acaule, pedunculis multifloris, foliis glabriusculis pinnati-sectis, segmentis pinnatifidis, lobis lanceolato-linearibus, rachi inter segmenta dentatâ, petalis retusis calyce duplò longioribus. \mathcal{L} in Occitaniæ et agri Ruscini rupibus apricis. — Gon! ill. t. 21. f. 1. *E. fœtidum* L'her! ined. n. 6. Flor. purpurascens. (v. v.)
 β . *crispum* (Lapeyr. abr. 390.) foliis villosioribus. (v. s.)
 γ . *lucidum* (Lapeyr. abr. 390.) foliis glabratiss. (v. s.)
4. *E. GLANDULOSUM* (Willd. sp. 3. p. 628.) acaule, pedunculis multifloris, foliis pubescenti-glandulosis bipinnatisectis, segmentis pinnatifidis, lobis lanceolato-linearibus, rachi inter segmenta dentatâ, petalis subæqualibus acutis calyce duplò longioribus. \mathcal{L} in Hispaniâ et Pyrenæis *E. macradenum* L'her! ger. t. 1. G. glandulosum Cav. diss. 5. t. 125. f. 2. G. radicans Lapeyr. pyr. t. 1. *E. graveolens* Lapeyr. abr. 390. (v. v.)
5. *E. TRICHOMANEFOLIUM* (L'her! ined. n. 3.) acaule, pedunculis 4-floris, foliis hirsutis subglandulosis bipinnatisectis, lobulis oblongo-linearibus, petalis obtusis calyce paulò majoribus. \mathcal{L} in Libano. Labillardière. Flores carnei lineati. (v. s.)
6. *E. CHRYSANTHUM* (L'her! ined. n. 2.) subacaule, pedunculis 3-4-floris, foliis adpressè sericeis bipinnatisectis, lobulis linearibus, petalis subrotundis calyce majoribus. \mathcal{L} in Parnasso et forsàn Olympo. Sibthorp! *E. absinthioides* fl. græc. t. 652. ex prod. 2. p. 34. distinctissimum floribus luteis. (v. s.)
7. *E. ABSINTHIÖIDES* (Willd. sp. 3. p. 627.) caulescens adscendens, pedunculis sub-4-floris, foliis subpubescentibus bipinnatisectis, lobulis linearibus acutis, rachi dentatâ, petalis obtusis calyce majoribus. \mathcal{L} in Armeniâ. — Tournef! cor. 20. n. 5. L'her! ined. n. 1. Calyx villosus-canescens. (v. s.)
8. *E. FUMARIÖIDES* (Stev! mem. soc. mosc. 4. p. 49.) caulescens adscendens, pedunculis sub-4-floris, foliis subpubescentibus bipinnatisectis, segmentis linearibus obtusis, rachi dentatâ, petalis emarginatis calyce majoribus. \mathcal{L} in Caucaso orient. ad rivulos. Petala violacea obscurius striata. An à priore differt? (v. s.)
9. *E. PULVERULENTUM* (Willd. sp. 3. p. 632.) caulescens, primò decumbens, dein erectum, pube brevi confertâ totum pulverulento-canum, pedunculis multifloris, petalis obtusis calyce paulò longioribus. \mathcal{L} An dñe hic confusæ nempe:
 α . *Tunetanum*, elatius, foliis bipinnati-sectis, lobulis tenuiter incis, pedunculis 7-8-floris. — in arenosis Caffæ. G. pulverulentum Desf! atl. 2. p. 111. *E. pulverulentum* L'her! ined. n. 4. (v. s.)
 β . *Hispanicum*, minus, foliis pinnatifidis, lobis incis dentatisve, pedunculis 4-5-floris. — in Hispaniâ ad Aranjuez, Orcehimve. G. pulverulentum Cav. diss. 5. p. 272. t. 125. f. 1. (v. s.)
10. *E. ANTHEMIDIÖFOLIUM* (Bieb. fl. taur. 2. p. 131.) caulescens ramoso-diffusum, pedunculis multifloris pube brevi subviscosâ hispidis, foliis cano-pubescentibus bipinnatisectis, lobulis linearibus, rachi dentatâ, petalis subemarginatis calyce longioribus. \mathcal{L} in Iberiâ occid. Steven! (v. s.)
11. *E. STEPHANIANUM* (Willd. sp. 3. p. 625.) caule villosa-ramoso pedunculis 2-5-floris, foliis glabriusculis bipinnatifidis, lobis decurrentibus, lobulis linearibus, petalis obtusis vix calyce longè mucronato longioribus. \mathcal{L} in Dahuriæ collibus et in desertis Kirghisorum. G. multifidum Patr! ined. Flores cærulei. Foliorum lobi inferiores ut in sequente ferè palmato-partiti. (v. s.)

12. *E. ALPINUM* (L'her! ger t. 3.) caule glabriusculo ramoso, pedunculis multifloris, foliis glabriusculis bipinnatifidis, rachi dentatâ, lobulis linearibus, petalis obtusis calyce longè mucronato longioribus. \mathcal{L} in montibus Italiæ? Græciæ. Cav. diss. 4. t. 96. f. 1. (v. s.)
13. *E. STEVENI* (Bieb. fl. taur. 2. p. 132.) caule diffuso subcanescente, pedunculis sub-3-floris, foliis glabriusculis pinnatisectis, segmentis linearibus alternis integris 3-fidisque, petalis obtusis calyce duplò longioribus. \mathcal{L} in promontorio Caucasico ad Stauropolim. Steven. (v. s.)
14. *E. STYLATUM* (Leman! in herb. DC.) caule glabriusculo diffuso, pedunculis 3-4-floris, foliis glabris bipinnati-sectis, lobulis linearibus, rachi integrâ calycibus rostrisque puberulis, stylo longissimo. \mathcal{L} Patr. ign. olim in hort. Lemonnier legi. Petalum unicum in specim. superstes calyce brevius. Species distinctissima habitu ferè Grieli. (v. s.)
15. *E. CRASSIFOLIUM* (Desf. fl. atl. 2. p. 111. *) caule ramoso diffuso puberulo, foliis crassis pinnatifido-lacinatis, lobulis linearibus, pedunculis multifloris, bracteis ovatis scariosis, aristis longissimis longè plumosis. \mathcal{L} in arenis Cafsæ. Sweet ger. 2. t. 111. Folia glabrata. (v. v.)
- β. salinarium* (Sibth! ined. in herb. L'her.) foliis pubescentibus. Juxta salinas Arnacæ in ins. Cypro. *E. crassifolium* L'her! ined. n. 5. (v. s.)
16. *E. LACINIATUM* (Cav. diss. 4. p. 228. t. 113. f. 3.) caule prostato folisque glabris bipinnati-sectis, lobulis linearibus acutis, imis sæpius trilobis, stipulis bracteis ovatis scariosis, pedunculis multifloris. (I) in arenosis maritimæ Africæ borealis à Cretâ ad Lusitaniam. — Desf! fl. atl. 2. p. 110. Variat foliis omnibus laciniatis (*G. laciniatum* L'her! ined. n. 17.) et foliis infer. trilobis (*G. diphyllum* L'her! ined. n. 18.). (v. s.)
17. *E. HIRTUM* (Willd. sp. 3. p. 650.) caule ramoso diffuso villosulo, foliis hirtis subbipinnatisectis, lobulis lanceolatis acutis, stipulis lanceolatis, pedunculis multifloris glanduloso-hirtis. \mathcal{L} in AEGYPTO. — Jacq. f. eclog. 1. p. 85. t. 58. *G. hirtum* Forsk. descr. 123.
18. *E. MILLEFOLIUM* (Willd. in H. B. et K. nov. gen. am. 5. p. 298.) caulibus procumbentibus diffusis hispidis, foliis bipinnatifidis, laciniis dentatis acuminatis subtus hispidis, stipulis ovatis, pedunculis multifloris. (I) ad radicem montis ignivomi Cotopaxi prope Mulalo Quitensiam. Fl. violacei. †.
19. *E. MORANENSE* (Willd. l. c.) caulibus procumbentibus hirtis, foliis bipinnatifidis acutis subtus hirtis, pedunculis multifloris. (I) prope Moran Mexicanorum. Fl. violacei. †.
20. *E. CICONIUM* (Willd. sp. 3. p. 629.) caule adscendente folisque subvillosis pinnatisectis, segmentis obtusis pinnatifidis dentatis, rachi inter segmenta dentatâ, pedunculis multifloris, petalis calycis longitudine, 2 submarginatis. (I) in ruderatis et agris regionis Mediterræ. Cav. diss. 4. t. 95. f. 2. Calycis sepala longè mucronata. (v. s.)
- β. erectum*, caule erecto. — in Tauriâ. (v. s.)
21. *E. CICUTARIUM* (Leman! in DC. fl. fr. 4. p. 840.) caule prostrato aut diffuso hirtulo, foliis pinnatisectis, segmentis sessilibus pinnatifidis incisis acutisve, pedunculis multifloris, petalis inæqualibus! (I) in Europæ totius, Africæ bor. et Orientis ruderatis.
- α. præcox* (Cav. diss. 5. t. 126. f. 2.) acaule, foliis patulis, segmentis subincisis, petalis cal. majoribus. — in apricis. (v. v.)
- β. pimpinellæfolium* (Cav. diss. 4. t. 126. f. 1.) caulescens demùm suberectum, foliis longè petiolatis, segmentis acutè incisis, petalis calici subæqualibus. — in pratis. (v. v.)
- γ. Chærophyllum* (Cav. diss. 4. t. 95. f. 1.) multicaule subprostratum, segmentis tenuiter pinnati-partitis, floribus pallidis aut albis. — in siccis lapidosis. (v. s.)
- δ. pilosum* (Thunb. fl. pat. ed. 2. p. 347.) multicaule subprostratum undi-

- quæ pilosum, segmentis tenuiter pinnati-partitis, floribus intense purpureis. — in arenosis. (v. v.)
1. *cicutæfolium* (Thuil. l. c. sub cicutario.) subcaulescens, segmentis oblongis obtusè incisis, floribus pallidis roseisve. — in apricis. (v. v.)
2. *bipinnatum* (Cav. diss. 5. t. 126. f. 3.) caulescens diffusum, segmentis in lobos lineares pinnati-partitis. — in Africâ bor. G. Numidicum Poir. barb. 2. p. 101. E. Petroselinum L'her. diss. n. 9. Valdè accedit ad var. *δ*. (v. s.)
22. E. ROMANUM (Willd. sp. 3. p. 630.) subacaule, foliis pinnatisectis, segmentis ovatis pinnatifidis, pedunculis multifloris, petalis calyce majoribus æqualibus. 2. secus vias Occitanicæ, Italiæ. Cav. diss. 4. t. 94. f. 2. — Barr. rar. t. 1245. Radix crassa intus subrubescens. (v. v.)
- β. *caucalisfolium* (Sweet ger. t. 6.) plantâ majore. 2. in solis pinguioribus et cultis. E. Alpinum hort. par! (v. v.)
23. E. MOSCHATUM (Willd. sp. 3. p. 631.) caule procumbente, foliis pinnatisectis, segmentis petiolulatis ovatis inæqualiter serrato-incisis, pedunculis multifloris pubescenti-subglandulosis. 1. in Europâ australi, Numidiâ, et etiam in Peru ad cap. B-Spei. — Jacq. h. vind. t. 55. Cav. diss. t. 94. f. 1. Variat glabrum et in caule costisque glanduloso-pubescentis. (v. v.)
24. E. TORDYLIÖIDES (L'her! ined. n. 14.) subacaule caulescensve, foliis villosis pinnatisectis, segmentis ovatis lobatis argutè dentatis, pedunculis multifloris longissimis. 2. in fissuris rupium Algeriæ. G. tordylioides Desf! atl. 2. p. 107. Calyces striati aristati. (v. s.)
25. E. BOTRYS (Bertol. am. it. p. 35.) caulescens setis substrigosis hispidum, foliis sinuato-pinnatifidis, lobis obtusis dentatis, pedunculis 2-4-floris, calycibus pubescentibus breviter mucronatis. 1. in Italiâ australi, Corsicâ, Mauritanîâ. E. gruinum β. Willd. sp. 3. p. 634. Ger. botrys Cav. diss. 4. t. 90. f. 2. Variat caule erecto et diffuso, foliis plus minùs incisis. Strigæ caulis partis infer. reflexæ. (v. s.)
- ** *Foliis subtrilobatis, aut indivisis.*
26. E. MURCICUM (Willd. sp. 3. p. 636.) caulibus diffusis glabris ramosis, foliis pinnatifidis subtripartitis, lobis obtusis dentatis, medio longiore, pedunculis multifloris, calycibus lineis striato. 1. in Murciâ et Africâ bor. G. Murcicum Cav. diss. 5. t. 126. f. 1. Affine E. botrydi. (v. s.)
27. E. GRUINUM (Willd. sp. 3. p. 633. excl. var. β.) caule erecto subglabro, foliis trisectis, segmentis inciso-dentatis, infimis divaricatis, medio longiore, pedunculis multifloris, calycibus striato-nervosis. 1. in Cretâ, Africâ bor. Hispaniâ. G. gruinum Lin. sp. 952. Cav. diss. 4. p. 217. t. 88. f. 2. (v. s.)
28. E. SEROTINUM (Stev! mem. ac. Petersb. 3 p. 297. t. 15. f. 2.) caulibus diffusis, foliis oppositis trisectis, segmentis lateralibus inciso-dentatis divaricatis, medio majore sublobato inciso-dentato, pedunculis multifloris. 2. in campis ad mare nigrum et ad Tyram. E. Ruthenicum Bieb. pl. rar. cent. t. 48. E. multicaule Link enum. 2. p. 184. Sweet ger. t. 137. Calyces longè aristati. (v. s.)
29. E. CHIUM (Willd. sp. 3. p. 634.) caule erecto subdiffuso, foliis glabris subcordatis superioribus 3-partitis, lobis inciso-dentatis, medio subtrifido, pedunculis multifloris. 1. in Archipelagi insulis et circa Neopolim. G. Chium Lin. sp. 951. Cav. diss. 4. t. 92. f. 1. (v. s.)
30. E. ASPLENIÖIDES (Willd. sp. 3. p. 635.) acaule villosio-pubescentis, pedunculis multifloris, foliis trisectis, segmentis obovatis obtusis inciso-dentatis, terminali sublobato. 2. in Africæ borealis montibus Shibæ. G. asplenioides Desf! fl. atl. 2. p. 109. t. 168. E. muticum L'her! ined. Calyces mutici. (v. s.)
31. E. HYMENODES (L'her! ger. t. 4.) caule erecto ramoso basi fruticoso, ramis annotinis molliter pilosis, pedunculis multifloris, foliis subtrilobis tripartitisve obtusissimis inciso-dentatis, stipulis bracteisque scariosis ovatis, caly-

- cibus muticis. \mathcal{L} in Atlantis rupium fissuris. *G. geifolium* Desf! fl. atl. 2. p. 108. *G. trifolium* Cav. diss. 4. t. 97. f. 3. *E. trilobatum* Jacq. ic. rar. t. 508. (v. v.)
32. *E. GUSSONI* (Tenor! prod. 39. Link enum. 2. p. 185.) caule erecto molliter piloso, pedunculis multifloris, foliis cordatis obtusis obtusè dentatis indivisis aut subtrilobis, stipulis bracteisque scariosis ovatis, calycibus aristatis. \mathcal{L} in pratis apricis regni Neapolitani. (v. s.)
33. *E. ARBORESCENS* (Willd. sp. 3. p. 638.) caule fruticoso erecto, foliis longè petiolatis cordatis sublobatis crenatis, stipulis ovatis subacutis, pedunculis multifloris. \mathfrak{J} in Africæ bor. montibus ad Cafsam. Facies Pelargonii. Ger. arborescens Desf! atl. 2. p. 110. (v. s.)
34. *E. MALACHOIDES* (Willd. sp. 3. p. 639.) caule herbaceo ramoso pilosiusculo, foliis cordatis indivisis trilobisve obtusis dentatis, pedunculis multifloris, petalis calycis longitudine. ① in Europà austr. Africâ bor. et Canariis. Cav. diss. 4. t. 91. f. 1. Corollæ ceruleæ. (v. v.)
Æ. ribifolium (Jacq. ic. rar. 3. t. 509.) foliis trilobis hispidulis, corollis purpureis. ① ad cap. Bonæ-Spei. (v. s.)
35. *E. INCARNATUM* (L'her! ger. t. 5.) caule fruticuloso folisque subglabris, infimis cordatis indivisis dentatis, ceteris 3-partitis, lobis cuneatis apice tridentatis, pedunculis multifloris. \mathfrak{J} ad cap. Bonæ-Spei. Cav. diss. 4. t. 91. f. 2. malè Sims. bot. mag. t. 261. Sweet. ger. t. 94. Delaun. herb. am. t. 11. Flores coccinei disco pallidiore. (v. v.)
36. *E. ARDUINUM* (Willd. sp. 3. p. 637.) caule, foliis cordatis 5-lobis crenatis obtusis, pedunculis multifloris. \mathcal{L} ad cap. Bonæ-Spei. *G. arduinum* Lin. sp. 952.
37. *E. GLAUCOPHYLLUM* (Ait. h. kew. ed. 1. v. 2. p. 416.) caule erecto folisque glabris oblongis sublobatis crenatis subcarnosis, pedunculis multifloris, aristis à medio ad apicem longè plumosis. ① in Ægypto prope Memphim. Cav. diss. 4. t. 92. f. 2. *G. crassifolium* Forsk. descr. 123. (v. s.)
38. *E. GUTTATUM* (Willd. sp. 3. p. 636.) caule prostrato, foliis cordatis oblongis subtrilobis dentatis incanis, pedunculis subtrifloris, calycibus longè aristatis. \mathcal{L} in arenosis Africæ bor. prope Belide. Ger. guttatum Desf! atl. 2. p. 113. t. 269. Petala magna violacea basi maculata obtusa. (v. s.)
39. *E. OXYRACHNUM* (Bieb. fl. taur. 2. p. 133.) caule prostrato, foliis cordatis oblongis subtripartitis crenatis subtus incanis, lobo extimo trifido, pedunculis subtrifloris. \mathcal{L} in Iberiæ orient. collibus siccis. Flores magn. *E. cicutariæ*, petalis oblongis. Affine *E. guttato*.
40. *E. HELIOTRPIOIDES* (Willd. sp. 3. p. 638.) caule suffruticoso hispido, foliis ovatis subrotundis crenatis villosis, pedunculis multifloris, aristis longissimis longè plumosis. \mathfrak{J} Patr. ign. Ger. heliotropioides Cav. diss. 4. p. 220. t. 113. f. 2.
41. *E. NERVULOSUM* (L'her! ined. n. 24.) caule diffuso brevi, foliis cordatis subindivisis dentatis crassis nervosis pubescenti-subincanis, stipulis ovatis membranaceis, pedunculis multifloris. \mathcal{L} in Siciliâ. — Bocc. mus. 2. t. 128. fig. dextr. Affine *E. laciniato*. (v. s.)
42. *E. MALOPOIDES* (Willd. sp. 3. p. 640.) caulescens erectum aut subcaule totum molliter cinereo-velutinum, foliis ovatis subcordatis crenatis interdum sublobatis, pedunculis 3-4-floris, aristis barbatis. \mathcal{L} in arenosis maritimis Ægypti, Algeriæ et Siciliæ. *G. malopoides* Cav. diss. 4. t. 90. f. 1. (v. s.)
Æ. Corsicum (Leman in DC. fl. fr. 4. p. 842.) pedunculis subbifloris, foliis magis tomentosis. \mathcal{L} in Corsicæ rupibus maritimis. (v. s.)
43. *E. LITTOREUM* (Leman! in DC. fl. fr. 4. p. 842.) caulescens diffuso-prostratum glabriusculum, foliis cordato-rotundis trilobis inæqualiter crenatis, pedunculis multifloris elongatis, aristis barbatis. \mathcal{L} in Narbonæ et Liguriæ arenosis maritimis. Sequenti valde affine. (v. s.)
44. *E. MARITIMUM* (Smith fl. brit. 2. p. 728.) caulescens diffuso-prostratum

glabriusculum, foliis cordatis ovatis inciso-crenatis pubescentibus, pedunculis uni aut sæpius bifloris, aristis imberbibus. \mathcal{U} in Angliæ et Galliæ maritimis. (v. s.)

ℓ. *Bocconi*, subcaulescens cæspitosum pumilum, foliis ovatis inciso-crenatis, pedunculis 1-rariis-2-floris, petalis vix calycis longitudine. \mathcal{U} in Corsicæ montibus. Saltzman. *Geranium minimum* chamædryoides Bocc. mus. 2. p. 160. t. 128. Flores pallidè rubentes. Planta nunc minima subcaulis, nunc ex cl. Saltzmann interdum pedalis. (v. s.)

45. E. REICHARDI, subcaule, foliis cordatis crenatis obtusis glabriusculis, pedunculis 1-floris, petalis calyce majoribus. \mathcal{U} in Majorcâ. G. Reichardi. Murr. comm. gætt. 1780. p. 11. t. 3. G. parvulum Scop. ins. 8. t. 3. f. B. — E. chamædryoides L'her! ger. t. 6. excl. syn. Bocc. et ideò nomine. (v. v.)

V. PELARGONIUM. L'her. ger. icon. et text. ined. Ait. h. kew. 2. p. 417. — Pelargoniceæ Sweet ger. 1. p. VIII.

Cal. 5-partitus, lacinia supremâ desinente in calcar seu tubulum nectariferum tenue sæcùs pedunculum decurrentem et huic adnatum. Pet. 5, rariùs 4, plus minùs irregularia. Filam. 10 inæqualia monadelphæ, 4-7 tantum fertilia, cetera castrata. Rostra seu styli introrsum barbati et more Erodii spiralliter contorti. — Species omnes, exceptis paucis passim indicatis, ad Caput Bonæ-Spei crescunt. — Genus vastissimum exhibens typos plurimos sat equidem inter se diversos, sed ut sectiones (nec genera) meo sensu habendas, ob proles hybridæ facili nascentes è mixturâ specierum sectionibus diversis pertinentium. Ex eadem causâ species pleræque in hortis natæ, inter se nimis affines et extricatu difficillimæ, hic partim ex Sweetii laboribus optimè inchoatis partim ex autopsiâ et habitu ordinatæ, sed stirpes plurimas in hortis obvias omisi, ne è phrasibus specificis icone non suffultis confusio nova oriatur.

Sect. I. HOAREA. Sweet ger. n. 18. et 72.

Petala 5 rariùs 2-4 oblongo-linearia, 2 super. parallela longè unguiculata medio abruptè reflexa. Stam. longè tubulosa petalorum inf. longitudine, 5 rariùs 2-4 antherifera, cetera sterilia recta aut apice incurva, 3 infer. fertilibus breviora. — Herbæ acaules, radice tuberoso-rapacæ, foliis radic. petiolatis.

* *Foliis oblongis integris lobatisve, lobis integris aut vix dentatis.*

1. P. LONGIFOLIUM (Jacq. ic. rar. 3. t. 518.) acaule, foliis lanceolatis integerrimis acutis glabris, senioribus pinnatifidis linearibus, umbellâ compositâ, floribus tetrandris, petalis obtusis lanceolatis, superioribus ovatis. \mathcal{U} G. acaule Burm. ger. 67. t. 2. Cav. diss. 4. t. 102. f. 1. Petala rosea. (v. s.)

2. P. LONGIFLORUM (Jacq. ic. rar. 3. t. 521.) acaule, foliis lanceolatis integerrimis acutis glabris, umbellâ compositâ 4-florâ, floribus tetrandris, petalis linearibus. \mathcal{U} Petala ochroleuca, sup. lineâ rubrâ notata.

ℓ. *depressum* (Jacq. ic. rar. 3. t. 520.) umbellâ 8-florâ, pedicellis demum subreflexis, filamentis 9. \mathcal{U} Pet. ochroleuca, sup. medio purpurea.

3. P. OVALIFOLIUM, foliis ovalibus aut ovali-oblongis obtusis planis aut margine involutis integerrimis hirsutis, umbellis simplicibus compositisve, petalis linearibus undulatis tortis. \mathcal{U} Petala alba, 2 super. ad basin medio lineâ pallidâ rubrâ; tubus nectarifer sessilis. An. ex Sweet, G. undulatum Andr. bot. rep. t. 292? Hoarea ovalifolia Sweet ger. t. 106.

4. P. RETICULATUM, acaule, foliis elliptico-lanceolatis oblongisve integerrimis pilosis margine revolutis, umbellâ compositâ, floribus pentandris, petalis lineari-spathulatis undulatis reflexis. \mathcal{U} Hoarea reticulata Sweet ger. t. 91. Petala rosea, lineis purp. reticulata, 2 sup. medio maculâ intensè purpureâ notata. — P. scapifolium L'her. ger. ined. n. 8?

5. *P. PARNASSIOIDES* (L'her! ger. ined. n. 1.) acaule, foliis ovatis obtusis integerrimis ciliatis, umbellâ compositâ, floribus pentandris, petalis lineari-subspathulatis. \mathcal{U} *G. ciliatum* Cav. diss. 4. p. 234. t. 118. f. 2. *P. ciliatum* Willd. sp. 3. p. 643. excl. L'her. syn. Pet. immaculata pallida. (v. s.)
6. *P. CILIATUM* (L'her! ger. t. 7. ined. n. 2.) acaule, foliis ovatis acutis integerrimis subciliatis, umbellâ compositâ, floribus pentandris, petalis lineari-subspathulatis. \mathcal{U} Petala carnea, 2 sup. medio maculata. (v. s. sine fl.)
 β . *hirtum* (Willd. sp. 3. p. 644.) foliis subtrilobis. *G. pilosum* Cav. diss. 5. p. 273. 6. t. 199. Folia ex L'her. mss. ficta, aut var. hortensis.
7. *P. PUNCTATUM* (Willd. sp. 3. p. 645.) acaule, foliis ovatis dentatis glabris, umbellâ compositâ, floribus diandris, petalis linearibus, 3 inferioribus dimidiò brevioribus. \mathcal{U} *G. punctatum* Andr. rep. t. 60. *P. punctatum* Pers. ench. n. 14 et 22. Petala ochroleuca, 2 sup. basi sanguineo-punctata.
8. *P. DIPETALUM* (L'her! ger. t. 43.) acaule, foliis ovatis integerrimis acutis glabris, umbellâ simplici, floribus pentandris dipetalis. \mathcal{U} Petioli pilis brevibus ciliati. Petala 2 super. obovata, dilutè purpurascens, basi maculata, inferiora nulla. (v. s.)
9. *P. RADICATUM* (Vent. malm. t. 65.) acaule, foliis ovali-oblongis integerrimis utrinquè acutis glabris ciliatis, umbellâ simplici, floribus pentandris, petalis lineari-oblongis retusis, tubo nectarifero calyce quadruplò longiori. \mathcal{U} *G. ciliatum* Andr. rep. t. 247. *P. concavifolium* Pers. ench. 2. p. 226. Hoarea radicata Sweet ger. t. 174. Petala flavescentia immaculata.
10. *P. SPATHULATUM* (Andr. rep. t. 152.) acaule, foliis oblongis subspathulatis obtusis glabris, umbellâ compositâ, floribus pentandris, petalis linearibus obtusis subrevolutis. \mathcal{U} Petala super. macula duplici lineari notata.
 β . *affine* (Andr. l. c. t. 282.) foliis ovato-lanceolatis acutis subciliatis, scapo gracili.
11. *P. RADIATUM* (Pers. ench. 2. p. 226.) acaule, foliis elliptico-spathulatis integerrimis glabris, umbellâ compositâ, floribus pentandris, petalis cuneiformibus. \mathcal{U} *G. radiatum* Andr. rep. t. 222. Petala flava, super. latiora basi maculâ duplici lineari notata.
12. *P. VIRGINEUM* (Pers. ench. 2. p. 226.) acaule, foliis elliptico-ovatis utrinquè acutis glabris, umbellâ subcompositâ, floribus pentandris, petalis lanceolato-cuneatis æqualibus undulatis. \mathcal{U} *G. undulatum* Andr. rep. t. 317.
13. *P. UNDULATUM* (Ait. h. kew. ed. 2. v. 4. p. 160.) acaule, foliis lineari-lanceolatis integerrimis ciliatis, umbellâ simplici, floribus pentandris, petalis undulatis subæqualibus. \mathcal{U} *G. undulatum* Andr. rep. t. 292. Pet. super. maculâ marginali notata.
14. *P. LINEARE* (Pers. ench. 2. p. 228.) acaule, foliis lineari-lanceolatis repandis, umbellâ subsimplici, floribus pentandris, petalis linearibus. \mathcal{U} *G. lineare* Andr. rep. t. 193. Petala longissima, flavescentia, omnia basi maculata.
 ** *Foliis sagittatis cordatis trilobis aut basi appendiculatis.*
15. *P. OXALOIDES* (Willd. sp. 3. p. 642.) acaule, foliis oblongis sagittatis integerrimis carnosis glabris, umbellâ compositâ. \mathcal{U} *G. oxaloides* Burm. ger. 71. t. 2. Cav. diss. 4. t. 97. f. 1. — *G. oxaloides* Andr. videtur diversum.
16. *P. CHELIDONIUM* (Houtt! pfl. syst. 8. p. 398. t. 61. f. 1.) acaule, foliis subrotundis basi truncatis acutis integerrimis pubescentibus, umbellâ compositâ. \mathcal{U} *P. Ficaria* Willd. sp. 3. p. 643. *P. deltoideum* L'her! ined. n. 111. (v. s.)
17. *P. VELUTINUM* (L'her! ger. ined. n. 82.) acaule? foliis profundè cordatis obtusissimis indivisis crenato-subsinuatis subtus cano-tomentosis, umbellâ compositâ. \mathcal{U} Burch! cat. n. 2828. Petala in sicco linearia subundulata atrata. (v. s.)
18. *P. BIFOLIUM* (Willd. sp. 3. p. 645.) acaule, foliis binis cordatis subacutis inciso-dentatis, umbellâ simplici. \mathcal{U} Burm. afr. t. 35. f. 1. *G. bifolium* Cav.

diss. 4. p. 254. t. 115. f. 3. Petala oblongo-cuneata, 2 sup. medio maculâ oblongâ notata. †.

19? *P. STIPULACEUM* (Willd. sp. 3. p. 655.) caule brevissimo stipularum rudimentis tecto, foliis subcordatis ovatis inciso-dentatis villosis, umbellâ simplici pauciflorâ. 2. *G. stipulaceum* Cav. diss. 4. p. 254. t. 122. f. 3. An flores pentandri? An hujus sectionis? An *Campylia*?

20? *P. ARTICULATUM* (Willd. sp. 3. p. 655.) caule brevissimo stipularum rudimentis tecto, foliis reniformibus 5-lobis villosis, lobis subtrilobis, umbellâ simplici pauciflorâ. 2. *G. articulatum* Cav. diss. 4. p. 252. t. 122. f. 1. Priori certè affine, sed dubium quoad staminum numerum.

21. *P. REVOLUTUM* (Pers. ench. 2. p. 226.) acaule, foliis cordatis obtusis nervosis integerrimis basi sæpe biauriculatis, umbellâ compositâ, involucri foliolis revolutis. 2. *G. revolutum* Andr. rep. t. 354. Flores purpurei lineati.

22. *P. AURICULATUM* (Willd. sp. 3. p. 643.) acaule, foliis oblongo-lanceolatis utrinquè acuminatis hirsutis margine ciliatis, nunc integerrimis, nunc sæpius basi appendicibus duabus oblongo-linearibus auctis, umbellâ compositâ. 2. *P. ciliatum* Jacq. ic. rar. 3. t. 519. Petala linearia albedo-rubrescentia.

23. *P. AURITUM* (Willd. sp. 3. p. 644.) acaule, foliis oblongis obtusis nunc integerrimis nunc basi appendicibus duabus oblongo-linearibus auctis, umbellâ subcompositâ. 2. *G. auritum* Linn. mant. 433. — Comm. hort. 2. t. 61. Flores ex rubro purpurascens.

24. *P. TRIFIDUM* (Willd. sp. 3. p. 646.) acaule, foliis tripartitis, laciniis lineari-cuneiformibus apice tridentatis, umbellâ simplici. 2. Burm. afr. t. 35. f. 2. *G. trifidum* Cav. diss. 4. t. 115. f. 1. Flores sanguinei.

25? *P. LACINIATUM* (Pers. ench. 2. p. 228.) acaule, foliis integris et apice inciso-lobatis, scapo flexuoso, umbellâ compositâ. 2. *G. laciniatum* Andr. rep. t. 131.

26. *P. HETEROPHYLLUM* (Jacq. ic. rar. 3. t. 516.) acaule, foliis ciliatis indivisis trilobis trisectisve, segmento medio trilobo, umbellâ compositâ. 2. Petala alba, 2 super. subretusa, basi maculâ sanguineâ notata.

27. *P. OXALIDIFOLIUM* (Pers. ench. 2. p. 227.) acaule, foliis ciliatis trisectis, segmentis ovatis obtusis, umbellâ compositâ. 2. *G. oxalidifolium* Andr. rep. t. 300. Petala ochroleuca, 2 snp. basi sanguineo maculata.

28. *P. NERVIFOLIUM* (Jacq. ic. rar. 3. t. 517.) acaule, foliis glabris trisectis, segmentis obtusis sublobatis nervosis subtus glaucis, scapis hispidis, umbellâ compositâ. 2. Petala oblongo-cuneata alba basi lineâ sanguineâ ramosâ picta.

29. *P. TRIPHYLLUM* (Jacq. ic. rar. 3. t. 515.) acaule, foliis glabris trisectis, segmentis obtusis crenatis, scapis petiolisque pubescentibus, umbellâ compositâ. 2. Petala linearia incarnata, 2 super. basi sanguineo maculata.

30. *P. REFLEXUM* (Pers. ench. 2. p. 227.) subacaule, foliis trisectis, segmentis lobato-incisis recurvis, umbellâ simplici, filamentis 2 super. stigmatibusque reflexis. 2. *G. reflexum* Andr. rep. t. 224. Petala albida.

31. *P. ROSEUM* (Ait. h. kev. ed. 2. v. 4. p. 181.) acaule, foliis inciso-lobatis tomentosis, umbellâ simplici confertâ, petalis 3 infer. multò minoribus. 2. *G. roseum* Andr. rep. t. 173. *P. condensatum* Pers. ench. 2. p. 227. Flores rosei unicolores.

*** *Foliis pinnatisectis, segmentis incisis multifidisve.*

32. *P. RAPACEUM* (Jacq. ic. rar. 3. t. 510.) acaule, foliis pilosis bipinnatisectis, lobulis linearibus subobtusis, petalis sup. refractis, infer. conniventibus. 2. *G. Selinum* Andr. rep. t. 239. Petala ex albo rosea, sup. subpunctata.

33. *P. NUTANS*, subacaule, foliis bipinnatisectis hirsutis, lobis pinnatifido-laciniatis multifidis linearibus subdentatis, umbellis capitatis congestis

- depressis, floribus nutantibus, petalis super. refractis, infer. concavis conniventibus. \mathcal{U} G. Selinum flavum Andr. ger. excl. foliis ex Sweet. Pel. rapaceum luteum Sims bot. mag. t. 1877. Hoarea nutans Sweet ger. t. 135.
34. P. CORDALIFLORUM (Sweet ger. t. 18. sub Hoarea.) acaule, foliis pilosis pinnatisectis. segmentis pinnatifidis trifidisve, lobulis linearibus acutis, petalis sup. refractis, inferioribus conniventibus. \mathcal{U} G. Selinum flavum Andr. excl. flor. ex Sweet. Petala pallide sulphurea, 2 sup. sanguineo maculata. An prioris var.?
35. P. BARBATUM (Jacq. ic. rar. 3. t. 513.) acaule, foliis pinnatisectis, segmentis trifidis, lobulis linearibus acuminatis apice barbatis, umbellâ compositâ, petalis linearibus obtusis. \mathcal{U} G. proliferum Burm. ger. 70. t. 2. Cav. diss. 4. t. 120. f. 3. G. barbatum Andr. rep. t. 303. Petala carnea, 2 sup. lineis duabus rubris medio picta.
36. P. FISSIFOLIUM (Pers. ench. 2. p. 227.) acaule, foliis pinnatisectis, segmentis trifidis apice incisus nudis, umbellâ simplici, petalis obtusis omnibus maculâ oblongâ notatis. \mathcal{U} G. fissifolium Andr. rep. t. 378. simile G. barbato.
37. P. SETOSUM (Sweet ger. t. 38. sub Hoarea.) acaule, foliis pinnatisectis pubescentibus, segmentis cuneatis apice 3-5-dentatis, dentibus apice setosis, umbellâ compositâ, petalis sup. refractis, inferioribus subconniventibus. \mathcal{U} Petala rosea, 2 sup. maculâ rubrâ notata.
38. P. BUBONIFOLIUM (Pers. ench. 2. p. 227.) acaule, foliis glabris pinnatisectis, segmentis inciso-lobatis acutis, umbellâ simplici, petalis emarginatis. \mathcal{U} G. bubonifolium Andr. rep. t. 328. Petala alba, 2. sup. basi maculâ purpureâ.
39. P. VIOLEFLORUM, subcaulescens, foliis pinnati-partitis trisectisve, segmentis oblongo-lanceolatis glabris integerrimis margine ciliatis apice acuminatis penicilliferis, inferioribus bifidis, petiolis hispidis, umbellâ compositâ, petalis reflexis, inferioribus multò minoribus. \mathcal{U} Flores albi. Cal. lobus superior erectus, ceteri reflexi. Hoarea violæflora Sweet ger. t. 123.
40. P. FLORIBUNDUM (Ait. h. kew. ed. 2 v. 4. p. 163.) acaule, foliis pinnatisectis, segmentis bipartitis, umbellâ compositâ. \mathcal{U} G. floribundum Andr. rep. t. 420. Pet. super. maculis 3 lunatis, infer. maculâ lineari notata.
41. P. PILOSUM (Pers. ench. 2. p. 227.) acaule, foliis pinnatisectis hirsutis, segmentis incisus multifidis, umbellâ simplici 4-6-florâ, petalis linearibus. \mathcal{U} G. pilosum Andr. rep. t. 249.
42. P. PENNIFORME (Pers. ench. 2. p. 227.) acaule, foliis pinnatipartitis, lacinii lanceolato-linearibus, umbellâ compositâ. \mathcal{U} G. laciniatum Andr. rep. t. 269. Petala flava basi rubra. Folia radicalia interdum integra. An potius P. longifolio affine?
43. P. PURPURASCENS (Pers. ench. 2. p. 227.) acaule, foliis lanceolato-linearibus integerrimis pinnatifidisque, umbellâ compositâ. \mathcal{U} G. laciniatum Andr. rep. t. 204. Pedicelli elongati. Flores purpurascens. Priori affine.
44. P. CENTAUROIDES (L'her! ger. ined. n. 12.) acaule, foliis hirsutis pinnatipartitis, lacinii remotis incisus integrisve, umbellâ simplici confertâ multiflorâ. \mathcal{U} Pedicelli elongati et flores purpurascens ut in priore, cui forsâ nimis affine. (v. s.)
45. P. HIRsutUM (Ait. h. kew. ed. 1. v. 2. p. 417.) acaule, foliis hirsutis ciliatis obovatis lanceolatisve integerrimis pinnatifidisve, stipulis petiolo adnatis, umbellâ compositâ. \mathcal{U} G. hirsutum Cav. diss. 4. p. 247. t. 101. f. 2. P. heterophyllum. L'her. ger. ined. n. 10. Petala carneo-albida venis sanguineis, super. maculâ magnâ atro-rubente notata.
46. P. SELINIFOLIUM, acaule, foliis pinnatisectis pilosis, segmentis pinnatifidis incisive, lobulis oblongo-lanceolatis subdentatis obtusiusculis, umbellâ compositâ, calycibus reflexis, petalis super. refractis, infer. patentibus. \mathcal{U}

Hybridum dicitur *P. malantheri* ab alterâ spec. ign. fœcundati. *Hoarea selinifolia* Sweet ger. t. 159.

47. *P. MELANANTHUM* (Jacq. ic. rar. 3. t. 514.) subacaule, foliis hirtis pinnati-partitis, lacinii ovali-oblongis obtusis subpinnatifidis dentatisve, umbellâ compositâ, petalis linearibus obtusis. \mathcal{U} *Hoarea melanantha* Sweet ger. t. 73. Folia primordialis ovata aut triloba. Petala fusco-atra. Cal. lacinia super. recta. Filam. sterilia apice incurva. Vix à *P. atro* differt. (v. v. in h. Walner.)
48. *P. dioicum* (Ait. h. kew. ed. 2. v. 4. p. 162.) acaule, foliis hispidis integris trisectisque, umbellâ compositâ, floribus dioicis. \mathcal{U} *G. melananthum* Andr. rep. t. 209. Petala atro-fusca.
49. *P. atrum* (L'her! ger. t. 44.) acaule, foliis puberulis aliis oblongis integris, aliis pinnati-partitis, umbellâ compositâ, calycis lacinia superiore erectâ, petalis linearibus, filamentis sterilibus apice incurvis. \mathcal{U} *Hoarea atra* Sweet ger. t. 72. Petala atro-fusca, ungue albidâ. (v. s.)
 β . *varium*, *Hoarea varia* Sweet ger. 2. t. 166. Hybridum dicitur ex *P. melanantho*, sed ex calyce et staminibus incurvis ad *P. atrum* pertinet.
50. *P. ATROSANGUINEUM*, acaule, foliis infer. ovatis, super. pinnati-lobatis utrinque hirsutis, lobis oppositis alternisque oblongo-ovatis obtusis integerrimis, umbellâ compositâ, petalis patentibus. \mathcal{U} Flores pulchrè atrosanguinei. Hybridum dicitur ex *P. vicifolium* à *P. melanantho* fœcundato. *Hoarea atrosanguinea* Sweet ger. t. 151.
51. *P. SWERTIANUM*, acaule, foliis infer. ovatis obliquis trisectisque, super. pinnatisectis pilosis, segmentis oppositis oblongo-ovatis obtusis, umbellâ compositâ, pedunculis elongatis filiformibus, tubo nectarifero calyce triplò longiore. \mathcal{U} Flores lætè purpurei venis superiorum intensioribus. Hybridum dicitur *P. vicifolium* à *P. atro* fœcundati. *Hoarea elegans* Sweet ger. 2. t. 132. Omnino diversum à n. 184.

Sect. II. DIMACRIA. *Lindl. in Sweet ger. n. 46.*

Petala 5 inæqualia, 2 super. conniventia apice divaricata. Stamina sepalis breviora, 5 fertilia, 2 inferioribus duplò longioribus porrectis, supremo brevissimo, 5 sterilia minima subæqualia. — Herbæ acaules radice tuberoso-rapacèâ, foliis petiolatis pinnatisectis.

* *Foliis impari-pinnatisectis, segmentis integris.*

52. *P. VICIFOLIUM* (L'her! ger. ined. n. 15.) acaule, foliis pinnatisectis villosis, segmentis ovatis 2-4-jugis, petalis subintegris planis. \mathcal{U} *G. pinnatum* Cav. diss. 4. t. 115. f. 2. non Lin. Dimacria pinnata Sweet ger. t. 46. Petala albo-subrosea, 2 super. lineolata punctis sanguineis basi maculata.
53. *P. BIPARTITUM*, acaule, foliis pinnatisectis pilosis, segmentis oppositis verticillatisque oblongo-lanceolatis bipartitis integrisque, umbellâ compositâ, petalis super. refractis basi maculatis, tubo nectarifero recurvo calyce duplò longiore. \mathcal{U} Flores ochroleuci, pet. 2. super. purpureis maculis notata. Hybridum dicitur *P. vicifolium* à *P. corydalifloro* fœcundati.
54. *P. FOLIOLOSUM*, acaule, foliis pinnatisectis, segmentis 15-20-jugis. \mathcal{U} *G. pinnatum* Andr. rep. t. 311. Dimacria foliolosa Sweet ger. n. 103. in adn. †.
55. *P. ASTRAGALIFOLIUM* (Pers. ench. 2. p. 227.) acaule, foliis pinnatisectis hirsutis, segmentis ellipticis multijugis, petalis undulatis basi tortis. \mathcal{U} *G. astragalifolium* Cav. diss. 4. t. 104. f. 2. *G. pinnatum* Linn. sp. 677. ex L'her! ger. ined. n. 14. — Comm. præl. 53. t. 3. Dimacria *astragalifolia* Sweet ger. t. 103. Petala albida angusta, 2 super. purpureo-maculata. *G. astragalifolium* Andr. rep. t. 190. forsân species fictitia ex cl. Sweet.
56. *P. CORONILLÆFOLIUM* (Pers. ench. 2. p. 227.) acaule, foliis pinnatisectis glabris, segmentis 1-2-jugis obovatis oblongisve. \mathcal{U} *G. coronillæfolium* Andr. rep. t. 305. Petala fulva, super. rubro maculata spathulato-retusa, infer. lanceolata.

57. *P. HERACLEIFOLIUM* (Lodd. cab. t. 437.) acaule, foliis pinnatisectis glabris, segmentis 2-3-jugis obovatis ultimis, confluentibus. 2 Petala obovato-cuneata atro-fusca basi et margine subflavida.

** *Foliis impari-pinnatisectis, segmentis lobatis multifidisve... An vera Dimacrice?*

58. *P. INCRASSATUM* (bot. mag. t. 761.) subacaule, foliis glabris pinnati-partitis, laciniis lobatis obtusis, scapo subramoso, petalis sup. obcordatis. 2 *G. incrassatum* Andr. rep. t. 246. Flores magni rosei venis obscurè rubris reticulatis picti.

59. *P. CARNEUM* (Jacq. ic. rar. 3. t. 512.) acaule, foliis glabris bipinnatisectis, lobis trifidis linearibus obtusiusculis, scapo simplici. 2 *G. pinuatifidum* Cav. diss. 4. p. 260. t. 120. f. 1. Petala albo-rosea venis rubris reticulatis picta.

Sect. III. CYNOSBATA.

Petala subovalia, inter se subæqualia, calyce duplò ferè longiora; stam. 10 erecta, 5 alterna antherifera. — Caules fruticosi erecti.

60. *P. MALVÆFOLIUM* (Jacq. f. ecl. 1. p. 145. t. 97.) caule basi fruticoso, ramis divaricato diffusis puberulis, foliis cordatis suborbiculatis 7-9-lobis grossè serratis utrinquè pubescentibus azonatis, umbellis 5-8-floris. 5 2 Petala cuneata carnea venis saturatoribus purpurea reticulata ferè ut in *P. carneo* et *incrassato*. Tubus nectarifer obsoletus.

61. *P. LATERTIUM* (Willd. enum. suppl. 47.) caule basi fruticoso, foliis cordatis 5-lobis pilosis zonatis, lobis apice acutè dentatis, umbellis 4-5-floris. 3 Petala lateritia non lineis striata minora quàm in *P. malvæfolio*. Vide florem in Jacq. ecl. 1. t. 97. fig. ult. qui pro varietate prioris habet.

62. *P. CYNOSBATIFOLIUM* (Willd. h. berol. t. 78.) caule fruticoso ramoso, foliis cordatis trilobis dentatis pilosis, lobo medio trilobo, pedunculis bifloris, tubo nectarifero brevissimo. 3 Flores intensè rosei ex ic., lilacini ex descr., petalo inferiore basi lineis ramosis striato.

Sect. IV. PERISTERA.

Petala inter se subæqualia calyci æqualia, aut paulò majora. Stam. 10, 5 longiora subæqualia fertilia aut uno castrato, 5 alterna brevissima sterilia dentiformia. — Herbæ caulescentes Geranii aut Erodii facie.

63. *P. COLUMBINUM* (Jacq. schœnb. 2. p. 4. t. 133.) multicaule diffuso-procumbens, foliis cordato subrotundis multipartitis, lobis trifidis, lobulis linearibus obtusis, pedunculis multifloris, floribus 4-antheris 2 *G. alceoides* Lin. sp. 948? Facies *Ger. columbini*. Petala purpurea oblongo-linearia.

64. *P. PROCUMBENS* (Pers. ench. 2. p. 229.) caulescens procumbens, foliis cordatis sublobatis crenato-dentatis, pedunculis subbifloris, floribus 4-antheris. 2? Patria ign. *G. procumbens* Andr. rep. t. 254. Petala parvula, 2 sup. albida, 3 inf. purpurascens, omnia mediò maculata. An à sequente satis differt?

65. *P. HUMIFUSUM* (Willd. enum. suppl. p. 47.) multicaule procumbens, foliis cordatis sub-3-partitis 5-lobisve dentatis, pedunculis 3-5-floris, floribus 4-5-antheris. ① Jacq. ecl. 1. t. 99. Sweet ger. t. 42. *G. decumbens* Balb. cat. h. taur. 1813. app. 1. p. 11. Petala parva, omnia mediò maculata, 2 sup. albida, 3 inf. purpurascens.

66. *P. CHAMÆDRYFOLIUM* (Jacq. ic. rar. 3. t. 523.) ramosissimum procumbens, foliis ellipticis obtusis incano-pubescentibus dentatis, pedunculis subbifloris, floribus 5-antheris. Petala calycis longitudine, alba, mediò sanguineo-maculata.

67. *P. AUSTRALE* (Willd. sp. 3. p. 675.) diffuso-procumbens, foliis cordatis sublobatis subtus villosis, pedunculis multifloris, floribus 5-antheris. 2? in Novâ-Hollandiâ. Jacq. f. ecl. 1. p. 149. t. 100. Petala rosea, 2 sup. maculis dendriticis rubris notata.

68. *P. ALTHÆOIDES* (L'her. ger. t. 10.) diffuso-decumbens, foliis cordato-ovatis villosis trilobis dentatis, summis sinuatis, umbellis multifloris, floribus 5-antheris. \mathcal{U} Jacq. coll. 4. p. 185. t. 21. f. 2. Ger. althæoides Cav. diss. 4. t. 123. f. 2. Pet. alba, medio rubro maculata, cuneiformia, calyci æqualia.

Sect. v. OTIDIA. *Lindl. in Sweet ger. p. 8. n. 98.*

Petala oblongo-linearia subæqualia calyce duplò circiter longiora, 2 sup. supernè basi auriculata. Stam. 10 erecta, 5 antherifera, 2 sup. spathulata aut subulata, 3 inf. breviora. — Caules fruticoso-carnosi. Folia alterna pinnatisecta carnosa. Flores albidì.

69. *P. CERATOPHYLLUM* (L'her. ger. t. 13.) caule fruticoso carnoso ramoso, foliis carnosìs pinnati-partitis, lobis lineari-teretibus integris ant apice 3-dentatis subcanaliculatis, pedunculis multifloris. \mathfrak{z} Curt. bot. mag. t. 315. Petala albidia lineari-lanceolata æqualia immaculata.

70. *P. DASYCAULON* (Sims bot. mag. t. 2029.) caule fruticoso-carnoso tuberculato, foliis carnosìs pinnatisectis, segmentis inciso-pinnatifidis apice subtrifidis, pedunculis 3-floris. \mathfrak{z} Patr. subdubia. *P. dasycaule* Haw. succ. 309. Petala linearia alba.

71. *P. FERULACEUM* (Willd. sp. 3. p. 687.) caule fruticoso-carnoso, foliis pinnatisectis, segmentis inciso-dentatis undulatis, pedunculis multifloris, petalis acutis longitudine calycis. \mathfrak{z} Burm. afr. t. 36. f. 1. *G. ferulaceum* Cav. diss. 4. t. 110. f. 2.

72. *P. CRITHMIFOLIUM* (Smith ic. pict. 1. t. 13.) caule fruticoso-carnoso, foliis carnosìs bipinnatisectis, lobis apice dilatatis incisìs, pedunculis multifloris paniculatis, petalis obtusis calyce longioribus, super. basi crispis. \mathfrak{z} *P. paniculatum* Jacq. schœnb. 2. t. 137. Petala alba, 2 basi punctis rubris notata.

73. *P. ALTERNANS* (Wendl. h. herrenh. 1. p. 14. t. 10.) caule fruticoso subcarnoso, ramis pilosis, foliis pinnatisectis, segmentis petiolulatis subalternis cuneiformibus apice inciso-dentatis, umbellis paucifloris. \mathfrak{z} *P. alternatim-pinnatum* Wendl. obs. 53. Petala alba, 2 sup. apice lineis rubris binis notata.

74. *P. CARNOSUM* (Ait. h. kew. 2. p. 421.) caule crasso carnoso basi suffruticoso, foliis glabris crassis sinuato-pinnatifidis, laciniis oblongis obtusis apice inciso-dentatis, umbellis multifloris, petalis linearibus, tubo nectarifero calyce sesquilingiori. \mathfrak{z} — Dill. elth. 1. p. 153. t. 127. f. 154. Ger. carnosum Lin. sp. 946. Cav. diss. t. 99. f. 1. *Otidia carnosata* Sweet ger. t. 98. Flores albidì. (v. s.)

Sect. VI. POLYACTIUM.

Cal. lacinie subæquales revolutæ. Pet. 5 subæqualia obovata. Stam. filamenta 10, 5 antherifera, 4 infer. longiora subulata, superius latius spathulatum apice reflexum, fertilia breviora apice incurva. — Petala omnia maculâ atrofusca maximâ notata vix flavo marginata.

75. *P. MULTIRADIATUM* (Wendl. coll. ic.) subcaulescens, foliis inferioribus pinnatisectis hirsutis, segmentis pinnatifidis, lobis oblongis obtusis inciso-dentatis, super. glabriusculis bipinnatifidis, umbellis 20-30-floris, tubo nectarifero calyce quadruplò longiori. \mathcal{U} Radix tuberosa. Sweet ger. t. 145. An *P. apiifolium* ad hanc sect. referendum? certè huic flore valde affine.

Sect. VII. ISOPETALUM. *Sweet ger. n. 126.*

Calycis lacinia superior basi in foveolam nectariferam nec in tubum desinens. Pet. 5 æqualia. Stam. filamenta 10 in tubum brevissimum monadelpha, 5-6 fertilia patentia apice incurva, sterilia inæqualia subulata incurva. — Frutex caule carnoso.

76. *P. COTYLEDONIS* (L'her! ger. t. 27.) caule crasso carnoso ramoso nudo, foliis cordatis subpeltatis rugosis pubescentibus subtus tomentosis reticulato-venosis, umbellis compositis. \mathfrak{z} in ins. Sanctæ-Helenæ. Ger. cotyledonis Lin.

mant. 569. Isopetalum cotyledonis Sweet ger. t. 126. Flores albi. Petala ovata. (v. s. sine fl.)

SECT. VIII. CAMPYLIA. Sweet ger. n. 43. et p. 8.

* Petala 5 inæqualia, 2 superiora majora in ungue subauriculata. Stam. filamenta 10 pilosa aut pubescentia, 5 fertilia erecta, 5 alterna sterilia quorum 2 superiora longiora recurvo-hamata. — Herbæ vix basi suffrutescentes ramosæ. Folia petiolata ovata aut oblonga dentata aut subincisa.

* Petalis ad unguem appendiculatis, staminibus 5 fertilibus erectis, 5 sterilibus quorum 2 sup. recurvo-hamatis. — Campylia. Lindl.

77. P. BLATTARIUM (Jacq. schœnbr. 2. p. 3. t. 131.) caule suffruticoso erecto, foliis ovato-rotundis obtusis incanis holosericeis dentatis, pedunculis 4-8-floris, petalis superioribus subrotundis, inferioribus oblongis. 2 5 Campylia blattaria Sweet ger. t. 88. Petala pallidè violacea. Stam. pubescentia.

78. P. DICHONDRAEFOLIUM, caule suffruticoso erecto, foliis reniformibus crenatis incanis, pedunculis 5-floris, petalis sup. obovatis, inf. oblongis. 2 5 Burch! cat. afr. austr. n. 3084. Valdè simile P. blattario et trichostemo, sed folia reniformia. (v. s.)

79. P. SCAPOSUM, subcaule ad collum squammoso-imbricatum, foliis ovatis obtusis dentatis utrinquè incano-tomentosis, scapis subradicalibus 1-floris. 2 5 Ger. glaucum Cav. diss. 4. n. 336. t. 103. f. 2. non Andr. P. tomentosum L'her! ger. ined. n. 80. non Jacq. (v. s.)

80. P. ERIOSTEMON (Jacq. schœnbr. 2. p. 4. t. 132.) caule suffruticoso erecto, foliis elliptico-subrotundis obtusis crenatis holosericeis, pedunculis 4-floris, petalis sup. obovatis acutè emarginatis. 2 5 Ger. ovatum Cav 4. t. 103. f. 3 P Ger. ovale Burm! cap. 19. Petala alba. Stam. pilosa. (v. s. sine fl.)

81. P. TRICHOSTEMON (Jacq. ic. rar. 3. t. 524.) caule suffruticoso adscendente, foliis elliptico-subrotundis crenatis holosericeis, pedunculis 3-floris, petalis sup. subrotundis, inf. oblongis. 2 5 Petala pallidè violacea ferè P. blattarii. Stam. pilosa.

82. P. VERBASCIFLORUM, caule suffruticoso adscendente, foliis subrotundo-ovatis obtusè duplicato-dentatis margine undulatis recurvis, stipulis acuminatis, umbellis 3-5-floris, petalis superioribus subrotundis, tubo nectarifero calyce triplò breviori. 2 5 Petala pallidè lilacina. Ger. verbasciflorum Andr. ger. ic. Campylia verbasciflora Sweet ger. 2. t. 157.

83. P. HOLOSERICEUM (Sweet ger. t. 75. sub Campylia.) caule suffruticoso erecto, foliis subrotundo-ovatis obtusis duplicato-dentatis holosericeis, pedunculis 5-6-floris, petalis sup. orbiculatis (atro-purpureis), inf. oblongis (pallidè lilacinis). 2 5 Stam. pubescentia.

84. P. OENOTHERÆ (Jacq. ic. rar. 3. t. 525.) caule herbaceo adscendente, foliis oblongo-lanceolatis obtusis dentatis incano-tomentosis, pedunculis 1-3-floris, petalis sup. obovatis vix inferioribus majoribus. 2 5 Flores roseo-violacei pentandri.

85. P. CORONOPIFOLIUM (Jacq. ic. rar. 3. t. 526.) caule suffruticoso adscendente, foliis lineari-lanceolatis apice inciso-dentatis subtus canescentibus, pedunculis 3-floris, petalis sup. obovato-oblongis latè in ungue appendiculatis. 2 5 Cor. pallidè violacæ unicolores.

86. P. CAPILLARE (Willd. sp. 3. p. 660.) caule suffruticoso brevi, foliis lanceolatis profundè pinnatifidis pubescentibus, pedunculis subbifloris. 2 5 Ger. capillare Cav. diss. 4. p. 258. t. 97. f. 1. Pel. mahernium L'her. ger. ined. n. 45. Simile coronopifolio ex Willd. Stam. ignota. (v. s.)

87. P. CANUM (Pers. euch. 2. p. 229.) caule suffruticoso, foliis ovatis plicatis serratis tomentosis, pedunculis 3-floris, petalis 2 sup. latissimis ovatis. 2 5 Campylia cana Sweet ger. t. 114. Ger. tomentosum Andr. rep. t. 115. non

aliorum. Filam. 2 sup. revoluta ciliata. Petala rosea. Tubus rectus calyce dimidiò brevior.

88. *P. CARINATUM* (Sweet ger. t. 21.) caule suffruticoso adscendente, foliis ovatis inæqualiter dentatis incisive, stipulis carinatis, pedunculis 2-4-floris, petalis sup. ovalibus undulatis subemarginatis. ⑤ Ger. tricolor ovalifolium Andr. ger. ic. Ger. ovatum Cav. diss. 4. t. 103. f. 3? Pet. 2 sup. rubro-purpurea, 3 inf. alba ut in sequentibus.

* * *Petalis super. suprâ unguem verrucosis, stam. tubo brevissimo, 5 fertilibus recurvo-patentibus, 5 sterilibus rectis.* — *Phymatanthus. Lindl.*

89. *P. ELATUM*, caule fruticoso erecto, foliis lanceolatis canescentibus incisodentatis, pedunculis subtrifloris, petalis superioribus majoribus basi sublævibus, inferioribus ovali-oblongis. ⑤ *Phymatanthus elatus* Sweet ger. t. 96. *Geranium tricolor arboreum* Andr. ger. ic. Flores ferè prioris. (v. v.)

90. *P. TRICOLOR* (Curt. bot. mag. t. 240.) caule suffruticoso erecto, foliis lanceolatis villosa-canescens inciso-dentatis subtrifidis, pedunculis subtrifloris, petalis super. subrotundis brevioribus basi bullato-verrucosis, inf. ovalibus. ④ ⑤ *Pel. violarium* Jacq. ic. rar. 3. t. 527. *Ger. tricolor* Andr. ger. ic. *Phymatanthus tricolor* Sweet ger. t. 43. Pet. 2 super. rubro-sanguinea ad unguem nigricantia, 3 inf. alba. (v. v.)

Sect. IX. MYRRHIDIUM.

Petala 4 aut rarissimè 5, 2 superiora maxima obovato-cuneata, sæpiùs lineis ramosis striata, 2-3 inf. multò minora oblongo-linearia. Stam. 10 tubo filamentisque rectis, sæpiùs 5 antherifera et 5 alterna sterilia, rariùs 7 antherifera. — Herbæ biennes aut perennes rariùs suffrutescentes, caulibus teretibus, foliis pinnatim rariùs ternatim sectis sæpè multifidis.

* 5-anthera, 4-petala.

91. *P. CANARIENSE* (Willd. hort. ber. t. 17.) caule suffruticoso, foliis 3-partitis, lobis apice dentatis obtusis, inf. obovatis, medio ovato sæpè trifido, pedunculis subbifloris. ⑤ in ins. Canariensibus. Petala alba, 2 sup. rubro-striata. (v. v.)

92. *P. BULLATUM* (Jacq. ic. rar. 3. t. 530. coll. 5. p. 124.) caule herbaceo bienni procumbente puberulo, foliis pinnati-partitis, lobis inciso-lobatis utrinquè subhispidulis, infer. majoribus, pedunculis bifloris. ② *P. myrrhifolium* ⑧ Willd. sp. 3. p. 661. Petala alba, 2 sup. rubro lineata. An sequentis var?

93. *P. MYRRHIFOLIUM* (Ait. h. kev. ed. 1. v. 2. p. 421.) caule herbaceo strigosiusculo adscendente, foliis utrinquè subhispidulis rigidulis pinnati-partitis, lobis inciso-serratis, infer. majoribus, pedunculis 2-3-floris ④? *Ger. myrrhifolium* Linn. sp. 949. *Pel. betonicum* Jacq. ic. rar. 3. t. 531. coll. 5. p. 127. Petala prioris, nunc 4 rariùs 5.

94. *P. CORIANDRIFOLIUM* (Jacq. ic. rar. 3. t. 528. coll. 5. p. 142.) caule herbaceo bienni subpuberulo, foliis bipinnatisectis glabris, lobis linearibus subpinnatifidis, pedunculis subtrifloris. ② *Ger. coriandrifolium* Cav. diss. 4. t. 116. f. 1. *Jenkinsonia coriandrifolia* Sweet ger. t. 34. ex ind. syst. — Petala alba, 2 sup. rubro-striata.

** 5-anthera, 5-petala.

95. *P. LACERUM* (Jacq. ic. rar. 3. t. 532. coll. 5. p. 122.) caule herbaceo hirsuto suberecto, foliis bipinnatifidis, laciniis lanceolatis obtusis apice dentatis, pedunculis 3-5-floris. ② Petala albo-rosea venis sanguineis. Pedicelli brevès ad basin usquè cuniculati.

*** 7-anthera, 4-petala.

96. *P. LONGICAULE* (Jacq. ic. rar. 3. t. 533. coll. 5. p. 125.) caule herbaceo bienni hirsuto, foliis pinnatipartitis, laciniis apice dentatis, infer. profundioribus, pedunculis 1-5-floris. ② Petala 4-5, albo-rosea venis rubris.

97. *P. ANEMONIFOLIUM* (Jacq. ic. rar. 3. t. 535. coll. 5. p. 133.) caule herbaceo

bienni hirsuto erecto, foliis pinnati-partitis subtus hirsutulis suprà glabris, lobis dentatis, pedunculis sub-5-floris. ② Petala rosea venis rubris.

98. *P. CAUCALIFOLIUM* (Jacq. ic. rar. 3. t. 529. coll. 5. p. 145.) caule herbaceo bienni hirsuto, foliis bipinnatisectis, lobis linearibus glabriusculis, pedunculis 1-floris. ② Petala alba vel carnea, venis rubris.

99. *P. MULTICAULE* (Jacq. ic. rar. 3. t. 534, coll. 5. p. 126.) caule herbaceo procumbente bienni glabro, foliis subbipinnatifidis dentatis, pedunculis multifloris capitatis. ② Petala pallidè violacea, 2 sup. venosa et maculata.

Sect. x. JENKINSONIA. Sweet ger. n. 79.

Petala 5, 2 sup. cæteris multò majora apice emarginata lineis coloratis striata, 3 inf. multò minor. Stamina 10 ascendentia apice patentia basi pilosa, 7-antherifera quorum 3 superiora breviora, 3 sterilia abbreviata subulata æquilonga. — Caules fruticosi. Flores ochroleuci majusculi.

100. *P. QUINATUM* (Curt. bot. mag. t. 547.) caule fruticoso flexuoso, foliis pubescentibus palmatim 5-fidis, lobis cuneatis apice 3-dentatis, pedunculis 1-2-floris, tubo nectarifero calyce maximo duplò longiore, petalis sup. apice emarginatis. 5 Ger. præmorsum Andr. t. 150. *Jenkinsonia quinata* Sweet ger. t. 79. Petala albida, 2 sup. venis atro-purpureis.

Sect. xi. CHORISMA. Lindl. in Sweet ger. n. 79.

Petala 4 rariùs 5, 2 superiora longè unguiculata majora, 2 infer. multò minora. Stam. in tubum longissimum declinatum medio geniculatum connata, 7 fertilia quorum 2 infer. libera, 3 sterilia abbreviata subulata æquilonga.

101. *P. TETRAGONUM* (L'her! ger. t. 23.) ramis tetragonis carnosis, foliis cordatis obtusè lobatis subdentatis. 5 Jacq. ic. rar. 1. t. 132. Roth abb. p. 51. t. 11 et 12. benè. Curt. bot. mag. t. 136. DC. pl. grass. t. 96. *Jenkinsonia tetragona* Sweet ger. t. 99. (v. v.)

β. *trigonum* (Scop. del. 1. p. 12. t. 5. sub Ger.) ramis trigonis. 5 Sæpè in iisdem individuis rami tri-et-tetragoni.

Sect. xii. PELARGIUM. — Pelargonium Lindl. in Sweet ger. n. 41.

Pet 5 inæqualia, 2 sup. approximata. Stam. 10 inæqualia, 7 antherifera, 3 sterilia subulata.

Series I. *CICONIA*, petalis unicoloribus, 2 sup. brevioribus et angustioribus, staminibus brevibus erectis, 2 infimis brevissimis antheris ferè sessilibus, caule fruticoso-carnoso. — *Ciconium* Sweet ger. 1. n. 13. in adn. et p. 9.

102. *P. ACETOSUM* (Ait h. kew. 2. p. 430.) foliis glaberrimis obovatis crenatis subcarnosis, pedunculis paucifloris, petalis linearibus. 5 Curt. bot. mag. t. 103. Ger. acetosum Lin. sp. 947. Cav. diss. 4. t. 104. f. 3. Petala ex roseo alba elongata. Stam. sæpius (an semper) 5 fertilia. (v. v.)

103. *P. SCANDENS* (Ehrh. Beitr. 7. p. 161. Willd. sp. 3. p. 666.) foliis subrotundis obsolete lobatis crenatis glabris zonatis, pedunculis multifloris, petalis linearibus latitudine laciniarum calycis. 5 Caulis non scandens sed rami flexuosi implexi. Ger. acetosum roseum Dum. Cours. bot. cult. 5. p. 16?

104. *P. PUMILUM* (Willd. enum. 704.) foliis subrotundis obsolete lobatis crenatis, junioribus subzonatis, pedunculis sub-4-floris, petalis linearibus. 5 Valde accedit ad *P. scandentem* sed pedale aut tantum sesquipedale.

105. *P. STENOPETALUM* (Ehrh. Beitr. 7. p. 161. Willd. sp. 3. p. 666) foliis subrotundis obsolete lobatis crenatis pubescentibus zonatis, pedunculis multifloris, petalis linearibus lobis calycinis angustioribus. 5 Petala saturatè coccinea *P. ribifolium* Dum. Cours. bot. cult. ed. 2. n. 58?

106. *P. HYBRIDUM* (Ait. h. kew. 2. p. 424.) foliis subrotundis obsolete lobatis crenatis glabris immaculatis, pedunculis multifloris, petalis lineari-cuneiformibus.

mibus laciniis calycinis latioribus. ♂ *G. hybridum* Lin. mant. 97. Cav. diss. 4. t. 105. f. 2. *P. coccineum* Ehrh. Beitr. 7. p. 162. *G. miniatum* Andr. ger. ic. *P. hybridum* Sweet ger. t. t. 63. Species propria nec hybrida. Petala coccinea. (v. v.)

♂. *roseum* (Ehrh. Beitr. 7. p. 161.) foliis cordato-reniformibus, petalis roseis. (v. s.)

107. *P. ZONALE* (Willd. sp. 3. p. 667.) foliis cordato-orbiculatis obsolete lobatis dentatis supernè zonatis, pedunculis multifloris, petalis cuneatis. ♂ *Ger. zonale* Lin. sp. 947. Cav. diss. 4. t. 98. f. 2. Flores variant rubicundi, sanguinei, coccinei, rosei, pallidi, albi. Folia sæpè albo et flavido variegata. (v. v.)

♀ *marginatum* (Cav. diss. 4. p. 230.) foliis crispo-rugosis non zonatis albo-marginatis, ramis rigidis erectis. ♂ Species propria et seminibus constans ex Dum. Cours. bot. cult. 5. p. 17. Bess. cat. h. crem. 1816. p. 100. Flores coccinei aut purpureo-rosei. (v. v. in hort. genev. Dufour.)

108. *P. INQUINANS* (Ait. h. kew. 2. p. 424.) foliis orbiculato-reniformibus subdivisus crenatis tomentoso-viscidis, pedunculis multifloris, petalis obovato-cuneatis. ♂ in ins. Sanctæ-Helenæ (L'her. mss.) et ad Cap. B.-Spei (Lin.) — Dill. h. elth. t. 125. f. 151. *Ger. inquinans* Lin. sp. 945. Cav. diss. 4. t. 106. f. 2. Folia tacta inquinant colore ferrugineo Petala chermesina et in var. carnea. (v. s.)

109. *P. RETINERVUM*, foliis cordato-reniformibus obsolete 5-lobis crenatis zonatis reticulato-venosis, stipulis cordatis, umbellis multifloris, petalis cuneatis, 3 infer. duplò majoribus. ♀ Flores aurantio-coccinei. Ciconium reticulatum Sweet ger. 2. t. 143.

110. *P. CERINUM*, foliis retundato-reniformibus obsolete crenulatis nervosis pubescentibus mollissimis, stipulis cordatis subdentatis, caule erecto, umbellis multifloris, petalis obovatis superioribus minoribus, tubo nectarifero calyce duplò longiore. ♂ Petala lætè purpureo-rubra vix basi lineata. Ciconium cerinum Sweet ger. t. 176. Stam. 10 breviter erecta, interdum omnia fertilia.

111. *P. HETEROGAMUM* (L'her! ger. t. 18.) foliis cordato-orbiculatis inciso-lobatis dentatis utrinquè pubescentibus, pedunculis multifloris, petalis oblongo-cuneatis. ♂ *Ger. quinquelobum* Lam. dict. 2. p. 671. Hybridum dicitur ex *P. zonali* et *alchimilloide*. Flores rosei aut coccinei. Stam. 6 fertilia. (v. s.)

112. *P. MONSTRUM* (Ait! h. kew. 2. p. 425.) foliis orbiculato-reniformibus obsolete lobatis subzonatis complicatis crispis utrinquè pubescentibus, pedunculis multifloris, floribus adgregatis, petalis lineari-cuneatis. ♂ Sweet ger. t. 13. Flores intensè rosei. Stam. 7 fertilia, 2 brevissima. (v. v. sine fl.)

Series II. *ISOPETALOÏDEA*, petalis nempè magnitudine subæqualibus.

* *Alchimilloïdea*, caulibus herbaceis, foliis cordatis palmati-lobis, petalis parvis.

113. *P. INODORUM* (Willd. h. ber. t. 34.) caulibus herbaceis diffusis, foliis cordato-ovatis obsolete lobatis obtusè dentatis ciliatis, umbellis multifloris capitatis, petalis calyci et inter se subæqualibus. ♀ in Novâ-Hollandiâ. Sweet ger. t. 56. Lag. nov. sp. n. 272. Pet. pallidè purpurascens, 2 sup. intensius maculata. Stam. fertilia ut in species Novæ-Holl. subexpansa ex Sweet. (v. s.)

114. *P. GLOMERATUM* (Jacq. f. ecl. 1. p. 146. t. 98.) caule suffruticoso diffuso, foliis cordatis sublobatis obtusè crenatis subtus villosis, umbellâ multiflorâ confertâ, petalis calyce majoribus inter se subæqualibus, tubo nectarifero brevissimo obsolete. ♀ in Novâ-Holl. Petala alba, 2 sup. rubro maculata. *Ger. glomeratum* Andr. ger. ic. *P. australe* ♂ Willd. enum. 707. *P. australe* Sweet ger. t. 68. non Willd. Jacq.

115. *P. ODORATISSIMUM* (Ait. h. kew. ed. 1. v. 2. p. 419.) caule carnosio brevissimo, ramis herbaceis longis diffusis, foliis subrotundo-cordatis mollissimis, umbellis sub-5-floris. ♀ — Dill. h. elth. t. 131. f. 138. *Ger. odoratissimum* Cav. diss. 4. t. 103. f. 1. Petala parva dilutè rosea subæqualia. (v. v.)

♂? *Africanum* (Cav. diss. 4. p. 242. t. 104. f. 1.) foliis superioribus subtrilobis.

116. *P. FRAGRANS* (Willd. h. ber. 1. t. 77.) caule basi suffruticoso, ramis divaricato-reflexis mollissimè pubescentibus, foliis subrotundo-cordatis subtrilobis obtusè dentatis mollissimis, pedunculis multifloris. ♀ *P. fragrans* Sweet ger. t. 172. Ger. odoratissimum erectum Andr. ger. ic. Petala alba, 2 sup. lineis ramosis sanguineis picta, calyce duplò longiora. (v. v.)
117. *P. GROSSULARIOIDES* (Ait. h. kew. 2. p. 42.) caulibus prostratis tetragonis glaberrimis, foliis cordatis subrotundis incisè dentatis, pedunculis subbifloris. ♀ *Herm. lugdb. t. 289. Ger. grossularioides* Lin. sp. 948. Cav. diss. 4. t. 119. f. 2. Flores minuti pallidè incanati. (v. v.)
118. *P. ANGEPS* (Ait. h. kew. 2. p. 40.) caulibus decumbentibus triquetro-ancipitibus glabris, foliis cordato-subrotundis obsolete lobatis dentatis, umbellis multifloris. ♀ *Jaqu. coll. 4. p. 184. t. 22. f. 3. Flores parvi, priore tamen majores, rubicundi.* (v. s.)
119. *P. ACUGNATICUM* (Pet. Th. fl. acugn. p. 44. t. 13.) caulibus herbaceis erectis, foliis glabris reniformibus obsolete lobatis dentato-crenatis, umbellis multifloris capitatis, petalis calycis pilosi acuminati longitudine. ♀ in ins. Tristan d'Acunna. Petala cuneata rubella. — *P. clandestinum* L'her! ger. ined. n. 29. ex Novâ-Zelandiâ ortum ab hoc non differre videtur. (v. s.)
120. *P. CLAVATUM* (L'her! ger. ined. n. 32.) caulibus herbaceis erectis puberulis, foliis reniformibus subdivisis pubescentibus crenato-dentatis, umbellis multifloris, tubo nectarifero clavato lobis calycinis acuminatis duplò longiore. ♀ ad cap. Bonæ-Spei. Sonnerat. (v. s.)
121. *P. DISTANS* (L'her! ger. ined. n. 35.) caulibus herbaceis erectis tetragonis glabris, foliis remotis subrotundis 5-lobis glabris summis inciso-laceris, umbellis multifloris confertis longissimè pedunculatis. ♀ ad cap. Bonæ-Spei. Sonnerat. (v. s.)
122. *P. TABULARE* (L'her! ger. t. 9.) caule adscendente piloso-hispido, foliis reniformibus 3-5-lobis obtusis apice dentatis glabrinusculis, pedunculis elongatis 2-4-floris. ♀ *Ger. Tabulare* Lin. sp. 947. non *Burm. G. elongatum* Cav. diss. 4. p. 233. t. 101. f. 3. Petala albidâ calyce duplò ferè longiora. (v. v.)
123. *P. ALCHIMILLOIDES* (Willd. sp. 3. p. 656.) caule diffuso villosa, foliis cordatis 5-lobis palmatis villosis, pedunculis paucifloris, stigmatibus sessilibus. ♀ *Herm. lugd-b. t. 283. Ger. alchimilloides* Lin. vir. 345. Cav. diss. 4. t. 98. f. 1. (v. v.)
- 124? *P. PARVULUM*, caule herbaceo multicauli erecto, foliis pubescenti villosis cordatis 3-partitis, lobis incisè, pedunculis subtrifloris, calycibus subacutis pubescentibus. ①? ad cap. Bonæ-Spei. Thunberg. *P. nanum* L'her! ger. n. 41. non Sweet. Flores parvuli. Planta facie *Erodii maritimi*. Hæc et sequens species etiamsi annuæ et herbacæ accedunt ad *Athamanthoidea*. (v. s.)
- 125? *P. SENEÇIOIDES* (L'her. ger. t. 11.) caule herbaceo erecto, foliis bipinnatifido-lacinatis glabris, pedunculis subtrifloris, involucris calycibusque obtusis. ① Flores albi, petalis 2 sup. ad basin maculatis. Stam. ignota. Locus in genere incertus.
- ** *Athamanthoidea*, caule suffruticoso, foliis pinnatisectis, lobis multifidis.
126. *P. MINIMUM* (Willd. sp. 3. p. 664.) caule vix suffruticoso erecto, ramis glabris herbaceis, foliis pinnatisectis, segmentis inferioribus pinnatifidis, summis integris lobulisque lineari-oblongis, umbellis submultifloris. ♀ *G. minimum* Cav. diss. 4. p. 260. t. 121. f. 3. *P. capnoides* L'her! ger. ined. n. 46. (v. s.)
127. *P. ATHAMANTHOIDES* (L'her! ger. ined. n. 47.) caule vix suffruticoso erecto, ramis subherbaceis angulatis, foliis pube adpressâ cinereo-tomentosis bipinnatisectis lacinatis, lobulis lineari-subulatis acutis, umbellis compositis. ♀ ad cap. Bonæ-Spei. Thunberg. (v. s.)

128. *P. ABROTANIFOLIUM* (Jacq. h. schoenbr. 2. t. 136.) caule suffruticoso, foliis cinereo-velutinis palmatim trisectis, lobis linearibus trifidis, umbellis paucifloris, calycibus subhispidis. ζ Ger. abrotanifolium Lin. t. suppl. 304. Cav. diss. 4. p. 256. t. 117. f. 1. (v. s.)
129. *P. FRUTICOSUM* (Willd. sp. 3. p. 689.) caule fruticoso, foliis glabris bipinnatim decompositis, lobis linearibus acutis, pedunculis bifloris, tubo nectarifero calyce duplò triplòve longiore. ζ Ger. fruticosum Cav. diss. 4. p. 163. t. 122. f. 2. Petala oblongo-cuneata dilutè violacea, 2 sup. majora.
130. *P. INCISUM* (Willd. sp. 3. p. 686.) caule fruticoso, foliis trisectis atroviridibus, lobis distantibus tripartito-laciniatis, umbellis 6-7-floris, petalis linearibus flaccidis, superioribus longioribus, tubo nectarifero subsessili calyce triplò longiore. ζ P. incisum Sweet ger. t. 93. G. incisum Andr. bot. rep. 67? malè. Petala albida 2 sup. medio maculà sanguineà. (v. v. in hort. Walner.)
131. *P. CANESCENS* (Sweet ger. n. 93. in adn.) caule fruticoso, foliis trisectis canescentibus, segmentis aggregatis tripartito-laciniatis obtusis, petalis linearibus rigidis, superioribus brevioribus, tubo nectarifero pedicellato calyce quadruplò longiore. ζ G. incisum Andr. bot. rep. 67? Priori affinis ex Sweet.
132. *P. ARTEMISIFOLIUM*, caule fruticoso, foliis glabris bipinnatim decompositis, lobis linearibus acutis, pedunculis 1-2-floris, tubo nectarifero calyce quintuplò longiore. ζ Petala linearia alba, 2 sup. approximata maculà lineari rubrà notata. Colui in hort. Monsp. et recepi siccum ex Burch. cat. afr. austr. n. 2710. P. fruticoso valdè affine. (v. s.)
133. *P. RAMOSISSIMUM* (Willd. sp. 3. p. 688.) caule fruticoso ramosissimo, foliis pubescenti-hirsutis pinnatisectis, segmentis distantibus inciso-pinnatifidis, umbellis paucifloris, calycibus obtusis. ζ Burm. afr. t. 34. f. 2. G. ramosissimum Cav. diss. 4. p. 260. Petala in meo specim. pallida, angustiora quàm in icoue Burm., 2 sup. medio inflexa ferè ut in Hoareà. (v. s.)
134. *P. TENUIFOLIUM* (L'her! ger. t. 12.) caule carnosò nudo erecto, foliis hirsutis bipinnatim decompositis, lobis lineari-subulatis, umbellis multifloris. ζ Pedunculi elongati graciles. Petala obovato-oblonga purpurea, 2 sup. basi intensiora. (v. s. sine fl.)
135. *P. HIRTUM* (Jacq. ic. rar. 3. t. 536. coll. 5. p. 144.) caule carnosò squamoso decumbente, foliis hirtis bipinnatim decompositis, lacinis linearibus obtusis, umbellis multifloris. ζ Sweet ger. t. 113. Ger. hirtum Cav. diss. 4. p. 258. t. 117. f. 2. Pet. rosea, 2 sup. latiora purpureo-maculata. Var. tenuifolii ex Thunb. prod. p. 116. et L'her. mss. Differt tantum caule decumbente foliorumque lobis brevioribus. (v. s.)
136. *P. CONFUSUM*, caule fruticoso, foliis glaucis glabris decompositis, lacinis cuneatis inciso-dentatis, infimis divaricatis, involuero polyphylo calycibusque ciliatis. ζ Proximum *P. abrotanifolium* et *tenuifolium*. — *P. sanguineum* Willd. enum. suppl. 48. non Sweet. An à *P. apiifolio* diversum?
137. *P. TRIPARTITUM* (Sweet ger. t. 115.) caule fruticoso, foliis trisectis carnosis inciso-dentatis glaucescentibus, segmentis subsessilibus cuneiformibus, medio tripartito elongato, umbellis plurifloris, tubo nectarifero pedicelli longitudinem adæquante calyce triplò longiore. ζ Pet. ochroleuca angusta, 2 sup. basi maculà purpureà elongatà.
 - α. umbellis 3-floris. Pel. tripartitum Willd. spec. 3. p. 683. P. trifidum Jacq. schoenb. 2. t. 134.
 - β. umbellis 5-6-floris. P. fragile Willd. spec. 3. p. 686. G. fragile Andr. rep. t. 37.
- 138? *P. SPINOSUM* (Willd. sp. 3. p. 681.) caule fruticoso, foliis cuneiformibus trifidis dentatis, petiolis stipulisque persistentibus spinulentibus, umbellis compositis paucifloris. ζ Paters. itin. ed. gall. t. 10. Flores ferè *P. tripartiti*, super. majora cuneata basi punctata apice paulò emarginata. An hujus dissim?

*** *Gibbosa, caule suffruticoso-carnoso, foliis tri aut pinnatisectis carnosis, petalis è flavo sordidè fuscis.*

139. *P. GIBBOSUM* (Willd. sp. 3. p. 684.) caule fruticoso carnosio nodis tumidis, foliis pinnatisectis, segmentis 1-2-jugis cum impari obtusis cuneatis inciso-dentatis, terminali trifido, umbellis multifloris. ♀ Sweet ger. t. 61. Ger. gibbosum Lin. sp. 946. Cav. diss. 4. t. 109. f. 1. Petala sordidè viridi-flava vespere odorata. (v. v.)

140. *P. APIFOLIUM* (Jacq. f. ecl. 1. t. 27.) caule fruticoso carnosio crasso, foliis geniculato-pinnatisectis, segmentis cuneatis pinnatifidis, lobis laciniato-incisus, umbellis multifloris. ♀ Petala margine sordidè flava, in disco fusco-atra. Valdè *P. multiradiato* simile, sed ex cl. Jacq. antheræ 7, et tubus necrifer vix conspicuus.

**** *Tristia, subacaulia, radice fasciculato-tuberosâ, foliis decompositè laciniatis, petalis è flavo sordidè fuscis.*

141. *P. FLAVUM* (Ait. h. kew. ed. 1. v. 2. p. 418.) subcaulescens, foliis decompositè laciniatis hirsutis, laciniis linearibus, umbellis multifloris. ♀ Ger. dancifolium Marr. goett. 1780. p. 13. t. 4. Cav. diss. 4. t. 120. f. 2. Ger. flavum Linn. f. suppl. 257. Petala sordidè flava noctu olentia. (v. s.)

β. *dancoides* (Jacq. ic. rar. 3. t. 522.) foliorum lobulis sublanceolatis, floribus pallidè flavis lineis rubris maculatis.

142. *P. APPENDICULATUM* (Willd. sp. 3. p. 651.) subacaule, foliis bipinnatisectis villosis, lobis linearibus, stipulis petiolo adnatis apice dilatato-ovatis, umbellis multifloris hirsutis. ♀ Ger. appendiculatum Cav. diss. 4. p. 262. t. 121. f. 2.

143. *P. FILIPENDULIFOLIUM* (Sweet ger. t. 85.) subacaule, foliis hirtellis pinnatisectis, segmentis bipinnatisectis, laciniis ovatis dentatis subacutis, umbellis multifloris subhirsutis. ♀ *P. triste* β Sims. hot. mag. t. 1641. Petala *P. tristis* cujus fortè varietas? (v. v.)

144. *P. TRISTE* (Ait. h. kew. 2. p. 418.) subacaule, foliis hirtis pinnatisectis, segmentis pinnatifidis, laciniis linearibus acutis, umbellis multifloris. ♀ Delaun. herb. am. t. 27. Ger. triste Corn. can. t. 110. Cav. diss. 4. t. 107. f. 1. Petala obovato-oblonga viridi-luteo-pallida fusco-maculata ad vesperam fragrantia. (v. v.)

145. *P. LOBATUM* (Willd. sp. 3. p. 650.) subacaule, foliis cordatis subtus tomentosis obtusè 3-5-lobatis sinuato-dentatis, scapo subdiviso, umbellis multifloris. ♀ Sweet ger. t. 51. Ger. lobatum Cav. diss. 4. p. 250. t. 114. f. 2. Petala obovato-oblonga nigricantia basi et margine flava. Stam. interdum 6, rariùs 9-10. Sweet. (v. s.)

***** *Fulgida, caule brevi aut subcarnoso, foliis sectis incisus dentatisve, petalis coccineis aut sanguineis.*

146. *P. ARDENS* (Sweet ger. t. 45.) subacaule, foliis molliter villosis cordatis ovato-oblongis 3-sectis 3-5-lobatisve, lobis obtusis sinuato-dentatis, scapo ramoso, umbellis multifloris. ♀ Ger. ardens Andr. ger. ic. Pel. ardens Lodd. bot. cab. t. 139. Petala obovata intensè purpuro-iguea, tubus neet. longissimus. Stam. sæpè 6. Hybridum dicitur ex lobato ♀ et fulgido ♂. (v. v. in hort. Walner.)

147. *P. SANGUINEUM* (Wendl. coll. 2. t. 53. Sweet ger. t. 76. non Willd.) caule carnosio-crasso, foliis pilosiusculis pinnatisectis, segmentis laciniato-pinnatifidis decurrentibus, lobis lineari-lanceolatis, umbellis multifloris, petalis obovatis suboblongis. ♀ Flores intensè purpuro-coccinei longe pedicellati. (v. v.)

148. *P. PATENS* (Sweet ger. t. 125.) caule carnosio suffruticoso erecto, foliis pilosulis trisectis, segmentis sessilibus inciso-dentatis obtusis, umbellis multifloris subsimplicibus, petalis ligulatis patentibus, calycibus reflexis

tubo nectarifero calyce duplò longiore. $\frac{1}{2}$ 5 Petala 3 inf. pallidè subrosea, 2 sup. lateritio-rosea subcoccinea. Natum ex semine P. fulgidi à sp. quâdam sect. Hoaræ ined. fecundati.

149. P. HOARÆFLORUM (Sweet ger. t. 133.) subcaulescens, scapo ramoso, foliis pinnatisectis pilosis canescentibus, segmentis alternis oppositisque pinnatifidis laciniatisque, lobis cuneatis inciso-dentatis, umbellis multifloris, petalis oblongo-subcuneatis, inferis concavis subconniventibus. $\frac{1}{2}$ $\frac{1}{2}$ Pet. ferè lateritia, 2 sup. nigro lineata. Anthæræ 7 effœtæ. Hybridum fortè è P. fulgido et nutante.
150. P. CRUENTUM (Sweet ger. t. 170.) subcaule, foliis pinnatifido-laciniatis pilosis, segmentis oblongo-spathulatis obtusis dentatis, umbellis compositis multifloris, petalis patentibus, infer. linearibus, tubo nectarifero calyce duplò longiore. $\frac{1}{2}$ $\frac{1}{2}$ Petala atro-anginea ferè P. melananthi sed antheræ 7. Dicitur hybridum P. fulgidi à P. melanantho fecundati.
151. P. AMOENUM (Sweet ger. t. 121.) subcaulescens, foliis pinnatisectis canescentibus, segmentis oppositis alternisque ovatis obtusis integris dentatisque, umbellâ compositâ, petalis patentibus oblongis, tubo nectarifero calyce duplò longiore. $\frac{1}{2}$ Pet. lateritia nigro lineata. Stam. 6-7 fertilia, 2 ut in Dimacriâ longiora. Hybridum è P. fulgido et P. viciefolio.
152. P. PYRETHRIFOLIUM (Sweet ger. t. 153.) ramosissimum, foliis cordatis palmato-bipinnatifidis scabris margine planis, segmentis oblongo-lanceolatis obtusis dentatis, pedunculis 2-3-floris, petalis infer. ligulatis, tubo nectarifero longitudine calycis. $\frac{1}{2}$ Petala lætè aurantio-rubra, sup. nigro lineata et maculata. Natum è semine P. fulgidi à P. radulâ fecundati.
153. P. FULGIDUM (Ait. h. kew. 2. p. 422.) caule fruticoso carnosio, foliis trisectis, segmentis sessilibus cuneatis inciso-dentatis, medio majore pinnatifido, umbellis multifloris sæpè geniculatis, pedicellis post anthesin subrefractis. $\frac{1}{2}$ G. fulgidum Lin. sp. 945. Cav. diss. 4. t. 116. f. 2. Pel. fulgidum Sweet ger. t. 69. Petala coccinea obovata. (v. v.)
154. P. POTTERI (Sweet ger. t. 147.) caule fruticoso-succulento, foliis cordatis trifidis multinerviis, segmentis lobatis inciso-dentatis, stipulis lanceolatis acutis, umbellis multifloris subpaniculatis, tubo nectarifero calyce sesquilongiore. $\frac{1}{2}$ Pet. obovata aurantio-coccinea, 2 sup. nigro-lineata et maculata. Natum è semine P. fulgidi à quodam ignoto fecundati.
155. P. SCINTILLANS (Sweet ger. t. 28.) caule fruticoso subcarnoso flexuoso adscendente, geniculis tumidis, foliis cuneatis ovatis aut basi subtruncatis inæqualiter dentatis incisive, umbellis multifloris, tubo nectarifero calycis longitudine. $\frac{1}{2}$ Petala coccinea, 2 sup. latiora lineis nigris ramosis picta. Hybridum è matre P. fulgido et patre ign.
156. P. IGNESCENS (Sweet ger. n. 2. et 55.) caule fruticoso subcarnoso, foliis cordatis trilobatis, laciniis dentatis, lateralibus bifidis, mediâ trilobâ, stipulis cordatis acuminatis subdentatis, umbellis 4-floris. $\frac{1}{2}$ Petala intensè rubro-coccinea, 2 sup. nigro maculata. Hybridum è matre P. fulgido et patre ignoto.
 - a. fertile (Sweet ger. t. 2.) petalis oblongioribus majoribus, tubo nectarifero calyce duplò longiore, antheris polliniferis. — Ger. ignescens majus Andr. ger. ic.
 - β. sterile (Sweet ger. t. 55.) petalis obovatis brevioribus, tubo nectarifero calyce vix longiore, antheris effœtis. — Ger. ignescens Andr. ger. ic. P. ignescens Lodd. bot. cab. t. 109. P. splendens Willd. h. berol. t. 76. (v. v.)
157. P. CONCOLOR (Sweet ger. t. 140.) caule suffruticoso ramoso-diffuso, foliis cordatis 5-lobis undulatis mollibus, stipulis cordatis, umbellis multifloris, tubo nectarifero calyce longiore. $\frac{1}{2}$ Petala rubro-purpurea, 2 sup. lineis nigricantibus. Hybridum dicitur è P. ignescente et P. capitato.
158. P. FLOCCOSUM (Sweet ger. t. 129.) caule fruticoso erecto, foliis cordatis

3-partitis undulatis utrinquē villosopilosis, segmentis sublobatis grossèquè obtusè dentatis, stipulis lanceolatis acutis, pedunculis paucifloris, tubo nectarifero calyce triplo breviorè. ♀ Petala coccineo-purpurea nigro-lineata, 2 sup. lineis intensioribus et maculâ nigrâ notata. Natum è semine P. ignescentis à P. Principissæ fecundati.

159. P. CHRYSANthemifolium (Sweet ger. t. 124.) caule fruticoso ramosissimo, foliis cordatis pinnatifido-sub 7-lobatis, sinubus rotundatis crenatis, stipulis oblato-cordatis subdentatis acutis, umbellis multifloris, petalis inferis linguatis, tubo nectarifero calyce reflexo longiore. ♀ Petala pulchrè rubro-purpurea, 2 sup. nigro-lineata submaculata. Natum è semine P. ignescentis à P. quercifolio fecundati.

160. P. SMITHII (Sweet ger. t. 110.) caule fruticoso ramis diffusis, foliis cordatis 5-lobis undulatis inæqualiter profundèquè dentatis rigidis, stipulis cordatis subdentatis, umbellis plurifloris, tubo nectarifero calyce parùm longiore. ♀ Pet. 3 inf. rosea, 2 sup. rubro-coccinea venis nigris lineata et medio maculata. Hybridum è P. ignescente secundato forsàn à P. Florido.

161. P. NANUM (Sweet ger. t. 102.) caule basi suffruticoso declinato, foliis cordatis trilobis inæqualiter obtusè dentatis, stipulis cordato-lanceolatis ciliatis, pedunculis sub 4-floris paniculatis. ♀ P. ignescens nanum Hoar. ex Sweet. Flores ferè prioris.

162. P. BENTINCKIANUM (hortul.) caule fruticoso-subcarnoso, foliis cordatis orbiculatis breviter et obtusissimè 5-7-lobis crenatis utrinquē mollissimè velutinis, umbellis multifloris velutinis, petalis obovatis subæqualibus, tubo nectarifero calyce quintuplo longiore. ♀ Petala intensè coccinea. Medium videtur inter fulgidum et inquinans. (v. v.)

163. P. BRACTEOSUM, caule fruticoso subcarnoso, foliis cordatis orbiculatis breviter et obtusè 5-7-lobis crenatis glabris, umbellis multifloris, petalis obovatis subæqualibus, tubo nectarifero calyce quadruplo longiore. ♀ Pedicelli et calyces atro-purpurascens. Petala intensè coccinea, striis paucis nigricantibus. Hybridum videtur è P. hybrido et fulgido. Stipulæ et bractea maximè reniformes. (v. v. in hort. Gen.)

***** Bicolora, caule suffruticoso, foliis lobatis hirsutis, petalis medio latè purpuro-maculatis.

164. P. QUINQUEVULNERUM (Willd. enum. 703.) caule suffruticoso, foliis hispidis trisectis, segmentis multifidis, lacinis lineari-lanceolatis serratis, umbellis multifloris. ♀ Delaun. herb. am. t. 28. Ger. quinquevulnerum Andr. rep. t. 114. Caulis spithameus. Petala atro-purpurea margine pallida. (v. v. in hort. Walner.)

165. P. BICOLOR (Ait. h. kew. 2. p. 425.) caule suffruticoso, foliis cordatis trifidis undulatis hirsutis obtusis dentatis, segmentis lateralibus trilobis, super. 5-lobis, stipulis reniformibus integris, umbellis confertis multifloris, calycibus reflexis. ♀ Ger. bicolor Jacq. h. vind. 3. t. 39. Cav. diss. 4. t. 111. f. 1. P. bicolor bot. mag. t. 201. Sweet ger. t. 97. Petala cuneato-oblonga distincta nunc pallida nunc sæpius atro-purpurea margine alba. (v. v.)

166. P. IMBRICATUM (Sweet ger. t. 65.) caule suffruticoso, foliis cordatis undulatis hirsutis subtrifidis, lobis obtusis dentatis, lateralibus bifidis, terminali lobato, stipulis reniformibus dentatis, umbellis confertis multifloris. ♀ G. bicolor var. magna Andr. ger. ic. Petala prioris sed duplo majora obovata imbricata. Hybridum è matre P. bicolor et forsàn patre P. obscuro.

167. P. OBSCURUM (Sweet ger. t. 89.) caule suffruticoso, foliis truncato-cordatis piloso-hirsutis concavis 5-lobis inæqualiter dentatis, sup. trilobis, stipulis reniformibus ciliatis, umbellis 4-floris, tubo nectarifero calyce parùm breviorè. ♀ P. maculatum obscurum Hoare ger. p. 21. Petala magna obovata, 2 sup. atro-purpurea margine alba, 3 inf. alba medio submaculata. Hybridum è P. maculato et quodam ignoto.

168? *P. ORNATUM* (Sweet ger. t. 39.) caule suffruticoso ramosissimo, foliis basi truncato-aut cuneato-cordatis trifidis undulatis inæqualiter dentatis hirsutis, stipulis latè cordatis, umbellis 3-4-floris, pedicellis fructiferis declinatis, calycibus revolutis. ♀ Petala alba, 2 sup. paulò latiora sanguineo mediò maculata et basi-lineata. Affine, ex cl. Sweet, *P. Hermannifolium*.

***** *Cortusina*, caule carnoso suffruticoso, foliis oblongis aut sæpius cordatis subincisis, stipulis lanceolatis patentibus acutis, radicibus tuberoso-fasciculatis. — Stam. 7, 6, rarius 5, et inde affinitas quædam cum *Otidia* et *Isopetalo*.

169? *P. PALLENS* (Sweet ger. t. 148.) caule suffruticoso carnoso decumbente, ramis floriferis divisis, foliis tripartitis pilosis, segmentis lateralibus minoribus lobato-dentatis, terminali elongato inciso-dentato, umbellis 4-5-floris, petalis patentibus inf. oblongis sup. subspathulatis, tubo nectarifero longissimo. ♀ 4 Petala ex albo ochroleuca. Ger. pallens Andr. ger. ic.

170? *P. FULCHELLUM* (Curt. bot. mag. t. 524. non Willd.) subacaule pilosiusculum, scapo diviso, foliis oblongis lobato-pinnatifidis, petiolis basi adnatis, umbellâ multiflorâ, tubo nectarifero gracili calyce triplò longiore. 4 5 Ger. pictum Andr. rep. t. 168. *P. pictum* Pers. ench. n. 18. nec n. 21. Petala alba, 2 sup. obovata medio maculâ parvâ subrotundâ, 3 inf. oblonga lineâ rubrâ picta.

171? *P. PICTUM* (Pers. ench. 2. p. 227.) subacaule, foliis cordatis oblongis subruncinatis dentatis tomentosis, scapo ramoso, umbellâ multiflorâ, involucro folioso. 4 5 Ger. pictum Andr. rep. t. 160. Pel. rubro-maculatum Poir. suppl. 2. p. 757. Petala alba rubro maculata.

172. *P. ECHINATUM* (Curt. bot. mag. t. 309.) caule crasso carnoso, foliis ovato-cordatis sublobatis creuatis subtus villosis, stipulis persistentibus spinescentibus, umbellis multifloris, tubo nectarifero gracili calyce duplò longiore. ♀ *P. hamatum* Jacq. h. schœnbr. 2. t. 138. *P. echinatum* Sweet ger. t. 154. Petala oblonga subcuneata alba, 2 sup. sanguineo-maculata. Antheræ 6. (v. s.)

173. *P. CRASSICAULE* (L'her! ger. t. 26.) caule carnoso ramoso lævi, foliis reniformibus obacuminatis dentatis utrinquè sericeis, umbellis multifloris, bracteis pedicello quadruplò brevioribus, petalis vix emarginatis. 5 5 Antheræ 5. Petala alba sup. imâ basi rubro-striata Tubus nectarifer, ex L'her. ic. et mss., nullus! An tenuissimus aut brevissimus? (v. s. sine fl.)

174. *P. PRIMULINUM* (Sweet ger. n. 14. in adn.) caule carnoso ramoso lævi, foliis reniformibus obacuminatis dentatis utrinquè sericeis, umbellis multifloris, bracteis dimidiò pedicellis brevioribus, petalis obcordatis. 5 Pel. crassicaule bot. mag. t. 477. Antheræ 7. Petala alba omnia medio sanguineo maculata. Flores odore *Primulæ veris*.

175. *P. CORTUSÆFOLIUM* (L'her. ger. t. 25.) caule crasso carnoso, foliis cordatis inciso-lobatis undulatis obtusè dentatis pubescentibus, umbellis multifloris, tubo nectarifero calyce quadruplò longiore. 5 Sweet ger. t. 14. Ger. cortusæfolium Andr. rep. t. 121. Petala obovata subobcordata ex albo subrosea, 2 sup. basi lineis striata, 3 inf. medio submaculata. Antheræ 6 ex L'her. mss. et Sweet.

176. *P. PARTICEPS* (Sweet ger. t. 49.) caule carnoso, foliis cordato-reniformibus leviter 5-lobis undulatis inæqualiter obtusè dentatis subtus tomentosis, stipulis persistentibus lanceolato-subulatis, umbellis multifloris, tubo nectarifero calyce triplò longiore. ♀ *P. cortusæfolium* var. *cortusoides* Andr. ger. ic. Petala intensè rosea subpurpurea. Hybridum è *P. reniforme* et *cortusæfolio*. Antheræ 6.

177. *P. SÆPEFLORENS* (Sweet ger. t. 58.) caule crasso flexuoso, foliis cordatis leviter 5-lobis crenatis planis subtus tomentosis, stipulis subulatis persistentibus, umbellis multifloris, tubo nectarifero subsessili calyce quadruplò longiore. ♀ *G. echinatum* semperflorens. Andr. ger. ic. Antheræ 6. Petala

obovato-oblonga retusa, pallidè purpurea medio submaculata. Hybridum è *P. reniformi* ♀ et verosim. echinato ♂.

178. *P. RENIFORME* (Curt. bot. mag. t. 493.) caule fruticoso flexuoso subcarnoso, foliis reniformibus crenato-dentatis subtus tomentosius, umbellâ 3-6-florâ, stipulis persistentibus basi dilatatis, tubo nectarifero subsessili calyce triplò longiore. § Sweet ger. t. 48. Ger. reniforme Andr. rep. t. 108. Petala purpurea medio submaculata obovata retusa. Antherae 6.

***** *Pinguifolia*, caule fruticoso carnosio, foliis peltatis cordatisve 5-lobis carnosius, tubo nectarifero longitudine pedicelli, stipulis lato-ovatis.

179. *P. PINGUIFOLIUM* (Sweet ger. t. 52.) caule fruticoso, ramis carnosius diffusis subangulatis, foliis reniformi-cordatis leviter 5-lobis subcrenatis carnosius puberulis, lobis obtusius muticis, umbellis multifloris. § G. hederinum fl. albo Andr. ger. ic. Petala obovato-oblonga pallidè rosea, 2 sup. purpureo-maculata. Forsan hybridum è *P. peltato* aut lateripede ♀ et reniforme ♂.

180. *P. LATERIPES* (L'her! ger. t. 24.) caule fruticoso, ramis carnosius teretibus, foliis cordatis 5-lobis subdentatis carnosius glabris, umbellis multifloris. § Petala dilutè purpurascens, 2 sup. lineis intensioribus, oblonga-subovata. (v. s.)

ℓ. *Walneri*, foliis obtusissimè lobatis, umbellis paucifloris. § Ferè medium inter lateripedem et peltatum. (v. v.)

γ. *albomarginatum*, foliis subcrispatis albo-marginatis. (v. v.)

181. *P. PELTATUM* (Ait. h. kew. 2. p. 427.) caule fruticoso, ramis carnosius angulatis, foliis peltatis 5-lobis integerrimis carnosius, umbellis paucifloris. § Ger. peltatum Lin. sp. 947. Cav. diss. 4. p. 232. t. 100. f. 1. *P. peltatum* Curt. bot. mag. t. 20. (v. v.)

β. *zonatum*, foliis zonâ fuscâ notatis. Haw. pl. succ. p. 307.

182. *P. SCUTATUM* (Sweet ger. t. 95.) caule fruticoso, ramis angulatis petiolisque puberulis, foliis carnosius subpeltatis 5-lobis glanduloso-crenatis pubescentibus, umbellis 7-8-floris, petalis patentissimis, filam. 7 antheriferis, 2 brevissimis, tubo nectarifero subsessili calyce triplò longiore. § Flores albi, pet. 2 sup. lineis 2 ramosis purpureis. Medium ferè inter *P. peltatum* et elegans, sed ex Sweet videtur species genuina.

Series III. *PLATYPETALA*, petalis 2 superioribus latioribus brevioribus obtusissimis.

183. *P. OVALE* (L'her! ger. t. 28. non Burm.) caule suffruticoso tortuoso debili prostrato, ramis petiolisque pedunculisque molliter hispidis, foliis ovalibus acutius dentatis canescentibus, umbellis sub-5-floris longè pedunculatis tubo nectarifero calyce brevior. § Petala purpurascens oblongo-obovata subæqualia, 2 sup. breviora latiora lineis intensius purpureis. Stam. 7 fertilia. (v. s.)

184. *P. ELEGANS* (Willd. sp. 3. p. 655.) caule suffruticoso erecto, foliis elliptico-subrotundis argutè serratis obtusius rigidis glabris, umbellis sub-5-floris, petalis omnibus obovatis, tubo nectarifero calyce brevior. § Ger. elegans Andr. rep. t. 28. *P. elegans* Sweet ger. t. 36. Petala alba 2 sup. basi sanguinea lineis intensioribus notata. Stam. 7 fertilia, filam. puberulis.

Series IV. *ANISOPETALA*, petalis 2 superioribus longioribus et latioribus, caulibus fruticosis.

§. 1. Glaucescens, foliis nempè glabris aut subglabris plus minùs glaucis.

* *Petalis albis*, sup. sæpiùs rubro lineatis, maculatisve.

185. *P. GLAUCUM* (L'her! ger. t. 29.) glaberrimum glaucum, foliis lanceolatis integerrimis acuminatis, pedunculis 1-2-floris, tubo nectarifero calyce quintuplò longiore. § Sweet ger. t. 57. Ger. glaucum Andr. ger. ic. Ger. lan-

- ceolatum Cav. diss. 4. t. 102. f. 2. bot. mag. t. 56. non Andr. Pet. alba, 2 sup. subrotunda basi sanguinea. (v. v.)
186. *P. DIVERSIFOLIUM* (Wendl. obs. 51.) glabrum glaucum, foliis lanceolatis integerrimis trisectisve, inferioribus dentatis, pedunculis sub-1-floris paniculatis. ♀ Flores prioris sed minores. Folia apice unbarbata.
187. *P. OXYPHYLLUM*, glaberrimum glaucum, foliis 3-foliolatis, foliolis integerrimis linearibus utrinque acuminatis, lateralibus brevioribus, pedicellis 1-floris, stipulis lineari-subulatis. ♂ *P. lævigatum* L'her! herb. et ger. ined. n. 91. non Willd. (v. s.)
188. *P. CUSPIDATUM* (Willd. enum. suppl. 47.) glaberrimum glaucescens, foliis ovatis acutis glaucis subinciso remotè serratis, pedunculis sub-3-floris. ♂ Affine *P. glauco* ex Willd.
189. *P. SORORIUM* (Willd. enum. suppl. 48.) glaberrimum, foliis profundè 3-partitis, lacinii acinaciformibus inciso-serratis, pedunculis 3-floris. ♂ Hybridum è *P. grandifloro* ♀ et *P. glauco* ♂, juxta Willd. Petala alba leviter incarnata, 2 sup. latiora lineis sanguineis ramosis duabus picta.
190. *P. LÆVIGATUM* (Willd. sp. 3. p. 685.) glaberrimum glaucum, foliis 3-sectis, segmentis 3-fidis cuneatis, lacinii lineari-lanceolatis, pedunculis subbifloris. ♂ *Ger. lævigatum* Lin. f. suppl. 306. Cav. diss. 4. t. 121. f. 1.
℘. compositum (L'her! ger. ined. n. 92.) foliis trisectis, segm. lateralibus indivisis, medio 3-partito subpinnatifido, summis 3-fidis. Tubus nectarifer longissimus ut in icone Cavan, et in specie sequente. (v. s.)
191. *P. GRANDIFLORUM* (Willd. sp. 3. p. 674.) glabrum glaucum, foliis 5-lobo-palmatifidis basi cordatis, lobis versus apicem dentatis, pedunculis 3-floris, tubo nectarifero calyce quadruplò longiore, petalis calyce triplò longioribus. ♂ *Ger. grandiflorum* Andr. rep. t. 12. non Lin. calyces et pedicelli subpioliusculi. Petala alba, 2 sup. cuneato-obovata basi sanguineo striata. (v. v.)
℘. roseum, flore roseo. — In hortis interdum occurrit sub nomine *P. nobilis*.
192. *P. VARIEGATUM* (Willd. sp. 3. p. 674.) glabrum glaucum, foliis 3-5-lobo-palmati-partitis, lacinii trifidis dentatis, stipulis ovato-cordatis acutis, pedunculis subbifloris, tubo nectarifero calyce quintuplò longiore, petalis calyce triplò longioribus. ♂ *Ger. variegatum* Lin. suppl. 305. Cav. diss. 4. p. 251. t. 118. f. 3. Petala 2 sup. lineis sanguineis ramosis picta. (v. s. sine fl.)
193. *P. ALBIFLORUM* (Spin. cat. 1818. p. 30?) subglabrum glaucescens, foliis 3-lobis, lobis obovatis apice acutè dentatis, petiolis pedicellis calycibusque puberulis, pedunculis bifloris, tubo nectarifero petalisque superioribus calyce subtriplo longioribus. ♂ *P. patulum* Spin. cat. h. S. Seb. Pet. alba, 2 sup. lineis sanguineis striata, basi valdè angustata. Priori affine sed flores dimidiò minores. Differt à *P. patulo* floribus albis nec roseis, tubo vix calycem excedente nec triplò longiore. (v. s.)
194. *P. CURTISIANUM* (Balb! et Spin! cat. hort.) puberulum glaucescens, foliis basi cuneatis trifidis, lobis dentatis, pedunculis bifloris, tubo nectarifero calyci æquali, petalis calyce duplò longioribus. ♂ Petala alba 2 sup. sanguineo-striata. (v. s.)
195. *P. COLLÆ* (Spin! cat. hort. 1818. p. 31.) puberulum glaucescens, foliis basi cuneatis trifidis, lobis acutè dentatis, pedunculis 4-5-floris ramiisque hispidis, tubo nectarifero calyce dimidiò brevior, petalis calyce duplò longioribus. ♂ Petala alba, 2 sup. lineis sanguineis ramosis picta. (v. s.)
196. *P. BELLARDII* (Spin! cat. hort. 1818. p. 30.) hispidulum glaucescens, foliis palmatis 5-7-fidis, lobis acutè dentatis, pedunculis 4-10-floris, tubo nectarifero calyce dimidiò brevior, petalis angustis calyce duplò vix longioribus. ♂ Petala alba immaculata. (v. s.)

** *Petalis roseis violaceisve, superioribus purpureo sæpè striatis.* (1)

197. *P. PATULUM* (Jacq. ic. rar. 3. t. 541. coll. 4. p. 187.) glabrum glaucum, foliis longissimè petiolatis cordato-reniformibus 3-5 fidis dentatis, pedunculis bifloris, tubo nectarifero calyce triplò longiore, petalis lanceolato-cuneatis, 2 sup. submarginatis. ♂ Stam. 6 fertilia. Petala rosea, 2 sup. circulo sanguineo picta.
198. *P. SANICULÆFOLIUM* (Willd. sp. 3. p. 673.) glabrum glaucum, foliis longè petiolatis cordato-subrotundis 5-fidis dentatis supernè zonatis, pedunculis 4-5-floris tubo nectarifero calycis longitudine, petalis calyce duplò longioribus. ♂ *P. cortusæfolium* Jacq. ic. rar. 3. t. 539. Ger. Tabulare Burm. ger. t. 1. Cav. diss. 4. t. 100. f. 2. *Geranium hederæfolium* Dum. Cours. bot. cult. 4. n. 62. *P. Hippocrepis* L'her! ger. ined. n. 94. Petala 2 sup. obovata magna violacea purpureo-striata, 3 inf. pallidè rosea ferè linearia. (v. s.)
199. *P. HEPATICIFOLIUM* (Spreng. nov. prov. p. 31.) glabrum glaucescens, foliis profundè trilobatis concavis supernè zonatis subtùs asperis, lobis cuneatis inciso-dentatis, pedunculis subbifloris. ♂ Pet. pallidè rosea, 2 sup. leviter triloba, 3 inf. linearia exalbida. Aff. priori.
200. *P. NOBILE* (hortul.) pilosiusculum subglaucescens, foliis cordatis 5-lobopalmatifidis, lobis obtusis versùs apicem grossè dentatis, pedunculis 3-4-floris, tubo nectarifero calycis longitudine, petalis calyce duplò longioribus. ♂ Petala pallidè rosea, 2 sup. purpureo-linearia. *P. nobile* Diet. nachtr. 6. p. 68. ex Link enum. h. ber. 2. p. 189? (v. v. in hort. Walner.)
℘. longècalcaratum, tubo nectarifero calyce duplò ferè longiore, flore majore. (v. v. in hort. Dufour.)
201. *P. BARNARDINUM* (Sweet ger. t. 127.) glaucescens subpilosulum, foliis 3-partitis glabriusculis, segmentis lateralibus bilobis, terminali 3-lobò dentato, umbellis 5-6-floris, tubo nectarifero calyce duplò longiore. ♂ Petala obovato-oblonga rosea nigro-linearata, 2 sup. colore intensiora. Natum è semine *P. grandiflori* à *P. fulgido* fecundati.
- 202? *P. OPULIFOLIUM* (Sweet ger. t. 53.) caule glabro, foliis pilosiusculis cordatis subcucullatis 5-lobis dentatis, stipulis cordatis acuminatis, pedunculis sub-4-floris, tubo nectarifero calyce paulò longiore. ♂ Flores ampli intensè rosei, petalis 2 sup. lineolatis. Hybridum fortè è *P. grandifloro* et quàdam specie cucullatà.
- 203? *P. FUSCATUM* (Jacq. ic. rar. 3. t. 540. coll. 4. p. 186.) subglabrum subglaucescens, foliis cordatis 5-lobis dentatis subtùs glaucis junioribus supernè zonatis, summis sub-5-partitis, pedunculis 4-floris, tubo nectarifero calycis longitudine. ♂ Pet. 2 sup. obovata lata intensè rosea purpureo-striata, 3 inf. linearia pallidè rosea.
- 204? *P. BEAUFORTIANUM* (Pers. ench. 2. p. 231.) glabrum subglaucescens, foliis acutè 5-lobis grossè et parcè serratis, supremis basi cuneatis subcucullatis, umbellis 4-floris pubescentibus, tubo nectarifero calyce duplò triplòve longiore. ♂ Sweet ger. t. 138. Ger. Beaufortianum Desf. arb. 2. p. 459. Ger. Beaufortiae Andr. ger. ic. Petala rosea 2 sup. venis atropurpureis picta. Hybridum fortè è *P. acerifolio* et variegato. (v. v. in hort. Bruce.)
- 205? *P. GHILINI* (Spin cat. 1818. p. 31.) subglaucescens glabrum, foliis subcuneatis 3-5-fidis, lobis ovatis acutè serratis rigidis, pedunculis 4-floris, tubo nectarifero calyce subæquali. ♂ Affine hinc *P. fuscato*, illinc *Beaufortiano*; ferè *P. opulifolii* varietas, sed minùs pilosum et calcar brevius. Petala roseo-purpurascencia, 2 sup. linearia. (v. v.)

(1) Series ferè media inter *P. grandiflorum* et *alchimilloideum*.

206. *P. GRIMALDIÆ* (Spin! cat. 1818. p. 31.) glaucescens glabratum, foliis basi subcuneatis trilobatis inæqualiter crenulatis cucullatis crispis, stipulis bracteisque cordato-ovatis acuminatis, umbellis 1-2-floris, tubo nectarifero calyce paulò longiore. ♂ Pet. roseo-carnea, 2 sup. majora obtusa lineis purpureis ramosis striata. A priore differt foliorum dentibus obtusis nec acutis. Hybridum videtur è sp. quâdam glaucescente et sp. quâdam crispâ.

§. 2. *Lineata, floribus albis aut vix roseis, petalis 2 sup. intensè rubro lineatis, foliis ovatis cordatis reniformibusve dentatis indivisis.*

* *Foliis ovatis.*

207. *P. PENICILLATUM* (Willd. h. berol. t. 37.) foliis ovatis inciso-serratis, junioribus retrorsum scabris, adultis subglabris, stipulis ovatis acuminatis, pedunculis 1-2-floris, tubo nectarifero calyce duplò ferè longiore. ♂ Ger. elegans Dum. Cours. bot. cult. 5. p. 382. Calyces apice pilis paucis penicillati. (v. v.)

208. *P. BOYLEÆ* (Sweet ger. t. 50.) foliis subrotundis, infer. subcuneatis, super. subrhombeis, obsolete 5-lobatis inæqualiter argutèque dentatis, stipulis lanceolatis acuminatis, umbellis 3-4-floris, tubo nectarifero calyce reflexo longiore. ♂ *P. formosissimum* Pers. ench. 2. p. 231. *P. formosum* Desf. arb. 2. p. 459. Horn. suppl. 75. (v. v.)

209. *P. BETULINUM* (Ait. h. kew. 2. p. 429.) foliis ovatis inæqualiter serratis glabriusculis, stipulis ovato-lanceolatis, pedunculis 2-4-floris, tubo nectarifero calyce brevior. ♂ Ger. betulinum Lin. sp. 946. — Burm. afr. t. 33. f. 1. 2. — bot. mag. t. 148. Petala alba aut rosea venis sanguineis striata. (v. v.)

210. *P. LINEATUM* (Sweet ger. t. 16.) foliis ovatis dentatis vix subtrilobis undulato-puberulis, stipulis ovatis parvis, pedunculis 1-2-floris, tubo nectarifero calycis longitudine. ♂ Petala albo-rosea venis sanguineis lineata.

211. *P. DUMOSUM* (Sweet ger. t. 19.) foliis ovatis obtusis dentatis subtrilobis pubescentibus, stipulis ovatis parvis, pedunculis 2-3-floris, pedicellis fructiferis geniculatis, tubo nectarifero vix calyce longiore. ♂ Pet. 3 inf. alba, 2 sup. medio et extus sanguinea, ad marginem alba substriata. (v. v. in hort. Dufour.)

** *Foliis basi cordatis.*

212. *P. OPTABILE* (Sweet ger. t. 62.) foliis cordatis 5-lobis inciso-dentatis scabris, stipulis acuminatis, umbellis 2-3-floris, calycibus reflexis longitudine tubi nectariferi. ♂ Hybridum forte è *P. penicillato* et *dumoso*, ex Sweet.

213. *P. VENUSTUM* (Sweet ger. t. 167.) foliis reniformibus sublobatis denticulatis utrinquè pilosis, stipulis cordatis ciliatis, pedunculis subpaniculatis 3-5-floris, calycibus reflexis tubi nectariferi longitudine. ♂ Pet. 2 sup. purpureo-lineata et maculata. Valdè simile priori. Hybridum dicitur *P. dumosi* à *P. Gardneriæ* (sp. ined.) fecundati.

214. *P. MELISSINUM* (Sweet ger. t. 5.) foliis cordatis 5-lobis undulato-plicatis dentatis, lobis inferioribus dilatatis lobatis, superioribus trilobis, stipulis dentatis ciliatis, umbellis 4-5-floris, calyce patente tubo nectarifero subbreuiore. ♂ Petala 2 sup. latè sanguineo-maculata, inf. alba linearia.

215. *P. HOAREANUM* (Sweet ger. t. 80.) foliis cordato-ovatis sublobatis angustatis rigidis glabris, marginibus undulatis cartilagineo-dentatis ciliatis, stipulis ovatis, pedunculis 1-2-floris, tubo nectarifero calyce duplò longiore. ♂ Petala ampla alba, 2 sup. majora sanguineo latè maculata et lineata.

216. *P. CANDIDUM* (Hoare in Sweet ger. t. 128.) foliis super. cordatis 3-lobis inæqualiter obtusè dentatis pilosis canescentibus, stipulis cordatis, umbellis plurifloris, pedunculis calycibusque villosissimis, tubo nectarifero calyce

longitudine. 5 Pet. alba. 2 sup. lineis purpureis notata. Hybrida hortensis progenies sed parentes ign.

217. *P. BAILEYANUM* (Sweet ger. t. 89.) foliis inf. reniformibus basi truncatis sublobatis inæqualiter argutè dentatis villosis, superioribus basi cuneatis sub-5-lobis, umbellis multifloris, tubo nectarifero longitudine calycis. 5 Petala ampla alba, 2 sup. obovato-cuneata intensè sanguineo maculata et lineata, 3 inf. oblonga.
218. *P. YOUNGII* (Sweet ger. t. 131.) foliis truncato-cordatis subtrilobis planis inæqualiter dentatis multinerviis utrinquè hirsutis margine undulatis, umbellis 4-floris, calycibus reflexis tubo nectarifero longitudine calycis. 5 Petala alba, 2 sup. fusco-purpureo-lineolata et latè maculata. Natum dicitur è semine *P. Waltsoni* à *P. maculato* fecundati.
219. *P. INVOLUCRATUM* (Sweet ger. t. 33.) foliis cordato-reniformibus rigidis subangulatis cartilagineo-dentatis, umbellis multifloris, bracteis magnis ovatis pedicello longioribus, calycibus reflexis, tubo nectarifero calyce breviorè 5 Flores ampli nunc albi nunc rosei, petalis sup. semper sanguineo-lineatis interdum etiam maculatis. Hybridum dicitur è *P. cucullato* et superbo.
220. *P. ROBINSONI* (Sweet ger. t. 150.) foliis cordatis acutis undulatis obsolete lobatis cartilagineo-dentatis utrinquè pubescentibus, stipulis cuneatis inæqualiter dentatis cuspidatis, umbellis multifloris, tubo nectarifero calycem subæquante. 5 Petala ampla alba subrosea, 2 super. nigro lineata cum macula purpurea. Hybridum forte è *P. involucrato* incarnato aut triumphante.
221. *P. PLATYPETALUM* (Sweet ger. t. 116.) foliis infer. cordatis obsolete lobatis inæqualiter dentatis hirsutis, super. basi cuneatis, umbellis plurifloris, petalis omnibus rotundato-obovatis, superioribus reflexis, tubo nectarifero calyce duplò breviorè. 5 Pet. 3 inf. alba, 2 sup. subrosea purpureo-atro-lineata submaculata. Affine hinc *P. eleganti*, illinc *Robinsoni* et omnibus lineatis.
222. *P. DIFFORME* (Sweet ger. t. 105.) foliis difformibus cordato-cuneatis, sublobatis cartilagineo-dentatis erosive, stipulis cordato-lanceolatis acuminatis dentatis-ciliatis, umbellis plurifloris, petalis obovatis, tubo nectarifero calyce breviorè. 5 Petala pallidè rosea, 2 sup. purpureo-latè maculata, omnia basi sublineata. Hybridum parentibus ignotis.
223. *P. MACRANTHUM* (Sweet ger. t. 83.) foliis inferioribus reniformibus obsolete 5-lobis cartilagineo-dentatis, super. profundè lobatis, umbellis multifloris, bracteis cordatis pedicello duplò brevioribus, tubo nectarifero longitudine calycis. 5 Flores 2-3-poll. diam. albi, petalis super. sanguineo-lineatis et maculatis. Affine *P. involucrato*.
224. *P. PANNIFOLIUM* (Sweet ger. t. 9.) foliis cordatis sublobatis inæqualiter dentatis villosis, umbellis 4-6-floris, bracteis lanceolatis brevissimis, tubo nectarifero calyce breviorè. 5 Affine *P. blando*. Flores pollicis diam. albi subrosei, petalis 2 sup. sanguineo-lineatis. (v. in hort. Dufour.)
225. *P. FULGURUM* (Sweet ger. t. 107.) foliis reniformibus cartilagineo-dentatis mollibus multinerviis, stipulis latis cordatis acuminatis, umbellis multifloris paniculatis, tubo nectarifero calyce multò breviorè. 5 Pet. 3 inf. roseo-alba venis rubentibus, 2 sup. roseo-purpurascencia atro-purpureo lineata et maculata. Hybridum forsàn è *P. striato* et florido.
226. *P. STRIATUM* (Sweet ger. t. 1.) foliis cordatis leviter 5-lobis dentatis pubescentibus, umbellis capitatis subsexflores, bracteis incis, tubo nectarifero pedicelli longitudine. 5 Petala obovata ferè æqualia alba venis sanguineis, imprimis in sup. duobus striata.
227. *P. NEWSHAMIANUM* (Sweet ger. t. 144.) foliis cordatis 3-lobis inæqualiter grossèque dentatis scabris, stipulis cordatis acutis subdentatis, pedunculis 2-3-floris, calycibus reflexis, tubo nectarifero calyce sesqui-longiore. 5 Pe-

tala alba, 2 sup. purpureo lineato-maculata. Natum è semine *P. lanceolati* à quodam ignoto fecundati.

228. *P. BLANDUM* (Sweet ger. t. 4. non Horn.) foliis cordatis leviter 5 lobis dentatis pubescentibus, floralibus bracteisque ovatis acutis, umbellis sub-6-floris, tubo nectarifero calyce parum longiore. ♀ Pet. 3 inf. elliptica alba, 2 sup. obovata sanguinea lineata et maculata.

§. 3. *Tomentosa, petalis albis angustis, foliis cordatis molliter tomentosis, stipulis pateptissimis.*

229. *P. TOMENTOSUM* (Jacq. ic. rar. 3. t. 537. coll. 5. p. 140.) caule fruticoso carnosio, ramis foliisque pedunculis hirsuto-tomentosis, foliis cordatis hastato-5-lobis mollissimis, umbellis multifloris paniculatis, tubo nectarifero calyce triplò ferè breviorè. ♂ Sweet ger. t. 168. Petala alba lineà brevissimà sanguineà colorata. Filam. sanguinea antheræ aurantiacæ. Folia trita Menthæ odora. (v. v.)

♀. *bicolorum*, petalis 2 superioribus lilacinis maculà ramosà notatis, infer. albis lineà purpureà medio instructis. *P. villosum* Spin! cat. 1818. p. 83. non Sweet. (v. s.)

230. *P. RIBIFOLIUM* (Jacq. ic. rar. 3. t. 538. coll. 5. p. 141.) caule fruticoso carnosio, ramis pedunculisque piloso-subhispidis, foliis cordatis hastato-5-lobis scabris, umbellis multifloris, tubo nectarifero calycis longitudine. ♂ Petala et filamenta alba; antheræ aurantiacæ.

§. 4. *Papilionacea, foliis cordatis planis dentatis, petalis inferioribus linearibus, superioribus purpurascentibus lineatis.*

231. *P. PAPILIONACEUM* (Ait. h. kew. 2. p. 423.) ramis foliis pedunculisque subpilosis, foliis cordatis subrotundo-angulatis dentatis, umbellis paniculatis multifloris, petalis infer. subulato-linearibus calyce brevioribus. ♂ Dill. h. elth. t. 128. f. 155. Ger. papilionaceum Lin. sp. 945. Cav. diss. 4. t. 112. f. 1. Pel. papilionaceum Sweet ger. t. 27. Pet. sup. purpurascentia basi alba medio subfusca, inf. albidà. (v. s.)

232. *P. CORDATUM* (Ait. h. kew. 2. p. 427.) foliis cordatis acutis dentatis planis subtus incanis pubescentibus, ramis pedunculisque subpilosis, umbellis multifloris paniculatis, petalis inferioribus subulato-linearibus acutis calyce longioribus, tubo nectarifero calyce paulò breviorè. ♂ G. cordifolium Cav. diss. 4. t. 117. f. 3. Pel. cordatum L'her! ger. t. 22. *P. cordifolium* bot. mag. t. 165. (v. s.)

233. *P. ERIOPHYLLUM* (Sweet ger. t. 141.) foliis cordatis obtusis obsolete lobatis obtusè denticulatis planis subtus incanis mollibus, umbellis plurifloris subdenticulatis, petalis inferis angusto-spathulatis, tubo nectarifero calyce parum longiore. ♀ *P. cordatum* ♀ tomentosum hortul. Hybridum è *P. cordato* et alt. ign.

234. *P. RUBRO-CINCTUM* (Link enum. h. ber. 2. p. 191.) foliis cordatis acutè crenulatis glaberrimis, stipulis linearibus reflexis, umbellis multifloris. ♂ affine *P. cordato*, sed folia glaberrima rubro cincta; stipulæ rubentes. †.

235. *P. CONDUPLICATUM* (Willd. enum. 705.) foliis subrotundo-ovatis basi truncato-subcordatis inciso-dentatis subundulatis subtus incanis pubescentibus, umbellis sub-4-floris, petalis infer. linearibus calyce longioribus. ♂ *P. cordifolium* hortul.

§. 5. *Purpurascentia, foliis cordatis aut cuneatis dentatis indicis aut sublobatis, lobis obtusis ad medium usque non divis, floribus purpurascentibus, petalis inf. oblongis obovatisve.*

* *Foliis subindivisis cucullatis.*

236. *P. CUCULLATUM* (Ait. h. kew. 2. p. 426.) foliis reniformibus cucullatis dentatis pubescentibus, ramis pedunculisque molliter hispidis, umbellis

- 5-floris, petalis infer. oblongis, tubo nectarifero calyce brevior. \S Seb. mus. 1. t. 26. f. 2. G. cucullatum Lin. sp. 946. Cav. diss. 4 p. 241. t. 106. f. 1. (v. v.)
237. *P. MULTINERVÆ* (Sweet ger. t. 17.) foliis orbiculato-reniformibus obsolete lobatis inæqualiter dentatis multinerviis subvillosis, ramis villosis, stipulis cordatis obliquis acuminatis, umbellis plurifloris, petalis inf. oblongis, tubo nectarifero longitudine calycis. \S Pet. sup. basi maculâ albâ lineisq. nigriscentibus striata et medio maculâ atro-purpureâ. Hybridum forte è *P. cucullato* et anguloso. (v. v.)
238. *P. SPECIOSUM* (Willd. enum. 706.) foliis suborbiculatis truncato-reniformibus acutè cartilagineo-dentatis multinerviis subpubescentibus, ramis puberulis, stipulis cordato-lanceolatis acuminatis, umbellis plurifloris, petalis inf. oblongis, tubo nectarifero calyce brevior. \S Pet. 2 sup. medio albida venis atro-purpureis ramosis striata. Folia subcucullata. Priori valdè affine. Hybridum fortè è *P. cucullato* et cardiifolio. (v. v. in hort. Wallner.)
239. *P. COCHLEATUM* (Willd. enum. suppl. 48.) foliis orbiculato-ovatis subcordatis concavis subangulosis serratis pubescentibus, umbellis 5-floris, tubo nectarifero longitudine calycis. \S *P. concavifolium* hortul.
240. *P. CARDIIFOLIUM* (Sweet ger. t. 15.) foliis cuneatis cucullato-plicatis sub-5-lobis cartilagineo-dentatis multinerviis pubescentibus, ramis puberulis, umbellis multifloris, tubo nectarifero calyce duplò longiore. \S Pet. 2 sup. ceteris vix majora medio maculâ atro-purpureâ notata sublineata.
241. *P. ACERIFOLIUM* (L'her! ger. t. 21. non Cav.) foliis basicuneatis integris apice palmatum 5-lobis dentatis multinerviis subvillosis, ramis pedunculisque molliter hispidis, umbellis sub-5-floris, stipulis cordato-ovatis, tubo nectarifero calyce subbreviore. \S G. citriodorum Cav. icon. 1. p. 6. t. 8. (v. s.)
242. *P. CUNEATUM* (Spin! cat. 1818. p. 31.) foliis basi cuneatis subcucullatis obtusè 3-5-lobis apice serrato-dentatis villosis multinerviis, ramis pedunculisque molliter hispidis, umbellis sub-5-floris, stipulis cordato-ovatis acutis, tubo nectarifero calyce dimidiò brevior. \S Petala purpurascencia angustiora et breviora quàm in *P. acerifolio* cujus verosim. progenies hybrida.
243. *P. ANGULOSUM* (Ait. h. kew. 2. p. 426.) foliis basi truncatis subcucullatis subrotundis obtusè 5-lobis dentatis pubescentibus, stipulis cordato-ovatis acuminatis, ramis pedunculisque molliter hispidis, umbellis plurimis 4-6-floris, tubo nectarifero calyce dimidiò brevior. \S — Dill. h. elth. t. 129. f. 156. Ger. acerifolium Cav. diss. 4. t. 112. f. 2. non L'her!
244. *P. BALBISIANUM* (Spin! cat. 1818. p. 30.) foliis basi truncatis subcucullatis subrotundis obtusè 5-lobis apice serrato-dentatis pubescentibus multinerviis, stipulis cordato-ovatis acuminatis, ramis pedunculisque molliter hispidis, umbellis plurimis 4-6-floris, tubo nectarifero calyce paulò longiore. \S Flores omninò *P. angulosi* aut *acerifolii* et verosim. hybrida progenies. (v. s.)
245. *P. ERUBESCENS* (Spin. cat. 1809. p. 27.) foliis longè petiolatis villosiusculis lobatis denticulatis basi rotundatis, margine et nervis rubellis, ramis villosis, umbellis 4-5-floris. \S Flores *P. acerifolii*.
246. *P. BARRINGTONI* (Willd. enum. 706.) foliis reniformibus obtusis cucullatis denticulatis utrinquè pilosis, umbellis multifloris. \S An à sequente satù diversa. Stirps in hortis sub hoc nomine mihi obvia vix à *P. cucullato* differt, nisi foliorum nervis subtus rubentibus, tubo nect. calycis ferè longitudine — *P. Torimauni* Dietr. beschr. gaertn. 3. aug. 235. Ab eo tantùm differt glabritie, sed mihi ignotum.
247. *P. DENNISIANUM* (Sweet ger. t. 20.) foliis cordato-reniformibus acutis concavis undulatis denticulatis, umbellis plurifloris, tubo nectarifero calyce duplò ferè brevior. \S Pet. 2 sup. purpurea medio subnigricantia lineata, 3 inf. pallidiora sublineata. Hybridum fortè ex *P. cucullato* et *Barringtonii* ex Sweet.

248. *P. CALYGINUM* (Sweet ger. t. 81.) foliis orbiculato-reniformibus subangulatis æqualiter denticulatis mollibus, umbellis plurifloris, calycibus inflatis, tubo nectarifero calyce triplò brevior. ♂ Variat petalis sup. atropurpureo maculatis lineatisve et floribus subplenis. Hybridum è *P. cucullato* ♀ et forsan fulgido ♂.
249. *P. WATSONI* (Link enum. 2. p. 192.) foliis cordatis rotundatis sublobatis dentato-crenatis margine undulatis, stipulis cordatis acutis subdentatis, umbellis plurifloris paniculatis, tubo nectarifero calyce reflexo subæquali. ♂ Petala roseo-purpurea, 2 sup. nigro lineata, purpureo maculata. Sweet ger. t. 130. *P. serotinum* Colv. cat. p. 22. Hybridum fortè *P. cucullati* et alt. sp. ign.
250. *P. FORMOSUM* (Sweet ger. t. 120. non Desf.) foliis rotundato-cordatis undulatis concavis rigidis simpliciter denticulatis subutis multinerviis, stipulis cordato-lanceolatis acuminatis, umbellis multifloris subpaniculatis, petalis inferis obovatis. ♂ Pet. inf. pallidè lilacina, sup. purpurascencia venis nigris. Hybridum fortè è *P. cucullato* et cordato, saltem iis affine.
251. *P. VILLOSUM* (Sweet ger. t. 100.) foliis inferioribus oblatis inæqualiter cartilagineo-dentatis villosis basi truncatis, superioribus rhomboideis, umbellis plurifloris, calycibus reflexis, tubo nectarifero calyce brevior. ♂ Petala purpurascencia, 2 sup. fusco-lineato-maculata. Hybridum fortè è *P. cardiifolii* et angulosi. Sweet.
252. *P. MOSTYNE* (Sweet ger. t. 10.) foliis reniformibus undulato-lobatis dentatis pedunculis calycibusque villosissimis, pedunculis paniculatis plurifloris, tubo nectarifero calyce subbrevior. ♂ Pet. 2 sup. medio atropurpurea lineata, 2 sup. oblonga pallida. Hybridum fortè è *P. cucullati* et cordati, sed folia magis lobata.
253. *P. SPECTABILE* (Sweet ger. t. 136.) foliis cordatis sublobatis undulatis inæqualiter grossèque dentatis, caule pedunculis calycibusque villosis, umbellis multifloris, bracteis ciliatis pedicellis longioribus, tubo nectarifero calycis circiter longitudine ♂ Petala ampla roseo-purpurea, 2 sup. nigro lineata et maculata. Hybridum dicitur è *P. cucullato* et ignescente.
254. *P. NERVOSUM* (Sweet ger. t. 47.) foliis reniformibus leviter 5-lobis subcucullatis inæqualiter grossèque dentatis multinerviis, stipulis cordatis acuminatis, umbellis plurifloris, petalis sup. revolutis, tubo nectarifero longitudine calycis. ♂ Pet. 2 maculà atropurpureâ latâ ad marginem lineatâ. Hybridum, sed par. ign.
255. *P. RIGESCENS* (Sweet ger. t. 112.) foliis rotundato-cordatis concavis subangulatis simpliciter denticulatis glabriusculis rigidis, junioribus zonatis, umbellis plurifloris, petalis inferis lingulatis, tubo nectarifero calycis longitudine. ♂ Petala inf. pallidè lilacina, sup. purpurea, venis nigricantibus ramosis. Fortè hybridum è *P. cordato* et alt. sp. ign.
256. *P. SOLUBILE* (Sweet ger. t. 24.) foliis orbiculato-reniformibus subcucullatis rigidis sublobatis dentatis, stipulis cordatis acuminatis, umbellis paniculatis subsexfloris, tubo nectarifero longitudine calycis. ♂ Petala purpurea, 2 sup. vix margine lineolata, medio notata maculâ atropurpureâ in centro albo obliquè striatâ. Affine *P. cochleato* et cucullato.
257. *P. COARCTATUM* (Sweet ger. t. 70.) foliis cordato-reniformibus leviter 5-lobis undulatis denticulatis coarctatis, caule humili, ramis brevissimis confertis, umbellis plurifloris, tubo nectarifero calyce paulò brevior. ♂ Petala purpurea, 2 sup. vix majora lineis duabus ramosis nigricantibus striata. Hybridum dicitur è *P. Barringtonii* et sp. alt. ign.
258. *P. ÆMULUM* (Sweet ger. t. 160.) foliis cordato-reniformibus sublobatis denticulatis concaviusculis, stipulis lanceolatis acutis, umbellis plurifloris, petalis omnibus obovatis, tubo nectarifero calyce reflexo parum brevior. ♂ Petala lætè purpurea, 2 sup. intensiora atropurpureo lineata. Hybridum fortè *P. cucullati* et Beaufortiani.

259. *P. LEPIDUM* (Sweet ger. t. 156.) foliis infer. cordatis sub-5-lobis undulatis inæqualiter crenato-dentatis hirsutis, super. ovatis basi cuneatis, stipulis cordato-ovatis acutis subdentatis, umbellis sub-4-floris, tubo nectarifero calyce reflexo multò brevior. 5 Priori et calycino valdè affine et verosim. hybridum.

** *Foliis lobatis planiusculis.*

260. *P. BARKSIANUM* (Sweet ger. t. 64.) foliis cordatis ovatis leviter et obtusè 5-lobis crenato-dentatis planis margine undulatis, caule petiolis calycibusque villosissimis, umbellis multifloris, tubo nectarifero calyce brevior. 5 Pet. pallidè purpurea, 2 sup. lineata submaculata. Hybridum dicitur è *P. quercifolio* et spec. quadam foliis orbiculatis.

261. *P. PULCHERRIMUM* (Sweet ger. t. 134.) foliis reniformibus planis obsolete lobatis margine undulatis inæqualiter dentatis basi truncatis, caule erecto ramosissimo, pedunculis plurifloris, tubo nectarifero longitudine calycis. 5 Petala roseo-purpurea sublineata, 2 sup. fusco maculata. Hybrida verosim. progenies, sed parentes ignoti.

262? *P. RIGIDUM* (Willd. sp. 3. p. 681.) foliis subrotundis basi truncatis leviter 3-lobis obtusis planis scaberrimis margine crispato-denticulatis, umbellis multifloris. 5 Frutex rigidus. Petala albida oblonga, 2 sup. basi lineis binis violaceis coloratis. An potiùs inter sect. *Hermannifolii*?

263? *P. ADULTERINUM* (L'her. ger. t. 34.) foliis cordatis obtusè trilobis undulatis villosis mollibus, pedunculis subbifloris, tubo nectarifero calyce dimidiò brevior. 5 Petala violacea, 2 sup. basi lineis ramosis atro-purpureis. (v. s.)

264. *P. OBTUSIFOLIUM* (Sweet ger. t. 25.) foliis subcordatis 3-lobis planis obtusissimis, lobis divaricatis inæqualiter dentatis, stipulis cordatis latis, pedunculis paucifloris, tubo nectarifero calyce longiore. 5 Petala rosea, 2 sup. intensiora lineis nigricantibus striata. Hybridum ex *adulterino* et sp. alt. ign.

265. *P. SEYMOURIAE* (Sweet ger. t. 37.) foliis basi truncatis (subcuneatis subcordatisve) trifidis undulatis, lobis latis obtusissimis denticulatis, pedunculis paucifloris, calycibus nervosis reflexis, tubo nectarifero calyce longiore. 5 Petala purpurea, infer. obtusissima, 2 sup. medio atro-purpureo maculata et striata. Hybridum forte è *P. solubile* et sp. alt. ign.

266. *P. ATROPURPUREUM* (Sweet ger. t. 152.) foliis truncato-subcordatis trilobis undulatis inæqualiter argutè dentatis, stipulis cordato-lanceolatis acuminatis subdentatis, umbellis plurifloris, petalis rotundato-obovatis, tubo nectarifero calyci æquali aut brevior. 5 Flores ferè *P. Seymouriae*. Natum dicitur è semine *P. Hoareani* à *P. Principissæ* fecundati, sed nimis differt et errorem quemdam suspicor.

267? *P. SEMITRILOBUM* (Jacq. h. schœnbr. 2. t. 130.) foliis basi truncatis subcuneatis 3-fidis planis hirtis, lobis divaricatis apice serratis, pedunculis subbifloris, petalis infer. linearibus. 5 Petala pallidè incarnata, 2 sup. basi sanguineo striata.

268. *P. VITIFOLIUM* (Ait. h. kew. 2. p. 425.) foliis cordatis trilobis scaberrimis obtusis dentatis, stipulis latis cordatis, caule erecto, umbellis multifloris capitatis, tubo nectarifero calyce triplò brevior. 5 — Dill. h. elth. 1. 126. f. 153. Ger. vitifolium Lin sp. 947. Cav. diss. 4. t. III. f. 2. Petala rosea, 2 sup. lineis atro-sanguineis striata. (v. v.)

269. *P. CAPITATUM* (Ait. h. kew. 2. p. 425.) foliis cordatis lobatis undatis molliter villosis dentatis, stipulis latis cordatis, caulibus diffusis, umbellis multifloris capitatis, tubo nectarifero calyce triplò brevior. 5 *G. capitatum* Lin. sp. 947. Cav. diss. 4. t. 105. f. 1. *G. Rosa hortul.* ob folia rosam spiciantia. Petala lætè purpurea, 2 sup. lineata. (v. v.)

270. *P. RUBENS* (Willd. enum. suppl. 48.) foliis subcordatis acutis levissimis

- 5-lobis serratis, umbellis 5-floris subcapitatis, pedicellis vix involucri longioribus, tubo nectarifero calyce quadruplò brevior. ♂ Petala pallidè violacea, 3 inf. lineà simplici sanguineà bifidà, 2 sup. lineis 4 ramosis picta. Wendl. coll. 2. t. 54.
271. *P. PRINCIPISSE* (Sweet ger. t. 139.) foliis cordato-reniformibus sublobatis inæqualiter mucronato-dentatis, stipulis acuminatis, umbellis plurifloris, petalis omnibus rotundato-obovatis, tubo nectarifero calyce triplò brevior. ♀ Pet. intensè purpurea, 2 sup. lineis nigris et maculà nigricante notata. Hortensis progenies sed parentes ignoti.
272. *P. HUSSEYANUM* (Sweet ger. t. 92.) foliis cordatis 5-lobis denticulatis, lobis rotundatis undulatis, umbellis multifloris capitatis, bracteis pedicellis longioribus, petalis omnibus obovatis, tubo nectarifero calyce triplò ferè brevior. ♂ Pet. amplissima purpurea, 3 inf. pallidiora, 2 sup. medio atropurpureo lineato-maculata. Hybridum videtur è *P. capitato* et *Barringtonii*.
273. *P. WELLSIANUM* (Sweet ger. t. 175.) foliis inferioribus cordatis 5-lobis obtusis inæqualiter dentatis planis, super. trilobis incisive acutis, umbellis plurifloris, calycibus reflexis, tubo nectarifero calyce sesquilongiore. ♂ Petala purpurea, 2 sup. basi alba ceterum nigro lineata et maculata. Dicitur hybridum è *P. Husseyano* à *P. ignescente* fecundati.
274. *P. CONCINNUM* (Sweet ger. t. 108.) foliis oblatis trifidis undulatis obtusè denticulatis glabrisculis basi truncatis, lobis lateralibus bilobis, medio trilobo, stipulis dentatis, umbellis sub-4-floris, tubo nectarifero longitudine calycis reflexi. ♂ Petala latè purpurea, 2 sup. nigro lineata et purpureo intensiore maculata. Hortense hybridum sed parent. ignoti.
275. *P. CORUSCANS* (Sweet ger. t. 173.) foliis cordatis sublobatis inæqualiter acutè dentatis utrinquè pubescentibus, umbellis plurifloris, petalis undulatis reflexis, tubo nectarifero calyce multò brevior. ♂ Petala ampla purpurea lineis nigris omnia lineata, 2 sup. maculà nigricante medio notata. Hybridum sed parentes ign.
276. *P. VERSICOLOR* (Sweet ger. t. 78.) foliis cordatis 5-lobis subvillosis, lobis rotundatis argutè duplicato-dentatis, stipulis cordatis acuminatis dentatis, pedunculis plurifloris, tubo nectarifero calyce duplò brevior. ♂ Petala obovata, 3 inf. carnea lineis intensioribus sublineata, 2 sup. purpurea lineis nigricantibus valdè striata. Folia summa basi subcuneata. Affine *P. bellulo* et *Daveyano*.
277. *P. THYNNÆ* (Sweet ger. t. 74.) foliis cordato-reniformibus 3-lobis planis rigidis glabrisculis, lobis obtusissimis inæqualiter dentatis, stipulis cordatis acuminatis ciliatis, umbellis plurifloris, tubo nectarifero calycis longitudine. ♂ Pet. magna 3 inf. pallidè violacea sublineata, 2 sup. purpurea lineis nigricantibus striata submaculata.
278. *P. BELLULUM* (Sweet ger. t. 60.) foliis cordatis subcuneatis undulatis 5-lobis pubescentibus, lobis obtusis inciso-dentatis, stipulis latis cordatis ciliatis, pedunculis sub 4-floris, tubo nectarifero calyce paulò brevior. ♂ Pet. purpurea lineis intensioribus striata, 3 inf. oblonga pallidiora. *P. rosetta* hortul.
279. *P. RUBASCENS* (Sweet ger. t. 30.) foliis cordatis subreniformibus 5-lobis undulatis inæqualiter profundè dentatis molliter villosis, stipulis ovatis acuminatis dentatis, umbellis plurifloris, tubo nectarifero calyce paulò longior. ♂ Flores ampli. Pet. 3 inferiora carnea oblongo-cuneata, 2 sup. obovata chermesino-coccinea lineis nigricantibus striata. Hybridum fortè è *P. cucullato* et ignescente.
280. *P. COLVILLII* (Sweet ger. t. 86.) foliis cordatis subreniformibus undulatis 7-lobis cartilagineo-dentatis rigidis, umbellis multifloris, petalis inferis ligulatis, calycibus reflexis, tubo nectarifero calyce longiore. ♂ Pet. chermesino-coccinea, 2 sup. lineis ramosis et maculà atropurpureà notata.

281. *P. TIBBISIANUM* (Sweet ger. t. 158.) foliis infer. cordatis trilobis undulatis inæqualiter grossèque dentatis, superioribus planis cuneatis, stipulis lanceolatis subdentatis, umbellis plurifloris, bracteis pedicellis longioribus, tubo nectarifero calyce parùm longiore. 5 Petala lætè purpureo-rosea, sup. nigro lineata. Hybridum *P. involuerati* à *P. ignescente* secundati.
282. *P. DAVATANUM* (Sweet ger. t. 32.) foliis cordato-reniformibus undulatis leviter 5-lobis dentatis molliter villosis, umbellis sub 4-floris, tubo nectarifero calyce paulò breviorè. 3 Petala omnia obovata, 3 inf. intensè rosea, 2 sup. obscurè punicea venis atro-purpureis notata. Medium inter *P. ignescentem* et *Barringtonii*.
283. *P. OBLATUM* (Sweet ger. t. 35.) foliis cordatis rotundato-5-7-lobatis apice recurvis basi inter lobos conduplicatis æqualiter dentatis villosis, umbellis multifloris, calycibus reflexis, tubo nectarifero subsessili calyce sesquolongiore. 3 Petala pallidè purpurea obovata, 2 sup. venis maculisque atro-purpureis notata. Hybridum forsàn è *P. capitato* et *eximio*.
284. *P. PAVONINUM* (Sweet ger. t. 40.) foliis cordato-cuneatis 5-lobis rigidis duplicato-dentatis, marginibus crispis incurvis, petiolis apice dilatatis canaliculatis marginibus inflexis, umbellis multifloris, tubo nectarifero calycis longitudine. 3 Pet. 3 inf. ligulata pallidè coccinea, 2 sup. obovata intensè coccinea medio maculà atro-purpureà, basi maculà albà, et lineis nigricantibus notata. Hybridum fortè è *P. ignescente* et solabili.
285. *P. EXIMIMUM* (Sweet ger. t. 26.) foliis cordatis rugosis undulatis 7-lobis inæqualiter dentatis villosis basi dilatatis, umbellis multifloris, calycibus reflexis, tubo calyce subbreviorè. 3 Pet. 3 inf. pallidè purpurea, 2 sup. intensiora, venis atro-purpureis confertis submaculata. Hybridum fortè è *P. capitati* et angusti.
286. *P. JENKINSONI* (Sweet ger. t. 154.) foliis infer. cordatis obsolete lobatis simpliciter denticulatis planis multinerviis, super. cuneato-ovatis acutiusculis, umbellis sub-4-floris, calycibus acuminatis, tubo nectarifero calyce parùm longiore. 5 Pet. 3 inf. pallidissimè rosea, 2 sup. latè purpureo-maculato maculata. Hortensis progenies è parentibus ignotis.
287. *P. BROWNII* (Sweet ger. t. 148.) foliis sub-5-lobis acutis argutè inæqualiter dentatis rigidis multinerviis basi subcuneatis, umbellis plurifloris, petalis omnibus obovatis, tubo nect. calyce parùm breviorè. 5 Pet. pallidè subroseo-lilacina, 2 sup. latè atro-purpureo-maculata. Progenies hortensis sed parentes ign. et affinitas dubia.
288. *P. FLORIDUM* (Sweet ger. t. 41.) foliis cordatis 5-lobis inæqualiter et argutè dentatis subvillosis, stipulis latis cordatis carinatis subdentatis, umbellis numerosis multifloris, tubo nectarifero longitudine calycis. 3 P autumnale hortul. Petala rosea, 3 inf. vix basi lineata, 2 sup. lineata et medio maculà latè atro-purpureà notata. Hybridum ex *acerifolio* et sp. alt. ign.
289. *P. ASPERIFOLIUM* (Sweet ger. t. 169) foliis cordatis profundè lobatis acutè serratis utrinque hirtis, umbellis sub-5-floris, calycibus reflexis, tubo nectarifero calyce sesquolongiore. 5 Pet. roseo-purpurea, 2 sup. venis nigro-purpureis. Ger. *asperifolium* Andr. ger. ic. An hybridum *P. obtusilobi*, *Beaufortiani* aut *grandiflori*?
290. *P. PECTINIFOLIUM* (Sweet ger. t. 66.) foliis reniformibus trilobis plicatis glabris obtusis, marginibus crispato-denticulatis, umbellis 3-4-floris, calycibus reflexis, tubo nectarifero calycis longitudine. 3 Pet. 3-inf. oblonga angusta obtusa carnea, 2 sup. obovata rosea basi lineis ramosis atro-purpureis striata.
291. *P. LOUSADIANUM* (Sweet ger. t. 44.) foliis infer. basi dilatatis cordatis trilobis, lobis sublobatis argutè dentatis, superioribus basi cuneatis subtrilobis, pedunculis 2-3-floris, calycibus reflexis, tubo nectarifero calyce duplò longiore. 3 Pet. 3 inf. ligulata obtusa carnea, 2 sup. obovata subrosea medio lineis et maculà atro-purpureis notata. Affine *P. Seymouriæ*.
292. *P. ATROFUSCUM* (Sweet ger. t. 82.) foliis superioribus breviter petiolatis

trilobis rigidis inæqualiter argutè dentatis basi cuneatis, umbellis plurifloris, calycibus reflexis tubo nectarifero calyce brevior. $\frac{3}{2}$ Pet. 3 inf. elliptico-oblonga obtusa pallidè purpurea, 2 superiora obovata intensè purpurea medio maculà atropurpureâ notata.

\S . 6. *Crispa, foliis lobatis, lobis apice acutè dentatis.*

293. *P. obtusilobum* (Willd. enum. p. 711.) foliis profundè trilobis, lobis rotundato-obtusis inæqualiter denticulatis, venis subtiùs calycibusque scabriusculis, stipulis cordatis, pedunculis 2-5-floris, tubo nectarifero calyce duplò ferè longiore. $\frac{3}{2}$ Sweet ger. t. 8. Affine *P. tricuspidato*. Petala 3 inf. rosea, 2 sup. intensiora lineis atropurpureis picta.

294. *P. Comptoniæ* (Sweet ger. t. 122.) foliis planis basi cuneatis vel truncato-cordatis trilobis inæqualiter obtusè denticulatis, lobis divaricatis obtusis, stipulis cordatis obtusiusculis, umbellis 3-4-floris, calycibus patentibus, tubo nectarifero longitudine calycis. $\frac{3}{2}$ Petala pallidè purpurea, 2 sup. fusco maculata. Natum è semine *P. obtusilobi* à sp. ign. fecundati.

295. *P. Scabroroviæ* (Sweet ger. t. 117.) foliis subtrifidis glabris ciliatis rigidis inæqualiter argutè dentatis, segmentis inferioribus divaricatis super. trilobis, stipulis cordatis acuminatis, umbellis 3-4-floris, tubo nectarifero calyce parùm brevior. $\frac{3}{2}$ Pet. 3 inf. pallidè rosea, 2 sup. basi nigro-purpurea, medio purpurea, margine subrosea. Hybridum fortè *P. obtusilobi* et sp. alt. ign.

296. *P. Lamberti* (Sweet ger. t. 104.) foliis cuneatis trifidis, lobis acutis lobatis inciso-serratis, caule ramosissimo, stipulis cordatis, pedunculis paucifloris, calycibus reflexis, tubo nectarifero calyce longiore. $\frac{3}{2}$ Pet. roseo-purpurea, 2 sup. basi intensius fusco maculata. Hybridum fortè è *P. acerifolio* aut scabro.

297. *P. tricuspidatum* (L'her! ger. t. 30.) foliis basi cuneatis trifidis, lobis acutis, medio productiore subserrato, costà subtus muricatà, pedunculis bifloris, tubo nectarifero calycis longitudine. $\frac{3}{2}$ Petala alba, 2 sup. purpureo maculata. Hybridum fortè è *P. hermännifolio* et scabro. (v. s.)

298. *P. trilobatum* (Schrad. h. goett. 1. p. 8. t. 2.) foliis basi cuneatis trilobis, lobis divaricatis apice inæqualiter et argutè serratis, lateralibus subbilobis, marginibus nervisque subtus puberulis, pedunculis subbifloris, tubo nectarifero calyce brevior. $\frac{3}{2}$ Petala pallidè rosea, 2 sup. venis purpureis striata. Medium inter *P. crispum* majus et adulterinum. *P. clandestinum* Spin! cat. 1818. p. 31. (v. v.)

299. *P. scabrum* (Ait! h. kew. 2. p. 430.) foliis basi cuneatis trifidis scabris, lobis lanceolatis laxè serratis, pedunculis 1-4 floris, tubo nectarifero calyce brevior. $\frac{3}{2}$ Ger. scabrum Lin. sp. 946. Cav. diss. 4. t. 108. f. 1. — Jacq. ic. rar. 3. t. 542. L'her! ger. t. 31. Petala alba aut rosea, 2 sup. lineata. (v. s.)

300. *P. hermännifolium* (Jacq. ic. rar. 3. t. 545.) foliis cuneiformibus distichis scabris plicatis apice truncatis inciso-dentatis, pedunculis bifloris brevibus, tubo nectarifero calyce paulò longiore. $\frac{3}{2}$ *G. hermännifolium* Lin. mant. 569. Berg! cap. 177. Petala alba vix subcarnea, 2 sup. lineis 2 purpureis basi striata. (v. s.)

301. *P. crispum* (Ait. h. kew. 2. p. 430.) foliis distichis subrotundo-carnosis basi subcuneatis trifidis undulato-plicatis scabris dentatis, pedunculis subbifloris, tubo nectarifero calycis longitudine. $\frac{3}{2}$ Ger. crispum Lin. mant. 257. Cav. diss. 4. t. 109. f. 2. *P. crispum* L'her! ger. t. 33. Petala purpurea, 2 sup. lineata. Folia odore Melissæ. (v. s.)

\mathcal{L} . *majus* (L'her! ger. t. 32.) foliis profundius trilobis majoribus, tubo nectarifero calyce sublongiore. (v. s.)

302. *P. uniflorum* (Spreng. nov. prov. 32.) foliis trilobis dentatis glaucescentibus supra nitidis, pedunculis 1-2-floris elongatis, petalis superioribus emarginatis. $\frac{3}{2}$ An var. *P. crispum*?

303. *P. EXSTIPULATUM* (Ait. h. kew. 2. p. 431.) foliis truncato-cordatis trilobatis dentatis cano-velutinis, stipulis subnullis, pedunculis 3-4-floris, tubo nectarifero calyce triplò longiore. $\frac{3}{5}$ Ger. *exstipulatum* Cav. diss. 4. t. 123. f. 1. — L'her! ger. t. 35. Petala pallidè violacea, sup. sublineata. (v. s.)
304. *P. MIXTUM* (Sweet ger. t. 71.) foliis subdistichis cordatis profundè 5-lobis undulato-plicatis rigidis dentatis, pedunculis 2-4-floris, calycibus reflexis, tubo nectarifero calyce sublongiore. $\frac{3}{5}$ Petala alba, 2 super. basi carnea sublineata, medio rubro maculata.
305. *P. PUSTULOSUM* (Sweet ger. t. 11.) foliis inf. profundè 3-lobis subtùs pustulato-scabris, lobis lateralibus divaricatis inæqualiter acutè dentatis, medio trilobo acutè dentato, caule diffuso, pedunculis subbifloris, tubo nectarifero calyce duplò ferè longiore. — Petala angusta alba vix subcarnea, 2 sup. medio rubro maculata.
306. *P. FRATERNUM* (Schränk pl. rar. mon. 1. t. 16.) foliis truncato-subobovatis 3-partitis, lobis cuneatis apice dentatis, lateralibus bifidis, medio trifido, pedunculis 4-5 floris, tubo nectarifero calyce breviorè. $\frac{3}{5}$ Petala calyce duplò triplòve majora oblonga alba subcarnea immaculata.
307. *P. PALLIDUM* (Willd. enum. 710.) foliis profundè trilobis, lobis divaricatis apice inæqualiter acutè dentatis subtùs margineque scabris, pedunculis sub-3-floris. $\frac{3}{5}$ Affine videtur sequenti.
308. *P. TERNATUM* (Jacq. ic. rar. 3. t. 544.) foliis trisectis cucullatis scabris, lobis cuneiformibus apice inciso-serratis, medio trifido, pedunculis 1-2-floris, tubo nectarifero calyce paulò longiore. $\frac{3}{5}$ Sweet ger. t. 165. Ger. *ternatum* Lin. suppl. 306. Cav. diss. 6. t. 107. f. 2. — bot. mag. t. 428. — Subglaucescens. Petala rosea, 2 sup. sanguineo-lineata. (v. s.)
- §. 7. *Radulæ, foliis ultrà medium partitis, lobis dentatis incisim pinatifidis, floribus purpurascens aut pallidis.*
309. *P. QUERCIFOLIUM* (Ait. h. kew. 2. p. 420. L'her! ger. t. 14.) foliis cordatis pinatifidis, sinibus rotundatis, lobis obtusis orenatis, ramulis petiolisque hispidis, umbellis submultifloris, filamentis apice adscendentibus, tubo nectarifero calyce sublongiore. $\frac{3}{5}$ G. *quercifolium* Lin. suppl. 306. Cav. diss. 4. t. 119. f. 1. G. *terebinthinaceum* Murr. goett. 1785. t. 4. Odor foliorum gravis. Petala roseo-purpurea, 2 sup. lineata. (v. v.)
- β. bipinnatifidum* (L'her. ger. t. 15.) foliis duplicato-pinnatifidis. (v. s.)
310. *P. LECHIANUM* (Spin! cat. 1818. p. 32.) foliis cordatis pinnatifidis, sinibus acutis, lobis obtusis dentatis, ramulis petiolisque molliter brevè velutinis, umbellis multifloris, tubo nectarifero calyce vix longiore. $\frac{3}{5}$ Petala roseo-albida, 2 sup. rosea venis atro-purpureis picta. An hybridum è *P. quercifolio* et *capitato*? (v. s.)
311. *P. AUGUSTUM* (hortul! — an Link enum. 2. p. 192?) foliis subcordatis 5-lobis subvillosis, lobis ovali-oblongis dentatis, terminali acuto, ramis pedunculisque villosis, umbellis 4-5-floris confertis, tubo nectarif. calyce dimidio ferè breviorè. $\frac{3}{5}$ Petala pallidè rosea, 2 sup. atro-purp. lineis striata. Flores ferè *P. Vandesii*. (v. v. in hort. Walluer.)
312. *P. BLANDFORDIANUM* (Sweet ger. t. 101.) foliis palmato-7-lobatis canescentibus, segmentis pinnatifido-incisis obtusè dentatis planis, umbellis multifloris, petalis superioribus obcordatis, tubo nectarifero calycis longitudine. $\frac{3}{5}$ G. *Blandfordium* Andr. ger. ic. Petala sordidè albida, 2 sup. maculis livido-purpureis. An hybridum è *P. graveolente* et *cortusæfolio* aut *echinato*? (v. v.)
313. *P. GRAVEOLENS* (Ait. h. kew. 2. p. 423.) foliis palmatis 7-lobatis, lobis oblongis obtusis dentatis margine revolutis, umbellis multifloris subcapitatis, tubo nectarifero calyce duplò breviorè. $\frac{3}{5}$ L'her. ger. t. 17. G. *terebinthinaceum* Cav. diss. 4. p. 250. t. 114. f. 1.

314. *P. PIPERATUM* (Spin! cat. 1818. p. 32.) foliis pubescentibus palmatim 5-7-lobis, lobis acuminatis inæqualiter serratis, umbellis 6-10-floris, tubo nectarifero calyce reflexo duplò brevior. ♂ Pet. purpurea, 2 sup. basi himaculata. Odor foliorum piperatus gratus. *P. verbenæfolium* Sweet ger. t. 149. (v. s.)
315. *P. GLUTINOSUM* (Ait. h. kew. 2. p. 426.) foliis cordatis hastato-5-angulis dentatis viscosis glabriusculis, umbellis 2-4-floris, tubo nectarifero calyce paulò longiore. ♂ L'her! ger. t. 20. Curt. bot. mag. t. 143. Jacq. ic. rar. t. 131. Ger. glutinosum Cav. diss. 4. t. 108. f. 1. *G. cratægifolium* Roth abb. t. 9. non Thunb. Petal. pallidè rosea aut albida purpuro-lineata, 2 super. lineato-maculata. (v. s.)
316. *P. VISCOSISSIMUM* (Sweet ger. t. 118.) foliis palmatim 5-7-lobatis canleque viscosis, segmentis planis sinuatis dentatisve apice recurvis, umbellis capitatis multifloris, petalis oblongis obtusis, calycibus obtusissimis, tubo nectarifero calyce parum longiore pedicelli basin subattingente. ♀ Pet. alba, 2 sup. lineis purpureis 2 subramosis notata.
317. *P. HISPIDUM* (Willd. sp. 3. p. 677.) foliis palmatifidis tomentosis hispidis, lobis acuminatis inciso-dentatis, umbellis paniculatis multifloris, tubo nectarifero calyce dimidiò brevior. ♂ *G. hispidum* Lin. suppl. 303. Cav. diss. 4. t. 110. f. 1. Flores parvi. (v. s.)
318. *P. VANDERIE* (Sweet ger. t. 7.) foliis ultrà medium palmatifidis, lobis 7-9 oblongis divisis dentatis scabris margine revolutis, umbellis plurifloris subcapitatis, tubo nectarifero calyce brevior. ♂ Flores ampli pallidè rosei, pet. 2 sup. lineis sanguineæ maculatis.
319. *P. RADULA* (Ait. h. kew. 2. p. 423.) foliis palmati-partitis scabris, lobis angustis pinnatifidis margine revolutis, laciniis linearibus, umbellis paucifloris, tubo nectarifero calyce triplò brevior. ♂ L'her! ger. t. 16. Ger. Radula Cav. diss. 4. t. 101. f. 1. *G. revolutum* Jacq. ic. rar. t. 133. (v. s.)
β. roseum (Willd. sp. 3. p. 679.) foliorum laciniis lineari-lanceolatis. — Folia rosæ centifoliæ odorem spirant. Hybridum è Radulâ ♀ et graveolente ♂. (v. s.)
320. *P. BALSAMEUM* (Jacq. ic. rar. 3. t. 543.) foliis palmatipartitis subscabris basi cuneatis, lobis lanceolatis remotè dentatis, umbellis paucifloris, tubo nectarifero brevissimo. ♂ Petala carnea oblonga, 2 sup. basi submaculata.
321. *P. ASPERUM* (Willd. sp. 3. p. 678.) foliis subpalmatilobis scabris, lobis 5-7 oblongis obtusis margine crispato-denticulatis, umbellis sub-5-floris capitatis, tubo nectarifero calyce brevior. ♂ Ger. Radula Roth abb. p. 51. t. 10. Involucri bractæ magnæ ovatæ acutæ. Pet. carnea, 2 sup. rubro lineata. Simillimum *P. graveolenti*. An satis ab eo differt?
322. *P. DENTICULATUM* (Jacq. schœnb. 2. t. 135.) foliis palmatipartitis viscosis glabris, lobis linearibus pinnatifidis repandis dentatis planiusculis, umbellis paucifloris, tubo nectarifero brevissimo, petalis superioribus bifido-emarginatis. ♂ Petala oblongo-cuneata carnea, 2 sup. maculis rubris 2-3 notata. Sweet ger. t. 109. (v. v.)
323. *P. JATROPHÆFOLIUM* (DC. cat. h. monsp. 1813.) foliis palmatipartitis viscosis glabris, lobis lanceolato-linearibus pinnatifidis, lobulis dentatis distantibus acuminatis, umbellis 4-floris, tubo nectarifero brevissimo, petalis superioribus obtusis. ♂ Medium ferè inter *P. denticulatum* et *quercifolium* var. pinnatifidum. (v. v.)
324. *P. DELPHINIFOLIUM* (Willd. enum. 708.) foliis scabris palmatim 5-lobis, lobis oblongis serratis, intermedio trilobo, umbellis paucifloris compositis, tubo nectarifero pedicello longiore. ♂ Petala incarnata, 2 sup. cuneata lineis binis et maculâ atro-sanguineâ notatis, 3 inf. linearia acuta. Affine, ex foliis, *P. balsameo*.

†. *Pelargonium non satis nota.** *Acaulia, ad Hoaream Dimacriamve referenda.*

325. *P. ANGUSTIFOLIUM* (Thunb. fl. cap. ed. Schult. 2. p. 514. sub Geranio.) acaule, foliis ellipticis integris glabris marginatis, scapo hirsuto umbellifero. ♀ Pet. lineari-obovata alba. *P. longifolio* affine.
326. *P. ENSATUM* (Thunb. l. c. sub Ger.) acaule, foliis ensiformi-obovatis hirsutis umbellâ proliferâ. ♀ Pet. alba lanceolato-obovata.
327. *P. SETIFERUM*, acaule, foliis subrotundis duplicato-serratis subtus pubescentibus, umbellis bifloris. ♀ Ger. setosum Thunb. l. c. 517.
328. *P. SIDOIDES*, acaule, foliis cordatis orbiculatis tomentosis, scapis bifidis umbelliferis. ♀ Ger. sidæfolium Thunb. l. c. 518. Pet. purpureo-nigricantia.
329. *P. HETEROLOBUM*, acaule, foliis trifidis pinnatisectisque subtus setosis, scapis bifidis sulcatis, umbellis 60-floris. ♀ Ger. heterophyllum Thunb. l. c. 515. Pet. alba obovata.
330. *P. ARENARIUM*, acaule, radice perpendiculari longâ, foliis pinnatifidis, lobis oblongis inciso-serratis, scapis 2-3 folio longioribus 4-5-floris. ♀ Ger. arenarium Burm. ger. n. 65. Cav. diss. 4. p. 259. Flores ex rubro violacei.
331. *P. TRILOBUM* (Thunb. l. c. 519. sub Ger.) acaule, foliis pinnatisectis, pinnis 5 ovatis ciliatis, umbellâ proliferâ. ♀ Pet. obovata sanguinea.
332. *P. Plicatum* (Thunb. l. c. 526. sub Ger.) acaule, foliis tomentosis pinnatisectis, pinnis plicatis, scapo umbellifero. ♀ Pet. pallidè incarnata, majora purpureo-maculata.
333. *P. ACAULE* (Thunb. l. c. 526. sub Ger.) acaule, foliis subbipinnatifidis hirsutis scapo umbellifero. ♀ Pet. albida.
334. *P. TOMENTELLUM*, caule brevissimo fruticoso, foliis bipinnatisectis tomentosis, umbellâ subtriflorâ. ♂ ♀ Ger. tomentosum Thunb. fl. cap. 2. p. 527.
335. *P. VILLIFERUM*, caule brevissimo, foliis tripinnatifidis villosis, umbellis multifloris. ♂ ♀ Ger. villosum Thunb. l. c. 528. Affine *P. hirtum*.

** *Herbacea.*

336. *P. APHANOIDES* (Thunb. l. c. 514.) foliis 5-partitis tomentosis, lobis incisis, pedunculis elongatis, umbellis 4-5-floris. ♀ Aff. *P. alchimilloidi* videtur.
337. *P. DONDÆFOLIUM* (Link enum. 2. p. 187.) caule petiolisque hirsutis, foliis 5-lobis anticè acutè-crenatis, pediculis longissimis, bracteis linearibus. ♀ Fol. infer. leviter lobata, sup. partita. Affine videtur *P. tabulari*.
338. *P. RUPHRASEUM* (Horn. h. hafn. 2. p. 634.) foliis cordatis trifidis incisis crenatis, petiolis longissimis, pedunculis paucifloris ♀.
339. *P. ACUMINATUM* (Thunb. l. c. 526.) caule herbaceo trigono, foliis inciso-pinnatifidis glabris, pedunculis bifloris. ♀ Pet. albida vel incarnata, majora purpureo maculata.
340. *P. FLEXUOSUM* (Thunb. l. c. 526.) caule suffrutescente erecto, foliis tomentosis trifidis, umbellis subtrifidis. ♂ ♀ Pet. albida, majora purpureo-maculata.

*** *Carnosa.*

341. *P. DISCIPES* (Haw. succ. p. 305.) caule carnoso ramoso arboreo, foliis cordatis peltatis pubescentibus variabiliter glauco-nitentibus, petiolis villosis basi exstipulatis, ramis subsimplicibus apice excepto defoliatis. ♂ Flores ignoti. An ad *Pelargia pinguifolia*? an ad *Isopetali* sect. referendum?
342. *P. TECTUM* (Thunb. l. c. 525.) caule fruticoso carnoso, foliis inciso-triternatis lanatis, umbellis multifloris. ♂ Pet. saturatè purpurea.
343. *P. PINGUE* (Thunb. l. c. 527.) caule fruticoso carnoso, foliis pinnatifidis hirtis, umbellis multifloris. ♂ Pet. albida.

**** *Fruticosa.*

344. *P. ERYTHRÆUM*, foliis oblongis serratis tomentosis, caule frutescente, pedunculis alternis subunifloris. ♂ Ger. rubens Thunb. l. c. 515.
345. *P. ELLIPTICUM* (Thunb. l. c. 525.) caule fruticoso, foliis ellipticis incanis, umbellâ subtriflorâ. ♂ Pet. sanguinea.
346. *P. SERRATUM* (Thunb. l. c. 517.) caulescens flexuoso-erectum, foliis subrotundis serratis, veniâ scabris, pedunculis subunifloris. ♂ Pet. obovata sanguinea.
347. *P. LANATUM* (Thunb. l. c. 518.) caule fruticoso hirsuto, foliis cordatis ovatis serratis subtus lanato-tomentosis. ♂ Pet. obovata, incarnata maculâ sanguineâ.
348. *P. SIDAIFOLIUM* (Willd. enum. 706.) foliis subrotundis cordatis subangulatis mucronato-denticulatis, umbellis sub-4-floris. ♂ Flores pallidè violacei. Affine *P. specioso*.
349. *P. TOTUM* (Thunb. l. c. p. 523.) caule fruticoso villosa, foliis 5-partitis scabris, lobis lacero-dentatis umbellis sub-4-floris. ♂ Pet. incarnata, majora purpureo maculata.
350. *P. SPURIUM* (Willd. enum. 709.) foliis reniformibus subdistichis levissimè subtrilobis obtusis inæqualiter dentatis undulatis, pedunculis 3-4-floris. ♂ Corollæ violacæ, pet. 2 sup. lineatis. Tubus nectarifer pedunculo longior. Affine forsân *P. scabro*.
351. *P. PANICULATUM* (Horn. suppl. 76. non Jacq.) foliis reniformibus pubescentibus 9-lobis, lobis rotundatis mucronato-dentatis, umbellis multifloris coarctato-paniculatis. ♂ An idem ac *P. formosum*.
352. *P. SUAVEOLENS* (Desf. arb. 2. p. 465.) villosum, foliis cordato-rotundatis crispis 5-lobis argutè denticulatis, pedunculis multifloris, pedicellis abbreviatis calycibusque villosissimis. ♂ Cor. purpurea. An affine *P. crispo*? — *P. suaveolens* Horn. h. bafn. 2. p. 640 videtur diversum.
353. *P. HORNEMANNI*, foliis cordato-reniformibus trifidis argutè dentatis glabris carnosis cucullatis, umbellis 4-floris. ♂ Habitus *P. tricuspidati*. *P. blandum* Horn. suppl. 76. non Sweet.
354. *P. CASTELLIANUM* (Spin cat. 1818. p. 30.) foliis basi cuneatis cucullatis angulosis dentatis scabris, ramis petiolisque moliter pilosis, umbellis multifloris. ♂ Petala pallidè rosea, 2 sup. rotunda basi maculâ purpureâ ramosâ, inf. linearia obtusa.
355. *P. GRATUM* (Willd. enum. p. 710.) foliis leviter trifidis inæqualiter acutè dentatis undulatis pilosis, pedunculis 2-4-floris. ♂ Odor plantæ citratus. Petala incarnata, 2 sup. obovata retusa lineis ramosis atro-sanguineis picta, 3 inf. lineari-cuneata.
356. *P. NOTRUM* (Willd. enum. p. 710.) foliis subrotundo-ovatis obtusis subtrifidis conduplicatis undulato-dentatis subtus pilosiusculis, pedunculis subbifloris, calycis laciniis erectis. ♂ Affine *P. grato* et ad *P. crispum* verosim. referendum.
357. *P. CONSANGUINEUM* (Willd. enum. 710.) foliis leviter trilobis planis obtusis, lobis divaricatis inæqualiter argutè dentatis, pedunculis subtrifloris. ♂ Affine *P. grato* sed folia inodora.
358. *P. PUBESCENS* (Spin cat. 1818. p. 32.) foliis trilobis inciso-dentatis pilosis margine ciliatis, ramis petiolis pedunculis calycibusque piloso-pubescentibus, pedunculis sub-5 floris. ♂ Folia citrato odore pollut. Petala carnea, sup. subrotunda lineis purpureis ramosis, inf. linearia.
359. *P. CUNEATUM* (Spin cat. 1818. p. 31.) foliis subvillosis cuneatis trilobis serratis, umbellis paucifloris, petalis calycis lobos subæquantibus. ♂ Petala rubro-purpurea, sup. rotundata lineis atro-purpureis, inf. linearia obtusa.
360. *P. WILDENOWII* (Link enum. 2. p. 190. non Dietr.) foliis subrotundo-

cuneatis levissimè trilobis undulato-denticulatis, ramis petiolis pedunculisque villosopilosis, umbellis 3-4-floris, tubo nectarifero calyce longiore. § *P. pulchellum* Willd. suppl. 47. non Ait. Petala alba, 3 inf. oblonga, 2 sup. medio maculâ lineisque sanguineis picta. Ad *Pelargia lineata* pertinere videtur.

361. *P. UNICOLORUM* (Willd. enum. suppl. 48.) foliis subrotundo-cuneatis leviter trilobis acutè undulato-dentatis, ramis petiolis pedunculisque pilosis, umbellis quadrifloris, tubo nectarifero calyce reflexo duplò breviorè. § Corolla sanguineo-violacea, petala 2 sup. basi venis saturatoribus instructa.
362. *P. ALNIFOLIUM* (Willd. enum. 710.) foliis ellipticis obtusis, floralibus obsolete subtrifidis inæqualiter dentatis basi subcuneatis integerrimis, pedunculis 1-4-floris. § Ad *P. betulinum* accedit. Corolla incarnata, petalis 2 sup. latioribus basi lineis atro-sanguineis ramosis pictis.
363. *P. AMPLISSIMUM* (Willd. enum. suppl. 48.) foliis planis glaberrimis semi-orbiculatis 7-lobis serratis basi leviter cordatis, pedunculis 2-5-floris, tubo nectarifero calyce reflexo duplò longiore. § Corolla magnitudine et colore *P. speciosi*. Aff. *P. grandifloro*?
364. *P. EGREGIUM* (Hort. belved. 71. Link enum. 2. p. 195.) foliis trisectopinnatifidis pubescentibus, lacinulis breviter acutatis, stipulis ovatis, umbellis trifloris, tubo nectarifero calycis longitudine. § Petala coccinea.
365. *P. FRAGRANTISSIMUM* (Link enum. 2 p. 192.) foliis basi subrotundatis lobatis, lobis mediis rotundatis, lateralibus acutatis, omnibus acutè serratis, tubo nectarifero calyce quadruplò breviorè. § Petala purpurascencia lineis ramosis purpureis. Stipulæ ovatæ reflexæ.
366. *P. LADANOMA* (Hoffmaus. in Dietr. nachr. 6. p. 68. ex Link. enum. 2. p. 192.) foliis subrotundis sublobatis acutè crenulatis, cor. lilacinis.
367. *P. EXSUL* (Hoffmans. et Link. l. c.) foliis tripartitis acutissimè dentato-incisis, stipulis subcordatis. *P. errans* hortul.
368. *P. MUNITUM* (Burch. cat. geog. n. 1240. voy. 1. p. 225.) glabrum, foliis bipinnatifidis, paniculâ dichotomâ spinoso-liguescente. — ad cap. B.-Spei prope Juk-River.
369. *P. DIVARICATUM* (Thunb. l. c. 525.) caule fruticoso erecto glabro, foliis multifido-pinnatifidis glabris, pedunculis 1-floris. § Pet. incarnata, majora purpureo maculata.

Ordo XLVII. TROPÆOLEÆ. *Juss. mem. mus.* 3. p. 447.

Calyx 5-partitus coloratus, laciniâ superiore basi calcaratâ, calcare libero intrâ florem hinc hiantē extrâ stamina, lobis nunc liberis, nunc inter se variē coalitis. Petala 5 calyci inserta, ejusdem lobis alterna, inæqualia, irregularia, 2 superiora sessilia remota, calcaris fauci infixa, 3 inferiora unguiculata minora interdum abortiva. Stamina 8, filamentis liberis ovarium arcuē cingentibus, disco (perigyno?) insertis, antheris terminalibus oblongis erectis bilocularibus rimâ duplici dehiscentibus. Carpella 3 arcuē connata in ovarium 1 trigonum. Styli 3 in 1 tristriatum filiformem connexi. Stigmata 3 acuta. Carpella 3 imo stylo seu fructus axi adnata, 1-locularia, 1-sperma. Semina magna, exalbuminosa, loculum suum implentia et hujus cavitati conformia. Embryo magnus; cotyledonibus 2, rectis, crassis, junioribus distinctis, dein arcuē conferruminatis et etiam cum spermodermate adhærentibus, imâ basi subdistinctis; radiculâ intrâ cotyledonum processus latente, tubercula 4 mox radicellas proferentiâ gerente. (Gaertn. fr. 1. p. 380. t. 79. Rich. ann. p. 90. Saint-Hil. ann. mus. 18. p. 461. t. 24.)

Herbæ Americanæ, sapore nasturtiano donatæ, glabriusculæ, teneræ, diffusæ aut volubiles. Folia alterna, non stipulacea, petiolata, peltinervia. Pedunculi axillares 1-flori. Characteres spermici è solo Tropæolo eruti.

Ordo Geraniaceis affinis ex calycis petalorumque structurâ et situ, differt staminibus liberis et numero cum petalis non congruentibus, floribus axillaribus et præsertim fructus et seminum structurâ. An tribus inter Geraniaceas? An ordo legitimus?

I. TROPÆOLUM. *Lin. gen. n.* 466.

Cal. 5-partitus, lobo sup. calcarato. Pet. 5 inæqualia, 3 inferiora minora aut evanida. Stam. 8 ab ipsâ basi libera. Carpella 3 suberosa reniformia indehiscentia hinc sulcata rotundata. Semina (conf. char. ord.)

§. 1. *Foliis peltinerviis integris lobatisve.*

1. *T. minus* (Lin. sp. 490.) foliis peltinerviis orbiculatis subrepandis, nervis apice mucronatis, petalis acuminato-setaceis. ① sub dio, ② in cald: — in Peruvîâ. Curt. bot. mag. t. 98. Schkuhr handb. t. 105. Variat flore pleno et tunc ②. Flores sesquipoll. diam., pallidiûs flavi. (v. v.)
2. *T. majus* (Lin. sp. 490.) foliis peltinerviis orbiculatis subquinquelobis, nervis apice non exsertis, petalis obtusis. ① ② in Peruvîâ undè in Europam allatum anno 1684. Curt. bot. mag. t. 23. Hern. mex. 161. ic. Variat etiam fl. pleno. (v. v.)

3. *T. HYBRIDUM* (Lin. mant. 64.) foliis peltinerviis orbiculatis subcucullatis 5-lobis, petalis variantibus. \mathcal{L} in horto Holmiensi à præced. semine ortum, ramis propagatum; potius varietas monstrosa.
4. *T. ADUNCUM* (Smith tour. 1. p. 158.) foliis peltinerviis subreniformibus 5-7-lobis, lobis integris mucronatis, petalis 2 super. lobatis mucronatis, 3 inf. minoribus fimbriatis, calcare adunco petalorum sup. longitudine. $\textcircled{1}$ in Peru, Mexico. *T. peregrinum* Jacq. schœnbr. t. 98. bot. rep. t. 617. bot. mag. t. 1351. bot. reg. t. 718. H. B. et Kunth nov. gen. am. 5. p. 252. non Lin. (v. v.)
5. *T. SMITHII*, foliis peltinerviis profundè palmatilobis, petalis omnibus lobatis fimbriatisque, calcare recto corollâ duplò longiore. — in Novâ-Granadâ. *T. peregrinum* Lin. excl. Feujll. syn. ex Smith in Rees cycl. 36. n. 4.
6. *T. DIPETALUM* (fl. per. 3. p. 77. t. 313.) foliis subpeltinerviis 5-7-lobis, lobis ovatis submucronatis subtus glaucis, petalis 2 spatulatis crenulatis. \mathcal{L} in Andium montibus. Feujll. per. 2. t. 42? et hinc forsân verum *T. peregrinum* Linn. sp. 940. (v. s.)
7. *T. BIGOLORUM* (fl. per. 1. c. t. 313.) foliis peltinerviis 7-lobis basi transversim truncatis, petalis inciso-ciliatis. — in Andium nemoribus humidis. Petala 2 superiora minora lutea, 3 majora puniceo-coccinea.
8. *T. TUBEROSUM* (flor. per. 3. t. 314.) foliis peltinerviis 5-lobis basi transversim truncatis glabris, petalis calycis ferè longitudine. \mathcal{L} in Peruviz præruptis. Radices tuberosæ post coctionem edules. Petala dentata ex Ruiz et Pav., integerrima ex H. B. et Kunth nov. gen. am. 5. p. 251.
9. *T. PUBESCENS* (H. B. et Kunth nov. gen. am. 5. p. 251.) foliis peltinerviis obsoletè acutè 5 lobis basi truncatis suprâ glabriusculis subtus pubescentipilosis, petalis lobato-2-aristatis calycem subæquantibus. $\textcircled{1}$ in sylvis prope Loxam Peruv.
10. *T. CILIATUM* (fl. per. 3. p. 77.) foliis peltinerviis 5-7-lobis, stipulis bracteolisque ciliatis, petalis integerrimis calycem subæquantibus. — in R. Chilensis sylvis.

§. 2. *Foliis peltatisectis.*

11. *T. PENTAPHYLLUM* (Lam. dict. 1. p. 605. ill. t. 277.) foliis peltatisectis, segmentis 5 ovalibus integris, petiolulatis, petalis omnibus sessilibus calyce brevioribus acutis integerrimis. \mathcal{L} in Monte-Video (v. s.)
12. *T. POLYPHYLLUM* (Cav. ic. 4. p. 65. t. 395.) foliis peltatisectis, segmentis 5-10 oblongis obovatisve subdentatis basi cuneatis, petalis unguiculatis calyce sublongioribus obtusis integerrimis. $\textcircled{1}$ in Chilensi monte Cordillerâ.
13. *T. PINNATUM* (And. bot. rep. v. 8. t. 535.) foliis subpeltatis, lobis obtusis inæqualibus, floribus pinnatis, petalis 5 cuneatis apice deutatis. $\textcircled{1}$ Patr. ign.; cult. in hort. Angliæ. — Charact. ex Andr. loc. cit.

II. MAGALLANA. Cav. ic. 4. p. 50.

Cal. 5-lobus calcaratus, lobis 2 profundè partitis, 3 ceteris in unicum tridentatum coalitis. Petala 5 inæqualia. Stam. 8 imâ basi subcoalita. Fructus 3-alatus abortu 1-locul, 1-spermus. Semina non satis nota.

1. *M. PORRIFOLIA* (Cav. ic. 4. p. 51. t. 374.) $\textcircled{1}$ in Americâ merid. ad Portum Desideratum juxta sepes scandens. Folia 3-partita, lobis linearibus integris.

Ordo XLVIII. BALSAMINEÆ. *A. Rich. dict. class. 2. p. 173.*
 — *Impatiens Lin. gen. n. 1008.* — *Balsamina Tourn. inst.*
418. t. 235. Juss. gen. 270. Gaertn. fr. 2. p. 151. t. 113.

Calyx 2-sepalus, sepalis parvis deciduis oppositis sæpè mucronatis per æstivationem imbricatis. Petala 4 hypogyna cruciata; 2 exteriora sepalis alterna apice callosa; superius fornicatum emarginatum, inferius integrum basi in calcar productum; 2 interiora prioribus alterna, magis petaloidea, inter se æqualia, sæpè bifida seu appendiculata. Stamina 5 hypogyna ovarium arcuè cingentia, filamentis brevibus apice incrassatis, antheris subconnatis, 3 inferiora petalis opposita anthera-constanter biloculares ovatas gerentia, 2 superiora antè petalum supremum orta, antheras nunc biloculares, nunc 1-loculares gerentia, undè in quibusdam staminis unci profundi bipartiti locum tenentia. Antheræ rimâ longitudinali dehiscences. Ovarium 1. Stylus 0. Stigmata 5 distincta aut in unicum coalita sessilia brevissima. Capsula oblonga aut ovata, 5-valvis, valvulis elasticè dissilientibus, placenta centralis apice in fila tenuia stigmatum basim attingentia, ut in Caryophylleis, producta, 5-angularis, angulis membranaceis suturas intervalvulares attingentibus et ideò capsula junior 5-locularis, apice suprâ placentam unilocularis. Semina placentæ adfixa, pendula, in quoque loculo plurima, ovato-oblonga, exalbuminosa. Embryo rectus, radiculâ superâ, cotyledonibus intus planis, extus convexis.

Herbæ teneræ. Folia alterna aut opposita exstipulata, simplicia, penninervia, dentata. Stipulæ nullæ. Pedunculi axillares. Flos ferè Fumariacearum. Capsula ferè Oxalidis. Embryo ferè Lini. Habitus proprius.

I. BALSAMINA. *Riv. irr. tetr. ic.*

Antheræ 5 biloculares. Stigmata 5 distincta. Capsula ovata, valvis ad maturitatem introrsum apice elasticè inflexis. Cotyledones crassæ. — Pedicelli semper 1-flori, solitarii aut aggregati. Flores in hortis faciliè pleni. Capsulæ puberulæ.

§. 1. *Pedicellis geminis ternis aggregatisve.*

1. *B. HORTENSIS* (Desp. dict. sc. nat. 3. p. 485.) pedicellis adgregatis, foliis lanceolatis, superioribus alternis, calcaribus flore brevioribus. ① in Indiâ orient. *Impatiens Balsamina* Lin. sp. 1328. Variat in hortis flore pleno, albo, roseo, rubro et ex his col. vario. Blackw. t. 583. Rheed. mal. g. t. 52? (v. v.)
2. *B. COCCINEA*, pedicellis adgregatis, foliis alternis oblongo-ovalibus serratis, petiolis multiglandulosis, calcaribus incurvis florem subæquantibus. ① in Indiâ orient. *Impatiens coccinea* Sims bot. mag. t. 1256.

12. *I. RACEMOSA*, pedunculis 7-8-floris folio ferè æqualibus racemosis, foliis glabris oblongis grossè dentatis utrinquè acuminatis, calcare recurvo tenui flore sublongiore. ① in Napanlià, Wallich. Flores flavi parvi. (v. s.)
13. *I. LEPTOCERAS*, pedunculis 6-8-floris erectis racemosis, foliis glabris oblongis utrinquè attenuatis serrato-mucronatis, calcare tenui recto longissimo. ① in Napanlià, Wallich.
- α. pedunculis folio longioribus. (v. s.)
- β. pedunculis foliis brevioribus. (v. s.)
14. *I. insignis*, pedunculis 8-20-floris erectis folio longioribus racemosis foliis, glabris oblongis utrinquè attenuatis, dentibus præsertim infimis mucronatis, calcare recto longissimo. ①? in Napanlià, Wallich. Flores purpurascens. (v. s.)

Ordo XLIX. OXALIDEÆ.

Cal. 5-sepalus 5-partitusve persistens æqualis. Pet. 5 hypogyna, æqualia, interdum infernè inter se cohærentia unguiculata, unguibus erectis, laminis patentibus, per æstivationem spiraliter contorta. Stam. 10, filamentis subulatis erectis basi sæpius monadelphis, 5 sepalis opposita exteriora breviora sæpè dentata, 5 petalis opposita interiora longiora. Antheræ biloculares, non adnatæ. Ovarium liberum 5-angulare 5-locul. Styli 5 filiformes, nunc staminibus omnibus breviores et *brevisissimi* dicti, nunc stam. exterioribus longiores interioribusque breviores et *intermedii* dicti, nunc omnibus stam. longiores et *longissimi* dicti. Stigmata penicilliformia, capitata aut subbifida. Capsula ovata aut oblonga sub-5-gona membranacea 5-locul. 5-10-valvis, ad angulos longitudinaliter dehiscens. Semina pauca, angulo loculamentorum centrali adfixa, ovata, striata, juniora inclusa intrà arillum carnosum demum ab apice elasticè dehiscentem et semen projicientem. Albumen cartilagineo-carnosum. Embryo inversus, longitudine albuminis, cotyl. foliaceis, radic. superâ longâ. — Suffrutices aut Herbæ. Folia alterna, rariùs opposita aut subverticillata, simplicia aut varè composita. — Ordo Geraniaceis olim adnexus, ad Rutaceas aut Zygophylleas potiùs accedit, ipso monente cl. Jussieo (mem. mus. 3. p. 448.), sed ab utroque satis diversus videtur habitu et characteribus.

I. AVERRHOA. *Lin. gen. n. 576. Lam. ill. t. 385. Corr. ann. mus. 8. p. 71. t. 2.*

Cal. sepala 5 inter se plus minùs basi coalita. Pet. 5. Stam. 5 petalis alterna, aut 10 alterna minora, basi in annulum brevem coalita. Ovarium angulatum. Styli 5 persistentes. Bacca magna oblonga 5-sulcata 5-loc. Semina in quoque loculo pauca; angulo centrali adnata. Embryo rectus in albumine carnosio. Cotyl. foliaceæ. — Arbores Indicæ. Folia alterna impari-pinnata, foliolis alternis. Flores racemoso-paniculati. Fructus edules. — Genus affine (ex cl. Correa) Rhamneis, (ex cl. Brown) potiùs Oxalideis. Folia A. Bilimbi tactu irritabilia ut in Biophyto.

1. A. CARAMBOLA (Lin. sp. 613.) calycibus glabris, petalorum limbo subrotundo, staminibus 5, fructibus acutangulis, seminibus arillatis. 5 in Indiâ orient. — Cav. diss. 7. t. 220. — Rumph. amb. 1. t. 35. (v. s.)

2. A. BILIMBI (Lin. sp. 613.) calycibus pubescentibus, petalorum limbo ovali-oblongo, staminibus 10, fructibus obtusangulis, seminibus exarillatis. 5 in Indiâ orient. — Cav. diss. 7. t. 219. — Rumph. amb. 1. t. 36. (v. s.)

II. BIOPHYTUM.

Cal. 5-sep. Pet. 5. Stam. 10 omninò libera, 5 ext. alterna breviora. Styli 5 apice emarginato-bifidi. Capsula ovato-globosa sub-5-gona. —

Herbæ annuæ, caule nudo, foliis ad apicem fasciculato-verticillatis abruptè pinnatis, foliolis multijugis oppositis, pedunculis apice umbellato-multifloris.

1. *B. SENSITIVUM*, pedunculis apice multifloris. ① in Indiæ orient. hortis et pratis. Folia valdè tactu irritabilia Mimosas simulant. — *Oxalis sensitiva*. Lin. sp. 622. Jacq. ox. n. 21. t. 73. f. 4. — Rumph. amb. 5. t. 104. f. 2. — Rheed mal. 9. t. 19. (v. s.)
2. *B. DENDROIDES*, pedunculis unifloris. ①? 3? in regno Novo-Granatensi inter Guaduas et Quebrada de la Carbonera. *Oxalis dendroïdes* H. B. et Kunth nov. gen. am. 5. p. 250. Caulis lignosus. Rachis foliorum pilos, inter quodque foliorum par, fasciculatos gerens. Corolla violacea. Ovarii loculi 1-spermi.

III. OXALIS. Lin. gen. n. 582. — *Oxys* Tourn. inst. t. 19.

Cal. 5-sepalus, sepalis liberis aut basi coalitis. Pet. 5. Stam. 10, filam. basi breviter inonadelphis, 5 ext. alternis brevioribus. Styli 5 apice penicilliformes aut capitati. Capsula pentagona oblonga aut cylindracea. — Herbæ perennes, caulescentes stipitatæ aut acaules, foliis variis sed nunquàm abruptè pinnatis.

§. 1. *Hedysaroïdæ*, pedunculis multifloris; caulibus sæpiùs suffrutescentis foliosis, foliis trifoliolatis, foliolis ovatis lanceolatisve nec obcordatis, medio petiolulato (1), ovarii loculis plerùmque 1-sperminis. — Omnes spec. ex Americâ calid. ortæ.

1. *O. PENTANTHA* (Jacq. ox. n. 1. t. 1.) caule erecto ramoso folioso, pedunculo umbellifero foliis brevioribus, foliis pilosis, foliolis lateralibus subrotundis emarginatis, intermedio obovato obtuso, stylis staminibus exter. brevioribus. 3 ad Caracas.

β. *Humboldtii*, stylis stamina longiora duplò superantibus. *O. pentantha* H. B. et Kunth nov. gen. am. 5. p. 247.* An species propria?

2. *O. PSORALIOIDES* (H. B. et Kunth nov. gen. am. 5. p. 246. t. 470.) caule erecto folioso, ramis tomentosis, pedunculo petiolo longiore apice bifido 6-8-floro umbellifero, foliis tomentosis subtus argenteo-sericeis, calycibus hirtis tomentosis, stylis intermediis. 3 prope Santa-Fe de Bogota? Fl. flavi?
3. *O. PLUMIERI* (Jacq. ox. n. 3.) caule erecto folioso, pedunculo umbellifero 4-floro foliorum longitudine, foliolis integris ovatis obtusiusculis. 3 in Americâ merid. — Plum. ed. Burm. t. 213. f. 1. *O. frutescens* Lin. sp. 624. Flores lutei. Capsulæ glabræ.
4. *O. NÉEII*, caule erecto folioso, pedunculo umbellifero 4-floro foliorum longitudine, foliolis villosis, lateralibus ovalibus emarginatis, medio ovato lanceolato, stylis intermediis. 3 ad Acapulco (Née) Media inter duas priores. (v. s.)
5. *O. BARRELIERI* (Jacq. ox. n. 4. t. 3.) caule erecto ramoso folioso, pedunculo longitudine foliorum bifido racemoso, foliolis ovato-lanceolatis, stylis longitudine staminum interiorum. 3 in Guianâ, Brasiliâ et ad Caracas. — Barr. ic. t. 1169. Petala albo-incarnata, 3 sup. basi flavo-maculata.
6. *O. FRUTICOSA* (Raddi mem. bras. add. p. 22.) caule erecto, pedunculis axillaribus paucifloris, petiolis foliaceis lanceolatis, foliolis subrotundo-ovatis ciliatis, medio longè petiolato. 3 in sepibus et nemoribus prope Rio-Janeiro. Flores aurei.
7. *O. HEDYSARIFOLIA* (Raddi l. c.) caule erecto folioso, pedunculis termina-

(1) Potiùs impari-pinnatis unijugis tunc forsàn dicendis.

libus paucifloris, foliolis ovatis utrinquè acutis petiolisque hirtis. — in herbosis montosis prope Rio-Janeiro.

8. *O. HEDYSAROIDES* (H. B. et Kunth nov. gen. am. 5. p. 247.) caule fruticoso folioso glabro, pedunculis apice bifidis multifloris, foliolis ovato-oblongis emarginatis glabris subtus glaucescentibus, lateralibus minoribus remotis, calycibus glabris. ♂ in frigidis et temperatis Andium Quitensium et prope Santa-Fe de Bogota. Affinis *O. Barrelieri*.

9. *O. GLAUCA* (H. B. et Kunth l. c. t. 471.) caule fruticoso folioso glabro, pedunculis apice bifidis 7-9-floris folii longitudine, foliolis ciliatis glabris glaucis, lateralibus subrotundis remotis, terminali ovato-elliptico, sepalis glabris apice maculâ notatis, stylis intermediis. ♂ in ripâ umbrosâ flum. Amazonum prope Tomependam. Flores albi.

10. *O. BORJENSIS* (H. B. et Kunth l. c.) caule fruticoso folioso glabro, pedunculis apice bifidis multifloris petioli longitudine, foliolis apice rotundatis glabris glaucis, lateralibus subrotundis remotis, terminali subovato-elliptico, sepalis puberulis emaculatis. ♂ in ripâ fl. Orinoci prope S.-Borja locis humidis umbrosis. Flores flavi.

11. *O. ANGUSTIFOLIA* (H. B. et Kunth l. c.) caule fruticoso folioso subglabro, pedunculis 3-4-floris petiolo longioribus, foliolis oblongo-linearibus glabris glaucis, lateralibus minoribus remotis, calycibus glabris emaculatis. ♂ in declivitate occid. Novæ-Hispaniæ prope la Venta del Peregrino. Flores flavi.

12. *O. RHOMBIFOLIA* (Jacq. ox. n. 2. t. 2.) caule erecto ramoso folioso, pedunculo umbellifero trifloro longitudine foliorum, foliolis pubescentibus rhombeo-ovatis sessilibus, stylis intermediis. — ad Caracas. Filam. interiora hispida.

§. 2. *Corniculatæ*. *caulibus basi non bulbosis herbaceis (rarissime suffrutescentibus) foliosis, pedunculis rarius 1-floris sæpius bi- aut multifloris, foliis palmato-trifoliatis, foliolis omnibus sessilibus obcordatis.*

13. *O. PEDUNCULARIS* (H. B. et Kunth nov. gen. 5. p. 239.) caule simplici carnosio folioso, pedunculis longissimis apice trifidis multifloris, foliolis subrotundo ovato ellipticis subtus pubescentibus, stylis intermediis. ① in temperatis montium Quitensium prope Mira et in convalli fl. Chambo. Pedunculi 7-9 poll. longi. Flores aurantiaci aut crocei. Radix fibrosa.

14. *O. PUBESCENS* (H. B. et Kunth l. c.) caule erecto basi ramoso molliter pubescente, pedunculis apice bifidis multifloris petiolo longioribus, foliolis subrotundo obovatis emarginatis utrinquè petiolis calycibusque pubescentibus, stylis stamina longiora æquantibus. 2℥ in Peruvîâ. Radix fibrosa. Flores flavi? Pedunculi 2-4 pollic.

15. *O. CRENATA* (Jacq. ox. n. 7.) caule erecto folioso, pedunculo umbellifero 5-6-floro foliis longiore, foliolis obovatis, petalis crenatis. ① in Peruvîâ. — Feuill. per. 3. p. 49. t. 24. Flores lutei purpuro-striati.

16. *O. PERENNANS* (Haw. misc. 181.) caulibus erectis foliosis flexuosis, pedunculis 2-3-floris petiolo sublongioribus, foliolis bilobo-obcordatis ciliatis, stylis stamina interiora paulò superantibus. 2℥ in Novâ-Hollandiâ. Flores extus fulvi, intus lutei.

17. *O. VERTICILLATA* (fl. mex. ic. ined.) caule erecto simplici glabro, foliis ternatis quaternatisve verticillatis, foliolis obcordatis, pedunculis 2-3-floris petioli longitudine. 2℥ in hortis Sancti-Angeli in Mexico.

18. *O. DILLENII* (Jacq. ox. n. 8.) caule erecto folioso subhirsuto, pedunculis umbelliferis 5-6-floris folio longioribus, foliolis obcordatis, petalis emarginatis, stylis stamina utraque superantibus. ① in Carolinâ et Guadalupâ. — Dill. elth. 2. t. 221. Flores flavi paulò majores quàm in sequente à quâ vix differt. (v. s.)

- β. *florida* (Saliab! prod. 322.) caule decumbente, pedunculis 2-3-floris.
O. glauca Raff. ined. *O. prostrata* Haw. misc. 183? Pedunculi inferiores petiolo longiores, supremi petiolo breviores. — An species propria? (v. s.)
19. *O. STRICTA* (Lin. sp. 624.) caule erecto folioso, pedunculis umbelliferis 2-6-floris folii circiter longitudine, foliolis obcordatis, petalis integris, stylis staminum inter. longitudine. (I) in Americæ bor. et Europæ ruderatis, sepius. Jacq. ox. n. g. t. 4. — fl. dan. t. 873. *O. ambigua* Sal. act. soc. lin. 2. p. 242. t. 23. f. 4. Variat caule subcumbente nunquam repente. Radix reptans. Flores flavi magnitudine sequentis. (v. v.)
20. *O. CORNICULATA* (Lin. sp. 624.) caule decumbente ramoso radicante, pedunculis subumbellatis petiolo brevioribus, foliolis obcordatis, petalis emarginatis, stylis longitudine staminum interiorum. 2℥ in Europæ, Teneriffæ, Japoniæ, Borboniæ, ins. Caribæarum, Mexici, et Amer. bor. cultis. Jacq. ox. n. 10. t. 5. *O. pusilla* Sal. act. soc. lin. 2. p. 243. t. 23. f. 5. Stipulæ ad basin petioli teretis. Flores flavi. (v. v.)
21. *O. VILLOSA* (Bieb! fl. taur. 1. p. 355.) undiquè hirsuta, caule repente ramoso, pedunculis bifloris petiolo longioribus, foliolis obcordatis, stylis longitudine staminum interiorum. (I) in Iberiâ. Priori nimis affinis et forsâ varietas? (v. s.)
22. *O. LYONI* (Pursh fl. bor. am. 1. p. 322.) tota sericeo-pilosa, caule ramoso decumbente, pedunculis 2-3-floris petiolo longioribus, foliolis bilobo-obcordatis, petalis cuneatis, capsulis tomentosis calyce lanceolato duplè longioribus. 2℥ in insulâ Cumberland Georgiæ Americ. Flores flavi. (v. s.)
23. *O. MICROPHYLLA* (Poir. suppl. 4. p. 248.) glabriuscula, caulibus procumbentibus, pedunculis bifloris petiolo longioribus, foliolis bilobo-obcordatis, petalis cuneatis, siliquis puberulis, stylis longitudine stam. inter. (I) in Novâ-Hollandiâ. (I) rubens Haw. misc. p. 183. Fl. flavi parvi. (v. s.)
24. *O. LOTOÏDES* (H. B. et Kunth nov. gen. am. 5. p. 241.) caulibus procumbentibus, ramis hirsuto-pilosis, pedunculis petiolo longioribus 3-4-floris, foliolis subrotundo-obovatis emarginatis adpresso-pilosis margine villosis subtus glaucescentibus, petiolis pilosis, calycibus pubescentibus, stylis intermediis. 2℥ in monte Quindiu. Flores flavi?
25. *O. MOLLIS* (H. B. et Kunth l. c.) caulibus ramosis repentibus petiolisque molliter villosis, pedunculis apice bifidis paucifloris petiolo longioribus, foliolis obovatis emarginatis pubescentibus subtus canescentibus, calycibus puberulis, stylis intermediis. 2℥ prope Almaguer in Audibus Popayanensium. Fl. flavi.
26. *O. SCANDENS* (H. B. et Kunth l. c.) caule ramoso scandente glabro, pedunculis elongatis dichotomo-4-fidis multifloris, foliolis obovatis emarginatis suprâ glabris subtus hirsuto-pubescentibus, petiolis villosis, calycibus glabris, stylis longissimis. 2℥ in monte Quindiu. Fl. magni *O. acetosellæ*, flavi.
27. *O. MEDICAGINEA* (H. B. et Kunth l. c.) caulibus ramosis procumbentibus glabris, pedunculis longissimis apice bifidis 4-12-floris, foliolis obovatis emarginatis ciliatis suprâ glabris subtus pubescentibus, petiolis glabrinsculis, calyce glabro, stylis intermediis. 2℥ in regno Novo-Granatensi? Flores albi?
28. *O. PILOSUSCULA* (H. B. et Kunth l. c.) caulibus subramosis basi repentibus petiolisque pilosis, pedunculis 1-floris pubescentibus, foliolis obcordatis ciliatis subtus adpresso-pilosiusculis, calycibus puberulis. 2℥ prope urbem Caracas. Fl. flavi.
29. *O. LUPULINA* (H. B. et Kunth l. c.) caulibus filiformibus procumbentibus subramosis glabriusculis, pedunculis 1-3-floris, foliolis subrotundo-obovatis emarginato-bilobis calycibusque glabris, stylis longissimis. 2℥ in frigidis Novæ-Granatæ prope Almaguer. Fl. flavi.
30. *O. ALBICANS* (H. B. et Kunth l. c.) caulibus cespitoso ramosis repentibus

puberulis, pedunculis 1-2-floris petiolo æqualibus, foliis obcordato-bilobis glaucescentibus violaceo-marginatis subtus petioli calycibusque puberulis, stylis longissimis. \mathcal{U} prope Moran Mexicanorum. Fl. flavi.

β . *sericea*, caulibus foliis pedunculis calycibus capsulisque sericeo-canescens. — prope Lactagunga Quitensium.

31. *O. FILIFORMIS* (H. B. et Kunth l. c. p. 245. t. 469.) caulibus filiformibus repentibus ramosis glabris, pedunculis 1-floris petiolo longioribus, foliis subrotundo-obovatis emarginato-bilobis margine pilosis subtus puberulis, petioli pubescentibus, calyce ciliato, stylis longissimis. \mathcal{U} in Andibus Novo-Granatensium prope Guadas. Fl. flavi?

32. *O. PARVIFOLIA*, caulibus ramosis repentibus glabris, ramulis foliosis, pedunculis 1-floris, foliis subrotundo-obovatis emarginato-bilobis ciliatis utrinque adpresso-pilosis, petioli calycibusque pubescentibus, stylis longissimis. \mathcal{H} in regno Quintensi. *O. microphylla* H. B. et Kunth l. c. non Poir. Priori simillima.

33. *O. REPENS* (Thunb. ox. n. 11. t. 1. f. 5.) caule folioso ramoso decumbente radicante, pedunculis subbifloris petioli longitudine, foliis obcordatis, stylis intermediis. — in Ceylonâ, Madagasc., cap. B.-Spei. — Jacq. ox. n. 11. t. 78. f. 1. *O. stricta* Houtt. pl. 6. t. 51. f. 2. Flores flavi. Filam. glabra. (v. s.)

34. *O. TUBEROSA* (Sav. in Lam. dict. 4. p. 684.) caule ramoso, pedunculis umbelliferis, foliis ovatis, radice tubercula plurima gerente. \mathcal{U} in Chili. Affinis *O. corniculatæ*. Mol. chil. ed. gall. p. 103. \uparrow .

35. *O. CONORHIZA* (Jacq. ox. n. 6.) caule erecto-folioso, pedunculo foliis longiore bifloro, foliis obcordatis, radice turbinatâ. \mathcal{U} in Paraguai. Flores ampli flavi. — Feuill. per. 2. p. 723. t. 24. \uparrow .

36. *O. ROSEA* (Jacq. ox. n. 5.) caule erecto folioso, pedunculis axillaribus folio quadruplò longioribus apice corymboso-racemosis, foliis obcordatis. — in Chili locis humidis. Sims bot. mag. t. 2415. — Feuill. obs. 2. p. 23. *O. racemosa*. Sav. in Lam. dict. 4. p. 684. Flores rosei.

37. *O. LATRIFLORA* (Jacq. h. schœnbr. 2. p. 41. t. 204.) caule ascendente subramoso basi nudo, pedunculis lateralibus apice umbellatis, foliis cuneatis emarginato-bilobis, stylis stam. exter. brevioribus. — ad cap. Bonæ-Spei. Flores purpurei. Filamenta hispida. Hæc et prior. ab *O. corniculatis* habitu recedunt.

\S . 3. *Sessilifloræ, caulibus basi bulbosis elongatis sparse foliosis, foliis sessilibus 3-foliolatis villosis non glanduliferis, pedunculis axillaribus unifloris.*

38. *O. MACROSTYLIS* (Jacq. ox. n. 22. t. 9.) caule erecto folioso ramoso, pedunculis folia multoties superantibus, bracteolis calyci approximatis, foliis lineari-cuneatis emarginatis, stylis staminibus inter. longioribus. \mathcal{U} ad cap. B.-Spei. Flores purpurascens tubo longo.

39. *O. TUBIFLORA* (Jacq. ox. n. 23. t. 10.) caule erecto subramoso folioso, pedunculis folia quadruplò superantibus, bracteolis calyci adpressis, foliis lineari-cuneatis obtusis, stylis stam. exter. brevioribus. \mathcal{U} ad cap. B.-Spei. Flores incarnuati tubo longo.

40. *O. CANESCENS* (Jacq. ox. n. 24. t. 11.) caule erecto subramoso folioso, pedunculis folia duplò superantibus, bracteolis calyci approximatis, foliis cuneatis subemarginatis, stylis stam. exter. brevioribus. \mathcal{U} ad cap. B.-Spei. Flores purpurasc. extus pallidi tubo longo. Var. prioris ex Willd. (v. v.)

41. *O. SECUNDA* (Jacq. ox. n. 68. t. 12.) caule declinato ramoso folioso, ramis secundis, foliis lineari-cuneiformibus, pediculis vix folia superantibus, stylis intermediis. \mathcal{U} ad cap. B.-Spei. Flores lilacini tubo longo. Filam. glabra.

42. *O. HIRTA* (Lin. sp. 623.) caule erecto subramoso folioso, foliis lineari-cuneatis retusis, pedunculis folia multoties superantibus, bracteolis à calyce

remotis, stylis Stam. inter. longioribus, Stam. edentulis eglandulos. \mathcal{U} ad cap. B.-Spei. — Jacq. ox. n. 26. t. 13. Flores lilacini fundo flavo tubo brevi.

\mathcal{E} . *brevipes*, pedunculo vix foliis longiore. \mathcal{U} Jacq. ox. t. 77. f. 3.

43. *O. HIRTELLA* (Jacq. ox. n. 27. t. 14.) caule erecto subramoso folioso, foliolis lineari-cuneatis retusis, pedunculis folia multoties superantibus, bracteolis à calyce remotis, stylis Stam. inter. longioribus, staminibus exter. denticulato-gibbis, pilis staminum glandulos. \mathcal{U} ad cap. B.-Spei. Flores pallide lilacini, fundo flavo, tubo brevi. *O. sessilifolia* Lin. mant. 241. *O. hirta* \mathcal{E} . Willd. sp. 2. p. 705.

44. *O. MULTIFLORA* (Jacq. ox. n. 28. t. 15.) caule erecto folioso ramosissimo, foliolis lineari-cuneatis obtusis, pedunculis folia multoties superantibus, bracteolis à calyce remotis, stylis Stam. exter. brevioribus. \mathcal{U} ad cap. B.-Spei. Cor. lilacinæ campanulatæ. Jacq. ic. rar. t. 472.

45. *O. RUBELLA* (Jacq. ox. n. 29. t. 16.) caule erecto folioso ramoso, foliolis lineari-cuneatis, pedunculis folia multoties superantibus, bracteis à calyce subremotis, stylis intermediis. \mathcal{U} ad cap. B.-Spei. Cor. rubellæ campanulatæ. Jacq. ic. rar. t. 471. Sims. bot. mag. t. 1031. (v. v.)

46. *O. ROSACEA* (Jacq. ox. n. 30. t. 17.) caule decumbente simpliciter folioso, foliolis oblongo-cuneiformibus, pedunculis folia multo superantibus, bracteolis à calyce distantibus, stylis intermediis. \mathcal{U} ad cap. B.-Spei. Cor. rubellæ subtus discolores. Stam. ext. denticulato-gibba. — bot. mag. t. 1698 (v. s.)

§. 4. *Caulifloræ, caulibus elongatis sparse foliosis, foliis saltem supremis petiolatis 3-5-foliolatis, pedunculis axillaribus unifloris. — Species inter se habitu dissimiles.*

47. *O. VIRGINEA* (Jacq. h. schœnbr. 3. t. 275.) caule erecto folioso subramoso, folii petiolatis, foliolis 3 lateralibus oblongis medio obovato-cuneato, pedicellis folio brevioribus, stylis intermediis, filam. edentulis. \mathcal{U} ad cap. B.-Spei. Flores albi.

48. *O. REPTATRIX* (Jacq. ox. n. 33. t. 20.) caule erecto brevi folioso, foliis longè petiolatis, foliolis 3 ovato-rotundis, pedunculis folio longioribus, stylis brevissimis, filam. dentatis. \mathcal{U} ad cap. Bonæ-Spei. Corollæ albidæ subincarnatæ.

49. *O. INCARNATA* (Lin. sp. 622.) caule erecto ramoso glabro folioso, foliis petiolatis hinc inde fasciculato-verticillatis, foliolis 3 obcordatis, pedunculis folii longitudine, stylis longissimis, stamin. dentatis. \mathcal{U} ad cap. Bonæ-Spei. Cor. albidæ subincarnatæ. — Jacq. h. vind. t. 71. (v. s.)

50. *O. DISTICHA* (Jacq. ox. n. 31. t. 18.) caule basi ramoso adscendente glabro folioso, foliis sparsis petiolatis, petiolis alato-stipulatis, foliis 3 obcordatis, pedunculis folio longioribus, stylis intermediis, filamentis dentatis. \mathcal{U} ad cap. B.-Spei. Cor. pallide flavæ.

51. *O. VENOSA* (Sav. in Lam. dict. 4. p. 681.) caule erecto hirsuto simplici folioso, foliis infer. breviter petiolatis sparsis, sup. longius petiolatis subverticillatis, foliolis cuneatis emarginatis subtus hispidæ, stipulis subulatis, pedunculis folio longioribus. \mathcal{U} in Africâ. Cor. venosæ violacæ, tubo flavo.

52. *O. EBRACTEATA* (Sav. l. c. p. 682.) caule simplici erecto hirsuto apice folioso, foliis petiolatis, foliolis oblongo-obcordatis, pilis glandulos in margine et costâ, pedunculis ebracteatis petioli longitudine. \mathcal{U} ad cap. B.-Spei. Flores flavi. Au eadem ac *O. disticha*?

53. *O. HETEROPHYLLA*, villosa, caule erecto basi ramoso, foliis infer. sessilibus 3-foliolatis, foliolis obcordatis, super. longè petiolatis foliolis bipartitis, pedunculis petiolo longioribus medio bibracteatis, stylis brevissimis. \mathcal{U} . ad cap. B.-Spei? Radix fibrosa. Flores purpurei. (v. 3.)

54. *O. QUINATA* (Sav. in Lam. dict. 4. p. 688.) caule adscendente superuè fo-

lioso, foliis petiolatis, foliolis 5 cuneiformibus subemarginatis, pedunculis petiolo longioribus medio bibracteatis. \mathcal{L} . ad cap. Bonæ-Spei. Flores purpureascentes tubo flavo.

§. 5. *Caprinæ, acaules aut stipite nudo apice tantum folioso donatæ, pedunculis uni-bi-aut sæpius-multifloris, foliis radicalibus petiolatis plurifoliolatis sæpius trifoliolatis.*

55. O. DECAPHYLLA (H. B. et Kunth nov. gen. am 5. p. 238. t. 468.) acaulis, foliolis 8-10 cuneatis apice bifidis subciliatis glabris, scapis 5-14-floris folio longioribus, sepalis obtusis apice bimaculatis, staminibus inter se subæqualibus stylos duplò superantibus. \mathcal{L} in planitie Mexicanà prope rupem El Peñon. Flores violacei magnitudine O. strictæ.

56. O. HERNANDESI (fl. mex. ic. ined.) acaulis, foliolis 9-11 oblongis villosis, scapo foliis longiore umbellifero 6-floro. \mathcal{L} in Mexico. — Hern. mex. 386. f. 3. Flores erecti pallidè violacei. An à priore certè distincta? †.

57. O. BURMANNI (Jacq. ox. n. 20.) caule brevissimo apice folioso, foliolis 5-6 lanceolatis glabris, scapo foliis longiore umbellifero 7-8-floro. \mathcal{L} ad cap. Bonæ-Spei. — Burm. afr. t. 29. Flores flavi.

58. O. TETRAPHYLLA (Cav. ic. 3. t. 237.) acaulis, foliolis 4 (rariùs 3) obcordatis glabris subtùs glaucescentibus, scapis umbelliferis 3-7-floris, stylis longissimis. \mathcal{L} in Mexico. Cor. purpureo-violaceæ. (v. s.)

59. O. SERICEA (Lin. f. suppl. 245.) acaulis, foliolis 3 obcordatis sericeis, scapo umbellifero foliis longiore, floribus nutantibus, stylis intermediis. \mathcal{L} ad cap. B.-Spei. — Jacq. ox. n. 13. t. 77. f. 1. Cor. flavæ.

60. O. ARTICULATA (Sav. in Lam. dict. 4. p. 686.) acaulis, foliolis 3 obcordatis sericeo-tomentosis, scapis umbelliferis 5-6-floris, calycibus apice callosis. \mathcal{L} in Monte-Video. Flores flavescentes. Radix nodoso-articulata.

61. O. MEGALORHIZA (Jacq. ox. n. 12.) acaulis, foliolis 3 obcordatis subtùs violaceis, scapo umbellifero foliis breviorè, radice crassà multicipite. \mathcal{L} in Peruvîâ — Feuille. per. 2. p. 734. t. 25. O. bicolor Sav. in Lam. dict. 4. p. 687. Flores flavi basi rubro-lineati.

62. O. VIRGOSA (Sav. in Lam. dict. 4. p. 685.) acaulis, foliolis 3 ovatis, scapis multifloris, floribus verticillatis. \mathcal{L} in Chili. Molina chil. p. 110. Scapi 5-pedales! †

63. O. NUDIFLORA (fl. mex. ic. ined.) acaulis, scapis petiolisque pilosis, foliolis 3 obcordatis, floribus umbellatis 5-6 involucro destitutis. \mathcal{L} in agris Mexici. Flores violacei erecti; alabastra nutantia. An à sequente sat diversa? †.

64. O. VIOLACEA (Lin. sp. 621.) acaulis, foliolis 3 obcordatis glabris, scapo umbellifero 3-9-floro, floribus breviter involucratiss nutantibus, stylis brevissimis. \mathcal{L} in umbrosis à Carolinà ad Canadàm. — Jacq. h. vind. t. 180. ox. n. 14. t. 80. f. 2. Flores violacei. Stam. hirta. (v. s.)

65. O. ELEGANS (H. B. et Kunth nov. gen. am. 5. p. 234. t. 466.) acaulis, foliolis 3 latè subrotundo-obovatis emarginatis subtùs violaceis petiolisque glabris, scapis longissimis 2-6-floris, sepalis acuminatis apice bimaculatis, stylis longissimis. \mathcal{L} prope Loxam Peruvianorum. Fl. violacei ampli. Bulbus solitarius.

66. O. JACQUINIANA (H. B. et Kunth l. c.) acaulis, foliolis 3 latè subrotundo-obovatis emarginato-subbilobis petiolisque glabris, scapis 3-6-floris, sepalis obtusis apice bimaculatis, stylis longissimis. \mathcal{L} in Mexico prope Real-del-Monte. Fl. violacei. Bulbus solitarius.

67. O. DEBILIS (H. B. et Kunth l. c.) acaulis, foliolis 3 subrotundis profundè emarginatis subtùs puberulis, scapis 6-floris bifidis petiolisque pilosis, sepalis obtusiusculis apice maculà notatis, stylis longissimis. \mathcal{L} inter la Venta Grande et urbem Caracas. Fl. magn. O. acetosellæ, violacei. Bulbi cespitosi.

68. O. SCHRADERTIANA (H. B. et Kunth l. c.) acaulis, foliolis 3 latè subrotundo-

- obovatis emarginatis glabris, petiolis pilosiusculis, scapis glabris 9-11-floris, sepalis obtusis apice maculâ notatis, stylis Stam. breviora subæquantibus. $\frac{1}{2}$ in monte Quindia. Fl. violacei magni. O. strictæ.
69. O. LATIFOLIA (H. B. et Kunth l. c. t. 467.) acaulis, foliolis 3 latis subdeltoideis emarginato-bilobis ciliatis glabris, petiolis puberulis, scapis 6-7-floris, sepalis obtusiusculis apice maculâ notatis, stylis intermediis. $\frac{1}{2}$ prope Campeche Mexican. Fl. violacei paulò minores quàm prioris. Bulbus solitarius.
70. O. GRANDIFOLIA acaulis, foliolis 3 subrotundo-obovatis profundè emarginatis ciliatis suprâ puberulis subtus pubescentibus, petiolis pilosis, scapis multifloris, sepalis obtusis apice maculâ notatis, stylis longissimis. $\frac{1}{2}$ in umbrosis humidis montis Cocollar Novæ-Andalusie. O. Macrophylla H. B. et Kunth l. c. non Horneman. — Bulbi cespitosi. Flores albi magni. O. strictæ.
71. O. CAPRINA (Lin. sp. 622.) acaulis, foliolis 3 obcordato-bilobis glabris, scapo umbellifero 2-4-floro, floribus erectis, stylis brevissimis. $\frac{1}{2}$ ad cap. B.-Spei. Flores cæruleo-incarnati fundo flavo. O. erecta Sav. in Lam. dict. 4. p. 685.
72. O. CERNUA (Thunb. diss. ox. n. 12. t. 2. f. 2.) acaulis subcaulescensve, foliolis 3 obcordato-bilobis glabris subciliatis, scapo umbellifero multifloro, floribus cernuis, stylis brevissimis. $\frac{1}{2}$ ad cap. B.-Spei. Corollæ flavæ. Stam. glabra. — Mill. ic. t. 195. f. 1. Jacq. ox. n. 16. t. 6. O. Pæ-capræ Sav. l. c. p. 685. Frequens in hortis sub nom. O. caprinæ. (v. v.)
73. O. COMPRESSA (Jacq. ox. n. 19. t. 78. f. 3.) acaulis, petiolo planiusculo, foliolis 3 obcordatis puberulis, scapo umbellifero 2-floro, sepalis integris, stylis longissimis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. lutea.
74. O. DENTATA (Jacq. ox. n. 17. t. 7.) acaulis, foliolis 3 obcordatis glabris subciliatis, scapo umbellifero 2-5-floro, sepalis apice tridentatis, stylis longissimis. $\frac{1}{2}$ ad cap. Bonæ-Spei. Cor. pallidè incarnatæ.
75. O. LIVIDA (Jacq. ox. n. 18. t. 8.) acaulis, foliolis 3 obcordato-bipartitis subtus livido-violaceis, scapo umbellifero 2-floro, stylis intermediis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. incarnatæ.
76. O. PURPURATA (Jacq. h. schæubr. t. 356.) subacaulis, foliolis 3 obcordatis ciliatis subtus sanguineis, scapo umbellifero 5-7-floro, stylis longissimis. $\frac{1}{2}$ ad cap. Bonæ-Spei. Flores albi aut subcarnei. Stam. vix connata. Petioli teretes.
77. O. MACROPHYLLA (Horn. h. hafn. 1. p. 428.) acaulis, foliolis 3 obcordatis glabris, scapo bifido foliis longiore subpiloso, floribus autè explicationem cernuis, stylis intermediis. $\frac{1}{2}$ ad cap. B.-Spei.
78. O. CORYMBOSA, acaulis, foliolis 3 glabriusculis latissimis obcordatis utrinquè subrotundis, scapis bifidis ramosis corymbosis multifloris. $\frac{1}{2}$ in ins. Borboniæ et Mauritiæ. Flores in meo specimine pleni, parvi, pallidi. Scapi 7-10-pollic. (v. s.)
- 79? O. LOBATA (Sims bot. mag. t. 2386.) acaulis glabra, pedunculis 1-floris folio longioribus, foliolis obcordatis, radice tuberosâ. $\frac{1}{2}$ in Chili. Flores flavi rubro subpunctati.
- §. 6. *Simplicifoliæ, acaules aut rariùs caulescentes, foliis simplicibus seu 1-foliolatis pubescentibus, petiolis immarginatis, scapis pluri-aut sæpiùs-1-floris.*
80. O. MANDIOCCANA (Baddil. mem. bras. p. 21.) caulescens erecta, foliis ovato-vel subcordato-acuminatis ciliatis, petiolis marginato-alatis pedunculisque florum ciliatis. $\frac{1}{2}$ in sylvis Mandioccæ montis prope Rio-Janeiro. (v. s. in h. Moric.)
81. O. ALIENA (Spreng. nene entd. 3. p. 58.) caulescens prostrata, foliis cordato-ovatis acutis villosis, pedunculis compressis alatis multifloris, staminibus

bus exterioribus stylos superantibus. — in Brasiliâ. An à priori differt? Fl. pallidi.

82. *O. PRIMULÆFOLIA* (Raddi mem. bras. add. p. 21.) acaulis, foliis oblongis obtusis, basi angustatis, scapis multifloris filiformibus semierectis prostratis — in sylvis mont. Mandioccæ prope Rio-Janeiro. Media inter Caprinas et Simplicifolias.
83. *O. MONOPHYLLA* (Lin. mant. 241.) acaulis, foliis ellipticis obtusis, scapo unifloro, filamentis glabris, stylis intermediis pilos glanduliferos gerentibus. $\frac{1}{2}$ ad cap. Bonæ-Spei. — Jacq. ox. n. 35. t. 79. f. 3. Thunb. ox. n. 1. t. 1. f. 1. Flores O. acetosellæ.
84. *O. LEPIDA* (Jacq. ox. n. 34. t. 21.) acaulis, foliis obovatis subretusis, scapo unifloro, stylis longissimis filamentisque pilos glandulosos gerentibus. $\frac{1}{2}$ ad cap. B.-Spei. Cor. albidæ.
85. *O. ROSTRATA* (Jacq. ox. n. 36. t. 22.) acaulis, foliis obovatis retusis, scapo unifloro, stylis brevissimis, filamentis pilos glandulosos gerentibus, interioribus ad apicem dorso rostratis et ex apice rostri antheriferis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. lilacino-albidæ.

§. 7. *Pteropodæ, acaules, foliis glabris bi-seu-trifoliolatis, petioliis alatis, scapis unifloris.*

86. *O. CRISPA* (Jacq. ox. n. 37. t. 23.) acaulis, foliolis 2 subrotundo-obovatis emarginatis, margine undulatis, stylis longissimis filamentisque piloso-glandulosis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. albæ magnæ. — Folium interdum 1-foliolatum.
87. *O. LEFORINA* (Jacq. ox. n. 39. t. 25.) acaulis, foliolis 2 ellipticis emarginatis margine cartilagineis denticulatis, stylis longissimis filamentisque piloso-glandulosis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. albæ margine rubicundæ.
88. *O. ASININA* (Jacq. ox. n. 38. t. 24.) acaulis, foliolis 2 lanceolatis margine cartilagineis denticulatis, stylis intermediis, filam. piloso-glandulosis. $\frac{1}{2}$ ad cap. B.-Spei. Flores flavi.
89. *O. LANCEÆFOLIA* (Jacq. ox. n. 40. t. 26.) acaulis, foliolis 2-3 margine cartilagineis scabris, stylis longissimis, filamentis glabris. $\frac{1}{2}$ ad cap. B.-Spei. Flores flavi.
89. *O. FABÆFOLIA* (Jacq. ox. n. 41. t. 27.) acaulis, foliolis 3 obovatis emarginatis mucronatis, stylis intermediis filamentisque piloso-glandulosis. $\frac{1}{2}$ ad cap. B.-Spei. Flores flavi.

§. 8. *Acetosellæ, acaules aut substipitatæ, foliis petiolatis trifoliolatis, petioliis immarginatis, foliolis subtus non glandulosis, scapis unifloris.*

* *Foliolis oblongis lanceolatisve.*

91. *O. LABURNIFOLIA* (Jacq. ox. n. 42. t. 28.) acaulis pubescens, foliolis lateralibus oblique oblongis, medio sublanceolato, scapis petiolos superantibus medio bibracteatis, stylis longissimis. $\frac{1}{2}$ ad cap. B.-Spei. Petioli rubri. Cor. flavæ. Aff. Pteropodis sed petiulus semiteres apterus.
92. *O. SANGUINEA* (Jacq. ox. n. 43. t. 29.) acaulis pubescens, foliolis oblongis obtusis, medio basi cuneato, scapis longitudine petiolorum infra medium bibracteatis, stylis intermediis. $\frac{1}{2}$ ad cap. B. Spei. Priori affinis. Petioli et pagina fol. inferior. sanguinei. Flor. flavi.
93. *O. RUBRO-FLAVA* (Jacq. ox. n. 65. t. 50.) subacaulis subhirsuta, foliolis lanceolato-oblongis obtusis, medio cuneato, scapis petiolo longioribus, infra medium bibracteatis, stylis brevissimis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. flava extus rubro marginata.
94. *O. TRICOLOR* (Jacq. ox. n. 63.) acaulis pubescens, foliolis oblongis obtusis,

medio-subcuneato, scapis petiolo longioribus, stylis intermediis, filamentis piloso-glandulosus. \mathcal{L} ad cap. B.-Spei.

α . *flava*, petalis flavis extus rubro-marginatis. — Jacq. ox. t. 47.

β . *alba*, petalis albis extus rubro-marginatis. — Jacq. ox. t. 48.

95. *O. CILIARIS* (Jacq. ox. n. 45. t. 30.) stipitata pubescens, caule inferè nudo, foliolis oblongis obtusis submarginatis, pedunculis petiolo longioribus immediatè sub calyce bibracteolatis, stylis longissimis. \mathcal{L} ad cap. B.-Spei. Cor. purpurea fundo flavo.

96. *O. ARGUATA* (Jacq. ox. n. 46. t. 31.) stipitata pubescens, caule inferè nudo decumbente, foliolis oblongis emarginatis, pedunculis petioli longitudine immediatè sub calyce bibracteolatis, stylis brevissimis. \mathcal{L} ad cap. B.-Spei. Cor. violacea.

97. *O. FLACCIDA* (Jacq. ox. n. 66. t. 51.) substipitata decumbens pubescens, foliolis oblongis retusis, medio cuneato, pedunculis folio duplò longioribus medio bibracteolatis, stylis brevissimis, filam. piloso-glandulosus. \mathcal{L} ad cap. B.-Spei. Petala alba extus rubro submarginata.

98. *O. FERRUGINATA* (Jacq. h. schœnbr. 3. t. 274.) subacaulis pubescens, foliolis obovatis retusis, pedunculis petiolo subbrevioribus medio bibracteolatis, stylis intermediis. \mathcal{L} ad cap. B.-Spei. Flores albi. Folia maculis ferrugineis picta.

99. *O. AMBIGUA* (Jacq. ox. n. 59. t. 43.) substipitata subhirsuta, foliolis obovato-oblongis obtusis, pedunculis petiolo subæqualibus medio bibracteolatis, stylis longissimis piloso-subglandulosus. \mathcal{L} ad cap. B.-Spei. Cor. alba fundo flavo, petalis extus rubro submarginatis. Sepala sub apice biglandulosa.

100. *O. UNDULATA* (Jacq. ox. n. 60. t. 44.) stipitata erecta subhirsuta, foliolis obovato-oblongis obtusis, pedunculis petiolo longioribus sub mediobi bracteolatis, stylis longissimis hirsutis. \mathcal{L} ad cap. B.-Spei. Flores præced. Sepala ovariaque glandulas plurimas gerentia.

101. *O. EXALTATA* (Jacq. ox. n. 64. t. 49.) subacaulis puberula, foliolis emarginatis, lateralibus ellipticis, medio obovato-cuneatis, pedunculis petiolo duplò longioribus medio bibracteolatis, stylis brevissimis. \mathcal{L} ad cap. B.-Spei. Petala alba extus rubro marginata.

102. *O. GLANDULOSA* (Jacq. ox. n. 62. t. 46.) substipitata puberula, foliolis oblongis obtusis, medio subcuneato, pedunculis petioli longitudine medio biglandulosus, stylis intermediis, filamentis pedunculis calycibusque piloso-glandulosus. \mathcal{L} ad cap. B.-Spei. Cor. albæ fundo flavo.

103. *O. FUSCATA* (Jacq. ox. n. 61. t. 45.) subacaulis puberula, foliolis obtusis lateralibus ovatis, medio cuneiformi, pedunculis petiolo duplò longioribus medio bibracteolatis, stylis longissimis piloso-subglandulosus. \mathcal{L} ad cap. B. Spei. Folia subtus sanguinea. Cor. albæ fundo flavo.

* * *Foliolis subrotundis subrhombeisve.*

104. *O. TRUNCATULA* (Jacq. ox. n. 76. t. 62.) acaulis hirsuta, foliolis triangularibus truncatis, scapis petiolo triplò longioribus medio bibracteolatis, stylis longissimis. \mathcal{L} ad cap. B.-Spei. Cor. lilacina.

105. *O. SUBRHUREA* (Jacq. ox. n. 77. t. 63.) acaulis puberula, foliolis subrotundis, pedunculis petioli longitudine ad basin bibracteolatis, stylis brevissimis, calycibus margine clavato-pilosis apice reflexis. \mathcal{L} Cor. pallidè flavæ. Folia subtus sanguinea.

106. *O. BREVISCAPA* (Jacq. ox. n. 72. t. 58.) acaulis puberula, foliolis subrotundis, petiolis planiusculis, pediculis petiolo duplò brevioribus infrà medium bibracteolatis, stylis intermediis. \mathcal{L} ad cap. B.-Spei. Cor. albæ fundo flavo. Folia subtus sanguineo-punctata.

107. *O. SPECIOSA* (Willd. sp. 2. p. 779.) acaulis puberula, foliolis subrotundis, scapis petiolo subæqualibus infrà medium bibracteolatis, stylis longissimis

sepalis pilos simplices et glandulosos mixtos gerentibus. \mathcal{U} ad cap. B.-Spei.

a. purpurea (Lin. sp. 621.) foliis subtus sanguineis, corollis roseo-purpureis. O. speciosa Jacq. ox. n. 74. t. 60.

ß. sugillata (Jacq. ox. n. 75 t. 61.) foliis adultis subtus totis et suprâ ad oras lividis, corollis albis extus subincarnatis.

γ. rigidula (Jacq. ox. n. 73. t. 59.) foliis utrinquë viridibus, corollis albis fundo flavido.

108. O. VARIABILIS (Jacq. ox. n. 67.) acaulis puberula, foliolis subrotundis, medio basi cuneato, pediculis folio æqualibus longioribusve infrâ medium bibracteolatis, stylis brevissimis, filamentis piloso glandulosis. \mathcal{U} ad cap. B.-Spei.

a. longiscapa, scapis folio duplò longioribus, cor. albis subincarnatis. — Jacq. t. 52.

ß. rubra (Jacq. t. 53.) scapis folio paulò longioribus, cor. roseo-purpureis.

γ. graciliflora (Jacq. n. 68. t. 54.) scapis petioli longitudine, cor. amplis albis, foliis subtus sanguineis.

δ. Simsii, scapis petioli longitudine, cor. amplis albis, foliis utrinquë viridibus. bot. mag. t. 1683. An cum var. γ . spec. propr. constituit?

109. O. PURPUREA (Willd. sp. 2. p. 778.) acaulis puberula, foliolis subrotundis, scapis folio longioribus infra medium bibracteolatis, stylis intermediis, filam. piloso-glandulosis. \mathcal{U} ad cap. B.-Spei. — Jacq. ox. n. 70. t. 56. Cor. purpureæ.

ß. laxula (Jacq. ox. n. 71. t. 57.) scapis petioli longitudine, cor. albis fundo flavido.

110. O. CONVEXULA (Jacq. ox. n. 69. t. 55.) glabra, stipite declinato infernè nudo, foliolis subrotundis punctatis, stipulis dilatatis acuminatis, pedunculis folio longioribus, bracteolis alternis, stylis intermediis, filam. piloso-glandulosis. \mathcal{U} ad cap. B.-Spei. Cor. roseæ fundo flavo.

111. O. HUMILIS (Thunb. prod. app. 190.) acaulis, foliolis subrotundis ciliatis, scapis unifloris. \mathcal{U} ad cap. B.-Spei. Bulbus ovatus. Petioli pubescentes prostrati. Scapi foliis paulò longiores, medio bibracteolati. Cor. tubus ampliatus flavescens, limbus purpureus. Thunb. fl. cap. ed. Schult. 2. p. 535, à quo eadem ac O. purpurea Lin. dicitur. †.

112. O. LÆVIGATA (Willd. enum. suppl. 26.) subacaulis, glaberrima, foliolis subrotundis, intermedio cuneato, scapis longitudine ferè petiolorum, stylis brevissimis. ① Patr. non indic. Cor. purpureæ. Link enum. 439.

*** *Foliolis obcordatis.*

113. O. PUNCTATA (Lin. f. suppl. 243.) acaulis, foliolis subrotundo-obcordatis punctatis glabris, petiolis scapisque puberulis, scapo foliis longiore suprâ medium bibracteolato, stylis brevissimis, filam. piloso-glandulosis, \mathcal{U} ad cap. B.-Spei. Petala pallidè incarnata, extus rubro marginata. — Jacq. ox. n. 82. t. 66.

114. O. STRUMOSA (Jacq. ox. n. 79. t. 64.) acaulis puberula, foliolis obcordatis, pediculis foliorum longitudine infrâ medium bibracteolatis, stylis brevissimis subapice strumosis, filam. piloso-glandulosis. \mathcal{U} ad cap. B.-Spei. Petala alba ungue flavo extus rubro-marginata. Fol. subt. sanguinea, sup. fusco maculata.

115. O. MARGINATA (Jacq. ox. n. 85. t. 68.) acaulis puberula, foliolis obcordatis subrotundis, scapis petiolo dimidiò ferè brevioribus medio bibracteolatis, calycibus clavato ciliatis apice reflexis, stylis intermediis. \mathcal{U} ad cap. B.-Spei. Flores albi. Folia subtus sanguinea viridi marginata.

116. O. PULCHELLA (Jacq. ox. n. 86. t. 69.) acaulis puberula, foliolis obcordatis subrotundis, scapis petiolo triplò brevioribus medio bibracteolatis, calycibus clavato-ciliatis apice reflexis, stylis longissimis. \mathcal{U} ad cap. B.-Spei. Flores albi subincarnati. Folia subtus sanguinea.

117. O. OBTUSA (Jacq. ox. n. 83. t. 79. f. 1.) acaulis dense puberula, foliolis obcordatis, scapo foliis longiore supra medium bibracteolato, calycibus obtusis, stylis intermediis. \mathcal{U} ad cap. B.-Spei. Cor. rubra. Filamenta glabra.

118. *O. LANATA* (Lin. f. suppl. 244. excl. syn.) acaulis villosa-lanata, foliolis obcordatis, scapo foliis longiore medio bibracteolato, calycibus acutis, stylis brevissimis. \mathcal{L} ad cap. B.-Spei. Jacq. ox. n. 81. t. 77. f. 2. Cor. alba. Filam. hirta.
119. *O. LUTEOLA* (Jacq. ox. n. 80. t. 65.) acaulis pilosa, foliolis obcordatis subretusis, scapis foliorum longitudine vix supra medium bibracteolatis, stylis brevissimis, filam. interioribus piloso-glandulosis. \mathcal{L} ad cap. B.-Spei. Flores flavi.
120. *O. FALLAX* (Jacq. ox. n. 84. t. 67.) acaulis pilosa, foliolis obcordatis subretusis, scapis foliis longioribus paulò supra medium bibracteolatis, stylis intermediis stam. interiora piloso-glandulosa subæquantibus. \mathcal{L} ad cap. B.-Spei. Flores flavi. Priori valdè affinis.
121. *O. MACROGONYA* (Jacq. ox. n. 87. t. 70.) subacaulis, pilosiuscula, foliolis obcordato-subrotundis, scapo foliis brevior paulò supra medium bibracteolato, stylis longissimis filamentisque piloso-glandulosis. \mathcal{L} ad cap. B.-Spei. Flores flavi. Sepala apice subtrès biglandulosa.
122. *O. MAGELLANICA* (Forst. comm. goett. 9. p. 33.) acaulis, foliolis obcordato-subrotundis carnosis lævibus, scapis pilosis apice bibracteolatis. \mathcal{L} in terrâ del Fuego locis irriguis. Flores albi *O. acetosellæ*.
123. *O. ACETOSELLA* (Lin. sp. 620.) acaulis, radice dentatâ repente, foliolis obcordatis puberulis, scapo foliis longiore supra medium bibracteolato, petalis ovalibus obtusis, stylis stam. interiora æquantibus aut superantibus. \mathcal{L} in Europæ umbrosis. — Fl. dan. t. 980. Jacq. ox. n. 91. t. 80. f. 1. Flores albi. (v. v.)
 β . *cærulea*, flore cærulescente. Tourn. inst. 88. (v. s.)
 γ . *subpurpurascens*, flore pallidè roseo aut purpurascente. (v. s.)
 δ . *parviflora* (Lejeun. spa. 2. p. 307.) floribus albis quadruplò minoribus pentandris — in sepibus circa Malmèdy.
124. *O. AMERICANA* (Bigl. in litt.) acaulis, radice dentatâ repente, foliolis obcordatis puberulis, scapo foliis longiore supra medium bibracteolato, petalis oblongis inæqualiter emarginatis, stylis stam. interiora vix superantibus. \mathcal{L} in Amer. bor. umbrosis. *O. acetosella* Michx. Pursh. Nutt. Flores albi. (v. s.)
125. *O. TINELLA* (Jacq. ox. n. 32. t. 19) substipitata glabriuscula, foliolis obcordatis, scapo foliis longiore supra medium bibracteolato, stylis brevissimis, filam. piloso-glandulosis. \mathcal{L} ad cap. B.-Spei. Cor. pallidè lilacinæ. Sepala apice biglandulosa.
126. *O. NATANS* (Lin. f. suppl. 243.) caule infernè nudo declinato, foliolis obcordatis glabris, pedunculis foliorum longitudine infrâ medium bibracteolatis, stylis brevissimis. \mathcal{L} in aquis ad. cap. B.-Spei. Thunb. ox. n. 4. t. 1. f. 4. Jacq. ox. n. 78. t. 76. f. 2. Cor. alba.
127. *O. LONGIFLORA* (Lin. sp. 621.) caule infernè nudo simplici, foliolis obcordato-bilobis, pedunculo unifloro. — in Virginiâ. \uparrow .
128. *O. CUENTATA* (Jacq. f. ecl. 1. p. 66. t. 45.) substipitata subhirsuta, foliolis obcordatis, pedunculis 1-floris folia paulò superantibus supra medium bibracteolatis, stylis intermediis. \mathcal{L} in Africâ? Flores purpurei. Filam. glabra, exteriora dentata.
129. *O. FILICAULIS* (Jacq. h. schænbr. 2. t. 205.) caule infernè nudo decumbente subramoso, foliolis obcordato-bilobis glabris, pedunculis folio longioribus medio bibracteolatis, stylis intermediis. \mathcal{L} ad cap. B.-Spei. Flores violacei. Filam. glabra, ext. dentata.
130. *O. BIFIDA* (Thunb. ox. n. 16. t. 1. f. 2.) caule infernè nudo erecto subramoso, foliolis obcordato-bilobis glabris, pedunculo foliis longiore paulò supra medium bibracteolato, stylis longissimis. \mathcal{L} ad cap. B.-Spei. — Jacq. ox. n. 89. t. 79. f. 4. Corolla violacea fundo flavo.

**** *Foliolis linearibus subcuneatis apice emarginatis aut retusis.*

131. *O. CUNEATA* (Jacq. ox. n. 55. t. 40.) caule declinato infernè nudo, foliis cuneatis emarginatis pilosis, pedicellis petioli longitudine ad apicem bibrac-

teolatis, stylis longissimis, filam. apice piloso-glandulosis. $\frac{1}{2}$ ad cap. B.-Spei. Flores albi fundo flavo.

132. O. CUNEIFOLIA (Jacq. ox. n. 56. t. 41.) caule declinato infernè nudo, foliis cuneatis emarginatis pilosis, pedicellis petioli longitudine ad apicem bibracteolatis, stylis brevissimis, filam. piloso-glandulosis. $\frac{1}{2}$ Flores albi fundo vix flavo. Priori valdè affinis.

133. O. PUSILLA (Jacq. ox. n. 57. t. 42.) caule brevi infernè nudo subdeclinato, foliolis linearibus emarginatis glabris, pedicellis petioli longitudine ad apicem bibracteolatis, stylis intermediis, filam. glabris. $\frac{1}{2}$ ad cap. B.-Spei. Planta vix digitalis. Flores albi subincarnati.

134. O. LINEARIS (Jacq. ox. n. 47. t. 32.) caule declinato infernè nudo folisque puberulis, foliolis linearibus emarginatis, pedicellis petiolo brevioribus apice summo bibracteolatis, stylis longissimis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. violacea fundo flavo. Filam. glabra edentula.

135. O. RECLINATA (Jacq. ox. n. 49. t. 34.) caule reclinato infernè nudo subramoso, foliolis linearibus subcuneatis emarginatis, pedicellis petiolo longioribus supra medium bibracteolatis petiolisque puberulis, stylis intermediis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. pallidè miniata. Filam. hirsutula.

136. O. GRACILIS (Jacq. ox. n. 48. t. 33.) caule declinato infernè nudo folisque glabris, foliolis linearibus apice subcallosis, pedicellis petiolo longioribus supra medium bibracteolatis, stylis longissimis filamentisque piloso-glandulosis. $\frac{1}{2}$ ad cap. B.-Spei. Cor. pallidè miniatæ. An ad sect. seq. referenda? O. versicolor γ . Willd. sp. 2. p. 792.

137. O. MINIATA (Jacq. ox. n. 50. t. 35.) caule declinato infernè nudo folisque glabris, foliolis linearibus apice subemarginatis, pediculis folio sublongioribus supra medium bibracteolatis, stylis brevissimis, filam. glabris. $\frac{1}{2}$ ad cap. B.-Spei. Flores miniati. Filam. edentula. O. elongata β Willd. sp. 2. p. 793.

§. 9. Adenophyllæ, caulibus nunc sparsè foliosis nunc apice tantum foliosis nunc subnullis, foliis petiolatis 3-5-foliolatis, foliolis linearibus apice subtus callos glandulosos gerentibus, pedunculis 1-floris.

138. O. MINUTA (Thunb. diss. n. 2. t. 2. f. 3.) acaulis, foliolis 3 linearilanceolatis acutis, scapo foliis longiore, stylis longissimis. ① ad cap. B.-Spei. — Jacq. ox. n. 44. t. 79. f. 2. An foliola glandulosa? An hujus sect.?

139. O. GLABRA (Thunb. ox. n. 17. t. 2. f. 1.) caule brevissimo infernè nudo erecto, foliolis 3 linearilanceiformibus emarginatis ciliatis subtus multiglandulosus, pedunculis folio longioribus, stylis longissimis. $\frac{1}{2}$ ad cap. Bonæ-Spei. — Jacq. ox. n. 58. t. 76. f. 3. Cor. purpureæ fundo flavo.

140. O. VERSICOLOR (Lin. sp. 622.) caule infernè nudo declinato, foliolis 3 linearibus emarginatis subtus biglandulosus, pedunculis folio longioribus, stylis longissimis filamentisque piloso-glandulosus. $\frac{1}{2}$ ad cap. B.-Spei. — Jacq. ox. n. 51. t. 36. Cor. intus albæ, extus coccineo-rubræ. (v. s.)

β . sylvestris, pedunculis folii longitudine, stylis hirsutis, filam. edentulis glabris. $\frac{1}{2}$ ad cap. Bonæ-Spei. — Jacq. ox. t. 77. f. 4. An species dist?

141. O. ELONGATA (Jacq. ox. n. 52. t. 37.) caule infernè nudo declinato, foliolis 3 linearibus emarginatis apice bicalliosis, pedunculis folio longioribus, stylis brevissimis. $\frac{1}{2}$ ad cap. B.-Spei. Petala alba vix margine flavo-purpurea, apice subemarginata.

β . amœna (Jacq. h. schœnbr. t. 206.) calycibus obtusis, floribus roseo-purpureis, petalis obtusis. (v. v.)

142. O. TENUIFOLIA (Jacq. ox. n. 53. t. 38.) caule erecto sparsè folioso, foliolis 3 linearibus emarginatis subtus pluriglandulosus, pedunculis folio longioribus, stylis brevissimis, filam. interioribus piloso-glandulosus. $\frac{1}{2}$ ad cap. B.-Spei. Lodd. bot. cab. t. 712. Cor. albæ margine rubentes.

143. O. POLYPHYLLA (Jacq. ox. n. 59. t. 39.) caule erecto subramoso apice folioso, foliolis 3 linearibus emarginatis subtus biglandulosus, pedunculis folio longioribus, stylis intermediis filamentisque piloso-glandulosus. $\frac{1}{2}$ ad cap. Bonæ-Spei. Cor. pallidè purpurascens. O. versicolor Jacq. ic. rar. t. 473. — Burm. afr. t. 27. f. 1.

144. *O. FILIFOLIA* (Jacq. h. schœnbr. 3. t. 273.) caule declinato infernè nudo, foliolis 3 linearibus apice integris calloso-glandulosi, pedunculis folio longioribus, stylis longissimis filamentisque interioribus piloso-glandulosi. $\frac{1}{2}$ ad cap. B.-Spei. Cor. roseo-purpurascens.

145. *O. PENTAPHYLLA* (Sims bot. mag. t. 1549.) caule suberecto infernè nudo, foliolis quinque linearibus apice subintegris 1-2-callosis-glandulosi, pedunculis folio longioribus, stylis intermediis. $\frac{1}{2}$ ad cap. B.-Spei. *O. digitata* Poir. suppl. 4. p. 254. excl. syn. et var. β . Flores roseo-incarnati. Petioli basi rubentes amplexicaules. (v. v.)

§. 10. *Palmatifoliæ, acaules aut stipite brevi basi nudo, foliis petiolatis palmatim peltatimve 5-13-foliolatis, foliolis eglandulosis, scapis unifloris.*

146. *O. MALLOROLBA* (Cav. ic. 4. p. 64. t. 393. f. 2.) acaulis, bulbis lanatis, foliolis 5 ovatis, scapis medio bracteolas 2 subulatas gerentibus. — in Americâ merid. Planta pusilla lanâ bulbi rufâ insignis.

147? *O. COMMENSONI* (Pers. ench. 1. p. 519.) caulibus brevissimis foliosis, foliolis 6 ovatis adpressè villosis, scapis folio longioribus medio bibracteolatis. — in Monte-Video. *O. sexenata* Sav. in Lam. dict. 4. p. 687. Flores flavi.

148. *O. LUPINIFOLIA* (Jacq. ox. n. 92. t. 72.) subacaulis, foliolis 7 lanceolatis acutiusculis glabris basi maculatis, petioli compressis, scapis petioli longitudine, stylis brevissimis, filam. piloso-glandulosi. $\frac{1}{2}$ ad cap. Bonæ-Spei. (v. v.)

149. *O. FLAVA* (Lin. sp. 621.) caule erecto brevi basi nudo, foliolis 6-7 glabris linearibus canaliculatis acutis, pedunculis petiolo sublongioribus, stylis brevissimis, filam. piloso-glandulosi. $\frac{1}{2}$ ad cap. B.-Spei. Jacq. ox. n. 93. t. 73. Burm. afr. t. 27. f. 4. Flores flavi.

150. *O. PECTINATA* (Jacq. ox. n. 95. t. 75.) subacaulis, foliolis 7 glabris lineari-lanceolatis obtusis, pedunculis folii longitudine, calycibus adpressis, stylis longissimis filamentisque piloso-glandulosi. $\frac{1}{2}$ ad cap. B.-Spei. — Burm. afr. t. 30. f. 1?

151. *O. FLABELLIFOLIA* (Jacq. ox. n. 94. t. 74.) subacaulis, foliolis 7-9 glabris linearibus emarginatis, pedunculis petiolo vix longioribus, calycibus apice reflexis, stylis intermediis filamentisque interioribus piloso-glandulosi. $\frac{1}{2}$ ad cap. B.-Spei. Cor. flavæ. (v. v.)

152. *O. TOMENTOSA* (Lin. f. suppl. 244.) subacaulis, foliolis 9-19 utrinque pubescenti-tomentosis lanceolato-cuneatis emarginatis, scapis petioli longitudine, stylis longissimis. $\frac{1}{2}$ ad cap. B.-Spei. — Jacq. ox. n. 96. t. 81. — Plum. t. 350. f. 3. Cor. albidæ.

153. *O. ENNEAPHYLLA* (Cav. ic. 5. p. 7. t. 411.) stipite basi repente interruptè squamoso, foliolis 9 glabris obcordato-cuneatis ferè bifidis, pedunculis longitudine foliorum, stylis longissimis. $\frac{1}{2}$ in ins. Maclovianis et Falklandianis. Cor. roseæ lineis lutescentibus.

154. *O. LACINIATA* (Cav. ic. 5. p. 7. t. 412.) stipite basi repente continuo-squamoso, foliolis 11-13 linearibus acutis glabris, scapis petiolo brevioribus, stylis longissimis. $\frac{1}{2}$ in Americâ merid. Cor. violacæ. Hæc et prior sectionem propriam ferè constituunt ob stipitem repentem squamosum et stigmata capitata.

IV. LEDOCARPUM. Desf. mem. mus. 4. p. 250.

Cal. persistens 5-partitus, bracteis 2-3-partitis cinctus. Pet. 5. Stam. 10 alterna breviora, filam. liberis persistentibus. Ovarium subrotundum. Styli 5 incrassati. Capsula 5-valvis, valvis introflexis, 5-locul. ∞ -sperma. — Frutex ramosus, foliis oppositis 3-partitis lineari-subulatis sericeis, floribus solitariis terminalibus.

1. *L. CHILOENSE* (Desf. l. c. t. 13.) $\text{\text{♂}}$ in Chili. *Balbisia verticillata* Cav. ic. ined. in bibl. Balb. (v. s.)

Ordo L. ZYGOPHYLLÆ. *Brown gen. rem. p. 13.*

Calyx sepalis 5 distinctis aut vix basi subconnexis. Petala 5, sepalis alterna, receptaculo inserta. Stam. 10 distincta, hypogyna, 5 sepalis, 5 petalis opposita. Ovarium unicum 5-locul. Stylus 5 in 1 coaliti, interdum apice summo subdistincti. Capsula carpellis 5 plus minus inter se et axi centrali adnatis constans, loculis ad angulum superius dehiscentibus, sæpius polyspermis, aut 1-spermis, nec cocculiferis nec arilliferis. Semina albuminosa aut sæpius exalbuminosa, embryone recto, radicula superâ, cotyledonibus foliaceis.

Herbæ frutices arboresve habitu vario, foliis basi stipulaceis, sæpius compositis, in Zygophylleis veris oppositis, alternis autem in Zygophylleis spuriis, hîc forsân laxiore nexu alligatis.

Ordo medius inter Oxalideas et Rutaceas; à prioribus distinguitur stylo unico nec multiplici, seminibus exarillatis, et foliis constantius stipulaceis et oppositis; à posterioribus differt carpellorum structurâ nempè absentîâ cocculorum elasticorum in Rutaceis veris adeò notabilium. Ab iis distinguitur etiam primâ fronte ob stipulas ad basin folii cujusque binas.

§. 1. ZYGOPHYLLÆ VERÆ OPPOSITIFOLIÆ.

I. TRIBULUS. *Tourn. inst. t. 141. Lin. gen. n. 532.*

Calyx 5-sepalus deciduus. Pet. 5 patentia. Stam. 10. Stylus o. Carpella 5 axi adnexa triangularia indehiscencia dura, extus in spinas tubercula aut alas exserta, intus transversè plurilocularia, rariùs 1-locularia. Semina in loculis solitaria horizontalia exalbuminosa. Cotyl. crassiusculæ. (Gaertn. fr. 1. t. 69.) — Herbæ diffusæ. Folia abruptè pinnata. Stipulæ membranacæ. Pedicelli axillares 1-flori solitarii. Flores sæpius flavi.

1. *T. CISTOIDES* (Lin. sp. 554.) foliis 8-jugis, foliolis subæqualibus subtus sericeis, pedicellis petioli longitudine. 2 in Americâ austr. — Jacq. schœnbr. t. 103. — Pluk. t. 67. f. 4. Capsula (ex Kunth) 5-cocca, coccis transversè 3-4-loc. (v. v.)
2. *T. ALBUS* (Poir. dict. 8. p. 45.) foliis sub-8-jugis, foliolis subæqualibus utrinquè villosis, pedicellis petiolo brevioribus. 3 in Senegaliâ. Flores albi, petalis vix calyce majoribus.
3. *T. TERRESTRIS* (Lin. sp. 554.) foliis 6-jugis, foliolis subæqualibus, pedicellis petiolo brevioribus, carpellis 4-cornutis. (1) in arenosis Europæ australis, Barbariæ, Senegaliæ, ins. Mauritiî. Variat foliolis villosis aut ferè glabris. — Lam. ill. t. 346. f. 1. Schkuhr handb. 1. t. 115. Capsula 5-cocca, coccis transversè 2-3-loc. (v. v.)

℞. *subinermis* (Fisch. cat. gor. 1808. p. 94.) hirsutior, carpellis subinermibus tuberculatis. — in Thibeto. An species?

4. *T. ALATUS* (Delil. ill. p. 44.) foliis 5-6-jugis, foliolis subæqualibus utrinquè adpressè villosis, pedicellis brevissimis, carpellis ecornutis margine alatis.

— in *Ægypto*. T. pentandrus Forsk. descr. 38? qui dicit flores 5-6-andros. Capsula 5-coeca, coccis transv. 2-loc. (v. s. sine fl.)

5. T. *LAXUGINOSUS* (Lin. sp. 553.) foliis 5-6-jugis, foliolis subæqualibus adpressè pubescenti-villosis, pedicellis folio brevioribus, carpellis bicornutis. — in *Ceylonâ*. — Burm. zeyl. 265. t. 106. f. 1. (v. s.)
6. T. *MAXIMUS* (Lin. sp. 553.) foliis 3-4-jugis, foliolis exterioribus majoribus, pedicellis folio brevioribus, carpellis inermibus in fructum 10-costatum 10-spermum connexis. ① in *Jamaicæ* et *Saucti-Thomasii* aridis. — Jacq. ic. rar. t. 462. Lam. ill. t. 346. f. 2. Capsula 10-coeca, coccis 1-loc. (v. s.)
7. T. *TRIUGATUS* (Nutt. gen. am. 1. p. 277.) foliis 3-jugis, foliolis ultimis majoribus, pedicellis....., carpellis cristato-muricatis 1-spermis. ① in cultis *Georgiæ* circa *Savannah*.

II. FAGONIA. Tourn. inst. t. 141. Lin. gen. n. 531.

Cal. 5-sepalus deciduus. Pet. 5, unguiculata. Stam. 10. Stigma 1. Capsula subrotunda, 5-angul., 5-locul., acuminata; loculis bivalvibus 1-spermis. Embryo rectus in axi albuminis carnosus. (Gaertn. fr. 2. p. 153. t. 113.) — *Herbæ*, stipulis sæpius geminis spinosis, pedicellis axillaribus solitariis 1-floris.

* Foliis trifoliolatis.

1. F. *CRETICA* (Lin. sp. 553.) foliis 3-foliolatis, foliolis lineari-lanceolatis planis glabris, stipulis spinosis, ovarii glabris. ① in montosis apricis *Crætæ*, *Mauritanicæ*, *Hispanicæ*. Lam. ill. t. 346. Caulis procumbens rarius erectus. F. erecta Mill. dict. n. 1. (v. v.)
2. F. *HISPANICA* (Lin. sp. 553.) foliis 3-foliolatis, stipulis non spinosis. ② in *Hispaniâ*. Planta dubia! An prioris varietas? Mill. dict. n. 2.
3. F. *ARABICA* (Lin. sp. 553.) foliis 3-foliolatis, foliolis linearibus mucronatis subtus convexis, stipulis spinosis, ovarii villosis. ③ in *Barbariâ*, *Ægypto*, *Arabiâ*. Spinæ longæ. Caulis ramosissimus. (v. s.)
4. F. *GLUTINOSA* (Delil! fl. eg. 86. t. 28. f. 2.) foliis 3-foliolatis, foliolis obovatis mucronatis, stipulis spinosis petiolo brevioribus, caulibus prostratis dichotomis fructibusque viscosis. — in desertis *Kahirinis*. F. scabra Forsk. descr. 88? (v. s. in h. Balbis.)
5. F. *MOLLIS* (Delil! fl. eg. 76. t. 27. f. 2.) foliis 3-foliolatis, foliolis ovalibus mucronatis villosis, stipulis spinosis petioli longitudine, caulibus fruticosis diffusis apice villosis, fructibus hispidis. ④ in desertis *Kahirinis*. (v. s.)
6. F. *LATIFOLIA* (Delil! fl. eg. 86. t. 28. f. 3.) foliis 3-foliolatis, foliolis lateralibus lanceolatis acutis, medio majore obovato-orbiculato, stipulis spinosis minimis. ① in *Ægypto* prope *Kahirum*. (v. s.)

** Foliis simplicibus.

7. F. *MYSOURENSIS* (Roth nov. sp. 215.) foliis simplicibus linearibus cartilagineo-cuspidatis, stipulis spinosis subulatis asperis folio longioribus pedicellis folio duplò brevioribus. — in *Mysore*.
8. F. *PERSICA*, foliis simplicibus ovalibus, stipulis spinosis subulatis folio brevioribus, pedicellis folio sublongioribus. ① in *Persiâ*. F. *Indica* Burm. fl. ind. p. 102. t. 34. f. 1.
9. F. *OLIVERII*, foliis simplicibus oblongo-linearibus, stipulis spinosis vix folii longitudine, pedicellis brevissimis, fructibus pubescentibus. ③ inter *Bagdad* et *Alep*. Olivier et Brug. Habitus ferè F. *Persicæ*, sed spinæ longiores et pedicelli brevissimi. (v. s.)
10. F. *BRUGUIERII*, foliis simplicibus oblongis cuspidatis, stipulis spinosis folio longioribus, ramis tetragonis, pedicellis brevissimis, fructibus pubescentibus. ④ inter *Bagdad* et *Alep*. Bruguère et Oliv. Radix lignosa. Caules à basi ramosi vix digitales. (v. s.)

III. LARREA. Cav. ic. 6. p. 39.

Cal. 5-sepalus. Pet. 5 unguiculata. Stam. 10, filam. intus ad basim appendice ovarium tegente aucta. Antheræ cordatæ. Styl. 1 pentagonus. Capsula carpellis quinque 1-loc. coalitis constans. Semina in loculis abortu solitaria, albuminosa. Embryo rectus. — Frutices floribus solitariis pedicellatis flavis. Genus vix à Zygophyllo differt, et species duæ posteriores forsan ad Zyg. removendæ.

1. L. NITIDA (Cav. ic. t. 559.) foliis impari-pinnatis glabris glutinosis, foliolis 5-7-jugis approximatis linearibus. ♂ in Bonariâ prope Mendoza. Flores flavi. Ovarium villosum. (v. s.)
2. L. DIVARICATA (Cav. ic. 6. t. 560. f. 1.) foliis sessilibus bilobis villosis, lobis profundis lanceolatis divaricatis. ♂ hab. cum priore : Ovarium et fructus pilis longis obsiti.
3. L. CUNEIFOLIA (Cav. ic. 6. t. 560. f. 2.) foliis subsessilibus villosis cuneatis apice bilobis, setulâ inter lobulos proeminente. ♀ cum prioribus.

IV. ZYGOPHYLLUM. Lin. gen. n. 530. — Fabago. Tourn. inst. t. 230.

Cal. 5-sepalus. Pet. 5. Stam. 10 filamentis intus ad basin appendice squamosâ auctis. Discus glandulosus nectarifer hypogynus. Stylus et stigma 1. Capsula oblonga 5-gona, 5-locul., 5-valvis, valvis demum separabilibus. Semina in quoque loculo plurima biserialia, angulo interno adfixa, exalbuminosa. (Gaertn. fr. 2. t. 112.) Herbæ aut Frutices. Folia sæpius carnosa, stipulis membranaceis stipata. Pedicelli axillares solitarii 1-flori.

* *Foliis simplicibus.*

1. Z. SIMPLEX (Lin. mant. 68.) foliis simplicibus sessilibus cylindricis. ① in desertis Egypti et Arabiæ frequens. Z. portulacoides Forsk. descr. 88. ic. t. 12. B. Flores flavi. Fructus oligospermi. (v. s.)
2. Z. CORDIFOLIUM (Lin. f. suppl. 232.) foliis simplicibus sessilibus subrotundis. ♀ ad cap. B.-Spei. Ait. h. kew. 2. p. 60.

** *Foliis bifoliolatis, foliolis planis glabris.*

3. Z. FABAGO (Lin. sp. 551.) foliis bifoliolatis petiolatis, foliolis obovatis, pedicellis erectis, calycibus glabris, petalis indivisis. ♀ in Syriâ, Tauriâ, Mauritanîâ. Lam. ill. t. 345. f. 1. Petala alba basi croceo-lateritia. (v. v.)
4. Z. FOETIDUM (Schr. et Vendl. sert. han. p. 17. t. 9.) foliis bifoliolatis petiolatis, foliolis obovatis, floribus nutantibus, petalis reflexis incisis. ♂ ad cap. B.-Spei. Petala aurantiaco-flava, basi maculâ purpureâ.
 - β. insuave (Sims. bot. mag. t. 372.) petalis oblongis basi maculâ fuscâ cordatâ, pedicellis fructiferis deflexis, petiolis apice setiferis.
 - γ. retrofractum (Thunb. prodr. 80.) foliolis ovatis, ramis retrofractis. — An plures hic confusæ species?
5. Z. DECUMBENS (Delil. fl. eg. 77. t. 27. f. 3.) foliis bifoliolatis petiolatis, foliolis oblongo-obovatis carnosissimis planis, pedicellis erectis, capsulis turbinatis sphaericis apice umbilicatis. ♀ in Egypto. Caules decumbentes basi perennantes. Rami fructiferi ob foliorum delapsum dichotomi videntur.
6. Z. BILLARDIERII, foliis bifoliolatis petiolatis, foliolis oblongis carnosissimis planis, capsulis turbinatis, caule herbaceo decumbente. — in Novæ-Hollandiæ terrâ Van-Leuwin. Flores ignoti. Fructus distinctissimi. Calyces elongati persistentes reflexi. (v. s.)
7. Z. FRUTICULOSUM, foliis bifoliolatis petiolatis, foliolis oblongis carnosissimis planis, pedicellis brevissimis, petali unguiculatis, staminibus exappendiculatis, caule fruticoso erecto. ♂ in Novæ-Hollandiæ littore orientali. Fructus ignoti. Flores parvi, flavi. (v. s.)

8. *Z. MACULATUM* (Ait. h. kew. 2. p. 60.) foliis bifoliolatis petiolatis, foliolis lineari-lanceolatis. $\frac{3}{5}$ ad cap. B.-Spei. Petala lutea, 3 super. maculâ cordatâ rubrâ ad basin.
9. *Z. MICROPHYLLUM* (Thunb. prodr. 80.) foliis bifoliolatis subsessilibus, foliolis ovatis lævibus minutis, capsulis retusob-emarginatis. $\frac{3}{5}$ ad cap. B.-Spei. Pedunculi capillares. Petala flava.
10. *Z. MORGSANA* (Lin. sp. 551.) foliis bifoliolatis breviter petiolatis, foliolis obovatis petiolis in setas spiniformes persistentes desinentibus, capsulis subinflatis 4-5-pteris. $\frac{3}{5}$ ad cap. B.-Spei. Flores nutantes flavi. Burm. afr. 7. t. 3. f. 2. — Dill. elth. t. 116. f. 141. (v. s.)
11. *Z. SESSILIFOLIUM* (Lin. sp. 552.) foliis bifoliolatis subsessilibus, foliolis lanceolato-ovalibus, petiolis in setas persistentes spiniformes desinentibus, capsulis globoso-depressis apteris. $\frac{3}{5}$ ad cap. B.-Spei. — Dill. h. elth. t. 116. f. 142. Burm. afr. 4. t. 2. f. 1. Flores prioris. Rami in specimin. quibusdam valde angulati, in aliis teretes. An duæ species confusæ? (v. s.)
12. *Z. SPINOSUM* (Lin. mant. 380.) foliis bifoliolatis subsessilibus, foliolis linearibus carnosius planiusculis, petiolis in setas spiniformes persistentes desinentibus. $\frac{3}{5}$ ad cap. B.-Spei. Flores et habitus duorum priorum. — Burm. afr. 5. t. 2. f. 2 exhibet plantam à nostrâ parum diversam, sed spinulis sparsis secus caulem donatam. In nostrâ spinæ sunt tantum reliquæ petiolorum. (v. s.)
13. *Z. CAPENSE* (Lam. dict. 2. p. 443.) foliis bifoliolatis sessilibus, foliis obovatis subrotundis carnosius. $\frac{3}{5}$ ad cap. B.-Spei. Flores rubri.
14. *Z. ÆSTUANS* (Lin. sp. 552.) foliis bifoliolatis subsessilibus, foliolis obovatis retusis, petiolis in setulas desinentibus, stipulis ex aliâ parte caulis binis, ex alterâ solitariis. — Surinami. Flores et fructus ignoti.

*** *Foliis bifoliolatis, foliolis teretibus.*

15. *Z. COCCINEUM* (Lin. sp. 551.) foliis bifoliolatis petiolatis, foliolis cylindricis carnosius lævibus, pedicellis erectis, petalis acuminatis, capsulis cylindraceis. $\frac{3}{5}$ in desertis siccis Mauritaniz, Ægypti. — Shaw. afr. f. 231. *Z. desertorum* Forsk. descr. 87. ic. t. 11. Flores coccinei. (v. s.)
16. *Z. ALBUM* (Lin. dec. 1. t. 8.) foliis bifoliolatis petiolatis, foliolis cylindraceo-clavatis carnosius arachnoideo-incanis, pedicellis erectis, petalis obtusis, capsulis cylindraceo-pentagonis. $\frac{3}{5}$ in Ægypto, Barbariâ, Cauariis. DC. pl. grass. t. 154. *Z. proliferum* Forsk. descr. 87. ic. t. 12. A. — Caulis procumbens. (v. v.)

**** *Foliis bifoliolatis, foliolis scabris.*

17. *Z. PROSTRATUM* (Thunb. prodr. app. 189.) foliis bifoliolatis scabris, caule decumbente, geniculis hirtis. — ad cap. B.-Spei. †.
18. *Z. TRIDENTATUM* (fl. mex. ic. ined.) foliis bifoliolatis subsessilibus, foliolis obovatis villosis, petalis staminumque squamis apice tridentatis. $\frac{3}{5}$ in regno Mexicano. Petala flava. Stam. filam. purpurea. Fructus hirsutissimus, forsâ 5-spermus.

†. *Species anomalæ trifoliolatæ, forsâ à genere removendæ.*

19. *Z? LANATUM* (Willd. sp. 2. p. 564.) foliis trifoliolatis, foliolis subtus pillosis, caule flexuoso geniculis lanato. — in Sierrâ-Leonâ. Stam. filamenta basi parum dilatata. Styli 5! Capsula basi dehiscens 5-loc., 5-sperma. Ergo Planta proprii generis!

V. GUAIACUM. Plum. gen. t. 17. Lin. gen. n. 518.

Cal. 5-partitus obtusus. Pet. 5. Stam. 10, filam. nudis aut subappendiculatis. Stylus et stigma 1. Capsula substipitata 5-locul., 5-angulata aut abortu 2-3-locul. Semina in loculis solitaria, axi adfixa, pendula.

Albumen cartilagineum rimulosum. Cotyl. crassiusculæ (Gaertn. fr. 2. t. 113.) — Arbores ligno duro, foliis abruptè pinnatis, pedunc. axillaribus 1-floris.

1. *G. DUBIUM* (Forst. prodr. n. 186.) foliis 1-jugis, foliolis oblongo-lanceolatis obtusis. 5 in ins. Tongatabu. †.
2. *G. OFFICINALE* (Lin. sp. 546.) foliis bijugis, foliolis obovatis ovalibusve obtusis. 5 in Jamaicâ, Hispaniolâ. Sloan. hist. t. 222. f. 3. Lam. ill. t. 342. Blackw. t. 350. — Capsulæ sæpè biangulatæ. (v. s.)
3. *G. VERTICALE* (Ort. dec. 8. p. 93.) foliis 3-6-jugis, sæpiùs 5-jugis, foliolis ovali-oblongis mucronulatis coriaceis, extimis subovatis ramulisque glaberrimis. 3 in Novâ-Hispaniâ et Sancto-Domingo. Flores cærulei, petalis verticalibus. (v. s.)
4. *G. SANCTUM* (Lin. sp. 546.) foliis 5-7-jugis, foliolis ovalibus obtusis mucronulatis, petiolis ramulisque subpubescentibus. 5 in Americæ ins. Sancti-Johannis de Porto-Ricco. (v. s.)
5. *G. ARBOREUM*, foliis 7-14-jugis, foliolis ovali-oblongis obtusis basi inæqualibus sæpiùs alternis, petiolis ramulisque subpubescentibus. 5 in sylvaticis Carthagenæ, Guadalupæ et circa Barcellonam Cumanensium. H. B. et Kunth nov. gen. am. 6. p. 11.* *Zygophyllum* arboreum Jacq. am. 130. t. 83. Racemi laxi. Petala unguiculata. Stam. basi breviter appendiculata. Capsula stipitata 5-ptera. Lignum durissimum. Flos *Zygophylli*. Habitus et fructus Guayaci. Vernaculè *Guyacan* dictum. (v. s. sine fl.)

VI. PORLIERIA. Ruiz et Pav. prod. p. 55. t. 9.

Cal. 4-sepalus æqualis. Pet. 4 conniventia. Stam. 8 dorso squamarum totidem disci inserta. Carpella 4 connata drupacea non alata. — Guayaco formâ et viribus valdè affinis.

1. *P. HYGROMETRICA* (Ruiz et Pav. syst. p. 94.) 3 in Peruviz et Chili ruderatis. Folia abruptè pinnata 7-8-juga, foliolis linearibus. Folia aperta aut clausa serenitatem et tempestatem denunciant. (Ruiz et Pav.) (v. s.)

§. 2. ZYGOPHYLLÆ SPURIZÆ ALTERNIFOLIÆ.

VII. CHITONIA. fl. mex. ic. ined.

Cal. 4-partitus deciduus. Pet. 4 orbiculata. Stam. 8. Ovarium quadrangulum. Stylus 1. Stigma peltato-hemisphæricum. Capsula 4-locul., 4-valvis, valvis carinatis alato-tetragona. Semina in quoque loculo 2 ovata apice truncata arillo carnosso involuta. — Folia alterna forsam stipulacea impari-pinnata 6-juga, foliolis ovatis petiolulatis. Pedicelli gemini 1-flori.

1. *C. MEXICANA* (Fl. mex. ic. ined.) — in Mexico. Flores magni roseo-purpurei. Semina nigra. Arillus ruberrimus.

VIII. BIEBERSTEINIA. Steph. mem. nat. mosc. 1. p. 89.

Cal. 5-sepalus persistens irregularis. Pet. 5, æstiv. contorta; unguiculata. Stam. 10 hypogyna, filam. subulatis subvillosis, antheris oscillatoriiis. Ovaria 5 apice distincta basi connata. Styli 5 è medio ovariorum orti simplices. Carpella 1-sperma intus arillata? — Herbæ perennes pilis glandulosis. Folia alterna impari-pinnatisecta, segmentis incis, stipulis petiolo adnatis. Racemus terminalis simplex. — An Rosacea? sed stamina hypogyna. An Geraniacea? sed fructus axis nullus. An Rutacea? sed folia basi stipulacea et fructus non satis definitus.

1. *B. ODORA* (Steph. l. c.) foliorum segmentis subrotundis inciso-dentatis, petalis integris. $\frac{1}{2}$ ad torrentem Tschuia montium Altaïcorum. (v. s.)
2. *B. MULTIFIDA*, foliorum segmentis multifidis, lobis linearibus, petalis dentatis. $\frac{1}{2}$ in Persiâ. Michaux. (v. s.)

IX. MELIANTHUS. *Tourn. inst. t. 245. Lin. gen. n. 795.*

Cal. magnus 5-fidus persistens inæqualis, laciniâ infimâ minore deorsum gibbâ et cucullatâ intus glandulam mellisfluam gerente. Pet. 5 ligulæformia, 4 inf. declinata, medio connexa, quintum minimum. Stam. 4 hypogyna, fil. 2. super. distinctis, 2 infer. basi connatis. Ovarium 4-striatum. Styl. 1. Stigma sub-4-fidum. Capsula membranacea 4-loba, 4-locul., loculis 2-ovulatis et abortu 1-spermis, compresso-alatis, angulo inter. dehiscentibus. Semina ovata nitida. Albumen carnosum (Gaertn. fr. 3. p. 158. t. 211.) Embryo lobis parvis, radiculâ longiore (Juss. gen. 298.) — Frutices Capenses; folia alterna eglandulosa impari-pinnata, foliolis dentatis, stipulis distinctis aut coalitis. Racemi multiflori spiciformes. — Genus ex calyce ad Tropæoleas, ex pericarpio et stipulis ad Zygophylleas accedens; forsân ordo proprius!

1. *M. MAJOR* (Lin. sp. 892.) foliis utrinquë glabris, stipulis maximis cum petiolo et inter se in unicam coalitis. $\frac{3}{4}$ ad cap. Bonæ-Spei in succulentis. — Lam. ill. t. 552. — Herm. lugd.-bat. t. 415. (v. v.)
2. *M. MINOR* (Lin. sp. 892. excl. syn. Comm.) foliis supernè glabris subtus incanis, stipulis distinctis. $\frac{3}{4}$ ad cap. B.-Spei. Racemi axillares elongati. Flores verticillati. Bractææ lineares attenuatæ ex Valh symb. 3. p. 85. (v. s. sine fl.)
3. *M. COMOSUS* (Valh symb. 3. p. 85.) foliis supernè villosis, subtus tomentosis, stipulis distinctis. $\frac{3}{4}$ ad cap. B.-Spei. — Comm. rar. t. 4. Racemi infra foliacei. Flores alterni. Bractææ cordatæ. Capsulæ incanæ. (v. v.)

X? BALANITES. *Delil. fl. eg. 77.*

Cal. 5-partitus. Pet. 5. Stam. 10, filam. subulatis. Discus glandulosus ovarium cingens. Ovarium 5-loc., 5-spermum. Drupa ovoidea acuta abortu 1-loc., 1-sperma, nucleo lignoso pentagono. Semen pendulum, testâ fibrosâ, endoplevrâ circa radiculam incrassatâ. Embryo rectus, radiculâ superâ, cotyl. semiovatis, plumulâ diphyllâ. — Arbores, foliis alternis bifoliolatis, spinis axillaribus, pedicellis 1-floris aggregatis, floribus parvis albidis. An hujus ordinis?

1. *B. ÆGYPTIACA* (Del! l. c. t. 28. f. 1.) in Africâ calidiore, culta in Egypto. — Agiholid. P Alp. æg. 20. t. 11. *Ximenia Ægyptiaca* Lin. sp. 1194. Semina gummi Senegal sæpè mixta germinarunt in hort. Genev. (v. s. et v. sine fl.)

Æ. ferox, putamine acutiore angustiore magis sulcato, foliolis subrotundis, spinis longioribus. *Ximenia ferox* Poir. dict. 8. p. 805. in Sante-Domingo ubi verosim. à Nigritis introducta. (v. s.)

Ordo LI. RUTACEÆ. *Juss. gen.* 296. *excl. sect. primâ.* — *Diosmeæ Brown gen. rem. p.* 13. — *Fraxinellæ, Diosmeæ et Xanthoxyleæ Nees et Mart. nov. act. bonn. xi. p.* 149, 180 et 184.

Calyx 3-4-5-sepalus, sepalis basi plus minùsve inter se coalitis et ideò dentatus fissus aut partitus. Petala rarissimè 0, sæpiùs tot quot sepala, sæpè unguiculata, distincta, in Cuspariearum tribu et in Correis nonnullis inter se coalita et in corollam monopetalam subaggregata. Discus carnosoglandulosus ovarium cingens, è receptaculo ortus, extùs petala, superius stamina gerens. Stam. sæpiùs sepalorum numero dupla, et tunc aut omnia fertilia aut alterna sterilia, rariùs (in *Pegano*) sepalis tripla aut in quibusdam iis æqualia et tunc adhuc nonnulla sterilia evadunt. Carpella tot quot sepala, interdum abortu pauciora, nunc distincta, nunc basi coalita, nunc omninò connata. Stylus è centro ovariorum ortus, unicus in stigmata tot quot ovaria divisus et sic stylis ex ovario quoque ortis et concretis compositus; in paucissimis (*Zanthoxylon* et *Aubertia*) styli ex ovariorum apice orti distincti remanent. Carpella in fructu sunt sæpiùs distincta, 1-locularia, dehiscencia, bivalvia, intùs cocculosa, nempè endocarpium mutatur in cocculum elasticum bivalve cartilagineum intrà tegumentum exterius inclusum. Semina angulo interno adfixa inversa. Embryo rectus compressus axilis, radicula superà (1), cotyledonibus foliaceis.

Herbæ sæpiùs perennes aut Frutices ferè omnes glandulis variis scatentes et odore vario pollentes. Folia alterna aut opposita, simplicia aut composita, omnia exstipulata. Flores varii. Ordo medius inter Thalamifloras et Calycifloras; inter priores ad *Zygophylleas* *Simarubeasque* proximè accedens; inter posteriores *Rhamneis* affinis.

Trib. I. DIOSMEÆ.

Petala 3-5 à basi libera, æqualia et corollam regularem constituentia. Semina albuminosa.

I. RUTA. *Tourn. inst. t.* 133. *Lin. gen. n.* 523.

Cal. persistens 4-partitus rariùs 3-5-partitus. Pet. totidem unguiculata subcochleata. Stam. petalorum numero dupla. Pori nectariferi ad basim ovarii tot quot stamina. Ovarium stipite crasso brevi im-

(1) In *Boroniâ* infera dicitur.

situm. Stylus 1. Capsula subglobosa divisa in tot loculos quot sunt petala. Semina angulo interno adfixa. Albumen carnosum. Embryo arcuatus. Radicula longa. Cotyledones lineares. — Herbæ 2 aut suffrutices graveolentes glaucescentes. Folia alterna. Flores corymbosi lutei, centrales sæpè 5-fidi.

§. 1. *Pinnatæ, foliis verè pinnatis, foliolis nempè lateralibus super petiolum articulatis, fructibus subcarnosis indehiscenibus.* RUTERIA. Medik. phil. bot.

1. R. PINNATA (Lin. f. suppl. 232.) caule arborescente, foliis impari-pinnatis, floralibus ad foliolum terminale reductis, petalis integris. 3 in Canariis ad rupes. bot. reg. t. 307. (v. v.)

§. 2. *Decompositæ, foliis variè sectis decompositisve, capsulis dehiscentibus.*

2. R. MONTANA (Clus. hist. 2. p. 136.) foliis suprâ-decompositis, lobis omnibus linearibus, petalis integris. 2 3 in siccis collinis lapidosis Europæ austr. et Africæ bor. R. legitima Jacq. ic. rar. 1. t. 76. R. sylvestris Mill. dict. n. 3. R. tennifolia Desf. atl. 1. p. 336. (v. v.)
3. R. GRAVEOLENS (Lin. sp. 548. excl. var. α et β.) foliis suprâ-decompositis, lobis oblongis, terminali obovato, petalis integris aut subdentatis. 2 3 in sterilibus Europæ australis et in hortis frequens. R. hortensis Mill. dict. n. 1. — Duham. arb. 2. t. 61. (v. v.)
4. R. BRACTEOSA, foliis suprâ-decompositis latitudine vix triplò longioribus, lobis oblongo-cuneatis subæqualibus, bracteis maximis subcordatis, petalis ciliatis. 3 in Siciliâ circa Panormum (Morici.), in insulâ Melo (d'Urv.). R. Chalepensis tenuifolia d'Urv! enum. p. 44. (v. s.)
5. R. DIVARICATA (Ten! cat. h. neap. 1819. p. 42. non Saltzm.) foliis suprâ-decompositis, lobis obovatis cuneatis æqualibus, petalis integris aut subdentatis. 2 3 in apricis lapidosis Italiæ australis, Græciæ, Tauriæ. R. intermedia quorundam et revera omninò media inter graveolentem et angustifoliam. An prioris var.? — Lam. ill. t. 345. f. 1. (v. s.)
6. R. ANGUSTIFOLIA (Pers. ench. 1. p. 464.) foliis suprâ-decompositis latitudine quadruplò longioribus, lobis oblongo-cuneatis subæqualibus, bracteis minimis ovatis, petalis ciliatis. 2 3 in apricis lapidosis Galliæ australis frequens. R. Chalepensis Vill. dauph. 4. p. 383. R. graveolens α Lin. sp. 548. R. Chalepensis β Lin. mant. 69. — Moris. ox. s. 5. t. 35. f. 8. — Sims bot. mag. t. 2311. Folia angusta valdè glauca. (v. v.)
7. R. MACROPHYLLA (Sol. in bot. mag. t. 2018.) foliis pinnatisectis, segmentis oblongis subpetiolulatis, terminali majore, infimis 3-5-sectis, petalis ciliatis. 3 in Africâ (Lin.) in agro Chalepensi (Tourn.) R. Chalepensis α Lin. mant. 69. (v. v.)
8. R. CRITHMIFOLIA (Morici! ined.) foliis suprâ-decompositis, lobis oblongo-linearibus subcuneatis subæqualibus, petalis integris aut subdentatis. 2 3 in agro Foro-juliensi prope Duinam. (Moriciand.) inter Montefiascone et Triestam in rupibus (J. De Candolle.) et in Oriente. (Olivier.) Flores omnes 4-fidi in specim. nostro. (v. s.)
9. R. CORSICA, foliis suprâ-decompositis lobis obovatis subæqualibus, racemis subsimplicibus, pedicellis elongatis 1-floris rigidè divaricatis, petalis integris. 2 3 in Corsicæ montibus. R. montana spinosa alba Bocc. mus. p. 70. t. 59. ic. bona sed flores plantæ nostræ lutei! R. divaricata Saltzm! in florâ 1821. p. 109. Corymbi ab ipsâ basi bifidi, ramis racemosis! veteribus ex ic. Bocc. spiuescentibus. (v. s.)

§. 3. Trifoliæ, foliis nempe aut 3 integerrimis approximatis aut 3-partiis lobis integerrimis, infimis supremisque solitariis indivisis.

10. R. PATAVINA (Lin. sp. 549.) foliis mediis ternatis linearibus basi attenuatis integerrimis, corymbo conferto, calycibus villosis. \mathcal{U} \S in monte Saxo-Nigro ad Arquam prope Patavium. — Mich. gen. t. 19. Folia interdum basi dente lobulove aucta. (v. s.)
11. R. DICHOTOMA, foliis mediis ternatis linearibus basi attenuatis integerrimis, corymbo laxè paniculato dichotomo, calycibus subciliatis. \mathcal{U} \S in Persiâ. (Michaux.) Filam. stam. basi valdè dilatata, persistentia. (v. s.)
12. R. PUBESCENS (Willd. enum. 446.) foliis plerisque ternatis lanceolatis pubescentibus, lateralibus brevissimis, corymbo multifloro, calycibus ovarisque villosis. \mathcal{U} in Hispaniâ prope Aranjuez et Alania. R. Patavina Poir. dict. 6. p. 335. excl. syn. Stam. filam. basi subciliata vix dilatata. Lam. ill. t. 345. f. 2. (v. s.)

§. 4. Integrifoliæ, foliis indivisis integris solitariis.

13. R. FRUTICULOSA (Labill! syr. 1. p. 13. t. 4.) foliis integris oblongis obtusiusculis calycibusque villosis, corymbo paucifloro, petalis oblongis, filamentis basi ciliatis, ovario glabro. \mathcal{U} \S ad Damascus. (v. s.)
14. R. TUBERCULATA (Forsk. descr. 86.) foliis integris pilosis subtus caulibus capsulisque tuberculosis, caulibus herbaceis, capsulis pilosisculis rotundè 5-lobis, loculis 2-spermis. \mathcal{U} in Ægypto et Nubiâ. Flores parvi. Corymbi dichotomi. Petala oblonga.
 - a. Forskahlii, foliis lineari-lanceolatis. (v. s.)
 - β. Montbretii, foliis obovato-cuneatis. (v. s.)
15. R. GLABRA, foliis integris oblongo-linearibus subtus caulibusque glandulosis utrinquè caulibus calycibus capsulisque glabris, petalis oblongo-linearibus. \mathcal{U} inter Mossul et Bagdad. Olivier. Corymbus dichotomus. Stam. filamenta ciliata. R. tuberculatæ affinis. (v. s.)
16. R. VILLOSA (Bieb. fl. taur. 1. p. 310.) foliis integris infimis oblongis ceteris linearibus caulibusque pubescentibus, calycibus ovarisque hirsutis, petalis oblongis. \mathcal{U} \S in Caucasi orient. et Iberiæ campis apricis. — Buxb. cent. 2. t. 28. f. 2. R. parviflora Desf. cor. tourn. t. 54. (v. s.)
17. R. LINIFOLIA (Lin. sp. 549.) foliis integris oblongo-lanceolatis basi attenuatis caulibusque glabriusculis, calycibus brevissimis subciliatis, ovaris villosis, petalis ovatis. \mathcal{U} \S in Hispaniâ prope Valentiam et in agro Tunetano. — Barr. ic. t. 1186. Bocc. mus. t. 73. f. 3. (v. s.)
18. R. BUAVEOLENS, foliis integris spathulato-lanceolatis glaucis glabriusculis, calycibus subciliatis, ovaris glaberrimis, petalis ovatis. \mathcal{U} \S in Tauriæ et Bessarabiæ apricis. R. linifolia Bieb. fl. taur. 1. p. 309. R. linifolia grandiflora Sims bot. mag. t. 2254. Flos odorem Primulæ officinalis spirat ex cl. Edwards. (v. s.)
 - β. congesta, floribus paucis congesto-approximatis. \mathcal{U} in monte Olympo. Cas. Rostan. Flos citri odorem spirat. (v. s.)
19. R. BUXBAUMII (Poir. dict. 6. p. 336.) foliis integris lanceolatis in petiolum longè angustatis subpuberulis, paniculæ ramis dichotomo-corymbosis divaricatis ad dichotomias compressis, stam. filamentis basi dilatatis et ciliatis. \mathcal{U} in agris post messem circa Rodosto ex Buxb., in Troade. (Olivier.) et in Syriâ. (Labill.) — Buxb. cent. 2. t. 28. f. 2. mal. Flores flavi, sæpè sicci virescentes. Petala oblonga. Ovarium glabriusculum. (v. s.)
20. R. ACUTIFOLIA, foliis integris oblongis subpetiolatis utrinquè acutis cauleque glabris, infimis suboppositis, paniculâ corymbosâ dichotomâ, calycibus acutissimis subciliatis, capsulis rotundè lobatis glabris. \mathcal{U} in Persiâ. (Michaux), inter Kermancha et Amadan. (Olivier.) (v. s.)

21. *R. LINEARIS*, foliis integris linearibus glabris subtus punctatis, corymbis paucifloris dichotomis, petalis lineari-oblongis, capsulis glabris trilobis. \mathcal{L} in Sibiria australi. (v. s.)
22. *R. THESIOIDES* (Fisch! in litt.) foliis integris lineari-lanceolatis glabris, caulibus apice puberulis, corymbo coarctato, sepalis subrotundis, petalis ovatis, stam. filamentis basi dilatatis et intus hirsutis. \mathcal{L} ad littus orientale maris Caspici. Mangischlag. Valdè accedit ad *R. Dahuricam*. (v. s.)
23. *R. DAHURICA*, foliis integris lineari-lanceolatis caulibusque glabris, corymbo paucifloro, sepalis staminibusque basi subciliatis, ovarii glabris. petalis oblongis. \mathcal{L} in Dahuriæ campis apricis. *Peganum Dahuricum* \neq Lin. sp. 638. — *Ruta* Gmel. fl. sib. 4. t. 68. f. 2. — *Amm. ruth.* n. 92. Petala pallidè flava interdum 6. Stamina numero petalorum dupla nec, ut in *Pegano*, tripla. Radix fusiformis. Caules à collo plurimi simplices. — Varietas flore albo (*Amm.* n. 91.) à Linnæo citata remanet ignota. (v. s.)
24. *R. ROSMARINIFOLIA* (Pers. ench. 1. p. 465.) foliis linearibus subtus punctatis, caule ramosissimo, radice fusiformi. \mathcal{L} in Hispaniâ, ex herb. Juss. \dagger .

II. PEGANUM. *Lin. gen. n. 601.*

Cal. 5-partitus persistens, laciniis longis sæpius basi dentatis. Pet. 5. Stam. petalorum numero tripla, filam. basi dilatatis. Stigma triquetrum. Capsula substipitata subsphærica, 3-loc., apice 3-valvis polysperma. Embryo rectus. — Flores albi. Folia multifida, lobis linearibus.

1. *R. HARMALA* (Lin. sp. 638.) \mathcal{L} in arenâ Madriti, et in Oriente. Lam. ill. t. 401. Bull. herb. t. 343. *Harmala multifida* Mœnch meth. 239. Flores solitarii pedunculati. (v. v.)
- β . *crithmifolium* (Retz obs. 2. p. 34.) caule perennante, calycibus edentulis \mathcal{Z} ad mare Caspium. Var. culturâ orta ex Bieb. fl. taur. 1. p. 364. — *P. crithmifolium* Georg. in h. Willd. videtur *Adonidis* spec., ex Stev. in litt.

III. DICTAMNUS. *Lin. gen. n. 522. Fraxinella. Tourn. inst. t. 243.*

Cal. deciduus 5-partitus. Pet. 5, subinæqualia. Stam. 10 declinata, filam. apice glanduloso-tuberculosus. Stylus declinatus longitudinaliter striatus. Stigma simplex. Capsula carpellis 5 internè connexis compressis 2-spermis. Cotyl. obovato-cuneatæ. (Gaertn. fr. 1. t. 69). — Herbæ. Folia alterna impari-pinnata exstipulata. Caules apice glandulosi. Racemi terminales.

1. *D. FRAXINELLA* (Pers. ench. 1. p. 464.) \mathcal{L} in Europâ australi. — *D. albus* Lin. sp. 548. Lam. ill. t. 344. f. 1.

a. purpurea, petiolo obscure marginato, petalis pallidè purpureis venis intensè rubris striatis. *D. Fraxinella* Link enum. 1. p. 398. (v. v.)

β . *alba*, petiolo vix evidentius marginato, petalis albis. *D. albus* Link l. c. (v. v.)

Stirps utraque variat sepalis petalisque foliaceis. DC. fl. fr. 4. p. 734. (v. s.)

IV. CALODENDRON. *Thunb. prod. 44.*

Cal. brevis deciduus 5-part. Pet. 5 lanceolata glandulosa. Stam. 10, 5 antherifera, 5 sterilia petaliformia. Capsula 5-angul., 5-locul., 5-valvis, loculis 2-spermis. — Arbor. Folia opposita simplicia integra. Paniculæ terminales.

1. *C. CAPENSE* (Thunb. l. c. Lam. journ. h. nat. 56. t. 3.) \mathcal{Z} ad cap. Bonæ-Spei. *Dictamnus Calodendron* Lam. ill. t. 344. f. 2. Poir. suppl. 2. p. 476. *Dict. Capensis* Liu. f. suppl. 232. *Pallasia Capensis* Houtt. (v. s.)

V. DIOSMA. *Lin. gen. n. 272. Lam. ill. t. 127.*

Cal. 5-partitus. Pet. 5 disco hypogyno? inserta, sessilia, sepalis alterna. Stam. 5 antherifera petalis alterna disco inserta, 5 sterilia, nunc staminiformia, nunc petaliformia, nunc squamiformia aut minima. Styl. 1. Carpella 5 (abortu 2-4) compressa intus conniventia et dehiscencia. Semina 2 in quoque loc., nitida. Cotyl. oblongæ planoconvexæ (Gaertn. fr. t. 94). Frutices sæpius ericoidei, omnes Capenses; foliis punctato-glandulosis integris aut subcrenatis. (1)

Sect. I. ADENANDRA. *Willd. enum. 256. — Glandulifolia Wendl. coll. 1. t. 10.*

Stam. petalis breviora, 5 sterilia staminiformia apice antherarum rudimenta gerentia. — Flores magni sæpius terminales. Folia alterna planiuscula.

1. D. UNIFLORA (Lin. sp. 287.) foliis oblongo-lanceolatis glabris subtus punctatis, margine subrevolutis, floribus solitariis terminalibus, calycibus ciliatis. ζ Schrad. sert. hann. 1. t. 8. bot. mag. t. 273. Hartogia uniflora Berg. cap. 71. Eriostemon uniflora Smith in Rees. cycl. v. 13. n. 4. D. acuminata Lodd. bot. cab. t. 493. — Variat interdum 4-petala 4-andra. in hac et in seq. stam. filamenta hirsuta (v. s.)

2. D. AMOENA (Lodd. bot. cab. t. 161.) foliis ovalibus glabris subtus punctatis, floribus terminalibus solitariis sessilibus, calycibus subciliatis, petalis orbiculatis submucronatis. ζ Ker. bot. reg. t. 553. Pet. margine lævia subtus rubentia. An ad hanc referendæ Adenandra glandulosa Licht. in Rœm. et Schult. syst. 5. p. 450, et Glandulifolia uniflora ovata Wendl. coll. 1. t. 33?

3. D. SPECIOSA (Sims bot. mag. t. 1271.) foliis oblongis subobovatis subtus punctatis margine glanduloso-crenulatis glabris subciliatis, petiolis basi biglandulosis, floribus terminalibus umbellatis, calycibus ciliatis, petalis margine glabris. ζ D. cistoides Lam. dict. 2. p. 288? Stamina glabra. (v. s.)

a. multiflora, floribus 1-12, ramis arcuatim adscendentibus.

β. pauciflora, floribus 1-2, ramis rectis.

γ. umbellata, foliis omnibus ciliatis, calycibus subglabris, petalis ciliatis. ζ Glandulifolia umbellata Wendl. coll. 1. t. 10. herb. amat. t. 163. — Au species propria?

4. D. VILLOSA (Thunb. prod. 43.) foliis oblongo-lanceolatis ciliatis subtusque pubescentibus et glandulosis, floribus terminalibus umbellatis, calycibus petalis staminibusque ciliatis. ζ Hartogia villosa Berg. cap. 70. (v. s.)

5. D. MARGINATA (Thunb. prodr. 43.) foliis glabris cordatis, inferioribus ovatis, superioribus lanceolatis, umbellis terminalibus, stam. sterilibus barbatis. ζ Adenandra marginata Rœm. et Schult. syst. 5. p. 452.

6. D. FRAGRANS (Sims bot. mag. t. 1519.) foliis oblongis glandulosis glabris subcrenulatis, pedicellis glutinosis aggregatis subterminalibus folio duplò ferè longioribus, petalis emarginatis calycibusque imberbibus. ζ Flores rosei. (v. s.)

7. D. CORIACEA, foliis ovatis coriaceis, floribus axillaribus solitariis in ramorum apicibus, laciniis calyciis obtusis crenulatis. ζ Adenandra coriacea Licht. in Rœm. et Schult. syst. 5. p. 452. An hujus sect.?

8. D. TETRAGONA (Thunb. fl. 1. p. 133. voy. 4. t. 5.) foliis ovatis carinatis ciliatis quadrifariam imbricatis, floribus terminalibus solitariis. ζ Bucco

(1) Plurimas Diosmæ species forsan novas posideo, sed eas non indicavi, quia ob descriptiones maucas et paucas extricatu sunt difficillimæ, et generis confusionem jam nimiam augere nolui.

tetragona Roem. et Schult. syst. 5. p. 444. *Adenandra tetragona* Sweet h. suburb. p. 45. — *D. tetragona* Lin. f. suppl. 155 forsā diversa?

Sect. II. *BAROSMA*. Willd. enum. 257. — *Parapetalifera* Wendl. coll. 1. p. 92.

Stam. petalis subæqualia, sterilia petaliformia. — Flores pedicellati axillares. — Folia opposita plana glabra.

9. *D. SERRATIFOLIA* (Lodd. bot. cab. t. 373.) foliis lineari-lanceolatis serrulatis glabris glandulosis, pedicellis solitariis, infer. foliola 2 supra medium gerentibus. 5 Burch. voy. 1. p. 476. ic. Par. serrata Wendl. coll. 1. t. 34. (v. s.)
10. *D. ONORATA*, foliis ovato-oblongis crenatis glabris glandulosis, pedicellis solitariis immediatè sub flore bibracteolatis. 5 Par. odorata Wendl. coll. 1. t. 15. Ramuli tetragoni. Pedicelli interdum biflori. *D. latifolia* Lodd. bot. cab. t. 290. (v. s.)
11. *D. OVATA* (Thunb. prodr. 43.) foliis ovalibus ovatis obovatisve glabris subtus ferrugineo-punctatis integris, pedicellis ebracteolatis sæpius geminis. 5 *D. ovata* Sims bot. mag. t. 1616. Bucco ovata Wendl. coll. t. 20. (v. s.)
12. *D. CRENATA*, foliis ovatis acutis punctatis margine glanduloso-serratis, pedicellis solitariis subfoliosis. 5 *Diosma crenata* Lodd. bot. cab. 5. t. 404. An eadem ac *D. crenata* Lin. et Thunb. quæ habet folia sparsa ovata? An potiùs ad *Agathosma* referenda?
13. *D. DIOICA* (Ker. bot. reg. t. 502.) foliis lanceolatis glabris oppositis ternatimve verticillatis, pedicellis ternatis folio dimidiò brevioribus, floribus abortu dioicis. 5 Stam. subserta. Petala purpurascencia. *D. linifolia* Lodd. bot. cab. t. 400. non Licht.
14. *D. BETULINA* (Thunb. fl. cap. 2. p. 139.) foliis obovato-cuneatis crenatis subtus punctatis, pedicellis solitariis in racemos approximatis squammas obovatas gerentibus. 5 *Hartogia betulina* Berg. cap. 67. Bucco betulina Roem. et Schult. syst. 5. p. 443. Folia interdum alterna ex Thunb. An potiùs ad *Agathosma* referenda?

Sect. III. *AGATHOSMA*. Willd. enum. 259. Bucco Wendl. coll. 1. p. 13.

Stam. petalis æqualia aut longiora per anthesin exserta, 5 sterilia petaliformia. — Folia alterna. Flores terminales capitato-umbellati.

- 15? *D. ORBICULARIS* (Thunb. fl. cap. 2. p. 146.) foliis orbiculatis ovatis reniformibusve glabris reflexis minimis, ramis villosis, floribus ad apices racemosis. 5 Flores albi. Stam. corollà duplò longiora, undè inter *Agathosmas* numeranda videtur.
16. *D. PULCHELLA* (Lin. sp. 288?) foliis ovatis obtusiusculis glanduloso-crenatis glabris, floribus axillaribus geminis in racemum terminalem aggregatis. 5 *Hartogia pulchella* Berg. cap. 69. Bucco pulchella Roem. et Schult. syst. 5. p. 442. — Sims bot. mag. t. 1357. Valdè accedit ad *D. betulinam* et crenatam. (v. s.)
- 17? *D. LATIFOLIA* (Lin. f. suppl. 154.) foliis ovatis crenatis pubescentibus, ramis subtomentosis, pedicellis lateralibus 1-floris ad apices ramorum racemosis. 5 Andr. rep. t. 33. *Barosma latifolia* Roem. et Schult. syst. 5. p. 449. Differt ut prior à *Barosmâ* foliis sparsis.
18. *D. OBTUSA* (Roem. et Schult. syst. 5. p. 438. sub Bucco.) foliis lanceolatis obtusis ciliatis patentibus, pedicellis terminalibus hirsutis capitato-umbellatis, ovario glabro. 5 *D. ciliata* Lam. dict. 2. p. 287. Lodd. bot. cab. t. 210. Flores pallide aut intensius purpurascences.
 - a. *ovata*, foliis ovatis confertis, superioribus pilos glandulosos margine et nervo medio subtus gerentibus. Wendl. coll. 1. t. 13. *Agathosma pubescens* Willd. enum. 259.

- β. oblonga* (Wendl. coll. 1. t. 14.) foliis oblongis, super. pilos glandulosos margine et nervo medio subtus gerentibus. (v. s.)
- γ. lanceolata* (Ker. bot. reg. t. 476.) foliis lanceolatis longius ciliatis, pedicellis gracilioribus. *Hartogia lanceolata* Lin. syst. ed. 12. v. 2. p. 625. (v. s.)
19. *D. CILIATA* (Lin. sp. 287.) foliis lanceolatis acuminatis ciliatis, subtus punctatis et in nervo medio piliferis, demum deflexis, pedicellis terminalibus capitato-umbellatis pilosiusculis, ovario apice hirsuto. *♂* Ker. bot. reg. t. 366. Flores albi. Fructus sæpius bicornis. (v. v.)
20. *D. PROLIFERA* (Wendl. coll. 3. t. 77. sub Bucco.) foliis lanceolatis adgregatis pilosis punctatis apice barbatis patulis, pedicellis terminalibus umbellatis pilosis, calycibus glabris. *♂* Flores albi.
21. *D. RUGOSA* (Thunb. fl. cap. 2. p. 138.) foliis ovatis rugosis subtus villosis reflexis, pediculis umbellatis terminalibus hirsutis, calycibus pubescentibus. *♂* *Agathosma rugosum* Link enum. 238.
22. *D. AMBIGUA* (Lodd. bot. cab. t. 461.) foliis lineari-lanceolatis acuminatis ciliatis, ramis erectis caulinis patulis, umbellis confertis terminalibus multifloris. *♂* Flores albi. Accedit ad *D. ciliatam*.
23. *D. ACUMINATA* (Wendl. coll. 1. t. 28. sub Bucco.) foliis ovatis subcordatis longè acuminatis ciliatis demum patentibus, pedunculis terminalibus umbellatis villosis, calycibus glabris glandulosis. *♂* *Agathosma acuminata* Willd. enum. 260. Flores pallide cærulescentes. (v. s.)
24. *D. BARBATA* (Spreng. pug. 1. p. 20. sub *Agathosma*.) foliis lanceolatis acutiusculis glabris imbricatis, floribus terminalibus fasciculatis, calycibus ciliatis, stam. abortivis apice barbatis. *♂* *D. barbigera* Thunb. prodr. 43?
25. *D. IMBRICATA* (Thunb. fl. cap. 2. p. 145.) foliis ovatis acuminatis confertis imbricatis punctatis ciliatis, capitulis terminalibus umbellatis, calycibus glabriusculis, petalis staminibusque basi barbatis. *♂* Bucco imbricata Wendl. coll. 1. t. 9. Petala pallide purpurea, limbo subrotundo. (v. s.)
26. *D. VESTITA* (Licht. in Rœm. et Schult. syst. 5. p. 447. sub Bucco.) foliis ovatis ovato-lanceolatisque ciliatis tetragono-imbricatis, capitulis terminalibus. *♂*
27. *D. SERPYLLACEA* (Licht. in Rœm. et Schult. syst. 5. p. 447. sub Bucco.) foliis lineari-lanceolatis margine revolutis ramisque hirtis, floribus terminalibus umbellato-capitatis. *♂*
28. *D. SPICATA* (Licht. in Rœm. et Schult. syst. 5. p. 448 sub Bucco.) foliis linearibus margine revolutis verticillato-confertis, floribus capitato-spicatis. *♂*
29. *D. LINIFOLIA* (Licht. in Rœm. et Schult. syst. 5. p. 448. sub Bucco.) foliis linearibus acutis planis glabris scabriusculis margine pellucido-punctatis, floribus umbellato-capitatis. *♂*
30. *D. CEREFOLIUM* (Vent! malm. t. 93.) foliis lanceolato-linearibus acutis subimbricatis ciliatis, capitulis terminalibus, pedicellis calycibusque villosis, ovariis glabris. *♂* Folia trita odorem cerefolii spirant. Flores albi parvi. Bucco *Cerefolium* Rœm. et Schult. syst. 5. p. 439. (v. s.)
31. *D. WENDLANDIANA*, foliis lineari-lanceolatis acutis canaliculatis adgregatis imbricatis glandulosis villosis, capitulis umbellatis terminalibus, pedicellis puberulis, calycibus glabriusculis. *♂* Bucco villosa Wendl. coll. 1. p. 14. t. 2. excl. syn. Thunb. (v. s.)
32. *D. ERECTA* (Wendl. coll. 1. p. 17. t. 3. sub Bucco.) foliis oblongis acutis canaliculatis erecto-imbricatis punctatis glabris junioribus subpubescentibus, capitulis terminalibus 2-3-floris, pedicellis pilosis, calycibus petalisque glabris. *♂* Petala ex albo subcærulea.
33. *D. HIRTA* (Vent! malm. t. 72.) foliis lineari-lanceolatis erecto-imbricatis

supernè subconcavis subtùs hirtis, corymbis terminalibus multifloris, ovario supernè biglanduloso. ζ .

α . floribus intensè purpureis. — Ker. bot. reg. t. 369. D. purpurea hortul.

β . floribus pallidè purpureis. Vent! l. c. Bucco Ventenatiana Rœm. et Schult. syst. 5. p. 442. (v. s.)

γ . floribus albis? — Lam. dict. 2. p. 286. ill. t. 127. f. 3. var. β . exsiccata albescit.

34. D. *HISPIDA* (Thunb. fl. cap. 2. p. 129.) foliis trigonis muticis villosis hispida patulis, umbellis terminalibus, pedicellis calycibusque pubescentibus. ζ Hartogia Capensis Lin. sp. 288. Bucco hispida Rœm. et Schult. syst. 5. p. 446. Flores albi.

35. D. *BREVI-FOLIA* (Lam. dict. 2. p. 285.) foliis sublanceolatis triquetris subtùs punctatis subciliatis erectiusculis, corymbis terminalibus capitatis, pedicellis hirtis brevibus. ζ Bucco brevifolia Rœm. et Schult. syst. 5. p. 446. Flores pallidè violacei. Affinis priori.

36? D. *REFLEXA* (Lodd. cat.) foliis linearibus acutis apice subreflexis margine et subtùs glandulis seriatis, ramis glabris. ζ Agathosma reflexum Link. enum. 238. Flos ignotus.

37? D. *BRUNIADÆ* (Link. enum. 237.) foliis linearibus tenuissimis acutis margine glandulis ciliisque sparsis, ramis hirtis, floribus terminalibus adgregatis. ζ Foliis accedit ad D. rubram, floribus ad Agathosmas. (Link.)

SECT. IV. DICHOSMA.

Stam. petalis subæqualia per anthesin subexserta. Stam. 5 fertilia, 5 sterilia nulla. Petala longè unguiculata in lobos lineares profundè bifida.

38. D. *BIFIDA* (Jacq. coll. 3. p. 278. t. 20. f. 1.) foliis lanceolatis mucronatis punctatis glabris imbricatis, pedunculis in capitulum terminalem subglobosum adgregatis. ζ Flores in sicca albentes.

SECT. V. EUDIOSMA. — Diosma. *Wendl. coll. 1. p. 1. Willd. enum. 257.*

Stam. petalis breviora, 5 fertilia, 5 sterilia evanida aut subsquammiformia. Petala subsessilia integra. — Flores terminales parvi.

* *Foliis oppositis.*

39. D. *OPPOSITIFOLIA* (Thunb. fl. cap. 2. p. 132.) foliis trigonis obtusis ciliatis floribus terminalibus subsessilibus. ζ . (v. s.)

40. D. *SUCCULENTA* (Wendl. coll. 1. t. 1.) foliis linearibus carinatis acutis crassiusculis ciliatis erectis, floribus terminalibus subsessilibus. ζ . (v. s.)

N. B. Confer an D. oppositifolia Lin. sit eadem ac Thunbergii? An D. pinifolia Fisch. nec non D. scabra et decussata Lam. dict. 2. p. 283. ad duas priores sint referendæ aut species distinctæ? An Icones Pluk. t. 279. f. 4. Comm. rar. t. 1. et Seb. thes. 2. t. 40. f. 5. inter se ut videntur diversæ, ad duas priores pertineant aut indicationes sint specierum oppositifoliarum nondum satis cognitarum?

41. D. *CUPRESSINA* (Thunb. fl. cap. 2. p. 134.) foliis oblongo-lanceolatis carinatis adpressis margine scabris, floribus terminalibus subsolitariis. ζ Brunia uniflora Linn. sp. 289. — Pluk. t. 279. f. 2. — Wendl. coll. 2. t. 61. — Lodd. cab. t. 303. (v. s.)

42. D. *LINEARIS* (Thunb. fl. cap. 2. p. 136.) foliis linearibus obtusis patulis, ramis pedicellisque glabris, floribus terminalibus subsolitariis. ζ .

** *Foliis alternis.*

43. D. *OBTUSATA* (Thunb. fl. cap. 2. p. 126.) foliis trigonis acutis punctatis, calyce petalisque ciliatis, ramulis pubescentibus. ζ Flores in apice ramorum solitarii rubri.

44. *D. PECTINATA* (Thunb. fl. cap. 2. p. 127.) foliis trigonis acutis punctatis ciliatis patentibus, floribus terminalibus, ramulis pubescentibus. *h* Rœm. et Schult. syst. 5. p. 453.
45. *D. SUBULATA* (Wendl. coll. 1. p. 31. t. 8.) foliis lineari-triquetris subulato-acuminatis ciliatis erectis interdum oppositis, floribus ramulorum terminalibus subcymosis. *h* Petala cœrulescentia calyce acuto glabro breviora. (v. s.)
46. *D. ALBA* (Thunb. fl. cap. 2. p. 126.) foliis linearibus carinatis mucronatis margine cartilagineis scabris, ramis glabris, floribus terminalibus subsolitariis. *h* Adenandra alba Rœm. et Schult. syst. 5. p. 451. Flores albi. Calyces et bracteæ ciliati.
47. *D. RUBRA* (Lin. sp. 287.) foliis linearibus subtrigonis carinatis mucronatis glabris molliterve ciliatis subtus bifariam punctatis, floribus terminalibus subumbellatis, calycibus glabris. *h* Ker. bot. reg. t. 563. *D. tenuifolium* Willd. enum. 258? *D. ericifolia* Andr. bot. rep. t. 451. Calyces rubescentes. Petala alba subrosea. (v. s.)
48. *D. HIRSUTA* (Thunb. fl. cap. 2. p. 128.) foliis linearibus carinatis mucronatis villosis - hirsutis, pedunculis terminalibus 1-floris subcorymbosis, ramis calycibusque hirtis. *h* Wendl. coll. 1. t. 27. Petala alba subcœrulescentia calyce longiora. (v. s.)
49. *D. BISULCA* (Thunb. prod. 1. p. 84.) foliis trigonis acutis hirsutis, racemis terminali - subumbellatis. *h* *D. bifurca* Willd. sp. 1. p. 1136. Folia acumine glanduloso subtus bisulca. Flores albi.
50. *D. VIRGATA* (Thunb. fl. cap. 2. p. 129.) foliis trigonis obtusis punctatis glabris, floribus subracemosis fastigiatis. *h* Flores albi glabri.
51. *D. ERICOIDES* (Thunb. fl. cap. 2. p. 130.) foliis trigonis obtusis glabris punctatis, floribus terminalibus subsolitariis. *h* — Pluk. t. 179. f. 3. — Mill. ic. t. 124. f. 2.
52. *D. ASPALATHOIDES* (Lam. dict. 2. p. 286.) foliis oblongis linearibusve acutis triquetris subtus punctatis glabris, apicibus extrorsum aduncis, floribus terminalibus subsolitariis. *h* Ad eadem ac præcedens ex cl. Poir? (v. s.)
53. *D. CAPITATA* (Lin. mant. 210.) foliis trigonis obtusis villosis - hispidis imbricatis, floribus spicato - capitatis. *h* *D. imbricatum* Dum. - Cours. bot. cult. ed. 2. v. 5. p. 405. (v. s.)
54. *D. LONGIFOLIA* (Wendl. coll. 1. p. 61. t. 19.) foliis linearibus cuspidatis longis glabris glandulosis, floribus ramulorum terminalibus paucissimis. *h* Petala alba subcœrulescentia.
55. *D. USTULATA* (Thunb. fl. cap. 2. p. 132.) foliis ovatis trigonis imbricatis glabris, flore terminali solitario. *h* Folia super. nigro-ustulata.
56. *D. UMBELLATA* (Thunb. fl. cap. 2. p. 133.) foliis trigonis obtusis punctatis ciliatis, umbellis multifloris terminalibus. *h* Pedunculi purpurei. Corollæ albæ.
- †. *Species non satis notæ, quoad sectionem dubiæ.*
57. *D. PUNCTATA* (Licht. in Rœm. et Schult. syst. 5. p. 461.) foliis orbiculatis integerrimis quintuplinerviis subtus punctatis. *h*
58. *D. CORDATA* (Mart. enum. h. erl. 67.) foliis subcordatis oblongis acutis ciliatis recurvato-imbricatis. *h*
59. *D. BARBIGERA* (Lin. f. suppl. 155.) foliis cordatis acutis glabris, umbellis terminalibus, petalis barbatis. *h* Stylus flore longior. Videtur *Agathosmæ* sp. Conf. cum *D. barbata*.
60. *D. SQUAMMOSA* (Willd. in Rœm. et Schult. syst. 5. p. 462.) foliis subrotundo-ovatis obtusis ciliatis minutis imbricatis, caule dichotomo, floribus terminalibus capitatis. *h*

61. *D. GLANDULOSA* (Thunb. fl. cap. 2. p. 145.) foliis ovatis acutis integris rugosis subtus villosis, umbellis terminalibus. 5.
62. *D. GRAVEOLENS* (Licht. in Rœm. et Schult. syst. 5. p. 461.) foliis ovatis integerrimis imbricatis glabris, floribus axillaribus pedunculatis solitariis. 5.
63. *D. LYCOPODIODES* (Willd. in Rœm. et Schult. syst. 5. p. 461.) foliis ovatis acutis tomentosis dense imbricatis, caule dichotomo, floribus terminalibus capitatis. 5.
64. *D. THYMIFOLIA* (Willd. in Rœm. et Schult. syst. 5. p. 462.) foliis ovatis ciliatis patentibus obtusis margine revolutis, pediculis terminalibus umbellatis glanduloso-pubescentibus. 5 Conf. cum *Linconia thymifolia*.
65. *D. CORYMBOSA* (Mont. in act. lund. 1. 2. n. 11.) foliis ovato-lanceolatis carnosiss integerrimis. 5.
66. *D. MYRSINITES* (Lam. ill. 2. p. 82.) foliis lanceolatis planis margine ciliato-hispidis, umbellis terminalibus, pedunculis glabris. 5 Conf. cum *D. marginata*. — Synon. Seab. (2. t. 96. f. 6.) male indicatum non reperio.
67. *D. PERFORATA* (Lam. ill. 2. p. 82.) foliis lanceolatis acuminatis subciliatis punctis marginalibus pellucidis subperforatis, umbellis terminalibus. 5.
68. *D. OBLONGA* (Thunb. fl. cap. 2. p. 139.) foliis oblongo-lanceolatis glanduloso-crenatis rugosis, floribus lateralibus. 5 *D. lanceolata* ß et γ. Thunb. prod.
69. *D. ROSMARINFOLIA* (Lam. ill. 2. p. 81.) foliis lanceolato-linearibus margine revolutis glabris, pedunculis axillaribus et terminalibus, capsulis obtusissimis. 5.
70. *D. STRICTA* (Willd. in Rœm. et Schult. syst. 5. p. 462.) foliis lineari-lanceolatis strictis pilis distantibus rigidis ciliatis, floribus capitatis. 5.
71. *D. TERETIFOLIA* (Link enum. 1. p. 237.) foliis linearibus supra convexis subtus canaliculatis obtusis ramisque pubescentibus. 5 Flores ignoti.
72. *D. TENUISSIMA* (Lodd. in Link enum. 1. p. 257.) foliis subtriquetris glabris remotiusculis vix 2 lin. longis. 5.
73. *D. THYROIDES* (Willd. in Rœm. et Schult. syst. 5. p. 462.) foliis lineari-subulatis glabris adpressis, ramis verticillatis, floribus terminalibus umbellatis, pedunculis pubescentibus. 5.
74. *D. PARVIFLORA* (Willd. in Rœm. et Schult. syst. 5. p. 462.) foliis subulato-triquetris acutis glabris, floribus umbellatis terminalibus, pedicellis pubescentibus. 5 Flores minuti albi.
- ††. *Species verosim. è genere excludenda, sed non satis nota.*
75. *D. ASIATICA* (Lour. coch. 1. p. 200.) foliis lanceolatis glabris alternis integris, racemis compositis subterminalibus. 5 in monte Hòn Chén Cochinchinæ. Differt à *Diosma* floribus flavis racemosis, antheris sessilibus, capsulis stipitatis distinctis, seminibus non calyptratis. Ergò genus diversissimum (*PSEUDIOSMA*) forsàn non *Rutaceum*.

VI. EMPLEURUM. Sol. in Ait. h. kew. 3. p. 340.

Flores abortu monoici polygamive. Cal. 4-fidus persistens. Pet. 0. Stam. 4, antheris filamentis longioribus apice glandulosis. Carpella 2 rarius solitaria stigmata cylindraceo dentiformi ad latus interioris instructa, demùm leguminiformia, bivalvia 1-loc. 2-ovulata et abortu 1-sperma. Semen nitidum. Cotyl. impares convolutæ (Gaertn. fr. 3. p. 157. t. 211.) — Fruticulus glaber glandulosus foliis lineari-lanceolatis serrulatis.

1. *E. SERRULATUM* (Sol. l. c. Smith exot. bot. 2. t. 63.) 5 ad cap. B.-Spei. Emp. Aitonii Gmel. syst. 1. p. 268. *Diosma uncapularis* Lin. f. suppl. 155. *D. ensata* Thunb. prod. 43. (v. s.)

VII. DIPLOLAENA. *Brown gen. rem. 14. Desf. mem. mus. 3. p. 449.*

Involucrum duplex multiflorum, exterius 5-lobum, interius 10-15-partitum exteriore longius. Flores sessiles involucello calyceve 5-paleaceo instructi. Pet. o. Stam. 10 infernè villosa. Annulus glandulosus ovarium cingens. Stylus 1. Stigma 5-dent. Carpella 5 bivalvia 1-sperma transversè striata. Semina oblongo-cylindracea, hilo laterali, ferè Correæ. — Caules fruticosi ramosi. Folia alterna ovalia punctatoglandulosa, ceterum foliis Correæ albæ similia. Capitula terminalia pediculata.

1. *D. GRANDIFLORA* (Desf! l. c. t. 19.) foliis ovalibus obtusis utrinquè incanis. ζ in arenosis Novæ-Holl. occidentalis in terrâ Endracht. Capitula 2 poll. diam. (v. s.)
2. *D. DAMPIERI* (Desf! l. c. t. 20.) foliis obovato-oblongis emarginatis superne viridibus glabris, subtus incanis. ζ cum priore. — Hanc aut priorem depinxit Dampier voy. 4. p. 141. t. 3. f. 3. (v. s.)

VIII. CORREÆ. *Smith tr. lin. soc. 4. p. — Mazeutoxeron Labill. voy. 2. p. 11.*

Cal. 4-dentatus persistens. Pet. 4 basi subconniventia aut in tubum longè coalita. Stam. 8 sub disco hypogyno? 8-glanduloso inserta. Ovar. 8-sulcatum. Styl. 1 persistens. Capsula 4-cocca, loc. truncatis compressis. Semina in loculis 2-3 nitida intus adfixa, cotyl. ovalibus extus convexis (Vent. malm. 13.) — Frutices, foliis oppositis integris pube squamosâ, Hippophaes more, obtectis, pedicellis 1-floris.

* *Brevifloræ, petalis plus minus distinctis semi-pollicem longis.*

1. *C. ALBA* (Andr. bot. rep. t. 18.) foliis ovatis subtus tomentosis, calycinis dentibus parvis acutis distantibus. ζ in Novâ-Hollandiâ. — Vent. malm. t. 13. *C. cotinifolia* Salisb. par. lond. t. 100. Flores albi aut intus carnei ut in ic. Salisb. Folia subtus albo aut rufo tomento vestita, superne subalbida. (v. v.)
 β . *rotundifolia*, foliis ovato-rotundis. *C. rufa* Vent. malm. 13. in adu. n. 2. (v. s.)

2. *C. RUFA* (Gaertn. f. fr. 3. p. 155. t. 210.) foliis ovatis suboblongis subtus tomentosis, calycinis dentibus latis obtusissimis. ζ in Novâ-Hollandiâ. Maz. rufam Labill! voy. 2. p. 11. t. 17. Folia subtus rufo-tomentosa superne obscurè viridia glabra. (v. s.)

** *Longifloræ, petalis omninò coalitis in corollam cylindraceam sesquipollicarem.*

3. *C. SPECIOSA* (bot. rep. t. 653.) foliis ovatis obtusis subtus ferrugineo-tomentosis, floribus erectis, calyce truncato. ζ in Novâ-Holl. orientali. — Ker. bot. reg. t. 26. Sims bot. mag. t. 1746. Lois. herb. am. t. 309. — *C. rubra* Smith exot. bot. 2. n. 26. *C. revoluta* Vent. l. c. n. 4? Corollæ rubræ. (v. s.)
4. *C. VIRENS* (Smith exot. bot. 2. p. 25. t. 72.) foliis ovato-oblongis cordatis, floribus pendulis, calyce acutè 4-denticulato. ζ in Nov.-Holl. orientali et merid. *C. viridiflora* Andr. bot. rep. t. 436. *C. reflexa* Pers. ench. 1. p. 419. Maz. reflexum Labill! 2. p. 66. t. 19. Cor. virescentes poll. sesquipollicisve longæ. (v. s.)

IX. PHEBALIUM. *Vent. malm. n. 102.*

Cal. 5-fidus. Pet. 5 sessilia calyci? imò inserta. (Vent.) extus cum calyce lepidota. Stam. 10, filam. glabris, antheris terminalibus. Styl. 1. Capsula calyce cincta, 5-loc., 5-valvis, oligosperma. — Frutices lepidoti, foliis alternis simplicibus integris, floribus corymbosis terminalibus.

1. *P. ANCEPS*, foliis lineari-lanceolatis obtusis, junioribus utrinquè lepidotis,

adultis subnudis, staminibus petalis brevioribus. ♀ in Novâ-Holl. orientali. An potius *Eriostemonis* species, sed flores lepidoti corymboso-terminales. (v. s.)

2. *P. squammulosum* (Vent. malm. t. 102.) foliis lineari-oblongis acutis superne glabris subtus lepidotis, staminibus exsertis. ♀ in montibus Novæ-Hollandiæ orientalis. (v. s.)

X. *CROWEA*. *Smith tr. lin. soc. 4. p. 220.*

Cal. 5-partitus. Pet. 5, sessilia, æstivatione quincunciali. Stam. 10, filam. basi planis, apice subulatis, pilis longis ciliatis, antheram adnatam gerentibus et superne apiculatis appendiculatisve. Stylus 1 brevissimus. Carpella 5 coalita in capsulam 5-angul., 5-locul. Sem. in loculis solitaria. — Frutices, ramis triquetris, foliis alternis lineari-lanceolatis integerrimis. Pedunculi 1-flori axillares. Flores purpurei extus vernicosi.

1. *C. SALIGNA* (Andr. bot. rep. t. 79) foliis lanceolatis, calycibus petalisque glabris, antheris in appendicem barbatam longè productis. ♀ in Novâ-Holl. orientali. Vent! malm. t. 7. (v. s.)

XI. *ERIOSTEMON*. *Smith in tr. lin. soc. 4. p. 221.*

Cal. 5 part. Pet. 5. Stam. 10, filam. hispidis ciliatis aut nudis, antheris terminalibus. Styl. 1 brevissimus. Carpella 5 basi coalita. Semina in loculis 2 aut abortu solitaria. Embryo subcurvatus, radic. longâ. — Frutices Arboresve nunc *Diosmis* nunc *Croweis*, nunc *Phebalis* affines, foliis alternis simplicibus, floribus axillaribus

* *Filamentis stam. hispidis ciliatisve, foliis calycibusque non lepidotis.*

1. *E. BUXIFOLIUM* (Smith in Rees cycl. 13. n. 2.) foliis ellipticis glabris, nervo medio prominulo et apice in mucronem validum producto, ramis pilosis teretibus, floribus axill. sessilibus glabris, filamentis hispidis. ♀ ad Port-Jackson. Habitus *Diosmæ*. An duæ var. distinguendæ?

α. foliis obovatis basi attenuatis crenato-glandulosis. (Smith 1 c.)

β. foliis latè ellipticis basi cordatis subamplexicaulibus sæpius integris glandulosis. Smith 1 c. (v. s.)

2. *E. SALICIFOLIUM* (Smith in Rees cycl. 13. n. 1.) foliis lineari-lanceolatis integerrimis glabris, ramulis triquetris, floribus axillaribus subsessilibus basi bracteolatis solitariis, calycibus petalisque extus canescentibus, filamentis hispidis. ♀ ad Port-Jackson. Habitus *Croweæ*. (v. s.)

3. *E. MYOPOROÏDES*, foliis lineari-lanceolatis acumine calloso mucronatis integerrimis glabris glanduloso-punctatis, ramulis teretibus, pedunculis axillaribus 3-fidis 3-floris, calycibus petalisque glabris, filamentis margine subciliatis. ♀ in Novâ Holl. orientali. (v. s.)

4. *E. LINEARIFOLIUM*, foliis linearibus obtusis integerrimis glabris glanduloso-punctatis, pedunculis axillaribus terminalibusque trifloris. ♀ in Novâ-Hollandiâ. Flores in meo specim. non explicati, sed affinitas cum priore summa genus indicat. (v. s.)

** *Filamentis staminum glabris, foliis calycibusque lepidotis.*

5. *E. SQUAMMEUM* (Labill! nov-holl. t. p. 111. t. 141.) foliis lanceolatis integris, subtus squamis argenteis subrufris onustis, ramulis angulatis, pedunculis axillaribus 3-fidis 3-floris, filamentis glabris. in Novâ-Hollandiâ. Habitus *Phebalii*. (v. s.)

†. *Species non satis nota.*

6. *E. LANCEOLATUM* (Gaertn. f. fr. 3. p. 154. t. 210.) foliis lanceolatis subrugosis. ♀ in Novâ-Holl. *E. australasium* Smith tr. soc. lin. 4. p. 221.

XII. PHILOTHECA. *Rudg. in tr. lin. soc. 11. p. 298.*

Cal. 5-partitus. Pet. 5. Stam. 10 usque ad medium in urceolum monadelphæ, in parte liberâ hirsuta, antheris oscillatoriis. Stylus 1. Carpella 5 et verosim. abortu 2-4 ovata 1-sperma intus coalita. — Flores terminales. Folia alterna integra.

1. *P. AUSTRALIS* (Rudg. l. c.) 5 in Novæ-Hollandiæ orientali. *Eriostemon salso-lifolia* Smith in Rees cycl. 13. n. 3. Fruticulus ericoideus, foliis numerosissimis linearibus subtus convexis subimbricatis. (v. s.)

XIII. BORONIA. *Smith tracts. p. 285.*

Cal. 4-fidus persistens. Pet. 4 ovata persistentia Stam. 8, rarius 4 sepalis opposita antherifera, 4 altera abortiva, filam. ciliatis incurvis. Styli 4 erecti approximati aut inter se coaliti. Carpella 4 bivalvia introrsum connata in capsulam 4-lobam 4-loc. Semina in loc. subsolitaria ovata compressa. Embryo rectus in album. carnosus, radícula infera ex Gaertn. f. fr. 3. p. 156. t. 211. — Frutices Novæ-Hollandiæ. Folia opposita. Pedunculi axill. Flores sæpius purpuracentes.

* *Foliis impari-pinnatis, foliolo terminali sessili.*

1. *B. PINNATA* (Smith l. c. t. 4.) foliis impari-pinnatis 2-4-jugis glaberrimis, foliolis linearibus acutis, pedunculis dichotomis, floribus 8-andris. 5 in Novæ-Holl. ad Port-Jackson. — Andr. bot. rep. t. 58. Vent! malm. t. 38. (v. s.)
2. *B. PSORALEOIDES*, foliis impari-pinnatis 1-2-jugis ramulisque glaberrimis punctato-glandulosis, foliolis linearibus obtusis, pedunculis brevibus 1-floris, floribus 4-andris. 5 in Novæ-Hollandiæ orâ meridionali. Flores parvi pallidi. (v. s.)
3. *B. TETRANDRA* (Labill! nov.-holl. 1. p. 98. t. 125.) foliis impari-pinnatis 4 5-jugis, foliolis linearibus obtusis glabris, ramis pilosis, pedicellis brevibus 1-floris, floribus 4-andris. 5 in Nov.-Holl. terrâ Van-Leuwin. (v. s.)
4. *B. PILOSA* (Labill! nov.-holl. 1. p. 97. t. 124.) foliis impari-pinnatis 2-5-jugis ramulisque pilosiusculis, foliolis linearibus approximatis, pedunculis 1-floris, floribus 8-andris. 5 in Nov.-Holl. Capite Van-Diemen. Petala pellucido-punctata. (v. s.)
5. *B. ALATA* (Smith tr. lin. soc. 8. p. 283.) foliis impari-pinnatis sub-5-jugis, rachi nervisque subtus pilosis, foliolis crenatis revolutis, pedunculis dichotomis, bracteis fimbriatis. 5 in Novæ-Holl. orâ occidentali. Petala superne albida; filam. obtusa, antheræ subtérminales.

** *Foliis simplicibus.*

6. *B. SERRULATA* (Smith tr. lin. soc. 8. p. 284. tracts. t. 5.) foliis trapeziformibus acutis anticè serrulatis glabris punctato-glandulosis, pedunculis aggregatis terminalibus. 5 in Novâ Holl. ad Port-Jackson. Flores rosei. Stam. filam. hispida, 4 longiora apice dilatata in discum subobcordatum medio intus orbiculatum. Stigma sessile capitatum sub-4-sulcum. (v. s.)
7. *B. CRENULATA* (Smith l. c. p. 284.) foliis obovatis mucronatis crenatis, pedicellis axillaribus terminalibusque 1-floris, filamentis apice obtusis glandulosis. 5 in Nov.-Holl. ad portum Regis-Georgii. Flores parvi rubri. Calyx fimbriatus.
8. *B. DENTICULATA* (Smith l. c. p. 284.) foliis linearibus retusis submucronatis denticulatis, pedunculis corymbosis, filamentis apice obtusis glandulosis. 5 in Nov.-Hollandiâ ad portum R.-Georgii. Petala pallidè rosea. Bracteolæ deciduæ.
9. *B. PARVIFLORA* (Smith l. c. p. 285. tracts. t. 6.) foliis obovato-lanceolatis obsolete crenatis, pediculis 1-floris bracteolatis, stam. filamentis apice

oblongis glandulosis. 5 in Nov.-Holl. prope Port-Jackson. Habitus B. denticulata sed minor.

10. B. TETRATHECOIDES, foliis linearibus integerrimis glabris utrinquē subacutis margine subrevolutis, pedunculis axillaribus 1-floris brevibus bibracteolatis, filamentis hispidis. 5 in Novâ-Holl. orientali. Tetrathea oppositifolia Pers. ench. 1. p. 419. (v. s.)
11. B. PILONEMA (Labill. nov.-holl. 1. p. 98. t. 126.) foliis oblongo-linearibus integerrimis utrinquē subacutis, pedunculis terminalibus solitariis ebracteolatis subturbinatis, stam. filamentis glabris apice abbreviatis. 5 in Nov.-Holl. Cap. Van-Diemen. (v. s.)
12. B. POLYGALÆFOLIA (Smith tr. lin. soc. 8. p. 285. tracts. t. 7.) foliis lineari-lanceolatis integerrimis, pedunculis axillaribus solitariis 1-floris, filamentis stam. apice abbreviatis. 4 in Novâ-Holl. ad Port-Jackson. Folia interdum ternatim verticillata aut alterna.
13. B. LEDIFOLIA (Gay! diss. las. p. 20.) foliis lineari-lanceolatis integerrimis subtus tomentosis, pedunculis axillaribus 1-floris medio bibracteolatis, filamentis stam. hispidis. 5 in Novâ-Holl. orientali. Lasiopetalum ledifolium Vent. malm. n. 59. in adn. Alabastra quadrata. (v. s.)

*** *Species dubia foliis variis.*

14. B? PARADOXA, foliis simplicibus trifoliolatis pinnatisve lanceolatis margine revolutis subtus hirsutis. 5 in Novâ-Holl. Eriostemon paradoxa Smith in Rees cycl. v. 13. n. 6. Flores axillares. Bracteolæ 2 in medio pedicelli. Calyx hirsutus. Pet 4. Habitus Boroniæ. (Smith.)

XIV. CYMINOSMA. Gaertn. fr. 1. p. 280. — Jambolifera. Lin. gen. n. 479. non Houtt. nec Gaertn. et excl. syn. Rumph. idèque nomine.

Cal. 4-sepalus, sepalis orbiculatis, 2 subminoribus. Pet. 4 lanceolata patentia. Stam. 8, alterua receptaculo, alterna patalorum basi inserta, basi pilosa. Antheræ versatiles. Discus carnosus octangularis circa ovarium. Stylus 1 erectus. (Koen. in Dryand. act. soc. lin. 2. p. 233.) Bacca 4 locul. loculamentis intus chartaceo-crustaceis, 1-spermis. Embryo radic. brevi, cotyl. ellipticis. (Gaertn. l. c.) — Arbusculæ; foliis oppositis integris, floribus corymbosis, fructibus foliisque odorem Cumini spirantibus. — Species malè cognitæ.

1. C. PEDUNCULATA, foliis lanceolato-ellipticis obtusis, petalis lineari-lanceolatis calyce triplo longioribus. 5 in Ceylonâ. Jambolifera Lin. fl. zeyl. 58. excl. syn. J. pedunculata Vahl symb. 3. p. 52. t. 61*. Dryand l. c.* — Perin-pandel Rheed. mal. 5. t. 15? — Gaertn. fr. 1. p. 281. in adn. — Jambolifera pedunculata Lour. coch. 1. p. 283 videtur diversa: ob folia emarginata, et forsân à genere aliena.
2. C. ANKOENDA (Gaertn. l. c. t. 58.) foliis, petalis oblongis extus pubescentibus calyce paulo longioribus. — in Ceylonâ. — Ankoenda Hermann. 73. Burm. fl. zeyl. 27.
3. C. ODORATA, foliis ovatis basi obliquè truncatis subtus punctatis, corymbis terminalibus racemosis, baccis (abortu?) 1-spermis. 5 in hortis Cochinchinæ. Jambolifera odorata Lour. coch. 1. p. 284.
4. C. RESINOSA, foliis oblongis, pedunculis axillaribus multifloris, baccis subrotundis 4-localaribus. 5 in dumetis Cochinchinæ. Jambolifera resinosa Lour. coch. 1. p. 284.

XV. ZIERIA. Smith tr. lin. soc. 4. p. 216.

Cal. 4-fidus. Pet 4, disco hypogyno inserta. Stam. 4 cal. lobis opposita, filam. glabris basi intus 1-glandulosis. Styl. 1. Stigma 4-lobum. Carpella 4 intus connexa in capsulam 4-lobam., 4-loc., lobis

divaricatis. Semina in loculis solitaria compresso-ovata. — Frutices aut arbores. Folia opposita petiolata, trifoliolata, summa simplicia. Pedunculi axillares sæpius trichotomi. Flores parvi albi.

1. *Z. LANCEOLATA* (Brown in Sm. cycl. n. 1.) ramulis pubescentibus, foliolis lanceolatis planis acutis pellucido-punctatis, pedunculis trichotomis longitudine foliorum. § in Novâ-Holland. orient. *Z. Smithii* Andr. bot. rep. t. 606. Sims bot. mag. t. 1395. Bonpl. nav. t. 24. (v. s.)
2. *Z. MACROPHYLLA* (Bonpl. nav. p. 64?) ramulis pulveraceis, foliolis oblongis utrinquè scutis planis glabris pellucido-punctatis, racemis trichotomis folio brevioribus. § in Novâ-Holl. orientali. (v. s.)
3. *Z. LÆVIGATA* (Smith in Rees cycl. n. 2.) ramulis glabris, foliolis linearibus margine revolutis glabris petiolo longioribus, pedunculis bis trifidis corymbosis folio brevioribus. § in Novâ-Holl. orient. ad Port-Jackson. — *Z. lævigata* Bonpl. nav. p. 64 quæ pedunculis folii longitudine gaudet videtur diversa. (v. s.)
4. *Z. CYTISOIDES* (Smith in Rees cycl. n. 4.) ramulis villosis, foliolis obovatis utrinquè villosis, pedunculis trifidis foliiferis. § in Novâ-Cambriâ australi.
5. *Z. ARBORESCENS* (Sims bot. mag. 1395. in adn.) caule arboreo, foliolis subtus tomentosis impunctatis, pedunculis petiolo longioribus. § in Novâ-Holl.
6. *Z. MICROPHYLLA* (Bonpl. nav. p. 64.) ramulis sericeis foliolis linearibus margine revolutis supernè glabris subtus sericeis, pedunculis subtrifloris. § in Novâ-Hollandiâ.
7. *Z. PAUCIFLORA* (Smith in Rees cycl. n. 3.) ramulis petiolisque pilosis, foliolis linearibus interdum margine revolutis, pedunculis 1-3-floris, cal. lobis lanceolatis acutis. § ad Port-Jackson.
8. *Z. PILOSA* (Rudg. tr. lin. soc. 10. p. 293. t. 17. f. 2.) foliolis lanceolatis subtus pilosis, pedunculis 1-floris. § in Novâ-Cambriâ austr. Foliola 3-4-lin. longa.
9. *Z. HIRSUTA*, ramis petiolis capsulisque hirsutis, foliolis oblongo-linearibus petiolo triplò longioribus subtus hirsutis margine subrevolutis, pedunculis 1-floris. § in Novâ-Holl. montibus cæruleis. (v. s.)

XVI. MELICOPE. *Forst. gen. n. 28.*

Cal. 4-partitus persistens. Pet. 4, unguibus in tubum tetragonum approximatis, limbis patentibus. Glandulæ 4 ovaria cingentes. Stam. 8 subulata. Ovaria 4 ovata supernè mediante stylo unico connexa. Stigma centro concavum. Carpella 4 subcoriacea divaricata supernè dehiscentia. Semina in carp. solitaria. Embryo radicula brevi, cotyl. ovatis.

1. *M. TERNATA* (Forst. l. c. Lam. ill. t. 294.) § in Novâ-Zelandiâ. Entoganum lævigatum Gaertn. fr. 1. p. 331. t. 68.

XVII. ELAPHRIUM. *Jacq. am. 105.*

Cal. 4-partitus deciduus. Pet. 4. Stam. 8 calycis longitudine. Stylus 1 brevis. Stigma bifidum. Capsula (carpelli unius abortu?) subglobosa 1-locul. 1-sperma. Semen pulpâ vestitum. — Arbores. Folia alterna impari-pinnata 4-5-juga petiolo sæpius marginato; racemi plures in ramorum apicibus. — Genus non satis notum forsân, suadente cl. Kunthio ad Amyridem referendum.

* *Petiolo marginato. Americanæ.*

1. *E. TOMENTOSUM* (Jacq. amer. 105. t. 71. f. 1-3.) foliis utrinquè tomentosis, foliolis ovatis dentatis, racemis paucifloris folio duplò brevioribus. § in

- saxosis glareosis Curaçao insulisque vicinis. Lam. ill. t. 304. f. 1. *Fagara octandra* Lin. mant. 40.
2. *E. COPALLIFERUM* (fl. mex. ic. ined.) foliis pubescentibus, foliolis ovatis dentatis, racemis folii ferè longitudine interruptis, floribus confertis brevissimè pedicellatis. 5 in Mexico. — Copalli Hern. mex. 45. f. 1.
3. *E. GLABRUM* (Jacq. am. 106. t. 71. f. 4.) foliis glaberrimis, foliolis oblongo-lanceolatis obtusè dentatis, racemis paucifloris folio duplò longioribus. 5 in Carthagenæ fruticetis maritimis. — Lam. ill. t. 304. f. 2. *Fagara Elaphrium* Willd. sp. 1. p. 668.
4. *E. PENICILLATUM* (fl. mex. ic. ined.) foliis subglabris, foliolis ovali-oblongis acuminatis grossè dentatis, petiolo basi piloso et pilorum fasciculo ad originem jugorum ornato, racemis paucifloris folio triplò ferè brevioribus. 5 in Mexico.
- ** *Petiolo tereti. Capenses.*
5. *E? INÆQUALE*, petiolis puberulis teretibus, foliolis petiolulatis ovatis basi inæqualibus irregulariter crenatis pellucido-punctatis glabris, racemis paucifloris folio triplò brevioribus. 5 ad cap. B.-Spei. — Burch. cat. n. 3082 et 3051. Habitus omninò *Fagaræ armatæ* sed petioli inermes. An genus proprium? An forsàn *Fagaræ* aliæ Capenses huc revocandæ? (v. s. comm. ab amic. Burchell.)
6. *E? CAPENSE*, foliis pinnatis, foliolis ovatis obtusis erectis subsessilibus crenatis, ramis aculeatis flexuosis. 5 ad cap. B.-Spei. Flores paniculati albi 8-andri. *Fagara Capensis* Thunb. fl. cap. 1. p. 516. H. B. et Kunth nov. gen. am. 6. p. 6. in adn. An, ex Kunthio, *Choisyæ* species?
- XVIII. *CHOISYA*. H. B. et Kunth nov. gen. am. 6. p. 4. — *Plenckia fl. mex. ic. ined. non Rafin.*

Flores hermaphroditi. Sepala 5 decidua. Pet. 5 brevi unguiculata. Stam. 10, libera. Ovarium 5-loc., loculis biovulatis. Discus nullus. Stylus 1. Stigma capitato-5-lobum. Capsula ovata 5-rostrata 5-sulcata 5-loc. — Frutex inermis. Folia opposita 3-foliolata glanduloso-punctata. Flores albi fasciculato-congesti.

1. *C. TERNATA* (H. B. et Kunth l. c. t. 513.) 5 in Mexico.

XIX. *EVODIA*. Forst. gen. t. 7. H. B. et Kunth nov. gen. am. 6. p. 1 et 6 in adn. St.-Hil. bull. philom. 1823. p. 129. non Gaertn.

Flores hermaphroditi. Calyx 4-5-fidus vel 4-5-partitus. Pet. 4-5. Glandulæ totidem (ex Kunth) aut nectarium cyathiforme (ex St.-Hil.) inter stamina et ovaria. Capsula 4-cocca, coccis basi tantum connexis bivalvibus 1-2-spermis. — Caules fruticosi. Folia opposita, foliolis 5 integerrimis glanduloso-punctatis. Corymbi axillares dichotomi.

1. *E. LATIFOLIA*, foliolis ovatis acutis subtus lanuginosis, paniculis axillaribus petioli longitudine. 5 5 in Amboinâ. *Ampacus latifolia* Rumph. 2. p. 186. t. 61. Foliola 8-12 poll. longa.
2. *E. TRIPHYLLA*, foliolis ovatis acuminatis subtus glabris, paniculis lateralibus brachiato-pyramidalis petioli circiter longitudine. 5 5 in Amboinâ et Philippinis. *Fagara triphylla* Lam. dict. 2. p. 447. Roxb. fl. ind. 1. p. 436. *Ampacus angustifolia* Rumph. 2. p. 188. t. 62. Foliola 5-6 poll. longa.
3. *E. OBTUSIFOLIA*, foliolis ovalibus obtusis utrinquè glabris, paniculis axillaribus petiolo ferè brevioribus. 5 in insulis Mauritianis. (v. s.)
4. *E. HORTENSIS* (Forst. gen. 14. t. 7.) foliolis lanceolatis ramulisque pubescentibus, paniculis axillaribus foliolisque petiolo longioribus. 5 in insulis Amicorum et nov. Hebridis. *Fagara Evodia* Lin. f. suppl. 125.
5. *E. FEBRIFUGA* (St.-Hil. bull. philom. 1823. p. 129.) foliolis lanceolato-

ellipticis subacuminatis, paniculâ terminali pubescente, nectario simplici.
5 in Brasiliâ.

XX. ZANTHOXYLUM. H. B. et Kunth nov. gen. am. 6. p. 1. — Zanthoxylum et Fagara. Lin. gen. n. 150 et 1109. — Xanthoxylum Smith. — Tenorea Rafin. — Aubertia Bory. — Langsdorfia Leand. in akad. mûnch. 1819. p. 229. — Ponlana Nees et Mart. nov. act. bonn. xi. p. 185.

Flores hermaphroditi aut abortu dioici aut monoici. Calyx 3-lobus sæpius 4-5-partitus. Petala tot quot calycis lobi, et iis alterna, rarò nulla. Stam. tot quot calycis lobi, iis opposita. Carpella tot quot cal. lobi, interdum basi connata aut libera, abortu sæpè ad minorem numerum reducta imò interdum solitaria, matura bivalvia 1-3-sperma. Semen nitidum (Gaertn. fr. 1. t. 68.) — Frutices arboresce sæpè aculeati. Folia sæpius alterna, sæpè punctato-glandulosa. — Genus forsàn in posterum denuò dividendum, sed characteres ex numero partium derumpti non sufficiunt.

* *Foliis simplicibus.*

1. Z? AUBERTIA, inerme, foliis oppositis ovatis obtusis integerrimis emarginatisve pellucido-punctatis, floribus axillaribus racemosis 4-fidis. 5 in ins. Borboniâ planitie excelsâ dictâ d'Afouge. Aubertia Borbonica Boryl'voy. afr. 1. p. 356. t. 18. (v. s.)
2. Z. PENTANOME, aculeatum, foliis alternis ovalibus utrinquè acuminatis integerrimis, paniculis terminalibus, floribus 5-andris 5-gynis. 5 in Mexico. Pentanome simplicifolia fl. mex. ic. ined.
3. Z. OCHROXYLUM, trunco aculeato, foliis ovatis punctato-pellucidis, floribus 5-andris 3-gynis. 5 in ins. Caribæis. Ochroxylum et Curtisia Schreb. gen. n. 508. et emend. p. 826. Fagara monophylla Lam. ill. n. 1643. †.

** *Foliis sæpius trifoliolatis.*

4. Z. TERNATUM (Swartz fl. ind. occ. 1. p. 570.) inerme, foliis 3-foliolatis, foliolis obovatis submarginatis nitidis subtus punctatis, racemis axillaribus compositis. 5 in ins. Dominicâ. Fagara ternata Swartz prodr. 33. Pet. Stam. et carpella 3. (v. s.)
5. Z. MONOGYNUM (St-Hil. bull. philom. 1823. p. 129.) inerme, foliis 3-foliolatis, foliolis rotundo-ellipticis breviter acuminatis integerrimis glabris petiolo communi nervoque medio puberalis, paniculis terminalibus, pistillo unico. 5 in Brasiliâ.
6. Z. PUNCTATUM (West St. Cr. 236.) spinosum, foliis trifoliolatis pinnatisque oblongis crenulatis subtus punctatis. 5 in ins. Sanctæ-Crucis.

*** *Foliis omnibus impari-pinnatis.*

7. Z. CULANTRILO (H. B. et Kunth nov. gen. am. 6. p. 2.) inerme, foliis 3-foliolatis aut impari-pinnatis, foliolis oblongis submarginatis crenatis, petiolo obsolete alato, pedunculis axillaribus ramosis petiolo parum longioribus. 5 in ripâ fl. Amazonum prope Jaen-de-Bracamoros.
8. Z. TRAGODES, aculeatum, foliis impari-pinnatis, foliolis cuneiformibus emarginatis, petiolo latè alato in articulos foliola ferè referentes solubili. 5 in ins. Sancti-Domingi. Schinus tragodes Linn. spec. ed. 1. p. 369. Fagara tragodes Jacq. am. 21. t. 14. Hayn. term. bot. t. 14. f. 2. (v. s. sine fl.)
9. Z. PTEROTA (H. B. et Kunth nov. gen. am. 6. p. 3.) aculeatum, foliis impari-pinnatis, foliolis obovatis subcrenatis, petiolo angustè marginato, aculeis 2 stipularibus uncinatis. 5 in Jamaicâ et Cubâ. Fagara Pterota Linn. amœn. 5. p. 393. Schinus Fagara Linn. sp. ed. 1. p. 389. Fagara lentiscifolia Willd. enum. 1. p. 166. ex Kunth. Spicæ axillares petiolo breviores. Stam. 4-5. Ovaria 2. (v. s.)
10. Z. PIPERITUM, aculeatum, foliis impari-pinnatis, foliolis oblongis basi inæ-

- qualibus crenatis, petiolo subulato articulato, aculeis stipularibus. \S in Japoniâ. Koempf. am. t. 893. *Fagara piperita* Linn. sp. 172. Thunb. fl. jap. 64?
11. *Z. AFFINE* (H. B. et Kunth nov. gen. am. 6. p. 3.) sparsè aculeatum, foliis impari-pinnatis, foliolis 7-9 oblongis obtusis glabris margine crenatis et glanduloso-punctatis, petiolo hirtello nudo, spicis axillaribus multifloris. \S in Mexico ad lacum Cuiseo. Valdè priori affine.
12. *Z. AVICENNÆ*, aculeatum, foliis impari-pinnatis, foliolis 9-13 lanceolatis subintegerrimis glabris petiolulatis, racemis paniculatis folio brevioribus \S in Chinâ. *Fagara Avicennæ* Lam. dict. 2. p. 445. Lob. ic. 2. t. 133. f. 2. ic. fr.
13. *Z. HETEROPHYLLUM* (Smith in Rees cycl. n. 14.) aculeatum, foliis impari-pinnatis, junioris arboris longissimis aculeatis sub-20 jugis, adultæ brevioribus latioribus inermis sub 4-jugis, petiolo tereti. \S in ins. Borboniz circa Gol. *Fagara heterophylla* Lam. dict. 2. p. 445. Maqueria Comm. Flores hermaphroditi. (v. s.)
14. *Z. SENEGALENSE*, aculeatum, foliis impari-pinnatis, foliolis obovatis integerrimis subemarginatis, petiolis teretibus nervisque foliorum aculeatis. \S in Senegal. *Fagara zanthoxyloides* Lam. dict. 2. p. 446. Flores 5-andri dioici. (v. s.)
15. *Z. THUNBERGII*, aculeatum, foliis impari-pinnatis, foliolis sessilibus ovatis tenuissimè crenatis, petiolis aculeatis vix submarginatis. \S ad cap. B.-Spei. *Fagara armata*. Thunb. fl. cap. 141. An potius ad *Elaphria Capensis* referendum? †.
16. *Z. MARTINICENSE*, foliis impari-pinnatis aculeatis, foliolis oblongis integerrimis alternis, floribus dioicis 5-andris, stigmatè peltato. \S in Martinicâ. *Fagara Martinicensis* Lam. ill. n. 1659. †.
17. *Z. PERROTETII*, foliis impari-pinnatis 7-9-jugis, foliolis sessilibus elliptico-oblongis acuminatis apice dentatis, petiolo immarginato ramisque inermibus, paniculâ subcorymbosâ. \S in Guianâ Gallicâ. (Perrotet.) *Carpella* matura 2 aut uno abortivo solitaria. Fl. ign. (v. s.)
18. *Z. EMARGINATUM* (Swartz fl. ind. occ. 1. p. 572.) inerme, foliis pinnatis 2-3-jugis, foliolis ovatis emarginatis villosis, racemis terminalibus. \S in montibus Jamaicæ. *Fagara emarginata* Swartz prod. 33. — Sloan. hist. 2. p. 24. t. 168. f. 4. Flores 3-andri 3-gyni. Folia nunc abruptè nunc impari-pinnata.
19. *Z. ACUMINATUM* (Swartz fl. ind. occ. 1. p. 575.) inerme, foliis pinnatis 3-4-jugis, foliolis ellipticis acuminatis integris nitidis coriaceis, cymis terminalibus. \S in Jamaicæ montosis. Carp. 3 basi juncta. *Fagara acuminata* Swartz prod. 33. Flores 3-andri 3-gyni.
20. *Z. SPINOSUM* (Swartz fl. ind. 1. p. 574.) foliis pinnatis 8-10-jugis, foliolis sessilibus ovatis acuminatis subtus petiolis ramisque spinosis, cymis terminalibus. \S in aridis montosis Jamaicæ. *Fagara spinosa* Swartz prod. 33. Pet. et Stam. 3. Styli o. Stigmata 3. (v. s.)
21. *Z? CORYMBOSUM*, foliis pinnatis 2-jugis, foliolis ovatis sessilibus acutis glabris integris, costâ subtus petiolisque aculeatis, floribus corymbosis. \S in ins. Caribæis. *Tenorea corymbosa* Rafin. spech. 1. p. 193. *Carpella* polysperma. †.
22. *Z. PIMPINELLOIDES*, foliis pinnatis multijugis, foliolis subrotundo-acuminatis nitidis, petiolis costisque aculeatis, corymbis terminalibus. \S in Sancto-Domingo. — Sloan. hist. 2. t. 174. f. 3-4? *Fagara pimpinelloides* Lam. ill. n. 1638. Cal. 5-fidus. Styli 3.
23. *Z. TRICARPUM* (Michx. am. 2. p. 335.) foliis impari-pinnatis 3-5-jugis, foliolis petiolulatis oblongo-ovalibus acuminatis serrulatis basi obliquis, petiolis ramisque aculeatis, paniculis terminalibus. \S in Carolinæ et Floridæ sylvis. *Fagara fraxinifolia* Lam. ill. 1. p. 334. — Cat. car. 1. t. 26. — Carp. 3. Sep. 5. *Kampmannia* Rafin. (v. s.)
24. *Z. FRAXINEUM* (Willd. sp. 4. p. 757.) foliis impari-pinnatis 4-5-jugis, fo-

liolis ovatis obsolete serrulatis basi æqualibus, petiolis teretibus inermibus, aculeis stipularibus, umbellis axillaribus. 5 in Americâ bor. Z. ramiflorum Michx! fl. 2. p. 235. Z. Clava-Herculis var. Lin. sp. 1455. Lam. dict. 2. p. 38. Z. Americanum Mill. dict. n. 2. Z. Caribæum Gaertn. non Lam. — Duham. arb. 1. t. 97. Flores dioici apetal. (v. v.)

25. Z. MITE (Willd. enum. 1013.) inerme, foliis impari-pinnatis subtus pubescentibus, floribus axillaribus. 5 in Americâ bor.

26. Z. HERMAPHRODITUM (Willd. sp. 4. p. 756.) aculeatum, foliis pinnatis 5-jugis, foliolis oblongis acuminatis integerrimis basi subæqualibus, petiolis inermibus, paniculis terminalibus. 5 in sylvis Guianæ. Pet. et Stam. 5. Carp. et Styli 3-5. Fagara pentandra Aubl. guian. 1. p. 78. t. 30. F. Guianensis Lam. dict. 2. p. 446.

27. Z. RIGIDUM (Humb. in Willd. sp. 2. p. 756. et H. B. et Kunth nov. gen. am. 6. p. 4.) aculeatum, foliis pinnatis 4-jugis, foliolis ellipticis integerrimis emarginatis mucronatis, venis subtus hirsutis, costâ mediâ petiolisque aculeatis. 5 in Americâ merid. Flores ign.

28. Z. JUGLANDIFOLIUM (Willd. sp. 2. p. 756.) aculeatum, foliis pinnatis, foliolis oblongis acuminatis obsolete serratis basi inæqualibus subpunctatis, petiolo subaculeato, paniculis terminalibus. 5 in Sancto-Domingo. — Pluk. t. 239. f. 6. Foliola petiolulata alterna.

æ Berterianum, foliis sessilibus sæpius oppositis, racemis petiolisque pubescentibus. 5. (v. s. ex cl. Bertero.)

29. Z. LANCEOLATUM (Poir. suppl. 2. p. 293.) aculeatum, foliis pinnatis 6-7-jugis, foliolis elliptico-lanceolatis membranaceis impunctatis, petiolis aculeatis venisque subtus hirsutis, paniculis terminalibus. 5 in Porto-Ricco ubi vulgò Cenizo dicitur et in Guadalupâ: Bertero. (v. s.)

30. Z. CLAVA-HERCULIS (Lin. sp. 1455. excl. syn.) aculeatum, foliis pinnatis 4-jugis, foliolis ovatis repando-dentatis basi inæqualibus sessilibus pellucido-punctatis glaberrimis, paniculis terminalibus. 5 in sylvestribus ins. Caribæarum. Z. Caribæum Lam. non Gaertn. Z. Carolinianum Gaertn. — Pluk. t. 239. f. 4. (v. s. sine fl.)

31. Z. AROMATICUM (Willd. sp. 4. p. 755.) aculeatum, foliis pinnatis 6-jugis, foliolis ovato-lanceolatis serratis petiolulatis basi inæqualibus pellucido-punctatis glabris, paniculis terminalibus et axillaribus. 5 in Sancto-Domingo. Poiteau. — Jacq. f. ecl. 1. p. 103. t. 70. Petiolus variat inermis et aculeatus. (v. s.)

32. Z. RHÖIFOLIUM (Lam. dict. 2. p. 39.) aculeatum, foliis pinnatis multijugis, foliolis lanceolatis serrulatis basi subæqualibus, petiolo pubescente subaculeato, paniculis lateralibus. 5 in Indiâ orient. — Pluk. t. 329. f. 1.

33. Z. OBTUSIFOLIUM (Poir. suppl. 2. p. 293.) aculeatum, foliis pinnatis sub-4-jugis, foliolis coriaceis obversè subovatis apice rotundatis, subtus petiolisque aculeatis glabris, paniculâ terminali densâ. 5 in Indiâ orient?

34. Z. ARMATUM (Roxb! cat. calc. 72.) aculeis rectis patentibus armatum, foliis pinnatis bijugis, foliolis oblongis utrinquè acuminatis integris, petiolo inermi, paniculis terminalibus aut subaxillaribus. 5 in Indiâ orient. Fl. dioici. (v. s. comm. à cl. Lambert.)

35. Z. ACANTHOPODIUM, aculeis rectis patentibus armatum, foliis impari-pinnatis 4-jugis, foliolis oblongis utrinquè acuminatis, costâ, petiolisque validè aculeatis, corymbis axillaribus brevissimis. 5 in Napauli. Habitus prioris. (v. s. comm. à cl. Wallich.)

36. Z. NITIDUM, ramis petiolis costisque aculeatis, foliis impari-pinnatis 2-3-jugis, foliolis oblongis lucidis remotè glanduloso-crenatis, apice elongatis emarginatis, racemis axillaribus fasciculatis. 5 in Chinâ. Fagara nitida Roxb. fl. ind. 1. p. 440. Fagara piperita Lour. fl. coch. 1. p. 101. ex Wall. in fl. ind. l. c. cum descr.

**** *Foliis abruptè pinnatis.*

37. *Z. SAPINDOIDES*, foliis abruptè pinnatis, foliolis ovatis, caule ramisque spinosissimis. 5 in Jamaicâ. — Browne jam. 207. t. 20. f. 2. *Sapindus spinosus* Linn. sp. 526. Licca-Tree Lunan. hort. jam. 1. p. 443.

38. *Z. RHETSA*, foliis abruptè pinnatis, foliolis lanceolatis falcatis lævibus integris, aculeis confertissimis incurvis, paniculis terminalibus. 5 in Indiâ orient. Fagara Rhetsa Roxb. fl. ind. 1. p. 438. Pet. et Stam. 4. Carp. 1.

†. *Species non satis notæ.*

39. *Z. HORRIDUM*, foliis alternis impari-pinnatis, foliolis ovatis crenatis, spinis ramorum spinosis. 5 in Japoniâ. Fagara horrida Thunb. act. soc. lin. 2. p. 329. Flores fructusque ignoti.

40. *Z. SPINIFEX*, foliis alternis subfasciculatis impari-pinnatis, petiolo subalato, foliolis ellipticis integris subemarginatis, ramis spinosis. 5 ad Caracas. Fagara spinifex Jacq. fragm. p. 10. t. 6. f. 2. Flor. ign.

41. *Z. BUDRUNGA*, arborum aculeis parvis incurvis armatum, foliis pinnatis, foliolis 5-6-jugis inæqualiter ovatis lanceolatis integris acuminatis, paniculis contractis. 5 in Indiâ orient. Fagara Budrunga Roxb. fl. ind. 1. p. 437. Pet. et stam. 4. Carpellum 1. An folia abruptè aut impari-pinnata?

42. *Z. ZEYLANICUM*, foliis , capsulis bilocularibus. 5 F. Lonn-Ankenda Gaertn. fr. 1. p. 334. t. 68. f. 9. Fagara Zeylanica Steud. nom. phan. 331.

43. *Z? OPPOSITIFOLIUM*, foliis oppositis impari-pinnatis 5-6-jugis ramisque inermibus glabris, petiolo subalato, foliis oblongo-ovalibus obtusis apice dentatis. 5 in Novâ-Hollandiâ. (v. s. sine fl. ex Mus. Par.)

XXI. *PILOCARPUS*. Vahl eclog. 1. p. 29. St.-Hil. bull. philom. 1823. p. 130. Nees et Mart. nov. act. bonn. xi. p. 176. t. 19. f. 1.

Cal. minimus 5-part. Pet. 5 basi latiuscula, apice uncinata. Stam. 5. Styli 5 breves, infra apicem ovariorum orti, apice coaliti in stigma 5 lobum. Carpella 5 minima, 1-sperma, approximata, gynophoro basi immersa et cum eo ovarium unicum simulantia. Coccæ 5 aut abortu pauciores bivalves, endocarpio crustaceo separabili. Cotyl. magnæ breviter biauriculatæ. — Frutices, foliis alternis simplicibus integerimis.

1. *P. RACEMOSA* (Vahl l. c. t. 10.) glaberrima, foliis confertis ellipticis apice rotundatis, floribus in racemum terminalem dispositis 5 in insulæ Montserrat montibus. — Plum. ed. Burm. p. 119. t. 127. Rami penduli. Ovula in carpellis 2 ex St.-Hil.

2. *P. SPICATA* (St.-Hil. l. c. p. 131.) glaberrima, foliis oblongis lanceolatis vel elliptico-lanceolatis acutiusculis, floribus spicatis subapproximatis brevissimè pedicellatis. 5 in Brasiliæ sylvis. P. parviflorus Nees et Mart. nov. act. bonn. xi. p. 177. t. 30.

3. *P. PAUCIFLORA* (St.-Hil. l. c.) foliis lanceolatis acutis vel acuminatis, petioliis incrassato-geniculatis, racemis paucifloris, rachi pedicellis bracteolisque puberulis. 5 in Brasiliâ.

XXII. *SPIRANTHERA*. St.-Hil. bull. philom. 1823. p. 130. — *Terpnanthus* Nees et Mart. nov. act. bonn. xi. p. 152 et 177. t. 19. f. K.

Cal. brevis cupulæformis profundè 5-dent. 5-gonus. Pet. 5 longa linearia angusta subfalcata, in unguem latiusculum attenuata. Stam. 5 petalis alterna, filam. filiformibus, anth. linearibus basi bifidis post anthesin spiraliter revolutis, connectivo crassiusculo. Styl. 1. Stigma 5-lobum. Nectarium cylindricum campanulatum ovarii stipitem cingens. Ovarium 5-lobum 5-loc., loculis 2-spermis, ovulo sup. adscen-

dente, inf. suspenso. — Frutex ramis patentibus, foliis 3-foliolatis, foliolis ovato-acuminatis crassiusculis subtus glaucis, floribus paniculato-corymbosis albo-roseis.

1. *S. ODORATISSIMA* (St.-Hil. l. c.) ♂ in Brasiliâ. *Terpnanthus jasminiodorus*. Nees et Mart. nov. act. bonn. xi. p. 178. t. 31.

XXIII. *ALMEIDEA*. St.-Hil. bull. philom. 1823. p. 129. — *Aruba* Nees et Mart. nov. act. bonn. xi. p. 152 et 172. t. 19. f. H. non Aubl. ex St.-Hil.

Cal. minimus 5-partitus deciduus. Pet. 5 calyce multoties longiora unguiculata spathulata æqualia erecta. Stam. 5 petalis alterna, fil. complanatis supra medium barbatis, anth. linearibus basi bifidis. Nectarium cupulæforme ovarium cingens. Styl. 1. Stigm. obtusè 5-lobum. Carpella 5 biovulata, (ovulo super. adscendente, inf. suspenso) axi centrali adfixo, demum abortu pauciora, libera, 1-sperma, ad umbilicum membranacea, parte membr. ruptili umbilico sem. adhærente arillæformi. Sem. reniforme. Embr. curvatus. Cotyled. magnæ, biauriculatæ, corrugatæ. Mucilago vix ulla inter cotyl. plicas. — Arbusculæ Brasilienses, foliis simplicibus sparsis petiolatis nitidis glanduloso-punctatis, floribus racemosis.

2. *A. LILACINA* (St.-Hil. l. c.) foliis lanceolatis, floribus thyrsoides, pedunculo puberulo, petalis obtusis. 5 in Brasiliâ.

2. *A. RUBRA* (St.-Hil. l. c.) foliis lanceolatis oblongis, floribus racemosis, pedunculo glabro, petalis obtusissimis. 5 in Brasiliâ.

3. *A. LONGIFOLIA* (St.-Hil. mss.) foliis lanceolatis oblongis, floribus racemosis, pedunculo pubescente, petalis obtusis. 5 in Brasiliâ.

4. *A. CÆRULEA* (St.-Hil. mss.) foliis oblongis basi attenuatis apice obtusis interdum emarginatis, racemis terminalibus, pedunculo glabro, petalis obtusis. 5 in Brasiliæ sylvis primævis ad fl. Ilhéos. *Aruba cærulea* Nees et Mart. nov. act. bonn. xi. p. 174. t. 27.

5. *A. ALBA* (St.-Hil. mss.) foliis obovato-cuneiformibus cuspidè obtuso, racemis terminalibus bifidis secundis, pedunculo nudo, petalis obtusis. ♂ in Brasiliâ ad viam Felisbertiam. *Aruba alba* Nees et Mart. nov. act. bonn. xi. p. 175. t. 28.

6. *A. ACUMINATA* (St.-Hil. mss.) foliis oblongo-lanceolatis utrinquè acutis, floribus terminalibus racemoso-paniculatis, pedunculo pubescente, petalis obtusis. 5 in Brasiliæ sylvis primævis ad præsidium S.-Joh.-Bapt. prov. Minas gener. Flores lilacini.

Trib. II. *CUSPARIÆ*. DC. mem. mus. 9. p. 141.

Petala sæpius inæqualia in corollam pseudo-gamopetalam approximata coalitave, nunc omnino coadunata, nunc lateribus tantum subadglutinata. Stamina varia, sæpè nonnulla sterilia. Urceolus carnosus ovarii basin cingens, stamina nec petala gerens. Stylus 1. Carpella 1-sperma. Semina (ex St.-Hil.) exalbuminosa. — Frutices rariùs herbæ.

XXIV. *MONNIERA*. Lin. gen. n. 850. Nees et Mart. nov. act. bonn. xi. p. 162. t. 18. f. C. — *Aubletia Pers. ench. 2. p. 638. non alior.*

Cal. 5-partitus, lobis dissimilibus. Pet. 5 in corollam irregularem bilabiata coalita. Stam. 5, filam. monadelphis tubo adnatis, 2 fertilibus, 3 sterilibus. Styl. 1. Carpella 5 aut abortu pauciora, basi subcoalita, bivalvia, 1-sperma, intus cocculifera. Seminis albumen tenue subcor-

neum, embryo inversus, radicula inflexa, cotyl. foliaceis, ex cl. Kunthio, reniformibus bifidis complicatis. — Herba villosa, foliis alternis 3-foliolatis, foliolis ovalibus utrinque acuminatis, cymis subterminalibus bifidis.

1. *M. TRIFOLIA* (Lin. sp. 986. Aubl. guian. 2. p. 730. t. 293. H. B. et Kunth nov. gen. am. 6. p. 9.*) (1) in fruticetis Cayennæ et Brasiliz. Jaborandi Marcg. bras. 36. ic. (v. s.)

XXV. *TICOREA*. Aubl. guian. 2. p. 689. DC. mem. mus. 9. p. 144. St.-Hil. bull. philom. 1823. p. 132. — *Ozophyllum* Schreb. gen. n. 1105. — *Sciuris* Nees et Mart. nov. act. bonn.

Cal. 5-dentatus. Pet. 5 in corollam monopetalam infundibuliformem adglutinata, tubo longo, limbo patente 5-fido. Stam. 5-8, quorum 2-6 sæpè sterilia connectivo sæpius producto, filamentis monadelphis tubo adglutinis, antheris liberis non appendiculatis. Stylus 1. Stigma 5-lobo-capitatum. Ovarium 5-lobum, loculis 2-spermis, ovulo sup. adscendente, inferiore suspenso. Urceolus carnosus truncatus ad basin ovarii. Carpella 5 obtusa, dorso carinata, reticulato-nervosa, endocarpio separabili, insidentia super stipitem crassum apice in discum 5-radiatum expansum. — Frutices glabri. Folia alterna, trifoliata, foliolis ovali-oblongis apice acuminatis. Corymbi pedunculati. Flores albi.

1. *T. FORTIDA* (Aubl. guian. 2. p. 689. t. 277.) foliolis 3 æqualiter subsessilibus, corymbo 6-7-floro, floribus secus ramos sessilibus, petalis calyce sextuplò longioribus. 5 in Guianæ sylvis. *Ozophyllum* foetidum Mart. in Mill. dict. O. trifoliolatum Willd. sp. 3. p. 585.
2. *T. PEDICELLATA* (DC. mem. mus. 9. p. 145. t. 8.) foliolis 3 æqualiter subsessilibus, corymbo 12-15-floro, floribus secus ramos pedicellatis, carpellis stipite insidentibus. 5 in Guianâ-Galicâ. (v. s. ex h. Mus. Par.)
3. *T. LONGIFLORA* (DC. mem. mus. 9. p. 145. t. 9.) foliolis lateralibus subpetiolatis, medio petiolo triplo longiore donato, corymbo 15-20-floro, floribus secus ramos subsessilibus, petalis calyce decies et ultra longioribus. 5 in Guianâ Gallicâ. (v. s. ex h. Mus. Par.)
4. *T. JASMINIFLORA* (St.-Hil. bull. philom. 1823. p. 132.) foliolis 3 ovato-lanceolatis acuminatis in petiolum attenuatis, paniculis laxiusculis staminibus 2-6 fertilibus. 5 in Brasiliâ prope Rio-Janeiro. *Sciuris* multiflora Nees et Mart. nov. act. bonn. xi. p. 155. t. 18. f. A.
5. *T. FEBRIFUGA* (St.-Hil. mss.) caule sæpius arboreo, foliis ternatis, foliolis lanceolatis acuminatis in petiolum attenuatis, paniculis coarctatis, staminibus 3-6 sterilibus. 5 in Brasilizæ minas geraes.
6. *T. BRACTEATA* (St.-Hil. mss.) foliolis 3 oblongis acuminatis basi attenuatis, panicula subspicata, multiflorâ glomeratâ erectâ, calyce 5-fido lobis acuminatis, stam. 2 fertilibus. 5 in Brasilizæ sylvis ad Arassatiba. *Sciuris* bracteata Nees et Mart. nov. act. bonn. xi. p. 156. t. 18. f. A. et t. 20. Bractezæ sub paniculæ ramis lanceolatæ subciliatæ.
7. *T. SIMPLICIFOLIA* (St.-Hil. mss.) foliis simplicibus oblongo-lanceolatis utrinque acuminatis racemis compositis multifloris erectis, stam. 2 fertilibus. 5 in Brasilizæ sylvis ad presidium S. Joh.-Bapt. Minarum generalium. *Sciuris* simplicifolia. Nees et Mart. nov. act. bonn. xi. p. 157.*

XXVI. *GALIPEA*. Aubl. guian. 2. p. 662. St.-Hil. bull. philom. 1823. p. 131. — *Galipea* et *Cusparia* DC. mem. mus. 9. 142 et 148. — *Cusparia* Humb. — *Bonplandia* Willd. non Cav. — *Angostura* Ræm. et Schult. — *Conchocarpus* Mik. Obontonia Vel.

Cal. brevis 5-dentatus. Pet. 5 in corollam hypocrateriformem coacta, seu valdè approximata, tubo brevi pentagono, lobis patentibus

acutis. Stam. 4-7 hypogyna, petalis subadhærentia, inæqualia, interdum omnia fertilia, sæpius 2 majora antherifera, 2-5 breviora sterilia. Nect. cupuliforme. Styli 5 in unicum mox coaliti et stigma 4-5-sulcum constituentes. Carpella 5 aut abortu pauciora biovulata obtusa cocculiformia sessilia, endocarpio separabili. Semina abortu solitaria. Cotyled. magnæ corrugatæ biauriculatæ. Frutices glabri; folia alterna simplicia aut plurifoliolata, foliolis oblongis acuminatis; pedunculi axillares multiflori.

* *Foliis compositis.*

1. *G. TRIFOLIATA* (Aubl. guian. 2. p. 662. t. 269.) foliis 3-foliolatis glabris, floribus corymbosis, pedunculo foliis brevior, calyce 5-dentato, staminibus sterilibus 2. $\frac{5}{2}$ ad ripas fluviorum Guianæ. Flores exigui virescentes.
2. *G. OSSANA* (DC. mem. mus. 9. p. 149. t. 10.) foliis 3-foliolatis punctatopellucidis glabris, floribus paniculatis, pedunculo foliis longiore, calyce 5-partito, staminibus sterilibus 5 $\frac{5}{2}$ circa Havanam. (v. s. comm. à cl. de la Ossa.)
3. *G. LASIOSTEMON* (St.-Hil. mss.) foliis 3-foliolatis glanduloso-punctatis, junioribus petiolis ramulisque pubescentibus, racemis subterminalibus erectis pubescentibus, petalis intus villosis extris calyceque pubescentibus, staminibus ciliato-villosis 3 fertilibus, 2 sterilibus. $\frac{5}{2}$ in Brasiliæ sylvis ad Ilhéos. *Lasiostemum* sylvestre Nees et Mart. nov. act. bonn. xi. p. 171. t. 19.
4. *G. CUSPARIA* (St.-Hil. mss.) foliis 3-foliolatis, racemis pedunculatis subterminalibus, calyce 5-dentato, staminibus sterilibus 3. $\frac{5}{2}$ in Americâ merid. *Cusparia* febrifuga Humb. tabl. geog. Bonplandia trifoliata Willd. act. acad. ber. 1802. p. 24. H. et B. pl. eq. 2. p. 59. t. 57. Kanth nov. gen. am. 6. p. 8. *Angostura* Cuspare Roem. et Schult. syst. 4. p. 183. *Cortex Angusturæ* offic.
5. *G. HETEROPHYLLA* (St.-Hil. bull. philom. 1823. p. 131.) foliis 3-4-5-foliolatis longè petiolatis, foliolis lanceolatis nervo medio subpubescente, racemis suprâ-axillaribus longè pedunculatis, stam. 2 sterilibus. $\frac{5}{2}$ in Brasiliâ.

** *Foliis simplicibus.*

6. *G. PENTAGYNA* (St.-Hil. mss.) foliis longis lanceolatis acutissimis glaberrimis, racemis in apice ramulorum axillaribus vel subextrâ-axill. compositis, pedunculis complanatis, stam. 3 sterilibus, stylis 5 distinctissimis. $\frac{5}{2}$ in Brasiliâ.
7. *G. RESINOSA* (St.-Hil. mss.) foliis oblongis utrinquè attenuatis pellucido-punctatis nitidis, petiolis glandulis prominentibus exasperatis, corymbis terminalibus pedunculatis, corollis subbilabiatis, stam. 5, fertilibus 2. $\frac{5}{2}$ in Brasiliâ ad viam Felisbertiam. *Ravja resinosa* Nees et Mart. nov. act. bonn. xi. p. 159. t. 19. f. E. et t. 23.
8. *G. PENTANDRA* (St.-Hil. l. c.) foliis lanceolatis acuminatis obtusis glabris, racemis axillaribus simplicibus paucifloris, pedicellis 3-bracteatis, stam. 5 fertilibus. $\frac{5}{2}$ in Brasiliâ.
9. *G. FONTANESIANA* (St.-Hil. l. c.) foliis lanceolatis oblongis acutiusculis glaberrimis, racemis terminalibus vel subextrâ-axillaribus basi vix ramosis, stam. 3 sterilibus, nectario 5-dentato. $\frac{5}{2}$ in Brasiliâ ad viam Felisbertiam. *Ravja racemosa* Nees et Mart. nov. act. bonn. 9. p. 169. t. 24. Flores albi.
10. *G. CANDOLLIANA* (St.-Hil. l. c.) foliis lanceolatis acuminatis glaberrimis, racemis subextrâ-axillaribus brevissimis, floribus confertis, stam. 3 sterilibus, nectario integro. $\frac{5}{2}$ in Brasiliâ.
11. *G. MACROPHYLLA* (St.-Hil. mss.) foliis ovali-oblongis petiolatis basi rotundatis subtus glandulis nigris opacis punctulatis, trunco simplici tenuissimo, florum racemis compositis interruptis, stam. 7, fertilibus 2. $\frac{5}{2}$ in Brasiliæ sylvis super montes graniticos circa Rio-Janeiro et in prov. Minarum. *Conchocarpus macrophyllus* Mikan. del Bras. 1. t. 2. Nees et Mart. nov. act.

- bonn. xi. p. 160. t. 18. f. B. *Raputia Conchocarpus* Schult. mant. 1. p. 126. (v. s.)
 12. G. *CUNEIFOLIA* (St.-Hil. mss.) foliis subpetiolatis lato-lanceolatis basi longè
 attenuato-cuneatis minutè pellucido-punctatis, racemis axillaribus elongatis
 interruptis folii longitudine, stam. 5, fertilibus 2. ♂ in Brasiliæ sylvis ad fluv.
 Peruaguacei et in prov. Bahiensi. Flores parvi albi.

XXVII. *ERYTHROCHITON*. Nees et Mart. nov. act. bonn. xi. p. 165. t. 18. f. D.

Cal. compresso-tubulosus 5-costatus bifidus, dentibus nempe in
 labia 2 concretis. Pet. 5 in cor. subhypocrateriformem concreta,
 limbo patente. Stam. 5 omnia fertilia, filam. basi in tubum carnosum
 connata. Styli 5 in 1 filiformem connata, stigmatè 5-sulcato. Carpella
 5 aggregata rugosa bivalvia 2-sperma. — Arbor parva caudice sim-
 plici, foliis alternis simplicibus oblongo-lanceolatis acutis basi at-
 tenuatis, pedunculis trigonis longis apice racemoso-spicatis. Flores magni
 bracteati cal. rubris, cor. albis.

1. E. *BRASILIENSIS* (Nees et Mart. l. c. p. 166. t. 22.) ♀ in Brasiliâ ad præsi-
 dium S.-Joh.-Bapt. prov. Minarum.

XXVIII. *DIGLOTTIS*. Nees et Mart. nov. act. bonn. xi. p. 170. t. 19. f. F.

Cal. campanulatus 5-fidus. Pet. 5 in corollam tubulosam ad medium
 usque cohærentia æqualia, limbo erecto. Stam. 5 breviter corollæ subad-
 hærentia, 2 fertilia apice appendiculata, 3 sterilia. Carpella 5 cupulâ
 carnosâ basi cincta 1-ovulata. Stylus brevissimus. — Frutex comâ
 densâ, foliis sparsis simplicibus obovato-oblongis apice rotundatis
 pellucido-punctatis, racemo brevi terminali subpaniculato.

1. D. *OBOVATA* (Nees et Mart. l. c. t. 25.) ♂ in sylvis Brasiliæ ad fluv. Xipoto.
 Calyx pubescens. Pet. acuta 3 lin. longa.

†. *Genera Rutaceis affinia non satis nota.*

XXIX. *BARRALDEIA*. Pet. Th. nov. gen. mad. p. 24. — *Baraultia* Steud.
 nom. phan. 101.

Cal. urceolaris 5-fidus. Pet. 5 minima bifida unguiculata laciniis
 calycinis inserta. Stam. 10. Filam. basi dilatata quorum 5 petalis op-
 posita longiora. Circulus glandulosus extrâ pistillum. Ovarium calycis
 tubo adhærens aut in eo immersum. Stylus 1 staminibus longior.
 Fructus ign. . . . — Frutex ramis oppositis articulatis. Folia oppo-
 sita glaberrima pellucido-punctata! subdentata. Pedunculi breves bi-
 tri-fidi; flores parvi globulosi. Alabastrum abundè resinoseum — Cl.
 autor ad *Rhamneas* hoc genus refert, sed ex foliis punctato-pellucidis et
 stam. 10 potius *Rutaceum* videtur, forsân *Calodendro* affine.

1. B. *MADAGASCARIENSIS*. ♂ in Madagascar.

XXX. *HORTIA*. Vand. in Rœm. script. bras. 188.

Cal. 5-dentatus persistens. Pet. 5 reflexa acuminata intus ad basim
 glanduloso-pilosa. Stam. 5 petalis alterna disco hypogyno? inserta.
 Ovarium 1 subrotundum 5-loc. 5-ovulatum. Stylus 0 aut brevissimus.
 Stigma capitatum. Caps. 5-loc., loculis 1-spermis. — Arbor, foliis
 alternis subcuneiformibus integris punctato-pellucidis!, pedunculis
 corymbosis. — Genus huc ex foliis punctatis revocatum, sed pericarpia
 non satis et semina minimè cognita.

1. H. *BRASILIANA* (Vand. l. c.) ♀ in Brasiliâ. (v. s. in h. Juss. specimen à
 Vandellio ipso sine nomine missum.)

Ordo LII. SIMARUBEÆ. *Rich. anal. fr. p. 21. DC. diss. ochn. ann. mus. 17. p. 323.*

Flores hermaphroditi aut rariùs abortu diclini. Calyx 4-5-partitus, persistens. Petala 4-5 hypogyna, erecta, decidua. Stam. petalis numero æqualia aut dupla, disco hypogyno inserta, libera. Ovarium in tot partes lobatum quot sunt petala. Stylus 1 filiformis basi ampliatus. Carpella tot quot petala axi articulatim inserta, capsularia, bivalvia, intus dehiscentia monosperma. Semina exalbuminosa pendula. Cotyledones 2 crassæ. Radicula brevis supera.

Arbores aut frutices è tropicis novi orbis regionibus ortæ, cortice amarissimo, succo lacteo, foliis alternis pinnatis exstipulatis.

Ordo hinc Rutaceis, illinc Ochnaceis affinis; ab Ochnaceis differt semine inverso, carpellis dehiscentibus, foliis pinnatis. Cl. Saint-Hilaire Simarubeas ut Rutacearum tribum propriam admittit forsàn non immeritò.

I. QUASSIA. *DC. l. c. p. 324.*

Flores hermaphroditi. Calyx parvus 5-partitus. Petala 5 erecta calyce triplò longiora. Stam. 10. Filam. basi interiore squamulà aucta. Stylus simplex.

1. Q. AMARA (Lin. f. suppl. 235.) 5 in Surinamo. Lodd. bot. cab. t. 172. Arbor glabra, foliis impari-pinnatis, petiolis alatis, racemis terminalibus, floribus magnis rubicundis. (v. s.)

II. SIMARUBA. *Aubl. guian. 2. p. 856.*

Flores abortu monoici aut polygami. Calyx parvus 5-partitus. Petala 5 calyce paulò majora. Stam. 5-10 basi squamulis aucta. Stylus apice partitus.

1. S. OFFICINALIS (DC. l. c. n. 1.) floribus monoicis, masculis decandris, stigmate 5-partito, foliis abruptè pinnatis, foliolis alternis subpetiolulatis subtus pubescentibus. 5 in arenosis Guianæ et ins. Caribæarum. Quassia Simaruba Linn. suppl. 234. Lam. ill. t. 343. f. 2. Simaruba amara Aubl. guian. t. 331. 332. (v. s.)
2. S. GLAUCA (DC. l. c. n. 2.) floribus monoicis, masculis decandris?, stigmate 5-partito, foliis abruptè pinnatis, foliolis alternis subpetiolulatis glabris glaucis. 5 in ins. Cubæ maritimis Humb. Bonpl. et Kunth nov. gen. am. 6. p. 16.
3. S. EXCELSA (DC. l. c. n. 3.) floribus polygamis pentandris paniculatis, stigmate trifido, foliis impari-pinnatis, foliolis oppositis petiolulatis. 5 in sylvis Jamaicæ. Quassia excelsa Swartz act. holm. 1788. p. 302. t. 8. Q. polygama Wright. act. edinb.

III. SIMABA. *Saint-Hil. bull. philom. 1823. p. 129.* — Simaba et Aruba. *Aubl. guian. 1. p. 400 et 293.* — Zwingera. *Schreb. gen. n. 1751.*

Flores hermaphroditi. Cal. parvus cupularis 5-part. vel 5-fidus 5-

dentatusve. Pet. 5 basi lata, calyce multoties longiora. Stam. 10, rarius 8, alia alterna petalis opposita, filam. apice glabris subulatisque in squamam villosam intus dilatatis. Styli 5 in 1 coaliti. Stigm. 5 brevia. Carpella 5 monosperma. — Frutices Americani.

1. *S. GUIANENSIS* (Aubl. guian. 1. p. 400. t. 153.) foliis impari-pinnatis, foliolis 1-2-jugis ovali-oblongis utrinque acuminatis apice emarginatis, racemis axillaribus. ♂ in Guianæ sylvis. Zwingera amara Willd. sp. 2. p. 569. Flores sæpius 4-fidi. (v. s.)
2. *S. ORINOCENSIS* (H. B. et Kunth nov. gen. am. 6. p. 18. t. 514. a et b.) foliis nunc 3-foliolatis, foliolis ex apice petioli æqualiter ortis, nunc abruptè pinnatis 1-3-jugis, foliolis oblongis obtusis, racemis terminalibus. 5 in arenosis calidis ad Orinocum prope Carichana.
3. *S. FLORIBUNDA* (Saint-Hil. bull. philom. 1823. p. 129.) foliis impari-pinnatis, foliolis lanceolato-ellipticis oblongis obtusiusculis glabris, paniculâ magnâ compositâ. ♂ in Brasiliâ.
4. *S. FERRUGINEA* (Saint-Hil. l. c.) foliis impari-pinnatis, foliolis ellipticis pubescentibus subtus nervosis, paniculâ compositâ sessili foliis brevior. ♂ in Brasiliâ.
5. *S. STAVEOLENS* (Saint-Hil. l. c.) foliis abruptè pinnatis, superioribus interdum simplicibus, foliolis ellipticis vel subrotundo-ellipticis glabris, floribus terminalibus racemosis, racemis compositis. 5 in Brasiliâ.
6. *S. TRICHILIOIDES* (Saint-Hil. l. c.) foliis impari vel abruptè pinnatis, foliolis ellipticis obtusissimis apice mucronatis nervosis supra pubescentibus subtus subtomentosus, paniculâ simplici folio multo majore. ♂ in Brasiliâ.
7. *S. ARUBA* (Saint-Hil. mss.) foliis 3-foliolatis, foliolis lanceolatis acuminatis, racemis terminalibus. 5 in Guianâ-Gallicâ. Aruba Guianensis Aubl. guian. 1. p. 293. t. 115.

IV? *RAPUTIA*. Aubl. guian. 2. p. 670. — *Pholidandra*. Neck. elem. n. 542. — *Sciuris*. Schreb. gen. n. 53.

Calyx brevis, 5 fidus. Pet. 5 in corollam tubulosam 5-fidam subbilabiatam coalita. Stam. 5 tubo adnata, 3 sterilia basi villosa, 2 antherifera basi bisquammosa. Stylus 1. Ovarium et stigma 3-5-lobum. Carpella 5 subrotunda bivalvia 1-sperma. Cotyled. seminis 2 crassæ. — Suffrutex glaber aromaticus. Folia opposita, 3-foliolata, foliolis punctato-pellucidis, ovali-oblongis obtusè acuminatis. Spicæ axillares. — Genus omnino medium inter Simabam et Galipeam. An Simarubeis, an Rutaceis adjungendum? Potius Simarubeis ob seminis fabricam si ab Aubletio rite descriptam.

1. *R. AROMATICA* (Aubl. l. c. t. 272. Lam. ill. t. 10.) foliolis 3 æqualiter sessilibus pellucido-punctatis. ♂ in Guianæ sylvis. Spicæ multifloræ, floribus distichis.
2. *R? HETEROPHYLLA* (DC. mem. mus. 9. p. 153.) foliis subtus glandulosis, aliis 1-foliolatis, aliis 3-foliolatis, foliolo medio longè petiolulato. 5 in Porto-Ricco. (Bertero.) (v. s. sine fl. et fr.)†.

Ordo LIII. OCHNACEÆ. DC. ann. mus. 17. p. 398. rec. mem. 1813. n. 5.

Calycis sepala 5 basi vix coalita persistentia æstivatione imbricata. Pet. hypogyna numero definita, nempe 5 sepalis alterna, rarius 10, caduca, patula, æstivatione imbricata. Stam. 5 petalis alterna, aut 10, aut indefinita, disco hypogyno inserta. Filam. sæpius persistentia. Antheræ biloculares, basi insertæ. Ovarium partitum in tot loculos quot sunt petala. Stylus 1 filiformis persistens basi ampliatus in discum subglobosum carnosum ovarii loculamenta gerentem et *gynobasim* dictum. Pericarpium loculamenta seu carpella tot quot petala, gynobasi grandifactæ articulatum inserta, circa stylum verticillata, 1-sperma, indehiscencia, subdrupacea. Semina exalbuminosa, embryone recto, radiculâ brevi, cotyledonibus 2 crassis.

Arbores fruticesve inter Tropicos crescentes glaberrimæ, succo aqueo foetæ. Folia alterna simplicia penninervia integra aut dentata, basi bistipulata. Flores racemosi, pedicellis medio aut infra medium articulatis, raro solitarii.

I. OCHNA. Schreb. gen. n. 354. — Ochnæ sp. Lin.

Petala 5-10. Stam. ∞, filam. filiformibus persistentibus, antheris linearibus ovatisve rimâ duplici ab apice ad basim dehiscentibus. — Gemmæ squamosæ. Flores infra folia è præcedentis anni ligno orti.

1. O. *OBTUSATA* (DC. l. c. n. 1. t. 1.) stigmatē capitato, floribus 8-10-petalis, foliis obovatis obtusissimis serratis. 5 in Indiâ orient. O. squarrosa Lin. sp. 781? (v. s.)
2. O. *LUCIDA* (DC. l. c. n. 2.) stigmatē capitato, floribus 7-10-petalis, foliis obovatis aut oblongo-ovatis acutis serratis. 5 in Indiâ orient.
 - a. foliis obovatis. O. lucida Lam. ill. t. 472. f. 1. (v. s.)
 - β. foliis oblongo-ovatis. O. squarrosa Rottb. act. dan. 2. p. 545. t. 6. excl. syn.
3. O. *NITIDA* (Thunb. prod. 67.) stigmatē capitato, floribus 5-petalis, foliis oblongis acutis serratis, racemis brevibus confertis. 5 in Indiâ orient. DC. l. c. n. 3. t. 2. (v. s. in h. Deless.)
4. O. *MULTIFLORA* (DC. l. c. n. 4. t. 3.) stigmatē capitato, floribus 5-petalis, foliis ovali-oblongis acuminatis subintegerrimis, racemis pedicellisque longissimis. 5 in Sierrâ-Leonâ. (v. s.)
5. O. *CILIATA* (Lam! dict. 4. p. 511.) stigmatē capitato breviter lobato, floribus 5-petalis, foliis ovali-oblongis ciliato-serratis, racemis brevibus paucifloris. 5 in Madagascar. DC. l. c. n. 6. t. 4. (v. s.)
6. O. *MADAGASCARIENSIS* (DC. l. c. n. 7.) stigmatē multipartito, floribus pentapetalis, foliis oblongis nitidis subserratis, petalis calyci æqualibus. 5 in insulâ Madagascar. (v. s. in h. Pet. Th.)
7. O. *MAURITIANA* (Lam! dict. 4. p. 522.) stigmatē multipartito, floribus 5-6-

†. *Genera Ōchnaceis affinia.*IV? ELVASIA. DC. *diss. och. n. iv.*

Cal. 4-partitus. Pet. 4. Stam. 8. Filam. longiusculis, antheris ovatis per rimas duas dehiscentibus. Pericarpium locul. 4. An hujus ordinis? †.

1. E. CALOPHYLLA (DC. l. c. n. i. t. 20.) 5 in Brasiliâ. Folia oblonga integerrima medio nervo et nervulis parallelis approximatis more Calophylli donata. (v. s. in h. Mus. Par.)

V? CASTELA. Turp. *ann. mus. 7. p. 78. t. 5.*

Cal. parvus 4-dentatus. Pet. 4 dentibus calycinis alterna, disco perigyno inserta. Stam. 8 cum petalis inserta, eorum longitudine. Ovarium 4-lobatum disco circulari impositum. Stylus 1. Lobi 4 drupacei 1-spermi è gynobasi carnosâ orti. Semen inversum. Radicula supera. Cotyl. 2 foliaceæ. Albumen carnosum. — Fruticuli, ramis alternis apice spinescentibus, foliis alternis integris, floribus parvis axillaribus. — An ordo proprius inter calycifloras, antè Celastrineas inserendus?

1. C. DEPRESSA (Turp. l. c. p. 79. t. 5. A.) foliis ovali-oblongis basi subcordatis sessilibus, spinis axillaribus. 3 in Sancto-Domingo inter montem Christi et S.-Yago.

2. C. ERECTA (Turp. l. c. p. 80. t. 5. B.) foliis lanceolatis petiolatis, spinis infra-axillaribus. 3 in Antigua. Richard.

Ordo LIV. CORIARIEÆ.

Flores aut hermaphroditi aut sextis unius abortu monoici dioice. Calyx (aut Perigonium) 1-sepalus campanulatus 10-fidus, lobis 5 exterioribus majoribus ovatis, interioribus 5 alternis minoribus callosis. Petala 0. Stamina 10, è toro exserta, 5 inter calycis lobos externos et ovarii angulos, 5 inter calycis lobos callosos et ovarii sulcos. Filamenta filiformia. Antheræ oblongæ biloculares. Ovarium toro crassiusculò insidens, 5-loculare, 5-angulare. Stylus 0. Stigmata 5 ex ovarii apice longa subulata. Carpella 5 ad maturitatem subdiscreta approximata, indehiscentia, 1-sperma, glandulosis lobis grandefactis cincta. Semen in carpello pendulum. Albumen 0. Embryo rectus, radiculâ superâ, cotyledonibus 2 carnosis.

Frutices ramis subtetragonis oppositis, utrinquè sæpè ternis, ramulo nempè minore ad quodque ramuli primitivi latus orto. Folia opposita, simplicia, 3-nervia, integra, ovata aut cordata. Gemmæ squammosæ. Florum racemi ramos ramulosve terminantes, basi foliosi, simplices: pedicelli oppositi aut ad apice racemi alterni, basi bracteati, medio sæpius bi-bracteolati.

Ordo unico genere notissimo constans, sed quoad locum in serie naturali dubius. Veteres referebant ad Terebinthaceas, sed stamina hypogyna et embryo rectus repugnant. Bern. Jussæus ad Atripliceas, sed, si petalorum absentiam prætermittas, vix alia cernitur analogia, A. L. Jussæus olim ad Malpighiaceas et partium numerus tantum repugnat. Ad Zygophylleas etiam accedunt plurimis notis.

I. CORIARIA. *Niss. act. par.* 1711. t. 12.

Char. idem. ac ordinis.

1. *C. MYRTIFOLIA* (Lin. sp. 1467.) foliis ovato-lanceolatis acutis triplinerviis subpetiolatis glabris, racemis erectiusculis. ♂ in sepibus et dumetis Europæ et Africæ Mediterraneæ Lam. ill. t. 822. Duham. arb. 1. t. 73. (v. v.)
2. *C. MICROPHYLLA* (Poir. dict. 6. p. 87.) foliis ovatis obtusiusculis 5-nerviis subsessilibus, racemis nutantibus. ♂ in Peruvîâ. (v. s.)
3. *C. SARMENTOSA* (Forst. prod. 377.) diffusa, foliis cordato-ovatis acuminatis integerrimis 5-nerviis subpetiolatis, racemis nutantibus. ♂ in Novâ-Zelandiâ.
4. *C. RUSCIFOLIA* (Lin. sp. 1467.) foliis cordato-ovatis acutis sessilibus multinerviis, racemis nutantibus cylindraceis. ♂ in Peruvîâ. — Feuill. per. 3. p. 17. t. 12. Folia sæpè in ramis præcipuis terna, in ramulis opposita.

5. *C. PHYLLICIFOLIA* (Humb. in Willd. sp. 4. p. 819.) foliis subcordato-oblongis acutis petiolatis trinerviis venosis, pedunculis hispidis. 5 in Peruvia.
6. *C. THYMIFOLIA* (Humb. in Willd. l. c.) foliis oblongis acutis petiolatis trinerviis, petiolis pedunculisque hispidis. 5 in Peruvia. Vix à priori differt ex Willd.
7. *C. ATROPURPUREA* (fl. mex. ic. ined.) foliis cordato-ovatis acutis sessilibus trinerviis, racemis terminalibus simplicibus cylindraceis. 5 in Mexico. Flos, ex icone, 5-petalus 5-stylusque videtur.

FINIS PRIMÆ PARTIS.

INDEX ORDINUM ET GENERUM,

PARTIS PRIMÆ.

Abatia,	Pagin 503	Apatelia,	526
Ablania,	516	Apeiba,	514
Abroma,	485	Aphragmus,	209
Abuta,	103	Aporetica,	610
Acer,	593	Aquilegia,	50
Aceratium,	519	ARABIDÆ,	132
ACERINEÆ,	593	Arabis,	142
Achlys,	112	Arenaria,	400
Aconitum,	56	Argemone,	120
Actæa,	64	Ascyrum,	555
Adansonia,	478	Asimina,	87
Adlumia,	126	Aspicarpa,	583
Adonis,	23	Assonia,	498
Adrastæa,	73	Astrapæa,	500
Ægle,	538	Atalantia,	535
Æsculus,	597	Aubrietia,	158
Æthionema,	208	AURANTIACEÆ,	535
Agdestis,	103	Averrhoa,	689
Agliaia,	537	Ayenia,	487
Aldrovanda,	319	Azara,	262
Alectryon,	617		
Alegria,	516	Badiera,	334
Alliaria,	196	Balanites,	708
Almeidea,	729	Balsamina,	685
Alsodeia,	313	BALSAMINEÆ,	<i>ibid.</i>
ALSODINEÆ,	312	Banara,	259
ALSINEÆ,	388	Bauffya,	355
Althæa,	436	Banisteria,	587
Altheria,	493	BANISTERIÆ,	584
ALYSSINEÆ,	156	Barbarea,	140
Alyssum,	160	Baraldeia,	732
Amirola,	616	Beatsonia,	350
AMPELIDEÆ,	627	BERBERIDEÆ,	105
Ampelopsis,	632	Berberis,	<i>ibid.</i>
Anastatica,	185	Bergera,	537
ANASTATICEÆ,	<i>ibid.</i>	Bergia,	390
ANCHONIEÆ,	212	Berrya,	517
Anchonium,	<i>ibid.</i>	Berteroæ,	158
Andreoskia,	190	Biebersteinia,	707
Androsæmum,	543	Billardiera,	345
Anemone,	16	Biophytum,	689
ANEMONEÆ,	10	Biscutella,	181
Anoda,	458	Bivonæa,	208
Anona,	83	Bixa,	259
ANONACEÆ,	<i>ibid.</i>	BIXINEÆ,	<i>ibid.</i>
Anthodon,	569	Blighia,	609
Antholoma,	585	Bocconia,	121
Antichorus,	504	Boleum,	223

BOMBACEÆ,	475	Cheiranthus,	135
Bombax,	478	Cheirostemon,	480
Boronia,	721	Chelidonium,	122
Boscia,	244	Cherleria,	421
Brachycarpæa,	236	Chitonia,	707
BRACHYCARPÆÆ,	235	CHLENACEÆ,	521
Brassica,	213	Chloromyron,	564
BRASSICÆÆ,	<i>ibid.</i>	Chloroxylon,	625
Braya,	141	Choisya,	724
Bredemeyera,	340	Chorisia,	480
Buffonia,	388	Chorisporea,	186
Bunchosia,	581	Christiania,	516
BUNIADEÆ,	229	Chrysopia,	563
Bunias,	<i>ibid.</i>	Cissampelos,	100
Burasia,	96	Cissus,	627
Bursaria,	347	CISTINEÆ,	263
Byblis,	319	Cistus,	263
Byrsonima,	579	Citrus,	539
Byttneria,	486	Clausena,	538
BYTTNERIACEÆ,	481	CLEMATIDÆÆ,	2
BYTTNERIÆÆ,	484	Clematis,	<i>ibid.</i>
		Cleome,	238
Cabomba,	112	CLEOMEÆÆ,	237
Cadaba,	244	Cleomella,	<i>ibid.</i>
Cakile,	185	Cleyera,	524
CALILINEÆ,	<i>ibid.</i>	Clusia,	538
Calepina,	225	CLUSIÆÆ,	557
Calodendron,	712	Clypeola,	165
CALOPHYLLEÆ,	561	Cocculus,	96
Calophyllum,	562	Cochlearia,	172
Caltha,	44	Cochlospermum,	527
Calypttrion,	288	Colbertia,	75
Camarea,	584	Columbia,	512
Camelina,	201	Comesperma,	334
CAMELINEÆÆ,	<i>ibid.</i>	Commersonia,	486
Camellia,	529	Conohoria,	312
CAMELLIÆÆ,	<i>ibid.</i>	Cookia,	537
Caudollea,	73	Coptis,	47
Canella,	563	Corchorus,	504
CAPPAREÆÆ,	242	Cordylocarpus,	186
CAPPARIDÆÆ,	237	Coriaria,	739
Capparis,	245	CORIARIÆÆ,	<i>ibid.</i>
Capsella,	177	Correa,	719
Carapa,	626	Corydalis,	126
Cardamine,	149	Cossignia,	614
Cardiospermum,	601	Crambe,	225
Carolinea,	478	Cratæva,	242
Carpodontos,	555	Cremolobus,	184
Carrichtera,	224	Cristaria,	453
Caryocar,	599	Crowea,	720
CARYOPHYLLEÆ,	351	CRUCIFERÆÆ,	151
Castela,	738	Cucubalus,	367
Caucanthus,	583	Cupania,	612
Cedrela,	624	Curatella,	70
CEDRELEÆÆ,	<i>ibid.</i>	CUSPARIÆÆ,	729
Ceranthra,	313	Cyminosma,	722
Cerastium,	414	Cysticapuos,	126
Ceratocephalus,	26		
Chamira,	231	Davilla,	69

Decadia,	520	Erythrochiton,	732
Delima,	69	ERYTHROSPERMEÆ,	257
DALLIÆ,	67	Erythrospermum,	<i>ibid.</i>
Delphinium,	51	ERYTHROXYLÆ,	573
Dentaria,	154	Erythroxyllum,	<i>ibid.</i>
Dianthus,	355	Espera,	517
Dicera,	520	EUCLIDÆ,	184
Diclytra,	125	Euclidium,	<i>ibid.</i>
Dictamnus,	712	Eucryphia,	556
Didesmus,	227	Eudema,	202
Diglottis,	732	Eunomia,	208
DILLENEÆ,	70	Euphoria,	611
Dillenia,	75	Eurya,	525
DILLENIAEÆ,	67	Furyale,	114
Dionæa,	320	Evodia,	724
Diosma,	713	Eystathes,	618
DROSMEÆ,	709		
Diphyllia,	110	Fagonia,	704
DIPLECOLOREÆ,	230	Farsetia,	157
Diplolæna,	719	Feronia,	538
Diplophractum,	514	Ficaria,	44
Dplotaxis,	221	Fissilia,	532
Discovium,	236	Flacourtia,	256
Dodonæa,	616	FLACOURTIANÆÆ,	255
DODONÆACEÆ,	615	FLACOURTIEÆ,	256
Doliocarpus,	69	Flindersia,	625
Dombeya,	498	Frankenia,	349
DOMBEYACEÆ,	497	FRANKENIAEÆ,	<i>ibid.</i>
Draba,	166	Freziera,	524
Drimys,	78	FREZIERÆÆ,	<i>ibid.</i>
Drosera,	317	Friesia,	520
DROSERACEÆ,	317	Fugosia,	457
Drosophyllum,	320	Fumaria,	129
Drymaria,	395	FUMARIAEÆ,	125
Drymis,	78		
Drypis,	388	Galipea,	730
Durio,	480	Calphimia,	582
		Garcinia,	560
Ekebergia,	623	GARCINIEÆ,	560
FLÆOCARPEÆ,	519	Garidella,	48
Elæocarpus,	<i>ibid.</i>	Caudichaudia,	584
Elaphrium,	723	GERANIAEÆ,	637
Elatine,	390	Geranium,	639
Elvasia,	738	Geruma,	619
Emplevrum,	718	Glaucium,	122
Euarthrocarpus,	228	Glossarrhen,	290
Enemion,	48	Glossostemon,	485
Enourea,	618	Gluta,	501
Epimedium,	110	Glycosmis,	538
Eranthis,	46	Godoya,	558
Eriodendron,	479	Gpethea,	501
Eriolæna,	501	Goldbachia,	212
Eriostemon,	720	Gomphia,	736
Erodium,	644	Gordonia,	528
Erophila,	172	GORDONIEÆ,	527
Eruca,	223	Gossypium,	456
Erucaria,	230	Gouffea,	388
EAUCARIEÆ,	<i>ibid.</i>	Grewia,	508
Erysimum,	196	Guaiaicum,	768

Guarea,	623	ILICINÆ,	77
Guatteria,	93	Illicium,	<i>ibid.</i>
Guazuma,	485	Impatiens,	686
Guichenotia,	489	Ingenhouzia,	474
GUTTIFERÆ,	557	Isatidæ,	209
Gynandropsis,	237	Isatis,	210
Gypsophila,	351	Isopyrum,	48
Gyrostemon,	515		
		Jeffersonia,	111
Hamadryas,	25	Jonhia,	571
Haronga,	541		
Heisteria,	532	Kadsura,	83
Helianthemum,	266	Keraudrenia,	489
Helicteres,	475	Kiggelaria,	257
Heliocarpus,	503	KROGGLARIÆ,	<i>ibid.</i>
Heliphila,	231	Kitaibelia,	436
HELIOPHILÆ,	<i>ibid.</i>	Kleinhowia,	488
HELLEBORÆ,	44	Knowltonia,	23
Helleborus,	46	Koelreuteria,	616
Hemistemma,	71	Krameria,	341
Hepatica,	22	Kydia,	500
Heritiera,	484		
Hermaunna,	493	Lacepedea,	571
HERMANNIÆ,	490	Laetia,	260
Hermupoa,	254	Lagunea,	474
Hesperis,	188	Lancretia,	555
Heteropteris,	591	Laplacea,	527
Heynea,	624	LAPLACÆ,	526
Hibbertia,	73	Larbrea,	395
Hibiscus,	446	Lardizabala,	95
HIPPOCASTANÆ,	597	LARDIZABALÆ,	<i>ibid.</i>
Hippocratea,	567	Larrea,	705
HIPPOCRATEACÆ,	<i>ibid.</i>	Lasianthera,	635
Hiptage,	583	LASIOFETALÆ,	488
HIPTAGÆ,	<i>ibid.</i>	Lasiopetalum,	489
Hiraa,	585	Lavatera,	438
Holosteum,	393	Lavradia,	314
Honckenya,	506	Lebretonia,	446
Hortia,	732	Lechea,	285
Hudsonia,	284	Ledocarpum,	702
Hugonia,	522	Leea,	635
Humiria,	619	LEEACÆ,	<i>ibid.</i>
Hutchinsia,	177	Leontice,	109
Hybanthus,	311	LEPIDINÆ,	202
Hydnocarpus,	257	Lepidium,	203
Hydrastis,	23	Leptaleum,	200
HYDROPHILTIDÆ,	112	Leptocarpea,	140
Hydropeltis,	<i>ibid.</i>	Leptolæna,	521
Hydropityon,	422	Lettsomia,	521
Hymenanthera,	314	Limonia,	531
Hymenella,	389	LINEÆ,	421
Hypocoum,	123	Linum,	<i>ibid.</i>
Hypelate,	614	Liriodendron,	8
HYPERICÆ,	543	Lopimia,	45
HYPERICINÆ,	541	Ludia,	26
Hypericum,	543	Luhea,	51
		Lunaria,	11
Iberis,	178	Luxemburgia,	31
Icacina,	534	Lychnis,	31

Macanea,	564	Morisonia,	254
Macoubea,	<i>ibid.</i>	Moronobeia,	563
Macropodium,	149	Mundia,	337
Mærua,	254	Muntingia,	514
Magallana,	684	Muraltia,	335
Magnolia,	79	Muricaria,	225
MAGNOLIACEÆ,	77	Murraja,	537
MAGNOLIÆ,	79	Myagrum,	212
Mahernia,	496	Myosurus,	25
Mahonia,	108	Myrodia,	477
Mahurea,	557		
Malachodendron,	528	Nandina,	109
Malachra,	440	Naravelia,	10
Malcomia,	186	Nasturtium,	137
Malope,	429	Negundo,	596
Malpighia,	577	Nelumbium,	113
MALPIGHIACEÆ,	<i>ibid.</i>	NELUMBONÆ,	<i>ibid.</i>
MALPIGHIÆ,	<i>ibid.</i>	Neslia,	202
Malva,	530	Neurolooma,	156
MALVACEÆ,	429	Niebuhria,	243
Malvaviscus,	445	Nigella,	48
Mammea,	561	Niota,	592
Marcgravia,	565	Noisettia,	290
MARCGRAVIACEÆ,	<i>ibid.</i>	Norantea,	566
MARCGRAVIÆ,	<i>ibid.</i>	NORANTEÆ,	<i>ibid.</i>
Marialva,	560.	Notoceras,	140
Marila,	558	NOTORHIZÆ,	186
Matayba,	609	Nymphæa,	114
Mathiola,	132	NYMPHÆACEÆ,	113
Matisia,	477	NYMPHÆÆ,	114
Mayna,	79		
Meconopsis,	120	Ochna,	735
Megacarpæa,	183	OCHNACEÆ,	<i>ibid.</i>
Melhania,	499	Ochrocarpos,	560
Melia,	621	Ochroma,	480
MELLACEÆ,	619	Ochthodium,	184
Melianthus,	708	OLACINÆÆ,	531
MELIÆ,	<i>ibid.</i>	Olaæ,	<i>ibid.</i>
Melicocca,	614	Ophelus,	478
Melicope,	723	Ortegaia,	388
Melicytus,	257	ORTHOPLOCEÆ,	213
Melochia,	490	OXALIDEÆ,	689
Meniocus,	165	Oxalis,	690
MENISPERMACEÆ,	95		
MENISPERMÆÆ,	96	Pachynema,	70
Menispermum,	102	Pæonia,	65
Menonvillea,	184	PÆONIACEÆ,	64
Mesua,	562	Palavia,	458
Michelia,	79	Papaver,	117
Micranthera,	560	PAPAVERACEÆ,	<i>ibid.</i>
Melhringia,	390	Parnassia,	320
Mollugo,	391	Patrisia,	255
Monniera,	729	PATRISIÆ,	<i>ibid.</i>
Monnina,	338	Paulonia,	604
Monodora,	87	PAULLINIÆ,	601
Monsonia,	638	Pavia,	598
Montezuma,	477	Pavonia,	442
Morettia,	185	Pedicellia,	618
Moricandia,	221	Peganum,	712


Pelargonium,	649	Ricotia,	157
Peltaria,	166	Riedleia,	490
Pentaloba,	314	Rinorea,	312
Pentapetes,	498	Romeria,	122
Periptera,	459	Romanzowia,	319
Peritoma,	237	Roridula,	320
Petrocallis,	166	Roumea,	256
Phebalium,	719	Ruizia,	497
Philotheca,	721	Ruta,	709
Physa,	393	RUTACEÆ,	<i>ibid.</i>
Physiphora,	314	Ruychia,	566
Pigea,	307	Ryanma,	255
Pilocarpus,	728		
Piparea,	316	Sagina,	389
PITTOSPOREÆ,	345	Salacia,	570
Pittosporum,	346	Salomonina,	333
Plagianthus,	477	Sandoricum,	621
Pleurandra,	71	Sanguinaria,	121
PLEURORHIZÆÆ,	132	SAPINDACEÆ,	601
PODOPHYLLACEÆ,	111	SAPINDEX,	607
PODOPHYLLÆÆ,	<i>ibid.</i>	Sapindus,	607
Podophyllum,	<i>ibid.</i>	Saponaria,	365
Polanisia,	242	Sarcocapnos,	129
Polygala,	321	Sarcocolla,	521
POLYGALEÆ,	<i>ibid.</i>	Saurauja,	525
Pombalia,	306	SAURAUJÆÆ,	<i>ibid.</i>
Porcelia,	88	SAUVAGEÆÆ,	315
Portleria,	707	Sauvagenia,	<i>ibid.</i>
Pourretia,	477	Savignya,	157
Prockia,	260	Schepperia,	245
Pselium,	100	Schivereckia,	160
Pseudaleia,	533	Schizandra,	104
Pseudaleioides,	<i>ibid.</i>	SCHIZANDRÆÆ,	<i>ibid.</i>
Psychine,	224	Schizolæna,	521
PSYCHINÆÆ,	<i>ibid.</i>	Schizopetalum,	236
Pteroneurum,	154	Schmidelia,	610
Pterospermum,	500	Schouwia,	224
Pugionium,	185	Securidaca,	340
		Senacia,	347
Quassia,	733	Senebiera,	202
Quivisia,	620	Senra,	457
		Seriugia,	488
Racaria,	618	Serjania,	602
Raddisia,	570	Sethia,	576
Radiola,	428	Sida,	459
RANUNCULACEÆ,	2	Silene,	367
RANUNCULÆÆ,	25	SILENÆÆ,	351
Ranunculus,	26	Simaba,	733
RAPHANÆÆ,	225	Simaruba,	<i>ibid.</i>
Raphanus,	228	SIMARUBEÆÆ,	<i>ibid.</i>
Rapistrum,	227	Sinapis,	217
Raputia,	734	Singana,	564
Ratonia,	618	SISTMBRÆÆ,	186
Recchia,	70	Sisymbrium,	190
Redoutea,	457	Sloanea,	515
Rheddia,	564	Sobolewska,	212
RHIZOBOLEÆ	599	Sodada,	245
Rhodolæna,	522	Solea,	306
Rhynchotheca,	637	Soulamea,	335

ORDINUM ET GENERUM.

747

Sparmannia,	503	TREMANDREÆ,	343
Spergula,	394	Tribulus,	703
Spergulastrum,	421	Trichilia,	622
Spermaxyrum,	532	TRICHILIEÆ,	<i>ibid.</i>
Spiranthera,	728	Trichoa,	103
SPINOLOSEÆ,	229	Tricuspidaria,	520
Spirospermum,	96	Trigonia,	571
Stadmannia,	615	Triopteris,	586
Stalagmitis,	562	Triphaca,	483
Stanleya,	200	Triphasia,	535
Stauntonia,	96	Tristellateia,	583
Stellaria,	396	Triumfetta,	506
Stenopetalum,	201	Troaletia,	499
Stephania,	253	Trollis,	45
Sterculia,	481	TROPÆOLEÆ,	683
STERCULIEÆ,	<i>ibid.</i>	Tropæolum,	<i>ibid.</i>
Sterigma,	212	Turraea,	620
Stevenia,	141	Turritia,	141
Stewartia,	528		
Stigmarota,	257	Unona,	88
Strigilia,	621	Urena,	441
Subularia,	235	Urvillea,	602
SUBULARIEÆ,	<i>ibid.</i>	Uvaia,	88
Succowia,	224		
Swietenia,	625	Valotinia,	618
SYMPHONIEÆ,	563	Valis,	520
		Vatia,	517
Talauma,	81	Velia,	387
Talisia,	609	Vell,	223
Tasmannia,	78	VELLEÆ,	<i>ibid.</i>
Tauscheria,	210	Venbatia,	527
Teesdalia,	178	Vesaria,	159
Temus,	77	Vicia,	291
Ternstroemia,	523	VIOLEÆ,	287
TERNSTROEMIEÆ,	<i>ibid.</i>	Viole,	288
TERNSTAGMIEÆ,	<i>ibid.</i>	VINAE,	627
Tetracera,	67	Vism,	542
Tetratheca,	343	Vismæ,	541
Tetractis,	16	Vitis,	633
Tetrapteris,	587		
Thalictrum,	11	Wall,	737
Thea,	330	Wall,	501
Theobroma,	484	WALNIEÆ,	<i>ibid.</i>
Thespesia,	455	Walthe,	492
Thlaspi,	175	Wickamia,	517
THLASPIDÆÆ,	175	Worm,	75
Thomasia,	489		
Thouinia,	612	Xanthymus,	562
Thryallis,	583	Ximen,	533
Thylachium,	254	Xylop,	92
Ticorea,	730		
Tilia,	512	Xanthoa,	65
TILIACEÆ,	503	Xanthom,	725
Tina,	614	Zilla,	224
Toulicia,	612	Zieria,	722
Trachytella,	70	ZILLEÆ,	224
Tremandra,	244	ZYGOLLEÆ,	703
		Zygoph,	705

FINIS INDICI





Stanford University Libraries



3 6105 011 548 125

582
G219

Candolle,
Prodromus

NAME

DATE

Botany Dept.
Botany department.

80 Sept 1881

Aug 30 1881

17

